Striking Antibody Evasion Manifested by the Omicron Variant of SARS-CoV-2

Lihong Liu[1]*, Sho Iketani[1,2]*, Yicheng Guo[1]*, Jasper F-W. Chan[3,4]*, Maple Wang[1]*, Liyuan

Liu[5]*, Yang Luo[1], Hin Chu[3,4], Yiming Huang[5], Manoj S. Nair[1], Jian Yu[1], Kenn K-H. Chik[4],

Terrence T-T. Yuen[3], Chaemin Yoon[3], Kelvin K-W. To[3,4], Honglin Chen[3,4], Michael T. Yin[1,6],

Magdalena E. Sobieszczyk[1,6], Yaoxing Huang[1], Harris H. Wang[5], Zizhang Sheng[1],

Kwok-Yung Yuen[3,4], David D. Ho[1,2,6^]

[1]Aaron Diamond AIDS Research Center, Columbia University Vagelos College of Physicians and

Surgeons, New York, NY 10032, USA

[2]Department of Microbiology and Immunology, Columbia University Vagelos College of

Physicians and Surgeons, New York, NY 10032, USA

[3]State Key Laboratory of Emerging Infectious Diseases, Carol Yu Centre for Infection,

Department of Microbiology, Li Ka Shing Faculty of Medicine, The University of Hong Kong,

Pokfulam, Hong Kong Special Administrative Region, China

[4]Centre for Virology, Vaccinology and Therapeutics, Hong Kong Science and Technology Park,

Hong Kong Special Administrative Region, China

[5]Department of Systems Biology, Columbia University Vagelos College of Physicians and

Surgeons, New York, NY 10032, USA

[6]Division of Infectious Diseases, Department of Medicine, Columbia University Vagelos College

of Physicians and Surgeons, New York, NY 10032, USA

* These authors contributed equally

^ Address correspondence to David D. Ho (dh2994@cumc.columbia.edu, 212-304-6101, 701 W

168th St, 11th Floor, New York, NY 10032)

Word count: 3309

Abstract

The Omicron (B.1.1.529) variant of SARS-CoV-2 was only recently detected in southern Africa,

but its subsequent spread has been extensive, both regionally and globally[1]. It is expected to

become dominant in the coming weeks[2], probably due to enhanced transmissibility. A striking

feature of this variant is the large number of spike mutations[3] that pose a threat to the efficacy of

current COVID-19 vaccines and antibody therapies[4]. This concern is amplified by the findings

from our study. We found B.1.1.529 to be markedly resistant to neutralization by serum not only

from convalescent patients, but also from individuals vaccinated with one of the four widely used

COVID-19 vaccines. Even serum from persons vaccinated and boosted with mRNA-based

vaccines exhibited substantially diminished neutralizing activity against B.1.1.529. By evaluating

a panel of monoclonal antibodies to all known epitope clusters on the spike protein, we noted that

the activity of 18 of the 19 antibodies tested were either abolished or impaired, including ones

currently authorized or approved for use in patients. In addition, we also identified four new spike

mutations (S371L, N440K, G446S, and Q493R) that confer greater antibody resistance to

B.1.1.529. The Omicron variant presents a serious threat to many existing COVID-19 vaccines

and therapies, compelling the development of new interventions that anticipate the evolutionary

trajectory of SARS-CoV-2.

Main text

The COVID-19 pandemic rages on, as the causative agent, SARS-CoV-2, continues to evolve. Many diverse viral variants have emerged (**Fig. 1a**), each characterized by mutations in the spike protein that raise concerns of both antibody evasion and enhanced transmission. The Beta (B.1.351) variant was found to be most refractory to antibody neutralization[4] and thus compromised the efficacy of vaccines[5-7] and therapeutic antibodies. The Alpha (B.1.1.7) variant became dominant globally in early 2021 due to an edge in transmission[8] only to be replaced by the Delta (B.1.617.2) variant, which exhibited even greater propensity to spread coupled with a moderate level of antibody resistance[9]. Then came the Omicron (B.1.1.529) variant, first detected in southern Africa in November 2021[3,10,11] (**Fig. 1a**). It has since spread rapidly in the region, as well as to over 60 countries, gaining traction even where the Delta variant is prevalent. The short doubling time (2-3 days) of Omicron cases suggests it could become dominant soon[2]. Moreover, its spike protein contains an alarming number of >30 mutations (**Fig. 1b and Extended Data Fig. 1**), including at least 15 in the receptor-binding domain (RBD), the principal target for neutralizing antibodies. These extensive spike mutations raise the specter that current vaccines and therapeutic antibodies would be greatly compromised. This concern is amplified by the findings we now report.

Serum neutralization of B.1.1.529

We first examined the neutralizing activity of serum collected in the Spring of 2020 from COVID-19 patients, who were presumably infected with the wild-type SARS-CoV-2 (see Methods). Samples from 10 individuals were tested for neutralization against both D614G (WT) and B.1.1.529 pseudoviruses. While robust titers were observed against D614G, a significant drop (>32-fold) in ID_{50} titers was observed against B.1.1.529, with only 2 samples showing titers above the limit of detection (LOD) (**Fig. 1c and Extended Data Fig. 2a**). We then assessed the neutralizing activity of sera from individuals who received one of the four widely used COVID-19 vaccines: BNT162b2 (Pfizer), mRNA-1273 (Moderna), Ad26.COV2.S (Johnson & Johnson), and ChAdOx1 nCoV-19 (AstraZeneca) (see Methods and **Extended Data Table 1**). In all cases, a substantial loss in neutralizing potency was observed against B.1.1.529 (**Fig. 1d and Extended Data Fig. 2b-f**). For the two mRNA-based vaccines, BNT162b2 and mRNA-1273, a >21-fold and >8.6-fold decrease in ID_{50} was seen, respectively. We note that, for these two groups, we

specifically chose samples with high titers such that the fold-change in titer could be better

quantified, so the difference in the number of samples having titers above the LOD (6/13 for

BNT162b2 versus 11/12 for mRNA-1273) may be favorably biased. Within the Ad26.COV2.S

and ChAdOx1 nCOV-19 groups, all samples were below the LOD against B.1.1.529, except for

two Ad26.COV2.S samples from patients with a previous history of SARS-CoV-2 infection (**Fig.**

1d). Collectively, these results suggest that individuals who were previously infected or fully

vaccinated remain at risk for B.1.1.529 infection.

Booster shots are now routinely administered in many countries 6 months after full vaccination.

Therefore, we also examined the serum neutralizing activity of individuals who had received three

homologous mRNA vaccinations (13 with BNT162b2 and 2 with mRNA-1273). Every sample

showed lower activity in neutralizing B.1.1.529, with a mean drop of 6.5-fold compared to WT

(**Fig. 1d**). Although all samples had titers above the LOD, the substantial loss in activity may still

pose a risk for B.1.1.529 infection despite the booster vaccination.

We then confirmed the above findings by testing a subset of the BNT162b2 and mRNA-1273

vaccinee serum samples using authentic SARS-CoV-2 isolates: wild type and B.1.1.529. Again,

a substantial decrease in neutralization of B.1.1.529 was observed, with mean drops of >6.0-fold

and >4.1-fold for the fully vaccinated group and the boosted group, respectively (**Fig. 1e**).

Monoclonal antibody neutralization of B.1.1.529

To understand the types of antibodies in serum that lost neutralizing activity against B.1.1.529, we

assessed the neutralization profile of 19 well-characterized monoclonal antibodies (mAbs) to the

spike protein, including 17 directed to RBD and 2 directed to the N-terminal domain (NTD). We

included mAbs that have been authorized or approved for clinical use, either individually or in

combination: REGN10987 (imdevimab)[12], REGN10933 (casirivimab)[12], COV2-2196

(tixagevimab)[13], COV2-2130 (cilgavimab)[13], LY-CoV555 (bamlanivimab)[14], CB6 (etesevimab)[15],

Brii-196 (amubarvimab)[16], Brii-198 (romlusevimab)[16], and S309 (sotrovimab)[17]. We also

included other mAbs of interest: 910-30[18], ADG-2[19], DH1047[20], S2X259[21], and our antibodies 1-

20, 2-15, 2-7, 4-18, 5-7, and 10-40[22-24]. The footprints of mAbs with structures available were

drawn in relation to the mutations found in B.1.1.529 RBD (**Fig. 2a**) and NTD (**Fig. 2b**). The risk

to each of the 4 classes[25] of RBD mAbs, as well as to the NTD mAbs, was immediately apparent.

Indeed, neutralization studies on B.1.1.529 pseudovirus showed that 18 of the 19 mAbs tested lost

neutralizing activity completely or partially (**Fig. 2c and Extended Data Fig. 3**). The potency of

class 1 and class 2 RBD mAbs all dropped by >100-fold, as did the more potent mAbs in RBD

class 3 (REGN10987, COV2-2130, and 2-7). The activity of S309 declined modestly, whereas

Brii-198 was spared. All mAbs in RBD class 4 lost neutralization potency against B.1.1.529 by at

least 10-fold, as did mAb directed to the antigenic supersite[26] (4-18) or the alternate site[23] (5-7) on

NTD. Strikingly, all four combination mAb drugs in clinical use lost substantial activity against

B.1.1.529, likely abolishing or impairing their efficacy in patients.

Approximately 10% of the B.1.1.529 viruses in GISAID[1] also contain an additional RBD mutation,

R346K, which is the defining mutation for the Mu (B.1.621) variant[27]. We therefore constructed

another pseudovirus (B.1.1.529+R346K) containing this mutation for additional testing using the

same panel of mAbs (**Fig. 2d**). The overall findings resembled those already shown in **Fig. 2c**,

with the exception that the neutralizing activities of S309 and Brii198 were further diminished or

abolished. In fact, the entire panel of antibodies was essentially rendered inactive against this

minor form of the Omicron variant.

The fold changes in IC_{50} of the mAbs against B.1.1.529 and B.1.1.529+R346K relative to D614G

are summarized in the first two rows of **Fig. 3a**. The remarkable loss of activity observed for all

classes of mAbs against B.1.1.529 suggest that perhaps the same is occurring in the serum of

convalescent patients and vaccinated individuals.

Key mutations conferring antibody resistance

To understand the specific B.1.1.529 mutations that confer antibody resistance, we next tested

individually the same panel of 19 mAbs against pseudoviruses for each of the 34 mutations

(excluding D614G) found in B.1.1.529 or B.1.1.529+R346K. Our findings not only confirmed the

role of known mutations at spike residues 142-145, 417, 484, and 501 in conferring resistance to

NTD or RBD (class 1 or class 2) antibodies[4] but also revealed several mutations that were

previously not known to have functional importance (**Fig. 3a and Extended Data Fig. 4**). Q493R

mediated resistance to CB6 (class 1) as well as to LY-CoV555 and 2-15 (class 2), findings that

could be explained by the abolishment of hydrogen bonds due to the long side chain of arginine
and induced steric clashes with CDRH3 in these antibodies (**Fig. 3b, left panels**). Both N440K
and G446S mediated resistance to REGN10987 and 2-7 (class 3), observations that could also be
explained by steric hindrance (**Fig. 3b, middle panels**). The most striking and perhaps unexpected
finding was that S371L broadly affected neutralization by mAbs in all 4 RBD classes (**Fig. 3a and**
Extended Data Fig. 4). While the precise mechanism of this resistance is unknown, in silico
modeling suggested two possibilities (**Fig. 3b, right panels**). First, in the RBD-down state,
mutating Ser to Leu results in an interference with the N343 glycan, thereby possibly altering its
conformation and affecting class 3 antibodies that typically bind this region. Second, in the RBD-
up state, S371L may alter the local conformation of the loop consisting of S371-S373-S375,
thereby affecting the binding of class 4 antibodies that generally target a portion of this loop (REF
10-40). It is not clear how class 1 and class 2 RBD mAbs are affected by this mutation.
150
Evolution of antibody resistance of SARS-CoV-2 variants
To gain insight into the antibody resistance of B.1.1.529 relative to previous SARS-CoV-2 variants,
we evaluated the neutralizing activity of the same panel of neutralizing mAbs against
pseudoviruses for B.1.1.7[8], B.1.526[28], B.1.429[29], B.1.617.2[9], P.1[30], and B.1.351[31]. It is evident
from these results (**Fig. 4** and **Extended Data Fig. 5**) that previous variants developed resistance
only to NTD antibodies and class 1 and class 2 RBD antibodies. Here B.1.1.529, with or without
R346K, has made a big mutational leap by becoming not only nearly completely resistant to class
1 and class 2 RBD antibodies, but also substantial resistance to both class 3 and class 4 RBD
antibodies. B.1.1.529 is now the most complete "escapee" from neutralization by currently
available antibodies.
161
Discussion
The Omicron variant struck fear almost as soon as it was detected to be spreading in South Africa.
That this new variant would transmit more readily has come true in the ensuing weeks[2]. The
extensive mutations found in its spike protein raised concerns that the efficacy of current COVID-
19 vaccines and antibody therapies might be compromised. Indeed, in this study, sera from
convalescent patients (**Fig. 1c**) and vaccinees (**Figs. 1d and 1e**) showed markedly reduced
neutralizing activity against B.1.1.529. These findings are in line with emerging clinical data on

the Omicron variant demonstrating higher rates of reinfection[11] and vaccine breakthroughs. In

fact, recent reports showed that the efficacy of two doses of BNT162b2 vaccine has dropped from

over 90% against the original SARS-CoV-2 strain to approximately 40% and 33% against

B.1.1.529 in the United Kingdom[32] and South Africa[33], respectively. Even a third booster shot

may not adequately protect against Omicron infection[32,34], but of course it is advisable to do so.

Vaccines that elicited lower neutralizing titers[35,36] are expected to fare worse against B.1.1.529.

The nature of the loss in serum neutralizing activity against B.1.1.529 could be discerned from our

findings on a panel of mAbs directed to the viral spike. The neutralizing activities of all four major

classes of RBD mAbs and two distinct classes of NTD mAbs are either abolished or impaired

(**Figs. 2c and 2d**). In addition to previously identified mutations that confer antibody resistance[4],

we have uncovered four new spike mutations with functional consequences. Q493R confers

resistance to some class 1 and class 2 RBD mAbs; N440K and G446S confer resistance to some

class 3 RBD mAbs; and S371L confers global resistance to many RBD mAbs via mechanisms that

are not yet apparent. While performing these mAb studies, we also observed that all the currently

authorized or approved mAb drugs are rendered weak or inactive by B.1.1.529 (**Figs. 2c and 3a**).

In fact, the Omicron variant that contains R346K seemingly flattens the antibody therapy

landscape for COVID-19 (**Fig. 2d and 3a**).

The scientific community has chased after SARS-CoV-2 variants for a year. As more and more

of them appeared, our interventions directed to the spike became increasingly ineffective. The

Omicron variant has now put an exclamation mark on this point. It is not too far-fetched to think

that this SARS-CoV-2 is now only a mutation or two away from being pan-resistant to current

antibodies, either monoclonal or polyclonal. We must devise strategies that anticipate the

evolutionary direction of the virus and develop agents that target better conserved viral elements.

Figure Legends

Fig. 1. Resistance of B.1.1.529 to neutralization by sera. a, Unrooted phylogenetic tree of

B.1.1.529 with other major SARS-CoV-2 variants. **b,** Key spike mutations found in the viruses

isolated in the major lineage of B.1.1.529 are denoted. **c,** Neutralization of D614G and B.1.1.529

pseudoviruses by convalescent patient sera. **d,** Neutralization of D614G and B.1.1.529

pseudoviruses by vaccinee sera. Within the four standard vaccination groups, individuals that were

vaccinated without documented infection are denoted as circles and individuals that were both

vaccinated and infected are denoted as triangles. Within the boosted group, Moderna vaccinees are

denoted as squares and Pfizer vaccinees are denoted as diamonds. **e,** Neutralization of authentic

D614G and B.1.1.529 viruses by vaccinee sera. Moderna vaccinees are denoted as squares and

Pfizer vaccinees are denoted as diamonds. For all panels, values above the symbols denote

geometric mean titer and the numbers in parentheses denote the sample size. *P* values were

determined by using a Wilcoxon matched-pairs signed-rank test (two-tailed).

Fig. 2. Resistance of B.1.1.529 to neutralization by monoclonal antibodies. a, Footprints of

RBD-directed antibodies, with mutations within B.1.1.529 highlighted in cyan. Approved or

authorized antibodies are bolded. The receptor binding motif (RBM) residues are highlighted in

yellow. **b,** Footprints of NTD-directed antibodies, with mutations within B.1.1.529 highlighted in

cyan. The NTD supersite residues are highlighted in light pink. **c,** Neutralization of D614G and

B.1.1.529 pseudoviruses by RBD-directed and NTD-directed mAbs. **d,** Neutralization D614G and

B.1.1.529+R346K pseudoviruses by RBD-directed and NTD-directed mAbs.

Fig. 3. Impact of individual mutations within B.1.1.529 against monoclonal antibodies. a,

Neutralization of pseudoviruses harboring single mutations found within B.1.1.529 by a panel of

19 monoclonal antibodies. Fold change relative to neutralization of D614G is denoted, with

resistance colored red and sensitization colored green. **b,** Modeling of critical mutations in

B.1.1.529 that affect antibody neutralization.

Fig. 4. Evolution of antibody resistance across SARS-CoV-2 variants. Neutralization of SARS-

CoV-2 variant pseudoviruses by a panel of 19 monoclonal antibodies. Fold change relative to

neutralization of D614G is denoted.

Fig. 1

Fig. 2

Fig. 3

a

Fold change in IC50 compared with WT	RBD mAbs																	NTD mAbs	
	Class 1				Class 2				Class 3					Class 4					
	CB6	Brii-196	1-20	910-30	REGN10933	COV2-2196	LY-CoV555	2-15	REGN10987	COV2-2130	S309	2-7	Brii-198	ADG-2	DH1047	10-40	S2X259	4-18	5-7
B.1.1.529	<-1000	-134	<-338	<-159	<-1000	-140	<-1000	<-1000	<-1000	-390	-6.1	-231	2.2	-43	-124	-11	-35	-125	-30
B.1.1.529 + R346K	<-761	-97	<-338	<-159	<-1000	-89	<-1000	<-1000	<-1000	<-988	-21	-109	<-32	-51	-167	-32	-16	-125	-33
A67V	1.1	1.0	-1.1	1.4	1.1	-1.0	1.1	1.1	1.1	1.2	-1.4	-1.1	-1.2	1.3	-1.3	-1.1	1.0	-1.6	-1.1
Del69-70	-1.4	-1.4	-1.6	-1.1	-1.8	-1.5	-1.4	-1.4	-1.7	-1.4	-2.2	-1.9	-2.3	-1.4	-3.3	-1.7	-1.3	-2.6	-9.4
T95I	-1.4	-2.0	-1.8	-1.7	-1.5	-1.6	-1.5	1.1	-2.0	-1.1	-2.3	-3.4	1.3	-2.5	-3.4	-1.9	-2.2	1.0	-9.5
G142D	-1.3	-1.4	-1.6	1.0	-1.6	-1.6	-1.7	-1.6	-1.9	-1.5	-2.9	-2.9	-1.5	-1.4	-2.8	-1.4	-1.5	<-125	-263
Del143-145	1.3	1.0	-1.2	1.4	1.3	1.6	1.3	1.5	1.1	-1.1	-1.9	1.2	-1.3	1.2	-2.0	-1.2	-1.2	<-125	-29
Del211	-2.4	-2.1	-1.6	-2.1	-1.5	-1.5	-1.4	-1.2	1.2	-1.2	-1.2	-1.3	-1.1	-1.9	-2.4	-1.6	-2.3	1.2	-9.1
L212I	-1.3	-1.8	-1.3	-1.6	-1.4	-1.4	-1.6	-1.3	-1.3	-1.4	-2.2	-1.9	-2.2	-1.7	-3.2	-2.0	-1.9	-7.2	-2.2
Ins214EPE	-2.4	-2.4	-2.2	-2.4	-2.8	-2.7	-2.3	-4.3	-3.0	-2.2	-3.0	-6.2	-2.7	-3.1	-2.9	-1.9	-3.3	-7.1	-15
G339D	-1.7	-1.6	-1.7	-1.4	-2.2	-1.7	-1.5	-1.4	-1.8	-1.6	-4.0	-1.9	-3.9	-1.6	-2.2	-1.5	-3.2	-4.5	-3.0
(R346K)	-1.5	-1.2	-1.3	1.0	-1.5	-1.3	-1.3	-1.4	-1.6	-2.9	-1.4	-1.0	-21	-1.1	-1.9	-1.2	-1.4	-1.4	-2.3
S371L	-19	-18	-15	-22	-10	-4.1	-2.9	-1.4	-25	-1.4	-12	-12	-17	-18	-49	-59	-23	-1.8	1.1
S373P	-1.9	-2.1	-1.6	-1.4	-1.9	-2.1	-2.0	-1.4	-1.9	-1.3	-2.3	-1.8	-2.5	-2.2	-5.1	-5.0	-2.8	-8.2	-5.0
S375F	1.7	1.6	1.6	1.5	2.1	1.9	1.9	2.6	1.2	1.5	-1.1	1.4	1.1	1.8	-1.8	-1.2	-1.6	-9.2	-1.6
K417N	<-761	-1.6	-2.3	<-158	-6.4	1.1	1.5	1.1	1.2	1.2	-1.8	1.5	-1.0	-1.1	-1.9	-1.5	-1.8	-5.3	-2.8
N440K	-1.4	-1.4	-1.6	-1.2	-1.7	-1.4	-1.4	-1.6	-246	-1.5	-2.3	-18	-1.6	1.1	-2.0	-1.3	-1.5	-4.3	-2.8
G446S	1.3	1.1	-1.1	1.2	-1.6	-1.1	-1.6	-3.0	-574	-3.7	-1.7	-50	-1.4	-1.6	-2.2	-1.4	-2.2	-3.9	-2.4
S477N	-1.8	-1.8	-1.7	-1.7	-2.4	-1.5	-1.5	-1.7	-2.9	-1.6	-1.9	-4.4	-2.4	-1.5	-2.3	-1.6	-2.2	-17	-5.1
T478K	1.2	1.1	1.4	1.6	1.3	-1.5	-1.4	-1.2	-1.6	1.1	-1.8	-2.6	-1.6	-1.2	-2.8	-1.3	-2.3	-3.3	-2.3
E484A	-2.8	-1.7	-1.8	-1.2	-4.8	-4.9	<-1000	<-1000	-1.6	-1.4	-1.4	-2.7	-1.9	-1.6	-1.5	-1.9	-1.9	-5.7	-2.9
Q493R	-16	-7.3	-3.2	2.9	-42	-4.2	<-1000	-705	-1.4	-1.1	-1.2	-1.9	-2.0	-1.6	-1.6	-1.6	-1.5	-4.0	-1.3
G496S	-1.3	1.3	1.1	1.1	1.0	1.1	1.0	-9.3	-6.2	-1.3	-1.4	1.4	-1.2	-1.2	-1.6	-1.1	-1.6	-2.6	-1.6
Q498R	-1.7	-1.2	1.1	1.4	-1.5	-1.1	-1.4	-1.0	-1.6	-1.4	-1.3	1.1	-1.2	2.4	-1.3	-1.2	-1.3	-1.5	-1.8
N501Y	-9.8	-1.2	-8.4	-16	-1.4	-1.5	-1.6	-1.2	-1.2	-1.1	-1.8	-1.5	-2.7	-1.8	-2.5	-1.9	-1.9	-20	-3.9
Y505H	-1.2	1.2	-1.3	-9.6	1.1	1.0	1.0	1.1	1.4	1.0	-1.4	1.7	1.1	1.3	-1.4	1.0	-1.2	-1.2	-1.1
T547K	-1.9	-2.0	-2.0	-1.9	-1.7	-1.3	-1.6	-1.7	-2.7	-1.6	-1.6	-4.3	-1.9	-1.7	-2.6	-1.5	-1.9	-2.7	-2.7
H655Y	-2.7	-3.1	-3.5	-2.7	-3.1	-2.0	-2.2	-8.6	-8.8	-1.7	-2.3	-13	-2.4	-2.1	-3.9	-3.3	-3.9	-23	-5.3
N679K	1.0	1.2	1.1	1.1	-1.1	-1.2	-1.2	-1.2	-1.9	-1.1	-1.3	-1.8	-1.7	-1.4	-2.4	-1.7	-1.7	-2.1	-2.7
P681H	-2.3	-2.1	-2.1	1.0	-2.4	-1.8	-2.2	-1.5	-1.5	-1.0	-1.6	-1.9	-1.5	-1.3	-2.3	-1.3	-1.3	-2.3	-2.4
N764K	-1.1	-1.5	-1.3	-1.1	-1.4	-1.4	-1.4	-2.1	-2.5	-1.5	-2.2	-4.3	-1.3	-1.4	-3.3	-2.1	-2.4	-2.3	-2.1
D769Y	1.3	1.1	1.0	1.2	-1.5	-1.0	-1.4	-1.4	-2.0	-1.3	-1.9	-2.5	-1.3	-1.1	-1.7	-1.2	-1.4	-3.1	-2.5
N856K	-10	-2.8	-1.3	-12	-2.2	-3.0	-1.1	-1.0	-1.4	-1.1	-1.2	-1.3	-2.3	-1.8	-4.4	-2.1	-2.5	-1.6	-1.9
Q954H	2.7	1.9	1.5	2.6	1.2	1.0	1.2	1.1	-1.1	1.2	-1.2	-1.4	-1.1	-1.1	-2.5	-1.0	-1.1	-2.3	-2.9
N969K	-5.4	-1.6	-1.1	-4.5	-1.3	-1.8	-1.1	-1.3	-1.6	-1.1	-1.4	-2.4	-1.4	-1.1	-2.3	-2.0	-2.4	-2.5	-2.0
L981F	3.2	3.3	2.1	4.6	2.4	2.5	2.2	1.9	1.3	2.5	-1.0	-1.5	8.6	2.8	1.1	2.0	2.1	-1.3	-1.5

Legend: >3 | <-3 | <-10 | <-100

b

Fig. 4

Fold change in IC50 compared with WT	RBD mAbs																	NTD mAbs	
	Class 1				Class 2				Class 3					Class 4					
	CB6	Brii-196	1-20	910-30	REGN10933	COV2-2196	LY-CoV555	2-15	REGN10987	COV2-2130	S309	2-7	Brii-198	ADG-2	DH1047	10-40	S2X259	4-18	5-7
B.1.1.7	-8.8	2.6	-5.2	-15	1.6	1.8	1.6	2.2	2.9	1.7	1.1	2.3	4.1	1.7	2.2	1.4	1.4	-5.1	-4.0
B.1.526	-1.0	1.1	-1.1	2.5	-4.5	-2.1	-590	-1329	1.8	1.2	2.9	1.8	-1.1	1.5	2.9	-2.2	1.4	4.5	-2.5
B.1.429	3.0	2.3	1.4	2.5	2.5	2.8	-590	-4.6	1.6	1.1	1.9	1.6	-2.4	2.0	2.9	1.3	3.3	-39	-59
B.1.617.2	2.1	1.2	-1.1	2.5	1.2	1.4	-590	-10	-1.8	-1.7	1.2	-1.1	-8.9	1.0	1.4	-1.8	-1.4	-39	-74
P.1	-196	2.2	-16	-60	-121	-2.0	-590	-1329	1.9	1.1	1.1	1.2	1.8	-1.0	3.0	-2.2	1.2	-39	-74
B.1.351	-196	2.0	-40	-60	-78	-2.5	-590	-1329	1.5	1.5	1.2	1.9	-1.5	1.0	3.0	-2.9	1.2	-39	-8.4
B.1.1.529	<-1000	-134	<-338	<-159	<-1000	-140	<-1000	<-1000	<-1000	-390	-6.1	-231	2.2	-43	-124	-11	-35	-125	-30
B.1.1.529 + R346K	<-761	-97	<-338	<-159	<-1000	-89	<-1000	<-1000	<-1000	<-988	-21	-109	<-32	-51	-167	-32	-16	-125	-33

Legend: | >3 | <-3 | <-10 | <-100 |

References

Shu, Y. & McCauley, J. GISAID: Global initiative on sharing all influenza data - from vision to reality. *Euro Surveill* **22**, doi:10.2807/1560-7917.ES.2017.22.13.30494 (2017).

Grabowski, F., Kochańczyk, M. & Lipniacki, T. Omicron strain spreads with the doubling time of 3.2—3.6 days in South Africa province of Gauteng that achieved herd immunity to Delta variant. *medRxiv*, doi:https://doi.org/10.1101/2021.12.08.21267494 (2021).

Scott, L. *et al.* Track Omicron's spread with molecular data. *Science*, eabn4543, doi:10.1126/science.abn4543 (2021).

Wang, P. *et al.* Antibody resistance of SARS-CoV-2 variants B.1.351 and B.1.1.7. *Nature* **593**, 130-135, doi:10.1038/s41586-021-03398-2 (2021).

Abu-Raddad, L. J., Chemaitelly, H., Butt, A. A. & National Study Group for, C.-V. Effectiveness of the BNT162b2 Covid-19 Vaccine against the B.1.1.7 and B.1.351 Variants. *N Engl J Med* **385**, 187-189, doi:10.1056/NEJMc2104974 (2021).

Madhi, S. A. *et al.* Efficacy of the ChAdOx1 nCoV-19 Covid-19 Vaccine against the B.1.351 Variant. *N Engl J Med* **384**, 1885-1898, doi:10.1056/NEJMoa2102214 (2021).

Sadoff, J. *et al.* Safety and Efficacy of Single-Dose Ad26.COV2.S Vaccine against Covid-19. *N Engl J Med* **384**, 2187-2201, doi:10.1056/NEJMoa2101544 (2021).

Davies, N. G. *et al.* Estimated transmissibility and impact of SARS-CoV-2 lineage B.1.1.7 in England. *Science* **372**, doi:10.1126/science.abg3055 (2021).

Mlcochova, P. *et al.* SARS-CoV-2 B.1.617.2 Delta variant replication and immune evasion. *Nature* **599**, 114-119, doi:10.1038/s41586-021-03944-y (2021).

Hadfield, J. *et al.* Nextstrain: real-time tracking of pathogen evolution. *Bioinformatics* **34**, 4121-4123, doi:10.1093/bioinformatics/bty407 (2018).

Pulliam, J. R. C. *et al.* Increased risk of SARS-CoV-2 reinfection associated with emergence of the Omicron variant in South Africa. *medRxiv*, doi:https://doi.org/10.1101/2021.11.11.21266068 (2021).

Hansen, J. *et al.* Studies in humanized mice and convalescent humans yield a SARS-CoV-2 antibody cocktail. *Science* **369**, 1010-1014, doi:10.1126/science.abd0827 (2020).

264   13   Zost, S. J. *et al.* Potently neutralizing and protective human antibodies against SARS-
265   CoV-2. *Nature* **584**, 443-449, doi:10.1038/s41586-020-2548-6 (2020).

266   14   Jones, B. E. *et al.* The neutralizing antibody, LY-CoV555, protects against SARS-CoV-2
267   infection in nonhuman primates. *Sci Transl Med* **13**, doi:10.1126/scitranslmed.abf1906
268   (2021).

269   15   Shi, R. *et al.* A human neutralizing antibody targets the receptor-binding site of SARS-
270   CoV-2. *Nature* **584**, 120-124, doi:10.1038/s41586-020-2381-y (2020).

271   16   Ju, B. *et al.* Human neutralizing antibodies elicited by SARS-CoV-2 infection. *Nature*
272   **584**, 115-119, doi:10.1038/s41586-020-2380-z (2020).

273   17   Pinto, D. *et al.* Cross-neutralization of SARS-CoV-2 by a human monoclonal SARS-CoV
274   antibody. *Nature* **583**, 290-295, doi:10.1038/s41586-020-2349-y (2020).

275   18   Banach, B. B. *et al.* Paired heavy- and light-chain signatures contribute to potent SARS-
276   CoV-2 neutralization in public antibody responses. *Cell Rep* **37**, 109771,
277   doi:10.1016/j.celrep.2021.109771 (2021).

278   19   Rappazzo, C. G. *et al.* Broad and potent activity against SARS-like viruses by an
279   engineered human monoclonal antibody. *Science* **371**, 823-829,
280   doi:10.1126/science.abf4830 (2021).

281   20   Li, D. *et al.* In vitro and in vivo functions of SARS-CoV-2 infection-enhancing and
282   neutralizing antibodies. *Cell* **184**, 4203-4219 e4232, doi:10.1016/j.cell.2021.06.021
283   (2021).

284   21   Tortorici, M. A. *et al.* Broad sarbecovirus neutralization by a human monoclonal
285   antibody. *Nature* **597**, 103-108, doi:10.1038/s41586-021-03817-4 (2021).

286   22   Liu, L. *et al.* Potent neutralizing antibodies against multiple epitopes on SARS-CoV-2
287   spike. *Nature* **584**, 450-456, doi:10.1038/s41586-020-2571-7 (2020).

288   23   Cerutti, G. *et al.* Neutralizing antibody 5-7 defines a distinct site of vulnerability in
289   SARS-CoV-2 spike N-terminal domain. *Cell Rep* **37**, 109928,
290   doi:10.1016/j.celrep.2021.109928 (2021).

291   24   Liu, L. *et al.* Isolation and comparative analysis of antibodies that broadly neutralize
292   sarbecoviruses. *bioRxiv*, doi:https://doi.org/10.1101/2021.12.11.472236 (2021).

293   25   Barnes, C. O. *et al.* SARS-CoV-2 neutralizing antibody structures inform therapeutic
294   strategies. *Nature* **588**, 682-687, doi:10.1038/s41586-020-2852-1 (2020).

26   Cerutti, G. *et al.* Potent SARS-CoV-2 neutralizing antibodies directed against spike N-
terminal domain target a single supersite. *Cell Host Microbe* **29**, 819-833 e817,
doi:10.1016/j.chom.2021.03.005 (2021).

27   Uriu, K. *et al.* Neutralization of the SARS-CoV-2 Mu Variant by Convalescent and
Vaccine Serum. *N Engl J Med*, doi:10.1056/NEJMc2114706 (2021).

28   Annavajhala, M. K. *et al.* Emergence and expansion of SARS-CoV-2 B.1.526 after
identification in New York. *Nature* **597**, 703-708, doi:10.1038/s41586-021-03908-2
(2021).

29   Zhang, W. *et al.* Emergence of a Novel SARS-CoV-2 Variant in Southern California.
JAMA **325**, 1324-1326, doi:10.1001/jama.2021.1612 (2021).

30   Faria, N. R. *et al.* Genomics and epidemiology of the P.1 SARS-CoV-2 lineage in
Manaus, Brazil. *Science* **372**, 815-821, doi:10.1126/science.abh2644 (2021).

31   Tegally, H. *et al.* Detection of a SARS-CoV-2 variant of concern in South Africa. *Nature*
592, 438-443, doi:10.1038/s41586-021-03402-9 (2021).

32   Andrews, N. *et al.* Effectiveness of COVID-19 vaccines against the Omicron (B.1.1.529)
variant of concern. *medRxiv*, doi:https://doi.org/10.1101/2021.12.14.21267615 (2021).

33   Wroughton, L. in *The Washington Post*   (2021).

34   Kuhlmann, C. *et al.* Breakthrough Infections with SARS-CoV-2 Omicron Variant
Despite Booster Dose of mRNA Vaccine. *SSRN*,
doi:https://dx.doi.org/10.2139/ssrn.3981711 (2021).

35   Garcia-Beltran, W. F. *et al.* mRNA-based COVID-19 vaccine boosters induce
neutralizing immunity against SARS-CoV-2 Omicron variant. *medRxiv*,
doi:https://doi.org/10.1101/2021.12.14.21267755 (2021).

36   Cromer, D. *et al.* Neutralising antibody titres as predictors of protection against SARS-
CoV-2 variants and the impact of boosting: a meta-analysis. *Lancet Microbe*,
doi:10.1016/S2666-5247(21)00267-6 (2021).

37   Baym, M. *et al.* Inexpensive multiplexed library preparation for megabase-sized
genomes. *PLoS One* **10**, e0128036, doi:10.1371/journal.pone.0128036 (2015).

38   Martin, M. Cutadapt Removes Adapter Sequences From High-Throughput Sequencing
Reads. *EMBnet.journal* **17**, 10, doi:10.14806/ej.17.1.200 (2011).

39 Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat Methods*
9, 357-359, doi:10.1038/nmeth.1923 (2012).
40 Robinson, J. T. *et al.* Integrative genomics viewer. *Nat Biotechnol* **29**, 24-26,
doi:10.1038/nbt.1754 (2011).
41 Chu, H. *et al.* Comparative tropism, replication kinetics, and cell damage profiling of
SARS-CoV-2 and SARS-CoV with implications for clinical manifestations,
transmissibility, and laboratory studies of COVID-19: an observational study. *Lancet*
Microbe **1**, e14-e23, doi:10.1016/S2666-5247(20)30004-5 (2020).
42 Krissinel, E. & Henrick, K. Inference of macromolecular assemblies from crystalline
state. *J Mol Biol* **372**, 774-797, doi:10.1016/j.jmb.2007.05.022 (2007).
335

Methods

Serum samples

Convalescent plasma samples were obtained from patients with documented SARS-CoV-2

infection approximately one month after recovery or later. These samples were collected at the

beginning of the pandemic in early 2020 at Columbia University Irving Medical Center, and

therefore are assumed to be infection by the wild-type strain of SARS-CoV-2[4]. Sera from

individuals who received two or three doses of mRNA-1273 or BNT162b2 vaccine were collected

at Columbia University Irving Medical Center at least two weeks after the final dose. Sera from

individuals who received one dose of Ad26.COV2.S or two doses of ChAdOx1 nCov-19 were

obtained from BEI Resources. Some individuals were also infected by SARS-CoV-2 in addition

to the vaccinations they received. Note that, whenever possible, we specifically chose samples

with high titers against the wild-type strain of SARS-CoV-2 such that the loss in activity against

B.1.1.529 could be better quantified, and therefore the titers observed here should be considered

in that context. All collections were conducted under protocols reviewed and approved by the

Institutional Review Board of Columbia University. Additional information for the vaccinee

samples can be found in **Extended Data Table 1**.

Monoclonal antibodies

Antibodies were expressed as previously described[22], by synthesis of VH and VL genes

(GenScript), transfection of Expi293 cells (Thermo Fisher), and affinity purification from the

supernatant by rProtein A Sepharose (GE). REGN10987, REGN10933, COV2-2196, and COV2-

2130 were provided by Regeneron Pharmaceuticals, Brii-196 and Brii-198 were provided by Brii

Biosciences, CB6 was provided by Baoshan Zhang and Peter Kwong (NIH), and 910-30 was

provided by Brandon DeKosky (MIT).

Variant SARS-CoV-2 spike plasmid construction

An in-house high-throughput template-guide gene synthesis approach was used to generate spike

genes with single or full mutations of B.1.1.529. Briefly, 5'-phosphorylated oligos with designed

mutations were annealed to the reverse strand of the wild-type spike gene construct and extended

by DNA polymerase. Extension products (forward-stranded fragments) were then ligated together

by Taq DNA ligase and subsequently amplified by PCR to generate variants of interest. To verify

the sequences of variants, NGS libraries were prepared following a low-volume Nextera

sequencing protocol[37] and sequenced on the Illumina Miseq platform (single-end mode with 50 bp

R1). Raw reads were processed by Cutadapt v2.1[38] with default setting to remove adapters and

then aligned to reference variants sequences using Bowtie2 v2.3.4[39] with default setting. Resulting

reads alignments were then visualized in Integrative Genomics Viewer[40] and subjected to manual

inspection to verify the fidelity of variants. Sequences of the oligos used in variants generation are

provided in **Extended Data Table 2**.

Pseudovirus production

Pseudoviruses were produced in the vesicular stomatitis virus (VSV) background, in which the

native glycoprotein was replaced by that of SARS-CoV-2 and its variants, as previously

described[24]. Briefly, HEK293T cells were transfected with a spike expression construct with PEI

(1 mg/mL) and cultured overnight at 37 °C under 5% CO_2, and then infected with VSV-G

pseudotyped ΔG-luciferase (G*ΔG-luciferase, Kerafast) one day post-transfection. Following 2 h

of infection, cells were washed three times, changed to fresh medium, and then cultured for

approximately another 24 h before supernatants were collected, centrifuged, and aliquoted to use

in assays.

Pseudovirus neutralization assay

All viruses were first titrated to normalize the viral input between assays. Heat-inactivated sera or

antibodies were first serially diluted in 96 well-plates in triplicate. Viruses were then added and

the virus-sample mixture was incubated at 37 °C for 1 h. Vero-E6 cells (ATCC) were then added

at a density of 3×10^4 cells per well and plates were incubated at 37 °C for approximately 10 h.

Luciferase activity was quantified by using the Luciferase Assay System (Promega) according to

the manufacturer's instructions. Neutralization curves and IC_{50} values were derived by fitting a

non-linear five-parameter dose-response curve to the data in GraphPad Prism version 9.2.

Authentic virus isolation and propagation

Authentic B.1.1.529 was isolated from a specimen from the respiratory tract of a COVID-19

patient in Hong Kong by Kwok-Yung Yuen and colleagues at the Department of Microbiology,

The University of Hong Kong. Isolation of wild-type SARS-CoV-2 was previously described[41].

Viruses were propagated in Vero-E6-TMPRSS2 cells and sequence confirmed by next-generation

sequencing prior to use.

Authentic virus neutralization assay

To measure neutralization of authentic SARS-CoV-2 viruses, Vero-E6-TMPRSS2 cells were first

seeded in 96 well-plates in cell culture media (DMEM + 10% FBS + 1% penicillin/streptomycin)

overnight at 37 °C under 5% CO_2 to establish a monolayer. The following day, sera or antibodies

were serially diluted in 96 well-plates in triplicate in DMEM + 2% FBS and then incubated with

0.01 MOI of wild-type SARS-CoV-2 or B.1.1.529 at 37 °C for 1 h. Afterwards, the mixture was

overlaid onto cells and further incubated at 37 °C under 5% CO_2 for approximately 72 h.

Cytopathic effects were then visually assessed in all wells and scored as either negative or positive

for infection by comparison to control uninfected or infected wells in a blinded manner.

Neutralization curves and IC_{50} values were derived by fitting a non-linear five-parameter dose-

response curve to the data in GraphPad Prism version 9.2.

Antibody footprint analysis and RBD mutagenesis analysis

The SARS-CoV-2 spike structure used for displaying epitope footprints and mutations within

emerging strains was downloaded from PDB (PDBID: 6ZGE). The structures of antibody-spike

complexes were also obtained from PDB (7L5B for 2-15, 6XDG for REGN10933 and

REGN10987, 7L2E for 4-18, 7RW2 for 5-7, 7C01 for CB6, 7KMG for LY-COV555, 7CDI for

Brii-196, 7KS9 for 910-30, 7LD1 for DH1047, 7RAL for S2X259, 7LSS for 2-7, and 6WPT for

S309). Interface residues were identified using PISA[42] using default parameters. The footprint for

each antibody was defined by the boundaries of all epitope residues. The border for each footprint

was then optimized by ImageMagick 7.0.10-31 (https://imagemagick.org). PyMOL 2.3.2 was used

to perform mutagenesis and to make structural plots (Schrödinger).

Acknowledgements

We are grateful Regeneron Pharmaceuticals, B. Zhang and P. Kwong (NIAID), and B. Dekosky (MIT) for antibodies. This study was supported by funding from the Gates Foundation, JPB Foundation, Andrew and Peggy Cherng, Samuel Yin, Carol Ludwig, David and Roger Wu, Health@InnoHK, and the National Science Foundation (MCB-2032259).

Author contributions

D.D.H. conceived this project. L.H.L., S.I., and M.W. conducted pseudovirus neutralization experiments. J.F-W.C., H.C., K.K-H.C., T.T-T.Y., C.Y., K.K-W.T., and H.C. conducted authentic virus neutralization experiments. Y.G. and Z.Z. conducted bioinformatic analyses. L.Y.L. and Y.M.H. constructed the spike expression plasmids. Y.L. managed the project. J.Y. expressed and purified antibodies. M.T.Y. and M.E.S. provided clinical samples. M.S.N. and Y.X.H. contributed to discussions. H.H.W., K-Y.Y., and D.D.H. directed and supervised the project. L.H.L., S.I., and D.D.H. analyzed the results and wrote the manuscript.

Competing interests

L.L., S.I., M.S.N., J.Y., Y.H., and D.D.H. are inventors on patent applications on some of the antibodies described in this manuscript.

Data and materials availability

All data are provided in the paper or the Supplementary Information.