Dynamic *in situ* confinement triggers ligand-free neuropeptide receptor

signaling

M. Florencia Sánchez[1], Marina S. Dietz[2], Ulrike Müller[3], Julian Weghuber[3,4], Karl

Gatterdam[1], Ralph Wieneke[1], Mike Heilemann[2], Peter Lanzerstorfer[3], Robert Tampé[1,*]

[1]Institute of Biochemistry, Biocenter, Goethe University Frankfurt, Max-von-Laue-Str. 9, 60438

Frankfurt am Main, Germany; [2]Institute of Physical and Theoretical Chemistry, Goethe University

Frankfurt, Max-von-Laue-Str. 7, 60438 Frankfurt am Main, Germany; [3]School of Engineering and

Environmental Sciences, University of Applied Sciences Upper Austria, Wels, Austria; [4]FFoQSI -

Austrian Competence Centre for Feed and Food Quality, Safety & Innovation, FFoQSI GmbH,

Technopark 1D, 3430 Tulln, Austria

*To whom correspondence may be addressed Email: tampe@em.uni-frankfurt.de

Abstract

Membrane receptors are central to cell-cell communication. Receptor clustering at the plasma membrane modulates physiological responses, and microscale receptor organization is critical for downstream signaling. Spatially restricted cluster formation of the neuropeptide Y_2 hormone receptor (Y_2R) was observed *in vivo*; however, the relevance of this confinement is not fully understood. Here, we controlled Y_2R clustering *in situ* by a multivalent chelator nanotool in prestructured matrices. Fast Y_2R enrichment in microscale arrays triggered a ligand-independent downstream signaling determined by an increase in cytosolic calcium, cell spreading and migration. We reveal that ligand-independent signaling by confinement differs from ligand-induced activation in the recruitment of arrestin-3 as downstream effector, which was recruited to confined regions only in presence of the ligand. The employed multivalent nanotool facilitated a dynamic receptor enrichment with high exchange in the confined regions, comparable to microscale condensates. This concept enables *in situ* organization of membrane receptors and the exploration of ligand-independent receptor signaling.

Keywords: G protein-coupled receptors / membrane organization / membrane proteins / phase segregation / receptor clustering

Introduction

Cells translate stimuli into biochemical signals through membrane receptors controlling multiple aspects of cell behavior, including migration (Kupperman *et al.*, 2000; Stallaert *et al.*, 2018), differentiation (Li & Rudensky, 2016; Luther & Cyster, 2001), apoptosis (Scott *et al.*, 2009), as well as infectious diseases and cancer (Boncompain *et al.*, 2019; Haqshenas & Doerig, 2019; Kawai & Akira, 2005; Pasquale, 2010; Pike *et al.*, 2018; Sebestyen *et al.*, 2020; Tsukiyama *et al.*, 2020). Receptors form dynamic assemblies, condensates, or clusters that modulate downstream signaling and thus the final physiological response. Upon activation, these receptors undergo transitions from freely diffusing monomers to less mobile nanoclusters and further to higher-order oligomers (Ojosnegros *et al.*, 2017; Su *et al.*, 2016). Recently, liquid-liquid phase separation (LLPS) has been considered as a general mechanism for higher-order receptor cluster formation. In these dynamic, liquid-like condensates driven by weak multivalent interactions, the likelihood of receptor activation increases as neighboring molecules can act together engaging ligands (Case *et al.*, 2019; Lyon *et al.*, 2021). In signal transduction, the mechanisms for receptor cluster formation and cluster behavior have become physiologically relevant topics. However, the role of mesoscale (hundreds of nanometers) receptor organization in signal transduction remains unsolved, mainly because concepts to trigger and analyze receptor clustering *in situ* are largely limited.

Nano- and microlithographic approaches have provided cell-compatible scaffolds to investigate confined ligand-receptor interactions (Manz & Groves, 2010; Nair *et al.*, 2011; Torres *et al.*, 2008). Various techniques, ranging from photolithography to polymer lift-off and microcontact printing (µCP), have yielded information on how topology and mobility of the stimulus regulate cell responses (Bernard *et al.*, 2000; Doh & Irvine, 2006; Mossman *et al.*, 2005; Sánchez *et al.*, 2018; Traub *et al.*, 2016). So far, these methods utilize receptor ligands or antibodies to induce a rather static cluster formation. However, fast dynamic signaling at the plasma membrane requires tools that enable receptor clustering in a fast, specific, and *in situ* intervention of receptor networks.

Heterotrimeric guanine nucleotide-binding protein (G protein)-coupled receptors (GPCR)

are key cell surface proteins that regulate a plethora of cellular responses to external stimuli

(Rosenbaum *et al.*, 2009; Venkatakrishnan *et al.*, 2013; Wootten *et al.*, 2018). The Y_2 receptor

(Y_2R) is one of the four human neuropeptide Y (NPY) receptor subtypes, which belong to the

rhodopsin-like (class A) GPCR superfamily (Parker & Balasubramaniam, 2008; Tang *et al.*,

2021). Y_2R is linked to many important physiological processes, such as fear extinction

(Méndez-Couz *et al.*, 2021), regulation of food intake (Huang *et al.*, 2014), and obesity

(Lafferty *et al.*, 2021). Y_2R activation by neuropeptide Y (NPY) has been shown to promote

cell migration and proliferation (Ekstrand *et al.*, 2003; Movafagh *et al.*, 2006). Recently, we

demonstrated that Y_2Rs respond to light-guided microscale clustering at spatially defined

locations (Sánchez *et al.*, 2021). Y_2Rs are activated independently of the presence of the

canonical ligand, evoking an increase in cytosolic calcium, a change in cell spreading

behavior, and a localized migratory pattern.

Here, we established a versatile concept for *in situ* receptor clustering using a multivalent

chelator nanotool (*tris N*-nitrilotriacetic acid, *tris*NTA), which displays high affinity for histidine

(His)-tagged proteins. The small size of the chelator tool in combination with a prestructured

platform enabled high-content receptor clustering with high spatiotemporal resolution. The

lateral organization of Y_2Rs in living cells was controlled within minutes in a non-invasive

("traceless") and ligand-free manner. Microscale receptor condensates with a high degree of

homogeneity in size and density were generated in the plasma membrane. Receptor

clustering triggered ligand-independent activation of signal transduction, as evidenced by an

increase in cytosolic calcium and cell motility. By evaluating arrestin-3 (Arr3) recruitment as a

downstream signaling event, we uncovered high Arr3 co-recruitment to the patterned areas

only in the presence of the canonical ligand, suggesting an Arr3-independent desensitization

mechanism for the ligand-independent response. Strikingly, analysis of the receptor mobility

revealed a dynamic receptor assembly with high exchange within the confined areas, showing

similarities to liquid-like condensates. The versatility of this generic approach may advance

studies of ligand-free receptor signaling by dynamic confinement.

Results

High-content receptor clustering in living cells

We developed a system to induce dynamic receptor assembly *in situ* based on a multivalent chelator *tris*NTA nanotool (Fig 1A), which is equipped with a biotin moiety (*tris*NTA$^{PEG12-B}$) (Fig 1B). *tris*NTA$^{PEG12-B}$ displays a high affinity for His$_6$-tagged proteins ($K_D \approx 10$ nM), resulting in a site-specific but reversible interaction (Gatterdam *et al.*, 2018). Microcontact printing is a widely used method to investigate protein-protein interactions in living cells (Bernard *et al.*, 2000; Torres *et al.*, 2008). However, patterned substrates with a generic structure allowing high-content analyses of cells are difficult to produce. We used a large-area perfluoropolyether (PFPE) elastomeric stamps inked with bovine serum albumin (BSA) to print 96-well size glass (Hager *et al.*, 2021; Lanzerstorfer *et al.*, 2020). Wells within these plates containing a BSA-structured matrix were functionalized with biotinylated BSA (biotin-BSA) and streptavidin (SA) (Fig 1A). Subsequent functionalization with the nanotool and His$_6$-tagged fluorescent proteins, resulted in well-resolved protein patterns that were analyzed by confocal laser scanning microscopy (CLSM). This confirmed the specificity of the nickel-loaded *tris*NTA chelator to capture His$_6$-tagged proteins in defined regions of 1 µm or 3 µm diameter (Fig 1C, D).

The ability to control the organization of membrane receptors in real time is important for dissecting the spatial complexity of cell signaling and the extracellular environment. To this end, we utilized a tetracycline-inducible (T-Rex) expression system to establish a monoclonal human cervical cancer HeLa cell line expressing low amounts of the Y$_2$R (~300,000 receptors/cell) (Sánchez *et al.*, 2021). Y$_2$Rs displayed an N-terminal His$_6$-tag to the extracellular space and a cytosolic, C-terminal monomeric Enhanced Green Fluorescent Protein (mEGFP) (His$_6$-Y$_2$R^{mEGFP}, in brief Y$_2$R). It is worth mentioning that these modifications do not affect receptor activity, selectivity, or ligand binding as demonstrated (Sánchez *et al.*, 2021). In addition, studies determined that Y$_2$R does not need the N-terminus for ligand binding (Lindner *et al.*, 2009). Y$_2$R-positive cells properly adhered to 1 µm and 3 µm SA-functionalized matrices and showed a homogeneous receptor distribution at the basal plasma

membrane (Fig 1E). Addition of *tris*NTA$^{PEG12-B}$ (100 nM final) triggered receptor assembly.

Within five minutes, all cells showed receptor patterns at the plasma membrane comparable

in size and density (Fig 1E, F, Fig EV1). Importantly, recruitment of soluble His$_6$-tagged GFP

proteins as well as Y$_2$Rs to 1 μm prestructured spots led to analogous intensity profiles,

reflecting that similar densities were obtained in both cases (Fig 1D, G). In contrast, cells

expressing Y$_2$Rs lacking the His$_6$-tag (Y$_2$R^{mEGFP}) showed no receptor clustering after addition

of *tris*NTA$^{PEG12-B}$ (Fig EV2), demonstrating the specificity of the His$_6$-tag/*tris*NTA interaction in

living cells. Remarkably, ten minutes after receptor clustering by the multivalent nanotool, the

Y$_2$R enrichment resulted in an integrated receptor density equivalent to that of cells cultured

on matrices functionalized with anti-His$_6$ antibodies. However, a tenfold higher antibody

concentration (1 μM) was required compared to the multivalent nanotool, demonstrating its

efficacy in capturing His$_6$-tagged membrane receptors (Fig EV3). The 3 μm diameter

nanotool-induced receptor clusters presented a ninefold increase in integrated density

compared to 1 μm arrays, consistent with the increase in pattern area (Fig EV3). Overall, our

approach enabled versatile *in situ* receptor organization with high specificity.

Ligand-independent receptor clustering triggers fast signaling

To follow receptor clustering *in situ*, Y$_2$R-expressing cells were cultured on microstructured

matrices with a diameter of 1 μm, the smallest pattern we can produce and analyze with high

accuracy. After addition of the multivalent nanotool, the receptor redistribution was tracked by

confocal microscopy. Receptor clustering occurred in the first minutes and increased steadily

within 10 min, resulting in a twofold increase in receptor density compared to the initial state

(Fig 2A, Movie 1). Considering the average cell area of 1,420 ± 50 μm^2 (n = 66 cells) and the

enrichment factor (twofold), we estimated a receptor density of ~400 receptors/μm^2 in the

patterned regions (~300 receptors per 1 μm spot), a value comparable to other receptor

studies utilizing Fluorescence Correlation Spectroscopy methods (Bag *et al.*, 2015; Chen *et*

al., 2009). The kinetic profile for Y$_2$R recruitment to the 1 μm spots followed a pseudo-first-

order assembly rate of 0.20 ± 0.03 min^{-1} (Fig 2A, C). Despite of the high affinity and kinetically

stable binding (k_{off} = 0.18 h^{-1}) (Gatterdam *et al.*, 2018), the His-tag/*tris*NTA system relies on

molecular multivalency, which enables competition of the binding sites with histidine, thus

making the process of receptor assembly reversible. Addition of histidine to patterned cells led

to the disassembly of the receptor clusters in a rapid and complete manner, demonstrating a

key advantage of the approach to investigate receptor dynamics (Fig 2B).

Upon receptor clustering by the multivalent interaction nanotool, we observed a fast change

in cell spreading and motility. Analysis of cell spreading revealed a 20% increase in the total

cell area concomitant to receptor assembly (Fig 2A, D, Fig 3A), indicating ligand-independent

receptor activation upon clustering. Cells cultured on matrices in the absence of SA as well as

cells expressing Y_2R^{mEGFP} (lacking a His$_6$-tag) on SA-matrices showed no significant change

in cell spreading upon addition of the *tris*NTA$^{PEG12-B}$ (Fig EV4). As calcium signals are widely

known to regulate cell motility, we monitored local calcium dynamics utilizing a far-red cell-

permeable Ca^{2+}-sensitive dye. By dual-color imaging, receptor assembly and the cytosolic

calcium concentration were simultaneously recorded in living cells over the matrices. Upon

addition of *tris*NTA$^{PEG12-B}$, receptor recruitment led to a twofold increase in cytosolic calcium

concentration with a rapid rise within two minutes (Fig 3B, C), contrary to control matrices

without SA (Fig EV5). This indicates that calcium signals are associated with the motility

change. To correlate ligand-free *versus* canonical receptor activation, the agonist porcine

neuropeptide Y (pNPY, K_D = 5.2 ± 2.0 nM) was added to pre-patterned cells. An increase in

cytosolic Ca^{2+} was detected, which reached a plateau after one minute, followed by a slow

decay of the signal over a period of four minutes (Fig 3B, C). Strikingly, more receptors were

recruited to the patterned areas after activation via the agonist, as reflected by the increase in

integrated density in the patterned regions (Fig 3D). These results indicate that ligand-

activated Y_2Rs diffuse into areas where increased receptor density is present and

preferentially localized there. Recently, specific clusters termed GPCR hot spots (40-300 nm)

were visualized at the plasma membrane of living cells (Calebiro & Jobin, 2019; Chavez-

Abiega *et al.*, 2019; Hilger *et al.*, 2018; Sungkaworn *et al.*, 2017). These hot spots represent

regions where signaling preferentially occurs and where both receptors and G proteins are

enriched. We hypothesize that the induced microscale clusters may also trigger the formation

of hot spots providing the ideal environment for recruitment of more active receptors and thus

amplification of the signal. By increasing the local effective receptor concentration, this

complex organization may increase both the speed and efficiency of receptor-G protein

coupling while enabling local signaling transduction.

Downstream signaling by ligand-free receptor clustering

We next explored the impact of receptor clustering on downstream signaling. GPCR

desensitization involves a complex series of events, e.g. receptor phosphorylation, arrestin-

mediated internalization, receptor recycling, and lysosomal degradation (Ziffert et al, 2020).

Short-term desensitization occurs within minutes and is primarily associated with arrestin

preventing G protein interaction with the GPCR. Arrestins bind to activated, phosphorylated

GPCRs and block receptor-G protein interaction by steric hindrance at the receptor-coupling

interface, while serving as adaptors for key components of the endocytic machinery and

numerous signaling proteins (Hilger et al, 2018, Wang et al, 2020). In the presence of high

concentrations of the canonical ligand, an Arr3-dependent internalization, subsequent

endosomal sorting, and recycling of Y_2R to the cell membrane were observed (Walther *et al.*,

2010; Wanka *et al.*, 2018). However, recent studies demonstrated a strong and persistent

activation of the $G_{\alpha i}$-pathway upon Y_2R activation, which depletes the intracellular G protein

repertoire before Arr3 binding can terminate signaling (Ziffert et al, 2020). To assess whether

ligand-free clustering leads to Arr3 recruitment, we transfected cells stably expressing the Y_2R

with Arr3[mCherry] (in brief Arr3) and monitored Arr3 recruitment in real-time by total internal

reflection fluorescence (TIRF) microscopy (Fig 4A). Upon microscale receptor confinement by

*tris*NTA[PEG12-B], we did not observe a significant increase in Arr3 recruitment by intensity-

contrast analysis of the patterned spots, whereas a significant Arr3 recruitment was detected

upon addition of the agonist pNPY (Fig 4B, C). Again, an increase in Y_2R density in the

clustered areas was detected upon addition of pNPY (Fig 4B, D). In a subsequent step, a

partial dissipation of the Y_2R/Arr3 assemblies by specific competition with histidine was

observed (Fig 4B, C). Interestingly, half of the intensity in the patterned regions was dissipated

in the presence of the pNPY, indicating that not all receptors within the cluster areas are

associated with the nanotool upon addition of the ligand. In addition, patterning of Arr3 was

also detected in cells on an anti-His$_6$ antibody matrix within the first minutes after addition of

pNPY (Fig EV6). In this case, we did not observe a significant change in receptor density upon

addition of the pNPY, indicating that the high degree of immobilization and large size of the

antibody might restrict the transient enrichment of active receptors into the clustered regions.

Overall, our results reflect a difference between Arr3 recruitment in the ligand-free mode

compared to ligand-activated state. These observations indirectly confirm a high-affinity

interaction between the Y$_2$R and Gα_i (Ziffert et al, 2020) and suggest an active recruitment of

G proteins that delay Arr3 recruitment and impair termination of G protein signaling.

Receptor diffusion and dynamic exchange in confined regions

GPCR signaling results from dynamic interactions among receptors, G proteins, and the

complex surrounding membrane environment, which confers flexibility and versatility on this

fundamental biological process. GPCR hot spots are highly heterogeneous in terms of

diffusion of receptors and G proteins, which rapidly switch among phases of rapid and slow

diffusion (Calebiro & Jobin, 2019; Sungkaworn *et al.*, 2017). First, we examined whether Y$_2$R

clustering affects lipid diffusion and distribution by labeling the membrane with the lipid-like

dye CellMask. We observed a homogeneous staining of the plasma membrane,

demonstrating that receptor confinement does not affect membrane organization (Fig EV7).

To determine diffusion coefficients (*D*), we performed fluorescence recovery after

photobleaching (FRAP) technique. *In situ* receptor clustering was triggered on Y$_2$R-expressing

cells cultured on SA-matrices by addition of *tris*NTA$^{PEG12-B}$ (100 nM, 10 min). In a subsequent

step, the membrane was labeled and square-shaped regions of interest (ROIs) covering four

1 µm-sized spots were photobleached. Fluorescence recovery was analyzed by a FRAP

simulation approach that enabled calculation of diffusion coefficients independent of bleaching

geometry (Blumenthal *et al.*, 2015). The diffusion coefficient of lipids obtained by FRAP

showed an average value of D_{lipid} = 0.66 ± 0.10 µm^2/s in agreement with literature values for

free Brownian lipid diffusion at the plasma membrane (Schwille *et al.*, 1999; Wawrezinieck *et*

al., 2005).

To evaluate the Y$_2$R mobility in the ligand-free induced clusters, we determined the receptor

lateral diffusion of Y$_2$R-expressing cells cultured on SA-matrices before and after addition of

trisNTA$^{PEG12-B}$ (100 nM, 10 min). A square-shaped ROI covering four 1 µm-sized spots and

recovery was photobleached. A significant decrease in the lateral diffusion of the Y$_2$R was

observed at the basal membrane of cells after receptor confinement by *tris*NTA$^{PEG12-B}$ (D_{before}

= 0.25 ± 0.08 µm^2/s *versus* D_{after} = 0.10 ± 0.03 µm^2/s) (Fig 5A, E). Surprisingly, the receptor

density showed a high recovery within ~3 min after photobleaching (Fig 5A, Fig EV8, Movie 2).

Notably, no significant difference in receptor mobile fraction (M_f) before and after addition of

the nanotool was observed (M_f = 0.80 ± 0.04) (Fig 5B). Importantly, FRAP analyses of cells

cultured on matrices functionalized with anti-His$_6$ antibodies presented a significant decrease

in the receptor diffusion as well as in the M_f at the clustered spots ($M_{f,anti-His6\ Ab}$ = 0.56 ± 0.08)

(Fig 5B, Fig EV9, Movie 3). In contrast to monovalent antibody immobilization, receptor

clustering by the multivalent interaction nanotool is dynamic, allowing fast exchange of His$_6$-

tagged receptors or competition by histidine. We rationalized that free receptors can diffuse

into the clustered spots and exchange with photobleached receptors at multivalent binding

sites, leading to a dynamic confinement. Our results point out that a high receptor exchange

is possible even in micrometer-sized clusters, an effect that likely depends on cluster size,

with larger clusters showing less recovery (Sánchez *et al.*, 2021).

Finally, we investigated the lateral receptor mobility with a higher spatiotemporal resolution

using imaging fluorescence correlation spectroscopy (imFCS). FCS is used to study the

diffusion of membrane proteins in living cells with single-molecule sensitivity (Fig 5C). These

multiplexed FCS measurements are realized by analyzing many pixels simultaneously using

a widefield setup (Harwardt et al, 2018, Kannan et al, 2006). Regions of interests (ROIs) on

Y$_2$R-expressing cells cultured on SA-matrices were analyzed before and after addition of

*tris*NTA$^{PEG12-B}$. Enrichment of Y$_2$R at the basal membrane was observed with total internal

reflection fluorescence (TIRF) microscopy (Fig 5D). Consistent with the FRAP measurements,

a decrease in the Y_2R diffusion coefficient was determined upon cluster formation (D_{before} =

0.32 ± 0.06 µm^2/s and D_{after} = 0.16 ± 0.05 µm^2/s). The receptor diffusion coefficient measured

before clustering is comparable to similar membrane proteins (Lippincott-Schwartz et al,

2001), demonstrating that the microstructured confinement does not affect the mobile fraction

of receptors. In contrast to FRAP, imFCS provides a two-dimensional diffusion map, which

enables the determination of local differences in the lateral diffusion coefficient of membrane

receptors with high precision. Quantitative analysis of the 1 µm cluster spots in the acquired

ROIs resulted in a lateral diffusion coefficient of D_{spots} = 0.14 ± 0.03 µm^2/s (Fig 5E). Taking into

consideration that imFCS detects only mobile particles, we determined a similar decrease in

lateral diffusion in the patterned regions (Fig EV9) for cells cultured on matrices functionalized

with anti-His$_6$ antibodies. Taken together, we unravel that in microscale clusters, associations

between His$_6$-tagged Y_2Rs and multivalent $tris$NTA$^{PEG12-B}$ resulted in decreased diffusion rate

but dynamic receptor exchange with unchanged mobile fraction, which is similar to the

behavior described upon ligand-activated receptor clustering. This plasticity of the confined

receptors likely has a relevant impact on the efficiency of hot spot formation and GPCR

signaling.

Discussion

We developed a versatile concept to cluster receptors *in situ* with minimal steric hindrance

and disturbance. Ligand-independent receptor activation by confinement was unraveled by

cytosolic calcium increase and changes in cell spreading and motility. Addition of the canonical

ligand led to a similar calcium increase and interestingly, an amplification of the signal was

observed in the clustered areas concomitant with an increased receptor density. Surprisingly,

we revealed a difference in triggering of downstream signaling for the ligand-free *versus*

ligand-activated receptors as evidenced by co-recruitment of Arr3 to the clustered spots only

occurring in the presence of the canonical ligand. These findings are consistent with previous

results demonstrating an Arr3-dependent internalization, subsequent endosomal sorting, and

receptor recycling to the cell membrane in the presence of high concentrations of NPY

(Walther *et al.*, 2010; Wanka *et al.*, 2018). We hypothesize that high-affinity $Y_2R/G\alpha_i$

interactions drive the initial cell response, cytosolic calcium increase, and cell motility. High

local receptor density in the spots increases the residence time of proximate $G\alpha_i$ proteins and

recruits further downstream effectors, which boost the probability of activation (Sánchez *et al.*,

2021). Further, $Y_2R/G\alpha_i$ interactions lead to persistent activation of the $G\alpha_i$ pathway, which

depletes the intracellular $G\alpha_i$ protein repertoire before Arr3 binding can terminate signaling

(Ziffert *et al.*, 2020). The time frame of imaging after addition of the nanotool (30 min) suggests

a long-lasting $G\alpha_i$ protein activation and favors the hypothesis of a mechanism for Y_2R

activation and desensitization that is limited to the cell membrane and partially independent of

Arr3 recruitment (Ziffert *et al.*, 2020).

Many transmembrane receptors form dynamic assemblies that promote downstream signal

transduction in cells. Recently, dynamic liquid-like microscale clusters were identified as

critical modulators of cell signaling (Case *et al.*, 2019; Lyon *et al.*, 2021). In our system, the

transient association between the nanotool and the receptors revealed the generation of a

dynamic platform for cell signaling, which may have characteristics akin liquid-like clusters

generated by LLPS. The dynamic exchange of molecules within induced Y_2R microscale

clusters may contribute to hot spots formation and final downstream signaling. This

characteristic together with the lower concentration required compared to established systems

with immobilized ligands or antibodies, highlight the relevance of this versatile approach along

with the developed nanotool. Multiscale analyses of the multivalent interactions between

receptor condensates, G proteins, the lipid environment and actin–myosin assemblies are

critical to confirm the cluster behavior and dynamics. *In vitro* reconstitution systems utilizing

lipid bilayers have proven phase-segregation in two dimensions (Banjade & Rosen, 2014;

Ditlev *et al.*, 2019). The comparison of a mobile system based on model membranes and our

established platform, together with advanced quantitative fluorescence microscopy

techniques such as fluorescence resonance energy transfer (FRET) and single-molecule

localization microscopy will help us to decode condensates behavior and ligand-independent

signaling. In summary, the developed tool and matrices allow the investigation of ligand-

independent receptor activation *in situ* facilitating the investigation of early key processes in

cell signaling.

Materials and Methods

Synthesis of *trisNTA*^{PEG12-B}: Cyclam-Lys-*tris*NTA (Gatterdam *et al.*, 2018) (5.0 mg, 4.8 µmol), Biotin-PEG$_{12}$-NHS (23.0 mg, 24.0 µmol) and DIPEA (12.2 µL, 72.0 µmol) were dissolved in 0.5 ml dry DMF and stirred for 2 h at RT. After reaction, the volatile components were removed by lyophilization. Raw product was purified by semi-preparative RP-HPLC (mobile phase A: H$_2$O + 0.1% TFA, B: CAN + 0.1% TFA; gradient 5% to 80% B in 20 min; MZ-PerfectSil, 300 ODS, 5 µm, 250 x 10 mm, flow 4 ml/min). A biotin moiety was integrated into the nanotool for immobilization to SA in the prestructured matrices. The PEG$_{12}$ linker between the biotin and the *tris*NTA unit increased the flexibility of the molecule. *tris*NTA$^{PEG12-B}$ was confirmed by liquid chromatography-mass spectrometry (LC-MS, Waters BioAccord System). Dataset was recorded with an ACQUITY UPLC I-Class Plus chromatography system and ACQUITY RDa Detector, which was set to a cone voltage of 25 V, capillary voltage of 1.2 kV and a desolvation temperature of 500 °C operating in positive ionization mode. For reverse-phase separation, an ACQUITY UPLC Peptide BEH C18 column (300 Å, 1.7 µm, 2.1 mm x 100 mm) was used (Fig EV10). The *tris*NTA chelator compounds were dissolved in HBS buffer (20 mM HEPES-NaOH pH 7.5, 150 mM NaCl) and incubated with tenfold excess of NiCl$_2$. After 30 min incubation at 4 °C, the excess of Ni(II) was separated by ultra-centrifugal filters (Amicon Ultra-0.5 ml; 3 kDa and 10 kDa MWCO).

Microcontact printing. Large-area microcontact printing was performed as described previously (Lanzerstorfer *et al.*, 2020) with modifications. In short, a field of a large-area PFPE elastomeric stamp (1 µm grid size), obtained by the EV-Group (St. Florian am Inn, Upper Austria, Austria), was cut out, and washed by flushing with ethanol (100%) and distilled water. After drying with nitrogen, the stamp was incubated in 50 ml BSA solution (1 mg/ml, Sigma-Aldrich) for 30 min followed by washing the stamp with phosphate-buffered saline (PBS) and distilled water. After drying with nitrogen, the stamp was placed with homogeneous pressure onto a clean epoxy-coated glass substrate (Schott Nexterion Slide E) and incubated overnight

at 4 °C. The next day, the stamp was stripped from the glass with a forceps, and the

microstructured glass was bonded to a 96-well plastic casting using an adhesive tape (3M)

and closed with an appropriate lid.

Functionalization of the prestructured matrices. BSA-prestructured wells were incubated

with biotin-BSA (0.1 mg/ml, Thermo Fisher Scientific) and SA (1 µM, Sigma-Aldrich) in PBS,

each for 1 h at RT. Incubated wells were washed thoroughly with PBS after each step, to

remove unbound biotin-BSA and SA. For binding of soluble His-tagged proteins, wells were

incubated with *tris*NTA$^{PEG12-B}$ (0.5 µM) in HBS buffer for 1 h at RT. For nickel-loading, the

prestructured matrices were sequentially incubated with imidazole (1 M, 2 min), EDTA

(100 mM, 2 min), and NiCl$_2$ (10 mM, 5 min). Wells were carefully washed after each step.

Finally, HBS buffer containing EDTA (50 µM) was used to remove the free, non-complexed

nickel ions. His$_6$-GFP (100 nM) previously expressed and purified was added to the wells and

incubated for 30 min at RT. Experiments were performed in biological replicas (N=5).

Cell culture. HeLa Flp-In^{TM} T-Rex^{TM} Y$_2$R cells (His$_6$-Y$_2$R^{mEGFP} or Y$_2$R^{mEGFP}) were generated as

previously described (Sánchez *et al.*, 2021) and cultured at 37 °C, 5% CO$_2$, and 95% humidity.

For culturing the stable cell line, high glucose Dulbecco's modified Eagle's medium (DMEM)

(Gibco/Thermo Fisher Scientific) was supplemented with 10% tetracycline-free fetal calf

serum (FCS, Bio&Sell), blasticidin S HCl (1 µg/ml, Thermo Fisher Scientific) and

hygromycin B (50 µg/ml, Thermo Fisher Scientific). To induce receptor expression, the cell

medium was replaced with fresh medium containing tetracycline (0.1 µg/ml, Fluka) 18 h before

imaging. The same concentration of tetracycline resulting in an efficient plasma membrane

targeting was used for all the experiments. The cells were regularly tested for mycoplasma

contamination.

Receptor confinement in *real-time* by *tris*NTA$^{PEG12-B}$. Cells expressing Y_2R (His$_6$-Y_2R^{mEGFP} or Y_2R^{mEGFP}) were trypsinized and allowed to adhere to SA prestructured matrices for 3 h or overnight. 15-18 h prior to the experiment, the cell medium was replaced with fresh medium containing tetracycline (0.1 µg/ml) to induce receptor expression. The cells were visualized by CLSM in live-cell imaging solution (LCIS, Thermo Fisher Scientific) at 37 °C. Cells were subsequently incubated with nickel-loaded *tris*NTA$^{PEG12-B}$ (100 nM) in LCIS for 10-15 min at 37 °C. Excess of unbound *tris*NTA$^{PEG12-B}$ was removed by washing with LCIS. For reversibility experiments, micropatterned cells were incubated with histidine (5 mM) in LCIS for 2 to 10 min followed by washing with LCIS. Experiments were performed in biological replicas (N=4).

Receptor confinement on antibody-micropatterned matrices. Wells prestructured with BSA were subsequently incubated with biotin-BSA (0.1 mg/ml), SA (1 µM), and a biotinylated anti-His$_6$ antibody (1 µM) (ab106261, Abcam) in PBS for 1 h at RT. Wells were washed thoroughly with PBS to remove unbound antibody. Cells expressing Y_2R were trypsinized and seeded onto the antibody patterns. After 3 h, cells were visualized by CLSM in LCIS at 37 °C. Experiments were performed in biological replicas (N=5).

Time-lapse calcium imaging. 18 h after seeding the cells onto prestructured SA-matrices, cells were incubated with BioTracker 609 Red Ca^{2+} AM dye (3 µM, Merck Millipore) in fresh medium for 30 min. The cell-membrane permeable dye is de-esterified by cellular esterases and remains trapped in the cytosol. After incubation with the Ca^{2+} dye, cells were rinsed three times with PBS and imaged by CLSM in LCIS at 37 °C. For investigation of Ca^{2+} signal, time-lapse images were taken (5 slices z-stacks, 45-s interval) before and after addition of *tris*NTA$^{PEG12-B}$. Fluorescence intensity (($\lambda_{ex/em}$ 590/609 nm) of the dye changes depending on the intracellular Ca^{2+} concentration. Maximum intensity projections of single channels were analyzed. The ImageJ ROI tool was used to define the areas of the image to be analyzed. We consider a ROI covering the complete cell contour. Mean gray value (F) were background

subtracted and normalized to the fluorescence in cells before F_0. Experiments were performed

in biological replicas (N=3).

Plasma membrane staining. Live-cell membrane staining was performed directly after

receptor assembly in living cells grown on prestructured matrices. CellMask[TM] deep red

plasma membrane stain (Thermo Fisher Scientific) was used according to manufacturer's

instruction. 1 µl of the stock solution (1000x dilution) was dissolved in 1 ml of warm LCIS (final

concentration 5 µg/ml) and subsequently added to the cells, incubated for 5 min at 37 °C, and

washed with LCIS before visualization. Experiments were performed in biological replicas

(N=3).

Confocal laser scanning microscopy. Images were recorded by using a CLSM Zeiss LSM

880 (Carl Zeiss) equipped with a Plan-Apochromat 63x/1.4 Oil DIC M-27 objective. Sequential

settings for dual-color imaging were used. Excitation wavelengths for the different

fluorophores: 488 nm (argon laser) for mEGFP; 594 nm for the Ca^{2+} dye; 633 nm (helium-

neon laser) for the plasma membrane dye. Signals were detected after appropriate filtering on

a photomultiplier. Intensities of channels were adjusted over the whole image for better

visualization of overlap and exported by Zen blue (version 2.3 lite, Zeiss). Detector

amplification, laser power, and pinhole were kept constant for all studies.

Image analysis. Fluorescence images were processed with Zen blue, ImageJ, and Fiji

software (Schindelin *et al.*, 2012; Schneider *et al.*, 2012). All images were background

subtracted. Integrated density, mean gray value and cell area were obtained with ImageJ.

Graphs were plotted with OriginPro.

Fluorescence recovery after photobleaching. FRAP experiments were conducted at the CLSM Zeiss LSM 880 using 63 x/1.4 Oil DIC objective. Rectangular-shaped regions (6-10 µm radius) were bleached within 10 s with high laser intensities. Fluorescence recovery was monitored by repetitively imaging an area containing the photobleached region at 0.1 frame/s for ~ 150 s. For the analysis, a simulation approach that allows computation of diffusion coefficients regardless of bleaching geometry used in the FRAP series was applied (Blumenthal *et al.*, 2015). The method is based on fitting a computer-simulated recovery to actual recovery data of a FRAP series. The algorithm accepts a multiple-frame TIFF file, representing the experiment as input, and simulates the (pure) diffusion of the fluorescent probes (*2D* random walk) starting with the first post-bleach frame of the actual data. Once the simulated recovery is finished, the algorithm fits the simulated data to the real one and extracts the diffusion coefficient. The algorithm iteratively creates a series of simulated images, where each frame corresponds to a single iteration. The intensity values are extracted from the (user indicated) bleached area of the simulated frames, thus determining the general shape of the recovery curve. The "time" axis at this stage is in arbitrary units (iterations). To extract the diffusion coefficient, the simulated recovery curve needs to be fitted to the real recovery curve, by appropriately stretching the "time" axis. The time between frames in the actual data set is obviously known, thus once overlapping optimally the simulated curve with the real one, the duration of one iteration, in *real-time* units, is determined. The diffusion coefficient of the simulated series is then calculated according to eq. 1,

where D_s is the simulation-extracted diffusion coefficient, l is the step of a molecule in each

$$D_s = \frac{l^2}{4t_i} \tag{1}$$

iteration of the simulation, corresponding to one pixel in the image (the pixel size is calibrated previously, by imaging a known calibration sample), and t_i is the time interval between steps (determined as explained). Technically, the simulation proceeds until a plateau is reached (equilibration of the fluorescence intensity in the bleached area). The number of data points in the simulated recovery is typically different (larger) than the number of experimental points. In

addition, the real experimental data may not have been acquired until equilibration of

fluorescence. To determine t_i, the algorithm scans a range of possible values for the total

duration represented by the simulation and calculates a value X^2 for the goodness-of-fit

between the simulated data and the real FRAP data. Total simulation duration is selected as

the one that produces the minimal X^2. Experiments were performed in biological replicas

(N=3).

imFCS measurements. imFCS measurements were performed as described earlier

(Harwardt *et al.*, 2018; Harwardt *et al.*, 2017). A home-built widefield setup with total internal

reflection fluorescence (TIRF) illumination was used for imFCS analysis. The experimental

setup was equipped with a 488 nm diode laser (100 mW, Obis, Coherent, USA). The excitation

light passes through an acousto-optical tunable filter (AA Opto-Electronic, Orsay, France) and

a telescope consisting of two achromatic lenses (Thorlabs, USA) with f = -40 mm and 750 mm.

A third achromatic lens (f = 400 mm, Thorlabs) directed the excitation light to the TIRF mirror

and had its focus on the back focal plane of the objective. The TIRF mirror was placed on a

motorized translation stage (25 mm, #MTS25/M-Z8, Thorlabs) controlled by a motion

controller (K-Cube Brushed DC Servo Motor Controller, # KDC101, Thorlabs) to switch

between widefield and TIRF illumination. The light entered an Eclipse Ti microscope (Nikon,

Japan), was reflected by a dichroic mirror (TIRF-Quad filter set 405/488/561/640 consisting of

a QuadLine Laser Clean-up ZET405/488/561/640x, QuadLine dichroic zt405/488/561/640rpc,

QuadLine rejection band ZET405/488/561/640 TIRF, all AHF Analysentechnik AG, Tübingen,

Germany), and was directed by an oil-immersion TIRF objective (UapoN 100xOTIRF, 1.49

Oil, Olympus, Japan) onto the sample. A nosepiece stage (IX2-NPS, Olympus) was used for

z plane adjustment and drift minimization. Emission light was collected by the same objective

and passed the dichroic mirror. In the detection path a TwinCam (Acal Bfi, Germany) with a

BrightLine HC 525/45 bandpass filter (AHF Analysentechnik AG) was implemented and the

signal was detected by a scientific complementary metal-oxide semiconductor (sCMOS)

camera (Zyla 4.2, Andor, Belfast, UK). Data were collected using the open-source software

μManager (Edelstein *et al.*, 2010). For data acquisition the following settings were applied:

24 W/cm^2 laser intensity, a bit depth of 16 bit, pixel readout rate of 540 MHz, frame time 4 ms,

4x4 binning, and 4,000 frames per film. For each film, a 40x25 pixel (or 40x20 pixel) region of

interest (ROI) was chosen and the measurement was performed with TIRF illumination to

observe membrane diffusion of Y$_2$R. In total 36 untreated cells, 24 cells with *tris*NTA$^{PEG12-B}$

immobilized receptors, and 26 cells with anti-His$_6$ antibody immobilized receptors were

measured. Each condition contains data from at least three independent measurement days

(N=3).

imFCS data analysis. Analysis of imFCS films was performed using the imFCS plugin

(version 1.52) (Sankaran *et al.*, 2010) for Fiji (Schindelin *et al.*, 2012). The following correlation

settings were chosen: emission wavelength = 515 nm, NA = 1.49, correlator scheme P = 16

and Q = 8, lateral PSF = 0.8, binning = 1, pixel size = 5.75 μm, magnification = 25 for 4x4

binning, and linear segment bleach correction with linear segments of 500 frames. Diffusion

coefficients were obtained for each pixel by fitting the correlation curves according to the

literature (Sankaran *et al.*, 2010). To compare the overall diffusion coefficients with those of

the patterned regions, ROIs were placed around patterned regions and analyzed separately.

For further analysis, the pixelwise diffusion coefficients for all measurements were imported

into OriginPro 2019 (OriginLab Corporation, Northampton, USA). For box plots of diffusion

coefficients, median diffusion coefficients were determined for each cell. Mean diffusion

coefficients per condition were obtained by averaging over the median diffusion coefficients

per measurement and calculating the standard error of the mean. Two-sample t-tests (α =

0.05) were applied to compare the diffusion coefficients for the different conditions. All

datasets were tested for normality using the Kolmogorov-Smirnov test (α = 0.05). Significance

was assigned as follows: $p > 0.05$ no significant difference between populations (n.s.), $p <$

0.05 significant difference (*), $p < 0.01$ significant difference (**), and $p < 0.001$ significant

difference (***). Two-dimensional maps of diffusion coefficients were generated also in

OriginPro. Diffusion coefficients were color coded from light yellow to dark red in the range of

0 to 0.5 µm²/s. Pixels that yielded correlation curves with diffusion coefficients higher than

0.5 µm²/s are presented in black. Pixels that yielded correlation curves which could not be

fitted by the imFCS plugin in Fiji are shown in light grey. To generate frequency distribution

plots, diffusion coefficients were log-transformed and binned in the interval between -5.3 and

1.0 with a bin size of 0.1 for each cell. Logarithmic diffusion coefficients were re-transformed,

frequency counts were averaged over all cells per condition and normalized. Frequency

counts were plotted logarithmically against diffusion coefficients. Errors bars represent

standard errors of the mean.

Arr3 recruitment upon receptor confinement. Microstructured surfaces were functionalized

with biotin-BSA and SA or SA and anti-His$_6$ antibody as described before. For transient co-

transfection with Arr3^{mCherry}, cells were sub-cultured the day before and then transfected with

the Arr3^{mCherry} plasmid using the TurboFect^{TM} transfection reagent (Thermo Fisher Scientific,

Waltham, MA, USA), according to the manufacturer's instructions and induced with

Tetracycline (0.1 µg/ml) 18 h before microscopy. Cells co-expressing His$_6$-Y$_2$R^{mGFP} and Arr3

were seeded onto the microstructured matrices and visualized by total internal reflection

fluorescence (TIRF) microscopy in LCIS at 37 °C after 3 to 4 h to ensure a homogeneous cell

membrane adhesion, which is a prerequisite for quantitative TIRF microscopy. For antibody

experiments, cells grown on prestructured matrices were incubated with pNPY (10 nM, Tocris)

in LCIS for 30 min at 37 °C. For *tris*NTA$^{PEG12-B}$ experiments, cells grown on SA-matrices were

subsequently incubated with nickel-loaded *tris*NTA$^{PEG12-B}$ (100 nM) and pNPY (10 nM) in LCIS

for 30 min at 37 °C. For reversibility, cells were incubated with histidine (5 mM) in LCIS for

30 min. Experiments were performed in biological replicas (N=2).

Arr3 imaging by TIRF microscopy. The detection system was set up on an epi-fluorescence

microscope (Nikon Eclipse Ti2). For selective fluorescence excitation of mGFP and mCherry,

a multi-laser engine (Toptica Photonics, Munich, Germany) was used at 488 and 561 nm,

respectively. The samples were illuminated in total internal reflection (TIR) configuration

(Nikon Ti-LAPP) using a 60x oil immersion objective (NA = 1.49, APON 60XO TIRF). After

appropriate filtering using standard filter sets, the fluorescence was imaged onto a sCMOS

camera (Zyla 4.2, Andor, Northern Ireland). The samples were mounted on an x-y-stage

(CMR-STG-MHIX2-motorized table, Märzhäuser, Germany), and scanning was supported by

a laser-guided automated Perfect Focus System (Nikon PFS).

Contrast Quantification and Statistical Analysis. Contrast analysis was performed as

described previously (Lanzerstorfer *et al.*, 2014; Lanzerstorfer *et al.*, 2020; Schütz *et al.*,

2017). Initial imaging recording was supported by the Nikon NIS Elements software. Images

were exported as TIFF frames and fluorescence contrast analysis was performed using the

Spotty framework (Borgmann *et al.*, 2012). The fluorescence contrast <c> was calculated as

$<c> = (F^+ - F^-)/(F^+ - F_{bg})$, where F^+ denotes the intensity of the inner pixels of the pattern. F^-

shows the intensity of the surrounding pixels of the micropattern, and F_{bg} the intensity of the

global background. Data are expressed as the means ± SEM. Comparisons of more than two

different groups were performed using one-way ANOVA, which was followed by Tukey's

multiple comparisons test in GraphPad Prism software (version 9.1.2).

Acknowledgements

We thank Christian Winter, Andrea Pott, Inga Nold, and all members of the Institute of Biochemistry (Goethe University Frankfurt) for discussion and comments. We thank Dr. Annette Beck-Sickinger (Leipzig University) for the Y_2 receptor construct and Dr. Alina Klein (Goethe University Frankfurt) for the generation of the Y_2R^{mEGFP} constructs with and without His_6-tag. We tank Dr. Cornelius Krasel (Philipps University of Marburg) for the $Arr3^{mCherry}$ construct. We also thank Christian Winter for the LC-MS analysis. This work was supported by the German Research Foundation (GRK 1986 (No. 237922874) to R.W. and R.T.; CRC 807 (No. 57566863) P16 to R.T.), LOEWE DynaMem P3 to R.W. and R.T., the Volkswagen Foundation (Az. 96 498 to R.W., Az. 96 497 to M.H., and Az. 96 496 to R.T.); the Christian-Doppler Forschungsgesellschaft (Josef Ressel Centre for Phytogenic Drug Research, the Austrian Science Fund (FWF, I4972-B) and the FH Upper Austria Center of Excellence for Technological Innovation in Medicine (TIMed Center) to U.M., J.W. and P.L..

Author contributions

M.F.S. performed the cell-based assays and imaging experiments. M.S.D. carried out the imFCS experiments and analyzed the data together with M.H.. U.M., P.L., and J.W. prepared the prestructured surfaces, performed the Arr3 recruitment assays and the intensity-contrast analysis. K.G. synthesized and characterized the chelator compound. M.F.S., R.W., and R.T. wrote the manuscript with contributions from all authors. R.T. conceived the study.

Competing interest. The authors declare no competing interest.

Data availability. Data and movies are available in the supplementary materials.

References

Bag N, Huang S, Wohland T (2015) Plasma Membrane Organization of Epidermal Growth

Factor Receptor in Resting and Ligand-Bound States. *Biophys J* **109:** 1925-36.

Banjade S, Rosen MK (2014) Phase transitions of multivalent proteins can promote

clustering of membrane receptors. *Elife* **3**

Bernard A, Renault JP, Michel B, Bosshard HR, Delamarche E (2000) Microcontact printing

of proteins. *Advanced Materials* **12:** 1067-70.

Blumenthal D, Goldstien L, Edidin M, Gheber LA (2015) Universal Approach to FRAP

Analysis of Arbitrary Bleaching Patterns. *Sci Rep* **5:** 11655.

Boncompain G, Herit F, Tessier S, Lescure A, Del Nery E, Gestraud P, Staropoli I, Fukata Y,

Fukata M, Brelot A, Niedergang F, Perez F (2019) Targeting CCR5 trafficking to inhibit

HIV-1 infection. *Sci Adv* **5:** eaax0821.

Borgmann D, Weghuber J, Schaller S, Jacak J, Winkler SM (2012) Identification of patterns

in microscopy images of biological samples using evolution strategies. In *Proceedings*

of the 24th European Modeling and Simulation Symposium.

Calebiro D, Jobin ML (2019) Hot spots for GPCR signaling: lessons from single-molecule

microscopy. *Curr Opin Cell Biol* **57:** 57-63.

Case LB, Ditlev JA, Rosen MK (2019) Regulation of Transmembrane Signaling by Phase

Separation. *Annu Rev Biophys* **48:** 465-94.

Chavez-Abiega S, Goedhart J, Bruggeman FJ (2019) Physical biology of GPCR signalling

dynamics inferred from fluorescence spectroscopy and imaging. *Curr Opin Struct Biol*

55: 204-11.

Chen Y, Munteanu AC, Huang YF, Phillips J, Zhu Z, Mavros M, Tan W (2009) Mapping

receptor density on live cells by using fluorescence correlation spectroscopy. *Chemistry*

15: 5327-36.

Ditlev JA, Vega AR, Koster DV, Su X, Tani T, Lakoduk AM, Vale RD, Mayor S, Jaqaman K,

Rosen MK (2019) A composition-dependent molecular clutch between T cell signaling

condensates and actin. *Elife* **8**

Doh J, Irvine DJ (2006) Immunological synapse arrays: patterned protein surfaces that

modulate immunological synapse structure formation in T cells. *Proc Natl Acad Sci U S*

A **103:** 5700-5.

Edelstein A, Amodaj N, Hoover K, Vale R, Stuurman N (2010) Computer control of

microscopes using µManager. *Curr Protoc Mol Biol* **Chapter 14:** Unit14.20.

Ekstrand AJ, Cao R, Bjorndahl M, Nystrom S, Jonsson-Rylander AC, Hassani H, Hallberg B,

Nordlander M, Cao Y (2003) Deletion of neuropeptide Y (NPY) 2 receptor in mice

results in blockage of NPY-induced angiogenesis and delayed wound healing. *Proc Natl*

Acad Sci U S A **100:** 6033-8.

Gatterdam K, Joest EF, Gatterdam V, Tampe R (2018) The scaffold design of trivalent
chelator heads dictates affinity and stability for labeling His-tagged proteins in vitro and
in cells. *Angew Chem Int Ed Engl* **57:** 12395-9.

Hager R, Müller U, Ollinger N, Weghuber J, Lanzerstorfer P (2021) Subcellular Dynamic
Immunopatterning of Cytosolic Protein Complexes on Microstructured Polymer
Substrates. *ACS Sens*

Haqshenas G, Doerig C (2019) Targeting of host cell receptor tyrosine kinases by
intracellular pathogens. *Sci Signal* **12**

Harwardt MIE, Dietz MS, Heilemann M, Wohland T (2018) SPT and Imaging FCS Provide
Complementary Information on the Dynamics of Plasma Membrane Molecules. *Biophys*
J **114:** 2432-43.

Harwardt MIE, Young P, Bleymüller WM, Meyer T, Karathanasis C, Niemann HH, Heilemann
M, Dietz MS (2017) Membrane dynamics of resting and internalin B-bound MET
receptor tyrosine kinase studied by single-molecule tracking. *FEBS Open Bio* **7:** 1422-
40.

Hilger D, Masureel M, Kobilka BK (2018) Structure and dynamics of GPCR signaling
complexes. *Nat Struct Mol Biol* **25:** 4-12.

Huang L, Tan HY, Fogarty MJ, Andrews ZB, Veldhuis JD, Herzog H, Steyn FJ, Chen C
(2014) Actions of NPY, and its Y1 and Y2 receptors on pulsatile growth hormone
secretion during the fed and fasted state. *J Neurosci* **34:** 16309-19.

Kawai T, Akira S (2005) Pathogen recognition with Toll-like receptors. *Curr Opin Immunol*
17: 338-44.

Kupperman E, An S, Osborne N, Waldron S, Stainier DY (2000) A sphingosine-1-phosphate
receptor regulates cell migration during vertebrate heart development. *Nature* **406:** 192-
5.

Lafferty RA, Flatt PR, Irwin N (2021) Established and emerging roles peptide YY (PYY) and
exploitation in obesity-diabetes. *Curr Opin Endocrinol Diabetes Obes* **28:** 253-61.

Lanzerstorfer P, Borgmann D, Schütz G, Winkler SM, Höglinger O, Weghuber J (2014)
Quantification and kinetic analysis of Grb2-EGFR interaction on micro-patterned
surfaces for the characterization of EGFR-modulating substances. *PLoS One* **9:**
e92151.

Lanzerstorfer P, Müller U, Gordiyenko K, Weghuber J, Niemeyer CM (2020) Highly Modular
Protein Micropatterning Sheds Light on the Role of Clathrin-Mediated Endocytosis for
the Quantitative Analysis of Protein-Protein Interactions in Live Cells. *Biomolecules* **10:**
540.

Li MO, Rudensky AY (2016) T cell receptor signalling in the control of regulatory T cell
differentiation and function. *Nat Rev Immunol* **16:** 220-33.

Lindner D, Walther C, Tennemann A, Beck-Sickinger AG (2009) Functional role of the
extracellular N-terminal domain of neuropeptide Y subfamily receptors in membrane
integration and agonist-stimulated internalization. *Cell Signal* **21:** 61-8.
Luther SA, Cyster JG (2001) Chemokines as regulators of T cell differentiation. *Nat Immunol*
2: 102-7.
Lyon AS, Peeples WB, Rosen MK (2021) A framework for understanding the functions of
biomolecular condensates across scales. *Nat Rev Mol Cell Biol* **22:** 215-35.
Manz BN, Groves JT (2010) Spatial organization and signal transduction at intercellular
junctions. *Nat Rev Mol Cell Biol* **11:** 342-52.
Méndez-Couz M, Manahan-Vaughan D, Silva AP, González-Pardo H, Arias JL, Conejo NM
(2021) Metaplastic contribution of neuropeptide Y receptors to spatial memory
acquisition. *Behav Brain Res* **396:** 112864.
Mossman KD, Campi G, Groves JT, Dustin ML (2005) Altered TCR signaling from
geometrically repatterned immunological synapses. *Science* **310:** 1191-3.
Movafagh S, Hobson JP, Spiegel S, Kleinman HK, Zukowska Z (2006) Neuropeptide Y
induces migration, proliferation, and tube formation of endothelial cells bimodally via Y1,
Y2, and Y5 receptors. *Faseb j* **20:** 1924-6.
Nair PM, Salaita K, Petit RS, Groves JT (2011) Using patterned supported lipid membranes
to investigate the role of receptor organization in intercellular signaling. *Nat Protoc* **6:**
523-39.
Ojosnegros S, Cutrale F, Rodríguez D, Otterstrom JJ, Chiu CL, Hortigüela V, Tarantino C,
Seriola A, Mieruszynski S, Martínez E, Lakadamyali M, Raya A, Fraser SE (2017) Eph-
ephrin signaling modulated by polymerization and condensation of receptors. *Proc Natl*
Acad Sci U S A **114:** 13188-93.
Parker SL, Balasubramaniam A (2008) Neuropeptide Y Y2 receptor in health and disease.
Br J Pharmacol **153:** 420-31.
Pasquale EB (2010) Eph receptors and ephrins in cancer: bidirectional signalling and
beyond. *Nat Rev Cancer* **10:** 165-80.
Pike R, Ortiz-Zapater E, Lumicisi B, Santis G, Parsons M (2018) KIF22 coordinates CAR
and EGFR dynamics to promote cancer cell proliferation. *Sci Signal* **11**
Rosenbaum DM, Rasmussen SG, Kobilka BK (2009) The structure and function of G-
protein-coupled receptors. *Nature* **459:** 356-63.
Sánchez MF, Els-Heindl S, Beck-Sickinger AG, Wieneke R, Tampé R (2021) Photoinduced
receptor confinement drives ligand-independent GPCR signaling. *Science* **371**
Sánchez MF, Murad F, Gülcüler Balta GS, Martin-Villalba A, García-Sáez AJ, Carrer DC
(2018) Early activation of CD95 is limited and localized to the cytotoxic synapse. *Febs j*
285: 2813-27.

Sankaran J, Shi X, Ho LY, Stelzer EH, Wohland T (2010) ImFCS: a software for imaging
FCS data analysis and visualization. *Opt Express* **18:** 25468-81.

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S,
Rueden C, Saalfeld S, Schmid B, Tinevez JY, White DJ, Hartenstein V, Eliceiri K,
Tomancak P, Cardona A (2012) Fiji: an open-source platform for biological-image
analysis. *Nat Methods* **9:** 676-82.

Schneider CA, Rasband WS, Eliceiri KW (2012) NIH Image to ImageJ: 25 years of image
analysis. *Nat Methods* **9:** 671-5.

Schütz GJ, Weghuber J, Lanzerstorfer P, Sevcsik E (2017) Protein Micropatterning Assay:
Quantitative Analysis of Protein-Protein Interactions. *Methods Mol Biol* **1550:** 261-70.

Schwille P, Korlach J, Webb WW (1999) Fluorescence correlation spectroscopy with single-
molecule sensitivity on cell and model membranes. *Cytometry* **36:** 176-82.

Scott FL, Stec B, Pop C, Dobaczewska MK, Lee JJ, Monosov E, Robinson H, Salvesen GS,
Schwarzenbacher R, Riedl SJ (2009) The Fas-FADD death domain complex structure
unravels signalling by receptor clustering. *Nature* **457:** 1019-22.

Sebestyen Z, Prinz I, Déchanet-Merville J, Silva-Santos B, Kuball J (2020) Translating
gammadelta (γδ) T cells and their receptors into cancer cell therapies. *Nat Rev Drug*
Discov **19:** 169-84.

Stallaert W, Brüggemann Y, Sabet O, Baak L, Gattiglio M, Bastiaens PIH (2018) Contact
inhibitory Eph signaling suppresses EGF-promoted cell migration by decoupling EGFR
activity from vesicular recycling. *Sci Signal* **11**

Su X, Ditlev JA, Hui E, Xing W, Banjade S, Okrut J, King DS, Taunton J, Rosen MK, Vale
RD (2016) Phase separation of signaling molecules promotes T cell receptor signal
transduction. *Science* **352:** 595-9.

Sungkaworn T, Jobin ML, Burnecki K, Weron A, Lohse MJ, Calebiro D (2017) Single-
molecule imaging reveals receptor-G protein interactions at cell surface hot spots.
Nature **550:** 543-7.

Tang T, Hartig C, Chen Q, Zhao W, Kaiser A, Zhang X, Zhang H, Qu H, Yi C, Ma L, Han S,
Zhao Q, Beck-Sickinger AG, Wu B (2021) Structural basis for ligand recognition of the
neuropeptide Y Y(2) receptor. *Nat Commun* **12:** 737.

Torres AJ, Wu M, Holowka D, Baird B (2008) Nanobiotechnology and cell biology: micro-
and nanofabricated surfaces to investigate receptor-mediated signaling. *Annu Rev*
Biophys **37:** 265-88.

Traub MC, Longsine W, Truskett VN (2016) Advances in Nanoimprint Lithography. *Annu*
Rev Chem Biomol Eng **7:** 583-604.

Tsukiyama T, Zou J, Kim J, Ogamino S, Shino Y, Masuda T, Merenda A, Matsumoto M,
Fujioka Y, Hirose T, Terai S, Takahashi H, Ishitani T, Nakayama KI, Ohba Y, Koo BK,

Hatakeyama S (2020) A phospho-switch controls RNF43-mediated degradation of Wnt
receptors to suppress tumorigenesis. *Nat Commun* **11:** 4586.

Venkatakrishnan AJ, Deupi X, Lebon G, Tate CG, Schertler GF, Babu MM (2013) Molecular
signatures of G-protein-coupled receptors. *Nature* **494:** 185-94.

Walther C, Nagel S, Gimenez LE, Mörl K, Gurevich VV, Beck-Sickinger AG (2010) Ligand-
induced internalization and recycling of the human neuropeptide Y2 receptor is
regulated by its carboxyl-terminal tail. *J Biol Chem* **285:** 41578-90.

Wanka L, Babilon S, Kaiser A, Mörl K, Beck-Sickinger AG (2018) Different mode of arrestin-
3 binding at the human Y(1) and Y(2) receptor. *Cell Signal* **50:** 58-71.

Wawrezinieck L, Rigneault H, Marguet D, Lenne PF (2005) Fluorescence correlation
spectroscopy diffusion laws to probe the submicron cell membrane organization.
Biophys J **89:** 4029-42.

Wootten D, Christopoulos A, Marti-Solano M, Babu MM, Sexton PM (2018) Mechanisms of
signalling and biased agonism in G protein-coupled receptors. *Nat Rev Mol Cell Biol* **19:**
638-53.

Ziffert I, Kaiser A, Babilon S, Mörl K, Beck-Sickinger AG (2020) Unusually persistent Gα(i)-
signaling of the neuropeptide Y(2) receptor depletes cellular G(i/o) pools and leads to a
G(i)-refractory state. *Cell Commun Signal* **18:** 49.

Figures

Figure 1. High-content ligand-free receptor confinement.

A Rational of the high-content ligand-free receptor clustering. Matrices prestructured with

BSA are stepwise functionalized with biotin-BSA and SA. Upon addition of the multivalent

nanotool *tris*NTA$^{PEG12-B}$, His$_6$-tagged receptors in HeLa cells are captured to the

prestructured regions via the His-tag/*tris*NTA interaction.

B Chemical structure of the *tris*NTA$^{PEG12-B}$.

C Variable size protein patterns generated by further functionalization of SA-matrices with the

nanotool followed by incubation with His$_6$-GFP (0.1 µM, 20 min). Images were acquired by

confocal laser scanning microscopy.

D Intensity profile of the 1 µm pattern (white line in C) reflects high specificity of the

interaction.

E Large-scale cell patterning in living cells occurred after addition of the *tris*NTA$^{PEG12-B}$

(100 nM, 10 min).

F Customized Y$_2$R assembly on 3 µm and 1 µm SA-prestructured matrices.

G Intensity profile of the 1 μm pattern (white line in f) showed an intensity comparable to a

soluble His$_6$-tagged protein. Scale bars: 10 μm.

Figure 2. Ligand-independent response triggered by *in situ* receptor confinement.

A Time-lapse imaging of Y_2R assembly. Y_2R expressing HeLa cells were allowed to adhere to prestructured SA-matrices for 3 h and visualized by CLSM in LCIS at 37 °C. Time-lapse images were recorded for 10 min immediately after addition of *tris*NTA$^{PEG12-B}$ (100 nM).

B Reversal of the interaction and disassembly of the clusters is demonstrated upon addition of histidine. Y_2R expressing cells were allowed to adhere to the SA-matrices for 3 h, and then receptor confinement was induced by addition of *tris*NTA$^{PEG12-B}$ (100 nM). Subsequently, cells were incubated with histidine (5 mM) for 2 to 10 min followed by washing.

C Receptor integrated density in the patterned regions increased mono-exponentially, leading to an assembly rate of 0.20 ± 0.03 min^{-1} and $\tau_{1/2}$= 5 min.

D Fast cell spreading upon cluster formation was observed. Cell area analysis before and 10 min after nanotool addition showed a 20% area increase, confirming an effect of the receptor clusters on cell response. Values for cell area were normalized with respect to the highest value. The mean ± SD (20 cells) is shown. **p≤ 0.01 for Tukey test. Scale bars: 10 µm.

Figure 3. Ligand-free receptor confinement triggers calcium signaling.

A   Representative images of His$_6$-Y$_2$R^{mEGFP} expressing cells over SA-prestructured matrices

before and in the presence of *tris*NTA$^{PEG12-B}$ demonstrated a significant change in cell area.

Scale bar: 20 µm upper panel, 10 µm lower panel.

B   Representative fluorescence images of the Y$_2$R (upper panel) and color-coded images of

the Ca^{2+} dye (lower panel). Y$_2$R expressing cells on SA-prestructured matrices were

incubated with BioTracker 609 Red Ca^{2+} AM dye (3 µM) for 30 min. After rinsing, cells were

immediately imaged by CLSM in LCIS at 37 °C. Addition of the nanotool showed a twofold

increase in cytosolic calcium.

C   Analysis of the mean gray value for Ca^{2+} signal before (F_0) and upon addition of

*tris*NTA$^{PEG12-B}$ (F) *versus* time. Time-lapse images were recorded with 45 s interval before

and after addition of *tris*NTA$^{PEG12-B}$ (100 nM) (5 slices z-stack per time-point). ROIs covering

the complete cell area were considered. The mean ± SD (10 cells) is shown. Scale bar:

10 µm.

D   Quantification of receptor integrated density in the nanotool patterned regions showed an

increase in the density of the receptors within the clusters after addition of the pNPY. The

mean ± SD is shown (113 1 µm ROI were analyzed). ***$p \leq 0.001$ for Tukey test.

Figure 4. Selective arrestin-3 recruitment upon ligand-induced receptor activation.

A Schematic representation of the experimental set-up. Cells co-expressing Y_2R and Arr3 were allowed to adhere to SA-prestructured matrices for 3 h and visualized by total internal reflection fluorescence (TIRF) microscopy in LCIS at 37 °C.

B Representative TIRF images of cells before and upon addition of $tris$NTA$^{PEG12-B}$ (100 nM, 30 min) and subsequent incubation with porcine neuropeptide pNPY (10 nM) and Histidine (5 mM) in LCIS for 30 min at 37 °C. Scale bar: 5 µm.

C Fluorescence contrast analysis demonstrated no significant recruitment of Arr3 upon $tris$NTA$^{PEG12-B}$ (1.4-fold with respect to the basal signal before, 100 nM, 30 min). Addition of pNPY increased Arr3 signal (3.6-fold, 10 nM, 30 min) confirming co-patterning of the downstream signaling molecule. Subsequent addition of histidine led to a decrease in the signal (2.3-fold, 5 mM, 30 min). Data was normalized with respect to the fluorescence intensity before clustering and it is expressed as the means ± SEM (60 cells for each condition were analyzed). Tukey's multiple comparisons test was applied (***p≤ 0.001)

D Quantification of the fluorescence contrast in the patterned regions for Y_2R confirmed receptor enrichment upon addition of $tris$NTA$^{PEG12-B}$, (2-fold with respect to the basal signal

806   before, 100 nM, 30 min) which further increased 4-fold upon addition of pNPY (10 nM, 30

807   min). Histidine addition led to a decrease in the signal (1.7-fold decrease compared to

808   pNPY, 5 mM, 30 min). Data was normalized with respect to the fluorescence intensity

809   before clustering and it is expressed as the means ± SEM (60 cells for each condition were

810   analyzed). Tukey's multiple comparisons test was applied (***$p \leq 0.001$).

Figure 5. Decrease of receptor mobility in confined regions.

A Confocal images of FRAP measurements upon Y_2R clustering induced either by the nanotool *in situ* or by an anti-His$_6$ antibody. Insets represent the bleached ROIs. Fast recovery of the clusters can be detected for the case of the multivalent nanotool.

B Quantification of the receptor mobile fraction for cell patterning by the *tris*NTA$^{PEG12-B}$ and anti-His$_6$ antibody demonstrated a significant decrease in the receptor mobile fraction for the antibody matrices suggesting a lower receptor exchange. The mean ± SD is shown. 9 cells before, 11 cells after *tris*NTA$^{PEG12-B}$ (45x 1 µm ROIs) and 5 cells on anti-His$_6$ antibody matrices (13x 1 µm ROIs) were analyzed. ***p≤ 0.001 for Tukey test.

C imFCS correlates fluorescence intensity fluctuations in single camera pixels with a large degree of statistics, providing accurate diffusion coefficients with high spatial and temporal resolution.

D Widefield image of a ROI at the plasma membrane of a living cell upon addition of the nanotool analyzed by imFCS (left). The analyses of numerous pixels simultaneously provide two-dimensional diffusion data that draw a picture of the mobility of membrane receptors and reveal local differences in the diffusion (right).

E Both techniques demonstrated a decrease in the lateral diffusion of the receptor at the

plasma membrane after addition of the nanotool. Analysis of 1 µm clusters within the entire

ROI led to a further decrease in the lateral diffusion coefficient. For imFCS analyses, two-

sample t-tests ($\alpha = 0.05$) were applied to compare the diffusion coefficients for the different

conditions. The mean ± SD is shown. 36 and 24 cells for the conditions before and after

addition of $tris$NTA$^{PEG12-B}$ were analyzed. For FRAP, the mean ± SD is shown (9 cells

before, 11 cells after $tris$NTA$^{PEG12-B}$, 41 1 µm ROIs were analyzed). ***p≤ 0.001 for Tukey

test. Scale bar: 10 µm (A), 1 µm (D).

Expanded View Figures

Figure EV1. Receptor confinement with high reliability.

A Representative confocal image of a cell patterned by *tris*NTA$^{PEG12-B}$ (100 nM).

B Automatic cluster analysis performed by ImageJ requires a "binary", black and white,

image. A threshold range is set to select the objects of interest apart from the background.

All pixels in the image whose values lie under the threshold are converted to white and all

pixels with values above the threshold are converted to black. Further selection of the

clusters according to area and roundness enable a large-scale analysis.

C Change in cluster area and integrated density of the receptor within different 96-well plates,

different months and cell stocks reflected a reliable and reproducible approach. The

average area in the clustered regions (0.77 ± 0.03 μm^2) and integrated density of ten

images in five different experiments (841x ROIs of 1 μm in total) is shown. Scale bar:

10 μm.

Figure EV2. Y_2 receptors lacking a His$_6$-tag do not cluster in confined areas.

A Cells expressing Y_2 receptors without N-terminal His$_6$-tag over SA-prestructured matrices

before and after addition of the nanotool. Within the timeframe of imaging, there was neither

a pattern formation nor a change in the integrated receptor density.

B Quantification of the integrated Y_2R density before and after addition of $tris$NTA$^{PEG12-B}$. The

mean ± SD (18 cells) is shown. **$p \leq 0.01$ for Tukey test. Scale bar: 50 µm.

Figure EV3. Receptor density correlates with the area of the prestructured regions.

A,B BSA-prestructured matrices (1 μm, A or 3 μm, B) were stepwise functionalized with biotin-BSA and SA. Y_2R expressing HeLa cells were allowed to adhere to the functionalized matrix for 3 h and immediately imaged by CLSM in live-cell imaging solution (LCIS) at 37 °C. Addition of *tris*NTA$^{PEG12-B}$ (100 nM) led to *in situ* receptor assembly.

C 1 μm BSA-prestructured matrices were stepwise functionalized with biotin-BSA, SA, and a biotinylated anti-His$_6$ antibody (αHis$_6$ Ab). Y_2R expressing cells were allowed to adhere to the functionalized matrix for 3 h and immediately imaged by CLSM in LCIS at 37 °C.

D Quantification of the receptor integrated density. *In situ* receptor confinement by *tris*NTA$^{PEG12-B}$ resulted in a receptor density, which is comparable to cells in contact with prestructured antibodies. For the 3 μm patterns, receptor density correlated with pattern area. The mean ± SD (38 to 132x ROIs of 1 μm) is shown. ***p≤ 0.001 for Tukey test. Scale bars: 10 μm.

Figure EV4. Changes in cell motility are exclusively triggered upon receptor clustering.

A,B Confocal images of cells expressing His$_6$-tagged Y$_2$R on matrices which do not contain

SA but only biotin-BSA. Addition of the *tris*NTA$^{PEG12-B}$ nanotool confirms no effect on cell

spreading and motility as shown in the quantification of the cell area (B). The mean ± SD

(31 cells) is shown. **p≤ 0.01 for Tukey test.

C Confocal images of cells expressing Y$_2$ receptors lacking the His$_6$-tag on SA-matrices do

not present significant changes in cells spreading upon addition of the nanotool.

D Quantification of the cell area before and after addition of *tris*NTA$^{PEG12-B}$ (100 nM). Values

for cell area were normalized with respect to the highest value. The mean ± SD (20 cells)

is shown. **p≤ 0.01 for Tukey test. Scale bar: 50 µm.

Figure EV5. Calcium signaling is a specific response upon clustering.

A   Representative fluorescence images of the Y_2R (upper panel) and color-coded images of the Ca^{2+} dye (lower panel). Y_2R expressing cells over prestructured matrices in absence of streptavidin were incubated with BioTracker 609 Red Ca^{2+} AM dye (3 µM) for 30 min. After rinsing, cells were immediately imaged by CLSM in LCIS at 37 °C. Addition of *tris*NTA[PEG12-B] showed neither clustering nor change in cytosolic calcium. Scale bar: 10 µm

B   Analysis of the mean gray value for Ca^{2+} signal before (F_0) and upon addition of *tris*NTA[PEG12-B] (F) *versus* time. Time-lapse images were recorded with 45 s interval before and after addition of *tris*NTA[PEG12-B] (100 nM) (5 slices z-stack per time-point). ROIs covering the complete cell area were considered. The mean ± SD (10 cells) is shown.

Figure EV6. Arrestin-3 recruitment on antibody-confined regions.

A Schematic representation of the experimental set-up. Cells co-expressing Y_2R and Arr3

were allowed to adhere to anti-His_6 antibody prestructured matrices for 3 h and visualized

by total internal reflection fluorescence (TIRF) microscopy in LCIS at 37 °C.

B Representative TIRF images of cells before and upon addition of pNPY (10 nM) in LCIS for

30 min at 37 °C. Scale bar: 10 μm.

C Quantification of the fluorescence contrast in the Y_2R-patterned regions showed no

significant change in receptor intensity yet a recruitment of Arr3 upon addition of pNPY

(twofold). Data are expressed as the means ± SEM (30 cells for each condition were

analyzed). Tukey's multiple comparisons test was applied (**p≤ 0.01).

Figure EV7. Lipid localization and dynamics after receptor confinement.

A Live-cell plasma membrane staining was performed 15 min after Y_2R assembly in living cells. 5 µg/ml CellMask staining solution was incubated for 5 min at 37 °C and washed with LCIS before visualization. Lipid distribution is not affected by receptor confinement as shown by the homogeneous staining of the membrane.

B Representative confocal images of FRAP measurements for the lipid dye. An image of the receptor channel confirmed the presence of the pattern.

C FRAP recovery for the lipid dye measured in the entire rectangular ROI or at the Y_2R cluster spots (region selected based on the receptor channel image) demonstrated a rapid recovery for the lipids.

D The analysis did not show any differences in lipid diffusion coefficients for the entire rectangular ROI or at the 1 µm clustered regions ($D_{\text{entire ROI}}$ = 0.66 ± 0.10 µm^2/s and D_{spots} = 0.67 ± 0.17 µm^2/s). The mean ± SD (6 cells, 15x 1 µm ROIs) is shown. **p≤ 0.01 for Tukey test.

E E Quantification of the mobile fraction (M_f) for FRAP measurements of the lipid dye reflected no significant difference. The mean ± SD (5 cells, 14x 1 µm ROIs) is shown. **p≤ 0.01 for Tukey test. Scale bars: 10 µm.

Figure EV8. Dynamic receptor exchange in confined clusters.

A Representative confocal images of FRAP measurements for Y_2R expressing cells on SA-

prestructured matrices 10 min after addition of the nanotool.

B FRAP recovery curves reflecting the entire bleached area or an analysis performed only in

the clustered 1 µm regions. The analysis is based on a simulation approach which fits a

computer-simulated recovery to actual recovery data of a FRAP series and determines the

diffusion coefficient regardless of bleaching geometry.

C Quantification of the receptor density in the confined regions showed 50% recovery

indicating a high exchange rate. The mean ± SD (6 cells, 15x 1 µm ROIs analyzed) is

shown. **p≤ 0.01 for Tukey test. Scale bar: 10 µm.

Figure EV9. Receptor mobility on antibody structured matrices.

A   Scheme representing the experimental set-up.

B   Representative widefield image (left) of a ROI at the plasma membrane of a living cell grown on prestructured matrices with an anti-His$_6$ antibody (αHis$_6$ Ab) analyzed by imFCS and the derived two-dimensional diffusion map (right).

C   Lateral diffusion of the receptor analyzed by FRAP and imFCS. Both techniques demonstrated a decrease in D at the plasma membrane (D_{before}= 0.32 ± 0.06 μm^2/s and 0.25 ± 0.08 μm^2/s; $D_{\text{anti-His6 Ab}}$ = 0.18 ± 0.06 and 0.19 ± 0.06 μm^2/s for imFCS and FRAP respectively), concurring with the values obtained for the measurements upon addition of the nanotool. Analysis of clustered regions (1 μm) within the selected ROIs led to a further decrease in the diffusion coefficient (D_{spots}= 0.15 ± 0.05 μm^2/s and 0.13 ± 0.03 μm^2/s for imFCS and FRAP, respectively). For imFCS measurements, two-sample t-tests (α = 0.05) were applied to compare the diffusion coefficients for the different conditions (***p≤ 0.001). The mean ± SD is shown. 36 and 26 cells for the conditions before and after addition of anti-His$_6$ antibody were analyzed. For FRAP, the mean ± SD is shown. Here, 9 cells before, 5 cells after addition of anti-His$_6$ antibody, 20x ROIs were examined. **p≤ 0.01 for Tukey test. Scale bar: 1 μm.

952 **Figure EV10. Multivalent nanotool *tris*NTA$^{PEG12-B}$ analyzed by LC-MS.**

953 A *tris*NTA$^{PEG12-B}$ chromatogram reflecting the purity of the synthesized nanotool.

954 B LC-MS of *tris*NTA$^{PEG12-B}$, yielding the experimental mass ($M_{exp.}$) of 1874.85 Da ($M_{theor.}$ =

955   1873.90 Da).