A mixture of innate cryoprotectants is key for cryopreservation of a drosophilid fly larva.

Lukáš Kučera[2£], Martin Moos[1£], Tomáš Štětina[1], Jaroslava Korbelová[1], Petr Vodrážka[1], Lauren Des Marteaux[1,#], Robert Grgac[1,3], Petr Hůla[1,3], Jan Rozsypal[1], Miloš Faltus[4], Petr Šimek[1], Radislav Sedlacek[2], Vladimír Koštál[1]*

[1] Institute of Entomology, Biology Centre, Czech Academy of Sciences, České Budějovice, Czech Republic
[2] Czech Centre of Phenogenomics, Institute of Molecular Genetics, Czech Academy of Sciences, Vestec, Czech Republic
[3] Faculty of Science, University of South Bohemia, České Budějovice, Czech Republic
[4] Crop Research Institute, Praha, Czech Republic
[#] Current Address: Agriculture and Agri-Food Canada, Harrow Research and Development Centre, Harrow, Ontario, Canada

*Corresponding author: Vladimír Koštál
                           e-mail: kostal@entu.cas.cz
                           phone: +420 387775324
                           full address:
                                    Institute of Entomology, Biology Centre, Czech
                                    Academy of Sciences, Branišovská 31,
                                    370 05 České Budějovice, Czech Republic

[£] These authors contributed equally to this work.

Abstract

Background: Organisms evolved biochemical strategies to cope with environmental stressors. For instance, insects that naturally tolerate internal freezing produce complex mixtures of multiple cryoprotectants (CPs). Better knowledge on composition of these mixtures, and on mechanisms of how the individual CPs interact, could inspire development of laboratory CP formulations optimized for cryopreservation of cells and other biological material.

Results:
Here we identify and quantify (using high resolution mass spectrometry) a range of putative CPs in larval tissues of a subarctic fly, *Chymomyza costata*, that survives long-term cryopreservation in liquid nitrogen. The CPs (proline, trehalose, glutamine, asparagine, glycine betaine, glycerophosphoethanolamine, glycerophosphocholine, and sarcosine) accumulate in hemolymph in a ratio of 313:108:55:26:6:4:3:0.5 mmol.L^{-1}. Using calorimetry, we show that the artificial mixtures, mimicking the concentrations of major CPs' in hemolymph of freeze-tolerant larvae, suppress the melting point of water and significantly reduce the ice fraction. We demonstrate in a bioassay that mixtures of CPs administered through the diet act synergistically rather than additively to enable cryopreservation of otherwise freeze-sensitive larvae. Using MALDI-MSI, we show that during slow extracellular freezing of whole larvae trehalose becomes concentrated in partially dehydrated hemolymph and stimulates transition to the amorphous glass phase. In contrast, proline moves to the boundary between extracellular ice and dehydrated hemolymph and tissues where it likely forms a layer of dense viscoelastic liquid.

Conclusion: Our results suggest that different components of innate cryoprotective mixtures of freeze-tolerant insect act in synergy during extracellular freezing. We propose that transitions to amorphous glass (stimulated by trehalose) and viscoelastic liquids (having proline as major component) may protect macromolecules and cells from thermomechanical shocks associated with freezing and transfer into and out of liquid nitrogen.

Keywords: Cryoprotectants, Cryopreservation, Freezing, Proline, Trehalose, NADES, Freeze tolerance, Insects, Fly

Background

Various organisms from bacteria to animals respond to environmental stressors such as heat, cold, freezing, drought, hypersalinity, or high hydrostatic pressure by accumulating a stereotypic set of cytoprotective compounds including amino acids, sugars, polyols, and methylamines [1, 2]. This is also true for insects that naturally tolerate extracellular freezing during overwintering in temperate and polar habitats [3, 4]. In such insects, the accumulation of low molecular weight carbohydrates and free amino acids (cryoprotectants; CPs) is considered a fundamental tenet of their freeze tolerance [4-7]. Extensive knowledge has been accumulated over many decades about *which* CPs are present in many different species [7-10], but our understanding of *how* they protect the insect tissues and cells is far from complete [6].

Here we focus on an important but previously neglected aspect of natural freeze tolerance: the components of complex *mixtures* of multiple CPs may contribute to freeze tolerance via additive or even synergistic mechanisms [4, 11, 12]. It is well known from practice in clinical medicine and biotechnology that combinations of CPs protect the viability of cryopreserved cells or materials better than single substances [13]. The mixtures may even have emergent properties, i.e. those beyond the summation of properties of the individual components. Relatively recently, the existence of specific mixtures, so-called NADES (Natural Deep Eutectic Systems) in various organisms was proposed [14, 15]. These mixtures consist of stereotypic sets of natural primary metabolites such as sugars, sugar alcohols, organic acids, amino acids, and amines, and are characterized by a much lower melting point than that of the individual components, mainly due to the formation of intermolecular hydrogen bonds [16]. These mixtures have been proposed to serve different physiological functions, including survival of organisms in extreme drought and cold [17-19] and have also been proposed as new agents for cell cryopreservation [20]. We believe that there may be major practical consequences of exploring the complex composition of insect CP mixtures, as they could aid in the development of applicable cryogenic techniques. Freeze-tolerant insects also offer the potential for the discovery of novel components, for the study of interactions between individual components, and for the analysis of the principles underlying survival after freezing in a complex organism composed of different cell types and tissues; a complexity that still poses a challenge for medical cryopreservation [21, 22].

It is reasonable to first analyze the suite of CPs accumulated by the most freeze-tolerant model species, and the larvae of the malt fly, *Chymomyza costata* (Diptera: Drosophilidae) are among the most cold-hardy animals known [23]. Overwintering (diapausing), cold-acclimated larvae can survive freezing of all osmotically active water (68% of total body water), have no apparent lower thermal limit while frozen, and even survive long-term cryopreservation in liquid nitrogen (LN$_2$) [24-26]. We have previously shown that high concentrations of the free amino acid proline acquired by diapausing *C. costata* larvae during cold acclimation are essential for survival of freezing and cryopreservation stress [26, 27]. Here we extend our previous results to include whole complex of CPs accumulated by these freeze-tolerant larvae. Using a combination of four chromatographic mass spectrometric (MS) analytical platforms, we established metabolite profiles of four different *C. costata* tissues in two contrasting larval phenotypes that differ greatly in freeze tolerance and cryopreservability in LN$_2$. By combining the results of the analytical chromatographic MS analysis with matrix-assisted laser desorption/ionization (MALDI) mass spectrometry imaging (MSI), we show that freeze-tolerant larvae accumulate putative CPs in all tissues, but especially in hemolymph. Using the same technique, we observed changes in CP localization during slow extracellular freezing associated with partial dehydration of hemolymph and tissues. We then used differential scanning calorimetry (DSC) to describe the thermal phase transition properties of artificial aqueous CP mixtures mimicking the concentrations of the five most abundant metabolites in the hemolymph. Finally, we show in a bioassay that mixtures of CPs administered through the diet work in a synergistic rather than an additive way to induce strong freeze tolerance and cryopreservability in otherwise freeze-sensitive larvae.

Results

Characterization of the CP mixture from *C. costata* larvae

We generated two acclimation/phenotypic variants of 3rd instar *C. costata* larvae: (i) freeze-sensitive larvae (FS), which develop directly to the pupal stage; (ii) freeze-tolerant larvae (FT), whose development is arrested at diapause and which undergo gradual cold acclimation (see Materials and Methods for more details).

We performed a targeted quantitative analysis of 49 metabolites in the hemolymph and tissues from two larval phenotypes using a combination of four different MS-based analytical platforms (Fig. 1a, Additional file 1: Fig. S1). For most metabolites, the pool in

the hemolymph was significantly larger than the pools in other tissues (Additional file: Fig. S2). The volume of hemolymph in one larva is approximately 200 nL (see Additional file 2: Table 1) representing only 10% of the total body volume or body mass (about 2 mg). Nevertheless, the hemolymph contained approximately 50% of the total metabolite pool in both FS and FT larvae (Additional file 2: Table S1). Entry into diapause and cold acclimation (associated with the acquisition of extreme freeze tolerance) induced accumulation of specific metabolites (putative CPs) in FT larvae (Additional file 1: Fig. S1). The sum of the molar concentrations of the 49 metabolites in FS hemolymph was 182 mmol.L^{-1}, but this had increased to 554 mmol.L^{-1} in FT hemolymph (Additional file 2: Table S1). Four compounds that contributed most to this increase were: proline (increase of 277 mmol.L^{-1}), trehalose (increase of 59 mmol.L^{-}), glutamine (increase of 22 mmol.L^{-1}), and asparagine (increase of 17 mmol.L^{-1}). The same four compounds also ranked first in order of concentration in FT hemolymph: proline (313 mmol.L^{-1}), trehalose (108 mmol.L^{-1}), glutamine (55 mmol.L^{-1}), and asparagine (26 mmol.L^{-1}). Glycine-betaine (hereafter 'betaine', 6.1 mmol.L^{-1}) was fifth in concentration, although it increased only moderately (by 1.7 mmol.L^{-1}) from FS to FT. In addition, three other compounds had statistically higher hemolymph concentrations in FT compared to FS larvae, although their concentrations in FT hemolymph were relatively low in absolute terms: glycerophosphoethanolamine (GPE, 4.0 mmol.L^{-1}), glycerophosphocholine (GPC, 2.9 mmol.L^{-1}), and sarcosine (0.5 mmol.L^{-1}). For the purposes of this article, we consider the above compounds as putative CPs.

Localization of CP mixture components in larval tissues

Using MALDI-MSI we localized select metabolites in larval sections. A longitudinal section through the FT larva provides the best overview of tissue structure (Fig. 1b). The highest intensity of the MALDI-TOF (TOF, Time Of Flight) signal for ATP is localized predominantly in muscle tissue (corresponding to its role as the most important intracellular phosphagen) (Fig. 1c). In contrast, the highest intensities of proline and trehalose signals are localized in the hemolymph (Fig. 1d, e) where they are also colocalized with the high-intensity signals of glutamine, asparagine, and GPE (Additional file 1: Fig. S3).

Figure 1: Metabolite profiles and localization in hemolymph and tissues of *C. costata* larvae.
(a) Forty-nine target metabolites were absolutely quantified in the hemolymph of freeze sensitive (FS) and freeze tolerant (FT) larvae using four MS-based analytical platforms. Each column represents a metabolite concentration (mean ± SD; *n* = 4 biological replicates, each containing hemolymph from 30 larvae). Differences between the means of FS and FT were assessed by unpaired t-tests corrected for multiple comparisons using the Holm-Sidak method with α = 0.01. Asterisks indicate significantly different means.
(b) Longitudinal section of a FT larva stained with hematoxylin and eosin, with tissues highlighted by different colors. (c-e) Intensities of the MALDI-TOF signals corresponding to the three analyzed compounds: **(c)** ATP; **(d)** proline; **(e)** trehalose. **(f)** While proline (green signal) is most evident in hemolymph, ATP (red signal) is mainly found in muscles. **(g)** Proline (green) and trehalose (red) co-localize in hemolymph (a yellow color results from the mixing of green and red). See Additional file 1: Fig. S3 for MALDI-MSI images of other compounds.

As it was difficult to obtain a sufficient number of high quality longitudinal sections, we used transverse sections through the middle region of the larvae for the statistical analysis of the relative intensities of the MALDI-TOF signals. Quantitative analysis of these signals is technically difficult for small metabolite molecules freely dissolved in biological aqueous solutions (Additional file 1: Fig. S4). Nevertheless, our analysis suggests that there are no differences in the overall localization patterns of the five putative CPs (proline, trehalose, glutamine, asparagine, and GPE) between FS and FT larvae. However, the signal intensities for all five CPs were much stronger in FT than in FS larvae (compare the *y*-axes of Additional file 1: Figs. S5 vs S6). The MALDI-MSI

thus confirms the results of the quantitative MS analysis (Fig. 1a) that the putative CPs are accumulated in all tissues, but especially in the hemolymph, of FT larvae (for details, see Additional file 3 containing all MALDI-MSI images).

Figure 2: Localization of proline and trehalose *in C. costata* FT larvae prior to and after slow freezing.
(a, b) Examples of transverse sections through the middle region of FT-larva (a) and FT-frozen-larva (b). The tissues are highlighted by different colors. Note the large masses of extracellular ice between the partially dehydrated tissues of FT-frozen-larva. **(c, d)** Relative proportions of total area occupied by different tissues or ice. Each column is a mean ± S.D. of five or six sections (FT or FT-frozen larvae, respectively). Means flanked by different letters are significantly different (one way ANOVA followed by Bonferroni's multiple comparison test). **(e, f)** Results of MALDI-TOF signal intensity quantification (arbitrary units, A.U. per pixel of 10x10 um) for three select metabolites (ATP, proline, trehalose). Each point is a mean ± S.D. of five or six sections (FT or FT-frozen larvae, respectively). Mean values flanked by different letters are significantly different (one way ANOVA followed by Bonferroni's multiple comparison test). **(g, h, i)** Examples of MALDI-TOF signal intensities for proline (green) and trehalose (red). Note that proline and trehalose signals markedly overlap (yellow color) in the hemolymph of FT larva (g), while they apparently de-localize in FT-frozen-larva (h, i). Trehalose concentrates in the partially dehydrated hemolymph, while proline migrates to the border between extracellular ice and partially dehydrated tissues of the FT-frozen larva.

During slow inoculative freezing to -30°C (rate of 0.1°C.min^{-1}; see Materials and Methods for the freezing protocol) large masses of extracellular ice crystals develop in between partially dehydrated pools of larval hemolymph and tissues (Fig. 2a-d). We observed a striking difference in the 'behavior' of two major components of the CP mixture during slow

extracellular freezing: while trehalose was concentrated in the pool of dehydrated hemolymph of FT larvae, a large quantity of proline apparently 'migrated' from the hemolymph to the ice (Fig. 2e, f) and formed thin 'layers' of concentrated proline at the boundary between the extracellular ice and the dehydrated hemolymph and tissues (Fig. 2g-i). In these layers, proline is colocalized with high-intensity signals from glutamine and asparagine, while trehalose is colocalized with GPE in the dehydrated hemolymph (Additional file 1: Figure S7; for details, see Additional file 3 containing all MALDI-MSI images).

Thermal phase transitions in artificial CP mixtures

To assess the influence of different CP mixture components on water binding and thermal phase transitions, we compared heating curves from differential scanning calorimetry (DSC) scans of aqueous solutions containing single components or different CP mixtures with real FT hemolymph. We focused on melting of bulk water (Fig. 3a) and glass transition (Fig. 3b). The full DSC analysis results can be found in Additional file 2: Table S2.

Figure 3: Thermal phase transitions in the aqueous solutions of CP mixtures and individual components.
(a) Example of a heating curves recorded by differential scanning calorimetry (DSC) of the artificial CP mix 1588 (for explanation, see text). Three technical replicates are shown in different colors. Results calculated by TA Universal Analysis 2000 software are shown for the 1st replicate only (green line). The analysis of melting endotherm shows that melting of bulk water starts at an m.p. of -9.28°C, while only 58.7% of the total water in the sample is osmotically active (OAW), i.e. melts (the enthalpy of melting transition of ΔH of 334 J.g^{-1} would

correspond to melting of 100% total water) while 41.3% of total water is osmotically inactive (OIW) or 'bound'. **(b)** A glass transition was observed at -48.5°C (T_g, inflection point) associated with change in specific thermal capacity ΔCp of 0.3289 J.(g.°C)$^{-1}$. **(c)** Linear associations of the melting point (m.p., red line) or the fraction of bound water (%OIW, blue line) with the osmolality of the solution. Each point represents a specific solution and all solutions are shown in detail in Additional file 2: Table S2. Points corresponding to water, FT hemolymph, and CP mixes 508 and 1588 are identified for clarity. The broken lines show 95% confidence intervals of the linear regressions. **(d, e)** Temperatures of the glass transition, Tg (d) and changes in thermal capacity, ΔCp (e) of the different solutions. Empty points in (d) or a missing column in (e) indicates that no glass transition was observed, while the violet points and orange columns show mean ± S.D. values of T_g or ΔCp, respectively, for solutions where a glass transition was observed (see Additional file 2: Table S2 for the full dataset).

We designed the CP mixtures according to the concentrations of the five most abundant (and commercially available / affordable) compounds in the FT larval hemolymph. CP mix 508 mimics the composition of the major CPs in the hemolymph of FT larvae prior to freezing, while CP mix 1588 takes into account a 3.125-fold increase in the concentration of all components resulting from extracellular freezing of osmotically active water (68%). In addition, we tested all components individually at concentrations equivalent to those estimated in FT hemolymph after extracellular freezing, and we also designed two reduced-composition mixes: Pro-Tre 1316 (proline 978 + trehalose 338) and Pro-Gln-Asn 1231 (proline 978 + glutamine 172 + asparagine 81) (see Additional file 2: Table S2 for details). We found that depression of the bulk water melting point, as well as the fraction of 'bound' water (unfrozen, osmotically inactive water, %OIW), correlate linearly with the molal concentration of a solution, regardless of whether it is a single component solution or a mixture. Even the values for real hemolymph collected from FT larvae fit well to the linear relationships (Fig. 3c).

Whenever trehalose was present in a mixture, a clear glass transition was observed (Additional file 1: Fig. S8). The temperatures of glass transition (T_g) in different trehalose-containing solutions varied from -29.6°C (trehalose 338) to -49.7°C (Pro-Tre 1316) (Fig. 3d). The change of thermal capacity (ΔCp) during de-glassing was slightly above 0.3 J.(g.°C)$^{-1}$ in the solutions containing a relatively high concentration of trehalose (338 mmol.kg^{-1}) but lower, ~ 0.1 J.(g.°C)$^{-1}$, in the solution containing a relatively low concentration of trehalose (CP mix 508 contains 108 mmol.kg^{-1} of trehalose) and in real FT hemolymph (which also contains 108 mmol.kg^{-1} of trehalose) (Fig. 3e). No glass transition was observed in the concentrated proline 978 solution (Additional file 1: Fig. S9). The presence of proline in a mixture with glutamine and asparagine (Pro-Gln-Asn 1231) resulted in the elimination of small glass transitions observed in single-component mixtures of glutamine or asparagine (Fig. 3d, e). We also observed minor exotherms in heating curves of glutamine and asparagine

at -22.8°C and -14.6°C, respectively (Additional file 1: Fig. S10). The presence of such exotherms was also eliminated by the addition of proline in the Pro-Gln-Asn 1231 mix (Additional file 2: Table S2).

CPs in mixture protect larvae from cryopreservation stress better than individual CPs

To evaluate the ability of a complex cryoprotective mixture to increase freeze tolerance compared to individual components of the CP mix, we fed FS larvae with CP-augmented diets and then performed freezing survival assays. We prepared the CP mix 508 diet by augmenting the standard larval diet with a CP mix 508 solution (see Additional file 2: Table S3). When FS larvae were fed with the CP mix 508 diet from hatching, more than 90% died (often as 1^{st} instars). We therefore exposed only 3^{rd} instar FS larvae to this diet and for only three days (at ages 17-19 days), which reduced larval mortality to 20%. The surviving larvae were significantly smaller at day 19 and pupariation was delayed by two days (Additional file 2: Table S4), indicating that even short-term exposure to the CP mix 508 appears to have some toxic effects.

A three-day exposure to the CP mix 508 diet significantly increased whole-body concentrations of most CP mix components: proline by 6.8-fold, trehalose by 1.7-fold, glutamine by 5.4-fold, and asparagine by 19.9-fold (see Additional file 2: Table S5 for more details). In the hemolymph, the concentrations of proline, glutamine, and asparagine reached high levels of 289, 80, and 49 $mmol.L^{-1}$, respectively, while trehalose and betaine remained relatively low at 29 and 7 $mmol.L^{-1}$, respectively (Fig. 4a, see Additional file 2: Table S5 for more details). The sum molarity of the five CPs in hemolymph reached 454 $mmol.L^{-1}$ in FS larvae fed with the CP mix 508 diet, which is broadly similar to the sum of 508 $mmol.L^{-1}$ seen in FT larvae.

No FS larvae that fed exclusively on the standard diet survived cryopreservation in LN_2. In striking contrast, 58% of the FS larvae survived cryopreservation after only three days of feeding with the CP mix 508 diet; 24.5% went on to pupate and 18.3% successfully metamorphosed to the adult stage (Fig. 4b).

Figure 4: Effects of CP-augmented diets on *C. costata* larvae.
(a) Change in the hemolymph concentration of five CPs in response to feeding 17 day-old FS larvae with a CP mix 508 diet for three days. Concentrations are compared with FS and FT larvae fed on a standard diet, using one way ANOVAs followed by Bonferroni's multiple comparison tests (columns flanked by different letters differ significantly). **(b)** Percent survival of larvae cryopreserved in LN_2 (note: only diets where at least one larva survived are shown in panel b). The 17 day-old FS larvae were fed different diets for 3 days, then exposed to LN_2 for 1 h, returned to the standard diet at a constant 18°C, and their survival (movements 24 h later), pupation, and adult emergence were observed over the next 14 days. The two numbers in parentheses (*n*, *N*) show: *n* = the number of fly generations, and *N* = the total number of larvae in all replicates. The mean survival for larvae fed a CP mix 508 diet versus a proline diet (Pro 313) were directly compared using a two-tailed Student's *t*-test.

The components of the CP mix 508 administered via the diet had much less (or no) effect on larval freeze tolerance when they acted in isolation: betaine and asparagine had absolutely no effect; glutamine and trehalose resulted in 1 and 2 larvae surviving (out of 100), respectively, but these larvae did not pupate; proline augmentation was most effective, resulting in 37.7%, 10.3% and 7.2% survival for larvae, pupae, and adults, respectively. When proline was excluded from the CP mix, the remaining components allowed only 4 larvae (out of 100) to survive LN_2 exposure, demonstrating that proline is the most important component. Nevertheless, the complete CP mix 508 was approximately 2-fold more effective at bolstering LN_2 survival than proline alone. This result was consistent at all three levels of survival (see *t*-tests in Fig. 4b) and also across five different generations of assayed insects (Additional file 2: Table S6). Simple addition of the individual effects of the three components (proline + trehalose + glutamine) was not sufficient to explain the effect of the complete CP mix 508. These results suggest a synergy between CP mix 508 components that should be investigated in the future.

Discussion

Different components of the fly's innate CP mixture behave differently during slow extracellular freezing

We identified the composition of the innate putative CP mixture which bathes the tissues of extremely freeze-tolerant *C. costata* larvae: proline, trehalose, glutamine, asparagine, betaine, GPE, GPC, and sarcosine, which occur in hemolymph in a stoichiometric ratio of 313:108:55:26:6:4:3:0.5 mmol.L^{-1}, respectively. It may not be surprising that the mixture is broadly similar to the cocktails known from other organisms exposed to various environmental stressors [1, 2, 14, 18]. The occurrence of the same classes of organic cytoprotectants and compatible osmolytes in different organisms is likely based on convergent evolution which favors accumulation of polar molecules with high solubility in water (high accumulation capacity) countered by low tendency to interact with charged groups of macromolecules and membranes (low toxicity) [28].

We performed a bioassay showing that supplementation of the larval diet with major components of the CP mixture in their native stoichiometric ratio changes freeze sensitive-larvae to freeze-tolerant/cryopreservable-larvae within just three days. This result convincingly demonstrates that the accumulation of CPs has a strong functional (adaptive) significance in *C. costata*. However, this adaptation has probably evolved to cope with conditions typical for larvae overwintering in thermally buffered microhabitats under the bark of fallen trees, often under a blanket of snow [29, 30]. Therefore, there may be different functional explanations for the roles of CPs in ecologically relevant situation in the field and for deep freezing and cryopreservation in LN$_2$ in the laboratory. Here we avoid the eco-physiological aspects of CP-based freeze tolerance in animals and insects in the field, as these have been discussed for animals [4, 6, 31, 32] and also specifically for *C. costata* [26] elsewhere. Instead, we limit further discussion to the potential relevance of our results for deep freezing and cryopreservation in the laboratory.

We applied the MALDI-MSI technique to localize CPs in larval tissues and, for the first time in cryobiology literature, to determine how this localization changes during slow extracellular freezing, i.e. under conditions that allow larvae to survive deep freezing and subsequent cryopreservation in LN$_2$. So far, concentrations of CPs in insects have mostly been reported in whole body samples for two general reasons: First, insect tissues are

small, making it technically difficult to estimate precise concentrations, and second, native CPs – being standard products of metabolism – have been considered to move freely across membranes [1]. In cases where CP analyses were performed on separate tissues [33-35], the CPs were generally found in all examined tissues. Here we have confirmed that FT larvae of *C. costata* accumulate putative CPs in all examined tissues but that the highest concentrations occur in the hemolymph.

One of the most salient findings of this study is the observation that the two major components of the CP mixture, proline and trehalose, 'behave' differently during slow extracellular freezing. The MALDI-MSI analysis suggests that the bulk of the trehalose molecules remain in their original location (before freezing) and concentrate in partially freeze-dehydrated hemolymph and tissues. In contrast, some of the proline molecules appear to move out of the hemolymph and concentrate in thin layers separating the extracellular ice crystals from freeze-dehydrated tissues and hemolymph. At this stage of research, we can only offer hypothetical/speculative explanations for the physico-chemical mechanisms responsible for de-localization of proline and trehalose and for its possible functional significance in a cryopreserved insect. We suggest that the mechanism behind the de-localization could be explained by differences between trehalose and proline in water solubility, propensity to induce glass transition, and mobility in highly viscous freeze-concentrated solutions.

Trehalose stimulates glass transition in partially freeze-dehydrated hemoplymph and tissues

We have previously shown that in gradually frozen FT larvae a glass transition of the residual solution occurs between -20°C and -30°C. Moreover, only larvae pre-frozen to temperatures below -30°C (i.e. below the temperature of glass transition) were able to survive abrupt submersion and cryopreservation in LN_2 and successfully develop into adults [26]. This result clearly demonstrated the importance of the glass transition for survival in LN_2. In our earlier work [26], we attributed glass formation to high concentrations of proline in accordance with results of the DSC analysis performed by [36]. However, this observation was challenged in later DSC studies [37, 38] and our own analysis confirms that proline at a concentration of 978 mmol.kg^{-1} does not induce glass transition, at least not at temperatures above -60°C. Instead, trehalose is known in physical chemistry as one of the strongest glass transition inducers in aqueous systems

[39-41]. Nicolajsen and Hvidt [42] reported that a water:trehalose mixture (%weight, 89.7:10.3) de-glasses at -30.5°C with a ΔCp of 0.38 J.(g.°C)$^{-1}$. These values agree well with our DSC analysis results showing de-glassing at -29.6°C with a ΔCp of 0.33 J.(g.°C)$^{-1}$ for the trehalose 338 solution (which is 88.7:11.3 %weight of water:trehalose). Accurate interpretation of the thermal behavior and glass transition parameters is a challenge for physical chemistry of simple binary mixtures such as trehalose:water [40-44]. This limitation becomes almost insurmountable for more complex mixtures and, especially, for biological solutions. Nevertheless, our DSC analysis suggests that trehalose rather than proline may be the component mainly responsible for the glass transition in larval body fluids: (i) all artificial CP mixtures containing trehalose showed clear glass transitions with broadly similar T_g and ΔCp parameters; (ii) the parameters were similar for real hemolymph collected from FT larvae (containing 108 mmol.L^{-1} trehalose) and for the artificial CP mix 508 (simulating composition of major CPs in FT hemolymph and also containing 108 mmol.kg^{-1} trehalose); (iii) no glass transition was observed in the concentrated proline 978 solution.

During glass transition, the viscosity of the supercooled liquid increases rapidly as the frequency, strength, and lifetime of hydrogen bonds in a system increases. In dilute systems, most hydrogen bonds are formed between water molecules, but as the concentration of solute molecules increases the interactions between water-trehalose and trehalose-trehalose become more frequent until trehalose molecules form large hydrogen-bonded clusters that exclude water [43, 44]. The increase in viscosity slows the molecular dynamics and macromolecules become trapped in an amorphous hydrogen-bonded matrix, which probably prevents undesirable transitions such as protein unfolding. In addition, trehalose can directly form hydrogen-bonds with macromolecules which probably further increases their stability in the amorphous glass [44, 45]. Trehalose molecules involved in glass formation are relative immobile. Moreover, large proportions of water molecules are 'bound', i.e. osmotically inactive in FT larval hemolymph (35.5%) or in whole-body *C. costata* (42% [26]), and therefore do not contribute much to trehalose solvation. As a result, trehalose can even locally reach the solubility limit (about 1.7 mol.kg^{-1} according to ALOGPS, http://www.vcclab.org/lab/alogps/), partly crystallize, and form a eutectic system with ice crystals. Both crystallization and glass formation would make trehalose molecules immobile, i.e. gradually concentrated in dehydrated tissues and hemolymph, which we observed.

Proline-rich viscoelastic liquid forms at the boundary between extracellular ice and dehydrated tissues

In contrast to trehalose, proline does not induce a glass transition [37, 38] and its solubility in water is enormous (it may reach over 15 mol.kg^{-1} [46, 47]). Nevertheless, physical chemistry shows that proline in high concentration also extensively hydrogen bonds with water and together they form a viscoelastic liquid [48-50]. The formation of dense, viscoelastic liquids is a guiding principle underlying the theory of NADES systems [14]. The components in a complex mixture forming NADES are connected to one another through a well-organized tridimensional system with optimum interactions via inter- and intramolecular hydrogen bonding [15, 16]. Through these interactions, NADES systems exhibit emergent properties: (i) the melting point is deeply suppressed (usually below -30°C); (ii) ice crystallization is completely (or at least strongly) inhibited; (iii) while the tendency to transition to amorphous glasses is enhanced [17, 19, 51]. However, work with NADES has also shown that the interactions between the NADES components are weakened with water dilution and even broken with water content exceeding 50 % weight, when each component recovers its own specific properties [16, 17, 52]. Our artificial CP mixtures have relatively high water content (between 79 and 99 %weight), as do the FT larval hemolymph (74.4 %weight) and whole larvae (69.3 %weight). Using DSC analysis, we show that these solutions exhibit significant suppression of the melting point (e.g. to -9.2°C in CP mix 1588, and to -4.5°C in FT hemolymph) and significant water-binding capacity (41.1 %OIW in CP mix 1588, and 35.5 %OIW in FT hemolymph). Both parameters (the decrease in melting point and the increase in OIW fraction) were linearly dependent on the osmolality of a solution (from 0 for water to 1,588 mmol.kg^{-1} for CP mix 1588). These results suggest that neither the CP mixes 508 and 1588 nor the FT hemolymph are 'deep eutectic systems' (DES, in the strict sense of [51]), but regular aqueous solutions with relatively low transition temperature from liquid to solid phase (LTTM, in the sense of [52]).

We suggest that a proline-dominated viscoelastic solution with NADES-like properties may form on a microscale level, locally, in slowly freeze-dehydrating larvae. As water molecules are osmotically driven from the gradually dehydrating hemolymph and tissues and migrate toward the extracellular ice during extracellular freezing, the highly soluble and relatively mobile proline molecules could move together with water until they reach a barrier of ice where they form a thin layer of concentrated proline, which we observed in

MALDI-MSI. Such a viscoelastic liquid could contribute to the stabilization of deeply frozen *C. costata* larvae by forming a rubber-like zone between extracellular ice crystals and freeze-dehydrated tissues, thus reducing the thermo-mechanical stresses associated with temperature fluctuations during immersion in LN_2 and rewarming [53].

Conclusions

Overall, our study reveals the composition of a specific mixture of putative cryoprotectants accumulated by the freeze-tolerant larvae of *C. costata*. We demonstrate in a bioassay that as little as three-days exposure to an artificial diet enriched with an artificial mixture of CPs allows an otherwise freeze-sensitive larva to survive not only deep freezing but also cryopreservation in LN_2. The components of the CP mixture act synergistically rather than additively to improve the freeze tolerance. These findings open avenues for functional experiments to better understand the mechanisms underlying cryoprotection and cryopreservation of complex multicellular organisms. Toward this goal, we demonstrate that the accumulated CPs suppress the melting point of body water, increase the fraction of osmotically inactive water, and significantly reduce the ice fraction. During slow extracellular freezing of whole larvae, trehalose becomes concentrated in the partially dehydrated hemolymph and tissues and stimulates glass transition, while proline moves to the boundary between extracellular ice and dehydrated tissues where it probably creates a layer of dense viscoelastic liquid. We suggest that these mechanisms contribute to larval extreme freeze tolerance and enable their cryopreservation by protecting their cells against thermomechanical shocks associated with freezing and transfers into and out of LN_2.

Materials and methods

Fly rearing and acclimations

A colony of *C. costata*, Sapporo strain [54], was reared on artificial diet in MIR 154 incubators (Sanyo Electric, Osaka, Japan) as described previously [55, 56]. Two phenotypic variants (FS and FT) at the 3rd larval instar were generated according to our earlier acclimation protocols [23, 26, 27] (Fig. 5a). The phenotypic variants FS and FT correspond to the variants LD and SDA, respectively, as described in our previous publications [23, 26, 27]. FS larvae have limited survival after freezing (35% survive to -5°C, 10% to -10°C, and none survive freezing to -20°C or below). In contrast, practically all FT larvae survive deep freezing to -30°C or even -70°C, and 42.5% survive six

months of cryopreservation in LN_2 [26]. In the present study, larvae were frozen according to a previously developed optimal freezing and cryopreservation protocol [26] schematically presented in Fig. 5b, and survival was assessed as the ability to pupate and metamorphose into the adult stage within 42 d after freezing (at a constant 18°C/LD).

Figure 5: Phenotypic/acclimation variants and freeze tolerance bioassay.
(a) Acclimation: *Chymomyza costata* flies were reared from eggs (time 0) to 3[rd] larval instars (time 3 weeks) at constant 18°C under one of two photoperiods: long-day (LD) or short day (SD). LD conditions promote direct development from larva to pupa. Mature LD-reared larvae are freeze sensitive (FS) and do not survive cryopreservation in LN_2. SD conditions induce entry into developmental arrest (diapause). Larvae reach their final weight after 6 weeks and are then transferred to constant darkness (DD) and low temperatures of 11°C followed by 4°C. Cold acclimation induces acquisition of freeze tolerance (FT) including the ability to survive in LN_2. **(b)** Optimal protocols for freezing and LN_2 cryopreservation according to [26]. The protocol consisted of five steps set in a Ministat 240 programmable cryostat (Huber, Offenburg, Germany). The addition of a small ice crystal at the beginning of step (ii.) induces inoculative internal freezing in larvae. The FT-frozen larvae underwent slow pre-freezing to -30°C associated with freeze dehydration of their tissues.

To collect the hemolymph, 30 larvae were gently pierced and torn on a piece of Parafilm, creating a large droplet of pooled hemolymph. This droplet was then extracted using a calibrated glass capillary (Drummond Sci., Broomall, PA, USA). To obtain fat body, muscle, and midgut tissues (including proventriculus and gastric caecae), the larvae were quickly dissected under binocular microscope in phosphate-buffered saline (PBS, 8.0 g NaCl; 0.2 g KCl; 1.8 g $Na_2HPO_4.2H_2O$; 0.24 g, KH_2PO_4 dissolved in 1 L of distilled water, pH 7.2). Dissection of a single larva took less than 2 min, and we took care to sample a similar part of the respective tissue from each individual. This allowed us to express each metabolite concentration as a 'pool per tissue'. Only in hemolymph samples, where we could measure the exact sample volume, we could also calculate the concentration in $mmol.L^{-1}$.

Extraction of metabolites and analytical MS platforms

Analyses were performed for whole *C. costata* larvae (pools of five larvae taken in four replicates) or dissected tissues (pools of 30 tissues in four replicates). Whole larvae were weighed to obtain fresh mass, immersed in LN_2, and stored at −80°C until analysis. Dissected tissues were collected in ice-cold extraction buffer (see below) and stored at −80°C until analysis. Frozen samples of whole larvae/tissues were melted on ice and homogenized in 400 µL of extraction buffer (methanol:acetonitrile:deionized water in a volume ratio of 2:2:1). The methanol and acetonitrile (Optima LC/MS) were purchased from Fisher Scientific (Pardubice, Czech Republic) and the deionized water was prepared using Direct Q 3UV (Merck, Prague, Czech Republic). Internal standards, *p*-fluoro-DL-phenylalanine, methyl α-D-glucopyranoside (both from Sigma-Aldrich, Saint Luis, MI, USA) were added to the extraction buffer, both at a final concentration of 200 nmol.mL^{-1}. Samples were homogenized using a TissueLyser LT (Qiagen, Hilden, Germany) set to 50 Hz for 5 min (with a rotor pre-chilled to −20°C). Homogenization and centrifugation (at 20,000 g for 5 min at 4°C) were repeated twice and the two supernatants were combined.

We performed targeted analyses of 49 select metabolites (Additional file 2: Table S1) using a combination of four different mass spectrometry-based (MS) analytical platforms. Three platforms have been described previously: ECF-LC/MS and ECF-GC/MS [57] and SILYL-GC/MS [58]. Further details on instrumentation and chromatographic columns can be found in Table S1. The fourth platform, HILIC-LC/MS [59], was performed using a high resolution Orbitrap mass spectrometer (HRMS) Q Exactive Plus coupled to a Dionex Ultimate 3000 liquid chromatograph and a Dionex open autosampler (all from ThermoFisher Scientific, San Jose, CA, USA). Full scan HRMS positive and negative ion mass spectra were acquired in separate runs with a mass range of 70-1000 Da at 70,000 resolution (at mass *m/z* 200). The HRMS settings were: Scan rate at ± 3 Hz, 3 x 106 automatic gain control (AGC) target, and maximum ion injection time (IT) 100 ms. The source ionization parameters were as follows: (+/-) 3000 kV spray voltage, 350°C capillary temperature, sheath gas at 60 au, auxiliary gas at 20 au, spare gas at 1 au, probe temperature 350°C, and S-Lens level at 60 au. For accurate mass identification, we used lock masses of 622.0290 Da for the positive ion mode and 301.9981 Da for the negative ion mode. Chromatographic separation of the metabolites was carried out on the SeQuant ZIC-pHILIC (150 mm x 4.6 mm i.d., 5 µm, Merck, Darmstadt, Germany), with a mobile

phase flow rate of 450 µL.min^{-1}, injection volume of 5 µL, and column temperature of 35°C. The mobile phase: A = acetonitrile (ThermoFisher Scientific), B = 20 mM aqueous ammonium carbonate (pH = 9.2 adjusted with NH$_4$OH, Sigma-Aldrich); gradient: 0 min, 20% B; 20 min, 80% B; 20.1 min, 95% B; 23.3 min, 95% B; 23.4 min, 20% B; 30.0 min 20% B. Data were collected and metabolites identified using the in-house Metabolite Mapper® platform, equipped with an internal metabolite database, in conjunction with XcaliburTM software (v2.1, ThermoFisher Scientific).

All 49 metabolites were identified against relevant standards (Sigma-Aldrich) and subjected to quantitative analysis using a standard calibration curve method. The analytical results were validated by simultaneously running blank samples (no larvae in the sample), standard biological quality control samples (the periodic analysis of a standardized larva/tissue sample – the pool of all samples), and quality control mixtures of amino acids (AAS18, Sigma-Aldrich).

Matrix-assisted laser desorption/ionization mass spectrometry imaging (MALDI-MSI)
The MALDI-MSI [60-62] was conducted in larvae of two contrasting phenotypes (FS and FT, Figure 5a) and also on FT-larvae that were slowly pre-frozen to -30°C (FT-frozen, Fig. 5b). Larvae were rinsed in water, incubated for 3 min in 2% gelatin, and then submerged into 5 mL of 10% gelatin in a plastic mold (Tissue-Tek Cryomold, Sakura, Mdesa, Czech Republic) placed on an ice-cold stage. The molds with solid gelatin were then immersed in isopentane pre-cooled to -140°C in LN$_2$ vapors for 3 min and transferred to -80°C for storage until further processing. Preparation of the FT-frozen larvae was modified as shown in Fig. 6.

Figure 6: Preparation of FT-frozen larvae for MALDI-TOF/MS analysis.
FT larvae were embedded in gelatin in a plastic mold atop a layer of moist cellulose to which a small ice crystal was added to initiate freezing. The mold had a hole in the bottom (0.5 cm in diameter) to allow ice crystal spreading from the moist cellulose to the gelatin. The whole setup was placed on an aluminum stage pre-cooled to 0°C in the programmable thermostat F32-ME (Julabo, Seelbach, Germany) and the optimal freezing protocol (Fig. 5b) was started. As soon as the temperature inside the gelatin reached -30°C, we immersed the mold into isopentane at -140°C. Using a K-type thermocouple attached to the PicoLog TC-08 datalogger (Pico technology, St. Neots, UK), we verified that the gelatin inside plastic mold was inoculated by external ice crystals at relatively mild subzero temperatures of -1°C to -2°C. We also verified that FT-frozen larvae survive this treatment and are able to move after re-warming.

For cryo-sectioning, the frozen molds were placed into a cryostat chamber (Leica CM1950, Germany) pre-cooled to -20°C for 1 h before cutting. Sections of 10 μm thickness were mounted on ITO glass (Bruker, Czech Republic), which was previously washed with hexane and then with 2-propanol. Sections of the FS and FT larval variants were thaw-mounted on warm glass (standard procedure MALDI-TOF/MSI), while sections of the FT-frozen larval variant were mounted on a gold-coated ITO glass pre-cooled to -20°C in order to preserve the situation as it developed during slow pre-freezing (extracellular ice formation and tissue freeze-dehydration). The sections were dried for 15 min in a desiccator at low air pressure (at room temperature for the FS and FT variants, while at -20°C for the FT-frozen variant), then sealed in plastic foil inside a plastic mailer, evacuated, and stored at -80°C until further processing. The ITO glasses with sections were transferred to room temperature, left to temper for 15 min, and dried in a desiccator for 15 min. Samples were then sprayed with a matrix solution of 9-aminoacridine hydrochloride, 7 mg/mL in 70% ethanol (v/v). A TM-Sprayer 3 (HTX-Technologies, USA) was used to cover the tissue with the matrix solution using the following settings: Nozzle temperature 50°C, 12 passes, flow 0.035 mL/min, nozzle velocity 1000 mm/min, track spacing 2 mm, HH pattern, gas pressure 10 psi, gas flow rate 2 L/min, drying time 0 sec, nozzle height 40 mm and propelled with 50% methanol (v/v). After spraying, the samples were dried in a desiccator for 15 min.

Spectral images were acquired using a rapifleX MALDI-TOF/TOF spectrometer (Bruker, Germany) in reflector negative mode with a 355 nm smartbeam™ 3D laser with a spatial resolution of 10 x 10 μm in the range of m/z 20-1000, at a constant laser fluence of 72% and laser frequency of 5 kHz. Two hundred images were accumulated from every position. The instruments were set up with Ion source 1 to 19.973 kV, PIE 2.664 kV, Lens 11.353 kV, Reflector 1 20.810 kV, Reflector 2 1.034 kV, and Reflector 3 8.577 kV. The pulsed Ion Extraction time was set to 90 ns and detector gain was 2302 V. Data were acquired with

digitizer speed 2.5 GS/s. Samples were measured in random order. Calibration was done externally using red phosphorus, achieving precision up to 5 ppm [63].

After MALDI-MSI, tissues on ITO glass were washed three times for 2 min with 70% ethanol and afterwards washed with water. The samples were then stained with hematoxylin & eosin (H&E) and scanned with Axio Scan.Z1 (Zeiss, Germany) operated by Zen 2 software (blue edition; Zeiss, Germany). Single raw data files on the intensities of the MALDI-TOF signals of individual m/z peaks were processed using SCiLS Lab software (version 2021c, SCiLS, Bruker, Germany). The raw data were smoothed with a convolution algorithm (width 20) and all subsequent computations and images rendering were done with Total Ion Count square normalization. We collected MALDI-TOF signal-intensity datasets for 5 transversal sections of FS and FT larvae and for 6 sections of FT-frozen larvae, plus one longitudinal section of an FT larva. The analysis of the longitudinal section is presented in Figs. 1b and Figs. S3, 4. For relative quantification of signal intensities and statistical analyses, the transversal sections were used. Galleries showing the complete results for all transversal sections are included in separate Additional file 3.

Regions (or tissues) of interest in the sections were manually delimited using H&E stained images in QuPath software (version 0.2.3, University of Edinburgh, United Kingdom) [64] and imported back into SCiLS software. The summary spectrum was exported into mMass software (version 5.5.0) [65], internally re-calibrated, and annotated based on m/z values with 50 ppm tolerance using an in-house modified database based on the HMDB metabolite database considering [M-H]- and [M+Cl]- adducts [66]. In addition, we verified the identities of m/z peaks for trehalose and proline in the experiment where FS larvae were fed diet augmented by $^{13}C6$ glucose (for details, see Figs. S11, S12). The $^{13}C6$ glucose was purchased from Sigma Aldrich (product ID 389374). Selected peaks were then analyzed using SCiLS software with mean spectral intensity scaled in arbitrary units (A.U.) for each region and treatment. These spectral intensity data were then statistically compared using ANOVA followed by Bonferroni's multiple comparison test (Prism, v. 6.07, GraphPad Software, San Diego, USA).

Differential scanning calorimetry (DSC)

Thermal analysis was performed using the Q2000 calorimeter (TA Instruments, New Castle, DE, USA) as previously described [67]. The analyzed solutions (see Additional file 2: Table

S2 for a complete list) were loaded and hermetically sealed into aluminum pans. The amount of loaded solution (ca. 5 µL) was controlled by weighing with precision to 0.01 mg using a Sartorius CP 225 D balance (Sartorius AG, Goettingen, Germany). The solutions were frozen and thawed according to the following protocol: (i) start at 25°C, (ii) cool to −90°C at 10°C.min^{-1}, (iii) hold for 5 min at −90°C, and (iv) warm to 25°C at 10°C.min^{-1}. An empty pan was used as a reference, and samples were run in technical triplicates (same solution analyzed thrice). DSC results were analyzed using TA Universal Analysis 2000 software (version 4.5A Build 4.5.0.5). The onset of exotherm on the cooling scan was taken as the supercooling point (temperature of ice crystallization). Two major thermal events were analyzed on heating scans above -60°C: (i) glass transition: an inflection point of the second-order phase transition was read as the temperature of vitrification (T_g). The change in specific heat capacity (ΔCp) was derived from the difference in heat flow between the onset and the end of the glass transition; (ii) melting of bulk water: an onset of melting endotherm (first-order phase transition) was read as the melting point (m.p.), while the enthalpy of melting (ΔH, calculated from the area under the endothermic peak) served to estimate the fraction of melted (i.e. crystallized) water using the standard heat of fusion for the ice/water transition of 334 J g^{-1}. The remaining fraction of unfrozen water was considered osmotically inactive (%OIW).

Larval feeding on CP-augmented diets

Seventeen day-old FS larvae (i.e. roughly mid-3rd instar, voracious feeding stage) were extracted from the standard diet and groups of 20 individuals were moved to one of a variety of specific diets: (i) fresh standard diet (control); (ii) 'CP mix 508' diet – a standard diet augmented with five select CPs (proline, trehalose, glutamine, asparagine, and betaine) in concentrations corresponding to those observed in FT larval hemolymph (i.e. 508 mmol.kg^{-1} water in total, see Additional file 2: Table S3); (iii) five different standard diets augmented with the select CPs individually; (iv) 'CP mix minus Pro 313' diet – a mixture of the four select CPs but without the 313 mmol.kg^{-1} proline. After three days exposure to these diets, the larvae were removed and used for the LN$_2$ survival bioassay (Fig. 5b).

Supplementary Information

(Additional file 1.pdf)

Supplementary Figure S1   Metabolomic profiles in *C. costata* tissues.

Supplementary Figure S2   Sizes of metabolite pools in *C. costata* tissues.

Acknowledgements

We thank Irena Vacková, Jana Járová, and Karolina Kowalska for maintenence of *C. costata* colonies, processing samples for metabolomics, and assistance during preparation of MALDI-MSI samples, respectively. This study was supported by a Grantová Agentura České Republiky (GAČR) grant no. 19-13381S to VK. Development of metabolomics platforms were supported by a GAČR grant no. 17-22276S to PŠ. The MALDI-MS analysis was supported by the Czech Academy of Science project no. RVO 68378050, Ministry of Education, Youth and Sports (MEYS) grant no. LM2018126, and MEYS in cooperation with European Structural and Investing Funds grant no. CZ.02.1.01/0.0/0.0/16_013/0001789 to RS. The DSC analysis was supported by the Ministry of Agriculture, grant number MZE RO0418 to MF.

Author contributions

Conceptualization:	VK
Experiments:	MM, LK, JK, PV, LDM, RG, PH, JR, TŠ, MF
Formal analysis:	VK, MM, LK
Funding acquisition.	VK, PŠ, RS, MF
Supervision and methodology validation:	VK, PŠ, RS, MF
Writing – original draft:	VK
Writing – review and editing:	all authors

References

1. Somero G: **Protons, osmolytes, and fitness of internal milieu for protein function**. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology* 1986, **251**(2):R197-R213.
2. Yancey PH: **Organic osmolytes as compatible, metabolic and counteracting cytoprotectants in high osmolarity and other stresses**. *J Exp Biol* 2005, **208**(15):2819-2830.
3. Lee REJ: **A primer on insect cold-tolerance**. In: *Low Temperature Biology of Insects.* Edited by Denlinger DL, Lee REJ: Cambridge University Press; 2010.
4. Storey KB, Storey JM: **Freeze tolerance in animals**. *Physiol Rev* 1988, **68**(1):27-84.
5. Storey KB: **Organic solutes in freezing tolerance**. *Comp Biochem Physiol A* 1997, **117**(3):319-326.
6. Toxopeus J, Sinclair BJ: **Mechanisms underlying insect freeze tolerance**. *Biological Reviews* 2018, **93**:1891-1914.
7. Storey KB, Storey JM: **Biochemistry of cryoprotectants**. In: *Insects at low temperature.* Springer; 1991: 64-93.
8. Asahina E: **Frost resistance in insects**. In: *Adv Insect Physiol.* vol. 6: Elsevier; 1970: 1-49.
9. Sømme L: **Supercooling and winter survival in terrestrial arthropods**. *CBPA* 1982, **73**(4):519-543.
10. Salt R: **Principles of insect cold-hardiness**. *Annu Rev Entomol* 1961, **6**(1):55-74.
11. Storey JM, Storey KB: **Winter survival of the gall fly larva, Eurosta solidaginis: profiles of fuel reserves and cryoprotectants in a natural population**. *J Insect Physiol* 1986, **32**(6):549-556.
12. Toxopeus J, Koštál V, Sinclair BJ: **Evidence for non-colligative function of small cryoprotectants in a freeze-tolerant insect**. *Proceedings of the Royal Society B* 2019, **286**(1899):20190050.
13. Elliott GD, Wang S, Fuller BJ: **Cryoprotectants: A review of the actions and applications of cryoprotective solutes that modulate cell recovery from ultra-low temperatures**. *Cryobiology* 2017, **76**:74-91.
14. Choi YH, van Spronsen J, Dai Y, Verberne M, Hollmann F, Arends IW, Witkamp G-J, Verpoorte R: **Are natural deep eutectic solvents the missing link in understanding cellular metabolism and physiology?** *Plant Physiol* 2011, **156**(4):1701-1705.
15. Dai Y, van Spronsen J, Witkamp G-J, Verpoorte R, Choi YH: **Ionic Liquids and Deep Eutectic Solvents in Natural Products Research: Mixtures of Solids as Extraction Solvents**. *J Nat Pro* 2013, **76**(11):2162-2173.
16. Dai Y, Witkamp G-J, Verpoorte R, Choi YH: **Tailoring properties of natural deep eutectic solvents with water to facilitate their applications**. *Food Chem* 2015, **187**:14-19.
17. Castro VIB, Craveiro R, Silva JM, Reis RL, Paiva A, C. Duarte AR: **Natural deep eutectic systems as alternative nontoxic cryoprotective agents**. *Cryobiology* 2018, **83**:15-26.

18. Gertrudes A, Craveiro R, Eltayari Z, Reis RL, Paiva A, Duarte ARC: **How Do Animals Survive Extreme Temperature Amplitudes? The Role of Natural Deep Eutectic Solvents**. *ACS Sustain Chem Eng* 2017, **5**(11):9542-9553.
19. Liu Y, Friesen JB, McAlpine JB, Lankin DC, Chen S-N, Pauli GF: **Natural Deep Eutectic Solvents: Properties, Applications, and Perspectives**. *J Nat Prod* 2018, **81**(3):679-690.
20. Hornberger K, Li R, Duarte ARC, Hubel A: **Natural deep eutectic systems for nature-inspired cryopreservation of cells**. *AIChE Journal* 2021, **67**(2):e17085.
21. Pegg D: **The current status of tissue cryopreservation**. *Cryo Lett* 2001, **22**(2):105-114.
22. Fahy GM, Wowk B, Wu J: **Cryopreservation of complex systems: the missing link in the regenerative medicine supply chain**. *Rejuvenation Res* 2006, **9**(2):279-291.
23. Des Marteaux LE, Hůla P, Koštál V: **Transcriptional analysis of insect extreme freeze tolerance**. *Proc Royal Soc B* 2019, **286**(1913):20192019.
24. Moon I, Fujikawa S, Shimada K: **Cryopreservation of *Chymomyza* larvae (Diptera: Drosophilidae) at-196°C with extracellular freezing**. *Cryo Lett* 1996, **17**:105-110.
25. Shimada K, Riihimaa A: **Cold acclimation, inoculative freezing and slow cooling: essential factors contributing to the freezing-tolerance in diapausing larvae of *Chymomyza costata* (Diptera: Drosophilidae)**. *Cryo Lett* 1988, **9**:5-10.
26. Rozsypal J, Moos M, Šimek P, Koštál V: **Thermal analysis of ice and glass transitions in insects that do and do not survive freezing**. *J Exp Biol* 2018, **221**:170464.
27. Koštál V, Zahradníčková H, Šimek P: **Hyperprolinemic larvae of the drosophilid fly, Chymomyza costata, survive cryopreservation in liquid nitrogen**. *Proc Natl Acad Sci* 2011, **108**(32):13041-13046.
28. Hochachka P, Somero G: **Biochemical Adaptation. Mechanism and process in physiological evolution**: Oxford University Press; 2002.
29. Band H, Band R: **Multiple overwintering mechanisms in Chymomyza amoena larvae (Diptera: Drosophilidae) and laboratory induction of freeze tolerance**. *Experientia* 1982, **38**(12):1448-1449.
30. Grimaldi D: **The Chymomyza aldrichii Species-Group (Diptera: Drosophilidae); Relationships, New Neotropical Species, and the Evolution of Some Sexual Traits**. *J N Y Entomol Soc* 1986:342-371.
31. Sinclair BJ: **Insect cold tolerance: How many kinds of frozen?** *Eur J Entomol* 1999, **96**:157-164.
32. Zachariassen KE: **Physiology of cold tolerance in insects**. *Physiol Rev* 1985, **65**(4):799-832.
33. Toxopeus J, Koštál V, Sinclair BJ: **Evidence for non-colligative function of small cryoprotectants in a freeze-tolerant insect**. *Proc Royal Soc B* 2019, **286**(1899):20190050.
34. Rozsypal J, Koštál V, Zahradníčková H, Šimek P: **Overwintering strategy and mechanisms of cold tolerance in the codling moth (Cydia pomonella)**. *PLoS One* 2013, **8**(4):e61745.
35. Koštál V, Renault D, Rozsypal J: **Seasonal changes of free amino acids and thermal hysteresis in overwintering heteropteran insect, Pyrrhocoris apterus**. *CBPA* 2011, **160**(2):245-251.
36. Rudolph AS, Crowe JH: **A calorimetric and infrared spectroscopic study of the stabilizing solute proline**. *Biophys J* 1986, **50**(3):423-430.
37. Rasmussen PH, Jørgensen B, Nielsen J: **Aqueous solutions of proline and NaCl studied by differential scanning calorimetry at subzero temperatures**. *Thermochim Acta* 1997, **303**(1):23-30.
38. Liu W, Huang Z, He X, Jiang P, Huo X, Lu Z, Liu B: **Impacts of trehalose and l-proline on the thermodynamic nonequilibrium phase change and thermal properties of normal saline**. *Cryobiology* 2020, **96**:92-98.
39. Green JL, Angell CA: **Phase relations and vitrification in saccharide-water solutions and the trehalose anomaly**. *J Phys Chem* 1989, **93**(8):2880-2882.
40. Chen T, Fowler A, Toner M: **Literature review: supplemented phase diagram of the trehalose-water binary mixture**. *Cryobiology* 2000, **40**(3):277-282.

41.   Cesaro A, De Giacomo O, Sussich F: **Water interplay in trehalose polymorphism**. *Food Chem* 2008, **106**(4):1318-1328.

42.   Nicolajsen H, Hvidt A: **Phase Behavior of the System Trehalose-NaCl-Water**. *Cryobiology* 1994, **31**(2):199-205.

43.   Weng L, Elliott GD: **Dynamic and thermodynamic characteristics associated with the glass transition of amorphous trehalose–water mixtures**. *Phys Chem Chem Phys* 2014, **16**(23):11555-11565.

44.   Olgenblum GI, Sapir L, Harries D: **Properties of aqueous trehalose mixtures: Glass transition and hydrogen bonding**. *J Chem Theory Comput* 2020, **16**(2):1249-1262.

45.   Crowe JH, Carpenter JF, Crowe LM: **The role of vitrification in anhydrobiosis**. *Annu Rev Physiol* 1998, **60**(1):73-103.

46.   Held C, Reschke T, Müller R, Kunz W, Sadowski G: **Measuring and modeling aqueous electrolyte/amino-acid solutions with ePC-SAFT**. *J Chem Thermodyn* 2014, **68**:1-12.

47.   Qiu J, Huang H, He H, Liu H, Hu S, Han J, Yi D, An M, Wang P: **Solubility Behavior and Data Modeling of l-Proline in Different Neat and Binary Solvent Systems**. *J Chem Eng Data* 2019, **64**(12):5920-5928.

48.   McLain SE, Soper AK, Terry AE, Watts A: **Structure and hydration of L-proline in aqueous solutions**. *J Phys Chem B* 2007, **111**(17):4568-4580.

49.   Troitzsch RZ, Vass H, Hossack WJ, Martyna GJ, Crain J: **Molecular mechanisms of cryoprotection in aqueous proline: light scattering and molecular dynamics simulations**. *J Phys Chem* 2008, **112**(14):4290-4297.

50.   de Molina PM, Alvarez F, Frick B, Wildes A, Arbe A, Colmenero J: **Investigation of the dynamics of aqueous proline solutions using neutron scattering and molecular dynamics simulations**. *Phys Chem Chem Phys* 2017, **19**(40):27739-27754.

51.   Martins MAR, Pinho SP, Coutinho JAP: **Insights into the Nature of Eutectic and Deep Eutectic Mixtures**. *J Solution Chem* 2019, **48**(7):962-982.

52.   Durand E, Lecomte J, Villeneuve P: **From green chemistry to nature: The versatile role of low transition temperature mixtures**. *Biochimie* 2016, **120**:119-123.

53.   Rubinsky B, Cravalho EG, Mikic B: **Thermal stresses in frozen organs**. *Cryobiology* 1980, **17**(1):66-73.

54.   Riihimaa AJ, Kimura MT: **A mutant strain of *Chymomyza costata* (Diptera: Drosophilidae) insensitive to diapause-inducing action of photoperiod**. *Physiol Entomol* 1988, **13**(4):441-445.

55.   Kostal V, Noguchi H, Shimada K, Hayakawa Y: **Developmental changes in dopamine levels in larvae of the fly Chymomyza costata: comparison between wild-type and mutant-nondiapause strains**. *J Insect Physiol* 1998, **44**(7-8):605-614.

56.   Lakovaara S: **Malt as a culture medium for Drosophila species**. *Drosophila Inf Serv* 1969, **44**:128.

57.   Štětina T, Hůla P, Moos M, Šimek P, Šmilauer P, Koštál V: **Recovery from supercooling, freezing, and cryopreservation stress in larvae of the drosophilid fly, *Chymomyza costata***. *Sci Rep* 2018, **8**(1):4414.

58.   Škodová-Sveráková I, Prokopchuk G, Peña-Diaz P, Záhonová K, Moos M, Horváth A, Šimek P, Lukeš J: **Unique dynamics of paramylon storage in the marine euglenozoan Diplonema papillatum**. *Protist* 2020, **171**(2):125717.

59.   Škodová-Sveráková I, Záhonová K, Juricová V, Danchenko M, Moos M, Baráth P, Prokopchuk G, Butenko A, Lukáčová V, Kohútová L *et al*: **Highly flexible metabolism of the marine euglenozoan protist Diplonema papillatum**. *BMC biology* 2021, **19**(1):251.

60.   Yang FY, Chen JH, Ruan QQ, Saqib HS, He WY, You MS: **Mass spectrometry imaging: An emerging technology for the analysis of metabolites in insects**. *Arch Insect Biochem Physiol* 2020, **103**(4):e21643.

61. Tuthill II BF, Searcy LA, Yost RA, Musselman LP: **Tissue-specific analysis of lipid species in Drosophila during overnutrition by UHPLC-MS/MS and MALDI-MSI**. *J Lip Res* 2020, **61**(3):275-290.
62. Khalil S, Pretzel J, Becker K, Spengler B: **High-resolution AP-SMALDI mass spectrometry imaging of Drosophila melanogaster**. *Int J Mass Spectrom* 2017, **416**:1-19.
63. Kolářová L, Prokeš L, Kučera L, Hampl A, Peña-Méndez E, Vaňhara P, Havel J: **Clusters of monoisotopic elements for calibration in (TOF) mass spectrometry**. *J Am Soc Mass Spectrom* 2016, **28**(3):419-427.
64. Bankhead P, Loughrey MB, Fernández JA, Dombrowski Y, McArt DG, Dunne PD, McQuaid S, Gray RT, Murray LJ, Coleman HG: **QuPath: Open source software for digital pathology image analysis**. *Sci Rep* 2017, **7**(1):1-7.
65. Niedermeyer TH, Strohalm M: **mMass as a software tool for the annotation of cyclic peptide tandem mass spectra**. *PLoS One* 2012, **7**(9):e44913.
66. Wishart DS, Feunang YD, Marcu A, Guo AC, Liang K, Vázquez-Fresno R, Sajed T, Johnson D, Li C, Karu N: **HMDB 4.0: the human metabolome database for 2018**. *Nucleic Acids Res* 2018, **46**(D1):D608-D617.
67. Pokorna E, Faltus M, Machova P, Zamecnik J, Fulin M: **Grey poplar explant acclimation to improve the dehydration tolerance and cryopreservation**. *Biol Plant* 2020, **64**(1):119-128.