1

An unexpected role for the conserved ADAM-family metalloprotease ADM-2 in

Caenorhabditis elegans **molting**

Braveen B. Joseph, Sarina Meadows, Phillip T. Edeen, Shaonil Binti, and David S. Fay*

Department of Molecular Biology, College of Agriculture and Natural Resources,

University of Wyoming, Laramie, WY 82071

*Corresponding author

Abstract (294/300)

Molting is a widespread developmental process in which the external extracellular matrix

(ECM), the cuticle, is remodeled to allow for organismal growth and environmental adaptation.

Studies in the nematode *Caenorhabditis elegans* have identified a diverse set of molting-

associated factors including signaling molecules, intracellular trafficking regulators, ECM

components, and ECM-modifying enzymes such as matrix metalloproteases. *C. elegans* NEKL-2

and NEKL-3, two conserved members of the NEK family of protein kinases, are essential for

molting and promote the endocytosis of environmental steroid-hormone precursors by the

epidermis. Steroids in turn drive the cyclic induction of many genes required for molting. Here

we report a novel role for the sole *C. elegans* ADAM–meltrin metalloprotease family member,

ADM-2, as a negative regulator of molting. Whereas loss of *adm-2* led to strong suppression of

molting defects in partial loss-of-function *nekl* mutants, overexpression of ADM-2 induced

molting defects in wild-type animals. CRISPR genome editing implicated the Zn-binding motif

within the metalloprotease domain as critical for mediating the effects of ADM-2 on molting.

ADM-2 is expressed in the epidermis, and its trafficking through the endo-lysosomal network

was disrupted after NEKL depletion. We also identified the epidermally expressed low-density

lipoprotein receptor–related protein, LRP-1, as a candidate target of ADM-2 regulation.

Whereas loss of ADM-2 activity led to the upregulation of LRP-1, ADM-2 overexpression caused

a reduction in LRP-1 abundance, suggesting that ADM-2 may function as a sheddase for LRP-1.

We propose that loss of *adm-2* suppresses molting defects in *nekl* mutants by eliminating a

negative regulator of LRP-1, thereby compensating for defects in the efficiency of LRP-1 and

sterol uptake. Our findings emphasize the importance of endocytic trafficking for both the

internalization of factors that promote molting and the removal of proteins that would

otherwise be deleterious to the molting process.

Author Summary (146/150)

The molecular and cellular features of molting in nematodes share many similarities with cellular and developmental processes that occur in mammals. This includes the degradation and reorganization of extracellular matrix materials that surround cells, as well as the intracellular machineries that allow cells to communicate and sample their environments. In the current study, we found an unexpected function for a conserved protein that degrades proteins on the external surface of cells. Rather than promoting molting through extracellular matrix reorganization, the ADM-2 protease can inhibit the molting process. This observation can be explained by data showing that ADM-2 negatively regulates LRP-1, a membrane protein that promotes molting by facilitating the uptake of molecular building blocks at the cell surface that are needed for molting-related signaling. Our data provide insights into the mechanisms controlling molting and link several conserved proteins to show how they function together during development.

Introduction

The cuticle of *Caenorhabditis elegans* is an external extracellular matrix (ECM) required for locomotion, body shape maintenance, and protection from the environment [1, 2]. During larval development *C. elegans* undergoes four molts, a specialized form of apical ECM remodeling, whereby a new cuticle is synthesized under the old cuticle, which is partially degraded and shed [1, 2]. *C. elegans* molting cycles are orchestrated by conserved steroid-hormone receptors, including NHR-23 (an ortholog of human RORC) and NHR-25 (an ortholog of human NR5A1), which collectively control the oscillation of hundreds of mRNAs [2-4]. The production of molting-specific steroid-hormone ligands is thought to be dependent in part on the uptake of environmental sterols by epidermally expressed LRP-1 (the homolog of human LRP2/megalin), a member of the low-density lipoprotein receptor–related protein family [5, 6]. Consistent with this, internalization of LRP-1 by clathrin-mediated endocytosis (CME) is essential for normal molting [2, 5-7].

We have previously shown that the *C. elegans* protein kinases NEKL-2 (an ortholog of human NEK8/9) and NEKL-3 (an ortholog of human NEK6/7) promote endocytosis of LRP-1 at the apical epidermal plasma membrane. Correspondingly, loss of either NEKL-2 or NEKL-3 function leads to a reduction or delay in the expression of molting genes, a failure to complete molting, and larval arrest [2, 8-12]. NEKL-2 and NEKL-3 (NEKLs) are members of the NIMA-related kinase (NEK) protein family, mammalian orthologs of which have been studied primarily in the context of cell cycle regulation and ciliogenesis [13-28]. *C. elegans* NEKLs bind to and co-localize with several conserved ankyrin-repeat proteins, MLT-2 (an ortholog of human ANKS6), MLT-3 (an ortholog of human ANKS3), and MLT-4 (an ortholog of human INVS), referred to here collectively as MLTs, which are essential for the proper localization of NEKLs [9]. Correspondingly, loss of MLT functions leads to molting defects that are identical to those observed with loss of the NEKLs [9]. NEKLs and MLTs form two distinct complexes (NEKL-2–MLT-2–MLT-4 and NEKL-3–MLT-3) and are expressed in a punctate pattern in the large epidermal syncytium known as hyp7, in which they are specifically required [2, 8, 9].

The cellular and physiological mechanisms by which NEKLs–MLTs impact the molting process through intracellular trafficking have yet to be fully explored. It is likely that NEKLs are

required for the uptake and processing of membrane cargo, including LRP-1, that is required for

molting. Using a forward-genetics suppressor approach [29], we previously found that loss of

AP2 clathrin-adapter subunits, as well as the AP2 allosteric activator FCHO-1 can suppress

molting and trafficking defects in NEKL mutants [11]. These and other studies revealed that

NEKLs control endocytosis in part by facilitating the uncoating of sub-apical clathrin-coated

vesicles and may affect additional trafficking processes through the regulation of actin via the

CDC-42–SID-3 (corresponding to the human CDC42–ACK1/2) pathway [10, 11].

Here we report suppression of *nekl* molting defects caused by loss of the conserved

ADM-2 transmembrane metalloprotease. Curiously, although proteases have previously been

implicated as positive regulators of molting, we find that ADM-2 exerts a negative influence on

the molting process. ADM-2 belongs to the ADAM (a disintegrin and metalloprotease) family of

metallopeptidases, which are members of the zinc protease superfamily [30, 31]. ADM-2 is the

sole member of the meltrin metalloprotease subfamily in *C. elegans* [32, 33], which in humans

consists of ADAM9 (Meltrin γ), ADAM12 (Meltrin α), ADAM19 (Meltrin β), and ADAM33 [34,

35]. Notably, knockouts of meltrin family members in mammals have generally not provided

clear insights into the roles of meltrins in vivo during development, which may in part be due to

genetic redundancies [32, 36, 37]. Here we show that, unlike AP2 and *fcho-1* mutants, loss of

ADM-2 function did not suppress *nekl*-associated trafficking defects. Rather, ADM-2 was itself

dependent on NEKL–MLTs for its uptake and endocytic processing from the epidermal surface

where it may act as a negative regulator of LRP-1. Our findings further suggest that loss of

ADM-2 may specifically bypass trafficking defects in weak loss-of-function *nekl* mutants by de-

repressing LRP-1. Thus, NEKLs may be required for both the internalization of positive and

negative regulators of the molting cycle.

Results

***nekl* molting defects are suppressed by reduced function of the ADM-2 metalloprotease**

We previously described an approach to identify genetic suppressors of partial loss-of-function

mutations in NEKL kinases [29]. Whereas strains homozygous for either *nekl-2(fd81)* or *nekl-*

3(gk894345) weak loss-of-function alleles are viable, *nekl-2(fd81)*; *nekl-3(gk894345)* double

Fig 1. Loss of ADM-2 function suppresses *nekl* molting defects

(A) DIC image of *nekl-2; nekl-3* double-mutant worms. The adult worm contains a rescuing extrachromosomal array *(nekl-3⁺; sur-5::GFP)*. An arrested larva is marked by the white arrow and enlarged in the inset; the white bracket indicates the constricted region containing a double cuticle. (B, C) DIC images of *nekl-2; nekl-3* double-mutant adult worms containing the *adm-2(fd130)* (B) and *adm-2(fd163)* (C) mutant alleles. Bar size in A = 100 μm (for A–C); in inset, 20 μm. (D) Schematic diagram of the *adm-2* locus. Solid black rectangles indicate exons; introns are demarcated by black lines. Locations of the *fd163*, *fd130*, *fd208*, and *fd243–fd247* alleles are indicated by arrows. Large deletion alleles *fd298–fd302 fd313, fd316–fd318, fd228–fd230*, and *fd235–fd237* are indicated by orange and blue lines. (E) Bar plot showing percentage of viable adult-stage *nekl-2; nekl-3* worms with the indicated *adm-2* alleles (or RNAi); + indicates wild-type *adm-2*. (F) Bar plot showing reversion of suppression in *nekl-2; nekl-3 adm-2(fd130)* mutants by fosmids expressing wild-type *adm-2* and by an *adm-2* cDNA fused to GFP (pDF417). Fosmid #1 and #2 indicate two independent extrachromosomal lines. (G) Bar plot showing failure to suppress molting defects in *nekl–mlt* hypomorphic mutants by *adm-2* null mutants [*nekl-2(fd91); adm-2(fd313)*, *nekl-3(sv3); adm-2(fd316)*, and *mlt-4(sv9); adm-2(fd317)*]. Error bars in E–G represent 95% confidence intervals. p-Values were determined using Fisher's exact test; ****p ≤ 0.0001.

mutants (hereafter referred to as *nekl-2; nekl-3* mutants) are synthetically lethal and exhibit

~98.5% larval arrest due to L2/L3 molting defects [9]. In the absence of a suppressor mutation,

propagation of *nekl-2; nekl-3* mutants requires the presence of a GFP-marked *nekl-2⁺* or *nekl-3⁺*

transgenic rescuing array (Fig 1A and 1E). In contrast, strains homozygous for the suppressor

alleles *fd130* or *fd163* of are ~80% viable and propagate in the absence of a rescuing array (Fig

1B, 1C, and 1E).

Using our protocols for whole-genome sequencing and bioinformatical analysis [29], we

identified the causal mutation corresponding to *fd130* to be a G-to-A transition in exon 10 of

adm-2/C04A11.4 (Fig 1D). *fd130* converts codon 494 (TG<u>G</u>; W) into a premature translational

termination signal (TG<u>A</u>; stop codon), resulting in the predicted truncation of the 952-amino-

acid protein. Correspondingly, the independently isolated allele *fd163* is a G-to-A transition in

the conserved 5' splice donor sequence in the first intron of *adm-2* (<u>G</u>T to <u>A</u>T) (Fig 1D). The

fd163 mutation is predicted to result in a stop codon immediately following R66. Using

CRISPR/Cas9 methods we isolated several additional *adm-2* alleles including *fd208*, a 1-bp

deletion that causes a frameshift after Y479, along with *fd229* and *fd230*, deletions that span

exons 4–10 and that result in frame shifts after S123 and T122, respectively (Fig 1D). Like *fd130*

and *fd163*, these alleles led to similarly robust suppression of molting defects in the *nekl-2;*

nekl-3 background and are predicted to result in strong or complete loss of ADM-2 function (Fig

1E).

Several additional pieces of evidence indicate that it is loss of ADM-2 function that leads to

suppression of *nekl-2; nekl-3* molting defects. (1) *adm-2* mutations that suppress these molting

defects (e.g., *fd130* and *fd163*) are fully recessive (see Materials and Methods). (2)

Extrachromosomal expression of a fosmid clone containing wild-type genomic ADM-2

sequences strongly reversed suppression in *nekl-2; nekl-3 adm-2* mutants (Fig 1F). (3) RNAi of

adm-2 led to significant suppression of molting defects in *nekl-2; nekl-3* mutants (Fig 1E). We

note that loss of *adm-2* in wild-type backgrounds, including a strong loss-of-function deletion

allele, *fd300*, did not appear to impair development, health, or fecundity, indicating that *adm-2*

is a non-essential gene (S1A Fig). Consistent with this, no phenotypes have been previously

ascribed to *adm-2* mutations.

ADM-2 is a member of the ADAM (a disintegrin and metalloprotease) family of

metallopeptidases, with its closest human homologs belonging to the meltrin subfamily

(ADAM9/12/19/33) (S1C Fig) [32, 34, 35]. Meltrins are notable for having functional proteases

that contain a histidine-coordinated zinc-binding site, which is also found in ADM-2 (S2 Fig). Like

other meltrins, ADM-2 contains an N-terminal cysteine switch, cysteine loop, and disintegrin

domain; a transmembrane domain; and several predicted SH3-binding sites in its cytoplasmic C

terminus (S2 Fig) [31, 38]. Although linked to a range of human diseases, individual loss-of-

function mutations in mouse meltrins have generally not produced robust developmental

defects, and no phenotypes have been associated with either of the two *Drosophila* meltrin

family members [32, 36, 37].

We note that WormBase annotates two *adm-2* isoforms that are identical through exon 18

(corresponding to aa A915) but differ at the 5' ends of their 19th (terminal) exons; *adm-2a* and

adm-2b are predicted to encode 952 and 929 aa proteins, respectively. The noncanonical 18th

intron acceptor site of *adm-2b* (5-'GCAAAAG-3') occurs 7 bp upstream of the corresponding

acceptor site of *adm-2a* (5'-ATTTCAG-3') and terminates translation 76 bp upstream of the stop

codon of *adm-2a*. The peptide regions corresponding to exon 19 of *adm-2a* (37 aa) and *adm-2b*

(14 aa) do not contain any known domains nor were homologies detected with other proteins.

To determine if other *C. elegans* ADAM family members may also contribute to molting control,

we tested ten other family members for their ability to suppress *nekl-2; nekl-3* molting defects

(S1B Fig). We failed to detect suppression after inhibition of each gene using RNAi (dsRNA)

injection methods, which were effective in promoting suppression when targeting *adm-2*. Thus,

the suppression of *nekl*-associated molting defects by *adm-2* is unique among the *C. elegans*

ADAM family members.

Suppression by ADM-2 occurs via a mechanism that is distinct from trafficking suppressors

Loss of AP2 clathrin-adapter subunits and loss of the AP2 activator FCHO-1 individually suppress

strong and/or null mutations in NEKLs and MLTs through their effects on CME [11]. We

therefore tested if an *adm-2* null mutation could also suppress strong loss-of-function alleles of

nekls and *mlts*. Notably, loss of *adm-2* was unable to restore viability to *nekl-2(fd91)*, *nekl-*

3(sv3), or *mlt-4(sv9)* strong loss-of-function alleles, which typically arrest as partially constricted

L2/L3 larvae (Fig 1G). These results suggest that the mechanisms underlying the suppression of

molting defects by *adm-2* and CME-associated trafficking factors may be distinct.

Fig 2. Loss of *adm-2* function does not correct the *nekl* trafficking defects
(A–C) Representative confocal images of 2-day-old adult worms expressing LRP-1::GFP in the apical hyp7 region of the epidermis. LRP-1 expression in the *nekl-3::AID* (A), *nekl-3::AID; fcho-1(ox477)* (B), and *nekl-3::AID; adm-2(fd318)* (C) mutant backgrounds in the absence (–) and presence (+) of auxin treatment. Bar in A = 5 μm (for A–C). (D) Dot plot showing the percentage of GFP-positive pixels within the apical plane of the worm epidermis for individuals of the specified genotypes and auxin treatment groups. Group means along with 95% confidence intervals (error bars) are indicated. p-Values were obtained by comparing means using an unpaired t-test: ****p ≤ 0.0001, *p ≤ 0.05.

To directly test if loss of ADM-2 can suppress CME defects in *nekls*, we examined the

localization of GFP-tagged LRP-1/megalin, an apical transmembrane cargo that is trafficked via

CME. Using the auxin-inducible degron (AID) system to remove NEKL-3 activity in 1-day-old

adult worms [11], we observed a dramatic accumulation of LRP-1::GFP at or near the apical

membrane (Fig 2A and 2D), consistent with our previous findings [11]. As anticipated, loss of

FCHO-1 partially corrected LRP-1::GFP mislocalization defects in NEKL-3::AID-depleted worms

(Fig 2B and 2D), consistent with the ability of *fcho-1* mutations to suppress *nekl*-associated

clathrin localization and mobility defects [11]. In contrast, complete loss of ADM-2 failed to

correct LRP-1::GFP defects in NEKL-3::AID-depleted adults and if anything showed slightly

enhanced apical LRP-1::GFP accumulation relative to the NEKL-3::AID-depleted worms (Fig 2C

and 2D; also see below). Collectively, these results indicate that ADM-2 does not suppress *nekl*

molting defects by correcting CME deficits and is therefore likely to act through a distinct

mechanism.

ADM-2 is expressed in multiple tissues including the epidermis

To gain insight into how ADM-2 may affect molting in *nekl* mutants, we examined endogenously

tagged *adm-2::mScarlet* and *adm-2::GFP* strains, in which the fluorescent marker was fused to

the C terminus of the ADM-2a cytoplasmic domain. Both CRISPR-tagged versions showed a

punctate pattern within hyp7, a large epidermal syncytium that encompasses most of the

central body region of the worm, including localization to small puncta near the apical

membrane (Fig 3A–E'). Notably, both NEKLs and MLTs are specifically expressed and required in

the hyp7 syncytium [8, 9]. We also detected some differences between the localization patterns

of ADM-2::mScarlet and ADM-2::GFP. In particular, ADM-2::mScarlet was observed in larger

vesicular and tubular-like structures throughout the epidermis, whereas these structures were

mostly absent in ADM-2::GFP worms (Fig 3A–D; also see below).

ADM-2 was also detectable in seam cells, a lateral epidermal syncytium that borders hyp7 along

the length of the animal (Fig 3E and 3E'); in the anterior epidermis (S3A and A'); and in a variety

of head, tail, and centrally located neurons (Fig 3E and 3E'; S3B–C and S3B'–C' Fig). In addition,

ADM-2 was observed in proximal oogenic cells of the hermaphrodite germline, with levels

increasing in maturing oocytes, where it was localized to the cytoplasm and plasma membrane

(Fig 3F and 3F'). Likewise, ADM-2 is expressed in fertilized oocytes (Fig 3F and 3F') and

throughout embryogenesis (S3D–F and S3D'–F' Fig). In contrast, ADM-2 was not detected in

Fig 3. Expression of ADM-2 in the epidermis and other tissues
(A–D and A') Representative confocal images of ADM-2 expression in the *C. elegans* hyp7 region. ADM-2::mScarlet (A, B) and ADM-2::GFP (C, D) expression at apical (A, C) and medial (B, D) planes. A' is the inset from panel A. Bar in B = 5 µm (for A–D); in inset A' = 5 µm. (E–G and E'–G') Representative DIC (E–G) and confocal (E'–G') images of ADM-2 expression. ADM-2::mScarlet in the hyp7 hypodermis; seam cells; sensory neurons; and ALM, DNC, and VNC neurons (E, E') and in spermatheca (G, G'). ADM-2::GFP expression in the distal germline, oocytes, and spermatheca (F, F'). Bar in E' = 50 µm (for E, E'); in G' = 10 µm (for G, G'); in F' = 50 µm (for F, F').

mature sperm cells but was expressed in myoepithelial cells of the hermaphrodite spermatheca

(Fig 3G and 3G'). We also note that ADM-2::GFP expression using a multicopy reporter under

the control of the *adm-2* promoter region showed strongest expression in neurons where it

accumulated at or near the plasma membrane (S3D Fig). These findings are consistent with

ADM-2 acting in the epidermis to affect molting, but they also suggest that ADM-2 may have

functions in other tissues.

ADM-2 is trafficked through endo-lysosomal compartments and is sensitive to NEKL activities

To determine the identity of ADM-2 puncta, vesicles, and tubular-like structures in the

epidermis, we performed colocalization experiments first using a CRISPR-tagged clathrin heavy

chain marker, GFP::CHC-1 [11]. Although statistically insignificant, rare colocalization between

ADM-2::mScarlet and apical clathrin was occasionally detected (Fig 4A–C', S4A Fig). We note

that endogenous plasma membrane–localized ADM-2::GFP and ADM-2::mScarlet both

presented with extremely faint signals within hyp7 (Fig 4A and 4C), making detection and

colocalization of this population difficult to assay; this suggests that ADM-2 may be rapidly

turned over at the plasma membrane, either by CME or through a CME-independent

mechanism. We note that mammalian ADAMs, including the meltrin family, are internalized via

CME [39-42]. In addition, we observed little or no colocalization between ADM-2::mScarlet and

medial GFP::CHC-1 clathrin-containing structures (S4C–E and S4C'–E' Fig), which may represent

clathrin-coated vesicles emanating from the trans Golgi or recycling endosomes.

In contrast, ADM-2::mScarlet exhibited partial colocalization with the endosomal marker

P$_{hyp7}$::*hgrs-1::GFP* in both the sub-apical and medial planes (Fig 4D–F', S4A–B, S4F–H and S4F'–

H' Fig). HGRS-1/HRS localizes to early endosomes and multivesicular bodies and is a component

of the ESCRT-0 complex, which, together with ESCRT-I–III, promotes cargo sorting and

lysosomal targeting [43-46]. Consistent with this, medial ADM-2::mScarlet showed strong co-

localization within the lysosomal marker LysoTracker Green during intermolts (Fig 4G–I), when

lysosomes appear roughly spherical, and during molting periods (Fig 4J–L), when lysosomes

acquire a tubular morphology [47]. Thus, following rapid uptake into endosomes, ADM-2 is

likely degraded by lysosomes, although some portion may be recycled back to the plasma

membrane. Degradation of ADM-2 by lysosomes is further supported by the relative absence of

medial ADM-2::GFP accumulation (Fig 3D), as GFP is acid sensitive and fluorescence is rapidly

quenched within maturing endosomes and lysosomes [47, 48].

Given our previous observations showing that NEKL–MLT proteins are required for normal

trafficking within hyp7 [8, 10, 11], we tested if depletion of NEKLs caused changes in the

abundance and subcellular localization of ADM-2. Notably, we observed total levels of ADM-

Fig 4. ADM-2 localization to endocytic compartments is affected by NEKL activity

(A–L) Co-localization analysis of ADM-2::mScarlet with GFP::CHC-1 (A–C), P$_{hyp7}$::HGRS-1::GFP (D–F), and LysoTracker Green (G–L) within apical (A–C) and sub-apical or medial (D–L) planes of hyp7. A'–F' are insets of A–F confocal images. White arrowheads in C' and F' indicate colocalized puncta. For LysoTracker studies (G–L), representative confocal images during intermolt (G–I) and molting (J–L) stages are shown. (M–P) Apical hyp7 ADM-2::mScarlet localization in auxin-treated wild-type (M), *nekl-2::AID* (N), and *nekl-3::AID* (O) 2-day-old adults with average mean intensity calculations (P). Group means along with 95% confidence intervals (error bars) are indicated. p-Values were obtained by comparing means using an unpaired t-test: ****p ≤ 0.0001, **p ≤ 0.01. Bar in M = 5 μm (for A–F and M–O); in F' = 5 μm (for insets A'–F'); in L = 5 μm (for G–L).

2::mScarlet to increase slightly in auxin-treated *nekl-2::AID* adults, with more robust changes

occurring in auxin-treated *nekl-3::AID* animals (Fig 4M–P). Consistent with this, partial

knockdown of MLT-3 in adults by RNAi led to modest increases in the levels of ADM-2::mScarlet

and GFP::CHC-1 (S5A–B Fig). In worms that had undergone NEKL::AID depletion, ADM-

2::mScarlet increased slightly in total levels and accumulated in large internal endocytic or

lysosome-like structures (Fig 5M–P). Collectively, these findings indicate that NEKL activities

impact ADM-2 trafficking within hyp7.

The protease domain of ADM-2 is critical for its function

Most ADAM family members, including the human meltrins, contain a catalytically active

metalloprotease domain that is distinguished by three conserved histidine residues

(HExxHxxGxxH) that coordinate the binding of zinc [49, 50]. To determine if the putative

metalloprotease activity of ADM-2 is critical for its influence on molting, we CRISPR-engineered

an ADM-2 variant in which the three conserved histidine residues within the predicted Zn-

binding domain were altered [Zn-MTP: H312–H322 (**HE**LG**H**TF**G**MD**H** > **DA**LA**Y**TF**R**MD**Y**)] (Fig 5A

and 5B). Notably, this variant displayed robust suppression of *nekl-2; nekl-3* molting defects,

indicating that the protease function of ADM-2 is central to its function in molting and that its

loss is sufficient for *nekl* suppression (Fig 5C).

To further assess ADM-2 functional domains, we tested CRISPR variants designed to disrupt a

predicted N-terminal furin-cleavage site (NT-Furin: R149–R152 [**R**KK**R** > **V**KK**V**]) and a predicted

disintegrin motif (DM: E388–G396 (EPGE**ECDCG** > EPGE**VLADP**) (Fig 5A and 5B). Both variants

led to relatively weak suppression of *nekl-2; nekl-3*, suggesting that these domains are less

critical with respect to the molting-associated functions of ADM-2 (Fig 5C). Interestingly, a

variant that affects a predicted cysteine loop within the greater disintegrin domain (CysL: C438–

P459 [CRAAIGICDL**DEYCNG**ETNDCP > CRAAIGICDL**QQNGDH**ETNDCP]) strongly suppressed *nekl-*

2; nekl-3 molting defects (Fig 5A–C). To better understand the basis for the observed

suppression patterns, we made use of the AlphaFold database [51, 52] to obtain the predicted

structure of ADM-2 and then performed homology modeling using the online Robetta protein

structure prediction service [53, 54] to model the effects of our mutations. Notably, whereas

the NT-furin and DM mutations were projected to have minimal impacts on the configuration

of histidine residues within the Zn-MTP domain (Fig 5D and 5E), the CysL variant was predicted

to substantially alter the position of His322, leading to an expected reduction in

metalloprotease activity (Fig 5F). Collectively, these data indicate that the protease domain of

ADM-2 is central to its impact on the molting process and that other N-terminal domains may

play more minor or indirect roles in this context.

Fig 5. Functional analysis of ADM-2 domains
(A) Schematic representation of predicted protein domains within ADM-2. PRO, prodomain; MTP, metalloprotease domain; DIS, disintegrin domain; CYS-R, cysteine repeat region; TM, transmembrane domain; NLS, predicted nuclear localization signal; DM, disintegrin motif; CysL, cysteine loop; SH3 binding (1–3) Src homology 3 binding domains. (B) Color-coded peptide sequence of ADM-2 corresponding to panel A. For additional details see S2 Fig. (C) Percentage of viable (non-molting-defective) *nekl-2; nekl-3* adults in the indicated CRISPR-derived *adm-2* mutant backgrounds. Error bars represent 95% confidence intervals. p-Values were determined using Fisher's exact test: ****p ≤ 0.0001; ns, not significant. (D–F) Predicted three-dimensional protein structures (for amino acid region 307–328) of wild-type ADM-2 (orange) superimposed on modeled structures for the Furin-1 (D; red), DM (E; cyan), and CysL (F; violet) mutant proteins. Three conserved histidine residues (His312, His316, His322) of the Zn-metalloprotease domain are represented as sticks. (F) A predicted 6.1-Å shift of the 'tele' nitrogen atom in the imidazole ring of His322 in the CysL mutant.

We also tested the importance of the ADM-2 C-terminal domain (R696–K952) using several

CRISPR-generated lines. Deletion of most of the C terminus (CT: ΔH718–M942) led to strong

suppression of *nekl-2; nekl-3* molting defects, whereas perturbation of individual SH3-binding

domains led to suppressive effects ranging from minimal (SH3-2: P839–V853

[**P**NVQ**PPP**V**PRP**S**DDV** > **G**NVQ**GAG**V**GAG**S**LLE**] and SH3-3: K874–K884 [**K**TL**P**L**PPP**L**PK** >

ITL**EL**GAG**L**GL**]) to moderate (SH3-1: V722–P731 [**VP**VRK**A**PPPP** > **EG**VLA**A**GAVG**]) (Fig 5A–C). We

also note the presence of a potential fourth SH3-binding domain in the region of P907 (Fig 5b,

S2 Fig). As expected, none of the C-terminal mutations led to predicted changes in the

configuration of histidine residues in the Zn-binding domain (S6 Fig). A role for the C-terminus

in mediating ADAM function is not unexpected given that this domain is proposed to be

important for the regulation of cell signaling and subcellular localization and contains

specialized motifs that are thought to be involved in the 'inside-out' regulation of ADAM

metalloprotease activity [30, 31, 33, 55].

Overexpression of ADM-2 causes molting defects

Given that loss of ADM-2 did not lead to defects in molting but was capable of suppressing

molting defects in *nekl-2; nekl-3* mutants, we hypothesized that ADM-2 may normally exert an

inhibitory effect on the molting process. To test if ADM-2 is a negative regulator of molting, we

generated strains carrying the *adm-2* cDNA under the control of a heat shock–inducible

promoter (P_{hsp-16}::*adm-*2 and P_{hsp-16}::*adm-2::GFP*. Strikingly, when subjected to heat shock

during larval development (i.e., overexpression of *adm-2*), we observed molting defects in

~50% of larvae carrying these heat-shock–ADM-2 transgenes (Fig 6A and 6D). In contrast,

molting defects were not observed in heat-shocked controls or in non-heat-shocked worms (Fig

6A); ADM-2::GFP expression was specifically detected in heat-shocked worms only (Fig 6B and

6C). In addition, we observed a low frequency of blister phenotypes in animals that

overexpressed ADM-2::GFP, a defect associated with detachment of the cuticle from the

epidermis (Fig 6E). These findings are consistent with ADM-2 playing an inhibitory role in

molting and suggest that ADM-2 may also influence attachment of the epidermis to the cuticle.

ADM-2 is a negative regulator of LRP-1/megalin

ADAM/meltrin family proteins function as "sheddases", cleaving target peptides that are

positioned at or near the outer leaflet of the plasma membrane [30, 31, 56-70]. Given our

above findings, we hypothesized that ADM-2 may function as a sheddase for one or more apical

Fig 6. ADM-2 is a negative regulator of molting and LRP-1

(A) Bar plot showing the percentage of molting defects in the specified strains in the presence and absence of the $P_{hsp-16}::adm-2::GFP$ transgene in the wild-type background and in the presence and absence of heat shock. Error bars represent 95% confidence intervals. p-Values were determined using Fisher's exact test: ****p ≤ 0.0001. (B,C) Merged DIC and fluorescence images of an adult worm carrying the $P_{hsp-16}::adm-2::GFP$ transgene in the absence of heat shock (B) and after heat shock (C). (D) Molting-defective larva after heat shock. White arrow in D indicates unshed old cuticle. (E) Rare blister phenotype in a worm after ADM-2::GFP induction. White arrowheads in E indicate blisters. Bar in C = 50 μm (for B and C); in D = 50 μm; in E = 50 μm. (F,G) Representative confocal images of 1-day-old adult wild-type (F) and *adm-2(fd300)* (G) worms expressing LRP-1::GFP in the hyp7 region of the hypodermis. Bar in F = 5 μm (for F,G). (H) Dot plot showing LRP-1::GFP mean intensity (a.u.) within the apical plane of the worm hypodermis for each individual worm of the specified genotype. (I–L) Representative confocal images of 1-day-old adult worms expressing LRP-1::GFP in the hyp7 region of the hypodermis. (I,J) LRP-1 expression in wild-type control worms. (K,L) LRP-1 expression in worms containing the $P_{hsp-16}::adm-2$ transgene. Worms are shown in the absence of heat shock (I,K) and after heat shock (J,L). Bar in I = 10 μm (for I–L). (M) Dot plot showing LRP-1::GFP mean intensity (a.u.) within the apical plane of the worm hypodermis for each individual worm of the specified genotype and heat shock conditions. In H and M, group means along with 95% confidence intervals (error bars) are indicated. p-Values were obtained by comparing means using an unpaired t-test: ****p ≤ 0.0001; ***p ≤ 0.001; ns, p ≥ 0.05. (N) Genetic model of ADM-2 function in *C. elegans* molting.

membrane proteins that positively regulate the molting process. As such, loss of ADM-2 may

lead to the upregulation or de-repression of these proteins, thereby alleviating molting defects

in weak *nekl* loss-of-function mutants. One candidate protein for regulation by ADM-2 is LRP-1,

a positive regulator of molting and a known target of NEKL trafficking (also see above) [71-79].

Notably, two mammalian homologs of LRP-1 (LRP1/LRP2) are negatively regulated by matrix

metalloproteinases and ADAM family members, including the meltrin ADAM12 [71-81]. To test

this possibility, we examined levels of LRP-1::GFP in *adm-2* null mutants and observed apical

levels of LRP-1::GFP to be ~1.6-fold higher in *adm-2* mutants relative to wild type (Fig 6F–H). In

contrast, loss of *adm-2* did not alter the levels of a clathrin heavy chain reporter at the apical

membrane (S7A–D Fig), indicating that ADM-2 does not generally affect CME.

To further examine the relationship between ADM-2 and LRP-1 we tested the effects of ADM-2

overexpression on LRP-1::GFP levels. One-day-old adult worms carrying the $P_{hsp-16}::adm-2$

transgene and LRP-1::GFP were heat shocked, and LRP-1::GFP levels were assayed ~2–3 hours

later. Whereas heat shock alone did not significantly alter LRP-1::GFP levels in the absence of

the $P_{HS}::adm-2$ transgene, levels of LRP-1::GFP were ~1.6-fold lower when worms containing

the $P_{HS}::adm-2$ transgene were heat shocked relative to non-heat-shocked controls (Fig 6I–M).

These striking reciprocal findings are consistent with ADM-2 functioning as a negative regulator

of LRP-1 and suggest that suppression by loss of functional ADM-2 may occur in part through

the upregulation or de-repression of cargo including LRP-1.

Discussion

In this study we demonstrated a unique role for the sole *C. elegans* meltrin family member,

ADM-2, as a negative regulator of the molting process. Whereas loss of ADM-2 did not overtly

impact the molting cycle, overexpression of ADM-2 during larval development did lead to

molting defects and larval arrest. Moreover, our previously described non-biased forward

genetic screen identified two independent loss-of-function alleles of *adm-2* as suppressors of

larval arrest and molting defects in *nekl-2; nekl-3* mutants [29]. Previous studies have

implicated proteases as positive regulators of molting including NAS-36, NAS-37, CPZ-1, and

SURO-1 [82-87]. Extracellular proteases have been suggested to be important for the

detachment and degradation of the old cuticle and may also play a dynamic role in ECM

remodeling during new cuticle synthesis [2]. Interestingly, roles for protease inhibitors in

molting have also been reported, as loss of the Kunitz domain–containing protease inhibitors

MLT-11 and BLI-5 lead to molting defects [83, 88, 89]. Protease inhibitors have been suggested

to be important for temporally or spatially restricting the activity of extracellular proteases

during the molting cycle. Consistent with this, epidermally expressed proteases and protease

inhibitors are highly enriched among genes that are transcriptionally regulated with the molting

cycle, indicating tight control of their proteolytic activity [90, 91].

Although we previously reported the suppression of *nekl* molting defects by mutations affecting

genes closely connected to CME, our data do not support a role for ADM-2 in the regulation of

intracellular trafficking per se. Rather, our findings are consistent with ADM-2 being a cargo of

CME, including during its passage through endosomes and its turnover in lysosomes. Moreover,

we observed increased levels of ADM-2 after NEKL knockdown along with the retention of

ADM-2 in intracellular compartments of the epidermis. Collectively these findings suggest that

loss of NEKL functions may lead to increased or aberrant ADM-2 activity, which may contribute

to molting defects in these mutants.

Notably, we identified LRP-1 as a candidate target of ADM-2 sheddase activity. Epidermal LRP-1

levels were increased in *adm-2* null mutants and were reduced when ADM-2 was

overexpressed. Moreover, alteration of the ADM-2 metalloprotease domain was sufficient to

mediate strong suppression of *nekl* molting defects, indicating that proteolytic/sheddase

activity is central to the effects of ADM-2 on molting. Notably, human LRP1 is a known target

for cleavage by ADAM10, ADAM12, and ADAM17 [71-79]. Moreover, after cleavage by ADAMs,

the solubilized extracellular domain of LRP1 retains its ability to bind apolipoproteins with high

affinity. This was proposed to decrease the effective concentration of ligand available for

internalization by membrane-bound LRP1, leading to a reduction in cholesterol internalization

[80]. In addition, matrix metalloproteases have been proposed to mediate the cleavage and

proteolysis of mammalian LRP2/megalin, although the identity of the protease(s) was not

determined [77-81]. Given that *C. elegans* LRP-1 is most similar to mammalian LRP2, our study

suggests that LRP2 may be regulated by meltrin family members. Together, our data are

consistent with ADM-2 playing a conserved role in the repression of LRP family members.

Our findings, together with published data from our lab and other research groups, point to a

working model in which epidermal intracellular trafficking may play two roles in the molting

process (Fig 6N). Based on previous studies, endocytosis is required for the internalization of

factors required for molting, such as steroid hormone precursors [5, 6], and may also be

important for recycling old cuticle components, such as collagens [47, 92]. This function is

consistent with our recent observation that loss of *nekls* leads to defects in the transcriptional

upregulation of those molting genes for which their expression depends on the activation of

nuclear hormone receptors [12]. In addition, endocytosis may also be required for the uptake

and degradation of cargo that would otherwise exert an inhibitory effect on molting. Impaired

endocytic trafficking in *nekl* mutants may simultaneously lead to a reduction in cholesterol

uptake via LRP-1 and an increase in the levels of a negative regulator of LRP-1, ADM-2 (Fig 6N).

In this model, loss of *adm-2* would lead to de-repression of LRP-1, and possibly other positive

effectors of molting, thereby compensating for a partial loss of NEKL trafficking functions. In

contrast, when NEKL functions are severely reduced, loss of ADM-2 would not be expected to

offset a more severe deficiency in the uptake of steroid precursors, consistent with our

observation that *adm-2* mutations do not suppress strong loss-of-function alleles in *nekls*. In

summary, our findings expand the roles for NEKLs and intracellular trafficking in the molting

process and implicate ADM-2 as a negative regulator of the molting process.

Materials and Methods

Strains. *C. elegans* strains were maintained according to standard protocols [93] and were

propagated at 22°C, unless stated otherwise. Strains used in this study are listed in S1 Table.

Transgenic Rescue. Fosmids containing rescuing sequences for *adm-2*/C04A11.4

(WRM0620dD12, WRM0632aG02, and WRM0610cA04, 2–6 ng/µl each + *sur-5::RFP* [pTG96],

50–100 ng/µl) were injected into strain WY1342. Stable strains (WY1386 and WY1388)

containing rescuing arrays for both *nekl-3* (*fdEx286*; GFP+) and *adm-2* (*fdEx315* or *fdEx356*;

RFP+) were scored to determine the percentage of viable RFP+ progeny.

Determination of dominant versus recessive alleles. To distinguish between dominant and

recessive alleles, we first crossed suppressed *nekl-2*(*fd81*); *nekl-3*(*gk894345*) *fd130*

hermaphrodites to WY1145 [*nekl-2*(*fd81*); *nekl-3*(*gk894345*); *fdEx286 (nekl-3$^+$ + sur-5::GFP)*]

males and scored for suppression of GFP$^-$ cross-progeny males. For *fd130* and *fd162*, 50/96 and

30/52 viable cross-progeny adult males were GFP$^-$, respectively, indicating that these mutations

are either dominant or on LGX. We next crossed *nekl-2*(*fd81*); *nekl-3*(*gk894345*) *fd130*

hermaphrodites to WY1232 [*nekl-2*(*fd81*); *nekl-3*(*gk894345*); *fdEx186 (nekl-3$^+$ + sur-5::GFP)*;

fdEx197 (sur-5::RFP)] males and scored for suppression of GFP$^-$ RFP$^+$ cross-progeny

hermaphrodites. In the case of cross-progeny *fd130/+* adult hermaphrodites, 62/62 were either

GFP$^+$ RFP$^-$ or GFP$^+$ RFP$^+$; no GFP$^-$ RFP$^+$ adults were observed. Similarly, 182/185 *fd162/+* adult

hermaphrodites were either GFP$^+$ RFP$^-$ or GFP$^+$ RFP$^+$, and only 3/185 were GFP$^-$ RFP$^+$. Given the

~98.5% penetrance of *nekl-2*(*fd81*); *nekl-3*(*gk894345*) larval lethality [29], our results indicate

that *fd130* and fd162 are fully recessive but are on LGX.

RNAi. Primers containing the binding motif for T7 RNA polymerase (5'-

TAATACGACTCACTATAGGGAGA-3') and corresponding to *adm-2* (5'-

GACCACAACAATGATACGGTCGAA-3'; 5'-CCTGGACACAATGCAGCATTTTGA-3'), *unc-71* (5'-

TGTCGTCGACGGTTCCGAAGA-3'; 5'-GCATCAGACAGACCAGGCATAG-3'), *adm-4* (5'-

ATGCATTCAATACACGTGTGA-3'; 5'-CTTCCTCTCCCAGATATATCGT-3'), *sup-17* (5'-

AGTGTCAACCTGGTCTTCCTG-3'; 5'-CTGTGCCCATTGTGTTAGAGTTTC-3'), *mig-17* (5'-

CTCAGCTACACAAGGAATGGC-3'; 5'-TTCGCACACGTTCTACAACA-3'), *tag-275* (5'-

TGTTCTCGCGTCATTCGTTGC-3'; 5'-ACTCGGTTTATTGGAACATTTGGC-3'), *F27D9.7* (5'-

CAACATTCTGTGCGATGCGGT-3'; 5'-TTAAATGGGCGCGACAGATCC-3'), *adt-1* (5'-

GTCAGTGCACTCACTGGACAT-3'; 5'-GGTTAGGCATGGCCTGAATCT-3'), *adt-2* (5'-

GAAGACGAAACCGAAGTCTGC-3'; 5'-TTACCTCCCCATGCAGCATTT-3'), *adt-3* (5'-

CAGGTATGTAACGGTGACTCCA-3'; 5'-CATTACACATGGTCCGGTTTC-3'), and *gon-1* (5'-

TGGATCACTGAAGATGTGTCT-3'; 5'- GCACTCCAATCAGTATTTCTC-3') were used to generate

dsRNA using standard methods [94]. After injection at 0.8–1.0 µg/µl into WY1145

hermaphrodites, F1 progeny were scored for adult viability. For RNAi feeding experiments, the

relevant bacterial strains were obtained from Geneservice and IPTG (8 mM) was added to

growth plates [95]. Worm strains were grown on *lin-35(RNAi)* plates for two generations to

increase RNAi susceptibility [96]. Second-generation fourth larval stage (L4) worms growing on

lin-35(RNAi) plates were transferred to experimental plates and were imaged after 48 hours.

RNAi feeding experiments were performed at 20°C.

ADM-2 CRISPR mutant alleles. Design of repair sequences containing introduced restriction

sites was facilitated using CRISPRcruncher [97]. For details on primers sequences see S2 File.

<u>Strong loss-of-function alleles.</u> Alleles *fd228–fd230* and *fd235–fd237* were generated using

guide dual sequences SB1 and SB2, PCR amplification primers SB3 and SB4, and the sequencing

primer SB6. In *fd228–fd230* and *fd235–fd237*, an ~3.2-kb region spanning *adm-2a* exons 3–9 is

deleted. *fd228* is an indel predicted to encode sequences through T122 of ADM-2a, followed by

eight divergent amino acids and a stop codon. *fd229* is an indel predicted to encode sequences

through S123 of ADM-2a, followed by a stop codon. *fd230* is an indel predicted to encode

sequences through T122 of ADM-2a, followed by a stop codon. *fd235* is an indel predicted to

encode sequences through T122 of ADM-2a, followed by 23 new amino acids and a stop codon.

fd236 is an indel predicted to encode sequences through F118 of ADM-2a, followed by six new

amino acids and a stop codon. *fd237* is an indel predicted to encode sequences through T122 of

ADM-2a, followed by a single new amino acid and a stop codon. Alleles *fd298–302* were

generated using guide sequences SB46 and SB47, PCR amplification primers SB48–SB51, and

the sequencing primer SB52. In *fd298–fd302*, *fd313*, *fd316*, and *fd317* an ~7.4-kb region

spanning *adm-2a* exons 1–19 is deleted. *fd298* is an indel that encodes sequences through T2 of

ADM-2a followed by four new amino acids and a stop codon. *fd299* is a deletion that encodes

sequences through M1 of ADM-2A, followed by three new amino acids and a stop codon. *fd300*

is an indel that encodes sequences through D3 of ADM-2a, followed by 14 new amino acids and

a stop codon. *fd301* and *fd302* are identical indels that encode sequences through D3 of ADM-

2a, followed by 12 new amino acids and a stop codon. *fd313* encodes sequences through T2 of

ADM-2a, followed by 40 new amino acids and a stop codon. *fd316* deletes the normal start

codon; an alternative ATG is predicted to encode 24 new amino acids followed by a stop codon.

481 *fd317* encodes sequences through D3 of ADM-2a, followed by 16 new amino acids and a stop

482 codon. *fd318* encodes sequences through D3 of ADM-2a, followed by 8 new amino acids and a

483 stop codon.

484 <u>*adm-2* metalloprotease mutation (Zn-binding domain)</u>. Alleles *fd243–fd247* were generated

485 using the guide sequence SB27, the repair template Rep2, PCR amplification primers SB28 and

486 SB29, and sequencing primers SB31 and SB32. *fd243–fd247* change the predicted ADM-2a Zn-

487 metalloprotease active site spanning H312–H322 (**HE**L**GH**TF**G**MD**H**) to **DA**L**AY**TF**R**MD**Y** (altered

488 aa are in bold). The Zn-metalloprotease consensus motif is HEXXHXUGUXH, where U is an

489 amino acid containing a bulky hydrophobic residue. The edited locus contains an introduced

490 BstBI site.

491 <u>*adm-2* disintegrin motif mutation.</u> Allele *fd322* was generated using guide sequence DF1, the

492 repair template Rep9, PCR amplification primers DF2 and DF3, and sequencing primers DF4 and

493 DF5. *fd322* changes the predicted ADM-2a disintegrin motif (DM) spanning E388–G396

494 (EPGE**ECDCG**) to EPGE**VLADP**. The edited locus contains introduced NheI and BamHI sites.

495 <u>*adm-2* cysteine loop mutation.</u> Alleles *fd324–fd325* were generated using guide sequence DF6,

496 the repair template Rep10, PCR amplification primers DF2 and DF3, and sequencing primers

497 DF4 and DF5. *fd324–fd325* change the predicted ADM-2a cysteine loop (CysL) spanning C438–

498 P459 (CRAAIGICDL**DEYCNG**ETNDCP) to CRAAIGICDL**QQNGDH**ETNDCP. The edited locus contains

499 an introduced PstI site.

500 <u>*adm-2* furin 1 mutation.</u> Alleles *fd288-fd290* were generated using guide sequence SB17, the

501 repair template Rep6, PCR amplification primers SB18 and SB19, and sequencing primers SB20

502 and SB21. *fd288-fd290* change the predicted ADM-2a furin-1 cleavage site spanning R149–R152

503 (**R**KK**R**) to **V**KK**V**. The edited locus contains an introduced BamHI site.

504 <u>*adm-2* SH3-binding domain 1 mutation.</u> Alleles *fd248–fd251* were generated using the guide

505 sequence SB32, the repair template Rep3, PCR amplification primers SB16 and SB34, and

506 sequencing primer SB35. *fd248–fd251* alter the predicted ADM-2a SH3-1 domain spanning

507 V722–P731 (**VP**V**RK**A**PPPP)** to **EG**V**LA**A**GAVG**. The edited locus contains an introduced XhoI site.

508 <u>*adm-2* SH3-binding domain 2 mutation.</u> Alleles *fd252–fd256* were generated using the guide

509 SB36, the repair template Rep4, PCR amplification primers SB37 and SB38, and sequencing

primers SB39 and SB40. *fd252–fd256* alter the predicted ADM-2a SH3-binding domain 2 domain

spanning P839–V853 (**P**NVQ**PPP**V**PRP**S**DDV**) to **G**NVQ**GAG**V**GAG**SLLE**. The edited locus contains

an introduced XhoI site.

adm-2 SH3-binding domain 3 mutation. Alleles *fd257–fd259* were generated using the guide

sequence SB41, the repair template Rep5, PCR amplification primers SB37 and SB38, and

sequencing primers SB39 and SB40. *fd257–fd259* alter the predicted ADM-2a SH3-binding

domain 3 spanning K874–K884 (**K**TL**P**L**PPP**LPK**) to **ITL**EL**GAG**LGL**. The edited locus contains an

introduced XhoI site.

adm-2 C-terminal (cytoplasmic domain) deletion. Alleles *fd231–fd234* were generated using the

guide sequences SB12 and SB13, the repair template Rep1, PCR amplification primers SB14 and

SB15, and sequencing primer SB16. *fd231–fd234* remove sequences from H718 to M942 of

ADM-2a and introduce a XhoI site.

ADM-2 expression plasmids and strains. *adm-2::GFP* and *adm-2::mScarlet* endogenously

tagged strains were made using CRISPR/Cas9 technology in collaboration with SunyBiotech

Corporation (China). Expression vectors were generated by amplifying the *adm-2* promoter

region from fosmid WRM0620dD12 using primers SB57 and SB58. After digestion with SphI and

SalI, the ~2.1-kb PCR product was inserted into pPD95.75 to create pDF403–pDF405. *adm-2*

cDNA was amplified from plasmid pDONR201 (Horizon Inc.) using primers SB59 and SB60.

Digestion and ligation of the ~2.8-kb PCR product and pDF403 with XmaI and KpnI generated

pDF417–pDF420, which were confirmed by sequencing. pDF417 (~100 ng/µl) was injected into

N2 worms with *sur-5::RFP* (~50 ng/µl) to obtain lines carrying extrachromosomal arrays

(*fdEx353*, *fdEx354*). CRISPR-tagged *adm-2::mScarlet* and *adm-2::GFP* strains contain codon

optimized fluorescent-reporter insertions just preceding the *adm-2a* stop codon (SUNY

biotech). In addition, sequences upstream of the stop codon contain the indicated (bold) silent

mutations (LGX 13695343–13695387).

TC**T**GAAGATGCAGCTGCAACCGAAGAAAAAGTAGATGTTCGCTC**C** (wild type)

TC**A**GAAGATGCAGCTGCAACCGAAGAAAAAGTAGATGTTCGCTC**G** (CRISPR tagged)

adm-2 and _adm-2::GFP_ heat shock strains. An ~4.6-kb _adm-2::GFP::unc-54_ 3'UTR cDNA

product was obtained by digesting pDF420 with XbaI and ApaI enzymes. Digestion of heat shock

vectors pPD49.78 and pPD49.83 was performed using NheI and ApaI to obtain an ~3-kb vector

backbone. Ligation of the _adm-2_ cDNA with vector backbones pPD49.83 and pPD49.78

generated pDF429–pDF431 and pDF432–pDF434, respectively. Likewise, an ~2.8-kb _adm-2::GFP_

product was obtained by digesting pDF420 with XbaI and KpnI enzymes. Digestion of heat shock

vectors pPD49.78 and pPD49.83 was performed using NheI and KpnI to obtain an ~3-kb vector

backbone. Ligation of _adm-2::GFP_ with vector backbones pPD49.83 and pPD49.78 generated

pDF423–pDF425 and pDF426–pDF428, respectively. pDF430 and pDF433 (50 ng/μl each) were

injected into N2 with pRF4 [_rol-6(gf)_] (~50 ng/μl) to obtain lines carrying extrachromosomal

arrays (N2: _fdEx373–375_). Likewise, pDF424 and pDF425 (50 ng/μl each) were injected into N2

with pRF4 [_rol-6(gf)_] (~50 ng/μl) to obtain lines carrying extrachromosomal arrays (_fdEx381,_

fdEx382).

Heat shock methods. For Fig 6A, all worm strains were synchronized using bleach.

Synchronized L1 worms were plated on NGM plates and grown at 20°C for 2 hours. The plates

were then incubated at 34°C for 4 hours, after which they were shifted back to 20°C for 20

hours. The plates were shifted again to 34°C for 4 hours and were subsequently grown at 20°C

for 20–24 hours before the percentage of molting-defective worms was determined. For Fig 6I–

L, 1-day-old adult worms grown at 20°C were heat shocked at 34°C for 4 hours and were shifted

to 20°C for 2–3 hours before imaging.

Protein domain identification and alignment tools. The following sites were used to identify

domains within ADM-2 and human ADAM homologs:

http://nls-mapper.iab.keio.ac.jp/cgi-bin/NLS_Mapper_form.cgi

https://www.ncbi.nlm.nih.gov/Structure/cdd/wrpsb.cgi

https://www.ebi.ac.uk/interpro/

https://prosite.expasy.org/

https://psort.hgc.jp/form2.html

http://www.cbs.dtu.dk/services/TMHMM/

http://www.cbs.dtu.dk/services/SignalP/

http://phobius.sbc.su.se/

http://tcoffee.crg.cat/apps/tcoffee/do:regular

https://embnet.vital-it.ch/software/BOX_form.html

Image acquisition. Fluorescence images in the following figure panels—Fig 2A–C; Fig 6F–G; S3G

Fig; and S7 Fig—were acquired using an Olympus IX81 inverted microscope with a Yokogawa

spinning-disc confocal head (CSU-X1). Excitation wavelengths were controlled using an acousto-

optical tunable filter (ILE4; Spectral Applied Research). MetaMorph 7.7 software (MetaMorph

Inc.) was used for image acquisition. z-Stack images were acquired using a 100×, 1.40 N.A. oil

objective. Fig 3A–G; Fig 3A'–G'; Fig 4A–L; Fig 6I–L; S3A–F Fig; S3A'–F' Fig; and S4C–H Fig were

acquired using an Olympus IX83 inverted microscope with a Yokogawa spinning-disc confocal

head (CSU-W1). z-Stack images were acquired using a 100×, 1.40 N.A. silicone oil objective.

cellSense3.1 software (Olympus corporation) was used for image acquisition. DIC images in the

panels Fig 1A–C and Fig 7B–E were acquired using a Nikon Eclipse epiflourescence microscope

and the cellSense3.1 software (Olympus corporation).

Image analysis. Mean intensity (measured in arbitary units, a.u.), percent of fluorescence-

positive pixels above threshold, and the colocalization analysis were performed using Fiji

software (NIH; available at https://imagej.net/Fiji/Downloads). For a given z-plane of interest,

rolling ball background subtraction was performed (radius = 50 pixels), and the polygon

selection tool was used to choose the region of hyp7 in which the mean intensity was

quantified (Fig 4, Fig 6, S5 Fig, and S7 Fig). The percentage of fluorescence-positive pixels for the

region of interest (Fig 2) was determined after thresholding, and the "Huang" thresholding

algorithm was used for strain comparisons. For colocalization, rolling ball background

subtraction was performed (radius = 25 pixels), followed by use of the mean filter (radius = 2

pixels) to minimize noise. Finally, the same thresholding algorithm was used for one particular

channel to obtain binary images to be used as masks (GFP::CHC-1 and HGRS-1::GFP—"Otsu";

ADM-2:mScarlet—"Isodata"). This binary mask was combined using the "AND" boolean

operation to the original image and the combined image was used for the colocalization

analysis. Data from Fig 6F and 6G (Olympus IX81) and Fig 6I–L (Olympus IX83) were obtained on

different confocal microscopes and thus differ in their mean intensity values.

Auxin treatment. Auxin (indole-3-acetic acid) was purchased from Alfa Aesar. A 100× stock

auxin solution (0.4 M) was made by dissolving 0.7 g of auxin in 10 ml of 100% ethanol. A

mixture of 25 µl of stock auxin solution and 225 µl of distilled water was added to plates

containing 1-day-old adult worms.

Protein 3D structure analysis. PDB file (Identifier : AF-G5EDW5-F) containing the three-

dimensional structure details of *C. elegans* ADM-2 was obtained from the AlphaFold database (

https://alphafold.ebi.ac.uk/) [51, 52]. Using the AlphaFold structure as template, homology

modeling was performed by the online Robetta structure prediction server

(https://robetta.bakerlab.org/) to obtain the predicted the three-dimensional structures of the

respective ADM-2 mutants [53, 54]. For modeling CM (Comparative modeling) option was used

and the number of models to sample was selected as 1. Other options remained unchanged.

The homology modeled three-dimensional structures were rendered, and was superimposed

onto the AlphaFold structure of ADM-2 using the PyMOL 2 software (The PyMOL Molecular

Graphics System, Version 2.0 Schrödinger, LLC.).

Statistical analyses. Statistical analyses were carried out using Prism software (GraphPad)

following established standards [98].

Acknowledgements

We thank Amy Fluet for editing this manuscript and Barth D. Grant for the HGRS-1 marker

strain. This project was supported by R35 GM136236 to DSF and by an Institutional

Development Award (IDeA) from the National Institute of General Medical Sciences of the

National Institutes of Health (P20GM103432).

Figure Legends

Fig 1. Loss of ADM-2 function suppresses *nekl* molting defects

(A) DIC image of *nekl-2; nekl-3* double-mutant worms. The adult worm contains a rescuing extrachromosomal array *(nekl-3+; sur-5::GFP)*. An arrested larva is marked by the white arrow and enlarged in the inset; the white bracket indicates the constricted region containing a double cuticle. (B, C) DIC images of *nekl-2; nekl-3* double-mutant adult worms containing the *adm-2(fd130)* (B) and *adm-2(fd163)* (C) mutant alleles. Bar size in A = 100 µm (for A–C); in inset, 20 µm. (D) Schematic diagram of the *adm-2* locus. Solid black rectangles indicate exons; introns are demarcated by black lines. Locations of the *fd163*, *fd130*, *fd208*, and *fd243–fd247* alleles are indicated by arrows. Large deletion alleles *fd298–fd302 fd313, fd316–fd318, fd228–fd230,* and *fd235–fd237* are indicated by orange and blue lines. (E) Bar plot showing percentage of viable adult-stage *nekl-2; nekl-3* worms with the indicated *adm-2* alleles (or RNAi); + indicates wild-type *adm-2*. (F) Bar plot showing reversion of suppression in *nekl-2; nekl-3 adm-2(fd130)* mutants by fosmids expressing wild-type *adm-2* and by an *adm-2* cDNA fused to GFP (pDF417). Fosmid #1 and #2 indicate two independent extrachromosomal lines. (G) Bar plot showing failure to suppress molting defects in *nekl–mlt* hypomorphic mutants by *adm-2* null mutants [*nekl-2(fd91); adm-2(fd313)*, *nekl-3(sv3); adm-2(fd316)*, and *mlt-4(sv9); adm-2(fd317)*]. Error bars in E–G represent 95% confidence intervals. p-Values were determined using Fisher's exact test; ****$p \leq 0.0001$.

Fig 2. Loss of *adm-2* function does not correct the *nekl* trafficking defects

(A–C) Representative confocal images of 2-day-old adult worms expressing LRP-1::GFP in the apical hyp7 region of the epidermis. LRP-1 expression in the *nekl-3::AID* (A), *nekl-3::AID; fcho-1(ox477)* (B), and *nekl-3::AID; adm-2(fd318)* (C) mutant backgrounds in the absence (–) and presence (+) of auxin treatment. Bar in A = 5 µm (for A–C). (D) Dot plot showing the percentage of GFP-positive pixels within the apical plane of the worm epidermis for individuals of the specified genotypes and auxin treatment groups. Group means along with 95% confidence intervals (error bars) are indicated. p-Values were obtained by comparing means using an unpaired t-test: ****$p \leq 0.0001$, *$p \leq 0.05$.

Fig 3. Expression of ADM-2 in the epidermis and other tissues

(A–D and A') Representative confocal images of ADM-2 expression in the *C. elegans* hyp7

region. ADM-2::mScarlet (A, B) and ADM-2::GFP (C, D) expression at apical (A, C) and medial (B,

D) planes. A' is the inset from panel A. Bar in B = 5 µm (for A–D); in inset A' = 5 µm. (E–G and E'–

G') Representative DIC (E–G) and confocal (E'–G') images of ADM-2 expression. ADM-

2::mScarlet in the hyp7 hypodermis; seam cells; sensory neurons; and ALM, DNC, and VNC

neurons (E, E') and in spermatheca (G, G'). ADM-2::GFP expression in the distal germline,

oocytes, and spermatheca (F, F'). Bar in E' = 50 µm (for E, E'); in G' = 10 µm (for G, G'); in F' = 50

µm (for F, F').

Fig 4. ADM-2 localization to endocytic compartments is affected by NEKL activity

(A–L) Co-localization analysis of ADM-2::mScarlet with GFP::CHC-1 (A–C), P_{hyp7}::HGRS-1::GFP

(D–F), and LysoTracker Green (G–L) within apical (A–C) and sub-apical or medial (D–L) planes of

hyp7. A'–F' are insets of A–F confocal images. White arrowheads in C' and F' indicate

colocalized puncta. For LysoTracker studies (G–L), representative confocal images during

intermolt (G–I) and molting (J–L) stages are shown. (M–P) Apical hyp7 ADM-2::mScarlet

localization in auxin-treated wild-type (M), *nekl-2::AID* (N), and *nekl-3::AID* (O) 2-day-old adults

with average mean intensity calculations (P). Group means along with 95% confidence intervals

(error bars) are indicated. p-Values were obtained by comparing means using an unpaired t-

test: ****p ≤ 0.0001, **p ≤ 0.01. Bar in M = 5 µm (for A–F and M–O); in F' = 5 µm (for insets A'–

F'); in L = 5 µm (for G–L).

Fig 5. Functional analysis of ADM-2 domains

(A) Schematic representation of predicted protein domains within ADM-2. PRO, prodomain;

MTP, metalloprotease domain; DIS, disintegrin domain; CYS-R, cysteine repeat region; TM,

transmembrane domain; NLS, predicted nuclear localization signal; DM, disintegrin motif; CysL,

cysteine loop; SH3 binding (1–3) Src homology 3 binding domains. (B) Color-coded peptide

sequence of ADM-2 corresponding to panel A. For additional details see S2 Fig. (C) Percentage

of viable (non-molting-defective) *nekl-2; nekl-3* adults in the indicated CRISPR-derived *adm-2*

mutant backgrounds. Error bars represent 95% confidence intervals. p-Values were determined

using Fisher's exact test: ****p ≤ 0.0001; ns, not significant. (D–F) Predicted three-dimensional

protein structures (for amino acid region 307–328) of wild-type ADM-2 (orange) superimposed

on modeled structures for the Furin-1 (D; red), DM (E; cyan), and CysL (F; violet) mutant

proteins. Three conserved histidine residues (His312, His316, His322) of the Zn-metalloprotease

domain are represented as sticks. (F) A predicted 6.1-Å shift of the 'tele' nitrogen atom in the

imidazole ring of His322 in the CysL mutant**.**

Fig 6. ADM-2 is a negative regulator of molting and LRP-1

(A) Bar plot showing the percentage of molting defects in the specified strains in the presence

and absence of the P_{hsp-16}::*adm-2*::*GFP* transgene in the wild-type background and in the

presence and absence of heat shock. Error bars represent 95% confidence intervals. p-Values

were determined using Fisher's exact test: ****p ≤ 0.0001. (B,C) Merged DIC and fluorescence

images of an adult worm carrying the P_{hsp-16}::*adm-2*::*GFP* transgene in the absence of heat

shock (B) and after heat shock (C). (D) Molting-defective larva after heat shock. White arrow in

D indicates unshed old cuticle. (E) Rare blister phenotype in a worm after ADM-2::GFP

induction. White arrowheads in E indicate blisters. Bar in C = 50 µm (for B and C); in D = 50 µm;

in E = 50 µm. (F,G) Representative confocal images of 1-day-old adult wild-type (F) and *adm-*

2(fd300) (G) worms expressing LRP-1::GFP in the hyp7 region of the hypodermis. Bar in F = 5 µm

(for F,G). (H) Dot plot showing LRP-1::GFP mean intensity (a.u.) within the apical plane of the

worm hypodermis for each individual worm of the specified genotype. (I–L) Representative

confocal images of 1-day-old adult worms expressing LRP-1::GFP in the hyp7 region of the

hypodermis. (I,J) LRP-1 expression in wild-type control worms. (K,L) LRP-1 expression in worms

containing the P_{hsp-16}::*adm-2* transgene. Worms are shown in the absence of heat shock (I,K)

and after heat shock (J,L). Bar in I = 10 µm (for I–L). (M) Dot plot showing LRP-1::GFP mean

intensity (a.u.) within the apical plane of the worm hypodermis for each individual worm of the

specified genotype and heat shock conditions. In H and M, group means along with 95%

confidence intervals (error bars) are indicated. p-Values were obtained by comparing means

using an unpaired t-test: ****p ≤ 0.0001; ***p ≤ 0.001; ns, p ≥ 0.05. (N) Genetic model of ADM-

2 function in *C. elegans* molting.

S1 Fig. Loss of other *C. elegans* ADAM family members does not suppress *nekl* defects

(A) Dot plot showing average brood sizes for 10 individual wild-type and *adm-2(fd300)* mutant

worms. (B) Bar plot showing the failure of most *C. elegans* ADAM family members to suppress

molting defects in *nekl-2; nekl-3* mutants. (C) Table showing ADM-2 *C. elegans* orthologs and

their corresponding human homologs. Error bars in A,B represent 95% confidence intervals. p-

Values were determined using an unpaired t-test (A) (ns, p ≥ 0.05) or Fisher's exact test (B):

****p ≤ 0.0001.

S2 Fig. Alignment of *C. elegans* ADM-2 with human ADAMs

Peptide alignment of *C. elegans* ADM-2 with human meltrin family members

(ADAM9/12/19/33). Predicted domains of ADM-2 are color coded. NLS, nuclear localization

domain.

S3 Fig. Additional ADM-2 expression

(A–F and A'–F') Representative DIC (A–F) and confocal (A'–F') images of ADM-2 expression

showing the anterior hypodermis (A, A'), nerve ring (B, B'), tail neurons (C, C'), and various

stages of embryonic development (D–F'). Bar in A' = 10 μm (for A, A'); in B' = 10 μm (for B, B');

in C' = 10 μm (for C, C'); in F' = 10 μm (for D–F'). (G) Representative confocal image of an L2

larva expressing multi-copy P_{adm-2}::ADM-2::GFP in the plasma membrane of head neurons. Bar

in G = 25 μm.

S4 Fig. Supplemental colocalization data

(A,B) Dot plots showing quantification of Mander's overlap coefficient for the overlap of ADM-

2::mScarlet with GFP::CHC-1 and P_{hyp7}::HGRS-1::GFP proteins within the apical (A) and medial

(B) planes. Mean values and 95% confidence intervals (error bars) are indicated. p-Values were

calculated using an unpaired test: ****p ≤ 0.0001, **p ≤ 0.01. (C–H) Representative confocal

images of GFP::CHC-1 (C), P_{hyp7}::HGRS-1::GFP (F), and ADM-2::mScarlet (D,G) within the hyp7

medial plane. C'–H' are insets of C–H confocal images. Bar in E= 10 µm (for C–H); in E' = 5 µm

(for insets C'–H'). (E',H') White arrows show ADM-2 large vesicular structures that do not

colocalize with GFP::CHC-1 and P_{hyp7}HGRS-1::GFP puncta, which are lysosomes. Cyan arrows

indicate vesicles containing ADM-2 that colocalize with GFP::CHC-1 and P_{hyp7}::HGRS-1::GFP.

S5 Fig. ADM-2 levels are increased slightly upon weak loss of *mlt-3*

(A,B) Dot plot showing the mean intensity (a.u.) of GFP::CHC-1 (A) and ADM-2::mScarlet (B)

expression in the presence of Control RNAi (i.e., empty vector) and *mlt-3* RNAi. Group means

along with 95% confidence intervals (error bars) are indicated. p-Values were obtained by

comparing means using an unpaired t-test: **p ≤ 0.01, *p ≤ 0.05.

S6 Fig. Mutations in the C-terminal region of ADM-2 do not cause conformational changes to

the metalloprotease active site

Three-dimensional protein structure of the region including amino acids 307–328 of wild-type

ADM-2 (orange) superimposed on modeled structures of SH3-1, SH3-2, and SH3-3 mutant

proteins.

S7 Fig. Clathrin expression is not perturbed by loss of ADM-2 function

(A,B) Representative confocal images of GFP::CHC-1 expression in the hyp7 region of the

hypodermis in the apical plane in wild-type (A) and *adm-2(fd300)* null mutant (B) 1-day-old

adult worms. Bar in A = 10 µm (for A,B). (C,D) Dot plots showing GFP::CHC-1 mean intensity

(a.u.) (C) and the percentage of GFP-positive pixels (D) within the apical plane for each

individual worm of the specified genotype. In C and D, group means along with 95% confidence

intervals (error bars) are indicated. p-Values were obtained by comparing means using an

unpaired t-test. ns, p > 0.05.

S1 Table. List of all the strains used in this study (pdf).

S1 File. Compilation or raw data used in this study (MS Excel).

S2 File. List of primers used in this study (MS Excel).

References

1.   Page AP, Johnstone IL. The cuticle. WormBook. 2007:1-15. Epub 2007/12/01. doi: 10.1895/wormbook.1.138.1. PubMed PMID: 18050497.

2.   Lazetic V, Fay DS. Molting in C. elegans. Worm. 2017;6(1):e1330246. Epub 2017/07/14. doi: 10.1080/21624054.2017.1330246. PubMed PMID: 28702275; PubMed Central PMCID: PMCPMC5501215.

3.   Antebi A. Nuclear hormone receptors in C. elegans. WormBook. 2006:1-13. Epub 2007/12/01. doi: 10.1895/wormbook.1.64.1. PubMed PMID: 18050471; PubMed Central PMCID: PMCPMC4781232.

4.   Gissendanner CR, Crossgrove K, Kraus KA, Maina CV, Sluder AE. Expression and function of conserved nuclear receptor genes in Caenorhabditis elegans. Dev Biol. 2004;266(2):399-416. Epub 2004/01/24. doi: 10.1016/j.ydbio.2003.10.014. PubMed PMID: 14738886.

5.   Kang YL, Yochem J, Bell L, Sorensen EB, Chen L, Conner SD. Caenorhabditis elegans reveals a FxNPxY-independent low-density lipoprotein receptor internalization mechanism mediated by epsin1. Mol Biol Cell. 2013;24(3):308-18. Epub 2012/12/18. doi: mbc.E12-02-0163 [pii]
10.1091/mbc.E12-02-0163. PubMed PMID: 23242996; PubMed Central PMCID: PMC3564534.

6.   Yochem J, Tuck S, Greenwald I, Han M. A gp330/megalin-related protein is required in the major epidermis of Caenorhabditis elegans for completion of molting. Development. 1999;126(3):597-606. Epub 1999/01/07. PubMed PMID: 9876188.

7.   Roudier N, Lefebvre C, Legouis R. CeVPS-27 is an endosomal protein required for the molting and the endocytic trafficking of the low-density lipoprotein receptor-related protein 1 in Caenorhabditis elegans. Traffic. 2005;6(8):695-705. Epub 2005/07/07. doi: 10.1111/j.1600-0854.2005.00309.x. PubMed PMID: 15998324.

8.   Yochem J, Lazetic V, Bell L, Chen L, Fay D. C. elegans NIMA-related kinases NEKL-2 and NEKL-3 are required for the completion of molting. Dev Biol. 2015;398(2):255-66. Epub 2014/12/20. doi: S0012-1606(14)00634-4 [pii]
10.1016/j.ydbio.2014.12.008. PubMed PMID: 25523392; PubMed Central PMCID: PMC4314388.

9.   Lazetic V, Fay DS. Conserved ankyrin repeat proteins and their NIMA kinase partners regulate extracellular matrix remodeling and intracellular trafficking in Caenorhabditis elegans. Genetics. 2017;205:273-93.

10.   Lazetic V, Joseph BB, Bernazzani SM, Fay DS. Actin organization and endocytic trafficking are controlled by a network linking NIMA-related kinases to the CDC-42-SID-3/ACK1 pathway. PLoS Genet. 2018;14(4):e1007313. Epub 2018/04/03. doi: 10.1371/journal.pgen.1007313. PubMed PMID: 29608564; PubMed Central PMCID: PMCPMC5897031.

11.   Joseph BB, Wang Y, Edeen P, Lazetic V, Grant BD, Fay DS. Control of clathrin-mediated endocytosis by NIMA family kinases. PLoS Genet. 2020;16(2):e1008633. Epub 2020/02/19. doi: 10.1371/journal.pgen.1008633. PubMed PMID: 32069276.

12.   Binti S.: Melinda RVJ, B.B.; Edeen, P., Miller, S.D.; Fay, D.S. A life cycle alteration can
correct molting defects in Caenorhabditis elegans. bioRxiv.
2021;https://doi.org/10.1101/2021.11.02.466921. Epub November 4, 2021.
13.   Oakley BR, Morris NR. A mutation in Aspergillus nidulans that blocks the transition from
interphase to prophase. J Cell Biol. 1983;96(4):1155-8. Epub 1983/04/01. doi:
10.1083/jcb.96.4.1155. PubMed PMID: 6339527; PubMed Central PMCID: PMCPMC2112314.
14.   Osmani SA, May GS, Morris NR. Regulation of the mRNA levels of nimA, a gene required
for the G2-M transition in Aspergillus nidulans. J Cell Biol. 1987;104(6):1495-504. Epub
1987/06/01. doi: 10.1083/jcb.104.6.1495. PubMed PMID: 3294854; PubMed Central PMCID:
PMCPMC2114495.
15.   Casey JP, Brennan K, Scheidel N, McGettigan P, Lavin PT, Carter S, et al. Recessive NEK9
mutation causes a lethal skeletal dysplasia with evidence of cell cycle and ciliary defects. Hum
Mol Genet. 2016;25(9):1824-35. Epub 2016/02/26. doi: 10.1093/hmg/ddw054. PubMed PMID:
26908619.
16.   Fry AM, O'Regan L, Sabir SR, Bayliss R. Cell cycle regulation by the NEK family of protein
kinases. J Cell Sci. 2012;125(Pt 19):4423-33. Epub 2012/11/08. doi: jcs.111195 [pii]
10.1242/jcs.111195. PubMed PMID: 23132929; PubMed Central PMCID: PMC3500863.
17.   Moniz L, Dutt P, Haider N, Stambolic V. Nek family of kinases in cell cycle, checkpoint
control and cancer. Cell Div. 2011;6:18. Epub 2011/11/02. doi: 1747-1028-6-18 [pii]
10.1186/1747-1028-6-18. PubMed PMID: 22040655; PubMed Central PMCID: PMC3222597.
18.   Osmani AH, McGuire SL, Osmani SA. Parallel activation of the NIMA and p34cdc2 cell
cycle-regulated protein kinases is required to initiate mitosis in A. nidulans. Cell.
1991;67(2):283-91. Epub 1991/10/18. doi: 10.1016/0092-8674(91)90180-7. PubMed PMID:
1913824.
19.   Manning DK, Sergeev M, van Heesbeen RG, Wong MD, Oh JH, Liu Y, et al. Loss of the
ciliary kinase Nek8 causes left-right asymmetry defects. J Am Soc Nephrol. 2013;24(1):100-12.
Epub 2013/01/01. doi: 10.1681/ASN.2012050490. PubMed PMID: 23274954; PubMed Central
PMCID: PMCPMC3537214.
20.   Quarmby LM, Mahjoub MR. Caught Nek-ing: cilia and centrioles. J Cell Sci. 2005;118(Pt
22):5161-9. Epub 2005/11/11. doi: 118/22/5161 [pii]
10.1242/jcs.02681. PubMed PMID: 16280549.
21.   Trapp ML, Galtseva A, Manning DK, Beier DR, Rosenblum ND, Quarmby LM. Defects in
ciliary localization of Nek8 is associated with cystogenesis. Pediatr Nephrol. 2008;23(3):377-87.
Epub 2008/01/15. doi: 10.1007/s00467-007-0692-y. PubMed PMID: 18189147.
22.   Otto EA, Trapp ML, Schultheiss UT, Helou J, Quarmby LM, Hildebrandt F. NEK8
mutations affect ciliary and centrosomal localization and may cause nephronophthisis. J Am Soc
Nephrol. 2008;19(3):587-92. Epub 2008/01/18. doi: ASN.2007040490 [pii]
10.1681/ASN.2007040490. PubMed PMID: 18199800; PubMed Central PMCID: PMC2391043.
23.   Walker C, Mojares E, Del Rio Hernandez A. Role of Extracellular Matrix in Development
and Cancer Progression. Int J Mol Sci. 2018;19(10). Epub 2018/10/06. doi:
10.3390/ijms19103028. PubMed PMID: 30287763; PubMed Central PMCID: PMCPMC6213383.
24.   Wang R, Song Y, Xu X, Wu Q, Liu C. The expression of Nek7, FoxM1, and Plk1 in
gallbladder cancer and their relationships to clinicopathologic features and survival. Clin Transl

Oncol. 2013;15(8):626-32. Epub 2013/01/30. doi: 10.1007/s12094-012-0978-9. PubMed PMID:
23359173.
25.   Choudhury AD, Schinzel AC, Cotter MB, Lis RT, Labella K, Lock YJ, et al. Castration
Resistance in Prostate Cancer Is Mediated by the Kinase NEK6. Cancer Res. 2017;77(3):753-65.
Epub 2016/12/03. doi: 10.1158/0008-5472.CAN-16-0455. PubMed PMID: 27899381; PubMed
Central PMCID: PMCPMC5290202.
26.   He Z, Ni X, Xia L, Shao Z. Overexpression of NIMA-related kinase 6 (NEK6) contributes to
malignant growth and dismal prognosis in Human Breast Cancer. Pathol Res Pract.
2018;214(10):1648-54. Epub 2018/08/30. doi: 10.1016/j.prp.2018.07.030. PubMed PMID:
30153958.
27.   Jee HJ, Kim AJ, Song N, Kim HJ, Kim M, Koh H, et al. Nek6 overexpression antagonizes
p53-induced senescence in human cancer cells. Cell Cycle. 2010;9(23):4703-10. Epub
2010/11/26. doi: 14059 [pii]. PubMed PMID: 21099361.
28.   Jee HJ, Kim HJ, Kim AJ, Song N, Kim M, Lee HJ, et al. The inhibition of Nek6 function
sensitizes human cancer cells to premature senescence upon serum reduction or anticancer
drug treatment. Cancer Lett. 2013;335(1):175-82. Epub 2013/02/19. doi: S0304-3835(13)00124-
9 [pii]
10.1016/j.canlet.2013.02.012. PubMed PMID: 23416273.
29.   Joseph BB, Blouin NA, Fay DS. Use of a Sibling Subtraction Method for Identifying Causal
Mutations in Caenorhabditis elegans by Whole-Genome Sequencing. G3 (Bethesda, Md).
2018;8(2):669-78. Epub 2017/12/15. doi: 10.1534/g3.117.300135. PubMed PMID: 29237702;
PubMed Central PMCID: PMCPMC5919755.
30.   Seals DF, Courtneidge SA. The ADAMs family of metalloproteases: multidomain proteins
with multiple functions. Genes Dev. 2003;17(1):7-30. Epub 2003/01/07. doi:
10.1101/gad.1039703. PubMed PMID: 12514095.
31.   Giebeler N, Zigrino P. A Disintegrin and Metalloprotease (ADAM): Historical Overview of
Their Functions. Toxins (Basel). 2016;8(4):122. Epub 2016/04/28. doi: 10.3390/toxins8040122.
PubMed PMID: 27120619; PubMed Central PMCID: PMCPMC4848645.
32.   Alfandari D, McCusker C, Cousin H. ADAM function in embryogenesis. Semin Cell Dev
Biol. 2009;20(2):153-63. Epub 2008/10/22. doi: 10.1016/j.semcdb.2008.09.006. PubMed PMID:
18935966; PubMed Central PMCID: PMCPMC2693894.
33.   Cousin H, Abbruzzese G, Kerdavid E, Gaultier A, Alfandari D. Translocation of the
cytoplasmic domain of ADAM13 to the nucleus is essential for Calpain8-a expression and cranial
neural crest cell migration. Dev Cell. 2011;20(2):256-63. Epub 2011/02/15. doi:
10.1016/j.devcel.2010.12.009. PubMed PMID: 21316592; PubMed Central PMCID:
PMCPMC3074609.
34.   Brocker CN, Vasiliou V, Nebert DW. Evolutionary divergence and functions of the ADAM
and ADAMTS gene families. Hum Genomics. 2009;4(1):43-55. Epub 2009/12/03. doi:
10.1186/1479-7364-4-1-43. PubMed PMID: 19951893; PubMed Central PMCID:
PMCPMC3500187.
35.   Yagami-Hiromasa T, Sato T, Kurisaki T, Kamijo K, Nabeshima Y, Fujisawa-Sehara A. A
metalloprotease-disintegrin participating in myoblast fusion. Nature. 1995;377(6550):652-6.
Epub 1995/10/19. doi: 10.1038/377652a0. PubMed PMID: 7566181.

36.   Weskamp G, Cai H, Brodie TA, Higashyama S, Manova K, Ludwig T, et al. Mice lacking the
metalloprotease-disintegrin MDC9 (ADAM9) have no evident major abnormalities during
development or adult life. Mol Cell Biol. 2002;22(5):1537-44. Epub 2002/02/13. doi:
10.1128/mcb.22.5.1537-1544.2002. PubMed PMID: 11839819; PubMed Central PMCID:
PMCPMC134708.
37.   Sahin U, Weskamp G, Kelly K, Zhou HM, Higashiyama S, Peschon J, et al. Distinct roles
for ADAM10 and ADAM17 in ectodomain shedding of six EGFR ligands. J Cell Biol.
2004;164(5):769-79. Epub 2004/03/03. doi: 10.1083/jcb.200307137. PubMed PMID: 14993236;
PubMed Central PMCID: PMCPMC2172154.
38.   Primakoff P, Myles DG. The ADAM gene family: surface proteins with adhesion and
protease activity. Trends Genet. 2000;16(2):83-7. Epub 2000/02/01. doi: 10.1016/s0168-
9525(99)01926-5. PubMed PMID: 10652535.
39.   Mygind KJ, Storiko T, Freiberg ML, Samsoe-Petersen J, Schwarz J, Andersen OM, et al.
Sorting nexin 9 (SNX9) regulates levels of the transmembrane ADAM9 at the cell surface. J Biol
Chem. 2018;293(21):8077-88. Epub 2018/04/07. doi: 10.1074/jbc.RA117.001077. PubMed
PMID: 29622675; PubMed Central PMCID: PMCPMC5971451.
40.   Stautz D, Leyme A, Grandal MV, Albrechtsen R, van Deurs B, Wewer U, et al. Cell-surface
metalloprotease ADAM12 is internalized by a clathrin- and Grb2-dependent mechanism. Traffic.
2012;13(11):1532-46. Epub 2012/08/14. doi: 10.1111/j.1600-0854.2012.01405.x. PubMed
PMID: 22882974.
41.   Lorenzen I, Lokau J, Korpys Y, Oldefest M, Flynn CM, Kunzel U, et al. Control of ADAM17
activity by regulation of its cellular localisation. Sci Rep. 2016;6:35067. Epub 2016/10/13. doi:
10.1038/srep35067. PubMed PMID: 27731361; PubMed Central PMCID: PMCPMC5059621.
42.   Marcello E, Saraceno C, Musardo S, Vara H, de la Fuente AG, Pelucchi S, et al.
Endocytosis of synaptic ADAM10 in neuronal plasticity and Alzheimer's disease. J Clin Invest.
2013;123(6):2523-38. Epub 2013/05/17. doi: 10.1172/JCI65401. PubMed PMID: 23676497;
PubMed Central PMCID: PMCPMC3668814.
43.   Kim DW, Sung H, Shin D, Shen H, Ahnn J, Lee SK, et al. Differential physiological roles of
ESCRT complexes in Caenorhabditis elegans. Mol Cells. 2011;31(6):585-92. Epub 2011/06/21.
doi: 10.1007/s10059-011-1045-z. PubMed PMID: 21688204; PubMed Central PMCID:
PMCPMC3887624.
44.   Hurley JH. The ESCRT complexes. Crit Rev Biochem Mol Biol. 2010;45(6):463-87. Epub
2010/07/27. doi: 10.3109/10409238.2010.502516. PubMed PMID: 20653365; PubMed Central
PMCID: PMCPMC2988974.
45.   Henne WM, Buchkovich NJ, Emr SD. The ESCRT pathway. Dev Cell. 2011;21(1):77-91.
Epub 2011/07/19. doi: 10.1016/j.devcel.2011.05.015. PubMed PMID: 21763610.
46.   Babst M. MVB vesicle formation: ESCRT-dependent, ESCRT-independent and everything
in between. Curr Opin Cell Biol. 2011;23(4):452-7. Epub 2011/05/17. doi:
10.1016/j.ceb.2011.04.008. PubMed PMID: 21570275; PubMed Central PMCID:
PMCPMC3148405.
47.   Miao R, Li M, Zhang Q, Yang C, Wang X. An ECM-to-Nucleus Signaling Pathway Activates
Lysosomes for C. elegans Larval Development. Dev Cell. 2020;52(1):21-37 e5. Epub 2019/11/19.
doi: 10.1016/j.devcel.2019.10.020. PubMed PMID: 31735670.

48.   Kneen M, Farinas J, Li Y, Verkman AS. Green fluorescent protein as a noninvasive
intracellular pH indicator. Biophys J. 1998;74(3):1591-9. Epub 1998/03/25. doi: 10.1016/S0006-
3495(98)77870-1. PubMed PMID: 9512054; PubMed Central PMCID: PMCPMC1299504.
49.   Bode W, Gomis-Ruth FX, Stockler W. Astacins, serralysins, snake venom and matrix
metalloproteinases exhibit identical zinc-binding environments (HEXXHXXGXXH and Met-turn)
and topologies and should be grouped into a common family, the 'metzincins'. FEBS Lett.
1993;331(1-2):134-40. Epub 1993/09/27. doi: 10.1016/0014-5793(93)80312-i. PubMed PMID:
8405391.
50.   Maskos K, Fernandez-Catalan C, Huber R, Bourenkov GP, Bartunik H, Ellestad GA, et al.
Crystal structure of the catalytic domain of human tumor necrosis factor-alpha-converting
enzyme. Proc Natl Acad Sci U S A. 1998;95(7):3408-12. Epub 1998/05/09. doi:
10.1073/pnas.95.7.3408. PubMed PMID: 9520379; PubMed Central PMCID: PMCPMC19849.
51.   Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, et al. Highly accurate
protein structure prediction with AlphaFold. Nature. 2021;596(7873):583-9. Epub 2021/07/16.
doi: 10.1038/s41586-021-03819-2. PubMed PMID: 34265844; PubMed Central PMCID:
PMCPMC8371605.
52.   Varadi M, Anyango S, Deshpande M, Nair S, Natassia C, Yordanova G, et al. AlphaFold
Protein Structure Database: massively expanding the structural coverage of protein-sequence
space with high-accuracy models. Nucleic Acids Res. 2021. Epub 2021/11/19. doi:
10.1093/nar/gkab1061. PubMed PMID: 34791371.
53.   Song Y, DiMaio F, Wang RY, Kim D, Miles C, Brunette T, et al. High-resolution
comparative modeling with RosettaCM. Structure. 2013;21(10):1735-42. Epub 2013/09/17. doi:
10.1016/j.str.2013.08.005. PubMed PMID: 24035711; PubMed Central PMCID:
PMCPMC3811137.
54.   Raman S, Vernon R, Thompson J, Tyka M, Sadreyev R, Pei J, et al. Structure prediction
for CASP8 with all-atom refinement using Rosetta. Proteins. 2009;77 Suppl 9:89-99. Epub
2009/08/25. doi: 10.1002/prot.22540. PubMed PMID: 19701941; PubMed Central PMCID:
PMCPMC3688471.
55.   Glassey B, Civetta A. Positive selection at reproductive ADAM genes with potential
intercellular binding activity. Mol Biol Evol. 2004;21(5):851-9. Epub 2004/02/14. doi:
10.1093/molbev/msh080. PubMed PMID: 14963094.
56.   Black RA, Rauch CT, Kozlosky CJ, Peschon JJ, Slack JL, Wolfson MF, et al. A
metalloproteinase disintegrin that releases tumour-necrosis factor-alpha from cells. Nature.
1997;385(6618):729-33. Epub 1997/02/20. doi: 10.1038/385729a0. PubMed PMID: 9034190.
57.   Moss ML, Jin SL, Milla ME, Bickett DM, Burkhart W, Carter HL, et al. Cloning of a
disintegrin metalloproteinase that processes precursor tumour-necrosis factor-alpha. Nature.
1997;385(6618):733-6. Epub 1997/02/20. doi: 10.1038/385733a0. PubMed PMID: 9034191.
58.   Lunn CA, Fan X, Dalie B, Miller K, Zavodny PJ, Narula SK, et al. Purification of ADAM 10
from bovine spleen as a TNFalpha convertase. FEBS Lett. 1997;400(3):333-5. Epub 1997/01/06.
doi: 10.1016/s0014-5793(96)01410-x. PubMed PMID: 9009225.
59.   Rosendahl MS, Ko SC, Long DL, Brewer MT, Rosenzweig B, Hedl E, et al. Identification
and characterization of a pro-tumor necrosis factor-alpha-processing enzyme from the ADAM
family of zinc metalloproteases. J Biol Chem. 1997;272(39):24588-93. Epub 1997/09/26. doi:
10.1074/jbc.272.39.24588. PubMed PMID: 9305925.

60.   Reddy P, Slack JL, Davis R, Cerretti DP, Kozlosky CJ, Blanton RA, et al. Functional analysis
of the domain structure of tumor necrosis factor-alpha converting enzyme. J Biol Chem.
2000;275(19):14608-14. Epub 2000/05/09. doi: 10.1074/jbc.275.19.14608. PubMed PMID:
10799547.
61.   Solomon KA, Pesti N, Wu G, Newton RC. Cutting edge: a dominant negative form of TNF-
alpha converting enzyme inhibits proTNF and TNFRII secretion. J Immunol. 1999;163(8):4105-8.
Epub 1999/10/08. PubMed PMID: 10510344.
62.   Zhang Z, Oliver P, Lancaster JR, Jr., Schwarzenberger PO, Joshi MS, Cork J, et al. Reactive
oxygen species mediate tumor necrosis factor alpha-converting, enzyme-dependent
ectodomain shedding induced by phorbol myristate acetate. FASEB J. 2001;15(2):303-5. Epub
2001/02/07. doi: 10.1096/fj.00-0371fje. PubMed PMID: 11156944.
63.   Althoff K, Reddy P, Voltz N, Rose-John S, Mullberg J. Shedding of interleukin-6 receptor
and tumor necrosis factor alpha. Contribution of the stalk sequence to the cleavage pattern of
transmembrane proteins. Eur J Biochem. 2000;267(9):2624-31. Epub 2000/04/28. doi:
10.1046/j.1432-1327.2000.01278.x. PubMed PMID: 10785383.
64.   Fischer OM, Hart S, Gschwind A, Ullrich A. EGFR signal transactivation in cancer cells.
Biochem Soc Trans. 2003;31(Pt 6):1203-8. Epub 2003/12/04. doi: 10.1042/bst0311203. PubMed
PMID: 14641026.
65.   Prenzel N, Zwick E, Daub H, Leserer M, Abraham R, Wallasch C, et al. EGF receptor
transactivation by G-protein-coupled receptors requires metalloproteinase cleavage of proHB-
EGF. Nature. 1999;402(6764):884-8. Epub 2000/01/06. doi: 10.1038/47260. PubMed PMID:
10622253.
66.   Yan Y, Shirakabe K, Werb Z. The metalloprotease Kuzbanian (ADAM10) mediates the
transactivation of EGF receptor by G protein-coupled receptors. J Cell Biol. 2002;158(2):221-6.
Epub 2002/07/18. doi: 10.1083/jcb.200112026. PubMed PMID: 12119356; PubMed Central
PMCID: PMCPMC2173139.
67.   Lemjabbar H, Basbaum C. Platelet-activating factor receptor and ADAM10 mediate
responses to Staphylococcus aureus in epithelial cells. Nat Med. 2002;8(1):41-6. Epub
2002/01/12. doi: 10.1038/nm0102-41. PubMed PMID: 11786905.
68.   Ohtsu H, Dempsey PJ, Eguchi S. ADAMs as mediators of EGF receptor transactivation by
G protein-coupled receptors. Am J Physiol Cell Physiol. 2006;291(1):C1-10. Epub 2006/06/14.
doi: 10.1152/ajpcell.00620.2005. PubMed PMID: 16769815.
69.   Kataoka H. EGFR ligands and their signaling scissors, ADAMs, as new molecular targets
for anticancer treatments. J Dermatol Sci. 2009;56(3):148-53. Epub 2009/11/10. doi:
10.1016/j.jdermsci.2009.10.002. PubMed PMID: 19896805.
70.   Brou C, Logeat F, Gupta N, Bessia C, LeBail O, Doedens JR, et al. A novel proteolytic
cleavage involved in Notch signaling: the role of the disintegrin-metalloprotease TACE. Mol Cell.
2000;5(2):207-16. Epub 2000/07/06. doi: 10.1016/s1097-2765(00)80417-7. PubMed PMID:
10882063.
71.   Selvais C, Gaide Chevronnay HP, Lemoine P, Dedieu S, Henriet P, Courtoy PJ, et al.
Metalloproteinase-dependent shedding of low-density lipoprotein receptor-related protein-1
ectodomain decreases endocytic clearance of endometrial matrix metalloproteinase-2 and -9 at
menstruation. Endocrinology. 2009;150(8):3792-9. Epub 2009/05/02. doi: 10.1210/en.2009-
0015. PubMed PMID: 19406945.

72.   Liu Q, Zhang J, Tran H, Verbeek MM, Reiss K, Estus S, et al. LRP1 shedding in human
brain: roles of ADAM10 and ADAM17. Mol Neurodegener. 2009;4:17. Epub 2009/04/18. doi:
10.1186/1750-1326-4-17. PubMed PMID: 19371428; PubMed Central PMCID:
PMCPMC2672942.
73.   Selvais C, D'Auria L, Tyteca D, Perrot G, Lemoine P, Troeberg L, et al. Cell cholesterol
modulates metalloproteinase-dependent shedding of low-density lipoprotein receptor-related
protein-1 (LRP-1) and clearance function. FASEB J. 2011;25(8):2770-81. Epub 2011/04/27. doi:
10.1096/fj.10-169508. PubMed PMID: 21518850; PubMed Central PMCID: PMCPMC3470721.
74.   Emonard H, Theret L, Bennasroune AH, Dedieu S. Regulation of LRP-1 expression: make
the point. Pathol Biol (Paris). 2014;62(2):84-90. Epub 2014/03/26. doi:
10.1016/j.patbio.2014.02.002. PubMed PMID: 24661974.
75.   Dekky B, Wahart A, Sartelet H, Fere M, Angiboust JF, Dedieu S, et al. Cellular Cholesterol
Distribution Influences Proteolytic Release of the LRP-1 Ectodomain. Front Pharmacol.
2016;7:25. Epub 2016/02/24. doi: 10.3389/fphar.2016.00025. PubMed PMID: 26903870;
PubMed Central PMCID: PMCPMC4751253.
76.   Calligaris M, Cuffaro D, Bonelli S, Spano DP, Rossello A, Nuti E, et al. Strategies to Target
ADAM17 in Disease: From its Discovery to the iRhom Revolution. Molecules. 2021;26(4). Epub
2021/02/14. doi: 10.3390/molecules26040944. PubMed PMID: 33579029; PubMed Central
PMCID: PMCPMC7916773.
77.   Biemesderfer D. Regulated intramembrane proteolysis of megalin: linking urinary
protein and gene regulation in proximal tubule? Kidney Int. 2006;69(10):1717-21. Epub
2006/03/25. doi: 10.1038/sj.ki.5000298. PubMed PMID: 16557231.
78.   Zou Z, Chung B, Nguyen T, Mentone S, Thomson B, Biemesderfer D. Linking receptor-
mediated endocytosis and cell signaling: evidence for regulated intramembrane proteolysis of
megalin in proximal tubule. J Biol Chem. 2004;279(33):34302-10. Epub 2004/06/08. doi:
10.1074/jbc.M405608200. PubMed PMID: 15180987.
79.   May P, Reddy YK, Herz J. Proteolytic processing of low density lipoprotein receptor-
related protein mediates regulated release of its intracellular domain. J Biol Chem.
2002;277(21):18736-43. Epub 2002/03/22. doi: 10.1074/jbc.M201979200. PubMed PMID:
11907044.
80.   Spuch C, Ortolano S, Navarro C. LRP-1 and LRP-2 receptors function in the membrane
neuron. Trafficking mechanisms and proteolytic processing in Alzheimer's disease. Front
Physiol. 2012;3:269. Epub 2012/08/31. doi: 10.3389/fphys.2012.00269. PubMed PMID:
22934024; PubMed Central PMCID: PMCPMC3429044.
81.   Quinn KA, Pye VJ, Dai YP, Chesterman CN, Owensby DA. Characterization of the soluble
form of the low density lipoprotein receptor-related protein (LRP). Exp Cell Res.
1999;251(2):433-41. Epub 1999/09/02. doi: 10.1006/excr.1999.4590. PubMed PMID:
10471328.
82.   Stepek G, McCormack G, Birnie AJ, Page AP. The astacin metalloprotease moulting
enzyme NAS-36 is required for normal cuticle ecdysis in free-living and parasitic nematodes.
Parasitology. 2011;138(2):237-48. Epub 2010/08/31. doi: 10.1017/S0031182010001113.
PubMed PMID: 20800010.
83.   Frand AR, Russel S, Ruvkun G. Functional genomic analysis of C. elegans molting. PLoS
Biol. 2005;3(10):e312. Epub 2005/08/27. doi: 05-PLBI-RA-0162R2 [pii]

10.1371/journal.pbio.0030312. PubMed PMID: 16122351; PubMed Central PMCID:
PMC1233573.
84.   Suzuki M, Sagoh N, Iwasaki H, Inoue H, Takahashi K. Metalloproteases with EGF, CUB,
and thrombospondin-1 domains function in molting of Caenorhabditis elegans. Biol Chem.
2004;385(6):565-8. Epub 2004/07/17. doi: 10.1515/BC.2004.069. PubMed PMID: 15255192.
85.   Davis MW, Birnie AJ, Chan AC, Page AP, Jorgensen EM. A conserved metalloprotease
mediates ecdysis in Caenorhabditis elegans. Development. 2004;131(23):6001-8. Epub
2004/11/13. doi: 10.1242/dev.01454. PubMed PMID: 15539494.
86.   Hashmi S, Zhang J, Oksov Y, Lustigman S. The Caenorhabditis elegans cathepsin Z-like
cysteine protease, Ce-CPZ-1, has a multifunctional role during the worms' development. J Biol
Chem. 2004;279(7):6035-45. Epub 2003/11/25. doi: 10.1074/jbc.M312346200. PubMed PMID:
14630920.
87.   Kim TH, Kim YJ, Cho JW, Shim J. A novel zinc-carboxypeptidase SURO-1 regulates cuticle
formation and body morphogenesis in Caenorhabditis elegans. FEBS Lett. 2011;585(1):121-7.
Epub 2010/11/26. doi: 10.1016/j.febslet.2010.11.020. PubMed PMID: 21094156.
88.   Stepek G, McCormack G, Page AP. The kunitz domain protein BLI-5 plays a functionally
conserved role in cuticle formation in a diverse range of nematodes. Mol Biochem Parasitol.
2010;169(1):1-11. Epub 2009/09/01. doi: 10.1016/j.molbiopara.2009.08.005. PubMed PMID:
19716386.
89.   Page AP, McCormack G, Birnie AJ. Biosynthesis and enzymology of the Caenorhabditis
elegans cuticle: identification and characterization of a novel serine protease inhibitor. Int J
Parasitol. 2006;36(6):681-9. Epub 2006/02/28. doi: 10.1016/j.ijpara.2006.01.004. PubMed
PMID: 16500660.
90.   Hendriks GJ, Gaidatzis D, Aeschimann F, Grosshans H. Extensive oscillatory gene
expression during C. elegans larval development. Mol Cell. 2014;53(3):380-92. Epub
2014/01/21. doi: 10.1016/j.molcel.2013.12.013. PubMed PMID: 24440504.
91.   Meeuse MW, Hauser YP, Morales Moya LJ, Hendriks GJ, Eglinger J, Bogaarts G, et al.
Developmental function and state transitions of a gene expression oscillator in Caenorhabditis
elegans. Mol Syst Biol. 2020;16(7):e9498. Epub 2020/07/21. doi: 10.15252/msb.20209498.
PubMed PMID: 32687264; PubMed Central PMCID: PMCPMC7370751.
92.   Roberts B, Clucas C, Johnstone IL. Loss of SEC-23 in Caenorhabditis elegans causes
defects in oogenesis, morphogenesis, and extracellular matrix secretion. Mol Biol Cell.
2003;14(11):4414-26. Epub 2003/10/11. doi: 10.1091/mbc.e03-03-0162. PubMed PMID:
14551256; PubMed Central PMCID: PMCPMC266761.
93.   Stiernagle T. Maintenance of C. elegans. WormBook. 2006:1-11. Epub 2007/12/01. doi:
10.1895/wormbook.1.101.1. PubMed PMID: 18050451; PubMed Central PMCID:
PMCPMC4781397.
94.   Fire A, Xu S, Montgomery MK, Kostas SA, Driver SE, Mello CC. Potent and specific
genetic interference by double-stranded RNA in Caenorhabditis elegans. Nature.
1998;391(6669):806-11. Epub 1998/03/05. doi: 10.1038/35888. PubMed PMID: 9486653.
95.   Ahringer J. Reverse Genetics, WormBook. 2005.
96.   Wang D, Kennedy S, Conte D, Jr., Kim JK, Gabel HW, Kamath RS, et al. Somatic
misexpression of germline P granules and enhanced RNA interference in retinoblastoma

pathway mutants. Nature. 2005;436(7050):593-7. Epub 2005/07/29. doi: 10.1038/nature04010.
PubMed PMID: 16049496.
97.   Fay SF, Fay DS, Chhatre VE. CRISPRcruncher: A tool for engineering restriction sites into
coding regions. MicroPubl Biol. 2021;2021. Epub 20210118. doi:
10.17912/micropub.biology.000343. PubMed PMID: 33490886; PubMed Central PMCID:
PMCPMC7816087.
98.   Fay DS, Gerow K. A biologist's guide to statistical thinking and analysis. WormBook.
2013:1-54. Epub 2013/08/03. doi: 10.1895/wormbook.1.159.1. PubMed PMID: 23908055;
PubMed Central PMCID: PMC3880567.
1124