Short Title:

Regulation of photosynthetic control

Title:

The algal PETC-Pro171-Leu suppresses electron transfer in the cytochrome b_6f complex under acidic

lumenal condition

Authors:

Shin-Ichiro Ozawa[a], Felix Buchert[b], Ruby Reuys[b], Michael Hippler[a, b], and Yuichiro Takahashi[c]

Author affiliations:

[a]Institute of Plant Science and Resources, Okayama University, Kurashiki 710-0046, Japan

[b]Institute of Plant Biology and Biotechnology, University of Münster, 48143 Münster, Germany

[c]Research Institute for Interdisciplinary Science, Okayama University, 700-8530, Japan

One sentence summary:

Change of PETC to P171L via site directed mutagenesis alters the pH dependency of the photosynthetic

control mechanism

Author Contributions:

S.O. and F.B. are co-first authors; S.O., F.B., M.H., and Y.T. conceived the original screening and

research plans; S.O., F.B., M.H., and Y.T. supervised the experiments; S.O., F.B., and R.R. designed

and performed the experiments; S.O., F.B., and R.R. analyzed the data; S.O., F.B., M.H., and Y.T.

conceived the project and wrote the article with contributions of all the authors; S.O., F.B., M.H., and

Y.T. supervised and completed the writing. S.O. agrees to serve as the author responsible for contact

and ensures communication.

Funding:

This work was supported by the Grants-in-Aid for Scientific Research from Japan Society for the

Promotion of Science KAKENHI JP21K06217 (to S.O.) and the JP16H06554 and 21H02510 (to Y.T.).

F.B. acknowledges funding from the Deutsche Forschungsgemeinschaft (BU 3426/3-1). M.H. also

acknowledges funding from the Deutsche Forschungsgemeinschaft (HI 739/13-2) and by the RECTOR

program (Okayama University, Japan).

Responsibilities of the Author for Contact:

Shin-Ichiro Ozawa

Institute of Plant Science and Resources, Okayama University, Okayama 710-0046, Japan

Phone: +81-86-434-1225, E-mail: OzwSh1r@okayama-u.ac.jp

Keywords:

photosynthesis, Q-cycle, cytochrome b_6f, photosynthetic control

Abstract

Linear photosynthetic electron flow (LEF) produces NADPH and generates a proton electrochemical

potential gradient across the thylakoid membrane used to synthesize ATP, both of which are required

for CO_2 fixation. As cellular demand for ATP and NADPH are variable, cyclic electron flow (CEF)

between PSI and cytochrome b_6f complex (b_6f) produces extra ATP. The b_6f regulates LEF and CEF via

photosynthetic control, which is a pH-dependent b_6f slowdown of plastoquinol oxidation at the lumenal

site. This protection mechanism is triggered at more alkaline lumen pH in the *pgr1* mutant of the vascular

plant *Arabidopsis thaliana*, carrying Pro194Leu in the b_6f Rieske Iron-sulfur protein. In this work, we

introduced *pgr1* mutation in the green alga *Chlamydomonas reinhardtii* (*PETC-P171L*). Consistent with

pgr1 phenotype, *PETC-P171L* displayed an impaired NPQ induction along with slower

photoautotrophic growth under high light conditions. Our data provides evidence that the ΔpH

component in *PETC-P171L* is dependent on oxygen availability. Only under low oxygen conditions the

ΔpH component was sufficient to trigger a phenotype in algal *PETC-P171L* where the mutant b_6f was

more restricted to oxidize the PQ pool and showed a diminished electron flow through the b_6f complex.

1 Introduction

Oxygenic photosynthesis fixes carbon dioxide and generates oxygen as a result of intra- and

intermolecular photochemical electron transfer reactions. The electron transfer pathways in oxygenic

photosynthesis either follows Linear Electron Flow (LEF) or, as an example of alternative passages,

Cyclic Electron Flow (CEF). Electrons stemming from water oxidation reduce $NADP^+$ during LEF, thus

providing the reducing power for CO_2 fixation in the Calvin Benson cycle. In the CEF, electrons from

reduced ferredoxin (Fd) are differently partitioned and redirected into the plastoquinone (PQ) pool. A

possible CEF route reinjects electrons into the cytochrome b_6f complex (b_6f) from the stromal side. In

LEF and CEF, the b_6f contributes to proton gradient formation across thylakoid membrane by

translocating protons from chloroplast stroma to thylakoid lumen. Light-driven charge separation, water

splitting as well as b_6f activity generate a membrane potential ($\Delta\Psi$) and a proton gradient (ΔpH),

respectively, and this proton motive force (*pmf*) drives ATP synthesis. During prolonged electron

transfer, the lumen will get further acidified, which in turn will slow down plastoquinol (PQH_2)

oxidation at the Qo-site of b_6f, and thereby establish photosynthetic control to protect photosystem I

(PSI) from acceptor side limitation (Rumberg and Siggel, 1969; Tikhonov et al., 1981; Nishio and

Whitmarsh, 1993). Lumen acidification is also important to induce Non-Photochemical Quenching

(NPQ) (Kanazawa and Kramer, 2002), which is the most rapid photoprotective response to excess light.

The fastest constituent of NPQ is qE, which operates at a time scale of seconds to minutes and regulates

the thermal dissipation of excess absorbed light energy, thereby providing effective photo-protection of

photosystem II (PSII). In vascular plants, the PSII protein PSBS is essential for qE (Li et al., 2002)

whereas qE induction in the green alga *Chlamydomonas reinhardtii* is mediated by LHCSR3, an ancient

light-harvesting protein that is missing in vascular plants (Peers et al., 2009).

Electron transfer through the b_6f is governed by the light-induced introduction of an electron hole

(positive charge) into the high-potential chain. The latter involves redox centers heme *f* and the [2Fe-

2S] type Rieske Iron Sulfur Cluster (Rieske ISC). The electron hole depends on photo-oxidized

plastocyanin and is successively passed from heme *f* via Rieske ISC to PQH_2. The Q-cycle within the

b_6f is responsible to generate a *pmf* across the thylakoid membrane, owing to a sophisticated electron

bifurcation. Two electrons are pulled from PQH_2 at the Q_o site, forming PQ within the membrane while

simultaneously releasing two H^+ into the lumen. During the PQH_2 oxidation at the Q_o site, the first

electron is transferred into the high-potential chain (Rieske ISC reducing heme f), and the second

electron into the low-potential chain (b_L heme reducing b_H heme, terminal acceptor is c_i heme). A second

PQH_2 oxidation at the Q_o site forms a reduced form of b_H / c_i heme pair which reduces a PQ bound at Q_i

site to complete the Q-cycle (note that alternative annotations for Q_o, Q_i, b_H and c_i can be found in

literatures). It was suggested that under LEF and CEF conditions two types of Q-cycle mechanism

operate in the b_6f with specific lumen acidification efficiencies per electron transferred via the high-

potential chain (Buchert et al., 2020).

The molecular structure of b_6f complex was first resolved in the green alga *Chlamydomonas reinhardtii*

and the cyanobacteria *Mastigocladus laminosus* independently (Kurisu et al., 2003; Stroebel et al., 2003).

The b_6f functions as a homodimer, and each monomer consists of four core subunits (PetA, PetB, PETC,

and PetD) and four stably associated small subunits (PetG, PetL, PETM, and PETN) to stabilize the

complex (Figure 1A). The cofactors and quinone binding pockets involved in electron transfer are found

in the b_6f core subunits which resemble the cytochrome bc_1 complex of the respiratory chain. The

structure-function similarities suggest that the hydrophilic domain of PETC subunit would move toward

PetA subunit in transferring the electrons from Rieske ISC to heme f.

In *Arabidopsis thaliana*, various photosynthetic mutants have been isolated and characterized by

forward genetics. One study employed ethyl methanesulfonate as mutagen to isolate the CE10-10-1

mutant which exhibited reduced NPQ with normal PSI/PSII quantum yields (Shikanai et al., 1999). A

follow-up study pinpointed the mutation as a proline-to-leucine substitution at position 194 of

Arabidopsis PETC, linking the NPQ defect to insufficient ΔpH formation in the light (Munekage et al.,

2001). The Arabidopsis PETC-P194L mutation was dubbed *pgr1* (*proton gradient regulation* 1).

It is known that green algal photosystems are protected from excess light illumination differently than

plants. For instance, algal state transitions redistribute excitation energy more effectively than in plants

(Delepelaire and Wollman, 1985). Moreover, flavodiiron protein was found in algae to protect PSI

acceptor side limitation induced by fluctuating light illumination (Jokel et al., 2018). Induction of qE

relies on LHCSR3, which requires a preceding stress period to trigger light to heat dissipation upon

lumen acidification (Peers et al., 2009; Bonente et al., 2011). Moreover, the algal PETO protein interacts

with b_6f and functions as a CEF effector under anoxic condition, which might point to a different CEF

regulation (Takahashi et al., 2016; Buchert et al., 2018).

Herein, we introduced the Arabidopsis *pgr1* mutation into *Chlamydomonas reinhardtii* (*PETC-P171L*)

to analyze the effects of the mutation on Photosynthetic Control in the green alga. The *PETC-P171L*

cytochrome b_6f showed a normal accumulation and electron flow activity in cells grown under oxic

conditions. However, the *PETC-P171L* displayed an impaired photoautotrophic growth and was less

competent in NPQ induction. Unlike Arabidopsis *pgr1*, the *PETC-P171L* phenotype was conditional

and revealed upon cellular stress treatment. In line, the electron transfer activity of the mutated b_6f was

significantly decreased in cells grown under anoxic conditions. Our experiments also revealed that the

ΔpH component in *PETC-P171L* is dependent on oxygen availability under light conditions that are

permissive under other circumstances. Under low oxygen condition the mutant b_6f was more restricted

to oxidize the PQ pool and limited electron flow.

Materials and methods

Strains and growth condition

The cells of *Chlamydomonas reinhardtii* were grown in Tris-acetate phosphate (TAP), Tris-Phosphate

(TP) medium, or minimum medium (MM) (Harris, 1988) under 50-100 μmol photons·m^{-2}·s^{-1} or 600

μmol photons·m^{-2}·s^{-1}. The previously isolated PETC gene disrupted strain *petC-Δ1* (mating type minus)

described in (de Vitry et al., 1999) was crossed with cc-125 (mating type plus) and the resultant PETC

deficient progeny was used as a recipient strain for transformation. Doubling time of the cell growth is

calculated by dividing 0.30 by specific growth rate obtained as the slope which is computed by linear

approximation for logarithmic growth curve in common logarithmic axis. Linear fitting was performed

at the highest coefficient of determination (R-square) with no restrictions by using built-in linear fitting

program of the Origin Pro 2021b version 9.8.5.212 (OriginLab corporation, USA). Refer the Appendix

1 for derivation of the formula to calculate the doubling time with specific growth rate. For *in vivo*

spectrophotometric analysis, cells were maintained at 20 μmol photons·m^{-2}·s^{-1} on agar-supplemented

TAP plates (Sueoka, 1960). When growing cells for experiments, liquid TP medium was used. The

cultures were grown at 40 μmol photons·m^{-2}·s^{-1} (16 h light/8 h dark) and were bubbled with sterile air

at 25°C while shaking at 120 rpm. About 300×10^6 cells were incubated in 400-mL TP in a FMT 150

photobioreactor vessel (Photon Systems Instruments, Czech Republic). Photobioreactor experiments

were carried out in the absence of air supply and comprised 3 days of 16-h photoperiods (sigmoidal light

and temperature gradients of 0-250 μmol photons·m^{-2}·s^{-1} from white-red LEDs and 18-28°C,

respectively, with 18°C constant in the dark).

Generation of transformation vectors

The BamHI fragment containing *PETC* gene (Cre11.g467689.t1.1, Chlamydomonas version 5.5) in the

pACR4.5 (de Vitry et al., 1999) was digested with BamHI and subsequently the resulting 4153 bp

fragment containing the *PETC* gene and 528 bp upstream of 5' -UTR of *PETC* was recloned into pBSSK$^-$

vector at BamHI site to yield pgcC plasmid. The mutation Pro171Leu in PETC was introduced by In-

Fusion^R (Clontech, TAKARA, Japan) with the DNA fragment between BbvCI and SnaBI sites which

was generated by PCR with If-BbvCI_P171L_gcC_F and If-SnaBI_gcC_R primers. The sequence of

this study's oligos is shown in Table 1. The DNA sequence of the inserted PCR product was validated by DNA sequencing with the primers used to generate the PCR product and we obtained pgcC-P171L plasmid.

We also prepared the DNA construct in which the *PETC* gene expression is controlled by Chlamydomonas *PSAD* promoter and UTRs. We amplified spliced *PETC* coding DNA sequence fragment by nested PCR with cDNA library as a template DNA. The 1st round of PCR from cDNA library with primers HK#1483 and HK#1484 generated a 731 bp DNA fragment. The following 2nd round of PCR from the 1000 times diluted 1st round of PCR product with primers HK#1485 and HK#1486. The resulting DNA fragment of the PCR was ligated into the pSL18 vector (Depège et al., 2003) digested with NdeI and XbaI. The cloned DNA sequence of spliced *PETC* coding DNA sequence was validated by DNA sequencing with HK#1491 and HK#1492 primers. To introduce Pro171Leu in PETC, the *PETC* coding DNA sequence was replaced by In-Fusion HD cloning kit at NdeI and XbaI site with the PCR product consisting of two overlapping DNA fragments generated by the following primers; one fragment is generated with If-paDcC-NdeI_F and ccC_mP171L_R while the other fragment is generated with ccC_mP171L_F and If-paDcC-XbaI_R. The replaced DNA sequence was validated by DNA sequencing with the primers used to generate PCR product and cC_cdn131-137_F and cC_cdn137-131_R, and we obtained pccC-P171L plasmid.

Nuclear gene transformation

Nuclear gene transformation was performed with NEPA21 (Nepagene, Japan) according to the previous research (Yamano et al., 2013). The plasmid DNA was linearized by digesting with ScaI and the digested DNA was delivered into the Δ*PETC*. Selection of transformants was performed on MM solid medium under continuous illumination (50-100 µmol photons·m^{-2}·s^{-1}).

SDS-PAGE

The whole cellular proteins were denatured by heating at 100°C for 2 minutes in the presence of SDS and DTT and the polypeptides were separated by SDS-PAGE. The polypeptides were transferred on nitrocellulose membrane for immunodetection. The signals were visualized with a CCD camera

1   equipped imagers (Amersham Imager, LAS-4000, GE healthcare, or Chemidoc, Biorad) and the signal

2   intensities were quantified with Multigauge ver. 3.0 (FujiFilm, Japan)

Low temperature fluorescence emission spectra

Cells were resuspended in TP medium at 2 µg Chl·mL^{-1} and incubated for overnight under illumination

at 300 µmol photons·m^{-2}·s^{-1} prior to measurements. After the incubation, culture was directly, state 1

inducible treatment, or state 2 inducible treatment at 25°C. For the case of state 1 inducible treatment,

cells were incubated under 300 µmol photons·m^{-2}·s^{-1} in the presence of 10 µM of 3-(3,4-

dichlorophenyl)-1,1-dimethylurea (DCMU). For the case of state 2 inducible treatment, cells were

incubated in the darkness in the presence of 100 mM glucose and 50 U·mL^{-1} of glucose oxidase. A

sample in a NMR tube was in liquid nitrogen, excited with 430 nm illumination, and emission spectrum

was recorded from 400 to 800 nm with FL-7000 (Hitachi, Japan).

In vivo **Spectroscopy**

TP-grown cells with sterile air supply were harvested after ~2-h in the light cycle. Grown cultures were

diluted ~6-fold at least once after inoculation and grown to a density of ~1×10^6 cells·mL^{-1} before

harvesting (5000 rpm, 5 min, 25°C). Cells were resuspended at 20 µg Chlorophyll·mL^{-1} in TP

supplemented with 20% (*w/v*) Ficoll. Oxic samples were resuspended throughout the measurements in

2 mL open cuvettes. For oxygen-deprived conditions, cells were supplemented with 50 mM glucose, 10

U glucose oxidase (Carl Roth, Germany) and 30 U catalase (Sigma-Aldrich, USA) in the cuvette, and

then overlaid with mineral oil for 20 min in the dark. The enzyme stock solutions to obtain anoxic

samples were freshly prepared. Cell incubation in the presence of 10 µM nigericin (from 10 mM

ethanolic stock) was carried out for 20 min in the light before the measurements. All samples were light-

adapted in the cuvette for 20 min before the measurements using LEDs emitting ~150 µmol photons·m^{-}

^{2}·s^{-1} of 630 nm actinic light (AL). Measurements of the electrochromic shift (ECS) and P700 were

obtained using a Joliot-type spectrophotometer (JTS-10, Biologic France in Figure 3; otherwise JTS-

150, Spectrologix USA) and are described in detail elsewhere (Buchert et al., 2020; Buchert et al., 2022).

In brief, all absorption changes were normalized to the ECS ΔI/I signals (520 nm – 546 nm) produced

after a saturating 6-ns laser flash in the presence of 1 mM hydroxylamine (HA) and 10 μM DCMU, which relate to the density of active PSI centers in the sample. Thus, the unit of 1 charge separation·PSI[-1] is produced when measuring the flash-induced ECS amplitude (rapidly established before the first detection at ~300-μs). This optical ruler was determined in separate measurements to compare sample activities regardless of the presence of HA and DCMU. The shown flash-induced ECS kinetics were obtained after lowering the flash intensity to 30-45% saturation to avoid double turnovers of the cytochrome b_6f complex. The flash-induced ECS kinetics are multiphasic and the b-phase, which is a ~10-ms ECS rise associated with charge separation activity via the b_6f, was deconvoluted to obtain the shown half-times. Deconvolution from the raw ECS kinetics involved subtraction of rate constants of a two-exponential decay function to fit the c-phase, which is associated with ATP synthase activity. The deconvoluted b-phase was quantified with a mono-exponential function (Buchert et al., 2020). The ECS was also deconvoluted in the dark pulse method (Joliot and Joliot, 2002; Nawrocki et al., 2019). Accordingly, AL-adapted cells experienced 30 s darkness and were re-illuminated with AL (Supplemental Figure 1). During this 10 s AL induction phase, activities of stable charge separations·s[-1]·PSI[-1] were measured by brief light shuttering at discrete time points (four technical replicates with 1 min AL in between; data assembly from two independent measurement sequences). For AL interruptions at $t = 0$ ms, the linear ECS slopes before (S_L from –4 ms to –1 ms) and after darkness (S_D from +1 ms to +5 ms) gave electron transfer rates before darkness ($S_L - S_D$). Rates at 1 ms illumination of light-adapted cells were calculated from separate slopes obtained during 2 ms AL after 30 s dark (15 replicates spaced by 5 s). P700 measurements (705 nm – 740 nm) are based on the method by (Klughammer and Schreiber, 1994), measuring AL-adapted cells. As shown in Supplemental Figure 2, the fractions of photooxidizable (by a 22 ms saturating pulse), pre-oxidized in AL and non-photooxidizable P700 were determined by referring to the fully oxidized P700 signals in PSII-inhibited HA/DCMU samples (Buchert et al., 2020; Buchert et al., 2022). Since the above-mentioned ECS slopes obtained during 2 ms AL are a function of the P700 oxidation rate in PSII-inhibited HA/DCMU samples, the obtained YI parameter in those samples served to calculate CEF (by multiplication with the P700 oxidation rate) as it a function of the concentration of reduced P700 in AL (Takahashi et al., 2013). Single-turnover measurements of the cytochrome b_6f complex were measured after a sub-saturating flash,

hitting 40-50% of PSI reaction centers (Buchert et al., 2020; Buchert et al., 2022). Using interference

filters, the kinetics of flash-induced cytochrome-*f* reduction (measured with 554-nm detection pulses)

was calculated using a baseline drawn between 546-nm and 573-nm after correcting for the spectral

overlap with the *b*-hemes by subtracting $0.27 \times$ (563-nm − 546-nm).

Chlorophyll fluorescence measurement

Cells were grown in liquid TP medium to $\sim 1 \times 10^6$ cells·mL^{-1} with gentle shaking at 25°C under

continuous light (50 μmol photons·m^{-2}·s^{-1}). The cells were harvested by centrifugation at 4000 rpm,

suspended in fresh liquid TP medium at 5 μg Chl·mL^{-1}, and placed in darkness for 30 min (growth light

control samples). Photosynthesis driven by 450 nm LEDs was measured with a Maxi-Imaging PAM

chlorophyll fluorometer (Heinz Walz, Germany). The effective photosynthetic active radiation in the

measuring chamber was determined with a Quantitherm PAR/Temp Sensor (Hansatech, England). Cell

aliquots were also incubated shaking for 4 h under high light illumination (500 μmol photons·m^{-2}·s^{-1}).

The maximal PSII quantum yield in dark-adapted samples and the PSII quantum yield during

illumination Y(II) were determined as (Fm−Fo)/Fm and (Fm'−F)/Fm', respectively. Fo and Fm of dark-

adapted samples are respective minimal (in the presence of detection light) and maximal fluorescence

yields (due to closed PSII centers after saturating pulse). F and Fm' are the fluorescence yields in light-

adapted cells obtained before and after a saturating pulse, respectively. The nonphotochemical

quenching NPQ = (Fm−Fm')/Fm' was also determined, as well as estimation of the plastoquinone pool

redox state using the (1−qL) parameter (Kramer et al., 2004). The latter required Fo' measurements

(minimal fluorescence in illuminated samples) and qL was calculated as ((Fm'−F)×Fo')/((Fm'−Fo')×F).

These parameters were either measured after 30 s during light curve experiments or at distinct time

points when a 10 min continuous illumination at 95 μmol photons·m^{-2}·s^{-1} was applied. Fluorescence

data in FMT 150 photobioreactor experiments was obtained from blue excitation lights, and Y(II) was

calculated as above using saturating pulses every 15 min. The NPQ values were obtained from

calculations with OD$_{680}$-corrected Fm and Fm' whereas for each 16-h photoperiod the respective Fm

was averaged over a 2-h dark period before the onset of light.

Results

The PETC protein consists of one transmembrane helix and a lumenal hydrophilic domain in which the

Rieske iron sulfur cluster (ISC) is present, and both domains are connected through a flexible hinge.

Based on the structural similarities with the Rieske subunit in bc_1 complex of respiratory chain,

photosynthetic electron transfer between Rieske ISC and heme f in PetA involves reversible movements

of the lumenal hydrophilic domain of PETC. The PETC amino acid sequence alignment shows that

chloroplast targeting transit peptide of *A. thaliana* is 23 amino acids longer than that of *C. reinhardtii*

while the mature protein sequence is highly conserved (Supplemental Figure 3). The four amino acids

for ligation of two iron atoms and a flexible hinge region are conserved. Moreover, the mutated proline

residue in Arabidopsis *pgr1* (Pro194) is conserved as Pro171 in *C. reinhardtii*. The location of this

proline is close to Rieske ISC and is on the surface of hydrophilic domain (Figure 1B, C, and D), and

its proximity to the Rieske ISC is conserved in the b_6f structures (Supplemental Figure 4).

Generation of *PETC-P171L* mutant in the green alga *Chlamydomonas reinhardtii*

The Arabidopsis *pgr1* mutant carrying a single amino acid alteration mutation in PETC (P194L) exhibits

a hypersensitive photosynthetic control and downregulates the electron transfer flow under strong light

illumination more strongly than wild type (Munekage et al., 2001). To characterize the corresponding

PETC mutation in the green alga *C. reinhardtii*, we generated a *PETC-P171L* strain in which proline

171 is substituted by leucine. In our complementation-based screening approach, *petC-Δ1* was selected

as a recipient strain since it has a frameshift upon TG base deletions within the 36th codon and does not

accumulate *PETC* transcripts, thus showing no spontaneous reversion (de Vitry et al., 1999). Since the

petC-Δ1 exhibited a reduced accumulation of PSI protein, we backcrossed the *petC-Δ1* to wild type

strain (cc-125) to assess whether the mutation of PETC gene affects the accumulation of PSI proteins.

We obtained several tetrads and analyzed the accumulation of PsaA and PETC by immunoblotting

(Supplemental Figure 5). The T_{15-3} progeny accumulates no PETC but PsaA normally and is referred to

as the ΔPETC for the subsequent transformation since the two phenotypes segregated. As described in

Materials and Methods, we generated two types of vectors for transforming *PETC* gene; one consists of

genomic DNA fragment around PETC gene and the other contains PETC cDNA fused with *PSAD*

promoter and UTRs. We delivered each transforming vector into the ΔPETC cells by electroporation

and obtained numerous colonies after screening under photoautotrophic condition (on minimum solid

medium in the light of 50 μmol photons·m^{-2}·s^{-1}). The resulting transformants recovered the accumulation

of PETC to wild-type level, indicating that both vectors efficiently complemented the ΔPETC phenotype

(Figure 2). Subsequently, we transformed ΔPETC with vectors containing the mutation PETC-P171L

and obtained complemented strains expressing the mutated PETC. Immunoblot analyses revealed that

the complemented PETC-P171L cells accumulated PETC as well as PetA at wild-type levels, suggesting

that the mutation does not affect the stability of b_6f complex.

The *PETC-P171L* accumulates more oxidizable P700 and displays slower onset of electron

transfer in anaerobic algae which relied on the lumen pH under these conditions.

To examine the *PETC-P171L* mutant *in vivo* we chose a moderate light intensity for the following

experiments (~160 μmol photons·m^{-2}·s^{-1} actinic red light). Moreover, *pmf* partitioning was modulated

on a metabolic level in light-adapted cells by lowering oxygen availability. A similar situation may

occur in the natural habitat of Chlamydomonas which shares it with respiring microbes. Accordingly,

the alga faces microoxic environments in which photosynthesis needs to be fine-tuned, which is

consistent with its versatile anoxic metabolism (Grossman et al., 2011). Figure 3 shows a spectroscopic

in vivo analysis of control and the *PETC-P171L* in oxic conditions and upon anaerobic treatment.

Mitochondrial congestion in anoxic samples coincides with a redox poise shown to trigger CEF (Alric,

2014) and a *pmf* re-partitioning towards a more pronounced ΔpH (Finazzi and Rappaport, 1998).

Figures 3A and 3B display the redox state of P700, the primary electron donor of PSI (see Materials and

Methods and Supplemental Figure 2 for representative kinetics). Under oxic conditions when PSII was

active, the P700 redox states were indistinguishable between control and *PETC-P171L* (Figure 3A).

About 20% of P700 were oxidized during actinic background light which is ascribed to similar donor

side limitation (YND) in both strains. P700 was able to be further photooxidized by a short saturating

light pulse, as shown by the yield of PSI (YI). A similar fraction (about 40%) of P700 was not oxidizable

due to acceptor side limitation (YNA). On the contrary, YND was remarkably large when PSII was

inhibited (HA+DCMU, Figure 3A) because electrons from PSII were suppressed. This coincided with

YI of ~0.2 in control, i.e., 20% of P700 remained reduced in the light-adapted state in absence of PSII activity and were only oxidized by saturating pulses. This finding suggests that this 20% of P700 is reduced by electrons through CEF. YI in controls was roughly twice as large as in *PETC-P171L*, which translates in lower CEF rates in the mutant (indicated as 12 electrons·s^{-1}·PSI^{-1} in parenthesis of Figure 3A).

In anoxic conditions the *PETC-P171L* established a larger YND and a lower YNA compared to control when PSII was active (Figure 3B). In the absence of PSII activity, control and *P171L* showed similar P700 parameters. This finding shows that CEF rates in the steady state were unaffected in both strains (shown as electrons·s^{-1}·PSI^{-1} in parenthesis of Figure 3B). Overall, we observed subtle differences on the level of light-adapted steady-state of P700. The differences were produced in the absence of PSII activity under oxic conditions (Figure 3A) and in the presence of PSII activity under anoxic conditions (Figure 3B). We concluded that the PSI donor side of the *PETC-P171L* is specifically limited in anoxic condition.

Next, we re-illuminated cells after a 30-s dark period. As mentioned above, a significant ΔpH is generated when algae are kept in the dark under anoxic conditions (Finazzi and Rappaport, 1998), which supposedly slows down the Qo-site activity in *PETC-P171L*. Panels C – E in Figure 3 tested this assumption. The electrochromic shift (ECS) of photosynthetic pigments allows to measure the membrane potential as the ECS signals linearly follow the electric *pmf* component (Junge and Witt, 1968; Witt, 1979). More specifically (see Materials and Methods), we measured electron transfer transients during the first 10-s of light in oxic (Figure 3C) and anoxic conditions (Figure 3D). The induction kinetics of total electron transfer rates were similar in both strains under oxic conditions (Figure 3C). The sum of LEF and CEF activities in the control strain was slightly higher at 4-s and 10-s of light (filled asterisks, Figure 3C). The kinetics were almost identical when PSII was inhibited, except for 100-ms of light where *PETC-P171L* produced slightly higher rates (open asterisk, Figure 3C). The induction of total electron transfer under anoxic conditions showed that activities were transiently less efficient in *PETC-P171L*, producing the largest differences within the first 25 ms of light (cf. filled symbols and asterisk in Figure 3D). When PSII was inhibited (open symbols Figure 3D), no differences were observed between both strains.

We noticed that light-adapted, anoxic cells produced different ECS kinetics when a sub-saturating flash was fired after 30-s darkness (Figure 3E). The measurements resolved the averaged b- and c-phases of the flash-induced ECS kinetics (reviewed in Bailleul et al., 2010). The non-resolved a-phase with an amplitude of ~ 0.6 charge separations$\cdot$PSI^{-1} relied on the charge separation activity of both photosystems. The b-phase, which is a ~ 10-ms ECS rise, involves charge separation activity via the b_6f and the c-phase depends on ATP synthase activity. The latter phase overlapped the b-phase since ATP synthase activity was driven by the pmf provided from photosystem activity. Under oxic conditions, the fast c-phase prevented the appearance of a b-phase (Supplemental Figure 6). Therefore, the detection of an apparent b-phase requires a slow c-phase in anoxic algae (Velthuys, 1978). The ECS kinetics in anoxic *PETC-P171L* displayed a flat b-phase and comparably slow c-phase kinetics (Figure 3E), suggesting a compromised reaction of b_6f electron transfer activity in *PETC-P171L*.

This observation prompted us to measure cytochrome-f reduction kinetics in single-turnover measurements. In line with the conditional phenotype of *PETC-P171L*, we did not observe a difference in oxic cells (Supplemental Figure 7). When cells were treated under anoxic conditions, we observed that the half-time of cytochrome-f reduction in the control strain was significantly faster than in *PETC-P171L* (43 ± 23 ms vs. 74 ± 8 ms, Figure 4A). To test whether this difference is linked to the ΔpH generated by anoxic algae in the dark (Finazzi and Rappaport, 1998), we took advantage of nigericin. The H^+/K^+ ionophore exchanges the osmotic pmf component (ΔpH) for the electric one ($\Delta\Psi$), and indeed it accelerated cytochrome-f reduction in both strains to the same level (Figure 4A). Although direct nigericin action may cause an electrogenic bias, the drug also restored the generation cytochrome b_6f-associated charge separation phase during the first 10-ms after the flash (this so-called b-phase is shown as solid lines in Supplemental Figure 8). The b-phase half-times in *PETC-P171L* were significantly slower compared to the control strain (Figure 4B). In contrast, the b-phase buildup was accelerated by nigericin treatment. However, it was slightly delayed in nigericin-treated *PETC-P171L* compared to the control strain. Although nigericin also accelerated the b-phase buildup in the control strain, the difference was not significant.

In the light of the nigericin effect on the single-turnover cytochrome-f kinetics and b-phase generation, we further examined the phenotype of anoxic *PETC-P171L* cells in multiple-turnover conditions. To do

so, we repeated the electron transfer (Figure 4C) and P700 experiments (Figure 4D) in the presence and absence of nigericin. Indeed, Figure 4C shows that nigericin action abolished the *PETC-P171L* electron transfer bottleneck in the first ~50-ms of light. Moreover, the steady state rates were elevated in both strains when nigericin was present. In these cells, P700 redox states were changed upon nigericin addition (Figure 4D). In the absence of the drug, the control strain established a larger YNA and thus a smaller YI than *PETC-P171L*. The YND parameter was lowered to similar levels after adding nigericin. Upon nigericin addition, *PETC-P171L* cells established YNA and YI like control. Moreover, samples were indistinguishable in the absence of PSII activity (HA+DCMU, Figure 4D).

Taken together, these spectroscopic findings reveal that the P700 redox states under the actinic light illumination are differently distributed in *PETC-P171L* cells when oxygen levels were low. The different P700 redox state in *PETC-P171L* relied on upstream events on the level of the cytochrome b_6f and, as shown in single-turnover measurements, the ΔpH component of the *pmf* could be identified as being one of the limiting factors in oxygen-deprived algae. In most experiments at ~160 μmol photons·m^{-2}·s^{-1} actinic red light, *PETC-P171L* performed like control strains when cells were well aerated, suggesting that the acidification of the lumen pH in the permissive light was insufficient to trigger the cytochrome b_6f phenotype. In the last two results sections, we will explore the *PETC-P171L* performance in the presence of oxygen at various light intensities.

The *PETC-P171L* mutant showed an impaired photoautotrophic growth rate under high light

We evaluated whether the *PETC-P171L* mutation has an impact on cellular growth over the course of days. The control and *PETC-P171L* cells grown in liquid medium were spotted on solid medium and the cell growth was observed (Figure 5A). The cells showed a similar photoheterotrophic growth under moderate light illumination on solid TAP medium. On the other hand, when cells were grown photoautotrophically on solid TP medium, *PETC-P171L* cells grew more slowly than control cells under illumination of 100 μmol photons·m^{-2}·s^{-1} and 600 μmol photons·m^{-2}·s^{-1}. Although the *PETC-P171L* cells were able to grow photoautotrophically, the relative growth rate of the mutant cells was apparently slower in the stronger light (Figure 5A). We also analyzed the photoautotrophic growth of control and *PETC-P171L* strains in liquid medium under illumination of 200 μmol photons·m^{-2}·s^{-1}. We monitored

the chlorophyll concentration of liquid culture as a measure of cell growth (Figure 5B). The doubling

times of control and *PETC-P171L* were estimated to be 9.4 and 14 hours, respectively, indicating that

PETC-P171L cells grow 1.5 times slower under 200 μmol photons·m^{-2}·s^{-1}.

The *PETC-P171L* shows normal state transitions but displays an impaired NPQ induction under strong light illumination and during CO$_2$ limitations

The slower photoautotrophic growth of *PETC-P171L* cells may be ascribed to higher photosensitivity

at an intensity of 200 μmol photons·m^{-2}·s^{-1}. One photoprotection strategy involves NPQ to which state

transitions are significantly contributing to in algae (Allorent et al., 2013). A functional Qo-site is

required to activate the STT7 kinase for catalyzing state transitions (Vener et al., 1997). Thus, the higher

photosensitivity in *PETC-P171L* cells may result from the point mutation disturbing state transitions.

Therefore, we measured fluorescence emission spectra at 77 K after the induction of state transitions.

Indeed, control and the *PETC-P171L* cells exhibited higher intensity of PSII core fluorescence emission

(peaking at 688 nm) in state 1 while the PSI core fluorescence emission (peaking at 716 nm) was higher

in state 2 (Figure 6A). This indicates that both state 1 and state 2 were fully induced and that the *PETC-P171L* mutation does not affect state transitions. Moreover, we tested whether the accumulation of the

LHCSR3 protein is affected since it is necessary to establish the energy-dependent NPQ in algae (Peers

et al., 2009). We observed that LHCSR3 was similarly induced after 4 h of strong light illumination in

control and *PETC-P171L* cells grown photoautotrophically (Figure 6B). Accordingly, the requirements

to establish NPQ are met in *PETC-P171L*.

Next, we monitored photosynthesis via chlorophyll fluorescence over a 16-h photoperiod to seek for a

link between a possible restriction in the *PETC-P171L* photosynthetic machinery and the growth

phenotype that we observed in Figure 5. Therefore, we cultivated the control strain and *PETC-P171L*

in photobioreactors in the absence of air supply to favor CO$_2$ limitation and induce NPQ (Kanazawa

and Kramer, 2002). We followed the effective PSII quantum yield (Y(II)) and NPQ for 72 h

(Supplemental Figure 9), and the progressive Y(II) decline at dawn was more pronounced in *PETC-P171L*. Figure 7 displays a typical 16-h photoperiod peaking at 250 μmol photons·m^{-2}·s^{-1}. The low

Y(II) of ~0.1 during elevated light levels was similar in both strains (Figure 7A). However, the initial

drop in Y(II) was more pronounced in *PETC-P171L* during the first 2 h of light, and the relaxation

from lower to higher Y(II) values at the end of the photoperiod was slightly delayed in *PETC-P171L*,

reaching stable Y(II) after ~1 h darkness. The associated NPQ values differed more significantly

beyond the first 2 h of the photoperiod (Figure 7B). They were strongly reduced to ~0.4 in *PETC-*

P171L. NPQ peaked at 1.1 in the control strain after 4 h of light, slowly decreasing to 0.8 at the end of

the light period. Moreover, NPQ relaxation after the end of the light period (after 41 h of culture time

in the figure) was slower in *PETC-P171L* than in the controls (~2 h vs 1 h). Our photobioreactor

findings are quantitatively in agreement with additional PAM fluorimetry measurements. Therein

(Supplemental Figure 10), we observed that high light treated *PETC-P171L* cells showed a lower

Fv/Fm and, especially in low light, a smaller Y(II) due to a compromised NPQ formation efficiency

that paralleled a more reduced PQ pool.

Taken together, the chlorophyll fluorescence parameters in *PETC-P171L* suggested a limitation of

photosynthesis on the level of cytochrome b_6f complex at elevated irradiances and/or when CO_2

availability is low.

Discussion

In Arabidposis, the *pgr1* mutant showed reduced NPQ owing to cessation of total electron transfer at elevated light intensities above ~200 µmol photons·m^{-2}·s^{-1}, most likely at the level of b_6f (Munekage et al., 2001; Okegawa et al., 2005). P700 redox measurements of isolated *pgr1* thylakoids further suggested that the pH-dependent inactivation of electron transfer via the b_6f was shifted towards more alkaline pH by about 1 unit (Jahns et al., 2002). Yet, direct b_6f measurements of the Arabidopsis *pgr1* mutant are missing in literature. In the current work we successfully generated a *PETC-P171L* strain in *C. reinhardtii,* harboring a genetic site-directed alteration corresponding to the *pgr1* mutation in *A. thaliana* (Shikanai et al., 1999). In *C. reinhardtii*, this mutation did neither affect b_6f accumulation, state transitions, nor genetic NPQ regulation via LHCSR induction. We directly show that, depending on the environmental conditions, a ΔpH-dependent slow-down of electron transfer through the algal *PETC-P171L* b_6f complex is established. This restriction most likely explains the observed growth defect in the mutants due to diminished electron transfer activity. In the algal system, under conditions where CO_2 limitation was initiated in sealed photobioreactors, the algal Rieske mutant showed a lower NPQ above ~100 µmol photons·m^{-2}·s^{-1} already.

The PETC-P171L mutation slows down the electron transfer through b_6f in lumenal pH acidification.

The previous studies observed strong NPQ reduction in the Arabidopsis *pgr1* mutant in high light. This phenotype linked to reduction of electron transfer in the mutant plants (Munekage et al., 2001; Okegawa et al., 2005; Yamamoto and Shikanai, 2019). Our measurements of photosynthetic control in *PETC-P171L* revealed similarities to the available data in vascular plants. Namely, the bottleneck of electron transfer in *PETC-P171L* was detected at the level of PSII via chlorophyll fluorescence (Figure 7) and, depending on the condition, at the level PSI (Figure 3B). Moreover, we ruled out that state transitions were affected by the Qo-site mutation (Figure 6A) and further evaluation of the photosynthetic machinery pinpointed the *PETC-P171L* defect to the b_6f complex (Figure 4). We identified a significant b_6f defect in *PETC-P171L* under anoxia which had an impact on cytochrome-*f* reduction, Q-cycle electrogenicity, faster electron flow cessation during photosynthetic induction and facilitated P700

oxidation in the light. This relates the *PETC-P171L* phenotype for the first time to a mechanistic defect

in the b_6f complex that is due to an altered pH-dependence within the mutant b_6f complex. Accordingly,

the addition of nigericin, exchanging ΔpH for ΔΨ, abolished the impact of the mutation. Anaerobiosis

increases the ΔpH contribution to the *pmf* in darkness, which depends on functional ATP synthase

(Finazzi and Rappaport, 1998; Rappaport et al., 1999). Oxygen deprivation caused a charge separation

deficiency of the b_6f in *PETC-P171L*, which is a consequence of delayed mutant Qo-site turnover.

However, our nigericin approach to address the ΔpH impact could not restore the same rapid charge

separation onset in *PETC-P171L* that we observed in the control strain (Figure 4B). Thus, the mutation

could affect the b_6f complex on multiple levels. In fact, a cooperativity of high- and low-potential chain

reactions has been documented in previous Rieske iron sulfur protein (ISP) mutants of the cytochrome

bc_1 complex from yeast (Snyder et al., 2000). Still, this coupling mechanism operating between Qo- and

Qi-site remains elusive. The authors mutated homologous Rieske ISP residues at position 157 and 159

(orange-colored rectangle in Supplemental Figure 3), not too far from proline 171 of *C. reinhardtii*

PETC, which eventually result in a more reducing redox midpoint potential (E_m) of the Rieske ISC and

a slowdown of *b*-heme reduction (Snyder et al., 2000). Thus, similar effects could be anticipated in

PETC-P171L provided that the mutation lowers the Rieske ISC E_m in b_6f as originally proposed

(Munekage et al., 2001; Jahns et al., 2002). However, the E_m has not been determined in *pgr1*. Previous

data indicated that the E_m of reconstituted recombinant *C. reinhardtii* Rieske ISC is pH-dependent with

a pK_{ox} of 6.2 and a pK_{red} of 8.2 (Soriano et al., 2002). The E_m changes in the pH-dependent region with

-40 mV/pH. Thus, lowering the Rieske ISC E_m via the PETC-P171L mutation could impact pK_{ox} and

pK_{red}, and might alter inter b_6f electron transfer as shown in yeast (Snyder et al., 2000). In an alternative

scenario, the PETC-P171L mutation might alter binding of PQH_2 to the Q_o-site and consecutively

change the pH-dependent redox properties and oxidation of PQH_2. The Rieske ISC is found in the

hydrophilic domain of PETC (Figure 1B) and, based on structural similarities with bc_1 complex

(Darrouzet et al., 2000), it is supposed to be a flexible hydrophilic domain to realize electron transfer

between redox centers Rieske ISC and heme *f*. Additionally, proline 171 mutated in this study is the

second proline of the "Proline-X-Proline" motif, which is a common structure at the beginning of turn

structure (MacArthur and Thornton, 1991). This motif is exclusively found in PETC protein and conserved among Rieske ISP (Supplemental Figure 3). Accordingly, this motif forms a typical hairpin structure and the substitution of the second proline to leucine may disturb conformation of PETC hydrophilic domain structure, causing the functional impact in the redox chemistry of b_6f.

Our findings revealed that the generated ΔpH in anaerobic Chlamydomonas during 30-s of darkness was sufficient to trigger severe photosynthetic control in *PETC-P171L*, preventing efficient electrogenicity via the low-potential chain due to aggravated cytochrome-f reduction. As the effect of nigericin on photosynthetic control are well established (Joliot and Johnson, 2011), our nigericin data also revealed a significant contribution of the ΔpH in anaerobic cells (Figures 3 and 4). Accordingly, b_6f regulation is important for PSI photoprotection as well as for qE quenching in a reducing cellular environment such as anaerobiosis. The differential regulation of pH dependent inactivation of electron transfer via the b_6f in oxic and anoxic conditions in *PETC-P171L* is surprising. This might suggest that depending on oxygen availability (or metabolic congestion), the regulation of photosynthetic control is distinct in *C. reinhardtii*. However, revealing a low-light *pgr1* phenotype may also occur in plants when metabolism becomes restricted, e.g. under drought stress conditions where a downregulated ATP synthase activity favors large ΔpH amplitudes (Kohzuma et al., 2009). How can this be rationalized? Three non-exclusive scenarios may occur: First, metabolic control of ATP synthase downregulates H^+ conductivity which theoretically facilitates a light-dependent ΔpH build up in anaerobic conditions. Second, sustaining a ΔpH in an ATP synthase-dependent manner is favored in anoxic algae even in darkness (Finazzi and Rappaport, 1998; Rappaport et al., 1999). Third, the regulation of b_6f electron transfer via the Q-cycle is distinct under oxic and anoxic conditions (Buchert et al., 2020; Buchert et al., 2022). Under anoxic conditions stromal electron intake into b_6f changes the Q-cycle from a canonical cycle during LEF to an alternative cycle during CEF, which would facilitate more efficient proton translocating into the lumen as compared with the canonical Q-cycle. In line, photosynthetic control would be faster activated under anoxia and revealed the *PETC-P171L* phenotype even in low-light conditions.

Taken together, *PETC-P171L* may be a useful tool for future investigations under various metabolic and environmental conditions to uncover details of the cooperative mechanism between Qo- and Qi-site. Moreover, *PETC-P171L* may help to elucidate the interplay of various *pmf*-shaping ion channels/antiporters involved to fine-tune *pmf* partitioning in algae in response to metabolic constraints.

PETC-P171L mutation retards photoautotrophic cell growth.

The growth phenotype of *pgr1* plants under different light intensities has not been addressed in literature (Shikanai et al., 1999; Munekage et al., 2001; Yamamoto and Shikanai, 2019). We found that *PETC-P171L* grew less efficiently under photoautotrophic condition at growth rates 1.5 times slower in the logarithmic phase than control (Figure 5 and Supplemental Figure 11). Despite similar amounts of LHCSR3 in the control and *PETC-P171L* cells, NPQ could not be fully established in *PETC-P171L* and PSII performance was low due to a more reduced PQ pool in the mutant (Figure 7, Supplemental Figure 10). This analysis shows the *PETC-P171L* mutation impacts cellular growth and underlines the importance of a balanced photosynthetic control in wild type cells (control strain), including qE regulation via acidification of the lumen. These data demonstrate the tight interconnection of pH-dependent regulation of photosynthetic control and qE driven quenching, which also requires expression of LHCSR3.

Taken together, *PETC-P171L* exhibits a reduced photosynthetic electron transfer rate because of its hypersensitive photosynthetic control mechanism. The light intensity threshold to trigger the hypersensitive effect in the mutant b_6f can be modulated by the metabolic state of the cell. The molecular machinery to establish photosynthetic control, energy quenching and balancing of light harvesting requires functional wild type b_6f strains. These results are consistent with Arabidopsis studies, indicating that the effect of the proline substitution in PETC on electron transfer is *per se* conserved between the two organisms. Moreover, our study proves that the *PETC-P171L* phenotype is directly linked to functional properties of the b_6f complex.

Acknowledgement

1   The *petC-Δ1* strain and pACR4.5 plasmid were kindly gifted by Catherine de Vitry. The cDNA library

2   of *C. reinhardtii* and the pSL18 plasmid expression vector were provided by Michel Goldschmidt-

3   Clermont. The antibodies against PsaA and PetA/PETC were kindly gifted by Kevin Redding and

4   Olivier Vallon, respectively.

Figures

Figure 1

The position of Rieske ISC and PETC-Pro171 in the Chlamydomonas b_6f complex

Structures of the dimeric b_6f from the *C. reinhardtii* (PDB; 1Q90). The flexible hinge of PETC is not

modeled. The Rieske ISCs are shown in space filling representation and of which ligand amino acids

(Cys134, His136, Cys152, and His155 of PETC) and Pro171 of PETC (colored in magenta) and four

hemes (heme c_i is yellow, heme b_H and heme b_L are blue, and heme f is orange) are shown in ball-stick

model. The PetA (orange), PetB (blue), PETC (gray), PetB (green), PetG (magenta), PetL (yellow),

PETM (cyan), and PETN (red) subunits are shown. **(A)** and **(C)**. Overall structure with the lumenal side

at bottom and the stromal side at the top. The position of one of the Rieske ISC is indicated with arrow.

(B) and **(D)**. The enlarged view around the Rieske iron sulfur cluster. **(C)** and **(D)**. Protein surface is

modeled. The subunits and cofactors except for PETC subunit are transparent in panel A and B.

Figure 2

Generation of *PETC-P171L* mutant

The accumulation of PSI (PsaA) and b_6f (PETC and PetA) proteins was analyzed by immunoblotting.

The cells were grown in TAP liquid medium under the 50 μmol photons·m^{-2}·s^{-1}. The Pro171Leu of

PETC (*PETC-P171L*) and wild type sequence of PETC (control), the PETC gene expressing with own

promoter and UTRs (genomic) or the spliced PETC gene expressing with PSAD promoter UTRs

(cDNA). Four independent clones from each transformant, a dilution series for the wild type (wild type

(%)), and ΔPETC (T$_{15-3}$ in Supplemental Figure 5) were also analyzed.

 Figure 3

The *PETC-P171L* mutation restricts photosynthetic performance of light- and dark-adapted cells

predominantly in anaerobic conditions.

P700 redox states were measured in **(A)** oxic and **(B)** anoxic light-adapted cells. The fractional yields

of P700 in the photo-oxidizable state (YI), the non-oxidizable form due to acceptor side limitation

(YNA), and the pre-oxidized P700 due to donor side limitation (YND) are shown (parameter-specific

one-way ANOVA, $n = 3$, $P < 0.05$). The oxic controls in absence of PSII activity (HA+DCMU) retained

a higher YI due to enhanced steady state CEF (expressed charge separations·s^{-1}·PSI^{-1} ± SD). In anoxic

conditions, *PETC-P171L* failed to establish the YNA due to larger YND. **(C)** After a 30-s dark period,

electron transfer rates over a 10-s photosynthetic induction phase were measured in oxic light-adapted

cells. The kinetics were largely indistinguishable (means ± SD are shown, $n = 3$). Total electron transfer

rates in the controls were enhanced at a later stage of illumination (filled asterisks, Student's t-test $P <$

0.05), and *PETC-P171L* showed transiently higher rates at 100-ms in absence of PSII activity (open

asterisk). **(D)** Total electron transfer in *PETC-P171L* was significantly less efficient at discrete time

points, especially in the initial ~25-ms (filled asterisk). A significant difference during induction

required PSII activity. **(E)** The electrochromic shift ~10-ms rising phase, associated with charge

1   separation activity upon a single b_6f turnover, was less pronounced in anoxic light-adapted *PETC-P171L*.

2   Sub-saturating flashes were fired on light-adapted cells after a 30-s dark period.

Figure 4

The cytochrome b_6f-related electron transfer restrictions in *PETC-P171L* cells under anaerobic conditions is overcome by diminishing the ΔpH via nigericin.

(A) The control and *PETC-P171L* cells displayed different cytochrome-*f* reduction half-times upon a single cytochrome b_6f turnover under anoxic conditions (means of n=4 ± SD are shown, miniscules on top indicate significances using One-Way ANOVA/Fisher-LSD, P > 0.05). The slower cytochrome-*f* reduction in *PETC-P171L* cells as well as control strain kinetics were accelerated to similar values upon nigericin (Nig) addition. **(B)** The events that affected cytochrome-*f* reduction also decreased the *PETC-P171L* electrogenic activity in the low-potential chain during the Q-cycle. The *b*-phase (for kinetics, see Supplemental Figure 8) also showed a longer half-time in *PETC-P171L* (means of n=4 ± SD are shown, miniscules on top indicate significances using One-Way ANOVA/Fisher-LSD, P > 0.05). Nigericin addition accelerated the buildup of the *b*-phase in the *PETC-P171L* strain. The drug accelerated the *b*-

phase insignificantly in the control. **(C)** After a 30-s dark period, electron transfer rates over a 10-s photosynthetic induction phase were measured in anoxic light-adapted cells (averaged kinetics ± SD, *n* = 3). Total electron transfer in *PETC-P171L* was significantly less efficient within the first ~50-ms (filled asterisk, Student's t-test *P* < 0.05). The different induction kinetics between the strains relied on the ΔpH and became similar after addition of the H^+/K^+ ionophore nigericin (Nig., 10 μM), although the control showed higher rates at discrete sampling intervals (open asterisk). In both strains the steady state rates were higher when Nig. was present. **(D)** On the level of co-dependent PSI parameters, the fractional yields of P700 in the photo-oxidizable state (YI), the non-oxidizable form due to acceptor side limitation (YNA), and the pre-oxidized P700 due to donor side limitation (YND) are shown (parameter-specific one-way ANOVA, *n* = 3, *P* < 0.05). The YI was larger in *PETC-P171L* owing to a diminished YNA. YND was lowered to similar levels after adding Nig. and diminishing ΔpH-dependent photosynthetic control. Moreover, further PSII inhibition (HA+DCMU) produced similar P700 yields.

1 **Figure 5**

A

B

The photoautotrophic growth rate is slowed down at low cell density in *PETC-P171L*.

(A) Cells were grown photoheterotrophically on solid TAP medium (50 μmol photons·m^{-2}·s^{-1}) or

photoautotrophically on solid TP medium (100 or 600 μmol photons·m^{-2}·s^{-1}). Three independent

clones of the control and *PETC-P171L* strain (*P171L*) were analyzed. **(B)** Photoautotrophic cell

growth is monitored by measuring Chlorophyll concentration of the TP liquid culture medium grown

under continuous light (200 μmol photons·m^{-2}·s^{-1}). Black line represents control and red line

represents *PETC-P171L (P171L)*. Three biological replicates were averaged (± standard errors).

Figure 6

The *PETC-P171L* mutation does neither affect induction of state-transitions, nor LHCSR3

expression induction and/or stability.

(**A**) The TP-grown Chlamydomonas cells acclimated under high light (300 μmol photons·m^{-2}·s^{-1}) were

resuspended with TP at 2 μg Chlorophyll·mL^{-1} and incubated in state 1 or state 2 inducible condition.

In the case of state 1 inducible treatment, cells were incubated under 200 μmol photons·m^{-2}·s^{-1} in the

presence of DCMU for 10 minutes. In the case of state 2 inducible treatment, cells were incubated in

the darkness for 10 minutes in the presence of 100 mM glucose and 50 U·mL^{-1} glucose oxidase. Samples

were excited with 430 nm and the emitted fluorescence were recorded from 600 to 800 nm in liquid

nitrogen. The fluorescence spectra were normalized at 688 nm. (**B**) The cells grown in TP medium were

shifted from low light (15 μmol photons·m^{-2}·s^{-1}) to stronger light (200 μmol photons·m^{-2}·s^{-1}) and

immune-chase experiment with the antibody against LHCSR3 protein was performed for cellular

proteins. Cells were harvested before shifting to higher light (0), after 4, 8, and 12 hours shifting to high

light. Ponceau staining panel served as loading control.

Figure 7

CO$_2$ limitation reveals the chlorophyll fluorescence phenotype of a limited photosynthetic capacity

in *PETC-P171L* at lower irradiances.

Averaged photobioreactor experiments (n = 3 ± SD) of non-aerated cultures are shown for control and

PETC-P171L cells. **(A)** The PSII quantum yield Y(II) and **(B)** NPQ show that *PETC-P171L*

performance is less efficient in the twilight and throughout the day, respectively. The second

photoperiod is shown (16-h light / 8-h darkness, PAR is µmol photons·m^{-2}·s^{-1}, for complete

experiments refer to Supplemental Figure 9).

References

Allorent G, Tokutsu R, Roach T, Peers G, Cardol P, Girard-Bascou J, Seigneurin-Berny D, Petroutsos
D, Kuntz M, Breyton C, Franck F, Wollman FA, Niyogi KK, Krieger-Liszkay A, Minagawa J,
Finazzi G (2013) A dual strategy to cope with high light in Chlamydomonas reinhardtii. Plant
Cell **25:** 545-557
Alric J (2014) Redox and ATP control of photosynthetic cyclic electron flow in Chlamydomonas
reinhardtii: (II) involvement of the PGR5-PGRL1 pathway under anaerobic conditions.
Biochim Biophys Acta **1837:** 825-834
Bailleul B, Cardol P, Breyton C, Finazzi G (2010) Electrochromism: a useful probe to study algal
photosynthesis. Photosynthesis Research **106:** 179-189
Bonente G, Ballottari M, Truong TB, Morosinotto T, Ahn TK, Fleming GR, Niyogi KK, Bassi R (2011)
Analysis of LhcSR3, a protein essential for feedback de-excitation in the green alga
Chlamydomonas reinhardtii. PLoS Biol **9:** e1000577
Buchert F, Hamon M, Gabelein P, Scholz M, Hippler M, Wollman FA (2018) The labile interactions of
cyclic electron flow effector proteins. J Biol Chem **293:** 17559-17573
Buchert F, Mosebach L, Gabelein P, Hippler M (2020) PGR5 is required for efficient Q cycle in the
cytochrome b6f complex during cyclic electron flow. Biochem J **477:** 1631-1650
Buchert F, Scholz M, Hippler M (2022) Electron transfer via cytochrome b6f complex displays
sensitivity to antimycin A upon STT7 kinase activation. Biochem J **479:** 111-127
Darrouzet E, Valkova-Valchanova M, Moser CC, Dutton PL, Daldal F (2000) Uncovering the [2Fe2S]
domain movement in cytochrome bc1 and its implications for energy conversion. Proc Natl
Acad Sci U S A **97:** 4567-4572
de Vitry C, Finazzi G, Baymann F, Kallas T (1999) Analysis of the nucleus-encoded and chloroplast-
targeted rieske protein by classic and site-directed mutagenesis of Chlamydomonas. Plant
Cell **11:** 2031-2044
Delepelaire P, Wollman FA (1985) Correlations between fluorescence and phosphorylation changes
in thylakoid membranes of Chlamydomonas reinhardtii in vivo: A kinetic analysis. Biochimica
et Biophysica Acta (BBA) - Bioenergetics **809:** 277-283
Depège N, Bellafiore S, Rochaix JD (2003) Role of chloroplast protein kinase Stt7 in LHCII
phosphorylation and state transition in Chlamydomonas. Science **299:** 1572-1575
Finazzi G, Rappaport F (1998) In vivo characterization of the electrochemical proton gradient
generated in darkness in green algae and its kinetic effects on cytochrome b6f turnover.
Biochemistry **37:** 9999-10005
Grossman AR, Catalanotti C, Yang W, Dubini A, Magneschi L, Subramanian V, Posewitz MC, Seibert
M (2011) Multiple facets of anoxic metabolism and hydrogen production in the unicellular
green alga Chlamydomonas reinhardtii. New Phytol **190:** 279-288
Harris EH (1988) The Chlamydomonas Sourcebook. Academic Press, San Diego
Jahns P, Graf M, Munekage Y, Shikanai T (2002) Single point mutation in the Rieske iron-sulfur
subunit of cytochrome b6/f leads to an altered pH dependence of plastoquinol oxidation in
Arabidopsis. FEBS Lett **519:** 99-102
Jokel M, Johnson X, Peltier G, Aro EM, Allahverdiyeva Y (2018) Hunting the main player enabling
Chlamydomonas reinhardtii growth under fluctuating light. Plant J **94:** 822-835
Joliot P, Johnson GN (2011) Regulation of cyclic and linear electron flow in higher plants. Proc Natl
Acad Sci U S A **108:** 13317-13322
Joliot P, Joliot A (2002) Cyclic electron transfer in plant leaf. Proc Natl Acad Sci U S A **99:** 10209-
10214
Junge W, Witt HT (1968) On the ion transport system of photosynthesis--investigations on a
molecular level. Z Naturforsch B **23:** 244-254
Kanazawa A, Kramer DM (2002) *In vivo* modulation of nonphotochemical exciton quenching (NPQ)
by regulation of the chloroplast ATP synthase. Proceedings of the National Academy of
Sciences, USA **99:** 12789-12794

Kanazawa A, Kramer DM (2002) In vivo modulation of nonphotochemical exciton quenching (NPQ) by regulation of the chloroplast ATP synthase. Proc Natl Acad Sci U S A **99:** 12789-12794

Klughammer C, Schreiber U (1994) An improved method, using saturating light pulses, for the determination of photosystem I quantum yield via P700$^+$-absorbance changes at 830 nm. Planta **192:** 261-268

Kohzuma K, Cruz JA, Akashi K, Hoshiyasu S, Munekage YN, Yokota A, Kramer DM (2009) The long-term responses of the photosynthetic proton circuit to drought. Plant Cell Environ **32:** 209-219

Kurisu G, Zhang H, Smith JL, Cramer WA (2003) Structure of the cytochrome b6f complex of oxygenic photosynthesis: tuning the cavity. Science **302:** 1009-1014

Li XP, Muller-Moule P, Gilmore AM, Niyogi KK (2002) PsbS-dependent enhancement of feedback de-excitation protects photosystem II from photoinhibition. Proc Natl Acad Sci U S A **99:** 15222-15227

MacArthur MW, Thornton JM (1991) Influence of proline residues on protein conformation. J Mol Biol **218:** 397-412

Munekage Y, Takeda S, Endo T, Jahns P, Hashimoto T, Shikanai T (2001) Cytochrome b(6)f mutation specifically affects thermal dissipation of absorbed light energy in Arabidopsis. Plant J **28:** 351-359

Nawrocki WJ, Bailleul B, Cardol P, Rappaport F, Wollman FA, Joliot P (2019) Maximal cyclic electron flow rate is independent of PGRL1 in Chlamydomonas. Biochim Biophys Acta Bioenerg **1860:** 425-432

Nishio JN, Whitmarsh J (1993) Dissipation of the Proton Electrochemical Potential in Intact Chloroplasts (II. The pH Gradient Monitored by Cytochrome f Reduction Kinetics). Plant Physiol **101:** 89-96

Okegawa Y, Tsuyama M, Kobayashi Y, Shikanai T (2005) The pgr1 mutation in the Rieske subunit of the cytochrome b6f complex does not affect PGR5-dependent cyclic electron transport around photosystem I. J Biol Chem **280:** 28332-28336

Peers G, Truong TB, Ostendorf E, Busch A, Elrad D, Grossman AR, Hippler M, Niyogi KK (2009) An ancient light-harvesting protein is critical for the regulation of algal photosynthesis. Nature **462:** 518-521

Rappaport F, Finazzi G, Pierre Y, Bennoun P (1999) A new electrochemical gradient generator in thylakoid membranes of green algae. Biochemistry **38:** 2040-2047

Rumberg B, Siggel U (1969) pH changes in the inner phase of the thylakoids during photosynthesis. Naturwissenschaften **56:** 130-132

Shikanai T, Munekage Y, Shimizu K, Endo T, Hashimoto T (1999) Identification and characterization of Arabidopsis mutants with reduced quenching of chlorophyll fluorescence. Plant Cell Physiol **40:** 1134-1142

Snyder CH, Gutierrez-Cirlos EB, Trumpower BL (2000) Evidence for a concerted mechanism of ubiquinol oxidation by the cytochrome bc1 complex. J Biol Chem **275:** 13535-13541

Soriano GM, Guo LW, De Vitry C, Kallas T, Cramer WA (2002) Electron transfer from the Rieske iron-sulfur protein (ISP) to cytochrome f in vitro. Is a guided trajectory of the ISP necessary for competent docking? J Biol Chem **277:** 41865-41871

Stroebel D, Choquet Y, Popot JL, Picot D (2003) An atypical haem in the cytochrome b$_6$f complex. Nature **426:** 413-418

Sueoka N (1960) MITOTIC REPLICATION OF DEOXYRIBONUCLEIC ACID IN CHLAMYDOMONAS REINHARDI. Proc Natl Acad Sci U S A **46:** 83-91

Takahashi H, Clowez S, Wollman FA, Vallon O, Rappaport F (2013) Cyclic electron flow is redox-controlled but independent of state transition. Nat Commun **4:** 1954

Takahashi H, Schmollinger S, Lee JH, Schroda M, Rappaport F, Wollman FA, Vallon O (2016) PETO Interacts with Other Effectors of Cyclic Electron Flow in Chlamydomonas. Mol Plant **9:** 558-568

Tikhonov AN, Khomutov GB, Ruuge EK, Blumenfeld LA (1981) Electron transport control in chloroplasts. Effects of photosynthetic control monitored by the intrathylakoid pH. Biochimica et Biophysica Acta (BBA) - Bioenergetics **637:** 321-333

Velthuys BR (1978) A third site of porton translocation in green plant photosynthetic electron transport. Proc Natl Acad Sci U S A **75:** 6031-6034

Vener AV, van Kan PJ, Rich PR, Ohad II, Andersson B (1997) Plastoquinol at the quinol oxidation site of reduced cytochrome bf mediates signal transduction between light and protein phosphorylation: Thylakoid protein kinase deactivation by a single-turnover flash. Proc Natl Acad Sci U S A **94:** 1585-1590

Witt HT (1979) Energy conversion in the functional membrane of photosynthesis. Analysis by light pulse and electric pulse methods. The central role of the electric field. Biochim Biophys Acta **505:** 355-427

Yamamoto H, Shikanai T (2019) PGR5-Dependent Cyclic Electron Flow Protects Photosystem I under Fluctuating Light at Donor and Acceptor Sides. Plant Physiol **179:** 588-600

Yamano T, Iguchi H, Fukuzawa H (2013) Rapid transformation of Chlamydomonas reinhardtii without cell-wall removal. J Biosci Bioeng **115:** 691-694