Role of immigrants and muzzle contacts in the uptake of a novel food by wild vervet

monkeys

Pooja Dongre *, #, 1, 2, Charlotte Canteloup #, 1, 2, Gaëlle Lanté 1, 3, Mathieu Cantat 2, Erica van

de Waal 1, 2

[1] Department of Ecology and Evolution, University of Lausanne, 1015 Lausanne,

Switzerland

[2] Inkawu Vervet Project, Mawana Game Reserve, KwaZulu Natal, 3115, South Africa

[3] University of Poitiers, France

[#] These authors contributed equally

*Correspondence to: pooja.dongre@unil.ch

ABSTRACT

The entry into and uptake of information in social groups is critical for behavioural

adaptation by long-lived species in rapidly changing environments. We exposed five groups

of wild vervet monkeys to a novel food to investigate innovation of processing and

consuming it. We observed whether dispersing males could introduce this innovation into

groups and compare uptake between groups. We report that immigrant males innovated in

two groups, and an infant innovated in one group. In two other groups, immigrant males

imported the innovation from another group. Uptake of the novel food was fastest in groups

where immigrant males ate first and younger individuals were most likely to acquire the

novel food. We also investigated the role of muzzle contact behaviour in information seeking

around the novel food. Muzzle contacts decreased in frequency over repeated exposures to

the novel food, were targeted towards knowledgeable adults the most, and juveniles and

naïve individuals initiated the most. Knowledgeable adult males were targeted but rarely

initiated muzzle contacts, whereas knowledgeable adult females were targeted and initiated

them, emphasising an adult sex difference in seeking social information. We highlight the

potential of immigrants as vectors of information, enabling rapid behavioural adaptation and

population level culture.

INTRODUCTION

To thrive in rapidly changing environments, including those induced by humans,

animals must respond quickly to relevant information about their surroundings [1]. Climate

change induced invasions, as well as the introduction of human artefacts into the

environment can affect different species in myriad ways, for example, bringing new threats,

disruptions, competition, or novel resource opportunities. Adaptive behavioural responses to

such changes can include effectively avoiding new predators, maintaining high competitive

ability and exploiting novel resources [1–4]. For long-lived species, behavioural responses

are crucial to adapt fast enough when circumstances change rapidly and genetic adaption

would not suffice. Whilst transmission mechanisms of genetic adaptation are well-

understood, our understanding of how behavioural adaptations arise and spread is murky,

and the role of individual heterogeneity in a group in this remains underexplored [5].

Research has identified two main classes of behavioural response to novel stimuli in

animals. These are neophobia and exploration [6,7]. Neophobia refers to the avoidance of

potentially risky novelty, which could include new predators or unknown toxins. Exploration,

on the other hand, involves behaviours that seek information about novel stimuli. Whilst

seeming to be opposites of one spectrum, neophobia and exploration are likely motivated by

different needs and served by different mechanisms, and it is possible to express high or low

levels of both, simultaneously [6]. Obtaining novel information directly from the environment

requires overcoming neophobia and engaging in exploration, tendencies for which may vary

between individuals [5], and which produces information or knowledge, potentially facilitating

innovation.

Kummer & Goodall [8] defined innovation as, "a solution to a novel problem, or a novel

solution to an old one", and "a new ecological discovery such as a food item not previously

part of the group". Behavioural innovations can therefore allow long-lived species to adapt

their behaviour quickly to changing circumstances, for example to exploit a novel resource

introduced into the current habitat (e.g. [9]). Nonetheless, if explorative conspecifics or

individuals that uniquely possess particular knowledge are present, one might rather learn

from them via social learning. Indeed, many studies of diverse species in captivity have

found that observing a conspecific eat a novel food reduces neophobic responses (reviewed

in [6]: e.g. marmosets [10], rooks [11], aye-ayes [12], as well as humans [13]). In the wild, a

study on jackdaws found the same [14], and similarly, wild baboons handled a novel food for

longer after seeing a demonstrator do so, though this was dependent on their personality

[15]. Further investigation is required in the wild, since the risk for foraging animals to ingest

toxins via unknown foods can be high, whilst this risk is diminished in captivity. Moreover,

the individual differences of the observed conspecifics may be important. In addition, in wild

groups, dispersing individuals can import information or behavioural innovations, upon

immigration, into new groups [16–19], thus facilitating the spread of information at the

population level, but experimental evidence focused on multiple groups is lacking.

Within wild groups, animals can use social information to guide foraging decisions.

Many social learning studies in primates have focused on visual access to information (e.g.

[15,18,20–23], and see review in [24]). However, for Old World monkeys, detailed olfactory

information, in a foraging context, may be acquired through muzzle contact behaviour – the

act of one individual bringing their muzzle into very close proximity with another's [25–28].

Indeed, previous studies found that, whilst foraging, muzzle contacts were most commonly

initiated by infants and juveniles towards adults [28,29], which supports the function in

information acquisition as young animals are still learning about their dietary repertoire, and

adults are likely the most reliable sources of information. Nord et al. [28] also suggest that

muzzle contacts are constrained by social tolerance, due to the necessary close proximity,

which may constrain information transmission. In the presence of novel resources, muzzle

contacts may be useful to adults as well as youngsters. Experimental research into this

mode of information transmission in the presence of a novel resource is now required.

Vervet monkeys (*Chlorocebus pygerythrus*) are a species that thrive in natural, urban

and agricultural habitats, and are widely distributed throughout eastern sub-Saharan Africa

[30]. This makes them ideal species in which to investigate adaptation to novel

environmental conditions. They live in multi-male multi-female troops, with philopatric

females and males dispersing multiple times in their lives, producing considerable adult

individual heterogeneity in groups. Furthermore, frequently dispersing males could serve as

vectors of information between groups, potentially facilitating behavioural adaptation to their

large geographical range [30]. In a previous study by our team [20], two groups (NH, KB) of

wild vervet monkeys were provided with a novel food that required extraction (peanuts in

shells) before consumption. The results supported social transmission of the opening

techniques used to extract peanuts, based on visual attention to demonstrators [20]. In the

present study, we presented this novel food to three more groups (AK, BD, LT) of the same

study population, to study five groups in total. Our first line of investigation here concerned 1)

innovation of extracting and eating the novel food. Second, utilizing natural dispersals of

males between our study groups, we investigated 2a) variation between groups in their

uptake of the novel food according to who innovated, or initiated eating, and 2b)

demographic (age, sex, rank) variation in in uptake of the novel food. Third, we analysed 3a)

the rate of muzzle contacts as groups' had more exposure and therefore increased

familiarity with the food; and 3b) whether knowledge of the food, age, sex and / or rank

influenced muzzle contact behaviour.

As questions 1) and 2a) were largely exploratory, we did not make specific predictions.

Regarding question 2b) we expected uptake to be more likely in younger monkeys, 2b)

during the first exposure, and over four exposures, due to previous work suggesting that

juveniles take more risks [31] and are less neophobic [32–35] than adults. Regarding

question 3a) we expected muzzle frequency to decrease as groups had more exposures to

the novel food due to its likely function in obtaining information acquisition. In 3b) we

expected an effect of knowledge, specifically for naïve monkeys to initiate more, and

knowledgeable monkeys to be targeted more. Similarly, we expected an effect of age, with

juveniles more likely initiators and adults more likely targets.

These combined approaches broaden our understanding of routes to behavioural

adaptation to exploit a novel resource in the wild, and the potential contribution to this of

dispersal.

RESULTS

We refer to the group AK differentially as AK_{19} and AK_{20}, representing their status in

2019 and 2020, respectively, as 40% of the group composition changed between years due

to dispersals, deaths and changes in age categories (infants that became juveniles; see

Table S1 and detailed description in Materials and Methods).

Across this experiment, a total of 81 vervet monkeys in five groups, learned to

successfully extract and eat peanuts during four exposures from the first eating event (Table

1).

1) Innovation

When presented with a novel food, in groups that had new immigrant males (AK_{20}, BD,

LT, NH), new immigrants were the first to extract and eat peanuts: at the first exposure in

BD, LT and NH, and at the second exposure in AK_{20}. Two of these males were previously

naïve to peanuts and individually innovated the solution (*Avo* in NH, *Bab* in LT); and two

imported the innovation from NH into their new groups (*Pro* in BD and *Yan* in AK_{20}; see Fig.

1). In all four cases, the males had been present in the group for less than three months

prior. In KB an infant (*Aar*) innovated at the third exposure (see Table S1), and in AK_{19} no

one innovated during their single exposure. There were no new immigrant males within the

last three months in AK_{19} at the time of their exposure to peanuts, nor in KB at any of their

exposures.

2a) Variation in the uptake of novel food according to who ate first

During the first exposure to peanuts in BD, LT and NH, when new immigrant males

initiated eating peanuts, we observed that the following percentages of these groups learned

to extract and eat peanuts during that exposure: BD: 31%; LT: 20%; NH: 9% (Fig. 2a). In the

first exposures in AK_{19} and KB, no monkeys learned to extract and eat peanuts. After an

immigrant male ate in AK_{20}, at the second exposure in that group (but the first exposure with

a new immigrant male, who was knowledgeable about peanuts) 30% of the group followed

during that exposure (Fig. 2a). When an infant innovated at the third exposure in KB, no

other group members followed during that exposure (Fig. 2a). After four exposures from the

first eating event in all groups, the percentages of each group extracting and eating peanuts

were: 95% in AK_{20}; 66% in BD; 21% in KB; 84% in LT; and 20% in NH (Fig. 2b).

2b) Demographic variation in the uptake of the novel food

Here we examined whether age, sex and rank predicted successfully extracting and

eating peanuts in the first exposure and over the course of four exposures. During the first

exposure, we found a significant main effect of age, with juveniles more likely to extract and

eat peanuts than both adults (GLMM, Tukey HSD: Estimate = 1.38, Standard error = 0.51, z

= 2.70, p = 0.012) and infants (Est. = 2.12, SE = 0.74, z = 2.85, p = 0.011), but no difference

between infants and adults (Est. = -0.74, SE = 0.70, z = -1.05, p = 0.538). Rank showed a

non-significant trend (Est. = -1.48, SE = 0.81, z = -1.83, p = 0.067) with high rankers more

likely than low rankers to extract and eat peanuts. There was no main effect of sex (Est. =

0.63, SE = 0.42, z = 1.50, p = 0.134). Over four exposures, we found a significant main

effect of age, with juveniles more likely to extract and eat peanuts than adults (Est. = 1.62,

SE = 0.54, z = 3.00, p = 0.008) and infants (Est. = 1.92, SE = 0.64, z = 3.02, p = 0.007), but

no difference between infants and adults (Est. = -0.30, SE = 0.54, z = -0.55, p = 0.845). We

also found a significant main effect of rank, with high rank individuals more likely than low

rank individuals to eat (GLMM: Est. = -2.66, SE = 0.80, z = -3.70, p < 0.001). We found no

main effect of sex (Est. = 0.27, SE = 0.41, z = 0.66, p = 0.510).

3a) Muzzle contact frequency in groups

Here, in the groups where large proportions of monkeys had started to eat peanuts

over four exposures (AK_{20}, BD and LT), we recorded 569 muzzle contacts in total. We found

that exposure number (1-4) had a significant effect on the rate of muzzle contact, with the

rate decreasing as groups had more exposures to peanuts (LME: Est. = -0.71, SE = 0.27, t =

-2.63, p = 0.030).

3b) Individual variation in muzzle contact behaviour

In the final analysis, with the simple model (see Materials and Methods) we found

significant main effects of both knowledge (Est. = 1.63, SE = 0.24, z = 6.67, p < 0.001) and

age (Est. -3.51, SE = 0.68, z = -5.18, p < 0.001) on individuals' status as initiators or targets

of muzzle contacts. Knowledgeable individuals and adults were more likely to be targets

rather than initiators of muzzle contact, compared to naïve individuals and juveniles/infants

respectively. We did not find significant main effects of sex or rank (sex: Est = 0.72, SE =

0.65, z = 1.12, p = 0.26; rank: Est. = 0.025, SE = 0.021, z = 1.19, p = 0.236).

When removing rank but adding interactions between knowledge and age and

knowledge and sex to the model, we found significant interactions between both, knowledge

and age (Est. = -1.79, SE = 0.52, z = -3.48, p < 0.001; Fig. 3A, left), and knowledge and sex

(Est. = 1.60, SE = 0.54, z = 2.93, p = 0.003; Fig. 3A, right). In Figure 3A (left) we can see

clearly the effect of knowledge on adults, but not on juveniles. In Figure 3A (right) we see

that the effect of knowledge on males and females is generally similar, but slightly stronger

in males.

In the three-way interaction model, the interaction between knowledge, age and sex

was not significant (Est. = -2.11, SE = 1.31, z = -1.62, p = 0.106; Fig. 3B).

DISCUSSION

By looking at the uptake of a novel food by members of five groups of wild vervet

monkeys, we found evidence for new immigrant males as fast innovators, and as vectors of

information between groups. We saw faster uptake of the novel food by groups when new

immigrant males, rather than infants or juveniles, ate first. We also found effects of age and

rank on uptake of the food. In addition, as groups had more exposures to the food, muzzle

contact behaviour decreased. Between dyads involved in muzzle contacts, both knowledge

and age, and interactions between knowledge and age, and knowledge and sex influenced

whether individuals were initiators or targets. Below we discuss the contributions of these

results to our perspectives on the potential value of dispersing individuals in transmission of

behavioural adaptations to novel circumstances around populations.

1) Innovation

Fast innovation to exploit a novel resource by new immigrants, in this case at the first

exposure to a novel food, could be linked to a dispersal state by physiological conditions

related to dispersal. Whilst the sample size of our study is low, the fact that no monkeys

innovated during the first exposures to peanuts in groups with no new immigrants supports

this too. Dispersal has indeed previously been associated with exploration and boldness (i.e.

low neophobia) in several taxa, with associated neurochemical variation, both within

(ontogenetically) and between individuals [36]. Moreover, evidence links lower serotonergic

activity with: earlier dispersal in rhesus macaques [37], greater social impulsivity in vervet

monkeys [38], and reduced harm avoidance in humans [39]; all of which together relate low

neophobia, or novelty seeking, with dispersal. Evidence does not however suggest that the

dispersing sex in wild vervet monkeys are more bold or explorative overall [40], and in the

present study long-term resident males did not show increased interest in the novel food. In

another population, long-term resident males also showed reduced responses to novel foods

compared to other age-sex classes [41]. We suggest rather that the unique individual, social

and environmental factors that prompt a male to disperse [42] may trigger a transitory

exploratory behavioural syndrome [43] that may subside again once males acquire more

secure residency in a group. Since dispersal inherently involves heightened risk, periods of

long-term residency would be well-served by a state characterised by reduced exploration

and increased neophobia to balance costs of risk-taking over the lifetime. The large variation

in risky predator inspection by adult male vervet monkeys, compared to adult females found

in [40] also supports this. Future work focused on the behaviour of dispersing individuals at

multiple time points, both proximal and distal to dispersal events, in this species, and others,

will help to more conclusively address this hypothesis. We highlight the need for researchers

to consider the nuances of life-history characteristics beyond simply splitting by broad age-

sex categories.

2a) Individual variation in uptake of a novel food according to who ate first

When immigrant males extracted and ate a food that was novel to their new groups,

i.e. during the first exposure to it (in BD, LT and NH), other monkeys quickly followed them in

doing so. As discussed above, two of these cases (LT, NH) involved innovation by the

immigrant males, and in the other (BD), the male had learned to extract and eat the food in

his previous group. Similarly, in AK, uptake of the novel food began precisely when a new

immigrant imported the innovation during the group's second exposure. In contrast, following

innovation by an infant in KB at their third exposure, no other individuals followed in

extracting and eating peanuts during this exposure. Over the three subsequent exposures

that followed these initial eating events, very few monkeys started to extract and eat peanuts

in KB, whereas in BD, LT and AK, where new immigrants ate first, large proportions of these groups learned to extract and eat peanuts. In NH however, this was not the case and is of great interest because though a new immigrant male innovated at the first exposure, he left the group before the second exposure. Closer inspection of our data revealed that, only juveniles had learned after the male in the first exposure and were thus the only knowledgeable individuals at the second exposure. Similarly, in KB, only juveniles ate after the infant. One interpretation of these results is that immigrant males were more effective in facilitating group members to overcome neophobia towards a novel food than infants or juveniles, which is in line with studies reporting age-biased social learning [18,20]. Nonetheless, in NH, more individuals did eventually start to eat during their fourth exposure, including a high-ranked adult female (after which it spread rapidly resulting in the data presented in [20]). In NH, the juveniles eating at the beginning were older than the infants and one year old who started eating in KB. It is possible that age-bias is less strong for older juveniles since they should have more reliable knowledge than their very young counterparts. Alternatively, this difference between NH and KB could be because the juveniles eating earlier in NH were of high rank, whereas the infants and juveniles who began to eat in KB were all of low rank. Indeed, rank-biased social learning has also been found in previous work in two groups (KB and NH) of this study population [20,44]. It is nonetheless likely that various interactions of socioecological factors affect the influence of juveniles in overcoming food neophobia in the wild.

Moreover, whilst previous experiments suggested that high-ranked adult philopatric females are preferred over high-ranked adult males as models to learn from [22], in the context of exploiting a novel resource, risk dynamics come into play. Adult females are likely to be the most risk averse age-sex category, due to great potential negative impact of risks on their inclusive fitness, especially when young dependent offspring are present or while being pregnant. This might limit their potential to discover new information that others can exploit. Under these conditions, adult males that are either in an exploratory dispersal state,

or that enter a group with knowledge of resources previously unknown to the group, or more

exploratory juveniles of high rank, may play important roles in generating and facilitating the

spread of behavioural adaptations to exploit novel resources and face rapid environmental

changes.

2b) Demographic variation in the uptake of the novel food

In the first exposure juveniles were most likely to eat, and over four exposures, both

infants and juveniles were more likely to eat than adults were. These results suggest that

younger animals overcome neophobia faster than adults, corresponding closely with results

regarding risk-taking in another population of vervet monkeys [31], and work in other species

suggesting that juveniles are overall more exploratory and less neophobic [32–35].

High rank individuals were more likely to eat over the whole experiment, probably

due to preferential access to the resource as it became more familiar.

3a) Muzzle contact frequency in groups

At the group level, frequency of muzzle contact decreased as groups had more

exposures to the novel food. This demonstrates the relevance of the behaviour in the context

of unknown or rarely encountered foraging items, because as the food became more familiar

to the monkeys, they sought olfactory information from their conspecifics less frequently.

This result concurs with findings from a similar study in wild olive baboons [25].

3b) Individual variation in muzzle contact behaviour

In individual monkeys, knowledge of the novel food predicted whether they were more

likely to be initiators or targets of muzzle contacts. Knowledgeable individuals were more

likely to be targets and naïve individuals were more likely to be initiators. This shows strong

support that, in the presence of a novel resource, individuals use this behaviour to seek

specific information from others who demonstrate having that information. We do not make

claims related to knowing what others know, but rather assume that seeing a group member

eating an unknown resource prompts the initiation of muzzle contact towards that individual.

This result further corroborates the finding in 3a) of decreasing muzzle contact frequency

with increased exposure to the resource, and the overall function of muzzle contact in

soliciting foraging information.

Age was also a significant predictor, with adults more likely to be targets, and juveniles

more likely to be initiators. This concurs with previous findings of younger vervet monkeys

targeting adults with muzzle contacts during natural foraging [28,29], as well as our team's

previous study where adults were preferentially observed eating this novel food [20]. This

suggests adults are generally preferred sources of information and juveniles are not.

Moreover, age also interacted with knowledge, whereby becoming knowledgeable increased

adults' likelihood of becoming targets rather than initiators much more than it did for

juveniles, and naïve adults were still more likely to be targets than knowledgeable juveniles

(Fig. 3A, left). This suggests that juveniles are *not* preferred sources of information, even

when knowledgeable regarding a new resource, which may impede information flow.

Nonetheless, as discussed in section 2b), this impediment may be overcome by older or

higher-ranked juveniles. Indeed, we can expect high-ranked juveniles to gain access to

preferred resources over lower-ranked adults as well as lower-ranked juveniles.

Nevertheless, any age-related impediment to information flow is likely an adaptive

mechanism as juveniles' greater curiosity combined with lack of experience is risky; but their

reliance on adults to gain information presents as an adaptive mitigation to this risk.

Intriguingly, whilst we found no significant main effect of sex, sex did interact with

knowledge. Inspection of the plot in Fig. 3A (right) shows that whilst gaining knowledge

affected both males and females similarly, this effect appears to be slightly stronger in

males. Bearing in mind that juveniles (both males and females) are very unlikely to be

targets, as shown by both, the main effect of age, and the interaction of knowledge and age

(Fig. 3A, left), we can assume that this stronger effect in males is driven by adult males.

Indeed, by plotting the three-way interaction between knowledge, age and sex (Fig. 3B),

whilst non-significant, we can see the effect of knowledge is strongest for adult males, but it

is slightly stronger for juvenile males than females too. This likely drives the interaction

between knowledge and sex in the two-way interaction model (Fig. 3B) too. Nonetheless, the

biggest difference remains between adults and juveniles, and is similar for both sexes.

What remains, however, is that knowledge made adult males almost exclusively

targets and not initiators (Fig. 3B). Adult females, on the other hand, did still initiate muzzle

contacts, as seen by their more intermediate probability of being targets or initiators. Given

that our results show strong support for the function of muzzle contact in acquiring

information about novel resources, this adult sex difference is particularly interesting. That

adult females still initiate muzzle contacts, even when knowledgeable, suggests that they still

seek social information even when they have first-hand experience. This is in line with the

assertion that they are the most risk-averse age-sex category, given the dramatic effects of

risk-taking on their inclusive fitness, particularly with dependent offspring around or if

pregnant. Moreover, given that they are philopatric, their social environment is likely to be a

dependable and consistent source of reliable information. Contrastingly, it seems that adult

males, once acquiring their own knowledge of a resource, will rely on this alone without

initiating more muzzle contacts. This implies that they very quickly rely on their own

information, which arguably reflects boldness or low neophobia. This may be related to their

status as dispersers, given that during dispersal they must fend for themselves and have no

group-mates to rely on. Alongside results from our previous study [20], where males were

more likely to be observed extracting and eating peanuts, this shows support for the

potential role of dispersing male vervet monkeys in generating and transmitting novel

information within groups, and transferring it between groups.

Ideas and Speculation

Taking all of the evidence together, from this and other studies, we propose a

prominent role of dispersing male vervet monkeys in generating and transmitting novel

behavioural adaptations between groups. Dispersing adult males are more likely to generate

novel information than adult females, given the associated risks of doing so, and they are

more likely to be learned from quickly than explorative juveniles. Moreover, they are the only

ones that can transmit information between groups, which can allow population level

adaptation. This could help facilitate the species' widespread distribution and fast adaptation

to human-modified environments, similar to the breadth of that of many macaque and

baboon species (e.g. [45,46]), also living in multi-male multi-female groups. Spotted hyenas

(*Crocuta crocuta*) also live in large multi-male multi-female groups, unique among large

carnivores to maintain a similarly large distribution, further motivating greater attention to this

proposed hypothesis of the impact of regular immigration to fast adaptation for socially

learning species in a changing world.

Conclusion

We add to the literature an experimental example of exploitation of a novel resource by

multiple groups, facilitated here by dispersers. Our results provide evidence that dispersing

individuals may promote the generation of new, environmentally relevant information and its

spread around populations – a factor that has been largely overlooked, despite the known

role of dispersal in gene flow [47]. We urge future research to investigate whether

physiological mechanisms exist that might underpin a transitory dispersal syndrome

characterised by heightened exploration and reduced neophobia that is triggered during, or

triggers, dispersal. Moreover, we suggest further research, in diverse species, into whether

dispersers transmit valuable information between groups, as this can have major

implications for population fitness, especially in the context of the rapid anthropogenic

change that most animal populations now face. This can have further implications for short-

and long-term conservation efforts, informing wildlife management tasked with, for example,

the maintenance of wildlife corridors beyond the borders of groups to allow intergroup

transfer, or establishing most appropriate group composition for long term adaptation and

survival of groups to be reintroduced to the wild.

MATERIALS AND METHODS

Experimental model and subject details

The study was conducted at the 'Inkawu Vervet Project' (IVP) in a 12000-hectares

private game reserve: Mawana (28°00.327S, 031°12.348E) in KwaZulu Natal province,

South Africa. The biome of the study site is described in [21].

Five groups of habituated wild vervet monkeys (*Chlorocebus pygerythrus*) took part

in the study: 'Ankhase' (AK), 'Baie Dankie' (BD), 'Kubu' (KB), 'Lemon Tree' (LT) and 'Noha'

(NH). Habituation began in 2010 in AK, BD, LT and NH, and in 2013 in KB. All observers

were trained to identify individuals by individual bodily and facial features (eye-rings, scars,

colour, shape etc.). During the study period, these stable groups comprised between 19 and

65 individuals including infants (Table 1). We refer to the group AK differentially as AK_{19} and

AK_{20}, representing their status in 2019 and 2020, respectively, as 40% of the group

composition changed between years due to dispersals, deaths and changes in age

categories (infants that became juveniles; see Table S1).

Ethics guidelines: Our study adhered to the "Guidelines for the use of animals in

research" of Association for Study of Animal Behaviour and was approved by the relevant

local authority, Ezemvelo KZN Wildlife, South Africa.

Hierarchy establishment

Agonistic interactions (aggressor behaviour: stare, chase, attack, hit, bite, take place;

victim behaviour: retreat, flee, leave, avoid, jump aside) were collected *ad libitum* [48] on all

adults and juveniles of each group. These data were collected throughout one year

preceding the experiment during all behavioural observation hours and during experiments

involving food provisions. Data were collected by CC, PD and different trained observers

from the IVP team. Before beginning data collection, observers had to pass an inter-

observer reliability test with Cohen's kappa > .80 [43] for each data category between two

observers. Data were collected on handheld computers (Palm Zire 22) using Pendragon

software version 5.1 and, from the end of August 2017, on tablets (Vodacom Smart Tab 2)

and smartphones (Runbo F1) equipped with the Pendragon version 8.

Individual hierarchical ranks were determined by the outcome of dyadic agonistic

interactions using Socprog software version 2.7 [44]. Linearity of hierarchies are reported in

supplementary material (SM2). Ranks were assessed by I&SI method [45].

Peanut exposures

We provided each group with a highly nutritious novel food that required extraction

before consumption – unshelled peanuts – in large quantities to avoid monopolisation by

single individuals. Experiments took place after sunrise when the monkeys were located at

their sleeping site during the dry, food-scarce South African winter, to maximise motivation to

engage in food-rewarded experiments.

CC ran field experiments during May-June 2018 in KB and NH, and PD led the

experiments during August-September 2019 in AK (AK$_{19}$), BD and LT, and May-June 2020

in AK (AK$_{20}$; Table S1). Figure 1 illustrates the relevant male immigrations into and

emigrations out of these groups. *Avo* left his natal group KB to immigrate into NH two weeks

before their first experiment in 2018. *Avo* never ate peanuts before the first exposure in NH.

KB had no new males since 2017. *Pro* originated from NH and learned to eat peanuts during

their experiment in 2018. He immigrated into BD three weeks before their first experiment in

2019. *Bab* immigrated into LT six weeks before their first experiment in 2019, from an

unhabituated group, though he was habituated to humans due to previous residence in two

habituated study groups. *Bab* never ate peanuts before LT's first exposure. In 2020, two

males, *Twe* and *Yan*, who were present in NH during peanut exposures in 2018, immigrated

into AK, six and ten weeks, respectively, before their experiment in 2020. *Twe* ate peanuts in

NH during peanut exposures that continued beyond the four presented here for another

study [19], and *Yan* had observed many others eating peanuts. *Yan* was the first to eat in AK

in 2020 (Fig. 1).

Peanuts were presented to all groups in clear rectangular plastic boxes (34 x 14 x 12

cm), containing 1 – 2.5 kg of unshelled peanuts. One box of peanuts was offered per

exposure in AK_{20}, BD and LT, which lasted up to one hour, after which the monkeys usually

left the vicinity. In AK_{19}, the exposure of one box of peanuts lasted approximately five

minutes, after which the monkeys left the vicinity. In KB and NH, two boxes were offered

during each exposure, and were topped up when they were empty, for a maximum duration

of 2 hrs 45 minutes. KB and NH had 10 exposures on 10 different days; AK_{20}, BD, and LT

had four exposures on four different days; and AK_{19} had a single exposure (Table S1). Here

we present results for each group from the first four exposures *from the first eating event* in

each group.

Reactions to and interactions with peanuts were recorded by three to five observers

using handheld JVC video cameras (EverioR Quad Proof GZ-R430BE) and cameras

mounted on a tripod. Observers narrated identities of monkeys interacting with peanuts for

later video coding.

Quantification and statistical analysis

Video coding

To extract the identities of individuals who successfully extracted and ate peanuts

from their shells during each exposure, PD coded videos of AK_{19-20}, BD and LT with the

Windows 10 default video software, and CC coded videos of KB and NH with Media Player

Classic Home Cinema software version 1.7.11.

To analyse muzzle contacts in groups where many individuals started eating the

novel food quickly, GL counted frequencies of muzzle contacts in videos of AK_{20}, BD and LT.

PD and MC identified individuals involved in all muzzle contacts in AK_{20}, BD and LT.

Data analysis

Demographic variation in the uptake of the novel food

We used two generalised linear mixed models (GLMMs) for data following a binomial

distribution to investigate demographic variation in who extracted and ate peanuts (question

2b) i) at the first exposure (AK_{19}, BD, KB, LT & NH) and ii) over the four exposures (AK_{19-20},

BD, KB, LT & NH). In each model, we considered age, sex and rank (normalised rank) as

fixed effects and group was included as random effect.

Muzzle contact frequency over repeated exposure to novel food

We investigated muzzle contact behaviour in groups where large proportions of the

groups started to extract and eat peanuts over the first four exposures (AK_{20}, BD, LT). We

ran a linear regression model to investigate the effect of exposure to the novel food on

muzzle contact rate (question 3a). The normality of the residuals was verified on the basis of

the QQ plot shape to use a linear model. Muzzle contact rate was calculated as the total

number of muzzle contacts observed within 2 m of the food, during each exposure, divided

by the total duration (in minutes) of each exposure. Muzzle contact rate was the outcome

variable, with exposure number (1-4) as the predictor. For BD and LT we excluded

exposures 4 and 3, respectively, due to circumstances resulting in very small proportions of

these groups present at these exposures. We ran the same model including these outlier

exposures and present these results in the supplementary material (SM3).

Factors influencing whether individuals are initiators and targets of muzzle contact

Finally, we ran three binomial GLMMs to investigate question 3b) factors affecting

whether individuals involved in muzzle contacts were initiators or targets. Due to very low

numbers of infants ever being targets of muzzle contacts, we merged the infant and juvenile age categories for this analysis. Our first simple model included individuals' knowledge of the novel food at the time of each muzzle contact (knowledgeable = previously succeeded to extract and eat peanuts; naïve = never previously succeeded to extract and eat peanuts) and age, sex and rank as fixed effects. Individual was included as a random effect. The second model was the same, but we removed rank and added interactions between: knowledge and age; and knowledge and sex. The third model was the same as the second, but we also added a three-way interaction between knowledge, age and sex. We interpreted interactions by visual inspection of plots made using R package 'interactions' [49].

Statistics were computed in R Studio (R version 4.0.3), and linear regression was done with the base R stats package [50]. GLMMs were done using the 'lme4' package [51].

All data are made available at: https://doi.org/10.5281/zenodo.5767598.

Acknowledgments

This study was supported by the Swiss National Science Foundation (PP03P3_170624 and PP00P3 198913), the Branco Weiss Fellowship—Society in Science granted to Erica van de Waal, and by the Fyssen Foundation and the Fondation des Treilles granted to Charlotte Canteloup. We are grateful to the van der Walt family for their permission to conduct the study on their land and to Arend van Blerk and Michael Henshall for their support in the field. We are particularly thankful to Mabia Biff Cera, Adam Cogan, Adwait Deshpande, Sashimi Wieprecht, Manon Kerréveur-Lavaud, Varun Manavazhi, Maria Teresa Martinez Navarrete, Tecla Mohr, Aurora Rozmaryn, Claudia Seminara and Luca Silvestri for their help in data collection. We thank Redouan Bshary, Sofia Forss, Rachel Harrison, Carel van Schaik and Andrew Whiten for comments on an earlier version of the manuscript.

Author contributions: conceptualization: P.D., C.C. and E.v.d.W.; Methodology: P.D., C.C.

and E.v.d.W; Investigation: P.D. and C.C.; Video coding: C. C., P. D., G. L., M. C.; Formal

Analysis: C.C., P.D.; Visualization: C.C., P.D.; Writing – Original draft; P.D.; Writing –

Reviews & Editing: P.D., C.C. and E.v.d.W.; Funding acquisition: E.v.d.W. and C.C.

Supervision: E.v.d.W.

Declaration of interests: The authors declare no competing interests

REFERENCES

1.   Barrett LP, Stanton LA, Benson-Amram S. 2019 The cognition of 'nuisance' species.
Anim. Behav. **147**, 167–177. (doi:10.1016/j.anbehav.2018.05.005)

2.   Jesmer BR *et al.* 2018 Is ungulate migration culturally transmitted? Evidence of social
learning from translocated animals. *Science (80-.).* **361**, 1023–1025.
(doi:10.1126/science.aat0985)

3.   Gruber T, Luncz L, Mörchen J, Schuppli C, Kendal RL, Hockings K. 2019 Cultural
change in animals: a flexible behavioural adaptation to human disturbance. *Palgrave
Commun.* **5**, 1–9. (doi:10.1057/s41599-019-0271-4)

4.   Sih A, Ferrari MCO, Harris DJ. 2011 Evolution and behavioural responses to human-
induced rapid environmental change. *Evol. Appl.* **4**, 367–387. (doi:10.1111/j.1752-
4571.2010.00166.x)

5.   Jolles JW, King AJ, Killen SS. 2020 The Role of Individual Heterogeneity in Collective
Animal Behaviour. *Trends Ecol. Evol.* **35**, 278–291. (doi:10.1016/j.tree.2019.11.001)

6.   Forss SIF, Koski SE, van Schaik CP. 2017 Explaining the Paradox of Neophobic
Explorers: The Social Information Hypothesis. *Int. J. Primatol.* **38**, 799–822.
(doi:10.1007/s10764-017-9984-7)

7.   Carter AJ, Marshall HH, Heinsohn R, Cowlishaw G. 2012 How not to measure
boldness: Novel object and antipredator responses are not the same in wild baboons.
Anim. Behav. **84**, 603–609. (doi:10.1016/j.anbehav.2012.06.015)

8.   Kummer H, Goodall J. 1985 Conditions of innovative behaviour in primates. *Proc. R.
Soc. B Biol. Sci.* **308**, 203–214.

9.   McLennan MR, Hockings KJ. 2015 Wild chimpanzees show group differences in
selection of agricultural crops. *Sci. Rep.* **4**, 5956. (doi:10.1038/srep05956)

10.   Yamamoto ME, De Araújo Lopes F. 2004 Effect of removal from the family group on
feeding behavior by captive Callithrix jacchus. *Int. J. Primatol.* **25**, 489–500.
(doi:10.1023/B:IJOP.0000019164.98756.9C)

11.   Dally JM, Clayton NS, Emery NJ. 2008 Social influences on foraging by rooks (Corvus
frugilegus). *Behaviour* **145**, 1101–1124. (doi:10.1163/156853908784474470)

12.   Krakauer EB. 2005 Developement of aye-aye (Daubentonia madagascariensis)
foraging skills: Independent exploration and social learning. Duke University.

13.   Addessi E, Galloway AT, Visalberghi E, Birch LL. 2005 Specific social influences on
529   the acceptance of novel foods in 2-5-year-old children. *Appetite* **45**, 264–271.
530   (doi:10.1016/J.APPET.2005.07.007)

531   14.   Greggor AL, McIvor GE, Clayton NS, Thornton A. 2016 Contagious risk taking: social
532   information and context influence wild jackdaws' responses to novelty and risk. *Sci.*
533   *Rep.* **6**, 27764. (doi:10.1038/srep27764)

534   15.   Carter AJ, Marshall HH, Heinsohn R, Cowlishaw G. 2014 Personality predicts the
535   propensity for social learning in a wild primate. *PeerJ* **2**, e283. (doi:10.7717/peerj.283)

536   16.   O'Malley RC, Wallauer W, Murray CM, Goodall J. 2012 The appearance and spread
537   of ant fishing among the Kasekela chimpanzees of Gombe: A possible case of
538   intercommunity cultural transmission. *Curr. Anthropol.* **53**, 650–663.
539   (doi:10.1086/666943)

540   17.   Biro D, Inoue-Nakamura N, Tonooka R, Yamakoshi G, Sousa C, Matsuzawa T. 2003
541   Cultural innovation and transmission of tool use in wild chimpanzees: Evidence from
542   field experiments. *Anim. Cogn.* **6**, 213–223. (doi:10.1007/s10071-003-0183-x)

543   18.   Barrett BJ, McElreath RL, Perry SE. 2017 Pay-off-biased social learning underlies the
544   diffusion of novel extractive foraging traditions in a wild primate. *Proc. R. Soc. B Biol.*
545   *Sci.* **284**, 20170358. (doi:10.1098/rspb.2017.0358)

546   19.   Perry S *et al.* 2003 Social conventions in wild white-faced capuchin monkeys:
547   Evidence for traditions in a neotropical primate. *Curr. Anthropol.* **44**, 241–268.
548   (doi:10.1086/345825)

549   20.   Canteloup C, Cera MB, Barrett BJ, van de Waal E. 2021 Processing of novel food
550   reveals payoff and rank-biased social learning in a wild primate. *Sci. Rep.* **11**, 1–13.
551   (doi:10.1038/s41598-021-88857-6)

552   21.   Bono AEJ, Whiten A, Schaik C Van, Krutzen M, Eichenberger F, Schnider A, van de
553   Waal E. 2018 Payoff- and sex-biased social learning interact in a wild primate
554   population. *Curr. Biol.* **28**, 2800–2805. (doi:10.1016/j.cub.2018.06.015)

555   22.   van de Waal E, Renevey N, Favre CM, Bshary R. 2010 Selective attention to
556   philopatric models causes directed social learning in wild vervet monkeys. *Proc. R.*
557   *Soc. B Biol. Sci.* **277**, 2105–2111. (doi:10.1098/rspb.2009.2260)

558   23.   Grampp M, Sueur C, van de Waal E, Botting J. 2019 Social attention biases in
559   juvenile wild vervet monkeys: implications for socialisation and social learning
560   processes. *Primates* **60**, 261–275. (doi:10.1007/s10329-019-00721-4)

561   24.   Whiten A. 2019 Social Learning: Peering Deeper into Ape Culture. *Curr. Biol.* **29**,
562   R845–R847. (doi:10.1016/j.cub.2019.07.003)

563   25.   Laidre ME. 2009 Informative Breath: Olfactory Cues Sought During Social Foraging
564   Among Old World Monkeys (Mandrillus sphinx, M. leucophaeus, and Papio anubis). *J.*
565   *Comp. Psychol.* **123**, 34–44. (doi:10.1037/a0013129)

566   26.   Drapier M, Chauvin C, Thierry B. 2002 Tonkean macaques (Macaca tonkeana) find
567   food sources from cues conveyed by group-mates. *Anim. Cogn.* **5**, 159–165.
568   (doi:10.1007/s10071-002-0145-8)

569   27.   Chauvin C, Thierry B. 2005 Tonkean macaques orient their food search from olfactory
570   cues conveyed by conspecifics. *Ethology* **111**, 301–310. (doi:10.1111/j.1439-
571   0310.2004.01066.x)

572   28.   Nord CM, Bonnell TR, Dostie MJ, Henzi SP, Barrett L. 2021 Tolerance of muzzle

contact underpins the acquisition of foraging information in vervet monkeys (Chlorocebus pygerythrus). *J. Comp. Psychol.* **135**, 349–359. (doi:10.1037/com0000258)

29. Lycett JE, Henzi PS. 1992 The relevance of sensory modality to the social transmission of information in vervet monkeys (Cercopithecus aethiops). *South African J. Sci.* **88**, 177–179.

30. Turner TR, Schmitt CA, Cramer JD. 2019 *Savanna Monkeys*. Cambridge University Press. (doi:10.1017/9781139019941)

31. Fairbanks LA. 1993 Risk-taking by juvenile vervet monkeys. *Behaviour* **124**, 57–72.

32. Benson-Amram S, Weldele ML, Holekamp KE. 2013 A comparison of innovative problem-solving abilities between wild and captive spotted hyaenas, Crocuta crocuta. *Anim. Behav.* **85**, 349–356. (doi:10.1016/j.anbehav.2012.11.003)

33. Bergman TJ, Kitchen DM. 2009 Comparing responses to novel objects in wild baboons (Papio ursinus) and geladas (Theropithecus gelada). *Anim. Cogn.* **12**, 63–73. (doi:10.1007/s10071-008-0171-2)

34. Miller R, Bugnyar T, Pölzl K, Schwab C. 2015 Differences in exploration behaviour in common ravens and carrion crows during development and across social context. *Behav. Ecol. Sociobiol.* **69**, 1209–1220. (doi:10.1007/s00265-015-1935-8)

35. Sherratt TN, Morand-Ferron J. 2018 The adaptive significance of age-dependent changes in the tendency of individuals to explore. *Anim. Behav.* **138**, 59–67. (doi:10.1016/j.anbehav.2018.01.025)

36. Cote J, Clobert J, Brodin T, Fogarty S, Sih A. 2010 Personality-dependent dispersal: Characterization, ontogeny and consequences for spatially structured populations. *Philos. Trans. R. Soc. B Biol. Sci.* **365**, 4065–4076. (doi:10.1098/rstb.2010.0176)

37. Kaplan JR, Fontenot MB, Berard J, Manuck SB, Mann JJ. 1995 Delayed dispersal and elevated monoaminergic activity in free-ranging rhesus monkeys. *Am. J. Primatol.* **35**, 229–234. (doi:10.1002/ajp.1350350305)

38. Fairbanks LA, Melega WP, Jorgensen MJ, Kaplan JR, Mcguire MT. 2001 Social impulsivity inversely associated with CSF 5-HIAA and fluoxetine exposure in vervet monkeys. *Neuropsychopharmacology* **24**, 370–378.

39. Hansenne M, Ansseau M. 1999 Harm avoidance and serotonin. *Biol. Psychol.* **51**, 77–81. (doi:10.1016/S0301-0511(99)00018-6)

40. Blaszczyk MB. 2017 Boldness towards novel objects predicts predator inspection in wild vervet monkeys. *Anim. Behav.* **123**, 91–100. (doi:10.1016/j.anbehav.2016.10.017)

41. Nord CM. 2021 The contexts of social learning in wild vervet monkeys. University of Lethbridge.

42. L'Allier S. 2020 Behavioral Endocrinology of Male Dispersal in Vervet Monkeys at Lake Nabugabo , Uganda. York University.

43. Sih A, Cote J, Evans M, Fogarty S, Pruitt J. 2012 Ecological implications of behavioural syndromes. *Ecol. Lett.* **15**, 278–289. (doi:10.1111/J.1461-0248.2011.01731.X)

44. Canteloup C, Hoppitt W, van de Waal E. 2020 Wild primates copy higher-ranked individuals in a social transmission experiment. *Nat. Commun.* **11**, 1–10. (doi:10.1038/s41467-019-14209-8)

45.　Fischer J *et al.* 2019 Insights into the evolution of social systems and species from
baboon studies. *Elife* **8**, 1–16. (doi:10.7554/eLife.50989)

46.　Thierry B. 2007 Unity in diversity: Lessons from macaque societies. *Evol. Anthropol.*
16, 224–238. (doi:10.1002/evan.20147)

47.　Greenwood PJ. 1980 Mating systems, philopatry and dispersal in birds and mammals.
Anim. Behav. **28**, 1140–1162.

48.　Altmann J. 1974 Observational Study of Behavior: Sampling Methods. *Behaviour* **49**,
227–267.

49.　Long JA. 2019 interactions: Comprehensive, User-Friendly Toolkit for Probing
Interactions.

50.　R Core Team. 2020 R: A language and environment for statistical computing.

51.　Bates D, Mächler M, Bolker B, Walker S. 2015 Fitting Linear Mixed-Effects Models
Using lme4. *J. Stat. Softw.* **67**, 1–48. (doi:10.18637/jss.v067.i01)

Tables and figures

Table 1. Cumulative numbers of monkeys eating at each exposure in each group, with a

total of 81 (of a possible 164) monkeys eating across the whole experiment.

Group	Group size	Exposure number						Total
		1	2	3	4	5	6	
AK	~20	0*	5**	13**	17**	19**	-	**19**
BD	65	19	25	29	32	-	-	**32**
KB	19	0	0	1	3	3	3	**3**
LT	25	5	13	16	21	-	-	**21**
NH	35	3	3	3	6	-	-	**6**
						Grand total =		**81**

NB. *AK in 2019; **AK in 2020

Figure 1. Aerial view of study area with coloured shapes showing a rough estimate of group

home-ranges for study groups AK, BD, KB, LT and NH. White arrows with annotations

represent relevant dispersals. Names of males and year of dispersal are shown. Black

outlined text indicates the immigrant innovators who were naïve to peanuts, solid white text

shows the immigrants that imported innovations, and white outlined text shows: parallel

dispersal with innovator (*Yan* 2018); that the innovator was habituated in a study group prior

to participation in this experiment (*Bab* 2016); or that the innovator left the study group (*Avo*

2018). Question mark shows that males dispersed to an unstudied group.

Figure 2. Uptake of extracting and eating peanuts in each group. A) Shows the proportion of

each group that started eating when the first eating event took place. Total numbers of

individuals, split by age, are shown below the x-axis. In NH the asterisk highlights the one

adult which was the innovator male that left after their first exposure, and was only followed

by juveniles learning to extract and eat. B) Shows the progression in each group over four

exposures from the first eating event. Solid squares represent when immigrant males were

the first to eat, and the open square shows when the infant was first to eat. Solid lines show

when there were adults present who had started to eat, whereas dashed lines show when

there were only juveniles and infants present that had already started to eat. Males who

were knowledgeable and imported innovations from other groups (*Pro* in BD and *Yan* in AK)

are excluded from totals in both panels (visualised in Microsoft Excel).

Figure 3. Plots illustrating significant interactions between the effects of A) left: knowledge

and age, and right: knowledge and sex on the likelihood of individuals being initiators rather

than targets of muzzle contact; and B) shows the (non-significant) three-way interaction

between knowledge, age and sex. Solid circles show predicted probabilities, and extending

bars show 95% confidence intervals (visualised using 'cat_plot' function from R package

'interactions' [49]). *N* is the number of muzzle contact interactions engaged in by each

age/sex category).