Title: Emergence of compensatory mutations reveal the importance of electrostatic

interactions between HIV-1 integrase and genomic RNA

Short title: Molecular basis of HIV-1 integrase-gRNA interactions

Authors: Christian Shema Mugisha[1], Tung Dinh[2], Abhishek Kumar[1], Kasyap Tenneti[1], Jenna E.

Eschbach[1], Keanu Davis[1], Robert Gifford[3], Mamuka Kvaratskhelia[2], Sebla B. Kutluay[1]

[1] Department of Molecular Microbiology, Washington University School of Medicine, Saint Louis,

MO 63110, USA

[2] Division of Infectious Diseases, University of Colorado School of Medicine, Aurora, CO 80045

[3] MRC-University of Glasgow Centre for Virus Research, 464 Bearsden Rd., Bearsden,

Glasgow G61 1QH, UK.

Correspondence: kutluay@wustl.edu

ABSTRACT

HIV-1 integrase (IN) has a non-catalytic function in virion maturation through its binding to the

viral RNA genome (gRNA). Allosteric integrase inhibitors (ALLINIs) and class II IN substitutions

inhibit IN-gRNA binding and result in non-infectious viruses marked by mislocalization of the

gRNA within virions. HIV-1 IN utilizes basic residues within its C-terminal domain (CTD) to bind

to the gRNA. However, the molecular nature of how these residues mediate gRNA binding and

whether other regions of IN are involved remain unknown. To address this, we have isolated

compensatory substitutions in the background of a class II IN mutant virus bearing

R269A/K273A substitutions within the IN-CTD. We found that the nearby D256N and D270N

compensatory substitutions restored the ability of IN to bind gRNA and led to the formation of

mature infectious virions. Reinstating the local positive charge of the IN-CTD through individual

D256R, D256K, D278R and D279R substitutions was sufficient to restore IN-RNA binding and

infectivity for the IN R269A/K273A as well as the IN R262A/R263A class II mutants. Structural

modeling suggested that compensatory substitutions in the D256 residue created an additional

interaction interface for gRNA binding. Finally, HIV-1 IN R269A/K273A, but not IN

R262A/R263A, bearing compensatory mutations was more sensitive to ALLINIs providing key

genetic evidence that specific IN residues required for RNA binding also influence ALLINI

activity. Taken together, our findings highlight the essential role of CTD in gRNA binding and

ALLINI sensitivity, and reveal the importance of pliable electrostatic interactions between the IN-

CTD and the gRNA.

IMPORTANCE

In addition to its catalytic function, HIV-1 integrase (IN) binds to the viral RNA genome (gRNA)

through positively charged residues within its C-terminal domain (CTD) and regulates proper

virion maturation. Here we show that compensatory mutations in nearby acidic residues (i.e.

D256N and D270N) restore the ability to bind gRNA for IN variants bearing substitutions in

these positively charged CTD residues. Similarly, charge reversals through individual D-to-R

and D-to-K substitutions at these positions enabled the respective IN mutants to bind gRNA and

restore virion infectivity. Further, we show that specific residues within the IN-CTD required for

RNA binding also influence sensitivity to allosteric integrase inhibitors, a class of novel IN-

targeting compounds that target the non-catalytic function of IN. Taken together, our findings

reveal the importance of electrostatic interactions in IN-gRNA binding and provide key evidence

for a crucial role of the IN-CTD in allosteric integrase inhibitor mechanism of action.

INTRODUCTION

A defining feature of retroviruses is the reverse transcription of the viral RNA genome (gRNA) and integration of the resultant linear viral DNA into the host chromosome, which establishes lifelong infection. The latter reaction is mediated by the viral integrase (IN) enzyme, which catalyzes 3' processing and DNA strand transfer reactions (1). The catalytic activity of HIV-1 IN has been successfully targeted by several integrase strand-transfer inhibitors (INSTIs) (2-7) that have become key components of frontline anti-retroviral therapy regimens due to their high efficacy and tolerance profiles (8-11). In addition, HIV-1 IN has a non-catalytic function in virus replication (12-16). Successful targeting of this second function can complement the existing antiviral regimens and substantially increase the barrier to INSTI resistance.

HIV-1 IN consists of three independently folded protein domains: the N-terminal domain (NTD) bears the conserved His and Cys residues (HHCC motif) that coordinate Zn^{2+} binding for 3-helix bundle formation; the catalytic core domain (CCD) adopts an RNase H fold and harbors the enzyme active site composed of an invariant DDE motif, and C-terminal domain (CTD) which adopts an SH3 fold (17, 18). Integration is facilitated by a cellular co-factor, lens epithelium-derived growth factor (LEDGF/p75), which binds tightly to a site within the CCD dimer interface (19, 20) and guides the preintegration complex to actively transcribed regions of the host chromosome (19-24). A group of pleotropic IN substitutions distributed throughout IN, collectively known as class II mutations, disrupt viral assembly (13, 16, 25-36), morphogenesis (12, 16, 27, 32-34, 37, 38) and reverse transcription in target cells (12-14, 16, 29, 31, 32, 34, 36-54) often without obstructing the catalytic activity of IN *in vitro* (13, 27, 28, 31, 40, 41, 44, 46, 55-57). A hallmark of class II IN mutant viruses is the mislocalization of the viral ribonucleoprotein complexes (vRNP) outside of the viral capsid (CA) lattice, a deformation which is often referred

to as eccentric morphology (12, 15, 16, 27, 32-34, 37, 38, 58-60). Although originally designed

to inhibit integration through preventing the binding of IN to LEDGF/p75 (58, 61-65), allosteric

integrase inhibitors (ALLINIs) potently inhibit proper virion maturation (12, 37, 61) and lead to

the formation of virions that display a similar eccentric morphology observed with class II IN

mutations (12).

We have recently shown that IN binds to the gRNA in mature virions to mediate proper

encapsidation of the viral ribonucleoproteins (vRNPs) inside the mature CA lattice (38). IN binds

with high affinity to the TAR hairpin present within the 5' and 3'UTRs, though multiple other

distinct locations on the gRNA without apparent secondary structure are also bound (38). A

parameter that appears to be critical for gRNA binding is functional tetramerization of IN.

Numerous class II IN mutations scattered throughout IN block IN-gRNA binding indirectly by

disrupting functional oligomerization of IN (15). Similar to these class II IN mutations, ALLINIs

are thought to interfere with IN-gRNA binding indirectly by inducing aberrant IN multimerization

(12, 38). ALLINIs bind to a pocket within the CCD of IN (12, 60, 61, 64, 66-68) and induce the

formation of open IN polymers by engaging the CTD of a nearby IN dimer (62). In contrast,

mutation of basic residues within IN-CTD (i.e. R262, R263, R269 and K273) inhibits IN-gRNA

binding without altering IN oligomerization in virions and *in vitro* (15, 38), suggesting that these

residues directly mediate IN binding to the gRNA.

While the involvement of IN-CTD in gRNA binding has been established, how the basic residues

within IN-CTD mediate recognition of the gRNA remain unknown. For example, it is possible

that the positively charged Lys and Arg residues interact with the negatively charged RNA

phosphate backbone in a non-specific or semi-specific manner, depending on the folding and

structure of the cognate RNA element, driven by electrostatic interactions (69-71). Alternatively,

these residues can mediate specific interactions with gRNA through H-bonding and van der

Waals contacts with individual nucleobases (69-71). Furthermore, it is possible that other

domains in IN also contribute to gRNA binding. For example, we have previously shown that

K34A substitution within the NTD blocks gRNA binding without impacting IN oligomerization

(15), suggesting its direct interaction with the gRNA.

To gain insight into the mode of IN-gRNA interactions, the non-infectious HIV-1$_{NL4\text{-}3 \text{ IN (R269A/K273A)}}$

class II IN mutant virus was serially passaged in T-cells until the acquisition of compensatory

mutations. Two compensatory substitutions, D256N and D270N, sequentially emerged within

the IN-CTD and allowed HIV-1$_{NL4\text{-}3 \text{ IN (R269A/K273A)}}$ to replicate with WT kinetics. Introduction of

these substitutions on the HIV-1$_{NL4\text{-}3 \text{ IN (R269A/K273A)}}$ virus backbone substantially increased virion

infectivity, restored IN-gRNA binding and proper virion morphogenesis. As the D-to-N mutations

resulted in loss of two negative charges, possibly overcoming the loss of two positive charges

with the R269A/K273A substitutions, we tested whether restoring the overall local charge of IN-

CTD would also restore IN-gRNA binding and virion infectivity. Indeed, the D256R as well as

other nearby charge reversal substitutions including D278R and D279R enhanced or restored

virion infectivity and RNA binding for the IN R269A/K273A mutant. We further extended these

findings to another class II IN mutant, R262A/R263A, which was similarly suppressed by the

D256R as well as D256K substitutions. Compensatory mutations did not affect the ability of IN

to multimerize *in vitro* or in virions, suggesting that they restored the RNA-binding ability of IN

directly. Structural modeling suggested that compensatory mutations created an additional

interaction interface for gRNA binding. Finally, the replication ability of the class II IN mutant

viruses bearing the compensatory substitutions allowed us to test the contribution of specific

CTD residues to ALLINI mechanism of action. Specifically, we found that viruses bearing

R269A/K273A, but not the R262A/R263A, substitutions were significantly more sensitive to

ALLINIs. Together, our findings highlight the essential role of IN-CTD in gRNA binding, reveal

an electrostatic component of IN-gRNA interactions, and provide key genetic evidence that

ALLINIs engage specific residues within the IN-CTD.

RESULTS

Compensatory IN D256N/D270N substitutions emerge in the background of HIV-1$_{\text{NL4-3 IN}}$

$_{\text{(R269A/K273A)}}$ class II IN mutant virus

The IN R269A/K273A class II substitutions obstruct IN-gRNA binding directly without interfering with IN multimerization and result in the formation of particles with eccentric morphology (15, 16). To gain insight into the molecular basis of how R269 and K273 residues mediate IN-gRNA binding, HIV-1$_{\text{NL4-3 IN (R269A/K273A)}}$ class II IN mutant virus was serially passaged in MT-4 cells until the emergence of compensatory mutations at the end of passage 3, which resulted in virus growth with WT replication kinetics (**Fig. 1A**). Deep sequencing of full-length gRNA isolated from virions across the three passages revealed that the IN R269A and K273A substitutions were retained while the nearby D256N and D270N mutations within IN-CTD were sequentially acquired at the end of passage 1 and passage2, respectively (**Fig. 1B**). Both mutations were fixed by the end of passage 2 and no other non-synonymous mutations were observed in IN or elsewhere on the gRNA.

The IN D256N and D270N mutations were introduced into the replication-competent pNL4-3 HIV-1 molecular clone bearing WT IN or IN R269A/K273A. Introduction of D256N and D270N substitutions in IN had no observable effect on Gag (Pr55) and Gag-Pol expression or processing in cells, particle release and virion-associated IN levels (**Fig. 1C, S1A-F**). Introduction of D256N and D270N mutations either individually or together (D2N) into HIV-1$_{\text{NL4-3}}$ bearing WT IN did not affect viral titers (**Fig. 1D, S1G**). Remarkably, while the individual D256N and D270N mutations introduced into the HIV-1$_{\text{NL4-3 IN (R269A/K273A)}}$ class II IN mutant virus increased virus titers by approximately 4-fold, the D2N substitutions increased virus titers by 30-fold (**Fig. 1D, S1G**). Despite no observable differences in multi-cycle replication kinetics (**Fig.**

1A), HIV-1$_{NL4-3\ IN\ (R269A/K273A/D2N)}$ virus was modestly (~4-fold) but significantly less infectious than

WT in single cycle replication assays (**Fig. 1D, S1G**). Overall, these results demonstrate that the

combination of D256N and D270N mutations is sufficient to substantially increase the replication

competency of the HIV-1$_{NL4-3\ IN\ (R269A/K273A)}$ class II IN mutant virus.

IN D256N/D270N substitutions restore IN-gRNA binding for the HIV-1$_{NL4-3\ IN\ (R269A/K273A)}$ virus

and lead to the formation of correctly matured virions

We next assessed whether the IN D256N and IN D270N substitutions rendered the HIV-1$_{NL4-3\ IN}$

$_{(R269A/K273A)}$ virus replication competent by restoring IN-gRNA binding and proper virion

maturation. To this end, IN-gRNA complexes were immunoprecipitated from UV-crosslinked

virions and visualized per CLIP protocol as described previously (38, 72, 73). IN-RNA

complexes were readily visible for viruses bearing WT IN and the D256N, D270N and D2N

substitutions introduced on the WT IN backbone did not impact RNA binding (**Fig. 2A, S2**).

Expectedly (15, 38), the IN R269A/K273A class II mutant had substantially lower levels of IN-

gRNA complexes isolated from virions (**Fig. 2A, S2**). The D256N substitution modestly

enhanced the ability of IN R269A/K273A to bind RNA, whereas the D270N substitution had no

observable impact (**Fig. 2A, S2**). In contrast, the D2N substitution substantially enhanced the

ability of IN R269A/K273A to bind RNA (**Fig. 2A, S2**). The virion morphology of WT and IN

mutant viruses was assessed by transmission electron microscopy (TEM). As expected, more

than 80% of WT particles had an electron-dense condensate that represents vRNPs inside the

CA lattice, whereas the majority of IN R269A/K273A class II mutant virions (~68%) had a clear

eccentric morphology (**Fig. 2B**). Consistent with effects on virus titers and RNA-binding, the

introduction of D2N substitutions largely restored the ability of the HIV-1$_{NL4-3\ IN\ (R269A/K273A)}$ virus to

form properly matured virions (**Fig. 2B**). Cumulatively, these data show that D256N and D270N

substitutions restore infectivity for the HIV-1$_{NL4-3\ IN\ (R269A/K273A)}$ class II IN mutant virus through

reestablishing gRNA binding and thereby enabling proper virion maturation.

A single charge reversal substitution alone enhances IN-gRNA binding and virion infectivity for the HIV-1$_{NL4-3\ IN\ (R269A/K273A)}$ virus

IN-CTD is decorated with several acidic and basic amino acids, and a long stretch of basic residues spanning 258-273 residues is notable (**Fig. 3A**). In effect, the D2N substitutions restored the overall local charge of this region, suggesting that electrostatic interactions may be a key parameter in gRNA binding. To test this hypothesis, we investigated whether restoring the overall charge of IN-CTD through other mutations would suppress the class II phenotype observed with HIV-1$_{NL4-3\ IN\ (R269A/K273A)}$. We focused our initial analysis on D256 and D270 residues for the following reasons: i) these amino acids were amenable to substitutions during virus passaging experiments; ii) D256N and D270N substitutions in the context of WT HIV-1 yielded infectious virions, suggesting that these compensatory mutations do not significantly contribute to the catalytic activity of IN thus allowing us to specifically probe their roles for IN-gRNA interactions and virion maturation. We introduced D256R, D270R and D256R/D270R (D2R) substitutions into the HIV-1$_{NL4-3\ IN\ R269A/K273A}$ backbone and transfected HEK293T cells with the resulting plasmids. Cell lysates and cell-free virions were then analyzed for Gag and Gag-Pol processing, particle release, and infectivity. Overall, D-to-R substitutions had no major effect on Gag or Gag-Pol expression and processing in cells, or virion release and virion-associated IN levels (**Fig. 3B, S3A-F**).

Remarkably, the D256R substitution alone increased the titers of HIV-1$_{NL4-3\ IN\ (R269A/K273A)}$ by 26-fold to a level comparable with the D2N substitution (**Fig. 3C, S3G**). In contrast, the D270R substitution did not increase virus titers and the D2R substitution increased the infectivity of HIV-1$_{NL4-3\ IN\ (R269A/K273A)}$ relatively modestly (~13-fold) compared to D256R and D2N substitutions (**Fig. 3C, S3G**). Both the D256R and the D2R substitutions restored IN-gRNA binding to WT levels (**Fig. 3D, S3H**), demonstrating that the increase in viral titers with the D256R and D2R substitutions correlates well with enhancement of IN-gRNA binding. To determine whether the

D256N/R and D256N/R substitutions affected the catalytic activity of IN, we conducted a

transcomplementation assay that relies on incorporation of class II IN mutants into a class I IN

mutant virus (HIV-1$_{NL4-3\ IN\ (D116N)}$) bearing a catalytically inactive IN through fusion to the viral Vpr

protein (74). In this assay, the RNA-binding defect of a class II IN mutant is complemented *in*

trans by the catalytically inactive IN, whereas the class II IN mutant can complement the

catalytic activity defect of the class I mutant. We found that none of the IN substitutions

impacted the catalytic activity of IN, except for the D2R substitution (**Fig. 3E**), suggesting that

the inability of the D270R mutation to restore infectivity to WT levels may be due to its adverse

effects on the catalytic activity of IN. Taken together, these findings indicate that restoring the

overall charge of IN-CTD through D256R mutation reestablishes virion infectivity and RNA

binding for the IN R269A/K273A class II mutant.

Charge reversal substitutions suppress the replication defect of a separate class II IN

mutant virus through restoring gRNA binding

Mutation of other basic residues within the IN-CTD, such as R262A/R263A, also directly inhibit

IN-gRNA binding without compromising functional oligomerization of IN (15, 38). If IN-gRNA

binding is mediated by electrostatic interactions, we reasoned that substitutions of D256 and

D270 residues in a way that restores the local positive charge should restore IN-gRNA binding

and infectivity in the background of HIV-1$_{NL4-3\ IN\ (R262A/R263A)}$. D256N, D256K, D256R, D270N and

D2N substitutions introduced into HIV-1$_{NL4-3\ IN\ (R262A/R263A)}$ did not have a major impact on Gag

and Gag-Pol expression, though a modest increase in distinct Gag processing intermediates

was observed (**Fig. 4A, C, S4A-D, S4G-J**). Notwithstanding, neither virion release nor virion-

associated IN levels were impacted (**Fig. 4A, C, S4E, S4K, S4L**). IN D256N and D2N

substitutions increased virion infectivity by 10- and 5-fold, respectively, whereas the D270N

substitution had no impact (**Fig. 4B, S4F**). In contrast, the D256K substitution increased viral

titers at a greater degree than D256N and the D256R substitution completely restored virion

infectivity (**Fig. 4D, S4M**). In line with the titer data, D256K significantly increased IN-gRNA

binding whereas the D256R completely restored it in the context of the HIV-1$_{NL4-3\ IN\ (R262A/R263A)}$

class II mutant virus (**Fig 4E, S4N**). D256N, D256K and D256R IN successfully

transcomplemented a class I IN mutant (D116N), suggesting that they did not distort the

catalytic activity of IN (**Fig 4F**).

Charge reversal substitutions at acidic residues other than D256 can suppress the class

II phenotype

Given the above findings, we next determined whether charge reversal substitutions at other

nearby acidic residues could restore IN-gRNA binding and infectivity for the HIV-1$_{NL4-3\ IN}$

$_{(R269A/K273A)}$ virus. Remarkably, the D279R substitution increased virus infectivity by 18-fold and

the D278R/D279R substitutions restored infectivity to WT levels (**Fig. 5A**). We also determined

whether substitutions of the original D256 and D270 residues into Ile would restore virus titers at

levels similar to the D-to-N substitutions. However, we found that neither D256I alone nor the

D256I/D270I (D2I) substitutions increased viral titers (**Fig. 5A**). Of note, these substitutions did

not impact Gag and Gag-Pol expression, processing, virion release or virion-associated IN

levels, though we noted the presence of an aberrantly processed IN with the D2I mutant (**Fig.**

5B). In line with the titer data, D279R substitution increased and the D2R substitution restored

IN-RNA binding for the HIV-1$_{NL4-3\ IN\ (R269A/K273A)}$ virus (**Fig. 5C**). Taken together, these results

demonstrate that restoring the local positive charge of IN-CTD is sufficient to restore virion

infectivity and RNA binding to two distinct class II IN mutants, which suggests the importance of

electrostatic interactions in mediating IN-gRNA binding.

Compensatory substitutions create an alternative interaction surface for RNA binding

We next used reported X-ray structure (62) and molecular modelling to visualize how class II

and compensatory IN mutations affect the electrostatic potential of the CTD surface. R269A and

K273A substitutions expectedly resulted in a substantial loss of a basic patch in IN (**Fig. 6A, B**).

Both the D256R and D256N substitutions resulted in a more positively charged surface distal

from the R269/K273A residues (**Fig. 6C, D**), suggesting that compensatory mutations likely

created an additional interaction interface for gRNA binding. A similar outcome was observed

when class II R262A/R263A and compensatory D256R changes were introduced in the CTD of

IN (**Fig. 6E-G**). While the D256R change is substantially distanced from R262/R263 and

R269/K273 residues, we note the following. R262, R263, R269, and K273 are positioned within

the same highly flexible C-terminal tail (aa 261-275), whereas D256 belongs to another, shorter

(aa 252-257) loop (**Fig. 6H**). The highly pliable nature of the tail and the loop could be crucial for

IN to optimally engage with cognate RNA as well as allow for emergence of compensatory

mutations at alternative sites positioned in these IN segments.

Compensatory mutations directly restore the ability of IN to bind vRNA without altering

functional IN oligomerization

As functional IN oligomerization is a prerequisite for RNA binding (15), we next examined how

the compensatory substitutions affected IN oligomerization. For in virion analysis, purified HIV-

1_{NL4-3} virions (WT and bearing the relevant IN substitutions) were treated with ethylene glycol bis

(succinimidyl succinate) (EGS) to covalently crosslink IN in situ and virus lysates were analyzed

by immunoblotting. As in WT virions, IN species that migrated at molecular weights consistent

with those of monomers, dimers, trimers, and tetramers were readily distinguished in HIV-1_{NL4-3}

IN (R269A/K273A) and HIV-1_{NL4-3} IN (R262A/R263A) viruses with additional compensatory mutations

(D256N/D270N, D256R, D256K, and D256R/D270R) but not with the canonical class II IN

mutant V165A that is unable to form functional oligomers (**Fig 7A-C** & (15)). Complementary in

vitro assessment of purified recombinant IN proteins by size exclusion chromatography (SEC)

revealed that D2N and D256R substitutions in the background of IN R269A/K273A or the IN

R262A/R263A class II mutants did not impact functional IN tetramerization (**Fig. 7D-H**).

We have previously shown that recombinant IN binds to TAR RNA with high affinity and

provides a nucleation point to bridge and condense RNA (38). We next examined the ability of

class II IN mutants bearing compensatory mutations to bind and bridge TAR RNA. Consistent

with findings from CLIP, the D2N and D256R substitutions also enhanced or restored the ability

of IN R269A/K273 and IN R262A/R263A mutants to bridge between RNA molecules (**Fig. 7I**).

Together, these data demonstrate that the compensatory mutations directly restore the ability of

IN to bind RNA without altering functional IN oligomerization.

Sensitivity to ALLINIs is determined by distinct residues within the CTD

ALLINIs potently disrupt proper virion maturation through inducing aberrant IN multimerization

and consequently inhibiting IN-gRNA binding (12, 38). Recent structural studies have shown

that the quinoline-based ALLINI, GSK-1264, can directly engage residues within the IN-CTD

through its tert-butoxy and carboxylic acid moieties and induce open IN polymers (62). These

findings are in line with previous biochemical and modeling studies that also showed the

involvement of the IN-CTD, in particular residues K264 and K266, in ALLINI-induced aberrant IN

multimerization (75, 76). Based on these prior findings, we wanted to test how adjacently

positioned IN R262A/R263A and R269A/K273A substitutions and the compensatory mutations

affected the antiviral activities of ALLINI.

To this end, we examined effects of representative quinoline-based ALLINIs BI-D and BI-B2,

which share many chemical features with GSK-1264, on the viruses bearing the class II and

compensatory IN mutations. While the titers of WT viruses only decreased at ALLINI

concentrations greater than 1 μM, the titers of HIV-1$_{NL4-3\ IN\ (R269A/K273A)}$ viruses bearing D2N,

D256R and D2R substitutions were reduced by 10-fold at ALLINI concentrations as low as 0.1

μM (**Fig. 8A, B**). In contrast, viruses bearing IN R262A/R263A/D256R were less sensitive to low

concentrations of ALLINIs (**Fig. 8D, E**), suggesting that the IN R269A and/or K273A, but not the

compensatory substitutions, underlie the increased sensitivity to ALLINIs. The increased

sensitivity to ALLINIs correlated well with inhibition of RNA binding. While 0.1 µM of BI-D didn't

affect the level of IN-gRNA binding in WT HIV-1, it significantly reduced IN-gRNA binding in

HIV-1$_{NL4-3\ IN\ (R269A/K273A)}$ but not HIV-1$_{NL4-3\ IN\ (R262A/K263A)}$ bearing compensatory mutations (**Fig.**

8C,F). Taken together, our findings provide key genetic, biochemical and virological evidence

that specific CTD residues required for RNA binding are also crucial for the ALLINI mechanism

of action.

Characterization of IN mutations present in latently infected cells

Persistence of HIV-1 in memory CD4+ T-cells as latent proviruses constitutes a major barrier to

HIV-1 cure. Although the majority of HIV-1 proviruses in these cells are defective (77), recent

evidence suggests that defective proviruses can be transcribed into RNAs that are spliced,

translated and can be recognized by HIV-1-specific cytotoxic T lymphocytes (78). We decided to

characterize IN mutations isolated from latently infected cells, given the possibility that class II

IN mutations existing in latently infected cells can result in the formation of defective particles

that may subsequently modulate immune responses. Though relatively uncommon, we found

the presence of R224Q, S230N, E246K and G272R substitutions in IN-CTD (**Fig. 9A**). Of note,

only the R224Q substitution resulted in loss of a positive charge, whereas the E246K and

G273R substitutions resulted in gain of positive charges. These mutations were introduced into

the NL4-3 proviral backbone with minimal effects on Gag expression and particle release (**Fig.**

9B). Although the E246K virus was significantly less infectious (**Fig. 9C**), we did not find any

evidence for loss of IN-gRNA binding (**Fig. 9D**), suggesting that this mutant likely displays a

class I phenotype. Thus, we conclude that the class II mutant viruses are rarely present in the

latently infected cells and therefore unlikely to contribute to chronic immune activation.

DISCUSSION

Class II IN mutations impair virion particle maturation by blocking IN-gRNA binding in virions

and those within the CTD, including R269A/K273A and R262A/R263A substitutions, impede IN-

gRNA binding without affecting functional oligomerization of IN (15, 38). During serial passaging

of the IN R269A/K273A class II mutant virus, the D256N and D270N substitutions emerged

sequentially. We also wondered whether mutations outside of IN, such as CA and NC, could

also arise, given that the IN R269A/K273A mutant is still catalytically active and compensatory

mutations in CA and NC could potentially provide alternative mechanisms for the proper

packaging of the gRNA within virions. However, we did not observe the emergence of such

substitutions showcasing the distinct role of IN-gRNA binding in proper virion maturation.

Furthermore, the emergence of compensatory substitutions only in CTD, but not NTD or CCD of

IN, highlights the importance of the IN-CTD in mediating direct RNA binding.

It is worth noting that D256N and D270N substitutions each arose through a single mutation

(D256N: GAC→AAC, D270N: GAU→AAU), and thus likely provided an easier pathway for

suppression than reverting back to R269 and K273, each of which would require two mutations.

Likewise, D-to-R and D-to-K substitutions require two and three mutations, respectively, and

thus likely had a higher barrier to arise in culture. Though other mutations in IN could in principle

restore the ability of IN to bind RNA, rise of such mutations was likely additionally constrained in

part by the necessity to maintain a catalytically active IN.

The IN-CTD is decorated by numerous acidic and basic residues (**Fig. 3A**). While the IN

R269A/K273A substitutions resulted in the loss of 2 positive charges, the compensatory D2N

substitutions restored the overall local charge within the CTD, suggesting that electrostatic

interactions contribute to IN-CTD binding to the gRNA. Consistent with this hypothesis, the

D256R, D278R and D279R substitutions that resulted in the gain of two positive charges locally

were all sufficient to substantially increase or restore IN-gRNA binding for the IN R269A/K273A

mutant.   Furthermore, these observations extended to another class II mutant, IN

R262A/R263A, whereby the D256R and to a lesser extent the D256K substitutions restored

RNA binding and subsequent infectivity. In contrast, the D2N substitutions did not enhance

RNA-binding or infectivity for the IN R262A/R263A mutant, demonstrating a degree of

context/localization dependency. In other words, it is possible that the proximity of the D270N to

R269A/K273A residues may explain why it was more effective in restoring RNA binding

particularly for this class II mutant but not for IN R262A/R263A. Cumulatively, these findings

strongly suggest an electrostatic component of IN-gRNA interactions. Inspection of the available

X-ray structure (62) indicates that all IN residues implicated in RNA binding by the present study

are positioned either in the highly flexible C-terminal tail (aa 261-275) or the 252-257 loop (**Fig.**

5H). The pliable nature of these CTD regions could be essential for allowing IN to optimally bind

to cognate gRNA.

Arg and Lys residues are commonly employed by numerous RNA-binding proteins, and Arg

residues are generally more heavily involved in interactions with all bases (70, 79-83). Though

the electrostatic interactions between the Arg/Lys residues of IN-CTD and gRNA imply a level of

non-specificity, IN binds to distinct locations on the gRNA in virions and displays high binding

affinity to structured elements, such as TAR (38). Thus, it is likely that IN-gRNA interactions are

mediated by both the non-specific interactions of the basic residues with the RNA phosphate

backbone and specific interactions with the cognate RNA. For example, recognition of the TAR

loop by Tat and the super elongation complex is based on a complex set of interactions: the

super elongation complex primarily reads out the structure as opposed to the sequence,

whereas the Tat arginine rich motif (ARM) interacts with the TAR bulge and the major groove

through electrostatic interactions with the RNA backbone and H-bonding with specific bases

(84, 85). Importantly, we note that while the present study elucidates crucial roles of the IN-CTD

basic residues, other yet to be identified IN amino acids are likely to also contribute to high

affinity binding of select gRNA segments. Structural studies of IN in complex with cognate RNA

oligonucleotides are needed to tease apart the specificity determinants for IN-gRNA

interactions.

RNA binding proteins commonly encode modular RNA-binding domains (i.e. RRMs and KH

domains), which form specific contacts with short degenerate sequences (86-88). Utilization of

multiple RRMs/KH domains is thought to create a much larger binding interface, which in turn

allows for specific, high affinity binding to target RNAs (86, 87, 89). Functionally important IN

tetramerization (15) may similarly enhance binding affinity and specificity to cognate gRNA

elements. In this respect, it is worth noting that CTDs of the two inner protomers in the

tetrameric intasome structure contact extensively with DNA, whereas the outer CTDs are

positioned in close proximity to the inner CTDs and only partially contribute DNA binding (90). It

remains to be seen whether the inner CTDs could also preferentially engage gRNA or CTD-

nucleic acid interactions within IN-gDNA and IN-gRNA complexes differ substantially.

ALLINIs selectively bind at the IN CCD dimer interface at the LEDGF/p75 binding pocket and

induce aberrant IN multimerization (12, 38), which is thought to underlie the mechanism by

which they inhibit IN-gRNA binding. Prior biochemical and structural studies suggested that

ALLINIs also engage the CTD of IN. For example, the quinoline-based ALLINI, GSK-1264, is

buried between the CTD of one IN dimer and the CCD of a nearby dimer resulting in the

formation of open, inactive IN polymers (62). Furthermore, mutation of Y226, W235, K264 and

K266 residues within the CTD prevents ALLINI-induced aberrant multimerization of IN (62, 75,

76). However, evidence for CTD involvement in relevant infection settings is lacking, given that

the mutation of the residues that mediate ALLINI induced hyper-multimerization of IN affect the

catalytic activity as well as RNA binding properties of IN (62, 91, 92). Our findings fill this critical

gap and provide key genetic evidence that ALLINI mechanism of action involves distinct

residues within the CTD, namely R269 and K273. The X-ray structure of GSK-1264 induced IN

polymers (62) reveals that R262, R263, K273A as well as D256 are distanced from the inhibitor

binding pocket, whereas the side chain of R269 points toward the inhibitor bound at the CTD-

CCD dimer interface (**Fig. 7G**). The guanidine group of R269 engages with Q168 of the CCD

from another IN subunit and is positioned within 3 Å of GSK-1264. Yet, there is no interaction

between R269 and the inhibitor as the positively charged guanidine group faces a hydrophobic

part of the benzodihydropyran moiety of GSK-1264 (**Fig. 7G**). Conversely, the R269A

substitution could provide more favorable hydrophobic environment for the CTD-ALLINI-CCD

interactions that may explain its increased sensitivity to ALLINIs.

Overall, our studies reveal that pliable electrostatic interactions play an important role in

mediating IN-CTD-gRNA interactions and demonstrate that CTD is a key determinant of both

RNA binding and ALLINI sensitivity. IN-gRNA binding is essential for HIV-1 virion morphological

maturation and infectivity, thus an excellent target for novel antiretroviral compounds. ALLINI-

mediated inhibition of the non-catalytic function of IN can complement existing INSTI-based

therapies and increase the barrier to drug resistance substantially. Of note, a highly potent and

safe pyrrolopyridine-based ALLINI, STP0404, has recently advanced to human trials (65).

Further structural characterization of IN-gRNA and IN-ALLINI complexes will be crucial to

determine the precise rules that govern IN-gRNA interactions and guide the development of

novel therapeutics that target IN-gRNA interactions.

MATERIALS AND METHODS

Plasmids and Compounds

IN mutations were introduced into the HIV-1$_{NL4-3}$ full-length proviral plasmid (pNL4-3) by overlap

extension PCR. Forward and reverse primers containing IN mutations were used in PCR

reactions with forward (with EcoRI restriction endonuclease site) and reverse (with AgeI

restriction endonuclease site) outer primers. The resulting fragments containing the desired

mutations were mixed at a 1:1 ratio and overlapped subsequently using the outer primer pairs.

The overlap fragments were digested with AgeI and EcoRI before cloning into pNL4-3 plasmids.

The pLR2P-VprIN plasmid expressing a Vpr-IN fusion protein has been previously described

(74). IN mutations were introduced in the pLR2P-VprIN plasmid using the QuikChange Site-

Directed Mutagenesis kit (Agilent Technologies). The presence of the desired mutations and the

absence of unwanted secondary changes was assessed by Sanger sequencing.

ALLINIs, BI-D and BI-B2, were synthesized by Aris Pharmaceuticals using previously published

protocols (93, 94). Compounds were dissolved in DMSO at a final concentration of 10 mM and

stored in -80C.

Cell lines and viruses

Hela-derived TZM-bL cells that express CD4/CXCR4/CCR5 receptor/coreceptors and bear an

HIV-1 LTR driven β-galactosidase reporter were obtained from the NIH AIDS Reagent Program.

HEK293T cells (ATCC CRL-11268) and TZM-bL cells were cultured in Dulbecco's modified

Eagle's medium supplemented with 10% fetal bovine serum. Human CD4+ T-cell line, MT4 (NIH

AIDS Reagents), were cultured in RPMI 1640 medium supplemented with 10% fetal bovine

serum. A derivative of MT4-LTR-GFP (MT4-GFP) indicator cells bearing an HIV-1 LTR driven

GFP reporter have been described before (95) and similarly maintained in RPMI 1640 medium

supplemented with 10% fetal bovine serum. All cell lines were obtained from the American Type

Culture Collection and NIH AIDS Reagents where short tandem repeat (STR) profiling was

performed. MT-4 T-cells were further authenticated by STR profiling at Washington University

School of Medicine Genome Engineering and iPSC center. The cell lines are regularly inspected

for mycoplasma contamination using MycoAlert mycoplasma detection kit (Lonza) and checked

for being free of any other contaminations.

For generation of HIV-1 stocks, HEK293T cells grown in 10-cm dishes were transfected with the

pNL4-3 and a vesicular stomatitis virus glycoprotein (VSV-G) expression plasmids at a ratio of

4:1 using polyethyleneimine (PolySciences, Warrington, PA). Cell culture supernatants

containing infectious virions were collected two days post-transfection, filtered, aliquoted and

stored at -80°C. For characterization of the effects of compensatory mutations on virus

replication, HEK293T cells were transfected in 24-well plates with pNL4-3-derived plasmids

similarly but without VSV-G pseudotyping. Cell lysates collected at 2 days post-transfection

were analyzed by immunoblotting as detailed below. In parallel, cell culture supernatants

containing virions were titered on TZM-bL cells using a β-galactosidase assay (Galactostar,

Thermo Fisher) per manufacturer's instructions. Virus replication in TZM-bL cells was limited to

a single-cycle via addition of dextran sulfate (50 µg/mL) 6-14 hours post-infection. An aliquot of

the viruses were also subjected to a qPCR-based reverse-transcriptase (RT) activity assay (96).

Virus titers obtained from infection of TZM-bL cells was normalized to particle numbers based

on values obtained from RT activity assays to determine the infectiousness (i.e. infectious

units/particle) of viruses. A fraction of the cell culture supernatant collected from transfected

HEK293T cells was also concentrated by centrifugation (16,100 x g, 4°C, 90min) on a 20%

sucrose cushion prepared in 1x SDS-PAGE sample buffer for analysis of virion-associated

proteins by immunoblotting as described below.

Analysis of compensatory mutations

Compensatory substitutions that arose in the background of HIV-1$_{NL4-3 \, IN \, (R269A/K273A)}$ class II IN

mutant virus were isolated by serial passaging as follows. In brief, one million MT4-GFP cells

were infected at a multiplicity of infection (MOI) 2 for the WT virus. An equivalent particle

number (as determined by a qPCR-based reverse transcriptase (RT) activity assay (96)) of the

HIV-1$_{NL4-3 \, IN \, (R269A/K273A)}$ virus was used as inoculum. Infected cells were split at a ratio of 1:5

every 3-4 days and infections were monitored by microscopy as well as flow cytometry of GFP

positive cells at the indicated intervals shown in Figure 1A. Cell culture supernatants containing

virions were also collected at the same time intervals for subsequent sequencing analysis. At

the end of each passage (i.e. when infections plateaued at >75%GFP+ cells), cell culture

supernatants containing infectious virions were collected. The resultant viruses were first titered

on MT4-GFP cells and used to infect MT4-GFP cells in the next passage at an MOI of 2 i.u./cell

for WT and an equivalent particle number for the class II IN mutant virus (or its derivative

bearing compensatory mutations). Aliquots of infected cells and viral particles in the cell culture

supernatants were collected over the duration of each passage as above.

For sequencing analysis of viruses bearing compensatory mutations, virions collected over the

three passages were concentrated on 20% sucrose cushions (prepared in 1X phosphate

buffered saline (PBS)) by ultracentrifugation (28000 rpm, 4°C, 90min, Beckman SW41 rotor).

Genomic RNA was isolated from pelleted virions using Trizol per manufacturer's instructions.

Extracted RNA was prepared for deep sequencing using the Illumina® TruSeq® Stranded Total

RNA library kit following the manufacturer's instructions but omitting the rRNA depletion step.

Resulting libraries were sequenced on an Illumina HiSeq 2000 platform at the Genome

Technology Access Center at Washington University School of Medicine. Sequencing reads

were mapped to the pNL4-3 reference viral genome allowing for 2 mismatches using the Bowtie

algorithm (i.e. -v 2, -m 10 parameters)(97) and the frequency of compensatory mutations was

determined thereafter.

Immunoblotting

Viral and cell lysates were resuspended in sodium dodecyl sulfate (SDS) sample buffer,

separated by electrophoresis on Bolt 4-12% Bis-Tris Plus gels (Life Technologies) and

transferred to nitrocellulose membranes (Hybond ECL, Amersham). The membranes were then

probed overnight at 4°C with a mouse monoclonal anti-HIV p24 antibody (183-H12-5C, NIH

AIDS reagents) or a mouse monoclonal anti-HIV integrase antibody (98) in Odyssey Blocking

Buffer (LI-COR). Membranes were probed with fluorophore-conjugated secondary antibodies

(LI-COR) and scanned using a LI-COR Odyssey system. IN and CA levels in virions were

quantified using Image Studio software (LI-COR).

Vpr-IN trans-complementation experiments

HIV-1$_{NL4-3}$ bearing the class I IN mutation D116N was trans-complemented with class II mutant

proteins as previously described (74). In brief, HEK293T cells plated in 24-well dishes were co-

transfected with full-length proviral plasmids expressing HIV-1$_{NL4-3\ IN\ (D116N)}$, VSV-G, and

derivatives of the pLR2P-VprIN plasmids bearing class II IN mutations (or the compensatory

mutations thereof) at a ratio of 6:1:3 using polyethyleneimine as described above. Cell-free

virions were collected from cell culture two days post-transfection. MT-4 T-cells were infected by

the resultant virus stocks and the integration capability of trans-complemented class II IN

mutants was measured by the yield of progeny virions in cell culture supernatants over a 6-day

period using the aforementioned reverse transcriptase (RT) activity assay as described before

(74). Briefly, MT-4 T-cells were incubated with virus inoculum in 96 V-bottom well plates for 4hr

at 37°C before washing away the inoculum and replacing it with fresh media. Right after the

addition of fresh media and over the ensuing 6 days, an aliquot of the media was collected and

the amount of virions in culture supernatant was quantified by measuring a qPCR-based RT

activity using assay (96), as also described above.

CLIP experiments

CLIP experiments were conducted as previously described (15, 72, 73, 99, 100). In short, cells

in 15 cm cell culture plates were transfected with 30 µg full-length proviral plasmid (pNL4-3)

DNA or derivatives carrying the IN mutations. 4-thiouridine (4SU) was added to the cell culture

media for 16hr before virus harvest. Cell culture supernatants containing virions were filtered

through 0.22 μm filters and pelleted by ultracentrifugation through a 20% sucrose cushion

prepared in 1X phosphate-buffered saline (PBS) using a Beckman SW32-Ti rotor at 28,000rpm

for 1.5hr at 4°C. The virus pellets were resuspended in 1XPBS and UV-crosslinked for two

consecutive times at an energy setting of 500 mJ in a Boekel UV-crosslinking chamber

equipped with UV368 nm bulbs. Following lysis in 1X RIPA buffer, IN-RNA complexes were

immunoprecipitated using a mouse monoclonal anti-IN antibody (98). Bound RNA was end-

labeled with γ-^{32}P-ATP and T4 polynucleotide kinase. The isolated protein-RNA complexes

were separated by SDS-PAGE, transferred to nitrocellulose membranes (Hybond ECL,

Amersham), and exposed to autoradiography films to visualize IN-RNA complexes. If needed,

following immunoblotting and quantitation of immunoprecipitated IN, equivalent amounts of

immunoprecipitated IN were re-run to accurately assess the ability of IN variants to bind RNA.

Autoradiographs were quantitated by the ImageStudio software (LI-COR). Lysates and

immunoprecipitates were also analyzed by immunoblotting using the aforementioned mouse

monoclonal antibodies against IN.

IN multimerization in virions

HEK293T cells grown in 10-cm dishes were transfected with 10 μg pNL4-3 plasmid DNA

bearing WT IN or the indicated *pol* mutations within IN coding sequence as above. Two days

post-transfection, cell-free virions in cell culture supernatants were pelleted through a 20%

sucrose cushion prepared in 1xPBS using a Beckman SW41-Ti rotor at 28,000 rpm for 1.5hr at

4°C. Pelleted virions were resuspended in 1xPBS and treated with a membrane-permeable

crosslinker, EGS (ThermoFisher Scientific), at a concentration of 1mM for 30 min at room

temperature. Crosslinking was stopped by the addition of SDS-PAGE sample buffer. The cross-

linked samples were then separated on 3-8% Tris-acetate gels and analyzed by immunoblotting

using a mouse monoclonal anti-IN antibody (98).

Virus production and transmission electron microscopy

HEK293T cells grown in 15-cm plates were transfected with 30 µg full-length proviral plasmid (pNL4-3) DNA containing WT IN or the indicated *pol* mutations within IN coding sequence as above. Two days post transfection, cell culture supernatants were filtered through 0.22 µm filters, and pelleted by ultracentrifugation using a Beckman SW32-Ti rotor at 28,000 rpm for 1.5 hr at 4°C. Fixative (2% paraformaldehyde/2.5% glutaraldehyde (Polysciences Inc., Warrington, PA) in 100 mM sodium cacodylate buffer, pH 7.2) was gently added to resulting pellets, and samples were incubated overnight at 4°C. Samples were washed in sodium cacodylate buffer and postfixed in 1% osmium tetroxide (Polysciences Inc.) for 1 hr. Samples were then rinsed extensively in dH_2O prior to en bloc staining with 1% aqueous uranyl acetate (Ted Pella Inc., Redding, CA) for 1 hr. After several rinses in dH_2O, samples were dehydrated in a graded series of ethanol and embedded in Eponate 12 resin (Ted Pella Inc.). Sections of 95 nm were cut with a Leica Ultracut UCT ultramicrotome (Leica Microsystems Inc., Bannockburn, IL), stained with uranyl acetate and lead citrate, and viewed on a JEOL 1200 EX transmission electron microscope (JEOL USA Inc., Peabody, MA) equipped with an AMT 8 megapixel digital camera and AMT Image Capture Engine V602 software (Advanced Microscopy Techniques, Woburn, MA). 100 virions from each sample was classified for virion morphology.

Size exclusion chromatography (SEC)

All of the indicated mutations were introduced into a plasmid backbone expressing His_6 tagged pNL4-3-derived IN by QuikChange site directed mutagenesis kit (Agilent) (66). His_6 tagged recombinant pNL4-3 WT and mutant IN molecules were expressed in BL21 (DE3) *E. coli* cells followed by nickel and heparin column purification as described previously (66, 101). Recombinant WT and mutant IN molecules were analyzed on Superdex 200 10/300 GL column (GE Healthcare) with running buffer containing 20 mM HEPES (pH 7.5), 1 M NaCl, 10% glycerol

and 5 mM BME at 0.3 mL/min flow rate. The proteins were diluted to 10 µM with the running

buffer and incubated for 1 h at 4°C followed by centrifugation at 10,000xg for 10 min. Multimeric

form determination was based on the standards including bovine thyroglobulin (670,000 Da),

bovine gamma-globulin (158,000 Da), chicken ovalbumin (44,000 Da), horse myoglobin (17,000

Da) and vitamin B12 (1,350 Da). Retention volumes for different oligomeric forms of IN were as

follows: tetramer ~12.5 mL, dimer ~14 mL, monomer ~15-16 mL.

Analysis of IN-RNA binding *in vitro*

To monitor IN-RNA interactions we utilized AlphaScreen-based assay (38), which allows to

monitor the ability of IN to bind and bridge between two TAR RNAs. Briefly, equal

concentrations (1 nM) of two synthetic TAR RNA oligonucleotides labeled either with biotin or

DIG were mixed and then streptavidin donor and anti-DIG acceptor beads at 0.02 mg/mL

concentration were supplied in a buffer containing 100 mM NaCl, 1 mM $MgCl_2$, 1 mM DTT, 1

mg/mL BSA, and 25 mM Tris (pH 7.4). After 2 hr incubation at 4°C, 320 nM IN was added to the

reaction mixture and incubated further for 1.5 hr at 4°C. AlphaScreen signals were recorded

with a PerkinElmer Life Sciences Enspire multimode plate reader.

Structural modeling of IN

Electrostatic potential maps of WT and mutant IN CTDs were created by Adaptive Poisson-

Boltzmann Solver (APBS) program (102) with macromolecular electrostatic calculations

performed in PyMOL. The published crystal structure (62) (PDB ID: 5HOT) was used as a

template. The calculation results are displayed as an electrostatic potential molecular surface.

The low, mid, and high range values are -5, 0, and 5, respectively.

Analysis of IN mutations from latently infected cells

Sequences identified from latently infected CD4+ T-cells (77) were downloaded from NCBI

GenBank based on their accession numbers (KF526120-KF526339). Sequences were imported

into the GLUE software framework (103, 104) and aligned. Multiple sequence alignments

(MSAs) containing subtype B sequences were constructed using MUSCLE, manually inspected

in AliView (105) and imported into a GLUE project database. Within GLUE, MSAs were

constrained to the pNL4-3 reference to establish a standardized coordinate space for the gene

being analyzed. Amino acid frequencies at each alignment position were summarized using

GLUE's amino-acid frequency calculation algorithm, which accounts for contingencies such as

missing data and incomplete codons.

ACKNOWLEDGEMENTS

This work was supported by NIH grants AI150497 (SBK), AI1508470 (U54 Center for HIV RNA

Studies, SBK, RG), R01 AI143649 (MK) and Milton Schlesinger Student Fellowship (CSM). We

thank all members of the Kutluay lab for critical suggestions and feedback and Wandy Beatty at

Washington University for assistance and expertise in transmission electron microscopy.

REFERENCES

1. Engelman AN. 2019. Multifaceted HIV integrase functionalities and therapeutic strategies for their inhibition. J Biol Chem doi:10.1074/jbc.REV119.006901.
2. Hazuda DJ, Felock P, Witmer M, Wolfe A, Stillmock K, Grobler JA, Espeseth A, Gabryelski L, Schleif W, Blau C, Miller MD. 2000. Inhibitors of strand transfer that prevent integration and inhibit HIV-1 replication in cells. Science 287:646-50.
3. Summa V, Petrocchi A, Bonelli F, Crescenzi B, Donghi M, Ferrara M, Fiore F, Gardelli C, Gonzalez Paz O, Hazuda DJ, Jones P, Kinzel O, Laufer R, Monteagudo E, Muraglia E, Nizi E, Orvieto F, Pace P, Pescatore G, Scarpelli R, Stillmock K, Witmer MV, Rowley M. 2008. Discovery of raltegravir, a potent, selective orally bioavailable HIV-integrase inhibitor for the treatment of HIV-AIDS infection. J Med Chem 51:5843-55.
4. Ramanathan S, Mathias AA, German P, Kearney BP. 2011. Clinical pharmacokinetic and pharmacodynamic profile of the HIV integrase inhibitor elvitegravir. Clin Pharmacokinet 50:229-44.
5. Min S, Song I, Borland J, Chen S, Lou Y, Fujiwara T, Piscitelli SC. 2010. Pharmacokinetics and safety of S/GSK1349572, a next-generation HIV integrase inhibitor, in healthy volunteers. Antimicrob Agents Chemother 54:254-8.
6. Tsiang M, Jones GS, Goldsmith J, Mulato A, Hansen D, Kan E, Tsai L, Bam RA, Stepan G, Stray KM, Niedziela-Majka A, Yant SR, Yu H, Kukolj G, Cihlar T, Lazerwith SE, White KL, Jin H. 2016. Antiviral Activity of Bictegravir (GS-9883), a Novel Potent HIV-1

Integrase Strand Transfer Inhibitor with an Improved Resistance Profile. Antimicrob
Agents Chemother 60:7086-7097.
7.   Smith SJ, Zhao XZ, Burke TR, Jr., Hughes SH. 2018. Efficacies of Cabotegravir and
Bictegravir against drug-resistant HIV-1 integrase mutants. Retrovirology 15:37.
8.   Rockstroh JK, DeJesus E, Lennox JL, Yazdanpanah Y, Saag MS, Wan H, Rodgers AJ,
Walker ML, Miller M, DiNubile MJ, Nguyen BY, Teppler H, Leavitt R, Sklar P,
Investigators S. 2013. Durable efficacy and safety of raltegravir versus efavirenz when
combined with tenofovir/emtricitabine in treatment-naive HIV-1-infected patients: final 5-
year results from STARTMRK. J Acquir Immune Defic Syndr 63:77-85.
9.   Taha H, Das A, Das S. 2015. Clinical effectiveness of dolutegravir in the treatment of
HIV/AIDS. Infect Drug Resist 8:339-52.
10.   Smith SJ, Zhao XZ, Passos DO, Lyumkis D, Burke TR, Hughes SH. 2020. HIV-1
Integrase Inhibitors that are active against Drug-Resistant Integrase Mutants. Antimicrob
Agents Chemother doi:10.1128/AAC.00611-20.
11.   Brooks KM, Sherman EM, Egelund EF, Brotherton A, Durham S, Badowski ME, Cluck
DB. 2019. Integrase Inhibitors: After 10 Years of Experience, Is the Best Yet to Come?
Pharmacotherapy 39:576-598.
12.   Jurado KA, Wang H, Slaughter A, Feng L, Kessl JJ, Koh Y, Wang W, Ballandras-Colas
A, Patel PA, Fuchs JR, Kvaratskhelia M, Engelman A. 2013. Allosteric integrase inhibitor
potency is determined through the inhibition of HIV-1 particle maturation. Proc Natl Acad
Sci U S A 110:8690-5.
13.   Lu R, Limon A, Devroe E, Silver PA, Cherepanov P, Engelman A. 2004. Class II
integrase mutants with changes in putative nuclear localization signals are primarily
blocked at a postnuclear entry step of human immunodeficiency virus type 1 replication.
J Virol 78:12735-46.
14.   Engelman A. 1999. In vivo analysis of retroviral integrase structure and function. Adv
Virus Res 52:411-26.
15.   Elliott JL, Eschbach JE, Koneru PC, Li W, Puray-Chavez M, Townsend D, Lawson DQ,
Engelman AN, Kvaratskhelia M, Kutluay SB. 2020. Integrase-RNA interactions
underscore the critical role of integrase in HIV-1 virion morphogenesis. Elife 9.
16.   Engelman A, Englund G, Orenstein JM, Martin MA, Craigie R. 1995. Multiple effects of
mutations in human immunodeficiency virus type 1 integrase on viral replication. J Virol
69:2729-36.
17.   Engelman A, Cherepanov P. 2014. Retroviral Integrase Structure and DNA
Recombination Mechanism. Microbiol Spectr 2.
18.   Engelman AN, Cherepanov P. 2017. Retroviral intasomes arising. Curr Opin Struct Biol
47:23-29.
19.   Craigie R, Bushman FD. 2012. HIV DNA integration. Cold Spring Harb Perspect Med
2:a006890.
20.   Engelman A, Cherepanov P. 2012. The structural biology of HIV-1: mechanistic and
therapeutic insights. Nat Rev Microbiol 10:279-90.
21.   Ciuffi A, Llano M, Poeschla E, Hoffmann C, Leipzig J, Shinn P, Ecker JR, Bushman F.
2005. A role for LEDGF/p75 in targeting HIV DNA integration. Nat Med 11:1287-9.
22.   Shun MC, Raghavendra NK, Vandegraaff N, Daigle JE, Hughes S, Kellam P,
Cherepanov P, Engelman A. 2007. LEDGF/p75 functions downstream from
preintegration complex formation to effect gene-specific HIV-1 integration. Genes Dev
21:1767-78.
23.   Schroder AR, Shinn P, Chen H, Berry C, Ecker JR, Bushman F. 2002. HIV-1 integration
in the human genome favors active genes and local hotspots. Cell 110:521-9.
24.   Christ F, Debyser Z. 2013. The LEDGF/p75 integrase interaction, a novel target for anti-
HIV therapy. Virology 435:102-9.

25. Ansari-Lari MA, Donehower LA, Gibbs RA. 1995. Analysis of human immunodeficiency
virus type 1 integrase mutants. Virology 213:680.
26. Bukovsky A, Gottlinger H. 1996. Lack of integrase can markedly affect human
immunodeficiency virus type 1 particle production in the presence of an active viral
protease. J Virol 70:6820-5.
27. Jenkins TM, Engelman A, Ghirlando R, Craigie R. 1996. A soluble active mutant of HIV-
1 integrase: involvement of both the core and carboxyl-terminal domains in
multimerization. J Biol Chem 271:7712-8.
28. Kalpana GV, Reicin A, Cheng GS, Sorin M, Paik S, Goff SP. 1999. Isolation and
characterization of an oligomerization-negative mutant of HIV-1 integrase. Virology
259:274-85.
29. Leavitt AD, Robles G, Alesandro N, Varmus HE. 1996. Human immunodeficiency virus
type 1 integrase mutants retain in vitro integrase activity yet fail to integrate viral DNA
efficiently during infection. J Virol 70:721-8.
30. Liao WH, Wang CT. 2004. Characterization of human immunodeficiency virus type 1
Pr160 gag-pol mutants with truncations downstream of the protease domain. Virology
329:180-8.
31. Lu R, Ghory HZ, Engelman A. 2005. Genetic analyses of conserved residues in the
carboxyl-terminal domain of human immunodeficiency virus type 1 integrase. J Virol
79:10356-68.
32. Nakamura T, Masuda T, Goto T, Sano K, Nakai M, Harada S. 1997. Lack of infectivity of
HIV-1 integrase zinc finger-like domain mutant with morphologically normal maturation.
Biochem Biophys Res Commun 239:715-22.
33. Quillent C, Borman AM, Paulous S, Dauguet C, Clavel F. 1996. Extensive regions of pol
are required for efficient human immunodeficiency virus polyprotein processing and
particle maturation. Virology 219:29-36.
34. Shin CG, Taddeo B, Haseltine WA, Farnet CM. 1994. Genetic analysis of the human
immunodeficiency virus type 1 integrase protein. J Virol 68:1633-42.
35. Taddeo B, Haseltine WA, Farnet CM. 1994. Integrase mutants of human
immunodeficiency virus type 1 with a specific defect in integration. J Virol 68:8401-5.
36. Wu X, Liu H, Xiao H, Conway JA, Hehl E, Kalpana GV, Prasad V, Kappes JC. 1999.
Human immunodeficiency virus type 1 integrase protein promotes reverse transcription
through specific interactions with the nucleoprotein reverse transcription complex. J Virol
73:2126-35.
37. Fontana J, Jurado KA, Cheng N, Ly NL, Fuchs JR, Gorelick RJ, Engelman AN, Steven
AC. 2015. Distribution and Redistribution of HIV-1 Nucleocapsid Protein in Immature,
Mature, and Integrase-Inhibited Virions: a Role for Integrase in Maturation. J Virol
89:9765-80.
38. Kessl JJ, Kutluay SB, Townsend D, Rebensburg S, Slaughter A, Larue RC, Shkriabai N,
Bakouche N, Fuchs JR, Bieniasz PD, Kvaratskhelia M. 2016. HIV-1 Integrase Binds the
Viral RNA Genome and Is Essential during Virion Morphogenesis. Cell 166:1257-1268
e12.
39. Ao Z, Fowke KR, Cohen EA, Yao X. 2005. Contribution of the C-terminal tri-lysine
regions of human immunodeficiency virus type 1 integrase for efficient reverse
transcription and viral DNA nuclear import. Retrovirology 2:62.
40. Busschots K, Voet A, De Maeyer M, Rain JC, Emiliani S, Benarous R, Desender L,
Debyser Z, Christ F. 2007. Identification of the LEDGF/p75 binding site in HIV-1
integrase. J Mol Biol 365:1480-92.
41. Engelman A, Liu Y, Chen H, Farzan M, Dyda F. 1997. Structure-based mutagenesis of
the catalytic domain of human immunodeficiency virus type 1 integrase. J Virol 71:3507-
14.

42.   Limon A, Devroe E, Lu R, Ghory HZ, Silver PA, Engelman A. 2002. Nuclear localization
of human immunodeficiency virus type 1 preintegration complexes (PICs): V165A and
R166A are pleiotropic integrase mutants primarily defective for integration, not PIC
nuclear import. J Virol 76:10598-607.
43.   Lloyd AG, Ng YS, Muesing MA, Simon V, Mulder LC. 2007. Characterization of HIV-1
integrase N-terminal mutant viruses. Virology 360:129-35.
44.   Lu R, Vandegraaff N, Cherepanov P, Engelman A. 2005. Lys-34, dispensable for
integrase catalysis, is required for preintegration complex function and human
immunodeficiency virus type 1 replication. J Virol 79:12584-91.
45.   Masuda T, Planelles V, Krogstad P, Chen IS. 1995. Genetic analysis of human
immunodeficiency virus type 1 integrase and the U3 att site: unusual phenotype of
mutants in the zinc finger-like domain. J Virol 69:6687-96.
46.   Rahman S, Lu R, Vandegraaff N, Cherepanov P, Engelman A. 2007. Structure-based
mutagenesis of the integrase-LEDGF/p75 interface uncouples a strict correlation
between in vitro protein binding and HIV-1 fitness. Virology 357:79-90.
47.   Riviere L, Darlix JL, Cimarelli A. 2010. Analysis of the viral elements required in the
nuclear import of HIV-1 DNA. J Virol 84:729-39.
48.   Tsurutani N, Kubo M, Maeda Y, Ohashi T, Yamamoto N, Kannagi M, Masuda T. 2000.
Identification of critical amino acid residues in human immunodeficiency virus type 1 IN
required for efficient proviral DNA formation at steps prior to integration in dividing and
nondividing cells. J Virol 74:4795-806.
49.   Wiskerchen M, Muesing MA. 1995. Human immunodeficiency virus type 1 integrase:
effects of mutations on viral ability to integrate, direct viral gene expression from
unintegrated viral DNA templates, and sustain viral propagation in primary cells. J Virol
69:376-86.
50.   Zhu K, Dobard C, Chow SA. 2004. Requirement for integrase during reverse
transcription of human immunodeficiency virus type 1 and the effect of cysteine
mutations of integrase on its interactions with reverse transcriptase. J Virol 78:5045-55.
51.   De Houwer S, Demeulemeester J, Thys W, Rocha S, Dirix L, Gijsbers R, Christ F,
Debyser Z. 2014. The HIV-1 integrase mutant R263A/K264A is 2-fold defective for TRN-
SR2 binding and viral nuclear import. J Biol Chem 289:25351-61.
52.   Johnson BC, Metifiot M, Ferris A, Pommier Y, Hughes SH. 2013. A homology model of
HIV-1 integrase and analysis of mutations designed to test the model. J Mol Biol
425:2133-46.
53.   Mohammed KD, Topper MB, Muesing MA. 2011. Sequential deletion of the integrase
(Gag-Pol) carboxyl terminus reveals distinct phenotypic classes of defective HIV-1. J
Virol 85:4654-66.
54.   Shehu-Xhilaga M, Hill M, Marshall JA, Kappes J, Crowe SM, Mak J. 2002. The
conformation of the mature dimeric human immunodeficiency virus type 1 RNA genome
requires packaging of pol protein. J Virol 76:4331-40.
55.   Engelman A, Craigie R. 1992. Identification of conserved amino acid residues critical for
human immunodeficiency virus type 1 integrase function in vitro. J Virol 66:6361-9.
56.   Lutzke RA, Plasterk RH. 1998. Structure-based mutational analysis of the C-terminal
DNA-binding domain of human immunodeficiency virus type 1 integrase: critical residues
for protein oligomerization and DNA binding. J Virol 72:4841-8.
57.   Lutzke RA, Vink C, Plasterk RH. 1994. Characterization of the minimal DNA-binding
domain of the HIV integrase protein. Nucleic Acids Res 22:4125-31.
58.   Balakrishnan M, Yant SR, Tsai L, O'Sullivan C, Bam RA, Tsai A, Niedziela-Majka A,
Stray KM, Sakowicz R, Cihlar T. 2013. Non-catalytic site HIV-1 integrase inhibitors
disrupt core maturation and induce a reverse transcription block in target cells. PLoS
One 8:e74163.

59. Desimmie BA, Schrijvers R, Demeulemeester J, Borrenberghs D, Weydert C, Thys W,
Vets S, Van Remoortel B, Hofkens J, De Rijck J, Hendrix J, Bannert N, Gijsbers R,
Christ F, Debyser Z. 2013. LEDGINs inhibit late stage HIV-1 replication by modulating
integrase multimerization in the virions. Retrovirology 10:57.
60. Sharma A, Slaughter A, Jena N, Feng L, Kessl JJ, Fadel HJ, Malani N, Male F, Wu L,
Poeschla E, Bushman FD, Fuchs JR, Kvaratskhelia M. 2014. A new class of
multimerization selective inhibitors of HIV-1 integrase. PLoS Pathog 10:e1004171.
61. Gupta K, Brady T, Dyer BM, Malani N, Hwang Y, Male F, Nolte RT, Wang L, Velthuisen
E, Jeffrey J, Van Duyne GD, Bushman FD. 2014. Allosteric inhibition of human
immunodeficiency virus integrase: late block during viral replication and abnormal
multimerization involving specific protein domains. J Biol Chem 289:20477-88.
62. Gupta K, Turkki V, Sherrill-Mix S, Hwang Y, Eilers G, Taylor L, McDanal C, Wang P,
Temelkoff D, Nolte RT, Velthuisen E, Jeffrey J, Van Duyne GD, Bushman FD. 2016.
Structural Basis for Inhibitor-Induced Aggregation of HIV Integrase. PLoS Biol
14:e1002584.
63. Christ F, Shaw S, Demeulemeester J, Desimmie BA, Marchand A, Butler S, Smets W,
Chaltin P, Westby M, Debyser Z, Pickford C. 2012. Small-molecule inhibitors of the
LEDGF/p75 binding site of integrase block HIV replication and modulate integrase
multimerization. Antimicrob Agents Chemother 56:4365-74.
64. Christ F, Voet A, Marchand A, Nicolet S, Desimmie BA, Marchand D, Bardiot D, Van der
Veken NJ, Van Remoortel B, Strelkov SV, De Maeyer M, Chaltin P, Debyser Z. 2010.
Rational design of small-molecule inhibitors of the LEDGF/p75-integrase interaction and
HIV replication. Nat Chem Biol 6:442-8.
65. Maehigashi T, Ahn S, Kim UI, Lindenberger J, Oo A, Koneru PC, Mahboubi B,
Engelman AN, Kvaratskhelia M, Kim K, Kim B. 2021. A highly potent and safe
pyrrolopyridine-based allosteric HIV-1 integrase inhibitor targeting host LEDGF/p75-
integrase interaction site. PLoS Pathog 17:e1009671.
66. Kessl JJ, Jena N, Koh Y, Taskent-Sezgin H, Slaughter A, Feng L, de Silva S, Wu L, Le
Grice SF, Engelman A, Fuchs JR, Kvaratskhelia M. 2012. Multimode, cooperative
mechanism of action of allosteric HIV-1 integrase inhibitors. J Biol Chem 287:16801-11.
67. Le Rouzic E, Bonnard D, Chasset S, Bruneau JM, Chevreuil F, Le Strat F, Nguyen J,
Beauvoir R, Amadori C, Brias J, Vomscheid S, Eiler S, Levy N, Delelis O, Deprez E,
Saib A, Zamborlini A, Emiliani S, Ruff M, Ledoussal B, Moreau F, Benarous R. 2013.
Dual inhibition of HIV-1 replication by integrase-LEDGF allosteric inhibitors is
predominant at the post-integration stage. Retrovirology 10:144.
68. Tsiang M, Jones GS, Niedziela-Majka A, Kan E, Lansdon EB, Huang W, Hung M,
Samuel D, Novikov N, Xu Y, Mitchell M, Guo H, Babaoglu K, Liu X, Geleziunas R,
Sakowicz R. 2012. New class of HIV-1 integrase (IN) inhibitors with a dual mode of
action. J Biol Chem 287:21189-203.
69. Serganov A, Patel DJ. 2008. Towards deciphering the principles underlying an mRNA
recognition code. Curr Opin Struct Biol 18:120-9.
70. Bahadur RP, Zacharias M, Janin J. 2008. Dissecting protein-RNA recognition sites.
Nucleic Acids Res 36:2705-16.
71. Stefl R, Skrisovska L, Allain FH. 2005. RNA sequence- and shape-dependent
recognition by proteins in the ribonucleoprotein particle. EMBO Rep 6:33-8.
72. Shema Mugisha C, Tenneti K, Kutluay SB. 2020. Clip for studying protein-RNA
interactions that regulate virus replication. Methods 183:84-92.
73. Bieniasz PD, Kutluay SB. 2018. CLIP-related methodologies and their application to
retrovirology. Retrovirology 15:35.

74.   Liu H, Wu X, Xiao H, Conway JA, Kappes JC. 1997. Incorporation of functional human
838   immunodeficiency virus type 1 integrase into virions independent of the Gag-Pol
839   precursor protein. J Virol 71:7704-10.
840   75.   Deng N, Hoyte A, Mansour YE, Mohamed MS, Fuchs JR, Engelman AN, Kvaratskhelia
841   M, Levy R. 2016. Allosteric HIV-1 integrase inhibitors promote aberrant protein
842   multimerization by directly mediating inter-subunit interactions: Structural and
843   thermodynamic modeling studies. Protein Sci 25:1911-1917.
844   76.   Shkriabai N, Dharmarajan V, Slaughter A, Kessl JJ, Larue RC, Feng L, Fuchs JR, Griffin
845   PR, Kvaratskhelia M. 2014. A critical role of the C-terminal segment for allosteric
846   inhibitor-induced aberrant multimerization of HIV-1 integrase. J Biol Chem 289:26430-
847   26440.
848   77.   Ho YC, Shan L, Hosmane NN, Wang J, Laskey SB, Rosenbloom DI, Lai J, Blankson JN,
849   Siliciano JD, Siliciano RF. 2013. Replication-competent noninduced proviruses in the
850   latent reservoir increase barrier to HIV-1 cure. Cell 155:540-51.
851   78.   Pollack RA, Jones RB, Pertea M, Bruner KM, Martin AR, Thomas AS, Capoferri AA, Beg
852   SA, Huang SH, Karandish S, Hao H, Halper-Stromberg E, Yong PC, Kovacs C, Benko
853   E, Siliciano RF, Ho YC. 2017. Defective HIV-1 Proviruses Are Expressed and Can Be
854   Recognized by Cytotoxic T Lymphocytes, which Shape the Proviral Landscape. Cell
855   Host Microbe 21:494-506 e4.
856   79.   Bayer TS, Booth LN, Knudsen SM, Ellington AD. 2005. Arginine-rich motifs present
857   multiple interfaces for specific binding by RNA. RNA 11:1848-57.
858   80.   Jones S, Daley DT, Luscombe NM, Berman HM, Thornton JM. 2001. Protein-RNA
859   interactions: a structural analysis. Nucleic Acids Res 29:943-54.
860   81.   Jeong E, Kim H, Lee SW, Han K. 2003. Discovering the interaction propensities of
861   amino acids and nucleotides from protein-RNA complexes. Mol Cells 16:161-7.
862   82.   Kim OT, Yura K, Go N. 2006. Amino acid residue doublet propensity in the protein-RNA
863   interface and its application to RNA interface prediction. Nucleic Acids Res 34:6450-60.
864   83.   Kruger DM, Neubacher S, Grossmann TN. 2018. Protein-RNA interactions: structural
865   characteristics and hotspot amino acids. RNA 24:1457-1465.
866   84.   Schulze-Gahmen U, Hurley JH. 2018. Structural mechanism for HIV-1 TAR loop
867   recognition by Tat and the super elongation complex. Proc Natl Acad Sci U S A
868   115:12973-12978.
869   85.   Schulze-Gahmen U, Echeverria I, Stjepanovic G, Bai Y, Lu H, Schneidman-Duhovny D,
870   Doudna JA, Zhou Q, Sali A, Hurley JH. 2016. Insights into HIV-1 proviral transcription
871   from integrative structure and dynamics of the Tat:AFF4:P-TEFb:TAR complex. Elife 5.
872   86.   Han SP, Tang YH, Smith R. 2010. Functional diversity of the hnRNPs: past, present and
873   perspectives. Biochem J 430:379-92.
874   87.   Jeong S. 2017. SR Proteins: Binders, Regulators, and Connectors of RNA. Mol Cells
875   40:1-9.
876   88.   Lunde BM, Moore C, Varani G. 2007. RNA-binding proteins: modular design for efficient
877   function. Nat Rev Mol Cell Biol 8:479-90.
878   89.   Ban T, Zhu JK, Melcher K, Xu HE. 2015. Structural mechanisms of RNA recognition:
879   sequence-specific and non-specific RNA-binding proteins and the Cas9-RNA-DNA
880   complex. Cell Mol Life Sci 72:1045-58.
881   90.   Passos DO, Li M, Yang R, Rebensburg SV, Ghirlando R, Jeon Y, Shkriabai N,
882   Kvaratskhelia M, Craigie R, Lyumkis D. 2017. Cryo-EM structures and atomic model of
883   the HIV-1 strand transfer complex intasome. Science 355:89-92.
884   91.   Kessl JJ, Sharma A, Kvaratskhelia M. 2016. Methods for the Analyses of Inhibitor-
885   Induced Aberrant Multimerization of HIV-1 Integrase. Methods Mol Biol 1354:149-64.
886   92.   Madison MK, Lawson DQ, Elliott J, Ozanturk AN, Koneru PC, Townsend D, Errando M,
887   Kvaratskhelia M, Kutluay SB. 2017. Allosteric HIV-1 Integrase Inhibitors Lead to

Premature Degradation of the Viral RNA Genome and Integrase in Target Cells. J Virol
91.

93.   Wang H, Jurado KA, Wu X, Shun MC, Li X, Ferris AL, Smith SJ, Patel PA, Fuchs JR,
Cherepanov P, Kvaratskhelia M, Hughes SH, Engelman A. 2012. HRP2 determines the
efficiency and specificity of HIV-1 integration in LEDGF/p75 knockout cells but does not
contribute to the antiviral activity of a potent LEDGF/p75-binding site integrase inhibitor.
Nucleic Acids Res 40:11518-30.

94.   Feng L, Sharma A, Slaughter A, Jena N, Koh Y, Shkriabai N, Larue RC, Patel PA,
Mitsuya H, Kessl JJ, Engelman A, Fuchs JR, Kvaratskhelia M. 2013. The A128T
resistance mutation reveals aberrant protein multimerization as the primary mechanism
of action of allosteric HIV-1 integrase inhibitors. J Biol Chem 288:15813-20.

95.   Kane M, Zang TM, Rihn SJ, Zhang F, Kueck T, Alim M, Schoggins J, Rice CM, Wilson
SJ, Bieniasz PD. 2016. Identification of Interferon-Stimulated Genes with Antiretroviral
Activity. Cell Host Microbe 20:392-405.

96.   Pizzato M, Erlwein O, Bonsall D, Kaye S, Muir D, McClure MO. 2009. A one-step SYBR
Green I-based product-enhanced reverse transcriptase assay for the quantitation of
retroviruses in cell culture supernatants. J Virol Methods 156:1-7.

97.   Langmead B, Trapnell C, Pop M, Salzberg SL. 2009. Ultrafast and memory-efficient
alignment of short DNA sequences to the human genome. Genome Biol 10:R25.

98.   Bouyac-Bertoia M, Dvorin JD, Fouchier RA, Jenkins Y, Meyer BE, Wu LI, Emerman M,
Malim MH. 2001. HIV-1 infection requires a functional integrase NLS. Mol Cell 7:1025-
35.

99.   Kutluay SB, Bieniasz PD. 2016. Analysis of HIV-1 Gag-RNA Interactions in Cells and
Virions by CLIP-seq. Methods Mol Biol 1354:119-31.

100. Kutluay SB, Zang T, Blanco-Melo D, Powell C, Jannain D, Errando M, Bieniasz PD.
2014. Global changes in the RNA binding specificity of HIV-1 gag regulate virion
genesis. Cell 159:1096-1109.

101. Cherepanov P. 2007. LEDGF/p75 interacts with divergent lentiviral integrases and
modulates their enzymatic activity in vitro. Nucleic Acids Res 35:113-24.

102. Jurrus E, Engel D, Star K, Monson K, Brandi J, Felberg LE, Brookes DH, Wilson L, Chen
J, Liles K, Chun M, Li P, Gohara DW, Dolinsky T, Konecny R, Koes DR, Nielsen JE,
Head-Gordon T, Geng W, Krasny R, Wei GW, Holst MJ, McCammon JA, Baker NA.
2018. Improvements to the APBS biomolecular solvation software suite. Protein Sci
27:112-128.

103. Singer JB, Thomson EC, Hughes J, Aranday-Cortes E, McLauchlan J, da Silva Filipe A,
Tong L, Manso CF, Gifford RJ, Robertson DL, Barnes E, Ansari MA, Mbisa JL, Bibby
DF, Bradshaw D, Smith D. 2019. Interpreting Viral Deep Sequencing Data with GLUE.
Viruses 11.

104. Singer JB, Thomson EC, McLauchlan J, Hughes J, Gifford RJ. 2018. GLUE: a flexible
software system for virus sequence data. BMC Bioinformatics 19:532.

105. Larsson A. 2014. AliView: a fast and lightweight alignment viewer and editor for large
datasets. Bioinformatics 30:3276-8.

Figure 1. D256N and D270N compensatory substitutions in HIV-1 IN suppress the replication defect of the HIV-1$_{NL4-3}$ $_{IN\ (R269A/K273A)}$ class II mutant virus. (A) MT4-LTR-GFP indicator cells were infected with HIV-1$_{NL4-3}$ at an MOI of 2 i.u./cell or an equivalent particle number of the HIV-1$_{NL4-3}$ $_{IN\ (R269A/K273A)}$ class II IN mutant virus (based on reverse transcriptase (RT) activity) as detailed in Materials and Methods. HIV-1$_{NL4-3}$ $_{IN\ (R269A/K273A)}$ viruses were serially passaged for three rounds until the emergence of compensatory mutations that allowed virus replication with WT kinetics. The graphs represent the percentage of GFP positive cells as assessed by flow cytometry over three passages at the indicated days post-infection (dpi). **(B)** HIV-1 genomic RNA was isolated from viruses collected from cell culture supernatants over the three passages (i.e. P1, P2, P3) and at the indicated days post-infection (i.e. D29, D40, etc.) as described in Materials and Methods. Heatmap shows the percentage of IN D256N and D270N substitutions at the indicated passages and days post-infection (dpi) as assessed by whole-genome deep sequencing. **(C)** HEK293T cells were transfected with full-length pNL4-3 expression plasmids carrying *pol* mutations coding for the IN D256N, D270N and D256N/D270N (D2N) substitutions introduced on the WT IN and IN R269A/K273A backbones. Cell lysates and cell culture supernatants containing virions were collected two days post-transfection and analyzed by immunoblotting for CA and IN. The image is representative of five independent experiments. See Fig. S1 for quantitative analysis of immunoblots. **(D)** HEK293T cells were

transfected as in (C) and cell culture supernatants containing viruses were titered on TZM-bl

indicator cells, whereby virus replication was limited to a single cycle by addition of dextran

sulfate (50 µg/mL). The titers were normalized relative to particle numbers as assessed by an

RT activity assay and are presented relative to WT (set to 1). Also see Fig. S1 for titer and RT

activity values prior to normalization. The columns represent the average of 5-6 independent

biological replicates and the error bars represent standard error of the mean (SEM)

(****$p<0.0001$, **$p<0.01$, *$p<0.05$ by one-way ANOVA multiple comparison test with Dunnett's

correction).

Figure 2. D256N and D270N substitutions restore IN-gRNA binding and accurate virion

maturation for the HIV-1$_{NL4-3\ IN\ (R269A/K273A)}$ class II mutant virus. (A) Autoradiogram of IN-RNA

adducts immunoprecipitated from virions bearing the indicated substitutions in IN. Immunoblots

below show immunoprecipitated (IP) IN or CA protein in lysates prior to immunoprecipitation.

Data shown are representative of three independent experiments. See also Fig. S2 for

quantitative analysis of autoradiographs. **(B)** Examination of virion maturation in WT and IN

mutant viruses by thin section electron microscopy (TEM). Data show quantification of virion

morphologies across 100 particles for each sample and replicate experiment. Data show the

average of two independent biological replicates and error bars represent the SEM (***$p<0.001$,

**$p<0.01$, ns=not significant by one-way ANOVA multiple comparison test with Dunnett's

correction).

Figure 3. Restoring the local net charge of IN-CTD restores RNA binding and infectivity

for the IN R269A/K273A class II mutant. (A) Schematic diagram of IN and sequence of CTD

residues with basic and acidic amino acids highlighted in blue and red, respectively. **(B, C)**

HEK293T cells were transfected with full-length proviral WT HIV-1$_{NL4-3}$ expression plasmid or its

derivatives carrying *pol* mutations encoding for D2N, D256R, D270R and D256R/D270R (D2R)

IN substitutions on the backbone of HIV-1$_{NL4-3\ IN\ (R269A/K273A)}$. **(B)** Cell lysates and purified virions

harvested two days post-transfection were analyzed by immunoblotting for CA and IN.

Representative image from five independent experiments is shown. See also Fig. S3A-E for

quantitative analysis of immunoblots. **(C)** Cell culture supernatants containing viruses were

titered on TZM-bl indicator cells, whereby virus replication was limited to a single cycle by

addition of dextran sulfate (50 µg/mL). The titers were normalized relative to particle numbers as

assessed by an RT activity assay and are presented relative to WT (set to 1). Also see Fig.

S3F, G for titer and RT activity values prior to normalization. The columns represent the

average of six independent experiments and the error bars represent SEM (****$p<0.0001$,

*$p<0.05$ by one-way ANOVA multiple comparison test with Dunnett's correction). **(D)**

Autoradiogram of IN-RNA adducts immunoprecipitated from HIV-1$_{NL4-3}$ virus particles bearing

WT IN or the indicated IN mutants. Immunoblot below shows immunoprecipitated (IP) IN

protein. Results are representative of three independent replicates. See Fig. S3H for

quantitative analysis of autoradiographs. **(E)** MT4 T-cells were infected with HIV-1 $_{NL4-3\ IN\ (D116N)}$

viruses that were trans-complemented with the indicated Vpr-IN mutant proteins as described in

Materials and Methods. Virion release was assessed by RT activity assays over the course of 5

days post-infection. Y-axis indicates fold increase in RT activity over day 0. Data are from 3

independent replicates, error bars show the SEM.

Figure 4. D256R and D256K substitutions restore IN-RNA binding and infectivity for the

HIV-1$_{NL4-3\ IN\ (R262A/R263A)}$ class II mutant virus. (A-E) HEK293T cells were transfected with

proviral HIV-1$_{NL4-3}$ expression plasmids carrying *pol* mutations for the indicated IN substitutions

on WT IN or IN R262A/R263A backbones. **(A, C)** Cell lysates and purified virions collected two

days post-transfection were analyzed by immunoblotting for CA and IN. See also Fig. S4 for

quantitative analyses of immunoblots. **(B, D)** Cell culture supernatants containing viruses were

titered on TZM-bl indicator cells, whereby virus replication was limited to a single cycle by

addition of dextran sulfate (50 µg/mL). The titers were normalized relative to particle numbers as

assessed by an RT activity assay and are presented relative to WT (set to 1). See also Fig.

S4F, L, M for titer and RT activity values prior to normalization. The columns represent the

average of four-to-six independent experiments, and the error bars represent SEM (**p<0.01,

*p<0.05, ns=not significant by one-way ANOVA multiple comparison test with Dunnett's

correction). **(E)** Autoradiogram of IN-RNA adducts immunoprecipitated from WT or IN mutant

HIV-1$_{NL4-3}$ virions. The amount of immunoprecipitated IN was assessed by the immunoblot

shown below. Immunoblots and CLIP autoradiographs are representative of three independent

replicates. See Fig. S4N for quantitative analysis of autoradiographs. **(F)** MT4 T-cells were

infected with HIV-1 $_{NL4-3\ IN\ (D116N)}$ viruses that were trans-complemented with the indicated Vpr-IN

mutant proteins as described in Materials and Methods. Virion release was assessed by RT

activity assays over the course of 5 days post-infection. Y-axis indicates fold increase in RT

activity over day 0. Data are from 3 independent replicates, error bars show the SEM.

Figure 5. D279R substitution increases and D278R/D279R substitutions restore IN-RNA

binding and infectivity for the HIV-1$_{NL4-3\ IN\ (R269A/K273A)}$ virus. HEK293T cells were transfected

with full-length pNL4-3 expression plasmids carrying mutations coding for the IN D279R,

D278R/D279R (D2R), D256I and D256I/D270I (D2I) substitutions introduced on the IN

R269A/K273A backbone. **(A)** Cell culture supernatants containing viruses were titered on TZM-

bl indicator cells, whereby virus replication was limited to a single cycle by addition of dextran

sulfate (50 µg/mL). The titers were normalized relative to particle numbers by an RT activity

assay and are presented relative to WT (set to 1). The columns represent the average of 4

independent biological replicates and the error bars represent standard error of the mean (SEM)

(*p<0.05 by one-way ANOVA multiple comparison test with Dunnett's correction). **(B)** Cell

lysates and purified virions collected two days post transfection were analyzed by

immunoblotting for CA and IN. The image is representative of four independent experiments.

(C) IN-RNA adducts immunoprecipitated from WT or IN mutant HIV-1$_{NL4-3}$ virions per CLIP

protocol were analyzed by autoradiography. Graph shows the quantification of IN-RNA adducts

from three independent biological replicates, error bars show the SEM.

Figure 6. Electrostatic potential maps of HIV-1 IN bearing class II and compensatory

mutations. (A-G) Electrostatic potential maps of the indicated HIV-1 IN mutants derived from

the crystal structure of Gupta et al. (62) is depicted. Results are displayed as an electrostatic

potential molecular surface. The low, mid, and high range values are -5, 0, and 5, respectively.

(**H**) A cartoon view of the CTD structure (62) is shown with the C-terminal tail (aa 261-275) and

the loop (aa 252-257) colored in cyan and orange, respectively. Side chains of indicated

residues are shown.

Figure 7. Assessing multimerization properties of IN in mutant viruses. (A) Purified WT or

IN mutant HIV-1$_{NL4-3}$ virions were treated with 1 mM EGS, and virus lysates analyzed by

immunoblotting using antibodies against IN following separation on 3-6% Tris-acetate gels as

detailed in Materials and Methods. The position of monomers (M), dimers (D), and tetramers (T)

are indicated by arrows in a representative western blot. **(B, C)** Quantification of IN

multimerization in virions from experiments conducted as in A. Error bars show the SEM from

three independent experiments. **(D-H)** Representative SEC traces for indicated recombinant IN

proteins. The X-axis indicates elution volume (mL) and Y-axis indicates the intensity of

absorbance (mAU). Tetramers (T), dimers (D), and monomers (M) are indicated. **(I)** Summary of

mutant INs bridging TAR RNA compared to WT IN. Alpha screen counts at 320 nM for each

protein is shown. The graphs show average values of three independent experiments and the

error bars indicate standard deviation.

Figure 8. Secondary site suppressors of the IN R269A/K273A mutant increase

susceptibility to ALLINIs. (A, B, D, E) Titers of viruses bearing the indicated substitutions in IN

produced from HEK293T cells at different concentrations of ALLINIs, BI-B2 or BI-D. WT or IN

mutant HIV-1$_{NL4-3}$ in cell culture supernatants were titered on TZM-bl indicator cells. The titer

values are represented relative to the mock control of each mutant (set to 1). Data are from two

independent biological replicates. **(C, F)** Autoradiogram of IN-RNA adducts immunoprecipitated

from WT or IN mutant HIV-1$_{NL4-3}$ virions produced from HEK293T cells in the presence of 0.1µM

of BI-D. The amount of immunoprecipitated IN protein was visualized by the immunoblot shown

below. Immunoblots and CLIP autoradiographs results are a representative of three

independent replicates. **(G)** The GSK-1264 binding pocket between CCD-CCD (brown) and

CTD (light blue) from different IN subunits is shown (62) (PDB ID: 5HOT). GSK-1264 is in green

with nitrogen and oxygen atoms colored blue and red, respectively. The side chain of R269 is

shown with nitrogen atoms colored in dark blue. Interactions between CTD R269 and CCD

Q168 are indicated by dashed lines.

Figure 9. Characterization of IN mutations present in latently infected CD4+ T-cells. (A)

Frequency of IN substitutions at amino acids 224, 230, 246 and 272 derived from HIV-1

sequences from latently infected CD4+ T-cells is shown. **(B)** HEK293T cells were transfected

with proviral HIV-1$_{NL4-3}$ expression plasmids carrying the R224Q, S230N, E246K, and G272R IN

mutations. Cell lysates and virions were purified two days post transfection and analyzed by

immunoblotting for CA and IN. The image is representative of two independent experiments. **(C)**

WT or IN mutant HIV-1L4-3 viruses in cell culture supernatants were titered on TZM-bl indicator

cells. The titers are presented relative to WT (set to 1). The columns represent the average of

three independent experiments and the error bars represent SEM (*$p<0.05$, by one-way ANOVA

multiple comparison test with Dunnett's correction). **(D)** Autoradiogram of IN-RNA adducts

immunoprecipitated from virions bearing the indicated substitutions in IN. Immunoblots below

show the amount of immunoprecipitated IN.

1078

1079

1080

1081

1082

1083

1084

SUPPLEMENTAL FIGURE LEGENDS

Figure S1. Effect of compensatory D256N and D270N substitutions on viral gene

expression, Gag processing, particle release and infectivity for the WT and HIV-1$_{NL4-3\ IN}$

$_{(R269A/K273A)}$ class II mutant virus. HEK293T cells were transfected with full-length proviral pNL4-

3 expression plasmids carrying mutations coding for the IN D256N, D270N and D256N/D270N

(D2N) substitutions introduced on the WT IN and IN R269A/K273A backbones as described in

Figure 1C. Cell lysates and purified virions were analyzed by immunoblotting for CA and IN.

Quantification of unprocessed Gag-Pr55 **(A)**, Gag processing intermediates **(B-D)**, virion-

associated IN levels **(E)** from 4 independent biological replicate experiments is shown. Error

bars show the SEM. **(F, G)** Cell culture supernatants containing viruses were collected and used

in an RT activity assay **(F)** or titered on TZM-bl indicator cells **(G)**, whereby virus replication was

limited to a single cycle by addition of dextran sulfate (50 μg/mL). The RT activity and titer

values were normalized relative to WT (set to 1).

Figure S2. D256N and D270N substitutions restore RNA binding for the HIV-1$_{NL4-3\ IN}$

$_{(R269A/K273A)}$ class II mutant virus. IN-RNA adducts were immunoprecipitated from virions

bearing the indicated substitutions in IN as in Fig. 2A. Autoradiography images were quantified

for IN-RNA adducts and presented relative to WT (set to 1). Data are derived from three

independent biological replicates with error bars showing the SEM. (****p<0.0001, ns=non-

significant by one-way ANOVA multiple comparison test with Dunnett's correction).

Figure S3. Effect of compensatory D256R and D270R substitutions on viral gene

expression, Gag processing, particle release, infectivity and IN-RNA binding for the WT

and HIV-1$_{NL4-3 IN (R269A/K273A)}$ class II mutant viruses. HEK293T cells were transfected with full-

length proviral pNL4-3 expression plasmids carrying mutations coding for the IN D2N, D256R,

D270R and D256R/D270R (D2R) substitutions introduced on the IN R269A/K273A backbone as

described in Fig. 3B, C. Cell lysates and purified virions were analyzed by immunoblotting for

CA and IN. Quantification of unprocessed Gag-Pr55 **(A)**, Gag processing intermediates **(B-D)**,

virion-associated IN levels **(E)** from 4 independent biological replicate experiments is shown.

Error bars show the SEM. **(F, G)** Cell culture supernatants containing viruses were collected

and used in an RT activity assay **(F)** or titered on TZM-bl indicator cells **(G)**, whereby virus

replication was limited to a single cycle by addition of dextran sulfate (50 µg/mL). The RT

activity and titer values were normalized relative to WT (set to 1). **(H)** IN-RNA adducts were

immunoprecipitated from virions bearing the indicated substitutions in IN as in Fig. 3D.

Autoradiography images were quantified for IN-RNA adducts and presented relative to WT (set

to 1). Data are derived from three independent biological replicates with error bars showing the

SEM. (***p<0.001, **p<0.01 by one-way ANOVA multiple comparison test with Dunnett's

correction).

Figure S4. Effect of compensatory substitutions on viral gene expression, Gag

processing, particle release, infectivity and IN-RNA binding for the WT and HIV-1$_{NL4-3 IN}$

$_{(R262A/R263A)}$ class II mutant viruses. HEK293T cells were transfected with full-length proviral

pNL4-3 expression plasmids carrying mutations coding for the IN D256N, D270N, D2N, D256K

and D256R substitutions introduced on the WT IN or IN R262/R263A backbones as described

in Fig. 4. Cell lysates and purified virions were analyzed by immunoblotting for CA and IN.

Quantification of unprocessed Gag-Pr55 **(A, G)**, Gag processing intermediates **(B, C, D, H, I, J)**,

virion-associated IN levels **(E, K)** from 4 independent biological replicates is shown. **(F, L, M)**

Cell culture supernatants containing viruses were collected and titered on TZM-bl indicator cells

(F, M) or used in an RT activity assay **(L)** to assess particle release. The RT activity and titer

values were normalized relative to WT (set to 1). **(N)** IN-RNA adducts were immunoprecipitated

from virions bearing the indicated substitutions in IN as in Fig. 4E. Autoradiography images

were quantified and presented relative to WT (set to 1). Data are derived from three

independent biological replicates with error bars showing the SEM. (****$p<0.0001$, **$p<0.01$,

*$p<0.05$, ns=not significant by one-way ANOVA multiple comparison test with Dunnett's

correction).

1137

Figure 1

Figure 2

A

B

C

D

E

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Figure 9