Title: Emergence of compensatory mutations reveal the importance of electrostatic

interactions between HIV-1 integrase and genomic RNA

Short title: Molecular basis of HIV-1 integrase-gRNA interactions

Authors: Christian Shema Mugisha[1], Tung Dinh[2], Kasyap Tenneti[1], Jenna E. Eschbach[1],

Keanu Davis[1], Robert Gifford[3], Mamuka Kvaratskhelia[2], Sebla B. Kutluay[1]

[1] Department of Molecular Microbiology, Washington University School of Medicine, Saint Louis,

MO 63110, USA

[2] Division of Infectious Diseases, University of Colorado School of Medicine, Aurora, CO 80045

[3] MRC-University of Glasgow Centre for Virus Research, 464 Bearsden Rd., Bearsden,
Glasgow G61 1QH, UK.
13

15 **Correspondence: kutluay@wustl.edu**

ABSTRACT

Independent of its catalytic activity, HIV-1 integrase (IN) enzyme regulates proper particle maturation by binding to and packaging the viral RNA genome (gRNA) inside the mature capsid lattice. Allosteric integrase inhibitors (ALLINIs) and class II IN substitutions inhibit the binding of IN to the gRNA and cause the formation of non-infectious virions characterized by mislocalization of the viral ribonucleoprotein complexes between the translucent conical capsid lattice and the viral lipid envelope. To gain insight into the molecular nature of IN-gRNA interactions, we have isolated compensatory substitutions in the background of a class II IN (R269A/K273A) variant that directly inhibits IN binding to the gRNA. We found that additional D256N and D270N substitutions in the C-terminal domain (CTD) of IN restored its ability to bind gRNA and led to the formation of infectious particles with correctly matured morphology. Furthermore, reinstating the overall positive electrostatic potential of the CTD through individual D256R or D256K substitutions was sufficient to restore IN-RNA binding and infectivity for the R269A/K273A as well as the R262A/R263A class II IN mutants. The compensatory mutations did not impact functional IN oligomerization, suggesting that they directly contributed to IN binding to the gRNA. Interestingly, HIV-1 IN R269A/K273A, but not IN R262A/R263A, bearing compensatory mutations was more sensitive to ALLINIs providing key genetic evidence that specific IN residues required for RNA binding also influence ALLINI activity. Structural modeling provided further insight into the molecular nature of IN-gRNA interactions and ALLINI mechanism of action. Taken together, our findings highlight an essential role of IN-gRNA interactions for proper virion maturation and reveal the importance of electrostatic interactions between the IN CTD and the gRNA.

AUTHOR SUMMARY

In addition to its well-defined catalytic function, HIV-1 integrase (IN) binds to the viral RNA genome

and regulates proper virion maturation. Inhibition of IN binding to the HIV-1 genome through

mutations of positively charged residues within the C-terminal domain (CTD, i.e. R269, K273)

results in non-infectious particles in which the viral genomes are mislocalized in improperly

matured virions. Here we have isolated compensatory mutations in the background of the class

II IN (R269A/K273A) mutant virus that restored the ability of IN to bind RNA. We found that

additional substitutions of nearby acidic residues (i.e. D256 and D270), which restored the overall

positive charge of the CTD, rescued the ability of IN to bind RNA and thus resulted in formation

of correctly matured, infectious virions. These compensatory substitutions also revealed the role

of specific residues within the CTD that determine sensitivity to allosteric integrase inhibitors

(ALLINIs), a class of compounds that indirectly target IN-RNA interactions. Taken together, our

findings reveal the importance of the electrostatic interactions between the IN CTD and the gRNA

and provide key genetic evidence for a crucial role of the CTD in antiviral activity of ALLINIs.

INTRODUCTION

A defining feature of retroviruses is the reverse transcription of the viral RNA genome (gRNA) and integration of the resultant linear viral DNA into a host chromosome, which establishes lifelong infection. The latter reaction is mediated by the viral integrase (IN) enzyme, which catalyzes 3' processing and DNA strand transfer reactions [1]. The catalytic activity of HIV-1 IN has been successfully targeted by several integrase strand-transfer inhibitors (INSTIs) [2-7] that have become key components of frontline anti-retroviral therapy regimens due to their high efficacy and tolerance profiles [8-11]. In addition, HIV-1 IN has an emerging non-catalytic function in virus replication [12-16]. Successful targeting of this second function can complement the existing antiviral regimens and substantially increase the barrier to INSTI resistance.

HIV-1 IN consists of three independently folded protein domains: the N-terminal domain (NTD) bears the conserved His and Cys residues (HHCC motif) that coordinate Zn^{2+} binding for 3-helix bundle formation; the catalytic core domain (CCD) adopts an RNase H fold and harbors the enzyme active site composed of an invariant DDE motif, and C-terminal domain (CTD) which adopts an SH3 fold [17, 18]. Integration is facilitated by a cellular co-factor, lens epithelium-derived growth factor (LEDGF/p75), which binds tightly to a site within the CCD dimer interface [19, 20] and guides the preintegration complex to actively transcribed regions of the host chromosome [19-24]. A group of pleotropic IN substitutions distributed throughout IN, collectively known as class II mutations, disrupt viral assembly [13, 16, 25-36], morphogenesis [12, 16, 27, 32-34, 37, 38] and reverse transcription in target cells [12-14, 16, 29, 31, 32, 34, 36-54] often without obstructing the catalytic activity of IN in vitro [13, 27, 28, 31, 40, 41, 44, 46, 55-57]. A hallmark of class II IN mutant viruses is the mislocalization of the viral ribonucleoprotein complexes (vRNP) outside of the viral capsid (CA) lattice, a deformation which is often referred to as eccentric morphology [12, 15, 16, 27, 32-34, 37, 38, 58-60]. Although originally designed to inhibit integration through preventing the binding of IN to LEDGF/p75 [58, 61-65], allosteric

integrase inhibitors (ALLINIs) potently inhibit proper virion maturation [12, 37, 61] and lead to the

formation of virions that display a similar eccentric morphology observed with class II IN mutations

[12].

We have recently shown that binding of IN to the gRNA in mature virions accounts for the non-

catalytic function of IN in virus replication and is required for proper encapsidation of the viral

ribonucleoproteins (vRNPs) inside the mature CA lattice [38]. Class II IN substitutions and

ALLINIs block IN-gRNA binding, thus causing formation of virions with an eccentric morphology

[15, 38]. IN binds to multiple distinct locations on the gRNA, including the TAR hairpin present

within the 5' and 3'UTRs of the gRNA and constitutes a high affinity binding site [38]. IN

preferentially binds to the gRNA in a tetrameric state, and many class II IN mutations block IN-

gRNA binding by disrupting the functional oligomerization of IN [15]. Mutation of basic residues

within IN-CTD (i.e. R262, R263, R269 and K273) inhibits IN-gRNA binding without altering

functional IN oligomerization in virions and in vitro [15, 38], suggesting that these residues are

directly involved in binding to the gRNA. On the other hand, the precise mechanism of how these

residues mediate recognition of specific sequence elements on the gRNA remains unknown. For

example, it is possible that the positively charged Lys and Arg residues interact with the negatively

charged RNA phosphate backbone in a non-specific or semi-specific manner, depending on the

folding and structure of the cognate RNA element, driven by electrostatic interactions [66-68].

Alternatively, these residues can mediate specific interactions with RNA targets through H-

bonding and van der Waals contacts with individual nucleobases [66-68].

To gain insight into the mode of IN-gRNA interactions, the non-infectious IN R269A/K273A class

II mutant virus was serially passaged in T-cells until the acquisition of compensatory mutations.

We found that two compensatory mutations, D256N and D270N, within the IN-CTD restored virion

infectivity, IN-RNA interactions and accurate virion morphogenesis. As the D-to-N mutations

resulted in loss of two negative charges, possibly overcoming the loss of two positive charges

with the R269A/K273A class II substitutions, we tested whether restoring the overall charge of

CTD through other mutations would restore IN-gRNA binding and virion infectivity. Indeed, the

D256R substitution alone restored virion infectivity and RNA binding for the IN R269A/K273A

mutant. We further extended these findings to another class II mutant, R262A/R263A, which was

similarly suppressed by the D256R as well as D256K substitutions. Compensatory mutations did

not affect the ability of IN to multimerize *in vitro* or in virions, suggesting that they restored the

RNA-binding ability of IN directly. Interestingly, the IN R269A/K273A, but not the IN

R262A/R263A, mutant viruses bearing the compensatory mutations had increased sensitivity to

ALLINIs, providing key genetic evidence that specific residues within the CTD, which are required

for RNA binding, also contribute to ALLINI mechanism of action. Together, our findings strongly

suggest that IN-RNA interactions are at least in part driven by electrostatic interactions between

the basic residues within IN-CTD and phosphate backbone of the gRNA and highlight that

ALLINIs engage CTD residues to inhibit RNA binding.

RESULTS

Compensatory IN D256N/D270N substitutions emerge in the background of the IN

R269A/K273A class II mutant virus

The R269A/K273A class II IN substitutions obstruct IN-gRNA binding directly without interfering

with IN multimerization and result in the formation of particles with eccentric morphology [15, 16].

To better understand the molecular basis of how R269 and K273 residues mediate IN-gRNA

binding, viruses bearing the IN R269A/K273A mutations were serially passaged in MT-4 cells until

the emergence of compensatory mutations at the end of passage 3, which completely restored

virion infectivity (**Fig. 1A**). Deep sequencing of full-length gRNA isolated from virions across the

three passages revealed that viruses retained the IN R269A/K273A substitutions while

sequentially acquiring D256N and D270N mutations in IN (**Fig. 1B**). The D256N substitution

emerged at the end of passage 1 whereas the D270N mutation emerged later in passage 2 (**Fig.**

1B). Both mutations were fixed by the end of passage 2 and no other mutations were observed

elsewhere on the viral genome.

The IN D256N and D270N mutations were introduced into the replication-competent pNL4-3

molecular clone bearing WT or R269A/K273A IN. Introduction of D256N and D270N substitutions

in IN had no observable effect on Gag (Pr55) expression or processing in cells or particle release

(**Fig. 1C**). Introduction of D256N and D270N mutations either individually or together (D2N) on

the WT IN backbone did not affect viral titers (**Fig. 1D**). Remarkably, while the individual D256N

and D270N mutations in the R269A/K273A IN backbone increased virus titers by 5-10-fold, the

D2N substitutions increased virus titers by 100-fold (**Fig. 1D**). Overall, these results demonstrate

that the combination of D256N and D270N mutations are sufficient to restore the replication

competency of the R269A/K273A class II IN mutant virus.

D256N/D270N substitutions restore IN-gRNA binding for the R269A/K273A class II IN

mutant viruses and lead to formation of correctly matured virions

We next assessed whether D256N and D270N substitutions rendered the R269A/K273A class II

IN mutant virus replication competent by restoring IN-gRNA binding and proper virion maturation.

To this end, IN-gRNA complexes were immunoprecipitated from UV-crosslinked virions and

visualized per CLIP protocol as described previously [38, 69, 70]. Equivalent amounts of

immunoprecipitated IN from WT and R269A/K273A IN viruses, or those additionally bearing the

IN D256N and D270N substitutions, were analyzed for their ability to bind RNA. IN-RNA

complexes were readily visible for WT viruses as well as D256N, D270N and D2N mutants

(introduced on the WT backbone), but not from the R269A/K273A class-II IN mutant viruses (**Fig.**

2A). Introduction of the D256N substitution on R269A/K273A IN backbone modestly enhanced

the ability of IN to bind RNA, whereas the D270N substitution had no observable impact (**Fig. 2A**).

In contrast, the D2N substitution substantially enhanced the ability of R269A/K273A IN to bind

RNA (**Fig. 2A**). The virion morphology of WT and IN mutant viruses was assessed by transmission

electron microscopy (TEM). As expected, more than 80% of WT particles had an electron-dense

condensate that represents vRNPs inside the CA lattice, whereas the majority of R269A/K273A

class II IN mutant virions (~68%) had a clear eccentric morphology (**Fig. 2B**). Consistent with

effects on virus titers and RNA-binding, the introduction of D256N/D270N substitutions restored

the ability of the R269A/K273A IN mutant to form properly mature virions (**Fig. 2B**). Cumulatively,

these data show that D256N and D270N IN substitutions restore infectivity for the R269A/K273A

class II IN mutant virus through reestablishing RNA binding and subsequently accurate virion

maturation.

Electrostatic interactions are required for IN-gRNA binding.

IN-CTD is decorated with several acidic and basic amino acids resulting in a net charge of +3

(**Fig. 3A**). The D2N substitutions in effect restored the overall charge of the IN R269A/K273A

CTD, suggesting that the net electrostatic charge of the IN-CTD may be a key parameter in gRNA

binding. To test this hypothesis, we investigated whether restoring the overall charge of IN-CTD

through other mutations would also restore RNA binding and infectivity for the R269A/K273A IN

virus. We focused our analysis on D256 and D270 residues for the following reasons: i) these

amino acids were amenable to substitutions during virus passaging experiments; ii) D256N and

D270N substitutions in the context of WT HIV-1 yielded infectious virions, suggesting that these

compensatory mutations do not significantly contribute to the catalytic activity of IN thus allowing

us to specifically probe their roles for IN-RNA interactions and virion maturation. To extend these

studies we introduced D256R, D270R and D256R/D270R (D2R) substitutions into the HIV-1$_{NL4-3}$

$_{IN R269A/K273A}$ backbone and transfected HEK293T cells with the resulting plasmids. Cell lysates and

cell-free virions were then analyzed for Gag processing, particle release, and infectivity. Overall,

D-to-R substitutions had no major effect on Gag (Pr55) expression or processing in cells (**Fig.**

3B).

Remarkably, the D256R substitution alone increased the titers of the R269A/K273A class II IN

mutant virus by 100-fold and to a level comparable to that of WT viruses (**Fig. 3C**). In contrast,

D270R substitution only had a modest impact on virus titers and the D2R substitution increased

virion infectivity at a modestly lower level than that of D256R and D2N (**Fig. 3C**). Importantly, the

D256R and the D2R substitutions completely restored IN-gRNA binding (**Fig. 3D**), demonstrating

that the increase in viral titers with the D256R and D2R substitutions correlates well with

enhancement of IN-gRNA binding. Further assessment of these substitutions with Vpr-IN

transcomplementation assays revealed that none of the IN substitutions impacted the catalytic

activity of IN, except for the D2R substitution (**Fig. 3E**), suggesting that the inability of the D270R

mutation to restore infectivity may be due to its adverse effects on the catalytic activity of IN.

Taken together, these findings indicate that restoring the overall charge of IN-CTD through D256R

mutation reestablishes virion infectivity and RNA binding for the IN R269A/K273A class II mutant.

D256R substitution fully restores virion infectivity and RNA binding for a separate class II

IN (R262A/R263A) mutant

We have previously shown that mutation of other basic residues within the IN-CTD (i.e.

R262A/R263A) also directly inhibit IN-gRNA binding without compromising functional

oligomerization of IN [15, 38]. We next wanted to extend our observations and test how

substitutions of D256 and D270 residues affect IN-gRNA binding and infectivity in the background

of the IN R262A/R263A mutant virus. Introducing the D256N, D270N and D2N compensatory

mutations on the IN R262A/R263A backbone did not affect Gag expression, processing, and

virion release (**Fig. 4A**). While the D256N substitution increased virion infectivity by 10-fold, the

D270N substitution had no impact and the D2N substitution had an intermediate phenotype (**Fig.**

4B), demonstrating the context dependency of these compensatory mutations. The D256K and

D256R substitutions introduced in the R262A/R263A IN backbone also had minimal effects on

Gag expression, processing, and virion release (**Fig. 4C**). In contrast, the D256K substitution

increased viral titers at a greater degree than D256N and the D256R substitution completely

restored virion infectivity (**Fig. 4D**). In line with the titer data, D256K significantly increased RNA

binding whereas the D256R completely restored it in the context of the IN R262A/R263A mutant

virus (**Fig 4E**). D256N, D256K and D256R IN successfully transcomplemented a class I IN mutant,

suggesting that they did not distort the catalytic activity of IN (**Fig 4F**). Taken together, these

findings indicate that restoring the overall positive charge of IN-CTD through D256R or D256K

mutations is sufficient to restore virion infectivity and RNA binding to class II IN mutants that

directly inhibit RNA binding.

We next used reported X-ray structure [62] and molecular modelling to visualize how class II and

compensatory IN mutations affect electrostatic potential of the CTD surface. Mutation of R269

and K273 residues expectedly resulted in a substantial loss of a basic patch in IN (**Fig. 5A, B**).

Both the D256R and D256N substitutions resulted in more positively charged surface distal from

the R269/K273A residues (**Fig. 5C, D**), suggesting that compensatory mutations likely created an

additional interacting interface with gRNA binding. A similar outcome was observed when class II

R262A/R263A and compensatory D256R changes were introduced in the CTD (**Fig. 5E-G**). While

the D256R change is substantially distanced from R262/R263 and R269/K273 residues, we note

the following. R262, R263, R269, and K273 are positioned within the same highly flexible C-

terminal tail (aa 261-275), whereas D256 belongs to another, shorter (aa 252-257) loop (**Fig. 5H**).

The highly pliable nature of the tail and the loop could be crucial for IN to optimally engage with

cognate RNA as well as allow for emergence of compensatory mutations at alternative sites

positioned in these IN segments.

Effects of the compensatory mutations on functional oligomerization of IN and IN-RNA

interactions.

As functional IN oligomerization is a prerequisite for RNA binding [15], we next examined how the

compensatory substitutions affected IN oligomerization. For in virion analysis, purified HIV-1$_{NL4-3}$

virions were treated with ethylene glycol bis (succinimidyl succinate) (EGS) to covalently crosslink

IN in situ and virus lysates were analyzed by immunoblotting. As in WT virions, IN species that

migrated at molecular weights consistent with those of monomers, dimers, trimers, and tetramers

were readily distinguished in R269A/K273A or R262A/R263A viruses with additional

compensatory mutations (D256N/D270N, D256R, D256K, and D256R/D270R) but not with the

canonical class II IN mutant V165A that is unable to form functional oligomers (**Fig 6A-C**).

Complementary in vitro assessment of purified recombinant IN proteins by size exclusion

chromatography (SEC) revealed that D2N and D256R substitutions in the background of

R269A/K273A or the R262A/R263A class II mutants did not impact functional IN tetramerization

(**Fig. 6D-H**).

We have previously shown that recombinant IN binds to TAR RNA with high affinity and provides

a nucleation point to bridge and condense RNA [38]. We next examined the ability of class II

mutant INs bearing compensatory mutations to bind and bridge TAR RNA. Consistent with

findings from CLIP, the D2N and D256R substitutions also enhanced or restored the ability of

R269A/K273 and R262A/R263A mutants to bridge between RNA molecules (**Fig. 6I**). Together,

these data demonstrate that the compensatory mutations directly restore the ability of IN to bind

RNA without altering functional IN oligomerization.

Sensitivity to ALLINIs is determined by distinct residues within the CTD

ALLINIs potently disrupt proper virion maturation through inducing aberrant IN multimerization

and consequently inhibiting IN-gRNA binding [12, 38]. Recent structural studies have shown that

the ALLINI, GSK-1264, can directly engage residues within the IN-CTD through its tert-butoxy

and carboxylic acid moieties and induce open polymers [62]. These findings are in line with

previous biochemical and modeling studies that also showed the involvement of the IN-CTD, in

particular residues K264 and K266, in ALLINI-induced aberrant IN multimerization [71, 72]. Based

on these prior findings, we wanted to test how adjacently positioned IN R262A/R263A and

R269A/K273A substitutions and the compensatory mutations affected ALLINI activities.

To this end, we examined effects of representative quinoline-based ALLINIs, BI-D or BI-B2, on

the viruses bearing the class II and compensatory mutations. While the titers of WT viruses only

decreased at ALLINI concentrations greater than 1 μM, the titers of IN R269A/K273A viruses

bearing D2N, D256R and D2R substitutions were significantly reduced by ALLINIs at

concentrations as low as 0.1 μM (**Fig. 7A, B**). In contrast, viruses bearing the

R262A/R263A/D256R IN were less sensitive to low concentrations of ALLINIs (**Fig. 7D, E**),

suggesting that the R269A and/or K273A mutations underlie the increased sensitivity to ALLINIs.

The increased sensitivity to ALLINIs correlated well with inhibition of RNA binding. While 0.1 μM

of the ALLINI, BI-D, did affect the level of IN-gRNA binding in WT viruses, it significantly reduced

IN-gRNA binding in viruses with class II compensatory mutations on the R269A/K273A but not

R262A/R263A backbone (**Fig. 7C,F**). Taken together, our findings provide key genetic and

virological evidence that specific CTD residues required for RNA binding are also crucial for the

ALLINI mechanism of action.

Characterization of IN mutations present in latently infected cells

Persistence of HIV-1 in memory CD4+ T-cells as latent proviruses constitutes a major barrier to

HIV-1 cure. Although the majority of HIV-1 proviruses in these cells are defective [73], recent

evidence suggests that defective proviruses can be transcribed into RNAs that are spliced,

translated and can be recognized by HIV-1-specific cytotoxic T lymphocytes [74]. We decided to

characterize IN mutations isolated from latently infected cells, given the possibility that class II IN

mutations existing in latently infected cells can result in the formation of defective particles that

may subsequently modulate immune responses. Though relatively uncommon, we found the

presence of R224Q, S230N, E246K and G272R substitutions in IN-CTD (**Fig. 8A**). Of note, only

the R224Q substitution resulted in loss of a positive charge, whereas the E246K and G273R

substitutions resulted in gain of positive charge. These mutations were introduced into the NL4-3

proviral backbone with minimal effects on Gag expression and particle release (**Fig. 8B**). Although

the E246K virus was significantly less infectious (**Fig. 8C**), we did not find any evidence for

inhibition of IN-gRNA binding (**Fig. 8D**), suggesting that this mutant likely displays a class I

phenotype. Thus, we conclude that the class II mutant viruses are rarely present in the latently

infected cells and therefore unlikely to contribute to chronic immune activation.

DISCUSSION

Class II IN mutations impair virion particle maturation by blocking IN-gRNA binding in virions and

those within the CTD, including R269A/K273A and R262A/R263A, impede IN-gRNA binding

without affecting functional oligomerization of IN [15, 38]. During serial passaging experiments to

identify compensatory mutations of the R269A/K273A class II IN mutant, viruses acquired D256N

and D270N IN mutations sequentially. We initially anticipated that mutations outside of IN, such

as CA and NC, could also arise, given that the IN R269A/K273A mutant is still catalytically active

and a compensatory mutation in CA or NC could presumably allow the proper packaging of the

gRNA within virions. On the other hand, we did not observe any such substitutions showcasing

the distinct role of IN:gRNA binding in proper virion maturation. Notably, D256N and D270N

substitutions each arose through a single mutation (D256N: GAC→AAC, D270N:GAU→AAU)

and thus likely provided an easier pathway for suppression than reverting back to R269 and K273,

each of which would require two mutations. Though other mutations in IN could in principle restore

the ability of IN to bind RNA, rise of such mutations was likely constrained in part by the necessity

to maintain a catalytically active IN.

IN CTD has a net positive charge of +3. While R269A/K273A class II mutations reduced the net

charge to +1, the introduction of D2N substitutions restored the overall charge back to a +3. This

observation suggested that the net electrostatic charge of the IN-CTD may be a key parameter

for IN-gRNA binding. Consistent with this hypothesis, the charge reversal by the D256R

substitution was sufficient to restore IN-gRNA binding for the R269A/K273A IN mutant. Though

these findings point to an electrostatic component of IN-RNA interactions, we cannot exclude the

possibility that the Asn and Arg residues also mediate H-bonding and van der Waals contacts

with distinct nucleobases in the cognate RNA molecules. Inspection of the available X-ray

structure [62] indicates that all IN residues implicated in RNA binding by the present study are

positioned either in the highly flexible C-terminal tail (aa 261-275) or the 252-257 loop (**Fig. 5H**).

The pliable nature of these CTD regions could be essential for allowing IN to optimally bind to

cognate gRNA.

These observations extended to another class II mutant, R262A/R263A, whereby the D256R and

to a lesser extent the D256K substitutions restored RNA binding. In contrast, the D2N

substitutions did not enhance RNA-binding or infectivity in this setting, demonstrating a degree of

context dependency. It is possible that the proximity of the D270N to R269A/K273A residues may

explain why it was more effective in restoring RNA binding for this class II mutant but not for IN

R262A/R263A. The role of Arg and Lys residues in RNA-binding proteins has been noted [67, 75-

78]. Interestingly, Arg residues are overall more heavily involved in interactions with all bases,

contributing with 16%–20% of all contacts while Lys residues only provide 3%–9%, which may

explain why the D256R substitution restored RNA binding to a greater extent [79]. Altogether, our

findings suggest that IN-gRNA binding is mediated in part by electrostatic interactions between

the basic residues in IN CTD and the negatively charged RNA backbone.

Though the electrostatic component of IN-RNA interactions imply a level of non-specificity, IN

bound to distinct locations on the gRNA and displays high binding affinity to structured elements,

such as the HIV-1 TAR element [38]. Thus, it is likely that IN-gRNA interactions are mediated by

both the non-specific interactions of the basic residues with the RNA phosphate backbone and

specific interactions with the cognate RNA. For example, recognition of the TAR loop by Tat and

the super elongation complex is based on a complex set of interactions that results primarily in

the readout of the structure as opposed to the sequence by SEC, and additional interaction of the

Tat arginine rich motif (ARM) with the TAR bulge and the major groove through electrostatic

interactions with the RNA phosphates, H-bonding with specific bases, as well as multiple van der

Waals contacts of Cyclin T with the flipped-out G32 TAR base to strengthen the TAR complex

[80, 81]. Structural studies of IN in complex with cognate RNA molecules are needed to tease

apart the specificity determinants for IN-RNA interactions.

RNA binding proteins commonly encode modular RNA-binding domains (i.e. RRMs and KH

domains), which form specific contacts with short degenerate sequences [82-84]. Utilization of

multiple RRMs/KH domains is thought to create a much larger binding interface, which allows

recognition of longer sequences on and enhanced affinity and specificity for target RNAs [82, 83,

85]. IN tetramerization is required for RNA binding in vitro [15] and may serve a similar purpose

through generation of a larger RNA-binding surface possibly for recognition of shape, RNA-

backbone and base-specific interactions as discussed above.

IN-gRNA binding is essential for HIV-1 virion morphological maturation and infectivity, thus an

excellent target for novel antiretroviral compounds. Clinically used HIV-1 IN inhibitors target the

catalytic activity of IN by blocking the strand transfer step of integration [2-7]. Despite high barriers

to resistance with the second-generation INSTIs, treatment continues to select for drug resistant

HIV-1 variants [86-91]. ALLINI-mediated inhibition of the non-catalytic function of IN can

complement existing INSTI-based therapies and increase the barrier to drug resistance

substantially. Although some ALLINIs displayed toxicity in animals, a highly potent and safe

pyrrolopyridine-based ALLINI, STP0404, has advanced to a human trial [65].

ALLINIs potently disrupt virion maturation indirectly through inducing aberrant IN multimerization

resulting in inhibition of IN-gRNA binding [12, 38]. Recent findings have shown that the ALLINI,

GSK-1264, is buried between the CTD of one IN dimer and the CCD of another dimer resulting in

the formation of open, inactive IN polymers [62]. Mutation of Y226, W235, K264 and K266

residues within the CTD prevents ALLINI-induced aberrant multimerization of IN [62, 71, 72].

However, the role of these residues in ALLINI-mediated inhibition cannot be easily assessed in

relevant infection settings given that their mutation alters the catalytic activity as well as RNA

binding properties of IN [62, 92, 93]. Our findings fill this gap and provide key genetic evidence

that ALLINI mechanism of action indeed involves the CTD. In particular, introducing the same

D256R change in the background of different class II substitutions similarly restored their

infectivity and thus enabled us to compare effects of R262/R263 vs R269A/K273A IN substitutions

on ALLINI activities. While HIV-1$_{IN(R262A/R263A/D256R)}$ and WT viruses were similarly susceptible to

ALLINIs (**Fig. 7A,B**), HIV-1$_{IN(R269A/K273A/D256R)}$ was substantially more sensitive to these inhibitors

(**Fig. 7D,E**). The X-ray structure of GSK-1264 induced IN polymers [62] reveals that R262, R263,

K273A as well as D256 are distanced from the inhibitor binding pocket, whereas the side chain

of R269 points toward the inhibitor bound at the CTD-CCD dimer interface (**Fig. 7G**).

The guanidine group of R269 engages with Q168 of the CCD from another IN subunit and is

positioned within 3 Å of GSK-1264. Yet, there is no interaction between R269 and the inhibitor as

the positively charged guanidine group faces a hydrophobic part of the benzodihydropyran moiety

of GSK-1264 (**Fig. 7G**). Conversely, the R269A substitution could provide more favorable

hydrophobic environment for the CTD-ALLINI-CCD interactions that may explain its increased

sensitivity to ALLINIs.

Overall, our studies reveal that electrostatic interactions play an important role in mediating IN-

gRNA interactions and demonstrate that CTD is a key determinant of ALLINI sensitivity. Structural

characterization of IN-gRNA complexes will be crucial to determine the precise mechanism of IN-

gRNA interactions. Such studies will also help better understand the ALLINI mechanism of action

and aid in the development of therapeutics that directly target IN-gRNA interactions.

MATERIALS AND METHODS

Plasmids

IN mutations were introduced into the HIV-1$_{NL4-3}$ full-length proviral plasmid (pNL4-3) by overlap

extension PCR. Forward and reverse primers containing IN mutations were used in PCR

reactions with anti-sense (with EcoRI restriction endonuclease site) and sense (with AgeI

restriction site) outer primers. The resulting fragments containing the desired mutations were

mixed at a 1:1 ratio and overlapped subsequently using the sense and antisense primer pairs.

The overlap fragments were digested with Age-I and EcoR-I before cloning into pNL4-3 plasmids.

The pLR2P-vprIN plasmid expressing a Vpr-IN fusion protein has been previously described [94].

IN mutations were introduced in the pLR2P-VprIN plasmid using the QuikChange Site-Directed

Mutagenesis kit (Agilent Technologies). The presence of the desired mutations and the absence

of unwanted secondary changes was assessed by Sanger sequencing.

Cell lines

All cell lines were obtained from the American Type Culture Collection and NIH AIDS Reagents

where STR profiling was performed. MT-4 cells were further STR profiled at Washington

University School of Medicine Genome Engineering and iPSC center. The cell lines are

consistently inspected for mycoplasma contamination using MycoAlert mycoplasma detection kit

(Lonza) and checked for being free of any other contaminations. HEK293T cells (ATCC CRL-

11268) and Hela-derived TZM-bl cells (NIH AIDS Reagent Program) were cultured in Dulbecco's

modified Eagle's medium supplemented with 10% fetal bovine serum. MT-4 cells were cultured

in in RPMI 1640 medium supplemented with 10% fetal bovine serum.

Analysis of compensatory mutations

Compensatory mutations of the R269A/K273A class-II IN mutation were isolated by serial

passaging. In this experiment, MT4 T-cells that express GFP under the control of HIV-1 LTR

(MT4-LTR-GFP), were infected with HIV-1pNL4-3 carrying the R269A/K273A class II IN mutation.

One million cells were infected at an MOI of 2 by WT cells, input of mutant virus was normalized

relative to particle numbers using a reverse transcriptase (RT) activity assay [95]. Infections were

monitored by FACS. At the end of each passage, virions were collected and wielded to infect cells

in the next passage while normalizing virus input before each infection. Aliquots of infected and

viral particles in the cell culture supernatants were collected over the duration of each passage.

Virions collected over the three passages were concentrated and gRNA was isolated from virions

using Trizol per manufacturer's instructions. Extracted RNA was prepared for deep sequencing

usingthe Illumina® TruSeq® Stranded Total RNA library prep workflow kit omitting the rRNA

depletion step. Resulting libraries were sequenced by an Illumina HiSeq 2000 platform.

Immunoblotting

Viral and cell lysates were resuspended in sodium dodecyl sulfate (SDS) sample buffer,

separated by electrophoresis on Bolt 4-12% Bis-Tris Plus gels (Life Technologies) and transferred

to nitrocellulose membranes. The membranes were then probed overnight at 4°C with a mouse

monoclonal anti-HIV p24 antibody (183-H12-5C, NIH AIDS reagents) or a mouse monoclonal

anti-HIV integrase antibody [96] in Odyssey Blocking Buffer (LI-COR). Membranes were probed

with fluorophore-conjugated secondary antibodies (LI-COR) and scanned using an LI-COR

Odyssey system. IN and CA levels in virions were quantified using Image Studio software (LI-
COR).

Vpr-IN trans-complementation experiments

Viruses bearing the class I IN mutation D116N were trans-complemented with class II mutant
proteins as previously described [94]; two-hundred thousand HEK293T cells were co-transfected
with a derivative of pNL4-3-derived plasmid bearing the IN D116N mutation, VSV-G, and
derivatives of the pLR2P-VprIN plasmids bearing class II IN mutations (or the compensatory
mutations thereof) at a ratio of 6:1:3. Cell-free virions were collected from cell culture two days
post-transfection. MT-4 cells were infected by the virions and the integration proficiency of trans-
complemented class II IN mutants was measured by the yield of progeny virions in cell culture
supernatants over a 6-day period as described before [94]. Briefly, MT-4 cells were incubated
with virus inoculum in 96 V-bottom well plates for 4hr at 37°C before washing away the inoculum
and replacing it with fresh media. Right after the addition of fresh media and over the ensuing 6
days, the number of virions in culture supernatant was quantified by measuring RT activity using
a Q-PCR-based assay [95].

CLIP experiments

CLIP experiments were conducted as previously described [15, 69, 70, 97, 98]. In short, cells in
15 cm cell culture plates were transfected with 30 µg full-length proviral plasmid (pNL4-3) DNA
containing the WT sequence or indicated IN mutations. 4-thiouridine (4SU) was added to the cell
culture media for 16hr before virus harvest. Cell culture supernatants were filtered through 0.22
µm filters and pelleted by ultracentrifugation through a 20% sucrose cushion using a Beckman
SW32-Ti rotor at 28,000rpm for 1.5hr at 4°C. The virus pellets were resuspended in phosphate-
buffered saline (PBS) and UV-crosslinked. Following lysis in 1X RIPA buffer, IN-RNA complexes
were immunoprecipitated using a mouse monoclonal anti-IN antibody [96] . Bound RNA was end-

labeled with γ-^{32}P-ATP and T4 polynucleotide kinase. The isolated protein-RNA complexes were

separated by SDS-PAGE, transferred to nitrocellulose membranes, and exposed to

autoradiography films to visualize IN-RNA complexes. Lysates and immunoprecipitates were also

analyzed by immunoblotting using antibodies against IN.

IN multimerization in virions

In 10 cm dishes, HEK293T cells were transfected with 10 µg pNL4-3 plasmid DNA containing the

WT sequence or indicated pol mutations within IN coding sequence. Two days post-transfection,

cell-free virions in cell culture supernatants were pelleted through a 20% sucrose gradient using

a Beckman SW41-Ti rotor at 28,000 rpm for 1.5hr at 4°C. Pelleted virions were resuspended in

1xPBS and treated with a membrane-permeable crosslinker, EGS (ThermoFisher Scientific), at a

concentration of 1mM for 30 min at room temperature. Crosslinking was stopped by the addition

of SDS sample buffer. The cross-linked samples were then separated on 3-8% Tris-acetate gels

and analyzed by immunoblotting using a mouse monoclonal anti-IN antibody [96].

Virus production and transmission electron microscopy

HEK293T cells in a 15 cm plate were transfected with 30 µg full-length proviral plasmid (pNL4-3)

DNA containing the WT sequence or indicated pol mutations within IN coding sequence. Two

days post transfection, cell culture supernatants were filtered through 0.22 µm filters, and pelleted

by ultracentrifugation using a Beckman SW32-Ti rotor at 28,000 rpm for 1.5 hr at 4°C. Fixative

(2% paraformaldehyde/2.5% glutaraldehyde (Polysciences Inc., Warrington, PA) in 100 mM

sodium cacodylate buffer, pH 7.2) was gently added to resulting pellets, and samples were

incubated overnight at 4°C. Samples were washed in sodium cacodylate buffer and postfixed in

1% osmium tetroxide (Polysciences Inc.) for 1 hr. Samples were then rinsed extensively in dH$_2$0

prior to en bloc staining with 1% aqueous uranyl acetate (Ted Pella Inc., Redding, CA) for 1

hr. After several rinses in dH$_2$0, samples were dehydrated in a graded series of ethanol and

embedded in Eponate 12 resin (Ted Pella Inc.). Sections of 95 nm were cut with a Leica Ultracut

UCT ultramicrotome (Leica Microsystems Inc., Bannockburn, IL), stained with uranyl acetate and

lead citrate, and viewed on a JEOL 1200 EX transmission electron microscope (JEOL USA Inc.,

Peabody, MA) equipped with an AMT 8 megapixel digital camera and AMT Image Capture Engine

V602 software (Advanced Microscopy Techniques, Woburn, MA).

Size exclusion chromatography (SEC)

All of the indicated mutations were introduced into a plasmid backbone expressing His_6 tagged

pNL4-3-derived IN by QuikChange site directed mutagenesis kit (Agilent) [99]. His_6 tagged

recombinant pNL4-3 WT and mutant Ins were expressed in BL21 (DE3) *E. coli* cells followed by

nickel and heparin column purification as described previously [99, 100]. Recombinant WT and

mutant Ins were analyzed on Superdex 200 10/300 GL column (GE Healthcare) with running

buffer containing 20 mM HEPES (pH 7.5), 1 M NaCl, 10% glycerol and 5 mM BME at 0.3 mL/min

flow rate. The proteins were diluted to 10 μM with the running buffer and incubated for 1 h at 4°C

followed by centrifugation at 10,000g for 10 min. Multimeric form determination was based on the

standards including bovine thyroglobulin (670,000 Da), bovine gamma-globulin (158,000 Da),

chicken ovalbumin (44,000 Da), horse myoglobin (17,000 Da) and vitamin B12 (1,350 Da).

Retention volumes for different oligomeric forms of IN were as follows: tetramer ~12.5 mL, dimer

~14 mL, monomer ~15-16 mL.

Analysis of IN-RNA binding in vitro

To monitor IN-RNA interactions we utilized AlphaScreen-based assay [38], which allows to

monitor the ability of IN to bind and bridge between two TAR RNAs. Briefly, equal concentrations

(1 nM) of two synthetic TAR RNA oligonucleotides labeled either with biotin or DIG were mixed

and then streptavidin donor and anti-DIG acceptor beads at 0.02 mg/mL concentration were

supplied in a buffer containing 100 mM NaCl, 1 mM $MgCl_2$, 1 mM DTT, 1 mg/mL BSA, and 25

mM Tris (pH 7.4). After 2 hr incubation at 4°C, 320 nM IN was added to the reaction mixture and incubated further for 1.5 hr at 4°C. AlphaScreen signals were recorded with a PerkinElmer Life Sciences Enspire multimode plate reader.

Structural modeling of IN

Electrostatic potential maps of WT and mutant IN CTDs were created by Adaptive Poisson-Boltzmann Solver (APBS) program [101] with macromolecular electrostatic calculations performed in PyMOL. The published crystal structure [62] (PDB ID: 5HOT) was used as a template. The calculation results are displayed as an electrostatic potential molecular surface. The low, mid, and high range values are -5, 0, and 5, respectively.

Analysis of IN mutations from latently infected cells

Sequences identified from latently infected CD4+ T-cells [73] were downloaded from NCBI GenBank based on their accession numbers (KF526120-KF526339). Sequences were imported into the GLUE software framework [102, 103] and aligned. Multiple sequence alignments (MSAs) containing subtype B and subtype C sequences were constructed using MUSCLE, manually inspected in AliView [104] and imported into a GLUE project database. Within GLUE, MSAs were constrained to the pNL4-3 reference to establish a standardized coordinate space for the gene being analyzed. Amino acid frequencies at each alignment position were summarized using GLUE's amino-acid frequency calculation algorithm, which accounts for contingencies such as missing data and incomplete codons.

ACKNOWLEDGEMENTS

This work was supported by NIH grants AI150497 (SBK), AI1508470 (U54 Center for HIV RNA Studies, SBK, RG), R01 AI143649 (MK) and Milton Schlesinger Student Fellowship (CSM). We thank all members of the Kutluay lab for critical suggestions and feedback.

REFERENCES

1.   Engelman AN. Multifaceted HIV integrase functionalities and therapeutic strategies for their inhibition. The Journal of biological chemistry. 2019. doi: 10.1074/jbc.REV119.006901. PubMed PMID: 31467082.

2.   Hazuda DJ, Felock P, Witmer M, Wolfe A, Stillmock K, Grobler JA, et al. Inhibitors of strand transfer that prevent integration and inhibit HIV-1 replication in cells. Science. 2000;287(5453):646-50. doi: 10.1126/science.287.5453.646. PubMed PMID: 10649997.

3.   Summa V, Petrocchi A, Bonelli F, Crescenzi B, Donghi M, Ferrara M, et al. Discovery of raltegravir, a potent, selective orally bioavailable HIV-integrase inhibitor for the treatment of HIV-AIDS infection. J Med Chem. 2008;51(18):5843-55. doi: 10.1021/jm800245z. PubMed PMID: 18763751.

4.   Ramanathan S, Mathias AA, German P, Kearney BP. Clinical pharmacokinetic and pharmacodynamic profile of the HIV integrase inhibitor elvitegravir. Clin Pharmacokinet. 2011;50(4):229-44. doi: 10.2165/11584570-000000000-00000. PubMed PMID: 21348537.

5.   Min S, Song I, Borland J, Chen S, Lou Y, Fujiwara T, et al. Pharmacokinetics and safety of S/GSK1349572, a next-generation HIV integrase inhibitor, in healthy volunteers. Antimicrobial agents and chemotherapy. 2010;54(1):254-8. doi: 10.1128/AAC.00842-09. PubMed PMID: 19884365; PubMed Central PMCID: PMCPMC2798521.

6.   Tsiang M, Jones GS, Goldsmith J, Mulato A, Hansen D, Kan E, et al. Antiviral Activity of Bictegravir (GS-9883), a Novel Potent HIV-1 Integrase Strand Transfer Inhibitor with an Improved Resistance Profile. Antimicrobial agents and chemotherapy. 2016;60(12):7086-97. doi: 10.1128/AAC.01474-16. PubMed PMID: 27645238; PubMed Central PMCID: PMCPMC5118987.

7.   Smith SJ, Zhao XZ, Burke TR, Jr., Hughes SH. Efficacies of Cabotegravir and Bictegravir against drug-resistant HIV-1 integrase mutants. Retrovirology. 2018;15(1):37. doi: 10.1186/s12977-018-0420-7. PubMed PMID: 29769116; PubMed Central PMCID: PMCPMC5956922.

8.   Rockstroh JK, DeJesus E, Lennox JL, Yazdanpanah Y, Saag MS, Wan H, et al. Durable efficacy and safety of raltegravir versus efavirenz when combined with tenofovir/emtricitabine in treatment-naive HIV-1-infected patients: final 5-year results from STARTMRK. J Acquir Immune Defic Syndr. 2013;63(1):77-85. doi: 10.1097/QAI.0b013e31828ace69. PubMed PMID: 23412015.

9.   Taha H, Das A, Das S. Clinical effectiveness of dolutegravir in the treatment of HIV/AIDS. Infect Drug Resist. 2015;8:339-52. Epub 2015/10/01. doi: 10.2147/IDR.S68396. PubMed PMID: 26491363; PubMed Central PMCID: PMCPMC4598225.

10.   Smith SJ, Zhao XZ, Passos DO, Lyumkis D, Burke TR, Hughes SH. HIV-1 Integrase Inhibitors that are active against Drug-Resistant Integrase Mutants. Antimicrob Agents Chemother. 2020. Epub 2020/06/29. doi: 10.1128/AAC.00611-20. PubMed PMID: 32601157.

11.   Brooks KM, Sherman EM, Egelund EF, Brotherton A, Durham S, Badowski ME, et al. Integrase Inhibitors: After 10 Years of Experience, Is the Best Yet to Come? Pharmacotherapy. 2019;39(5):576-98. doi: 10.1002/phar.2246. PubMed PMID: 30860610.

12.   Jurado KA, Wang H, Slaughter A, Feng L, Kessl JJ, Koh Y, et al. Allosteric integrase inhibitor potency is determined through the inhibition of HIV-1 particle maturation. Proceedings of the National Academy of Sciences of the United States of America. 2013;110(21):8690-5. doi: 10.1073/pnas.1300703110. PubMed PMID: 23610442; PubMed Central PMCID: PMC3666754.

13.   Lu R, Limon A, Devroe E, Silver PA, Cherepanov P, Engelman A. Class II integrase mutants with changes in putative nuclear localization signals are primarily blocked at a postnuclear entry step of human immunodeficiency virus type 1 replication. J Virol. 2004;78(23):12735-46. doi: 10.1128/JVI.78.23.12735-12746.2004. PubMed PMID: 15542626; PubMed Central PMCID: PMCPMC525011.

14.   Engelman A. In vivo analysis of retroviral integrase structure and function. Adv Virus
Res. 1999;52:411-26. Epub 1999/06/29. PubMed PMID: 10384245.
15.   Elliott JL, Eschbach JE, Koneru PC, Li W, Puray-Chavez M, Townsend D, et al.
Integrase-RNA interactions underscore the critical role of integrase in HIV-1 virion
morphogenesis. Elife. 2020;9. Epub 2020/09/23. doi: 10.7554/eLife.54311. PubMed PMID:
32960169; PubMed Central PMCID: PMCPMC7671690.
16.   Engelman A, Englund G, Orenstein JM, Martin MA, Craigie R. Multiple effects of
mutations in human immunodeficiency virus type 1 integrase on viral replication. J Virol.
1995;69(5):2729-36. Epub 1995/05/01. PubMed PMID: 7535863; PubMed Central PMCID:
PMC188965.
17.   Engelman A, Cherepanov P. Retroviral Integrase Structure and DNA Recombination
Mechanism. Microbiol Spectr. 2014;2(6). doi: 10.1128/microbiolspec.MDNA3-0024-2014.
PubMed PMID: 26104441.
18.   Engelman AN, Cherepanov P. Retroviral intasomes arising. Curr Opin Struct Biol.
2017;47:23-9. doi: 10.1016/j.sbi.2017.04.005. PubMed PMID: 28458055; PubMed Central
PMCID: PMCPMC5660667.
19.   Craigie R, Bushman FD. HIV DNA integration. Cold Spring Harb Perspect Med.
2012;2(7):a006890. Epub 2012/07/05. doi: 10.1101/cshperspect.a006890. PubMed PMID:
22762018; PubMed Central PMCID: PMCPMC3385939.
20.   Engelman A, Cherepanov P. The structural biology of HIV-1: mechanistic and
therapeutic insights. Nat Rev Microbiol. 2012;10(4):279-90. doi: 10.1038/nrmicro2747. PubMed
PMID: 22421880; PubMed Central PMCID: PMC3588166.
21.   Ciuffi A, Llano M, Poeschla E, Hoffmann C, Leipzig J, Shinn P, et al. A role for
LEDGF/p75 in targeting HIV DNA integration. Nat Med. 2005;11(12):1287-9. doi:
10.1038/nm1329. PubMed PMID: 16311605.
22.   Shun MC, Raghavendra NK, Vandegraaff N, Daigle JE, Hughes S, Kellam P, et al.
LEDGF/p75 functions downstream from preintegration complex formation to effect gene-specific
HIV-1 integration. Genes Dev. 2007;21(14):1767-78. doi: 10.1101/gad.1565107. PubMed PMID:
17639082; PubMed Central PMCID: PMC1920171.
23.   Schroder AR, Shinn P, Chen H, Berry C, Ecker JR, Bushman F. HIV-1 integration in the
human genome favors active genes and local hotspots. Cell. 2002;110(4):521-9. Epub
2002/08/31. PubMed PMID: 12202041.
24.   Christ F, Debyser Z. The LEDGF/p75 integrase interaction, a novel target for anti-HIV
therapy. Virology. 2013;435(1):102-9. Epub 2012/12/12. doi: 10.1016/j.virol.2012.09.033.
PubMed PMID: 23217620.
25.   Ansari-Lari MA, Donehower LA, Gibbs RA. Analysis of human immunodeficiency virus
type 1 integrase mutants. Virology. 1995;213(2):680. PubMed PMID: 7491794.
26.   Bukovsky A, Gottlinger H. Lack of integrase can markedly affect human
immunodeficiency virus type 1 particle production in the presence of an active viral protease. J
Virol. 1996;70(10):6820-5. Epub 1996/10/01. PubMed PMID: 8794322; PubMed Central
PMCID: PMC190728.
27.   Jenkins TM, Engelman A, Ghirlando R, Craigie R. A soluble active mutant of HIV-1
integrase: involvement of both the core and carboxyl-terminal domains in multimerization. The
Journal of biological chemistry. 1996;271(13):7712-8. PubMed PMID: 8631811.
28.   Kalpana GV, Reicin A, Cheng GS, Sorin M, Paik S, Goff SP. Isolation and
characterization of an oligomerization-negative mutant of HIV-1 integrase. Virology.
1999;259(2):274-85. doi: 10.1006/viro.1999.9767. PubMed PMID: 10388652.
29.   Leavitt AD, Robles G, Alesandro N, Varmus HE. Human immunodeficiency virus type 1
integrase mutants retain in vitro integrase activity yet fail to integrate viral DNA efficiently during
infection. J Virol. 1996;70(2):721-8. PubMed PMID: 8551608; PubMed Central PMCID:
PMCPMC189872.

30.   Liao WH, Wang CT. Characterization of human immunodeficiency virus type 1 Pr160
gag-pol mutants with truncations downstream of the protease domain. Virology.
2004;329(1):180-8. doi: 10.1016/j.virol.2004.08.010. PubMed PMID: 15476885.
31.   Lu R, Ghory HZ, Engelman A. Genetic analyses of conserved residues in the carboxyl-
terminal domain of human immunodeficiency virus type 1 integrase. J Virol. 2005;79(16):10356-
68. doi: 10.1128/JVI.79.16.10356-10368.2005. PubMed PMID: 16051828; PubMed Central
PMCID: PMCPMC1182625.
32.   Nakamura T, Masuda T, Goto T, Sano K, Nakai M, Harada S. Lack of infectivity of HIV-1
integrase zinc finger-like domain mutant with morphologically normal maturation. Biochem
Biophys Res Commun. 1997;239(3):715-22. doi: 10.1006/bbrc.1997.7541. PubMed PMID:
9367835.
33.   Quillent C, Borman AM, Paulous S, Dauguet C, Clavel F. Extensive regions of pol are
required for efficient human immunodeficiency virus polyprotein processing and particle
maturation. Virology. 1996;219(1):29-36. doi: 10.1006/viro.1996.0219. PubMed PMID: 8623542.
34.   Shin CG, Taddeo B, Haseltine WA, Farnet CM. Genetic analysis of the human
immunodeficiency virus type 1 integrase protein. J Virol. 1994;68(3):1633-42. PubMed PMID:
8107224; PubMed Central PMCID: PMCPMC236621.
35.   Taddeo B, Haseltine WA, Farnet CM. Integrase mutants of human immunodeficiency
virus type 1 with a specific defect in integration. J Virol. 1994;68(12):8401-5. PubMed PMID:
7966634; PubMed Central PMCID: PMCPMC237311.
36.   Wu X, Liu H, Xiao H, Conway JA, Hehl E, Kalpana GV, et al. Human immunodeficiency
virus type 1 integrase protein promotes reverse transcription through specific interactions with
the nucleoprotein reverse transcription complex. J Virol. 1999;73(3):2126-35. PubMed PMID:
9971795; PubMed Central PMCID: PMCPMC104457.
37.   Fontana J, Jurado KA, Cheng N, Ly NL, Fuchs JR, Gorelick RJ, et al. Distribution and
Redistribution of HIV-1 Nucleocapsid Protein in Immature, Mature, and Integrase-Inhibited
Virions: a Role for Integrase in Maturation. J Virol. 2015;89(19):9765-80. doi:
10.1128/JVI.01522-15. PubMed PMID: 26178982; PubMed Central PMCID: PMCPMC4577894.
38.   Kessl JJ, Kutluay SB, Townsend D, Rebensburg S, Slaughter A, Larue RC, et al. HIV-1
Integrase Binds the Viral RNA Genome and Is Essential during Virion Morphogenesis. Cell.
2016;166(5):1257-68 e12. doi: 10.1016/j.cell.2016.07.044. PubMed PMID: 27565348; PubMed
Central PMCID: PMCPMC5003418.
39.   Ao Z, Fowke KR, Cohen EA, Yao X. Contribution of the C-terminal tri-lysine regions of
human immunodeficiency virus type 1 integrase for efficient reverse transcription and viral DNA
nuclear import. Retrovirology. 2005;2:62. doi: 10.1186/1742-4690-2-62. PubMed PMID:
16232319; PubMed Central PMCID: PMCPMC1277849.
40.   Busschots K, Voet A, De Maeyer M, Rain JC, Emiliani S, Benarous R, et al. Identification
of the LEDGF/p75 binding site in HIV-1 integrase. Journal of molecular biology.
2007;365(5):1480-92. doi: 10.1016/j.jmb.2006.10.094. PubMed PMID: 17137594.
41.   Engelman A, Liu Y, Chen H, Farzan M, Dyda F. Structure-based mutagenesis of the
catalytic domain of human immunodeficiency virus type 1 integrase. J Virol. 1997;71(5):3507-
14. PubMed PMID: 9094622; PubMed Central PMCID: PMCPMC191497.
42.   Limon A, Devroe E, Lu R, Ghory HZ, Silver PA, Engelman A. Nuclear localization of
human immunodeficiency virus type 1 preintegration complexes (PICs): V165A and R166A are
pleiotropic integrase mutants primarily defective for integration, not PIC nuclear import. J Virol.
2002;76(21):10598-607. doi: 10.1128/jvi.76.21.10598-10607.2002. PubMed PMID: 12368302;
PubMed Central PMCID: PMCPMC136649.
43.   Lloyd AG, Ng YS, Muesing MA, Simon V, Mulder LC. Characterization of HIV-1
integrase N-terminal mutant viruses. Virology. 2007;360(1):129-35. doi:
10.1016/j.virol.2006.10.007. PubMed PMID: 17109911; PubMed Central PMCID:
PMCPMC2094525.

44.   Lu R, Vandegraaff N, Cherepanov P, Engelman A. Lys-34, dispensable for integrase
catalysis, is required for preintegration complex function and human immunodeficiency virus
type 1 replication. J Virol. 2005;79(19):12584-91. doi: 10.1128/JVI.79.19.12584-12591.2005.
PubMed PMID: 16160186; PubMed Central PMCID: PMCPMC1211547.
45.   Masuda T, Planelles V, Krogstad P, Chen IS. Genetic analysis of human
immunodeficiency virus type 1 integrase and the U3 att site: unusual phenotype of mutants in
the zinc finger-like domain. J Virol. 1995;69(11):6687-96. PubMed PMID: 7474078; PubMed
Central PMCID: PMCPMC189578.
46.   Rahman S, Lu R, Vandegraaff N, Cherepanov P, Engelman A. Structure-based
mutagenesis of the integrase-LEDGF/p75 interface uncouples a strict correlation between in
vitro protein binding and HIV-1 fitness. Virology. 2007;357(1):79-90. doi:
10.1016/j.virol.2006.08.011. PubMed PMID: 16959283.
47.   Riviere L, Darlix JL, Cimarelli A. Analysis of the viral elements required in the nuclear
import of HIV-1 DNA. J Virol. 2010;84(2):729-39. doi: 10.1128/JVI.01952-09. PubMed PMID:
19889772; PubMed Central PMCID: PMCPMC2798343.
48.   Tsurutani N, Kubo M, Maeda Y, Ohashi T, Yamamoto N, Kannagi M, et al. Identification
of critical amino acid residues in human immunodeficiency virus type 1 IN required for efficient
proviral DNA formation at steps prior to integration in dividing and nondividing cells. J Virol.
2000;74(10):4795-806. doi: 10.1128/jvi.74.10.4795-4806.2000. PubMed PMID: 10775618;
PubMed Central PMCID: PMCPMC112002.
49.   Wiskerchen M, Muesing MA. Human immunodeficiency virus type 1 integrase: effects of
mutations on viral ability to integrate, direct viral gene expression from unintegrated viral DNA
templates, and sustain viral propagation in primary cells. J Virol. 1995;69(1):376-86. PubMed
PMID: 7983732; PubMed Central PMCID: PMCPMC188585.
50.   Zhu K, Dobard C, Chow SA. Requirement for integrase during reverse transcription of
human immunodeficiency virus type 1 and the effect of cysteine mutations of integrase on its
interactions with reverse transcriptase. J Virol. 2004;78(10):5045-55. doi:
10.1128/jvi.78.10.5045-5055.2004. PubMed PMID: 15113886; PubMed Central PMCID:
PMCPMC400327.
51.   De Houwer S, Demeulemeester J, Thys W, Rocha S, Dirix L, Gijsbers R, et al. The HIV-
1 integrase mutant R263A/K264A is 2-fold defective for TRN-SR2 binding and viral nuclear
import. The Journal of biological chemistry. 2014;289(36):25351-61. doi:
10.1074/jbc.M113.533281. PubMed PMID: 25063804; PubMed Central PMCID:
PMCPMC4155696.
52.   Johnson BC, Metifiot M, Ferris A, Pommier Y, Hughes SH. A homology model of HIV-1
integrase and analysis of mutations designed to test the model. Journal of molecular biology.
2013;425(12):2133-46. doi: 10.1016/j.jmb.2013.03.027. PubMed PMID: 23542006.
53.   Mohammed KD, Topper MB, Muesing MA. Sequential deletion of the integrase (Gag-
Pol) carboxyl terminus reveals distinct phenotypic classes of defective HIV-1. J Virol.
2011;85(10):4654-66. doi: 10.1128/JVI.02374-10. PubMed PMID: 21367893; PubMed Central
PMCID: PMC3126176.
54.   Shehu-Xhilaga M, Hill M, Marshall JA, Kappes J, Crowe SM, Mak J. The conformation of
the mature dimeric human immunodeficiency virus type 1 RNA genome requires packaging of
pol protein. J Virol. 2002;76(9):4331-40. doi: 10.1128/jvi.76.9.4331-4340.2002. PubMed PMID:
11932399; PubMed Central PMCID: PMCPMC155102.
55.   Engelman A, Craigie R. Identification of conserved amino acid residues critical for
human immunodeficiency virus type 1 integrase function in vitro. J Virol. 1992;66(11):6361-9.
PubMed PMID: 1404595; PubMed Central PMCID: PMCPMC240128.
56.   Lutzke RA, Plasterk RH. Structure-based mutational analysis of the C-terminal DNA-
binding domain of human immunodeficiency virus type 1 integrase: critical residues for protein

oligomerization and DNA binding. J Virol. 1998;72(6):4841-8. PubMed PMID: 9573250; PubMed Central PMCID: PMCPMC110031.

57.   Lutzke RA, Vink C, Plasterk RH. Characterization of the minimal DNA-binding domain of the HIV integrase protein. Nucleic Acids Res. 1994;22(20):4125-31. doi: 10.1093/nar/22.20.4125. PubMed PMID: 7937137; PubMed Central PMCID: PMCPMC331899.

58.   Balakrishnan M, Yant SR, Tsai L, O'Sullivan C, Bam RA, Tsai A, et al. Non-catalytic site HIV-1 integrase inhibitors disrupt core maturation and induce a reverse transcription block in target cells. PloS one. 2013;8(9):e74163. Epub 2013/09/17. doi: 10.1371/journal.pone.0074163 PONE-D-13-26050 [pii]. PubMed PMID: 24040198; PubMed Central PMCID: PMC3767657.

59.   Desimmie BA, Schrijvers R, Demeulemeester J, Borrenberghs D, Weydert C, Thys W, et al. LEDGINs inhibit late stage HIV-1 replication by modulating integrase multimerization in the virions. Retrovirology. 2013;10:57. Epub 2013/06/01. doi: 1742-4690-10-57 [pii] 10.1186/1742-4690-10-57. PubMed PMID: 23721378; PubMed Central PMCID: PMC3671127.

60.   Sharma A, Slaughter A, Jena N, Feng L, Kessl JJ, Fadel HJ, et al. A new class of multimerization selective inhibitors of HIV-1 integrase. PLoS pathogens. 2014;10(5):e1004171. doi: 10.1371/journal.ppat.1004171. PubMed PMID: 24874515; PubMed Central PMCID: PMC4038613.

61.   Gupta K, Brady T, Dyer BM, Malani N, Hwang Y, Male F, et al. Allosteric inhibition of human immunodeficiency virus integrase: late block during viral replication and abnormal multimerization involving specific protein domains. The Journal of biological chemistry. 2014;289(30):20477-88. doi: 10.1074/jbc.M114.551119. PubMed PMID: 24904063; PubMed Central PMCID: PMC4110260.

62.   Gupta K, Turkki V, Sherrill-Mix S, Hwang Y, Eilers G, Taylor L, et al. Structural Basis for Inhibitor-Induced Aggregation of HIV Integrase. PLoS Biol. 2016;14(12):e1002584. doi: 10.1371/journal.pbio.1002584. PubMed PMID: 27935939; PubMed Central PMCID: PMCPMC5147827 contained in this manuscript to improve therapeutics for HIV infection. The other authors declare that they have no conflict of interest.

63.   Christ F, Shaw S, Demeulemeester J, Desimmie BA, Marchand A, Butler S, et al. Small-molecule inhibitors of the LEDGF/p75 binding site of integrase block HIV replication and modulate integrase multimerization. Antimicrobial agents and chemotherapy. 2012;56(8):4365-74. doi: 10.1128/AAC.00717-12. PubMed PMID: 22664975; PubMed Central PMCID: PMC3421592.

64.   Christ F, Voet A, Marchand A, Nicolet S, Desimmie BA, Marchand D, et al. Rational design of small-molecule inhibitors of the LEDGF/p75-integrase interaction and HIV replication. Nature chemical biology. 2010;6(6):442-8. doi: 10.1038/nchembio.370. PubMed PMID: 20473303.

65.   Maehigashi T, Ahn S, Kim UI, Lindenberger J, Oo A, Koneru PC, et al. A highly potent and safe pyrrolopyridine-based allosteric HIV-1 integrase inhibitor targeting host LEDGF/p75-integrase interaction site. PLoS pathogens. 2021;17(7):e1009671. Epub 2021/07/23. doi: 10.1371/journal.ppat.1009671. PubMed PMID: 34293041; PubMed Central PMCID: PMCPMC8297771 following competing interests: S.A., U.K., and K.K. are employees of ST Pharm, B.K. is consultant of ST Pharm, and A.N.E. consults for ViiV Healthcare, Co.. All other authors declare no competing interests.

66.   Serganov A, Patel DJ. Towards deciphering the principles underlying an mRNA recognition code. Curr Opin Struct Biol. 2008;18(1):120-9. Epub 2008/02/08. doi: 10.1016/j.sbi.2007.12.006. PubMed PMID: 18255277; PubMed Central PMCID: PMCPMC3744885.

67.   Bahadur RP, Zacharias M, Janin J. Dissecting protein-RNA recognition sites. Nucleic Acids Res. 2008;36(8):2705-16. Epub 2008/03/21. doi: 10.1093/nar/gkn102. PubMed PMID: 18353859; PubMed Central PMCID: PMCPMC2377425.

68.   Stefl R, Skrisovska L, Allain FH. RNA sequence- and shape-dependent recognition by
proteins in the ribonucleoprotein particle. EMBO Rep. 2005;6(1):33-8. Epub 2005/01/12. doi:
10.1038/sj.embor.7400325. PubMed PMID: 15643449; PubMed Central PMCID:
PMCPMC1299235.
69.   Shema Mugisha C, Tenneti K, Kutluay SB. Clip for studying protein-RNA interactions
that regulate virus replication. Methods. 2020;183:84-92. Epub 2019/11/26. doi:
10.1016/j.ymeth.2019.11.011. PubMed PMID: 31765715; PubMed Central PMCID:
PMCPMC6963001.
70.   Bieniasz PD, Kutluay SB. CLIP-related methodologies and their application to
retrovirology. Retrovirology. 2018;15(1):35. Epub 2018/05/03. doi: 10.1186/s12977-018-0417-2.
PubMed PMID: 29716635; PubMed Central PMCID: PMCPMC5930818.
71.   Deng N, Hoyte A, Mansour YE, Mohamed MS, Fuchs JR, Engelman AN, et al. Allosteric
HIV-1 integrase inhibitors promote aberrant protein multimerization by directly mediating inter-
subunit interactions: Structural and thermodynamic modeling studies. Protein Sci.
2016;25(11):1911-7. Epub 2016/10/26. doi: 10.1002/pro.2997. PubMed PMID: 27503276;
PubMed Central PMCID: PMCPMC5079246.
72.   Shkriabai N, Dharmarajan V, Slaughter A, Kessl JJ, Larue RC, Feng L, et al. A critical
role of the C-terminal segment for allosteric inhibitor-induced aberrant multimerization of HIV-1
integrase. J Biol Chem. 2014;289(38):26430-40. Epub 2014/08/15. doi:
10.1074/jbc.M114.589572. PubMed PMID: 25118283; PubMed Central PMCID:
PMCPMC4176244.
73.   Ho YC, Shan L, Hosmane NN, Wang J, Laskey SB, Rosenbloom DI, et al. Replication-
competent noninduced proviruses in the latent reservoir increase barrier to HIV-1 cure. Cell.
2013;155(3):540-51. Epub 2013/11/19. doi: 10.1016/j.cell.2013.09.020. PubMed PMID:
24243014; PubMed Central PMCID: PMCPMC3896327.
74.   Pollack RA, Jones RB, Pertea M, Bruner KM, Martin AR, Thomas AS, et al. Defective
HIV-1 Proviruses Are Expressed and Can Be Recognized by Cytotoxic T Lymphocytes, which
Shape the Proviral Landscape. Cell Host Microbe. 2017;21(4):494-506 e4. Epub 2017/04/14.
doi: 10.1016/j.chom.2017.03.008. PubMed PMID: 28407485; PubMed Central PMCID:
PMCPMC5433942.
75.   Bayer TS, Booth LN, Knudsen SM, Ellington AD. Arginine-rich motifs present multiple
interfaces for specific binding by RNA. RNA. 2005;11(12):1848-57. Epub 2005/11/30. doi:
10.1261/rna.2167605. PubMed PMID: 16314457; PubMed Central PMCID: PMCPMC1370873.
76.   Jones S, Daley DT, Luscombe NM, Berman HM, Thornton JM. Protein-RNA interactions:
a structural analysis. Nucleic Acids Res. 2001;29(4):943-54. Epub 2001/02/13. doi:
10.1093/nar/29.4.943. PubMed PMID: 11160927; PubMed Central PMCID: PMCPMC29619.
77.   Jeong E, Kim H, Lee SW, Han K. Discovering the interaction propensities of amino acids
and nucleotides from protein-RNA complexes. Mol Cells. 2003;16(2):161-7. Epub 2003/12/04.
PubMed PMID: 14651256.
78.   Kim OT, Yura K, Go N. Amino acid residue doublet propensity in the protein-RNA
interface and its application to RNA interface prediction. Nucleic Acids Res. 2006;34(22):6450-
60. Epub 2006/11/30. doi: 10.1093/nar/gkl819. PubMed PMID: 17130160; PubMed Central
PMCID: PMCPMC1761430.
79.   Kruger DM, Neubacher S, Grossmann TN. Protein-RNA interactions: structural
characteristics and hotspot amino acids. RNA. 2018;24(11):1457-65. Epub 2018/08/11. doi:
10.1261/rna.066464.118. PubMed PMID: 30093489; PubMed Central PMCID:
PMCPMC6191724.
80.   Schulze-Gahmen U, Hurley JH. Structural mechanism for HIV-1 TAR loop recognition by
Tat and the super elongation complex. Proceedings of the National Academy of Sciences of the
United States of America. 2018;115(51):12973-8. Epub 2018/12/06. doi:

10.1073/pnas.1806438115. PubMed PMID: 30514815; PubMed Central PMCID:
PMCPMC6305006.
81.   Schulze-Gahmen U, Echeverria I, Stjepanovic G, Bai Y, Lu H, Schneidman-Duhovny D,
et al. Insights into HIV-1 proviral transcription from integrative structure and dynamics of the
Tat:AFF4:P-TEFb:TAR complex. Elife. 2016;5. Epub 2016/10/21. doi: 10.7554/eLife.15910.
PubMed PMID: 27731797; PubMed Central PMCID: PMCPMC5072841.
82.   Han SP, Tang YH, Smith R. Functional diversity of the hnRNPs: past, present and
perspectives. Biochem J. 2010;430(3):379-92. doi: 10.1042/BJ20100396. PubMed PMID:
20795951.
83.   Jeong S. SR Proteins: Binders, Regulators, and Connectors of RNA. Mol Cells.
2017;40(1):1-9. doi: 10.14348/molcells.2017.2319. PubMed PMID: 28152302; PubMed Central
PMCID: PMCPMC5303883.
84.   Lunde BM, Moore C, Varani G. RNA-binding proteins: modular design for efficient
function. Nat Rev Mol Cell Biol. 2007;8(6):479-90. Epub 2007/05/03. doi: 10.1038/nrm2178.
PubMed PMID: 17473849; PubMed Central PMCID: PMCPMC5507177.
85.   Ban T, Zhu JK, Melcher K, Xu HE. Structural mechanisms of RNA recognition:
sequence-specific and non-specific RNA-binding proteins and the Cas9-RNA-DNA complex.
Cell Mol Life Sci. 2015;72(6):1045-58. Epub 2014/11/30. doi: 10.1007/s00018-014-1779-9.
PubMed PMID: 25432705.
86.   Wijting IEA, Lungu C, Rijnders BJA, van der Ende ME, Pham HT, Mesplede T, et al.
HIV-1 Resistance Dynamics in Patients With Virologic Failure to Dolutegravir Maintenance
Monotherapy. J Infect Dis. 2018;218(5):688-97. doi: 10.1093/infdis/jiy176. PubMed PMID:
29617822.
87.   Radzio-Basu J, Council O, Cong ME, Ruone S, Newton A, Wei X, et al. Drug resistance
emergence in macaques administered cabotegravir long-acting for pre-exposure prophylaxis
during acute SHIV infection. Nat Commun. 2019;10(1):2005. Epub 2019/05/01. doi:
10.1038/s41467-019-10047-w. PubMed PMID: 31043606; PubMed Central PMCID:
PMCPMC6494879.
88.   Anstett K, Brenner B, Mesplede T, Wainberg MA. HIV drug resistance against strand
transfer integrase inhibitors. Retrovirology. 2017;14(1):36. doi: 10.1186/s12977-017-0360-7.
PubMed PMID: 28583191; PubMed Central PMCID: PMCPMC5460515.
89.   Rossouw TM, Hitchcock S, Botes M. The end of the line? A case of drug resistance to
third-line antiretroviral therapy. South Afr J HIV Med. 2016;17(1):454. Epub 2016/05/31. doi:
10.4102/sajhivmed.v17i1.454. PubMed PMID: 29568612; PubMed Central PMCID:
PMCPMC5843205.
90.   Zhang WW, Cheung PK, Oliveira N, Robbins MA, Harrigan PR, Shahid A. Accumulation
of Multiple Mutations In Vivo Confers Cross-Resistance to New and Existing Integrase
Inhibitors. J Infect Dis. 2018;218(11):1773-6. doi: 10.1093/infdis/jiy428. PubMed PMID:
30010985.
91.   Ndashimye E, Li Y, Reyes PS, Avino M, Olabode AS, Kityo CM, et al. High-level
resistance to bictegravir and cabotegravir in subtype A- and D-infected HIV-1 patients failing
raltegravir with multiple resistance mutations. J Antimicrob Chemother. 2021. Epub 2021/08/29.
doi: 10.1093/jac/dkab276. PubMed PMID: 34453542.
92.   Kessl JJ, Sharma A, Kvaratskhelia M. Methods for the Analyses of Inhibitor-Induced
Aberrant Multimerization of HIV-1 Integrase. Methods Mol Biol. 2016;1354:149-64. doi:
10.1007/978-1-4939-3046-3_10. PubMed PMID: 26714710; PubMed Central PMCID:
PMCPMC4793731.
93.   Madison MK, Lawson DQ, Elliott J, Ozanturk AN, Koneru PC, Townsend D, et al.
Allosteric HIV-1 Integrase Inhibitors Lead to Premature Degradation of the Viral RNA Genome
and Integrase in Target Cells. J Virol. 2017;91(17). doi: 10.1128/JVI.00821-17. PubMed PMID:
28615207; PubMed Central PMCID: PMCPMC5553177.

94.   Liu H, Wu X, Xiao H, Conway JA, Kappes JC. Incorporation of functional human
immunodeficiency virus type 1 integrase into virions independent of the Gag-Pol precursor
protein. J Virol. 1997;71(10):7704-10. PubMed PMID: 9311854; PubMed Central PMCID:
PMCPMC192121.
95.   Pizzato M, Erlwein O, Bonsall D, Kaye S, Muir D, McClure MO. A one-step SYBR Green
I-based product-enhanced reverse transcriptase assay for the quantitation of retroviruses in cell
culture supernatants. J Virol Methods. 2009;156(1-2):1-7. doi: 10.1016/j.jviromet.2008.10.012.
PubMed PMID: 19022294.
96.   Bouyac-Bertoia M, Dvorin JD, Fouchier RA, Jenkins Y, Meyer BE, Wu LI, et al. HIV-1
infection requires a functional integrase NLS. Molecular cell. 2001;7(5):1025-35. PubMed PMID:
11389849.
97.   Kutluay SB, Bieniasz PD. Analysis of HIV-1 Gag-RNA Interactions in Cells and Virions
by CLIP-seq. Methods Mol Biol. 2016;1354:119-31. doi: 10.1007/978-1-4939-3046-3_8.
PubMed PMID: 26714708; PubMed Central PMCID: PMCPMC6548315.
98.   Kutluay SB, Zang T, Blanco-Melo D, Powell C, Jannain D, Errando M, et al. Global
changes in the RNA binding specificity of HIV-1 gag regulate virion genesis. Cell.
2014;159(5):1096-109. Epub 2014/11/25. doi: 10.1016/j.cell.2014.09.057. PubMed PMID:
25416948; PubMed Central PMCID: PMCPMC4247003.
99.   Kessl JJ, Jena N, Koh Y, Taskent-Sezgin H, Slaughter A, Feng L, et al. Multimode,
cooperative mechanism of action of allosteric HIV-1 integrase inhibitors. The Journal of
biological chemistry. 2012;287(20):16801-11. doi: 10.1074/jbc.M112.354373. PubMed PMID:
22437836; PubMed Central PMCID: PMC3351293.
100.   Cherepanov P. LEDGF/p75 interacts with divergent lentiviral integrases and modulates
their enzymatic activity in vitro. Nucleic Acids Res. 2007;35(1):113-24. doi: 10.1093/nar/gkl885.
PubMed PMID: 17158150; PubMed Central PMCID: PMCPMC1802576.
101.   Jurrus E, Engel D, Star K, Monson K, Brandi J, Felberg LE, et al. Improvements to the
APBS biomolecular solvation software suite. Protein Sci. 2018;27(1):112-28. Epub 2017/08/25.
doi: 10.1002/pro.3280. PubMed PMID: 28836357; PubMed Central PMCID: PMCPMC5734301.
102.   Singer JB, Thomson EC, Hughes J, Aranday-Cortes E, McLauchlan J, da Silva Filipe A,
et al. Interpreting Viral Deep Sequencing Data with GLUE. Viruses. 2019;11(4). Epub
2019/04/17. doi: 10.3390/v11040323. PubMed PMID: 30987147; PubMed Central PMCID:
PMCPMC6520954.
103.   Singer JB, Thomson EC, McLauchlan J, Hughes J, Gifford RJ. GLUE: a flexible software
system for virus sequence data. BMC Bioinformatics. 2018;19(1):532. Epub 2018/12/20. doi:
10.1186/s12859-018-2459-9. PubMed PMID: 30563445; PubMed Central PMCID:
PMCPMC6299651.
104.   Larsson A. AliView: a fast and lightweight alignment viewer and editor for large datasets.
Bioinformatics. 2014;30(22):3276-8. Epub 2014/08/07. doi: 10.1093/bioinformatics/btu531.
PubMed PMID: 25095880; PubMed Central PMCID: PMCPMC4221126.
916
917
918
919

    FIGURE LEGENDS:

Figure 1. D256N and D270N substitutions in HIV-1 IN suppresses the replication defect of

R269A/K273A class-II IN mutant virus. (A) MT4-LTR-GFP cells were infected with equal particle

numbers of either WT or R269A/K273A IN mutant HIV-1 (NL4-3). The R269A/K273A IN viruses

were serially passaged for three times until the emergence of compensatory mutations. The

graphs represent the percentage of GFP positive cells as assessed by FACS over three passages

at the indicated days post-infection (dpi). **(B)** HIV-1 genomic RNA was isolated from viruses

collected from cell culture supernatants over the three passages (i.e. P1, P2, P3) and at the

indicated days post-infection (i.e. D29, D40, etc.). Whole-genome deep sequencing revealed the

acquisition of D256N and D270N compensatory mutations. Heatmap shows the percentage of

mutations at the indicated passages and days post-infection (dpi). **(C)** HEK293T cells were

transfected with full-length pNL4-3 expression plasmids carrying the D256N, D270N and

D256N/D270N (D2N) IN mutations introduced on the WT IN and IN R269A/K273A backbones.

Cell lysates and virions were purified two days post transfection and analyzed by immunoblotting

for CA and IN. The image is representative of five independent experiments. **(D)** HEK293T cells

were transfected as in C and cell culture supernatants containing viruses were titered on TZM-bl

indicator cells. The titers are presented relative to WT (set to 1). The columns represent the

average of five independent experiments and the error bars represent SEM (****$p<0.0001$, by

one-way ANOVA with Dunnett's multiple comparison test).

Figure 2. D256N and D270N substitutions restore IN-gRNA binding and accurate virion

maturation for the R269A/K273A class II IN mutant virus. (A) Autoradiogram of IN-RNA

adducts immunoprecipitated from virions bearing the indicated substitutions in IN. Immunoblots

below show the amount of IP'ed IN or CA protein in lysates. Data shown are representative of at

least three independent experiments. **(B)** Examination of virion maturation in WT and mutant IN

viruses by thin section electron microscopy (TEM). The graph quantifies virion morphologies;

each column is the average of two independent experiments and error bars represent the SEM

(****p<0.0001, by repeated measures one-way ANOVA).

Figure 3. Restoring the net charge of IN-CTD restores RNA binding and infectivity for the

R269A/K273A class II mutant. (A) Schematic diagram of IN and sequence of CTD residues with

basic and acidic amino acids highlighted in blue and red, respectively. **(B, C)** HEK293T cells were

transfected with full-length proviral HIV-1$_{NL4-3}$ expression plasmids carrying *pol* mutations

encoding for the indicated IN substitutions. **(B)** Cell lysates and purified virions were harvested

two days post transfection and analyzed by immunoblotting for CA and IN. Representative image

of one of five independent experiments is shown. **(C)** Infectious titers of WT or IN mutant HIV-

1$_{NL4-3}$ viruses in cell culture supernatants were determined on TZM-bl indicator cells and

normalized relative to particle number based on RT activity. Titer values are expressed relative

to WT (set to 1). The columns represent the average of four independent experiments and the

error bars represent SEM (****p<0.0001, by one-way ANOVA with Dunnett's multiple comparison

test). **(D)** Autoradiogram of IN-RNA adducts IP'ed from WT or IN mutant HIV-1$_{NL4-3}$ virions. The

amount of immunoprecipitated IN protein was assessed by the immunoblot below. Results are

representative of four independent replicates. **(E)** A representative growth curve of HIV-1 $_{NL4-3 \ IN}$

$_{(D116N)}$ viruses that were trans-complemented with the indicated Vpr-IN mutant proteins in cell

culture. Y-axis indicates fold increase in virion yield over day 0 as measured by RT activity in

culture supernatants. Error bars show SEM from 3 independent many replicates.

Figure 4. D256R and D256K substitutions restore IN-RNA binding and infectivity for the

HIV-1$_{NL4-3 \ IN(R262A/R263A)}$ virus. (A-E) HEK293T cells were transfected with proviral HIV-1$_{NL4-3}$

expression plasmids carrying *pol* mutations for the indicated IN substitutions. **(A, C)** Cell lysates

and virions were purified two days post transfection and analyzed by immunoblotting for CA and

IN. **(B, D)** WT or IN mutant HIV-1$_{NL4-3}$ viruses in cell culture supernatants were titered on TZM-bl

indicator cells. The titer values are presented relative to WT (set to 1). The columns represent the

average of two-three independent experiments, and the error bars represent SEM (****p<0.0001,

by one-way ANOVA with Dunnett's multiple comparisons). **(E)** Autoradiogram of IN-RNA adducts

immunoprecipitated from WT or IN mutant HIV-1$_{NL4-3}$ virions. The amount of immunoprecipitated

IN was assessed by the immunoblot shown below. Immunoblots and CLIP autoradiographs are

representative of four independent replicates. **(F)** A representative growth curve of HIV-1 $_{NL4-3\ IN}$

$_{(D116N)}$ viruses that were trans-complemented with the indicated IN mutant proteins in cell culture.

Y-axis indicates fold increase in virion yield over day 0 as measured by RT activity in culture

supernatants. Error bars show SEM from 3 independent many replicates.

Figure 5. Electrostatic potential maps of HIV-1 IN bearing class II and compensatory

mutations. (A-G) Electrostatic potential maps of the indicated HIV-1 IN mutants derived from the

crystal structure of Gupta et al. [62] is depicted. Calculation results are displayed as an

electrostatic potential molecular surface. The low, mid, and high range values are -5, 0, and 5,

respectively. (**H**) A cartoon view of the CTD structure [62] is shown with the C-terminal tail (aa

261-275) and the loop (aa 252-257) colored in cyan and orange, respectively. Side chains of

indicated residues are shown.

Figure 6. Assessing multimerization properties of IN in mutant viruses. (A) Purified WT or

IN mutant HIV-1$_{NL4-3}$ virions were treated with 1 mM EGS, and virus lysates analyzed by

immunoblotting using antibodies against IN following separation on 6% Tris-acetate gels. The

position of monomers (M), dimers (D), and tetramers (T) are indicated by arrows in a

representative western blot. **(B,C)** Quantification of IN multimerization in virions from experiments

conducted as in A. Error bars show the SEM from three independent experiments. **(D-I)**

Biochemical analysis of IN mutants multimerization and their interactions with HIV-1 TAR RNA.

(D-H) Representative SEC traces for indicated recombinant IN proteins. The X-axis indicates

elution volume (mL) and Y-axis indicates the intensity of absorbance (mAU). Tetramers (T),

dimers (D), and monomers (M) are indicated. **(I)** Summary of mutant INs bridging TAR RNA

compared to WT IN. Alpha screen counts at 320 nM for each protein is shown. The graphs show

average values of three independent experiments and the error bars indicate standard deviation.

Figure 7. Secondary site suppressors of the IN R269A/K273A mutant increase

susceptibility to ALLINIs. (A, B, D, E) Titers of viruses bearing the indicated substitutions in IN

produced from HEK293T cells at different concentrations of ALLINIs, BI-B2 or BI-D. WT or IN

mutant HIV-1$_{NL4-3}$ in cell culture supernatants were titered on TZM-bl indicator cells. The titer

values are represented relative to the mock control of each mutant (set to 1). Data are from two

independent biological replicates. **(C, F)** Autoradiogram of IN-RNA adducts immunoprecipitated

from WT or IN mutant HIV-1$_{NL4-3}$ virions produced from HEK293T cells in the presence of 0.1μM

of BI-D. The amount of immunoprecipitated IN protein was visualized by the immunoblot shown

below. Immunoblots and CLIP autoradiographs results are a representative of three independent

replicates. **(G)** The GSK-1264 binding pocket between CCD-CCD (brown) and CTD (light blue)

from different IN subunits is shown [62] (PDB ID: 5HOT). GSK-1264 is in green with nitrogen and

oxygen atoms colored blue and red, respectively. The side chain of R269 is shown with nitrogen

atoms colored in dark blue. Interactions between CTD R269 and CCD Q168 are indicated by

dashed lines.

Figure 8. Characterization of IN mutations present in latently infected CD4+ T-cells. (A)

Alignment from Gifford- (B) HEK293T cells were transfected with proviral HIV-1$_{NL4-3}$ expression

plasmids carrying the R224Q, S230N, E246K, and G272R IN mutations. Cell lysates and virions

were purified two days post transfection and analyzed by immunoblotting for CA and IN. The

image is representative of two independent experiments. **(C)** WT or IN mutant HIV-1L4-3 viruses

in cell culture supernatants were titered on TZM-bl indicator cells. The titers are presented relative

to WT (set to 1). The columns represent the average of three independent experiments and the

error bars represent SEM (****p<0.0001, by one-way ANOVA with Dunnett's multiple comparison

test). **(D)** Autoradiogram of IN-RNA adducts immunoprecipitated from virions bearing the

indicated substitutions in IN. Immunoblots below show the amount of immunoprecipitated IN.

Figure 1

Figure 2

A

HIV-1 IN

| 1 | NTD | 50 | CCD | 212 | CTD | 288 |

269   273

FRVYYRDSRDPVWKGPAKLLWKGEGAVVIQDNSDIKVVPRRKAKIIRDYGKQMAGDDCVASRQDED

B

IN R269A/K273A

WT - D2N D256R D270R D2R

51-

39-

28-

α-CA (cells)

α-CA (Virions)

α-IN

C

Relative infectivity

WT - D2N D256R D270R D2R

R269A/K273A
IN Backbone

D

IN R269A/K273A

WT - D2N D256R D2R

51-

39-

28-

IN-RNA

α-IN (IP)

E

Particle Release

Fold change

dpi

WT
D116N
R269A/K273A
D256N
D256R   R269A/K273A IN
D270N
D2N
D2R

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

A

IN codon 224
- 2.04% C
- 2.04% E
- 14.29% Q
- 81.63% R

Total=49

IN codon 230
- 2.08% L
- 10.42% N
- 87.50% S

Total=48

IN codon 246
- 97.87% E
- 2.13% K

Total=47

IN codon 272
- 88.00% G
- 12.00% R

Total=50

■ WT codon

B

WT, R269A/K273A, R224Q, S230N, E246K, G272R

51-
39-
28-

α--CA (cells)

α-CA (V)

C

Relative infectivity

*

WT, R269A/K273A, R224Q, S230N, E246K, G272R

D

WT, R269A/K273A, R224Q, S230N, E246K, G272R

51-
39-
28-

IN-RNA

α-IN (IP)

Figure 8