Changes in blood cell deformability in Chorea-Acanthocytosis and effects of treatment with dasatinib or lithium

Felix Reichel[1,2], Martin Kräter[1,2], Kevin Peikert[3,4], Hannes Glaß[3], Philipp Rosendahl[2], Maik Herbig[1,2], Alejandro Rivera Prieto[2], Alexander Kihm[5], Giel Bosman[6], Lars Kaestner[5,7], Andreas Hermann[3,4,8,9], Jochen Guck[1,2*]

[1]Max-Planck-Institut für die Physik des Lichts and Max-Planck-Zentrum für Physik und Medizin, Erlangen, Germany

[2]Biotechnology Center, Center for Molecular and Cellular Bioengineering, Technische Universität Dresden, Dresden, Germany

[3]Translational Neurodegeneration Section "Albrecht Kossel", Department of Neurology, University Medical Center Rostock, University of Rostock, Rostock, Germany

[4]Division for Neurodegenerative Diseases, Department of Neurology, Technische Universität Dresden, Dresden, Germany

[5]Department of Experimental Physics, Saarland University, Campus E2 6, Saarbrücken, Germany

[6]Department of Biochemistry, Radboud UMC, Nijmegen, The Netherlands

[7]Theoretical Medicine and Biosciences, Saarland University, Building 61.4, 66421 Homburg, Germany

[8]Deutsches Zentrum für Neurodegenerative Erkrankungen (DZNE), Rostock/Greifswald, Rostock, Germany

[9]Center for Regenerative Therapies Dresden, Technische Universität Dresden, Dresden, Germany

* Correspondence:
Jochen Guck
Jochen.guck@mpl.mpg.de

Keywords: chorea-acanthocytosis, blood cell deformability, real-time deformability cytometry, dasatinib, lithium, cell mechanics

Abstract

Misshaped red blood cells (RBCs), characterized by thorn-like protrusions known as acanthocytes, are
a key diagnostic feature in Chorea-Acanthocytosis (ChAc), a rare neurodegenerative disorder. The
altered RBC morphology likely influences their biomechanical properties which are crucial for the
cells to pass the microvasculature. Here, we investigated blood cell deformability of 5 ChAc patients
compared to healthy controls during up to one-year individual off-label treatment with the tyrosine
kinases inhibitor dasatinib or several weeks with lithium. Measurements with two microfluidic
techniques allowed us to assess RBC deformability under different shear stresses. Furthermore, we
characterized leukocyte stiffness at high shear stresses. The results show that blood cell deformability
– including both RBCs and leukocytes - in general is altered in ChAc patients compared to healthy
donors. Therefore, this study shows for the first time an impairment of leukocyte properties in ChAc.
During treatment with dasatinib or lithium, we observe alterations in RBC deformability and a stiffness
increase for leukocytes. The hematological phenotype of ChAc patients hints at a reorganization of the
cytoskeleton in blood cells which partly explains the altered mechanical properties observed here.
These findings highlight the need for a systematic assessment of the contribution of impaired blood
cell mechanics to the clinical manifestation of ChAc.

1 Introduction

Chorea-acanthocytosis/VPS13A disease (ChAc) is a rare monogenetic neurodegenerative disease of
the young adulthood affecting multiple systems other than the central nervous system, e.g., the red
blood cells (RBCs) (1–3). ChAc is characterized by various movement disorders (due to a degeneration
mainly of striatal neurons), epilepsy, cognitive decline and misshaped spiky RBCs, the latter being
referred to as acanthocytes (1,2,4,5). The autosomal-recessive condition is caused by loss of function
mutations in the *vacuolar protein sorting 13 homolog A* (*VPS13A*) gene (6–10). Therefore, the term
"VPS13A disease" has been recommended to replace the historical, more descriptive terminology (11).
A disease-modifying therapy has not been established so far.

It is still unclear, how exactly the loss of function of VPS13A leads to the manifestation of the cellular
and clinical phenotype. VPS13A localizes at different membrane contact sites and is most probably
involved in non-vesicular lipid transport as it is assumed also for the other members of the VPS13
protein family (for review, see Leonzino, Reinisch and De Camilli, 2021) (12–15). Two downstream
mechanisms are considered to be important drivers of ChAc pathophysiology: decreased
phosphoinositide-3-kinase (PI3K) signaling and increased activity of Src family tyrosine kinase Lyn
(for review, see Peikert et al. 2018 (2)).

Active Lyn kinase accumulates in ChAc RBCs and hyperphosphorylates membrane proteins such as
band 3 protein. Since band 3 is a structural protein, linking the cytoskeleton to the plasma membrane,
it is likely to be causally involved in the genesis of the spiky morphology. Accumulation of active Lyn
kinase was also found to be related to impaired autophagic flux (16,17). These phenotypes could be
reversed by *in vitro* treatment with tyrosine kinase inhibitors (TKI) such as the Src kinase family
inhibitor dasatinib (16). In a Vps13a knockout mouse model, the more specific Lyn kinase inhibitor
nilotinib improved both the hematological and neurological phenotypes by improving autophagy and
preventing neuroinflammation (18). In their recent work, Peikert et al., 2021 (19) report on single
individual treatment approaches targeting Lyn kinase with dasatinib in ChAc patients. They showed
that initially reduced F-actin signal, increased osmotic fragility and impaired autophagy were partially
restored in ChAc RBCs.

Furthermore, disturbed signaling via Phosphoinositide-3-kinase (PI3K) has been also identified as
relevant in the pathophysiology of ChAc patients. Reduced signaling via the PI3K-Rac1-PAK pathway
was reported to lead to disordered actin polymerization (20). Additionally, in other cells, altered PI3K
signaling caused a decreased ORAI1 expression and store-operated Ca^{2+} entry (SOCE), subsequently
disturbed Ca^{2+} homeostasis and apoptosis which could be partly reversed by *in vitro* lithium treatment
(21–23).

The rather late onset of the disease at the age of early adulthood and the slow progression rate (1,3,24)
lead to the assumption of a dependency between the altered RBC properties and neuronal degeneration:
Changes in the mechanical properties of the RBCs and hence reduced passage in micro-vessels and
capillaries may impair oxygen supply in certain areas of the brain that could over time accumulate to
tissue alterations leading to the symptoms described (25).

Altered RBC morphology is often linked to a change in cell mechanics which may lead to impaired
blood flow (26,27). Until now, the effect of acanthocytes in ChAc patients on the pathophysiology of
the disease remains elusive. Here, we describe, for the first time, altered deformability of blood cells
from ChAc patients including white blood cells (WBCs). In a second step, we observed these
parameters during individual off-label treatments with dasatinib or lithium, each targeting one of the
signaling pathways believed to be involved in the genesis of the RBC phenotype.

Deformability was measured using two techniques that asses the response to different ranges of shear
stresses acting on the cells: shape analysis of RBCs at low shear stress (0.1-3 Pa) (28) and real-time
fluorescence and deformability cytometry (RT-FDC) (29,30) at high shear stress (ca. 100 Pa). RBC
shape analysis only assesses the deformability of RBCs because the stresses are too low to deform
WBCs and the throughput is too low to measure a sufficient number of WBCs. In contrast, RT-FDC
was used to characterize the mechanics of both, RBCs and WBCs. In RT-FDC on RBCs, the
interpretation of the deformation data is not as straight forward as for the RBC shape analysis because
the RBCs undergo a non-trivial shape change here. We chose to use both methods to have a
comprehensive image of the blood cell deformability at different stresses. We demonstrate that the
deformability of all blood cells is affected in ChAc patients and this is further modulated by the
treatments. This highlights that monitoring blood cell mechanical properties of ChAc patients during
the course of disease and possible treatments can increase our understanding of this disease.

2   Materials and Methods

2.1   Cell source and reagents

We included 5 ChAc patients in this study for whom diagnosis was confirmed by Western blot
(absence of chorein/VPS13A protein) and genetic testing (31). The clinical parameters of the patients
are listed in Table S1 and complete blood count, RBC indices and hemolytic parameters are listed in
Table S2. Further information for P1-P3 can be found in Peikert et al., 2021 (19). Patients and healthy
control blood donors were enrolled in ongoing studies on the pathogenesis and natural history of
neurodegenerative diseases approved by the institutional review board of the Technische Universität
Dresden, Germany (EK 45022009, EK 78022015). The ChAc patients were treated with dasatinib (P1-
P3) or lithium (P4-P5) in the context of an individual off-label therapy based on the above described
preclinical evidence. Standard dose of 100 mg dasatinib per day was administered orally, as it was
lithium for which serum lithium concentration target range was defined as 0.6-0.8 mmol/l. Treatment
started after the blood for the baseline measurements was taken.

2.2   RBC shape analysis at low shear stress

The distribution of shapes for RBCs flowing through a narrow capillary at a fixed flow condition is
directly linked to the distribution of shear moduli within the population. Thus, changes in the shape
distribution are directly linked to a change of cell mechanics in the sample. This approach and the
experimental design are described in detail in Reichel et al., 2019 (28). Here, cells flow through a
roughly 5 mm long measurement channel that is 10 µm wide, and their shapes and dynamics are tracked
at the end of the channel over a length of 435 µm. Cells were recorded with the same setup used for
real-time deformability cytometry (29) using a different microfluidic chip. Examples of cells flowing
through the region of interest are shown in figure 1A,B and in supplementary videos S1-S8.

The shape analysis introduced in Reichel et al., 2019 (28), comprising the RBC shapes tumblers, tank-
treaders, parachutes and multilobes, cannot directly be used for samples containing acanthocytes
because their shape at rest differs from that of healthy RBCs. Further, acanthocytes show shapes not
observed for healthy cells (discocytes or normocytes) when deformed by the flow in the channel. For
this study, cells flowing through a square channel with a width of 10 µm at cell velocities ranging up
to 3 mm/s were characterized by their membrane morphology as normal looking (referred to as
normocytes) or as acanthocytes if they showed thorn-like protrusions. We did not further distinguish
between acanthocytes and echinocytes but argue that the mechanical properties of both cell types
should be very close (32). Furthermore, cells were classified as either deformed by the flow or still
maintaining their resting shape (undeformed). This classification was done by eye. We chose the
channel dimension of 10×10 µm because in such channels, RBCs undergo a sharp transition from
undeformed discocyte to parachute shape with a low probability of other transient shapes (28) which
makes it well suited to describe the deformed state of the cells. A change of the cell mechanical
properties should manifest in a change of the fraction of deformed cells in the sample and that the
velocity at which the majority of RBCs get deformed is shifted to higher values. For examples see
figure 1A,B.

The classification was done by eye from the recorded videos of cells passing the region of interest.
This resulted in a curve with probabilities to find cells in a deformed state for a given range of
velocities. The resulting curve for healthy RBCs is given in figure 2A (data from Reichel et al., 2019
(28), 3 control samples). To characterize differences to ChAc patient cells and changes on the cells
during treatment, a limited exponential growth function $P = 1 - e^{-\lambda \cdot v}$, with cell velocity v, was fitted
to the data. With this function, the probability to deform at $v = 0$ is zero and for very high velocities
the probability converges to 100% which motivated this function because cells will eventually deform
at high velocities. The cell deformability is characterized with the *growth rate* λ which describes how
quickly the cells reach a deformed state. The fitted curve for the control values is also depicted in
figure 2A. A higher growth rate would indicate that cells are more deformable.

The shape analysis method is not suited to measure the deformability of WBCs, because the stresses
acting the cells are too low and also the throughput of the method is not high enough to capture a
sufficient number of cells.

2.3   Real-time fluorescence and deformability cytometry (RT-FDC)

The setup of real-time fluorescence and deformability cytometry (RT-FDC) is described in detail in
Otto et al., 2015 (29) and Rosendahl et al., 2018 (30). In brief, flow is introduced into a microfluidic
chip with the help of syringe pumps and cells get deformed by hydrodynamic stresses as they pass a
narrow constriction, still larger than their own size. The chip is mounted on an inverted microscope
(Axiovert 200M, ZEISS, Oberkochen, Germany) and images are obtained at the end of the constriction

with a CMOS camera (Mikrotron, Unterschleissheim, Germany). Camera and syringe pump are
controlled via the measuring software ShapeIn (Zellmechanik Dresden, Dresden, Germany). ShapeIn
analyzes the recorded images in real-time and computes a contour and projected area for each cell. The
deformation parameter used in RT-FDC is defined as 1-circularity and is calculated from the contour's
area and perimeter using: $\text{deformation} = 1 - 2 \cdot \sqrt{\pi \cdot \text{Area}}/\text{perimeter}$, illustrated on the example of
a deformed RBC in figure 1C. As introduced by Toepfner et al., 2018 (33), the data from RT-FDC
experiments can be used to distinguish between different types of blood cells. These strategies were
used here to selectively study the deformation and mechanics of RBCs, lymphocytes, and myelocytes
from ChAc patients (see also SI text and fig. S5A,B).

Because the stresses required to deform RBCs are smaller than for leukocytes, both were measured
under different conditions. For measurements on leukocytes, citrate blood was resuspended in a
viscosity-adjusted measurement buffer (phosphate saline buffer without Mg^{2+} and Ca^{2+} (PBS−)
containing 0.5% (w/v) methyl cellulose (4000 cPs, Alfa Aesar 036718.22, CAS# 9004-67-5); adjusted
in HAAKE Falling Ball Viscometer type C (Thermo Fisher Scientific, Dreieich, Germany) using ball
number 3 to a viscosity of 15 mPa s) at a ratio of 1:20. The increased viscosity of the buffer also
increases the stresses acting on the cells inside the channel. Measurements were performed in channels
with 20×20 µm cross-section at a flow rate of 0.08 µl/s. Example images of a deformed lymphocyte
and myelocyte are shown in figure 1D and E, respectively. To decouple effects of the cell size on the
deformation, the Young's modulus was computed for leukocytes as described in Mokbel et al., 2017
(34). Since the derivation of Young's moduli for this system is only valid for initially spherical cells
(34,35), it cannot be calculated for RBCs.

RBC measurements were performed on an RT-FDC setup with a fluorescence module described in
detail in Rosendahl et al., 2018 (30). Whole blood was resuspended in the same measurement buffer
described above at a ratio of 1:200. The buffer was complemented with 2.5 µM syto13 nucleic acid
stain (Thermo Fisher Scientific, Dreieich, Germany) to mark for the reticulocytes in the sample.
Fluorescence was excited with a 60 mW, 488 nm laser at 8% power (OBIS 488-nm LS 60 mW,
Coherent Deutschland) and the signal was measured with an avalanche photodiode (MiniSM10035;
SensL Corporate, Cork, Ireland). Measurements were performed in channels with a 20×20 µm cross-
section at a flow rate of 0.02 µl/s. The control data for the RT-FDC measurements on RBCs was not
measured for this study but taken from 10 control samples for Rosendahl et al., 2018 (30) which were
measured under the same conditions (data shown in figure S4E,F).

2.4 Data analysis and data availability

186 Data was analyzed and plotted using custom python scripts. A detailed documentation can be found
187 here: https://gitlab.gwdg.de/freiche/changes-in-blood-cell-deformability-in-chorea-acanthocytosis-
188 and-effects-of-treatment-with-dasatinib-or-lithium. A collection of the raw data files is on figshare:
189 https://doi.org/10.6084/m9.figshare.c.5793482.

3 Results

3.1 Red blood cell deformability in Chorea-Acanthocytosis

To investigate differences in the deformability of red blood cells (RBCs) from chorea-acanthocytosis
(ChAc) patients compared to that of healthy donors, we performed shape-analysis and real-time
fluorescence and deformability cytometry (RT-FDC) measurements on the blood of 5 patients (P1-P5)
and compared it to measurements of healthy controls. The results are depicted in figure 1A,B and 1F,G.
Every velocity bin of 0.25 mm/s in figure 1A and B includes approx. 100 cells for each patient. The

shape-probability curves of normocytes (fig. 1A) and acanthocytes (fig. 1B) show that RBCs from
ChAc patients are less likely to deform in channel flow. It should be mentioned that also control
samples can include acanthocyte-like cells but to a much lesser extent than in patient samples. The
analysis for the control samples only included normocytes. Acanthocytes are clearly less likely to
deform due to flow even for higher flow velocities.

Analysis of the exponential growth rate (see fig. 2A) of control and ChAc normocytes is depicted in
figure 2B and showed that the growth rate for control normocytes was higher which indicated that the
cells are were deformable. This is in line with findings from Rabe et al., 2021 (32) which reported
similar results for ChAc normocytes. A comparison for acanthocytes was not possible because they
were not present in sufficient numbers in control samples.

The RT-FDC measurements showed that the RBCs of ChAc patients had a slightly larger projected
area than those from healthy donors (fig. 1F, effect size of +1.89 μm^2, p=0.0036 computed with linear
mixed effect models (36)). No significant difference between control and ChAc RBCs' deformation
could be detected (p=0.1382, fig. 1G).

Figure 1: Blood cell deformability of ChAc patients compared to healthy control blood. (A) Example
shapes of normal, discocyte RBCs (normocytes) in an **(i)** undeformed state and **(ii)** deformed by flow during
low shear stress flow in a 10 µm channel. The diagram shows the probability to find normocyte cells from the
sample in a deformed state at a given cell velocity in the channel for the acanthocyte patients and the mean curve
from 3 control measurements. Error bars represent SEM. scale bar represents 10 µm **(B)** Example shapes of
acanthocytes in an **(i)** undeformed state and **(ii)** deformed by flow. Below is the shape probability diagram for
the acanthocytes and the control curves. **(C)** Example shape of an RBC in RT-FDC and representation how the
deformation parameter is computed. All scale bars represent 10 µm. **(D)** Example image of a lymphocyte in RT-
FDC. **(E)** Example image of a myelocyte. **(F)**, **(G)** Projected area within the contour and deformation of RBCs
of the ChAc patients from RT-FDC measurements vs. pooled data from 10 control measurements (full data in
fig. S4). White dots represent the median value; grey box in the violin shows inter-quartile range (IQR) and
extended lines 1.5×IQR. Dashed grey line shows the median control value. P-values were calculated with
linear-mixed effect models as described in Herbig et al., 2018 (36). Numbers on top of the plots indicate the
number of observations per violin. **(H)** Young's modulus of lymphocytes from ChAc patients vs. control (n=1).
(I) Young's modulus of myelocytes from ChAc patients vs. control (n=1).

3.2 White blood cell deformability in Chorea-Acanthocytosis

In RT-FDC measurements on whole blood it is possible to distinguish different cell types (see materials
and methods, SI text and Töpfner et al., 2018 (33)). Thus, we investigated mechanical differences also
between the leukocytes of ChAc patients and one healthy donor. We distinguished, mainly by size,
between all lymphocytes and cells resulting from the myeloid lineage without RBCs (myelocytes),
comprising all mono- and granulocytes. The myelocyte fraction mainly consisted of neutrophil
granulocytes (>80%).

Since white blood cells are spherical at rest, we can employ a model that maps the projected area and
the deformation to an apparent Young's modulus (34), to see differences in stiffness before, during and
after the treatments. As depicted in figure 1H,I, both lymphocytes and myelocytes of ChAc patients
were significantly stiffer than their healthy counterparts before treatment onset (lymphocytes: +240 Pa,
p=0.0186, myelocytes: +150 Pa, p=0.0070; effect sizes and p-values determined by linear mixed effects
model analysis (36)). The projected area and deformation data used to compute the Young's modulus
is shown in figure S5C,D.

3.3 Dasatinib and lithium treatment effect on RBC deformability

3.3.1 RBC shape analysis during dasatinib or lithium treatment

To study the effect of dasatinib or lithium on RBC deformability of ChAc patients, we monitored RBCs
during treatment with shape analysis measurements and RT-FDC.

To compare shape-analysis data over the treatment time, a limited exponential growth was fitted to the
data (see fig. 2A) and the fitted growth rate was used to detect changes in the deformability during the
treatment time. The results for the dasatinib treatment are shown in figure 2C,D, those of the lithium
in figure 3. Even though one might get the impression that the growth rate showed a slight increase
with dasatinib treatment time for both normocytes and acanthocytes, the overall RBC deformability
remained unaffected by the treatment. The results pooled for all RBCs are shown in figure S1 and the
curves from all measurements are given in figure S2.

The acanthocyte count over the treatment is given in figure 2E and shows that the fraction of
acanthocytes remained constant within a certain range of fluctuation without a systematic trend for
both treatments, which was in line with the clinical data (19).

To check if dasatinib had a short-term effect on RBC deformability, RBCs from P3 were treated with
dasatinib *in vitro* and the shapes in channel flow were analyzed (for details, see SI text). The results
are shown in figure 2F. There was no significant difference in the deformability between control- and
dasatinib-treated RBCs. This indicated that the observed effects on cell deformability were likely
caused by long-term treatment effects, e.g., acting on erythropoiesis or indirect systemic effects.

For the lithium treatment, normocyte growth rate even increased above control values during the
treatment and recovered to, or below, pre-treatment levels after the treatment was discontinued.
Acanthocytes showed a slight increase in growth rate during lithium treatment. This indicated that the
RBCs got softer during lithium treatment but RT-FDC results showed that the cells also got larger with
the treatment (see fig. 4) which would make the cells more likely to deform in the channel if the overall
mechanics are unchanged.

Figure 2: Dasatinib treatment effect on RBC deformability. (A) Illustration of the limited exponential growth function fit, used to characterize the shape probability curves presented on the control data. **(B)** Fitted growth rate for normocytes of control samples vs samples from ChAc patients. P-value calculated by Welch's t-test (p = 0.009). **(C)** Fitted growth rate λ of the shape probability curves from ChAc patients as a function of treatment time with dasatinib for normocytes and **(D)** acanthocytes respectively. Dashed lines represent datapoints taken after treatment was stopped. Errorbars represent standard error of the fit, calculated from the covariance matrix. Gray region shows the respective value from healthy donors without treatment (Mean ± SD). **(E)** Percentage of acanthocytes found in each sample during shape analysis measurements for patients P1-P5. Dashed lines indicate datapoints taken after the respective treatment already stopped. **(F)** Fraction of healthy shapes (hs) and acanthocytes (ac) over cell velocity for untreated and *in-vitro* dasatinib treated RBCs from patient P3.

279 **Figure 3: Lithium treatment effect on RBC deformability.** Fitted growth rate λ of the shape probability
280 curves from ChAc patients as a function of treatment time with lithium for normocytes and acanthocytes
281 respectively. Dashed lines represent datapoints taken after treatment was stopped. Errorbars represent standard
282 error of the fit, calculated from the covariance matrix. Gray region shows the respective value from healthy
283 donors without treatment (Mean ± SD).

3.3.2 RBC deformation in RT-FDC during dasatinib or lithium treatment

RT-FDC results for the dasatinib treatment are shown in figure 4A-C and for the lithium treatment in
figure 4D-F. It can be seen that during treatment with dasatinib, the RBC deformation parameter
dropped for all patients between 50-100 days after treatment start, while the projected area remained
mainly constant with the exception of some outliers.

For the lithium treatment, the RT-FDC results showed that the deformation parameter decreased with
treatment time which was accompanied by an increase in cell size. After the treatment, the deformation
and area values returned almost to the control values for P5 while the values for P4 did not fully
recover.

The interpretation of deformation values for RBCs in RT-FDC cannot directly be linked to cell
mechanics because they undergo a non-trivial shape transition from discocyte to tear-drop shape (28).
A decrease of the deformation value does not necessarily mean that the cells got stiffer. To back this
up, we artificially stiffened RBCs from healthy donors with glutaraldehyde (37,38) or diamide (39)
(see SI text and fig. S3) and saw that an increasing concentration of the reagents led to an increase of
the deformation when working at low concentrations. At high concentrations the cell shape got fixed
which led to lower deformation values.

Since reticulocytes have been reported to have a higher projected area and lower deformation compared
to the whole RBC population (30), we speculated that above mentioned results might be due to
increased levels of reticulocytes in ChAc patients. The results for reticulocytes, labelled with syto13
nucleic acid stain, together with all RBCs are given in figure S4A-D. The patients' reticulocytes indeed
had higher projected area and lower deformation if compared to the whole RBC population which did
not differ from healthy controls. Qualitatively, reticulocyte area and deformation changed over
treatment time like the total RBC population. The fraction of reticulocytes in the blood was initially
higher compared to controls and slightly decreased during the dasatinib treatment closer to values from
healthy donors (fig. 4C). A reduction of the reticulocyte count with dasatinib treatment was also
reported for VPS13a[-/-] mice phenocopying human ChAc (18). During lithium treatment P4 showed a
decrease during and after treatment, while P5 showed a slight increase but values were mostly in the
range expected for healthy individuals (0.5-2.5%).

Figure 4: RBC deformation, size and reticulocyte count in RT-FDC during treatment with dasatinib or lithium. (A-C) RBC median deformation, projected area and reticulocyte fraction measured with RT-FDC before, during and after treatment with dasatinib. **(D-F)** RBC median deformation, projected area and reticulocyte fraction measured with RT-FDC before, during and after treatment with lithium. Dashed lines indicate time points after the treatment was stopped, dashed gray lines indicate control values without treatment. The red region in the reticulocyte plots indicates the range for healthy individuals reported in the literature of 0.5-2.5%.

3.4 ChAc Leukocytes stiffened during treatment with dasatinib

The Young's moduli of lymphocytes and myelocytes in dasatinib treatment are shown in figure 5A,B. The lithium data is shown in figure 5C,D. During treatment with dasatinib, we observed a further increase of the Young's modulus for all three patients which also persisted for the lymphocytes shortly after the treatment stopped. For myelocytes, the median Young's modulus decreased for patients P1 and P2 shortly after the treatment but kept increasing for P3.

During treatment with lithium both patients showed an increase in lymphocyte and myelocyte stiffness. After the treatment was stopped, we observed a further increase for patient P4 and a drop back to the pre-treatment level for patient P5 approximately one year after the treatment stopped. The area and deformation data used to compute the Young's modulus is given in figure S5E,F.

Figure 5: Leukocyte stiffness in RT-FDC during treatment with Dasatinib or lithium. (A) Median Young's
modulus of ChAc patients' lymphocytes during dasatinib treatment. **(B)** Median Young's modulus of ChAc
patients' myelocytes during dasatinib treatment. **(C)** Median Young's modulus of ChAc patients' lymphocytes
during lithium treatment. **(D)** Median Young's modulus of ChAc patients' myelocytes during lithium treatment.

4   Discussion

Our study highlights that the deformability, not only of RBCs but of all blood cell types, was altered
in ChAc patients. Shape analysis results showed a decreased deformability of ChAc RBCs compared
to controls. The decreased deformability can be directly inferred from the reduced number of deformed
cells at any cell velocity which was also reflected by a smaller growth rate. A reduced deformability
of ChAc normocytes, deduced from cell shapes, has recently also been reported by Rabe et al., 2021
(32).

In RT-FDC, we did not observe a change in the deformation parameter but ChAc RBCs had a larger
projected area. Because deformation and size are always correlated in RT-FDC (35), this is also an
indicator for a change of RBC deformability, but the interpretation of this behavior is not straight
forward. To see how the deformation parameter changes when RBC stiffness is controllably changed,
we treated RBCs *in vitro* with glutaraldehyde or diamide. At low concentrations, we observed an
increase of the deformation parameter which shows that higher deformation parameter can also occur
for stiffer RBCs.

While we observed larger RBCs in ChAc patients, values for the mean corpuscular volume (MCV)
reported in Peikert et al., 2021 (19) (see also Table S2) did not show an increased RBC volume for
patients P1-P3. Sizes reported in RT-FDC measurements are based on the projected cross-sectional
area of the deformed cells. Simulations of the flow fields inside the channel suggest that the stresses
near the centerline can be approximated as axisymmetric (40). Therefore, it is fair to assume that the
RBCs take up an axisymmetric shape as well and the projected area is directly correlated to cell size.
In automated cell counters, the MCV is determined, e.g., by impedance or light scattering
measurements which require a constant deformability of RBCs for a reliable result. Thus, MCV is not
a direct measurement of single cell sizes. Here, we provide a direct observation of sizes based on single
cell imaging.

Both, shape analysis and RT-FDC indicated a change of RBC deformability during treatment with
dasatinib. Shape analysis experiments showed a slight increase of the growth rate with treatment time,
which indicated an increased deformability of both normocytes and acanthocytes. In RT-FDC, we
observed a sharp drop of the deformation parameter after 50-100 treatment days while the size
fluctuated around a constant value. Again, for RBCs a direct link between the deformation parameter
in RT-FDC and cell mechanics is not trivial and a decrease in deformation does not necessarily mean
that the cells got stiffer (see fig. S3). Even though we do not have certainty of the direction of the
change, we claim that RBC mechanics was altered during dasatinib treatment.

During lithium treatment, we also observed an increasing growth rate in the shape analysis which
indicated increased RBC deformability. In RT-FDC, we saw a decrease of deformation accompanied
by an increase in projected area. An increase in projected area and decrease in deformation hints at a
volume increase and a reduced surface area-to-volume ratio, which leads to an increased sphericity or
effective rounding of the cells. Rounder shapes lead to smaller deformation values, assuming that this
effect dominates any stiffness changes of the cells. The increased cell size could also explain the
increased deformability in the shape analysis experiments because larger cells experience higher
stresses and are therefore more likely to deform. All in all, these findings indicate that lithium affects
RBC properties in ChAc patients but because of the scarcity of the data here, one should be careful
with the interpretation.

Data from Peikert et al., 2021 indicated that with dasatinib treatment, F-actin became more localized
at the cortex (19). These modifications in the actin network likely contribute to changes in RBC
deformability (41) that we observed here.

Treatment with dasatinib led to a decrease of the reticulocyte fraction in the blood of ChAc patients.
The same effect was observed in a mouse model phenocopying human ChAc (18). Peikert et al., 2021
reported an abundance of the autophagy initiator Ulk1 in ChAc patients RBCs, which decreased with
dasatinib treatment (19). Ulk1 is connected to autophagy in erythropoiesis and could influence the rate
of reticulocyte production (42), providing a mechanistic link to the reticulocyte count.

While an impaired RBC deformability was described before for patients with other forms of
neuroacanthocytosis (43) this is, to our knowledge, the first time that also patients' leukocyte stiffness
was studied. Lymphocytes and myelocytes were both stiffer in ChAc patients compared to cells from
healthy donors. This is of interest as VPS13A, the gene which is known to have a loss-of-function
mutation in ChAc, is especially expressed in human monocytes and B-, and helper T-cells (44). Since
VPS13A is known to cause structural reorganizations of the cytoskeleton from studies on other cell
types (17,20,45), it is likely that we observed these as altered mechanical properties in leukocytes.
Another feature in ChAc blood is an increased expression of Lyn kinase, which is known to set the
threshold for B-cell activation. Mechanical changes of lymphocytes after activation have been reported
before (33,46). Another mechanism that could lead to altered leukocyte mechanics is an increase in
interleukin-1 beta (IL-1b) that was observed in the brain of Vps13a knockout mice (18). IL-1b not only
plays a key role in autoinflammation but is also known to cause re-organization of the actin
cytoskeleton in different cell types (47,48).

Young's moduli of both lymphocytes and myelocytes increased even further away from control values
during treatment with dasatinib or lithium. A stiffening was previously also observed for leukemic cells
during treatment with different chemotherapeutic agents for acute lymphoblastic leukemia
(dexamethasone and daunorubicin), such as dasatinib (49). Dasatinib interferes with all the pathways
described above and likely induces further mechanical changes in leukocytes. The inhibited activity of

Lyn during dasatinib treatment could cause a change in lymphocyte activation levels leading to altered
mechanics but the exact mechanisms for the observed leukocyte stiffening with dasatinib treatment
remain rather speculative at this point.

Overall, our results showed that the mechanical phenotype of all blood cell types in ChAc patients
were impaired but did not change towards regimes of healthy donors during the treatments. Clinical
data in Peikert et al., 2021 (19) hints on slightly improved hematological features but these did not
manifest in an improved neurological phenotype. This demonstrates that the techniques that were
utilized here are suited to measure a treatment response.

In general, decreased blood cell deformability often leads to clinical symptoms due to decreased cell
survival and secondary pathology (25–27,50). Especially, an impaired oxygen transport by altered
blood cell deformability should be considered in neurological diseases (51). While clinical descriptions
of ChAc patients focus on neurological symptoms, we highlight in this study that the effects on blood
deformability should not be neglected and could contribute to the clinical manifestation (through the
mechanisms stated above) and thus methods that are able to monitor this (as shown here) should be
considered as read outs for clinical trials. Our results on the impaired mechanics of leucocytes
especially highlights that the focus of ChAc research should not only be on RBCs but also include
other blood cell types.

5   Conflict of Interest

422 The authors declare that the research was conducted in the absence of any commercial or financial
423 relationships that could be construed as a potential conflict of interest.

6   Author Contributions

425 Conceptualization: F.R., M.K., K.P., G.B., A.H., and J.G.; Experiments–shape analysis: F.R. and
426 A.R.P., Experiments–RT-FDC: F.R., P.R., M.H., and M.K., Experiments–in-vitro shape analysis:
427 A.K.; Formal analysis: F.R., P.R., A.K., and M.K.; Supervision: L.K.,A.H., and J.G.; Writing–
428 original draft, K.P. and F.R..; Writing–review and editing, F.R., M.K., K.P., G.B., L.K., A.H., and
429 J.G.; All authors have read and agreed to the published version of the manuscript.

7   Funding

K.P. was supported by the Else Kröner clinician scientist program and the MeDDrive program (TU
Dresden, Germany), as well as by the Rostock Academy for Clinician Scientists (RACS) and the
FORUN program (University of Rostock, Germany). A.H. was supported by the "Hermann und Lilly
Schilling-Stiftung für medizinische Forschung im Stifterverband". Expenses for the off-label
dasatinib prescription were gratefully covered in part by the Center for Regenerative Therapies
Dresden (CRTD, TU Dresden, Germany) and the public healthcare insurance of the patients.

8   Acknowledgments

438 The authors thank all patients and their families for giving consent for publication of the data and the
439 healthy control subjects for participation in this study. We are especially grateful to Glenn (†) and
440 Ginger Irvine as the founders of the Advocacy for Neuroacanthocytosis Patients

441 (www.naadvocacy.org) and to Susan Wagner and Joy Willard-Williford as representatives of the NA
442 Advocacy USA (www.naadvocacyusa.org).

9   Supplementary Material

444 Supplementary Information, Figures, Table and Videos will be available online.

10   Data Availability statement

446 Raw   data   files   and   initial   analysis   scripts   are   available   on   figshare.com:
447 https://doi.org/10.6084/m9.figshare.c.5793482. Analysis documentation and creation of figure plots is
448 documented on gitlab: https://gitlab.gwdg.de/freiche/changes-in-blood-cell-deformability-in-chorea-
449 acanthocytosis-and-effects-of-treatment-with-dasatinib-or-lithium.

11   References

1.   Jung HH, Danek A, Walker RH. Neuroacanthocytosis Syndromes. Orphanet J Rare Dis.
(2011) 6(1):68. doi: 10.1186/1750-1172-6-68

2.   Peikert K, Danek A, Hermann A. Current state of knowledge in Chorea-Acanthocytosis as
core Neuroacanthocytosis syndrome. Eur J Med Genet. (2018) 61(11):699–705. doi:
10.1016/j.ejmg.2017.12.007

3.   Walker RH. Chorea-Acanthocytosis. In: Reference Module in Neuroscience and Biobehavioral
Psychology. Elsevier; (2017.). doi: 10.1016/B978-0-12-809324-5.00521-6

4.   Levine IM, Estes JW, Looney JM. Hereditary Neurological Disease With Acanthocytosis: A
New Syndrome. Arch Neurol. (1968) 19(4):403–9. doi:
10.1001/archneur.1968.00480040069007

5.   Critchley EM, Clark DB, Wikler A. Acanthocytosis and neurological disorder without
betalipoproteinemia. Arch Neurol. (1968) 18(2):134–40. doi:
10.1001/archneur.1968.00470320036004

6.   Danek A, Bader B, Velayos-Baeza A, Walker RH. Autosomal recessive transmission of
chorea-acanthocytosis confirmed. Acta Neuropathol. (2012) 123(6):905–6. doi:
10.1007/s00401-012-0971-y

7.   Dobson-Stone C, Danek A, Rampoldi L, Hardie R, Chalmers R, Wood N, et al. Mutational
spectrum of the CHAC gene in patients with chorea-acanthocytosis. Eur J Hum Genet. (2002)
10(11):773–81. doi: 10.1038/sj.ejhg.5200866

8.   Rampoldi L, Dobson-Stone C, Rubio JP, Danek A, Chalmers RM, Wood NW, et al. A
conserved sorting-associated protein is mutant in chorea-acanthocytosis. Nat Genet. (2001)
28(2):119–20. doi: 10.1038/88821

9.   Ueno S, Maruki Y, Nakamura M, Tomemori Y, Kamae K, Tanabe H, et al. The gene encoding
a newly discovered protein, chorein, is mutated in chorea-acanthocytosis. Nat Genet. (2001)
28(2):121–2. doi: 10.1038/88825

10. Nishida Y, Nakamura M, Urata Y, Kasamo K, Hiwatashi H, Yokoyama I, et al. Novel
pathogenic VPS13A gene mutations in Japanese patients with chorea-acanthocytosis. Neurol
Genet. (2019) 5(3):e332. doi: 10.1212/NXG.0000000000000332

11. Walker RH, Danek A. "Neuroacanthocytosis" – Overdue for a Taxonomic Update. Tremor
and Other Hyperkinetic Movements. (2021) 11(1):1–6. doi: 10.5334/tohm.583

12. Leonzino M, Reinisch KM, De Camilli P. Insights into VPS13 properties and function reveal a
new mechanism of eukaryotic lipid transport. Biochim Biophys Acta - Mol Cell Biol Lipids.
(2021) 1866(10):159003. doi: 10.1016/j.bbalip.2021.159003

13. Gao M, Yang H. VPS13: A lipid transfer protein making contacts at multiple cellular
locations. J Cell Biol. (2018) 217(10):3322–4. doi: 10.1083/jcb.201808151

14. Kumar N, Leonzino M, Hancock-Cerutti W, Horenkamp FA, Li P, Lees JA, et al. VPS13A
and VPS13C are lipid transport proteins differentially localized at ER contact sites. J Cell Biol.
(2018) 217(10):3625–39. doi: 10.1083/jcb.201807019

15. Yeshaw WM, van der Zwaag M, Pinto F, Lahaye LL, Faber AIE, Gómez-Sánchez R, et al.
Human VPS13A is associated with multiple organelles and influences mitochondrial
morphology and lipid droplet motility. Elife. (2019) 8. doi: 10.7554/eLife.43561

16. Lupo F, Tibaldi E, Matte A, Sharma AK, Brunati AM, Alper SL, et al. A new molecular link
between defective autophagy and erythroid abnormalities in chorea-acanthocytosis. Blood.
(2016) 128(25):2976–87. doi: 10.1182/blood-2016-07-727321

17. De Franceschi L, Tomelleri C, Matte A, Brunati AM, Bovee-Geurts PH, Bertoldi M, et al.
Erythrocyte membrane changes of chorea-acanthocytosis are the result of altered Lyn kinase
activity. Blood. (2011) 118(20):5652–63. doi: 10.1182/blood-2011-05-355339

18. Peikert K, Federti E, Matte A, Constantin G, Pietronigro EC, Fabene PF, et al. Therapeutic
targeting of Lyn kinase to treat chorea-acanthocytosis. Acta Neuropathol Commun. (2021)
9(1):81. doi: 10.1186/s40478-021-01181-y

19. Peikert K, Glaß H, Federti E, Matte A, Pelzl L, Akgün K, et al. Targeting Lyn Kinase in
Chorea-Acanthocytosis: A Translational Treatment Approach in a Rare Disease. J Pers Med.
(2021) 11(5):392. doi: 10.3390/jpm11050392

20. Föller M, Hermann A, Gu S, Alesutan I, Qadri SM, Borst O, et al. Chorein-sensitive
polymerization of cortical actin and suicidal cell death in chorea-acanthocytosis. FASEB J.
(2012) 26(4):1526–34. doi: 10.1096/fj.11-198317

21. Pelzl L, Elsir B, Sahu I, Bissinger R, Singh Y, Sukkar B, et al. Lithium Sensitivity of Store
Operated Ca2+ Entry and Survival of Fibroblasts Isolated from Chorea-Acanthocytosis
Patients. Cell Physiol Biochem. (2017) 42(5):2066–77. doi: 10.1159/000479901

22. Pelzl L, Hauser S, Elsir B, Sukkar B, Sahu I, Singh Y, et al. Lithium Sensitive ORAI1
Expression, Store Operated Ca2+ Entry and Suicidal Death of Neurons in Chorea-
Acanthocytosis. Sci Rep. (2017) 7(1):6457. doi: 10.1038/s41598-017-06451-1

23.   Yu W, Honisch S, Schmidt S, Yan J, Schmid E, Alkahtani S, et al. Chorein Sensitive Orai1
Expression and Store Operated Ca2+ Entry in Rhabdomyosarcoma Cells. Cell Physiol
Biochem. (2016) 40(5):1141–52. doi: 10.1159/000453168

24.   Rampoldi L, Danek A, Monaco AP. Clinical features and molecular bases of
neuroacanthocytosis. J Mol Med. (2002) 80(8):475–91. doi: 10.1007/s00109-002-0349-z

25.   Adjobo-Hermans MJW, Cluitmans JCA, Bosman GJCGM. Neuroacanthocytosis:
Observations, Theories and Perspectives on the Origin and Significance of Acanthocytes.
Tremor and Other Hyperkinetic Movements. (2015) 5:328. doi: 10.5334/tohm.271

26.   An X, Mohandas N. Disorders of red cell membrane. Br J Haematol. (2008) 141:367–75. doi:
10.1111/j.1365-2141.2008.07091.x

27.   Da Costa L, Galimand J, Fenneteau O, Mohandas N. Hereditary spherocytosis, elliptocytosis,
and other red cell membrane disorders. Blood Rev. (2013) 27(4):167–78. doi:
10.1016/j.blre.2013.04.003

28.   Reichel F, Mauer J, Nawaz AA, Gompper G, Guck J, Fedosov DA. High-Throughput
Microfluidic Characterization of Erythrocyte Shapes and Mechanical Variability. Biophys J.
(2019) 117(1):14–24. doi: 10.1016/j.bpj.2019.05.022

29.   Otto O, Rosendahl P, Mietke A, Golfier S, Herold C, Klaue D, et al. Real-time deformability
cytometry: on-the-fly cell mechanical phenotyping. Nat Methods. (2015) 12(3):199–202. doi:
10.1038/nmeth.3281

30.   Rosendahl P, Plak K, Jacobi A, Kraeter M, Toepfner N, Otto O, et al. Real-time fluorescence
and deformability cytometry. Nat Methods. (2018) 15(5):355–8. doi: 10.1038/nmeth.4639

31.   Dobson-Stone C, Velayos-Baeza A, Filippone LA, Westbury S, Storch A, Erdmann T, et al.
Chorein detection for the diagnosis of chorea-acanthocytosis. Ann Neurol. (2004) 56(2):299–
302. doi: 10.1002/ana.20200

32.   Rabe A, Kihm A, Darras A, Peikert K, Simionato G, Dasanna AK, et al. The Erythrocyte
Sedimentation Rate and Its Relation to Cell Shape and Rigidity of Red Blood Cells from
Chorea-Acanthocytosis Patients in an Off-Label Treatment with Dasatinib. Biomolecules.
(2021) 11(5):727. doi: 10.3390/biom11050727

33.   Toepfner N, Herold C, Otto O, Rosendahl P, Jacobi A, Kräter M, et al. Detection of human
disease conditions by single-cell morpho-rheological phenotyping of blood. Elife. (2018)
7:e29213. doi: 10.7554/eLife.29213

34.   Mokbel M, Mokbel D, Mietke A, Träber N, Girardo S, Otto O, et al. Numerical Simulation of
Real-Time Deformability Cytometry To Extract Cell Mechanical Properties. ACS Biomater
Sci Eng. (2017) 3(11):2962–73. doi: 10.1021/acsbiomaterials.6b00558

35.   Mietke A, Otto O, Girardo S, Rosendahl P, Taubenberger A, Golfier S, et al. Extracting Cell
Stiffness from Real-Time Deformability Cytometry: Theory and Experiment. Biophys J.
(2015) 109(10):2023–36. doi: 10.1016/j.bpj.2015.09.006

36.   Herbig M, Mietke A, Müller P, Otto O. Statistics for real-time deformability cytometry:
552   Clustering, dimensionality reduction, and significance testing. Biomicrofluidics. (2018)
553   12(4):042214. doi: 10.1063/1.5027197

554   37.   Holmes D, Whyte G, Bailey J, Vergara-Irigaray N, Ekpenyong A, Guck J, et al. Separation of
555   blood cells with differing deformability using deterministic lateral displacement. Interface
556   Focus. (2014) 4(6):20140011. doi: 10.1098/rsfs.2014.0011

557   38.   Forsyth AM, Wan J, Ristenpart WD, Stone HA. The dynamic behavior of chemically
558   "stiffened" red blood cells in microchannel flows. Microvasc Res. (2010) 80(1):37–43. doi:
559   10.1016/j.mvr.2010.03.008

560   39.   Sinha A, Chu TTT, Dao M, Chandramohanadas R. Single-cell evaluation of red blood cell bio-
561   mechanical and nano-structural alterations upon chemically induced oxidative stress. Sci Rep.
562   (2015) 5(1):9768. doi: 10.1038/srep09768

563   40.   Panhwar MH, Czerwinski F, Dabbiru VAS, Komaragiri Y, Fregin B, Biedenweg D, et al.
564   High-throughput cell and spheroid mechanics in virtual fluidic channels. Nat Commun. (2020)
565   11(1):2190. doi: 10.1038/s41467-020-15813-9

566   41.   Gokhin DS, Nowak RB, Khoory JA, Piedra A de la, Ghiran IC, Fowler VM. Dynamic actin
567   filaments control the mechanical behavior of the human red blood cell membrane. Pollard TD,
568   editor. Mol Biol Cell. (2015) 26(9):1699–710. doi: 10.1091/mbc.E14-12-1583

569   42.   Kundu M, Lindsten T, Yang C-Y, Wu J, Zhao F, Zhang J, et al. Ulk1 plays a critical role in
570   the autophagic clearance of mitochondria and ribosomes during reticulocyte maturation.
571   Blood. (2008) 112(4):1493–502. doi: 10.1182/blood-2008-02-137398

572   43.   Cluitmans JCA, Tomelleri C, Yapici Z, Dinkla S, Bovee-Geurts P, Chokkalingam V, et al.
573   Abnormal Red Cell Structure and Function in Neuroacanthocytosis. Connes P, editor. PLoS
574   One. (2015) 10(5):e0125580. doi: 10.1371/journal.pone.0125580

575   44.   Samaras P, Schmidt T, Frejno M, Gessulat S, Reinecke M, Jarzab A, et al. ProteomicsDB: a
576   multi-omics and multi-organism resource for life science research. Nucleic Acids Res. (2019)
577   48(D1):D1153–63. doi: 10.1093/nar/gkz974

578   45.   Rzepnikowska W, Flis K, Muñoz-Braceras S, Menezes R, Escalante R, Zoladek T. Yeast and
579   other lower eukaryotic organisms for studies of Vps13 proteins in health and disease. Traffic.
580   (2017) 18(11):711–9. doi: 10.1111/tra.12523

581   46.   Zhang X, Kim T-H, Thauland TJ, Li H, Majedi FS, Ly C, et al. Unraveling the
582   mechanobiology of immune cells. Curr Opin Biotechnol. (2020) 66:236–45. doi:
583   10.1016/j.copbio.2020.09.004

584   47.   Franco-Barraza J, Valdivia-Silva JE, Zamudio-Meza H, Castillo A, García-Zepeda EA,
585   Benítez-Bribiesca L, et al. Actin Cytoskeleton Participation in the Onset of IL-1β Induction of
586   an Invasive Mesenchymal-like Phenotype in Epithelial MCF-7 Cells. Arch Med Res. (2010)
587   41(3):170–81. doi: 10.1016/j.arcmed.2010.04.010

48.   Lie PPY, Cheng CY, Mruk DD. The biology of interleukin-1: emerging concepts in the

regulation of the actin cytoskeleton and cell junction dynamics. Cell Mol Life Sci. (2012)
69(4):487–500. doi: 10.1007/s00018-011-0760-0

49. Lam WA, Rosenbluth MJ, Fletcher DA. Chemotherapy exposure increases leukemia cell
stiffness. Blood. (2007) 109(8):3505–8. doi: 10.1182/blood-2006-08-043570

50. Brown CD, Ghali HS, Zhao Z, Thomas LL, Friedman EA. Association of reduced red blood
cell deformability and diabetic nephropathy. Kidney Int. (2005) 67(1):295–300. doi:
10.1111/j.1523-1755.2005.00082.x

51. Kikuchi Y, Da Q-W, Fujino T. Variation in Red Blood Cell Deformability and Possible
Consequences for Oxygen Transport to Tissue. Microvasc Res. (1994) 47(2):222–31. doi:
10.1006/mvre.1994.1017

RBC shape analysis at low shear stress (0.1-1 Pa)

RT-FDC (shear stress = 100-400 Pa)

A — Normocytes

i) ii)

Probability for deformed shape vs Velocity [mm/s]

P1, P2, P3, P4, P5, CTRL

B — Acanthocytes

i) ii)

Probability for deformed shape vs Velocity [mm/s]

P1, P2, P3, P4, P5, CTRL

Red blood cells

C

$$\text{Deformation} = 1 - \frac{2 \cdot \sqrt{\pi \cdot \text{Area}}}{\text{Perimeter}}$$

F Projected area [µm²]

129114 103938 16718 16637 15555 8862

p = 0.0036

CTRL P1 P2 P3 P4 P5 — Patient

G Deformation

p = 0.1382

CTRL P1 P2 P3 P4 P5 — Patient

White blood cells

D **E**

H Lymphocytes

Young's modulus (stiffness) [kPa]

385 236 213 175 159 149

p = 0.0186

CTRL P1 P2 P3 P4 P5 — Patient

I Myelocytes

Young's modulus (stiffness) [kPa]

495 363 331 217 207 160

p = 0.0070

CTRL P1 P2 P3 P4 P5 — Patient