Changes in blood cell deformability in Chorea-Acanthocytosis and effects of treatment with dasatinib or lithium

Felix Reichel[1,2], Martin Kräter[1,2], Kevin Peikert[3,4], Hannes Glaß[3], Philipp Rosendahl[2], Maik
Herbig[1,2], Alejandro Rivera Prieto[2], Alexander Kihm[5], Giel Bosman[6], Lars Kaestner[5,7],
Andreas Hermann[3,4,8,9], Jochen Guck[1,2]*

[1]Max-Planck-Institut für die Physik des Lichts and Max-Planck-Zentrum für Physik und Medizin,
Erlangen, Germany

[2]Biotechnology Center, Center for Molecular and Cellular Bioengineering, Technische Universität
Dresden, Dresden, Germany

[3]Translational Neurodegeneration Section "Albrecht Kossel", Department of Neurology, University
Medical Center Rostock, University of Rostock, Rostock, Germany

[4]Division for Neurodegenerative Diseases, Department of Neurology, Technische Universität
Dresden, Dresden, Germany

[5]Department of Experimental Physics, Saarland University, Campus E2 6, Saarbrücken, Germany

[6]Department of Biochemistry, Radboud UMC, Nijmegen, The Netherlands

[7]Theoretical Medicine and Biosciences, Saarland University, Building 61.4, 66421 Homburg,
Germany

[8]German Center for Neurodegenerative Diseases (DZNE), Dresden, Germany

[9]Center for Regenerative Therapies Dresden, Technische Universität Dresden, Dresden, Germany

* Correspondence:
Jochen Guck
Jochen.guck@mpl.mpg.de

Keywords: chorea-acanthocytosis, blood cell deformability, real-time deformability cytometry,
dasatinib, lithium, cell mechanics (Min.5-Max. 8)

Abstract

Misshaped red blood cells (RBCs), characterized by thorn-like protrusions known as acanthocytes, are
a key diagnostic feature in Chorea-Acanthocytosis (ChAc), a rare neurodegenerative disorder. The
altered RBC morphology likely influences their biomechanical properties which are crucial for the
cells to pass the microvasculature. Here, we investigated blood cell deformability of 5 ChAc patients
compared to healthy controls during up to one-year individual off-label treatment with the tyrosine
kinases inhibitor dasatinib or several weeks with lithium. Measurements with two microfluidic
techniques allowed us to assess RBC deformability under different shear stresses. Furthermore, we
characterized leukocyte stiffness at high shear stresses. The results show that blood cell deformability
– including both RBCs and leukocytes - in general is altered in ChAc patients compared to healthy
donors. Therefore, this study shows for the first time an impairment of leukocyte properties in ChAc.
During treatment with dasatinib or lithium, we observe alterations in RBC deformability and a stiffness
increase for leukocytes. The hematological phenotype of ChAc patients hints at a reorganization of the
cytoskeleton in blood cells which partly explains the altered mechanical properties observed here.
These findings highlight the need for a systematic assessment of the contribution of impaired blood
cell mechanics to the clinical manifestation of ChAc.

1   Introduction

Chorea-acanthocytosis/VPS13A disease (ChAc) is a rare monogenetic neurodegenerative disease of
the young adulthood affecting multiple systems other than the central nervous system, e.g., the red
blood cells (RBCs) (1–3). ChAc is characterized by various movement disorders (due to a degeneration
mainly of striatal neurons), epilepsy, cognitive decline and misshaped spiky RBCs, the latter being
referred to as acanthocytes (1,2,4,5). The autosomal-recessive condition is caused by loss of function
mutations in the *vacuolar protein sorting 13 homolog A* (*VPS13A*) gene (6–10). Therefore, the term
"VPS13A disease" has been recommended to replace the historical, more descriptive terminology (11).
A disease-modifying therapy has not been established so far.

It is still unclear, how exactly the loss of function of VPS13A leads to the manifestation of the cellular
and clinical phenotype. VPS13A localizes at different membrane contact sites and is most probably
involved in non-vesicular lipid transport as it is assumed also for the other members of the VPS13
protein family (for review, see Leonzino, Reinisch and De Camilli, 2021) (12–15). Two downstream
mechanisms are considered to be important drivers of ChAc pathophysiology: decreased
phosphoinositide-3-kinase (PI3K) signaling and increased activity of Src family tyrosine kinase Lyn
(for review, see Peikert et al. 2018 (2)).

Active Lyn kinase accumulates in ChAc RBCs and hyperphosphorylates membrane proteins such as
band 3 protein. Since band 3 is a structural protein, linking the cytoskeleton to the plasma membrane,
it is likely to be causally involved in the genesis of the spiky morphology. Accumulation of active Lyn
kinase was also found to be related to impaired autophagic flux (16,17). These phenotypes could be
reversed by *in vitro* treatment with tyrosine kinase inhibitors (TKI) such as the Src kinase family
inhibitor dasatinib (16). In a Vps13a knockout mouse model, the more specific Lyn kinase inhibitor
nilotinib improved both the hematological and neurological phenotypes by improving autophagy and
preventing neuroinflammation (18). In their recent work, Peikert et al., 2021 (19) report on single
individual treatment approaches targeting Lyn kinase with dasatinib in ChAc patients. They showed
that initially reduced F-actin signal, increased osmotic fragility and impaired autophagy were partially
restored in ChAc RBCs.

Furthermore, disturbed signaling via Phosphoinositide-3-kinase (PI3K) has been also identified as
relevant in the pathophysiology of ChAc patients. Reduced signaling via the PI3K-Rac1-PAK pathway
was reported to lead to disordered actin polymerization (20). Additionally, in other cells, altered PI3K
signaling caused a decreased ORAI1 expression and store-operated Ca^{2+} entry (SOCE), subsequently
disturbed Ca^{2+} homeostasis and apoptosis which could be partly reversed by *in vitro* lithium treatment
(21–23).

The rather late onset of the disease at the age of early adulthood and the slow progression rate (1,3,24)
lead to the assumption of a dependency between the altered RBC properties and neuronal degeneration:
Changes in the mechanical properties of the RBCs and hence reduced passage in micro-vessels and
capillaries may impair oxygen supply in certain areas of the brain that could over time accumulate to
tissue alterations leading to the symptoms described (25).

Altered RBC morphology is often linked to a change in cell mechanics which may lead to impaired
blood flow (26,27). Until now, the effect of acanthocytes in ChAc patients on the pathophysiology of
the disease remains elusive. Here, we describe, for the first time, altered deformability of blood cells
from ChAc patients including white blood cells. In a second step, we observed these parameters during
individual off-label treatments with dasatinib or lithium, each targeting one of the signaling pathways
believed to be involved in the genesis of the RBC phenotype.

Deformability was measured using two techniques that asses the response to different ranges of shear
stresses acting on the cells: shape analysis of RBCs at low shear stress (0.1-3 Pa) (28) and real-time
fluorescence and deformability cytometry (RT-FDC) (29,30) at high shear stress (ca. 100 Pa). The first
only assesses the deformability of RBCs while RT-FDC was also used to characterize the mechanics
of leukocytes. We demonstrate that the deformability of all blood cells is affected in ChAc patients and
this is further modulated by the treatments. This highlights that monitoring blood cell mechanical
properties of ChAc patients during the course of disease and possible treatments can increase our
understanding of this disease.

2   Materials and Methods

2.1   Cell source and reagents

We included 5 ChAc patients in this study for whom diagnosis was confirmed by Western blot
(absence of chorein/VPS13A protein) and genetic testing (31). The clinical parameters of the patients
are listed in Table S1 and for P1-P3 further information can be found in Peikert et al., 2021 (19).
Patients and healthy control blood donors were enrolled in ongoing studies on the pathogenesis and
natural history of neurodegenerative diseases approved by the institutional review board of the
Technische Universität Dresden, Germany (EK 45022009, EK 78022015). The ChAc patients were
treated with dasatinib (P1-P3) or lithium (P4-P5) in the context of an individual off-label therapy based
on the above described preclinical evidence. Standard dose of 100 mg dasatinib per day was
administered orally, as it was lithium for which serum lithium concentration target range was defined
as 0.6-0.8 mmol/l. Treatment started after the blood for the baseline measurements was taken.

2.2   RBC shape analysis at low shear stress

The distribution of shapes for RBCs flowing through a narrow capillary at a fixed flow condition is
directly linked to the distribution of shear moduli within the population. Thus, changes in the shape
distribution are directly linked to a change of cell mechanics in the sample. This approach and the
experimental design are described in detail in Reichel et al., 2019 (28). Here, cells flow through a

roughly 5 mm long measurement channel that is 10-15 µm wide, and their shapes and dynamics are
tracked at the end of the channel over a length of 435 µm. Cells were recorded with the same setup
used for real-time deformability cytometry (29) using a different microfluidic chip. Examples of cells
flowing through the region of interest are shown in figure 1A,B and in supplementary videos S1-S8.

The concepts introduced in Reichel et al., 2019 (28) cannot directly be used for samples containing
acanthocytes because their shape at rest differs from that of healthy RBCs. Further, acanthocytes show
shapes not observed for healthy cells (discocytes or normocytes) when deformed by the flow in the
channel. For this study, cells flowing through a square channel with a width of 10 µm at cell velocities
ranging up to 3 mm/s were characterized by their membrane morphology as normal looking (referred
to as normocytes) or as acanthocytes if they showed thorn-like protrusions. Furthermore, cells were
classified as either deformed by the flow or still maintaining their resting shape (undeformed). This
classification was done by eye. A change of the cell mechanical properties should manifest in a change
of the fraction of deformed cells in the sample and that the velocity at which the majority of RBCs get
deformed is shifted to higher values.

For examples see figure 1A,B. The classification was done by eye from the recorded videos of cells
passing the region of interest. This resulted in a curve with probabilities to find cells in a deformed
state for a given range of velocities. The resulting curve for healthy RBCs is given in fig. 2A (data
from Reichel et al., 2019 (28)). To characterize differences to ChAc patient cells and changes on the
cells during treatment, a logistic growth function $A/(1 + e^{-k(v-v_0)})$, with cell velocity v, was fitted to
the data and the fit parameters A, v_0, and k were used to make statements of the progression of the
curves during treatment (see fig 2A).

2.3   Real-time fluorescence and deformability cytometry (RT-FDC)

The setup of real-time fluorescence and deformability cytometry (RT-FDC) is described in detail in
Otto et al., 2015 (29) and Rosendahl et al., 2018 (30). In brief, flow is introduced into a microfluidic
chip with the help of syringe pumps and cells get deformed by hydrodynamic stresses as they pass a
narrow constriction, still larger than their own size. The chip is mounted on an inverted microscope
(Axiovert 200M, ZEISS, Oberkochen, Germany) and images are obtained at the end of the constriction
with a CMOS camera (Mikrotron, Unterschleissheim, Germany). Camera and syringe pump are
controlled via the measuring software ShapeIn (Zellmechanik Dresden, Dresden, Germany). ShapeIn
analyzes the recorded images in real-time and computes a contour and projected area for each cell. The
deformation parameter used in RT-FDC is defined as 1-circularity and is calculated from the contour's
area and perimeter using: $\text{deformation} = 1 - 2 \cdot \sqrt{\pi \cdot \text{Area}}/\text{perimeter}$, illustrated on the example of
a deformed RBC in figure 1C. As introduced by Toepfner et al., 2018 (32), the data from RT-FDC
experiments can be used to distinguish between different types of blood cells. These strategies were
used here to selectively study the deformation and mechanics of RBCs, lymphocytes, and myelocytes
from ChAc patients (see also SI text).

Because the stresses required to deform RBCs are smaller than for leukocytes, both were measured
under different conditions. For measurements on leukocytes, citrate blood was resuspended in a
viscosity-adjusted measurement buffer (phosphate saline buffer without Mg^{2+} and Ca^{2+} (PBS−)
containing 0.5% (w/v) methyl cellulose (4000 cPs, Alfa Aesar 036718.22, CAS# 9004-67-5); adjusted
in HAAKE Falling Ball Viscometer type C (Thermo Fisher Scientific, Dreieich, Germany) using ball
number 3 to a viscosity of 15 mPa s) at a ratio of 1:20. The increased viscosity of the buffer also
increases the stresses acting on the cells inside the channel. Measurements were performed in channels
with 20×20 µm cross-section at a flow rate of 0.08 µl/s. Example images of a deformed lymphocyte

and myelocyte are shown in figure 1D and E, respectively. To decouple effects of the cell size on the
deformation, the Young's modulus was computed for leukocytes as described in Mokbel et al., 2017
(33). Since the derivation of Young's moduli for this system is only valid for initially spherical cells
(33,34), it cannot be calculated for RBCs.

RBC measurements were performed on an RT-FDC setup with a fluorescence module described in
detail in Rosendahl et al., 2018 (30). Whole blood was resuspended in the same measurement buffer
described above at a ratio of 1:200. The buffer was complemented with 2.5 µM syto13 nucleic acid
stain (Thermo Fisher Scientific, Dreieich, Germany) to mark for the reticulocytes in the sample.
Fluorescence was excited with a 60 mW, 488 nm laser at 8 % power (OBIS 488-nm LS 60 mW,
Coherent Deutschland) and the signal was measured with an avalanche photodiode (MiniSM10035;
SensL Corporate, Cork, Ireland). Measurements were performed in channels with a 20×20 µm cross-
section at a flow rate of 0.02 µl/s.

2.4 Data analysis and data availability

Data was analyzed and plotted using custom python scripts. For details, see SI text.

The raw experiment data is available in the deformability cytometry open repository, DCOR, under
https://dcor.mpl.mpg.de. Scripts used to analyze the data are available in the supplementary material
or by request to the authors.

3 Results

3.1 Red blood cell deformability in Chorea-Acanthocytosis

To investigate differences in the deformability of red blood cells (RBCs) from chorea-acanthocytosis
(ChAc) patients compared to that of healthy donors, we performed shape-analysis and real-time
fluorescence and deformability cytometry (RT-FDC) measurements on the blood of 5 patients (P1-P5)
and compared it with measurements of healthy controls. The results are depicted in figure 1A,B and
1F,G. Every velocity bin of 0.25 mm/s in figure 1A and B includes approx. 100 cells for each patient.
The shape-probability curves of normocytes (fig. 1A) and acanthocytes (fig. 1B) show that RBCs from
ChAc patients are less likely to deform in channel flow. It should be mentioned that also control
samples can include acanthocyte-like cells but to a much lesser extent than in patient samples.
Acanthocytes are clearly less likely to deform due to flow even for higher flow velocities.

The RT-FDC measurements show that the RBCs of ChAc patients have a slightly larger projected area
than those from healthy donors (fig. 1F, effect size of +1.89 µm^2, p=0.0036 computed with linear mixed
effect models (35)). No significant difference between control and ChAc RBCs' deformation could be
detected (p=0.1382, fig. 1G).

Figure 1: Blood cell deformability of ChAc patients compared to healthy control blood. **(A)** Example shapes of normal, discocyte RBCs (normocytes) in an **(i)** undeformed state and **(ii)** deformed by flow during low shear stress flow in a 10 µm channel. The diagram shows the probability to find normocyte cells from the sample in a deformed state at a given cell velocity in the channel for the acanthocyte patients and the mean curve from 3 control measurements. Error bars represent SEM. Dotted line shows a logistic-growth function fit to the control data (see fig. 2A) **(B)** Example shapes of acanthocytes in an **(i)** undeformed state and **(ii)** deformed by flow. Below is the shape probability diagram for the acanthocytes and the control curves. **(C)** Example shape of an RBC in RT-FDC and representation how the deformation parameter is computed. All scale bars represent 10 µm. **(D)** Example image of a lymphocyte in RT-FDC. **(E)** Example image of a myelocyte. **(F), (G)** Projected area within the contour and deformation of RBCs of the ChAc patients from RT-FDC measurements vs. pooled data from 10 control measurements (full data in fig. S4). White dots represent the median value; grey box in the violin shows inter-quartile range (IQR) and extended lines 1.5×IQR. Dashed grey line shows the median control value. P-values were calculated with linear-mixed effect models as described in Herbig et al., 2018 (35). **(H)** Young's modulus of lymphocytes from ChAc patients vs. control. **(I)** Young's modulus of myelocytes from ChAc patients vs. control.

3.2 White blood cell deformability in Chorea-Acanthocytosis

In RT-FDC measurements on whole blood it is possible to distinguish different cell types (see materials and methods, SI text and Töpfner et al., 2018 (32)). Thus, we investigated mechanical differences also between the leukocytes of ChAc patients and healthy donors. We distinguish, mainly by size, between all lymphocytes and cells resulting from the myeloid lineage (myelocytes), comprising all mono- and granulocytes. The myelocyte fraction mainly consists of neutrophil granulocytes (> 80%).

Since white blood cells are spherical at rest, we can employ a model that correlates the projected area and the deformation to an apparent Young's modulus (33), to see differences in stiffness before, during and after the treatments. It can be seen (fig. 1H,I) that both lymphocytes and myelocytes of ChAc patients are significantly stiffer than their healthy counterparts before treatment onset (lymphocytes: +240 Pa, p=0.0186, myelocytes: +150 Pa, p=0.0070; effect sizes and p-values determined by linear mixed effects model analysis (35)).

3.3 Dasatinib and lithium treatment effect on RBC deformability

3.3.1 RBC shape analysis during dasatinib or lithium treatment

To study the effect of dasatinib or lithium on RBC deformability of ChAc patients, we monitored RBCs during treatment with shape analysis measurements and RT-FDC.

To compare shape-analysis data over the treatment time, a logistic-growth function was fitted to the data (see fig. 2A) and the fitting parameters were used to detect changes in the deformability during the treatment time. The results for the dasatinib treatment are shown in figure 2B-D. The lithium results are depicted in figure 3. The plateauing probability A (fig. 2B) for normocytes and acanthocytes increases slightly during treatment with dasatinib and drops again after treatment. The parameter estimates the probability of deformed cells at very high velocities ($v \to \infty$) and is close to 100% for control blood, meaning nearly all RBCs should deform at high velocities. This indicates that RBCs become more deformable during dasatinib treatment. The results for the other fitting parameters do not show a clear trend. The results pooled for all cells are shown in figure S1 and the curves from all measurements are given in figure S2.

The acanthocyte count over the treatment is given in figure 2E and shows that the fraction of acanthocytes remains constant within a certain range of fluctuation without a systematic trend for both treatments, which is in line with the clinical data (19).

To check if dasatinib has a short-term effect on RBC deformability, RBCs from P3 were treated with dasatinib *in vitro* and the shapes in channel flow were analyzed (for details, see SI text). The results are shown in figure 2F. There was no significant difference in the deformability between control- and dasatinib-treated RBCs. This indicates that the observed effects on cell deformability are likely caused by long-term treatment effects, e.g., acting on erythropoiesis or indirect systemic effects.

During treatment with lithium, we also observed an apparent increase of the plateauing probability with and even after treatment time for normocytes and acanthocytes (fig. 3A). Transition velocity (fig. 3B) and growth rate (fig. 3C) both decreased with treatment time, but the scarcity of the datapoints makes it difficult to draw any conclusions from these results.

Figure 2: Dasatinib treatment effect on RBC deformability. (A) Illustration of the fitting parameters used to characterize the shape probability curves presented on the control data. **(B)** Fitted plateauing probability A of the shape probability curves from ChAc patients as a function of treatment time with dasatinib for normocytes and acanthocytes respectively. Dashed lines represent datapoints taken after treatment was stopped. Dashed grey lines show the respective value from healthy donors without treatment. **(C)** Fitted transition velocity v_0 for the probability to find deformed cells over treatment time. **(D)** Fitted growth rate k of the probability over treatment time. **(E)** Percentage of acanthocytes found in each sample during shape analysis measurements for patients P1-P5. Dashed lines indicate datapoints taken after the respective treatment already stopped. **(F)** Fraction of healthy shapes (hs) and acanthocytes (ac) over cell velocity for untreated and *in-vitro* dasatinib treated RBCs from patient P3.

Figure 3: Lithium treatment effect on RBC deformability. (one column figure) **(A)** Fitted plateauing probability A of the shape probability curves from ChAc patients as a function of treatment time with lithium for normocytes and acanthocytes respectively. Dashed lines represent datapoints taken after treatment was stopped. Dashed grey lines show the respective value from healthy donors without treatment. **(B)** Fitted transition velocity v_0 for the probability to find deformed cells over treatment time. **(C)** Fitted growth rate k of the probability over treatment time.

3.3.2 RBC deformation in RT-FDC during dasatinib or lithium treatment

RT-FDC results for the dasatinib treatment are shown in figure 4A-C and for the lithium treatment in
figure 4D-F. It can be seen that during treatment with dasatinib, the RBC deformation parameter drops
for all patients between 50-100 days after treatment start, while the projected area remains mainly
constant with the exception of some outliers.

For the lithium treatment, the RT-FDC results show that the deformation parameter decreases with
treatment time which is accompanied by an increase in cell size. After the treatment, the deformation
and area values return almost to the control values for P5 while the values for P4 do not fully recover.

The interpretation of deformation values for RBCs in RT-FDC cannot directly be linked to cell
mechanics because they undergo a non-trivial shape transition from discocyte to tear-drop shape (28).
A decrease of the deformation value does not necessarily mean that the cells got stiffer. To back this
up, we artificially stiffened RBCs from healthy donors with glutaraldehyde (36,37) or diamide (38)
(see SI text and fig. S3) and saw that an increasing concentration of the reagents led to an increase of
the deformation when working at low concentrations. At high concentrations the cell shape gets fixed
which leads to lower deformation values.

Since reticulocytes have been reported to have a higher projected area and lower deformation
compared to the whole RBC population (30), we speculated that above mentioned results might be due
to increased levels of reticulocytes in ChAc patients. The results for reticulocytes, labelled with syto13
nucleic acid stain, together with all RBCs are given in figure S4A-D. The patients' reticulocytes indeed

had higher projected area and lower deformation if compared to the whole RBC population which did
not differ from healthy controls. Qualitatively, reticulocyte area and deformation change over treatment
time like the total RBC population. The fraction of reticulocytes in the blood is initially higher
compared to controls and slightly decreases during the dasatinib treatment closer to values from healthy
donors (fig. 4C). A reduction of the reticulocyte count with dasatinib treatment was also reported for
VPS13a$^{-/-}$ mice phenocopying human ChAc (18). During lithium treatment P4 showed a decrease
during and after treatment, while P5 showed a slight increase but values were mostly in the range
expected for healthy individuals (0.5-2.5%).

Figure 4: RBC deformation, size and reticulocyte count in RT-FDC during treatment with Dasatinib or
lithium. (A-C) RBC median deformation, projected area and reticulocyte fraction measured with RT-FDC
before, during and after treatment with dasatinib. **(D-F)** RBC median deformation, projected area and
reticulocyte fraction measured with RT-FDC before, during and after treatment with lithium. Dashed lines
indicate time points after the treatment was stopped, dashed gray lines indicate control values without treatment.
The red region in the reticulocyte plots indicates the range for healthy individuals reported in the literature of
0.5-2.5%.

3.4   ChAc Leukocytes stiffen during treatment with dasatinib

The Young's moduli of lymphocytes and myelocytes in dasatinib treatment are shown in figure 5A,B.
The lithium data is shown in figure 5C,D. During treatment with dasatinib, we observe a further
increase of the Young's modulus for all three patients which also persists for the lymphocytes shortly
after the treatment stopped. For myelocytes, the median Young's modulus decreases for patients P1
and P2 shortly after the treatment but keeps increasing for P3.

During treatment with lithium both patients show an increase in lymphocyte and myelocyte stiffness.
After the treatment stopped, we observe a further increase for patient P4 and a drop back to the pre-
treatment level for patient P5 approximately one year after the treatment stopped. The area and
deformation data used to compute the Young's modulus is given in figure S5C,D.

Figure 5: Leukocyte stiffness in RT-FDC during treatment with Dasatinib or lithium. (one column figure) **(A)** Median Young's modulus of ChAc patients' lymphocytes during dasatinib treatment. **(B)** Median Young's modulus of ChAc patients' myelocytes during dasatinib treatment. **(C)** Median Young's modulus of ChAc patients' lymphocytes during lithium treatment. **(D)** Median Young's modulus of ChAc patients' myelocytes during lithium treatment.

4   Discussion

Our study highlights that the deformability, not only of RBCs, but of all blood cell types was altered in ChAc patients. Shape analysis results hinted at a decreased deformability of ChAc RBCs compared to controls. The decreased deformability can be directly inferred from the reduced number of deformed cells at any cell velocity. A reduced deformability of ChAc RBCs, deduced from cell shapes, has recently also been reported by Rabe et al., 2021 (39).

In RT-FDC, we did not observe a change in the deformation parameter but ChAc RBCs had a bigger projected area. Because deformation and size are always correlated in RT-FDC (34), this is also an indicator for a change of RBC deformability, but the interpretation of this behavior is not straight forward. To see how the deformation parameter changes when RBC stiffness is controllably changed, we treated RBCs *in vitro* with glutaraldehyde or diamide. At low concentrations, we observed an increase of the deformation parameter which shows that higher deformation parameter can also occur for stiffer RBCs.

While we observed bigger RBCs in ChAc patients, values for the mean corpuscular volume (MCV) reported in Peikert et al., 2021 (19) did not show an increased RBC volume for patients P1-P3. Sizes reported in RT-FDC measurements are based on the projected cross-sectional area of the deformed cells. Simulations of the flow fields inside the channel suggest that the stresses near the centerline can be approximated as axisymmetric (40). Therefore, it is fair to assume that the RBCs take up an axisymmetric shape as well and the projected area is directly correlated to cell size. In automated cell counters, the MCV is determined, e.g., by impedance or light scattering measurements which require a constant deformability of RBCs for a reliable result. So, MCV is not a direct measurement of single cell sizes. Here, we provide a direct observation of sizes based on single cell imaging.

Shape analysis and RT-FDC both indicated a change of RBC deformability during treatment with dasatinib. Shape analysis experiments showed an increase of the plateau probability parameter with treatment time, which showed that more deformed cells are observed at high cell velocities, hinting at an increased deformability of both normocytes and acanthocytes. In RT-FDC, we observed a sharp drop of the deformation parameter after 50-100 treatment days while the size fluctuated around a constant value. Again, for RBCs a direct link between the deformation parameter in RT-FDC and cell mechanics is not trivial and a decrease in deformation does not necessarily mean that the cells got

stiffer (see fig. S3). Even though we do not have certainty of the direction of the change, we claim that
RBC mechanics was altered during dasatinib treatment.

The sharp drop in deformation for the RBCs occurred at different timepoints for P3 than for P1 and
P2. A possible correlation to this can be found in the levels of Lyn kinase and the autophagy marker
p62 for the patients reported in Peikert et al., 2021 (19). Lyn kinase levels for P3 dropped directly after
treatment start and stayed on this level during the treatment while the levels for P1 and P2 had a sudden
rise after 10 weeks of treatment. On the other hand, the levels of p62 stayed rather constant for P1 and
P2 during the treatment while they were rising above control level for P3 after 10 weeks. Patient
incompliance during treatment could be a possible explanation for these effects.

During lithium treatment, we also observed an increasing plateauing probability in the shape analysis
accompanied with a decrease of the transition velocity, both indicators for increased RBC
deformability. In RT-FDC we saw a decrease of deformation accompanied by an increase in projected
area. An increase in projected area and decrease in deformation hints at a volume increase and a
reduced surface area-to-volume ratio, which leads to an increased sphericity or effective rounding of
the cells. Rounder shapes lead to smaller deformation values, assuming that this effect dominates any
stiffness changes of the cells. All in all, these findings indicate that lithium affects RBC properties in
ChAc patients but because of the scarcity of the data here, one should be careful with the interpretation.

Data from Peikert et al., 2021 indicated that with 12asatinib treatment, F-actin became more localized
at the cortex (19). These modifications in the actin network likely contribute to changes in RBC
deformability (41) that we observed here.

Treatment with dasatinib led to a decrease of the reticulocyte fraction in the blood of ChAc patients.
The same effect was observed in a mouse model phenocopying human ChAc (18). Peikert et al., 2021
report an abundance of the autophagy initiator Ulk1 in ChAc patients RBCs, which decreased with
dasatinib treatment (19). Ulk1 is connected to autophagy in erythropoiesis and could influence the rate
of reticulocyte production (42), providing a mechanistic link to the reticulocyte count.

While an impaired RBC deformability was described before for patients with other forms of
neuroacanthocytosis (43) this is, to our knowledge, the first time that also patients' leukocyte stiffness
was studied. Lymphocytes and myelocytes were both stiffer in ChAc patients compared to cells from
healthy donors. This is of interest as VPS13A, the gene which is known to have a loss-of-function
mutation in ChAc, is especially expressed in human monocytes and B-, and helper T-cells (44). Since
VPS13A is known to cause structural reorganizations of the cytoskeleton from studies on other cell
types (17,20,45), it is likely that we observed these as altered mechanical properties in leukocytes.
Another feature in ChAc blood is an increased expression of Lyn kinase, which is known to set the
threshold for B-cell activation. Mechanical changes of lymphocytes after activation have been reported
before (32,46). Another mechanism that could lead to altered leukocyte mechanics is an increase in
interleukin-1 beta (IL-1b) that was observed in the brain of Vps13a knockout mice (18). IL-1b not only
plays a key role in autoinflammation but is also known to cause re-organization of the actin
cytoskeleton in different cell types (47,48).

Young's moduli of both lymphocytes and myelocytes increased even further away from control values
during treatment with dasatinib or lithium. A stiffening was previously also observed for leukemic cells
during treatment with different chemotherapeutic agents for acute lymphoblastic leukemia
(dexamethasone and daunorubicin), such as dasatinib (49). Dasatinib interferes with all the pathways
described above and likely induces further mechanical changes in leukocytes. The inhibited activity of

Lyn during dasatinib treatment could cause a change in lymphocyte activation levels leading to altered
mechanics but the exact mechanisms for the observed leukocyte stiffening with dasatinib treatment
remain rather speculative at this point.

Overall, our results show that the mechanical phenotype of all blood cell types in ChAc patients were
impaired but did not change towards regimes of healthy donors during the treatments. This
demonstrates that the techniques that were utilized here are suited to measure a treatment response.
Clinical data in Peikert et al., 2021 (19) hints on slightly improved hematological features but these did
not manifest in an improved neurological phenotype.

In general, decreased blood cell deformability often leads to clinical symptoms due to decreased cell
survival and secondary pathology (25–27,50). Especially, an impaired oxygen transport by altered
blood cell deformability should be considered in neurological diseases (51). While clinical descriptions
of ChAc patients focus on neurological symptoms, we highlight in this study that the effects on blood
deformability should not be neglected and could contribute to the clinical manifestation (through the
mechanisms stated above) and thus methods that are able to monitor this (as shown here) should be
considered as read outs for clinical trials. Our results on the impaired mechanics of leucocytes
especially highlights that the focus of ChAc research should not only be on RBCs but also include
other blood cell types.

5   Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial
relationships that could be construed as a potential conflict of interest.

6   Author Contributions

Conceptualization: F.R., M.K., K.P., G.B., A.H., and J.G.; Experiments–shape analysis: F.R. and
A.R.P., Experiments–RT-FDC: F.R., P.R., M.H., and M.K., Experiments–in-vitro shape analysis:
A.K.; Formal analysis: F.R., P.R., A.K., and M.K.; Supervision: L.K.,A.H., and J.G.; Writing–
original draft, K.P. and F.R..; Writing–review and editing, F.R., M.K., K.P., G.B., L.K., A.H., and
J.G.; All authors have read and agreed to the published version of the manuscript.

7   Funding

K.P. was supported by the Else Kröner clinician scientist program and the MeDDrive program (TU
Dresden, Germany), as well as by the Rostock Academy for Clinician Scientists (RACS) and the
FORUN program (University of Rostock, Germany). A.H. was supported by the "Hermann und Lilly
Schilling-Stiftung für medizinische Forschung im Stifterverband". Expenses for the off-label
dasatinib prescription were gratefully covered in part by the Center for Regenerative Therapies
Dresden (CRTD, TU Dresden, Germany) and the public healthcare insurance of the patients.

8   Acknowledgments

The authors thank all patients and their families for giving consent for publication of the data and the
healthy control subjects for participation in this study. We are especially grateful to Glenn (†) and

Ginger Irvine as the founders of the Advocacy for Neuroacanthocytosis Patients
(www.naadvocacy.org) and to Susan Wagner and Joy Willard-Williford as representatives of the NA
Advocacy USA (www.naadvocacyusa.org).

9   Supplementary Material

447 Supplementary Information, Figures, Table and Videos will be available online.

10   Data Availability statement

450 The raw experiment data is available in the deformability cytometry open repository, DCOR, under
451 https://dcor.mpl.mpg.de. Scripts used to analyze the data are available in the supplementary material
452 or by request to the authors.

11   References

1.   Jung HH, Danek A, Walker RH. Neuroacanthocytosis Syndromes. Orphanet J Rare Dis.
(2011) 6(1):68. doi: 10.1186/1750-1172-6-68

2.   Peikert K, Danek A, Hermann A. Current state of knowledge in Chorea-Acanthocytosis as
core Neuroacanthocytosis syndrome. Eur J Med Genet. (2018) 61(11):699–705. doi:
10.1016/j.ejmg.2017.12.007

3.   Walker RH. Chorea-Acanthocytosis. In: Reference Module in Neuroscience and Biobehavioral
Psychology. Elsevier; (2017.). doi: 10.1016/B978-0-12-809324-5.00521-6

4.   Levine IM, Estes JW, Looney JM. Hereditary Neurological Disease With Acanthocytosis: A
New Syndrome. Arch Neurol. (1968) 19(4):403–9. doi:
10.1001/archneur.1968.00480040069007

5.   Critchley EMR, Clark DB, Wikler A. Acanthocytosis Without. Arch Neurol. (1968)
18(2):134–40. doi: 10.1001/archneur.1968.00470320036004

6.   Danek A, Bader B, Velayos-Baeza A, Walker RH. Autosomal recessive transmission of
chorea-acanthocytosis confirmed. Acta Neuropathol. (2012) 123(6):905–6. doi:
10.1007/s00401-012-0971-y

7.   Dobson-Stone C, Danek A, Rampoldi L, Hardie R, Chalmers R, Wood N, et al. Mutational
spectrum of the CHAC gene in patients with chorea-acanthocytosis. Eur J Hum Genet. (2002)
10(11):773–81. doi: 10.1038/sj.ejhg.5200866

8.   Rampoldi L, Dobson-Stone C, Rubio JP, Danek A, Chalmers RM, Wood NW, et al. A
conserved sorting-associated protein is mutant in chorea-acanthocytosis. Nat Genet. (2001)
28(2):119–20. doi: 10.1038/88821

9.   Ueno S, Maruki Y, Nakamura M, Tomemori Y, Kamae K, Tanabe H, et al. The gene encoding
a newly discovered protein, chorein, is mutated in chorea-acanthocytosis. Nat Genet. (2001)
28(2):121–2. doi: 10.1038/88825

10.   Nishida Y, Nakamura M, Urata Y, Kasamo K, Hiwatashi H, Yokoyama I, et al. Novel
pathogenic VPS13A gene mutations in Japanese patients with chorea-acanthocytosis. Neurol
Genet. (2019) 5(3):e332. doi: 10.1212/NXG.0000000000000332

11.   Walker RH, Danek A. "Neuroacanthocytosis" – Overdue for a Taxonomic Update. Tremor
and Other Hyperkinetic Movements. (2021) 11(1):1–6. doi: 10.5334/tohm.583

12.   Leonzino M, Reinisch KM, De Camilli P. Insights into VPS13 properties and function reveal a
new mechanism of eukaryotic lipid transport. Biochim Biophys Acta - Mol Cell Biol Lipids.
(2021) 1866(10):159003. doi: 10.1016/j.bbalip.2021.159003

13.   Gao M, Yang H. VPS13: A lipid transfer protein making contacts at multiple cellular
locations. J Cell Biol. (2018) 217(10):3322–4. doi: 10.1083/jcb.201808151

14.   Kumar N, Leonzino M, Hancock-Cerutti W, Horenkamp FA, Li P, Lees JA, et al. VPS13A
and VPS13C are lipid transport proteins differentially localized at ER contact sites. J Cell Biol.
(2018) 217(10):3625–39. doi: 10.1083/jcb.201807019

15.   Yeshaw WM, van der Zwaag M, Pinto F, Lahaye LL, Faber AIE, Gómez-Sánchez R, et al.
Human VPS13A is associated with multiple organelles and influences mitochondrial
morphology and lipid droplet motility. Elife. (2019) 8. doi: 10.7554/eLife.43561

16.   Lupo F, Tibaldi E, Matte A, Sharma AK, Brunati AM, Alper SL, et al. A new molecular link
between defective autophagy and erythroid abnormalities in chorea-acanthocytosis. Blood.
(2016) 128(25):2976–87. doi: 10.1182/blood-2016-07-727321

17.   De Franceschi L, Tomelleri C, Matte A, Brunati AM, Bovee-Geurts PH, Bertoldi M, et al.
Erythrocyte membrane changes of chorea-acanthocytosis are the result of altered Lyn kinase
activity. Blood. (2011) 118(20):5652–63. doi: 10.1182/blood-2011-05-355339

18.   Peikert K, Federti E, Matte A, Constantin G, Pietronigro EC, Fabene PF, et al. Therapeutic
targeting of Lyn kinase to treat chorea-acanthocytosis. Acta Neuropathol Commun. (2021)
9(1):81. doi: 10.1186/s40478-021-01181-y

19.   Peikert K, Glaß H, Federti E, Matte A, Pelzl L, Akgün K, et al. Targeting Lyn Kinase in
Chorea-Acanthocytosis: A Translational Treatment Approach in a Rare Disease. J Pers Med.
(2021) 11(5):392. doi: 10.3390/jpm11050392

20.   Föller M, Hermann A, Gu S, Alesutan I, Qadri SM, Borst O, et al. Chorein-sensitive
polymerization of cortical actin and suicidal cell death in chorea-acanthocytosis. FASEB J.
(2012) 26(4):1526–34. doi: 10.1096/fj.11-198317

21.   Pelzl L, Elsir B, Sahu I, Bissinger R, Singh Y, Sukkar B, et al. Lithium Sensitivity of Store
Operated Ca2+ Entry and Survival of Fibroblasts Isolated from Chorea-Acanthocytosis
Patients. Cell Physiol Biochem. (2017) 42(5):2066–77. doi: 10.1159/000479901

22.   Pelzl L, Hauser S, Elsir B, Sukkar B, Sahu I, Singh Y, et al. Lithium Sensitive ORAI1
Expression, Store Operated Ca2+ Entry and Suicidal Death of Neurons in Chorea-
Acanthocytosis. Sci Rep. (2017) 7(1):6457. doi: 10.1038/s41598-017-06451-1

23.   Yu W, Honisch S, Schmidt S, Yan J, Schmid E, Alkahtani S, et al. Chorein Sensitive Orai1
Expression and Store Operated Ca2+ Entry in Rhabdomyosarcoma Cells. Cell Physiol
Biochem. (2016) 40(5):1141–52. doi: 10.1159/000453168

24.   Rampoldi L, Danek A, Monaco AP. Clinical features and molecular bases of
neuroacanthocytosis. J Mol Med. (2002) 80(8):475–91. doi: 10.1007/s00109-002-0349-z

25.   Adjobo-Hermans MJW, Cluitmans JCA, Bosman GJCGM. Neuroacanthocytosis:
Observations, Theories and Perspectives on the Origin and Significance of Acanthocytes.
Tremor and Other Hyperkinetic Movements. (2015) 5:328. doi: 10.5334/tohm.271

26.   An X, Mohandas N. Disorders of red cell membrane. Br J Haematol. (2008) 141:367–75. doi:
10.1111/j.1365-2141.2008.07091.x

27.   Da Costa L, Galimand J, Fenneteau O, Mohandas N. Hereditary spherocytosis, elliptocytosis,
and other red cell membrane disorders. Blood Rev. (2013) 27(4):167–78. doi:
10.1016/j.blre.2013.04.003

28.   Reichel F, Mauer J, Nawaz AA, Gompper G, Guck J, Fedosov DA. High-Throughput
Microfluidic Characterization of Erythrocyte Shapes and Mechanical Variability. Biophys J.
(2019) 117(1):14–24. doi: 10.1016/j.bpj.2019.05.022

29.   Otto O, Rosendahl P, Mietke A, Golfier S, Herold C, Klaue D, et al. Real-time deformability
cytometry: on-the-fly cell mechanical phenotyping. Nat Methods. (2015) 12(3):199–202. doi:
10.1038/nmeth.3281

30.   Rosendahl P, Plak K, Jacobi A, Kraeter M, Toepfner N, Otto O, et al. Real-time fluorescence
and deformability cytometry. Nat Methods. (2018) 15(5):355–8. doi: 10.1038/nmeth.4639

31.   Dobson-Stone C, Velayos-Baeza A, Filippone LA, Westbury S, Storch A, Erdmann T, et al.
Chorein detection for the diagnosis of chorea-acanthocytosis. Ann Neurol. (2004) 56(2):299–
302. doi: 10.1002/ana.20200

32.   Toepfner N, Herold C, Otto O, Rosendahl P, Jacobi A, Kräter M, et al. Detection of human
disease conditions by single-cell morpho-rheological phenotyping of blood. Elife. (2018)
7:e29213. doi: 10.7554/eLife.29213

33.   Mokbel M, Mokbel D, Mietke A, Träber N, Girardo S, Otto O, et al. Numerical Simulation of
Real-Time Deformability Cytometry To Extract Cell Mechanical Properties. ACS Biomater
Sci Eng. (2017) 3(11):2962–73. doi: 10.1021/acsbiomaterials.6b00558

34.   Mietke A, Otto O, Girardo S, Rosendahl P, Taubenberger A, Golfier S, et al. Extracting Cell
Stiffness from Real-Time Deformability Cytometry: Theory and Experiment. Biophys J.
(2015) 109(10):2023–36. doi: 10.1016/j.bpj.2015.09.006

35.   Herbig M, Mietke A, Müller P, Otto O. Statistics for real-time deformability cytometry:

Clustering, dimensionality reduction, and significance testing. Biomicrofluidics. (2018)
12(4):042214. doi: 10.1063/1.5027197

36. Holmes D, Whyte G, Bailey J, Vergara-Irigaray N, Ekpenyong A, Guck J, et al. Separation of
blood cells with differing deformability using deterministic lateral displacement. Interface
Focus. (2014) 4(6):20140011. doi: 10.1098/rsfs.2014.0011

37. Forsyth AM, Wan J, Ristenpart WD, Stone HA. The dynamic behavior of chemically
"stiffened" red blood cells in microchannel flows. Microvasc Res. (2010) 80(1):37–43. doi:
10.1016/j.mvr.2010.03.008

38. Sinha A, Chu TTT, Dao M, Chandramohanadas R. Single-cell evaluation of red blood cell bio-
mechanical and nano-structural alterations upon chemically induced oxidative stress. Sci Rep.
(2015) 5(1):9768. doi: 10.1038/srep09768

39. Rabe A, Kihm A, Darras A, Peikert K, Simionato G, Dasanna AK, et al. The Erythrocyte
Sedimentation Rate and Its Relation to Cell Shape and Rigidity of Red Blood Cells from
Chorea-Acanthocytosis Patients in an Off-Label Treatment with Dasatinib. Biomolecules.
(2021) 11(5):727. doi: 10.3390/biom11050727

40. Panhwar MH, Czerwinski F, Dabbiru VAS, Komaragiri Y, Fregin B, Biedenweg D, et al.
High-throughput cell and spheroid mechanics in virtual fluidic channels. Nat Commun. (2020)
11(1):2190. doi: 10.1038/s41467-020-15813-9

41. Gokhin DS, Nowak RB, Khoory JA, Piedra A de la, Ghiran IC, Fowler VM. Dynamic actin
filaments control the mechanical behavior of the human red blood cell membrane. Pollard TD,
editor. Mol Biol Cell. (2015) 26(9):1699–710. doi: 10.1091/mbc.E14-12-1583

42. Kundu M, Lindsten T, Yang C-Y, Wu J, Zhao F, Zhang J, et al. Ulk1 plays a critical role in
the autophagic clearance of mitochondria and ribosomes during reticulocyte maturation.
Blood. (2008) 112(4):1493–502. doi: 10.1182/blood-2008-02-137398

43. Cluitmans JCA, Tomelleri C, Yapici Z, Dinkla S, Bovee-Geurts P, Chokkalingam V, et al.
Abnormal Red Cell Structure and Function in Neuroacanthocytosis. Connes P, editor. PLoS
One. (2015) 10(5):e0125580. doi: 10.1371/journal.pone.0125580

44. Samaras P, Schmidt T, Frejno M, Gessulat S, Reinecke M, Jarzab A, et al. ProteomicsDB: a
multi-omics and multi-organism resource for life science research. Nucleic Acids Res. (2019)
48(D1):D1153–63. doi: 10.1093/nar/gkz974

45. Rzepnikowska W, Flis K, Muñoz-Braceras S, Menezes R, Escalante R, Zoladek T. Yeast and
other lower eukaryotic organisms for studies of Vps13 proteins in health and disease. Traffic.
(2017) 18(11):711–9. doi: 10.1111/tra.12523

46. Zhang X, Kim T-H, Thauland TJ, Li H, Majedi FS, Ly C, et al. Unraveling the
mechanobiology of immune cells. Curr Opin Biotechnol. (2020) 66:236–45. doi:
10.1016/j.copbio.2020.09.004

47. Franco-Barraza J, Valdivia-Silva JE, Zamudio-Meza H, Castillo A, García-Zepeda EA,
Benítez-Bribiesca L, et al. Actin Cytoskeleton Participation in the Onset of IL-1β Induction of

an Invasive Mesenchymal-like Phenotype in Epithelial MCF-7 Cells. Arch Med Res. (2010)
41(3):170–81. doi: 10.1016/j.arcmed.2010.04.010

48.   Lie PPY, Cheng CY, Mruk DD. The biology of interleukin-1: emerging concepts in the
regulation of the actin cytoskeleton and cell junction dynamics. Cell Mol Life Sci. (2012)
69(4):487–500. doi: 10.1007/s00018-011-0760-0

49.   Lam WA, Rosenbluth MJ, Fletcher DA. Chemotherapy exposure increases leukemia cell
stiffness. Blood. (2007) 109(8):3505–8. doi: 10.1182/blood-2006-08-043570

50.   Brown CD, Ghali HS, Zhao Z, Thomas LL, Friedman EA. Association of reduced red blood
cell deformability and diabetic nephropathy. Kidney Int. (2005) 67(1):295–300. doi:
10.1111/j.1523-1755.2005.00082.x

51.   Kikuchi Y, Da Q-W, Fujino T. Variation in Red Blood Cell Deformability and Possible
Consequences for Oxygen Transport to Tissue. Microvasc Res. (1994) 47(2):222–31. doi:
10.1006/mvre.1994.1017

RBC shape analysis at low shear stress (0.1-1 Pa)

RT-FDC (shear stress = 100-400 Pa)

Red blood cells

White blood cells

A
Normocytes

i) ii)

Probability for deformed shape vs Velocity [mm/s]

P1 P2 P3 P4 P5 CTRL CTRL fit

B
Acanthocytes

i) ii)

Probability for deformed shape vs Velocity [mm/s]

P1 P2 P3 P4 P5 CTRL CTRL fit

C

$$\text{Deformation} = 1 - \frac{2 \cdot \sqrt{\pi \cdot \text{Area}}}{\text{Perimeter}}$$

D

E

F

p = 0.0036

Projected area [µm²] vs Patient (CTRL, P1, P2, P3, P4, P5)

G

p = 0.1382

Deformation vs Patient (CTRL, P1, P2, P3, P4, P5)

H
Lymphocytes

p = 0.0186

Young's Modulus [kPa] vs Patient (CTRL, P1, P2, P3, P4, P5)

I
Myelocytes

p = 0.0070

Young's Modulus [kPa] vs Patient (CTRL, P1, P2, P3, P4, P5)

A Probability for deformed cell

Logistic growth function:

$$Prob = \frac{A}{1 + e^{-k(v-v_0)}}$$

A = Plateau probability

v_0 = transition velocity

k = growth rate