Title

Prolonged nicotine exposure reduces aversion to the drug in mice by altering nicotinic transmission in the interpeduncular nucleus.

Authors

Sarah Mondoloni[1], Claire Nguyen[1], Eléonore Vicq[1,2], Joachim Jehl[1,2], Romain Durand-de Cuttoli[1], Nicolas Torquet[1], Stefania Tolu[1], Stéphanie Pons[3], Uwe Maskos[3], Fabio Marti[1,2], Philippe Faure[1,2*] and Alexandre Mourot[1,2*]

Affiliations

1. Sorbonne Université, Inserm, CNRS, Neuroscience Paris Seine - Institut de Biologie Paris Seine (NPS - IBPS), 75005 Paris, France.

2. Brain Plasticity Unit, CNRS, ESPCI Paris, PSL Research University, 75005 Paris, France

3. Institut Pasteur, Unité Neurobiologie intégrative des systèmes cholinergiques, Département de neuroscience, 75724 Paris cedex, France.

* equal contributions

Correspondence to: almourot@gmail.com or phfaure@gmail.com

Keywords

nicotinic acetylcholine receptors, nicotine addiction, two-bottle choice paradigm, drug aversion, medial habenula-interpeduncular nucleus pathway

Abstract

Nicotine intake is likely to result from a balance between the rewarding and aversive

properties of the drug, yet the individual differences in neural activity that control

aversion to nicotine and their adaptation during the addiction process remain largely

unknown. Using a two-bottle choice experiment, we observed a high heterogeneity in

nicotine-drinking profiles in isogenic adult male mice, with about half of the mice

persisting in consuming nicotine even at high concentrations, whereas the other half

stopped consuming. We found that nicotine intake was negatively correlated with

nicotine-evoked currents in the interpeduncular nucleus (IPN), and that prolonged

exposure to nicotine, by weakening this response, decreased aversion to the drug, and

hence boosted consumption. Lastly, using knock-out mice and local gene re-

expression, we identified $\beta 4$-containing nicotinic acetylcholine receptors of IPN

neurons as the molecular and cellular correlates of nicotine aversion. Collectively, our

results identify the IPN as a substrate of individual variabilities and adaptations in

nicotine consumption.

Introduction

Nicotine remains one of the most-widely used addictive substance in the world, and even though cigarette smoking is overall decreasing, the use of new products such as electronic cigarettes has risen dramatically in recent years [1]. Nicotine administration induces a series of effects, ranging from pleasant (i.e. appetitive, rewarding, reinforcing, anxiolytic…) to noxious (i.e. anxiogenic, aversive…) [2,3]. These multifaceted effects have been described both in humans and in rodents. They greatly depend on the dose of nicotine administered, show substantial inter-individual variability, and are believed to be key in the regulation of nicotine intake and in the maintenance of addiction [2-4]. Understanding the variable effects of nicotine at the molecular and circuit levels is therefore fundamental to progress in the pathophysiology of nicotine addiction, and to develop efficient smoking-cessation therapies.

Nicotine mediates its physiological effects by activating nicotinic acetylcholine receptors (nAChRs), pentameric ligand-gated ion channels encoded by a large multigene family [2]. There are nine nAChR α (α2-10) and three β (β2-4) subunits expressed in the brain, which can assemble to form homo-pentamers or hetero-pentamers with various localizations and functions [5,6]. Initiation of consumption and reinforcement to nicotine involve the mesolimbic dopamine reward circuit, which originates in the ventral tegmental area (VTA) [7]. Nicotine primarily acts on this circuit by activating α4β2 nAChRs, a receptor subtype that displays high affinity for the drug [7-10]. Interestingly, an acute injection of nicotine also inhibits a subset of VTA dopamine neurons that project to the amygdala [11], and this results in elevated anxiety in mice,

illustrating the heterogeneity of the brain reward circuit, and the complexity of nicotine

dependence.

Another important pathway in the neurobiology of nicotine addiction is the medial

habenulo-interpeduncular (MHb-IPN) axis [12-15]. This pathway is deeply implicated in

the regulation of aversive physiological states such as fear and anxiety [16-18]. It is

believed to directly mediate aversion to high doses of nicotine [12,13], to trigger the

affective (anxiety) and somatic symptoms following nicotine withdrawal [19-22] and to be

involved in relapse to nicotine-seeking [23]. Strikingly, neurons of the MHb-IPN axis

express the highest density and largest diversity of nAChRs in the brain, notably the

rare α5, α3 and β4 subunits [6]. These are encoded by the CHRNA5-A3-B4 gene cluster,

of which some sequence variants are associated with a high risk of addiction in humans

[24,25]. The α3 and β4 subunits are virtually absent in the VTA, or in other parts of the

brain. The α3β4 nAChR shows lower affinity for nicotine than the α4β2 subtype, which

has contributed to the widely accepted idea that nicotine is rewarding at low doses

because it activates primarily α4β2 receptors of the VTA, while it is aversive at high

doses because only then does it activate α3β4 nAChRs of the MHb-IPN axis [12,14].

Nevertheless, this hypothesis of different sensitivity to nicotine in different circuits is

based on indirect evidence from c-fos quantification or brain slice physiology

experiments, and does not take into account the possible adaptive changes in the

circuits that result from repeated exposure to nicotine. Therefore, recording the

physiological response of IPN neurons to nicotine *in vivo*, both in naive animals and

after prolonged exposure to nicotine, remains a prerequisite to understand the

mechanism by which smokers develop tolerance to the aversive effects of nicotine.

A distinct feature of addiction is that, overall, only some individuals lose control over

their drug use, progressively shifting to compulsive drug intake [26-30]. About a third to

half of those who have tried smoking tobacco become regular users [31]. Individual

differences in the sensitivities of the VTA and MHb-IPN systems, and in their respective

adaptations during chronic tobacco use, could contribute to the vulnerability to nicotine

and to the severity of the addiction process [14,32-35]. Yet, the neural mechanism that

makes individuals more prone to maintain nicotine consumption versus durably stop

are unclear. In addition, whether the MHb-IPN pathway respond differently to nicotine

in individuals with and without a history of nicotine use is largely unknown. Here we

used isogenic mice and electrophysiology (*ex vivo* and *in vivo*) to study the neuronal

correlates of inter-individual variabilities in nicotine consumption behavior, and their

adaptation after chronic exposure to the drug.

Results

Heterogeneity in nicotine consumption in isogenic wild-type mice.

We used a continuous access, two-bottle choice nicotine-drinking test to assess

consumption profiles in wild-type (WT) C57Bl6 male mice single-housed in their home

cage (Fig. 1A). In this test, animals have continuous and concurrent access to two

bottles containing a solution of either 2% saccharine (vehicle) or nicotine plus 2%

saccharine (to mask the bitter taste of nicotine). After a 4-day habituation period with

water in both bottles, nicotine concentration was progressively increased in one bottle

across 16 days, from 10 to 200 μg/ml (4 days at each concentration), while alternating

the side of the nicotine-containing solution every other day to control for side bias (Fig.

1B). Consumption from each bottle was measured every minute. We found that the

daily nicotine intake increased throughout the paradigm, to stabilize at about 10

mg/kg/day on average for the highest nicotine concentration tested (Fig. 1C). Overall,

the percent of nicotine consumption, i.e., the nicotine solution intake relative to the total

fluid intake, was initially close to 50%, and decreased below that value for nicotine

concentrations above 50 μg/ml (Fig. 1D). These results match what was observed in

previous studies using C57Bl6 male mice, notably the fact that these mice rarely show

over 50% nicotine consumption, whether the water is supplemented with saccharine

or not [36,37]. The decreased percent nicotine consumption observed at the population

level in the course of the task suggests that mice adapt their behavior to reduce their

number of visits to the nicotine-containing bottle. We indeed observed that mice rapidly

(within a day) adjusted their nicotine intake when the concentration of nicotine

increases (Fig. S1A), resulting in titration of the nicotine dose, as previously reported

[12,34].

We noticed some disparity between mice and decided to analyze more precisely

nicotine consumption profiles for each individual. In particular, some mice abruptly

reduced their intake right after an increase in nicotine concentration in the bottle, or

had very low nicotine intake throughout the entire task (intake never exceeded 2

mg/kg/day) (Fig. 1E). We classified these mice (17/35) as "avoiders". The other half of

the mice (18/35), in contrast, displayed a continuous increase in nicotine intake, or

eventually reached a titration plateau in their consumption. These mice were classified

as "non-avoiders". Another way of looking at these distinct consumption profiles is to

quantify the differences in intake between two consecutive days (positive differences

indicate increased intake, while negative differences indicate decreased intake). The

distinction between the two phenotypic groups appears clearly when we plot, for each

mouse, the minimum and maximum values of the difference in intake between two

consecutive days (Fig. 1E, right). The group of avoiders was characterized either by a

minimum difference in intake that was negative (mice that reduce their intake abruptly),

or by both a minimum and a maximum difference in intake that were close to zero (very

low intake throughout the task). In contrast, the non-avoider mice were characterized

by a minimum difference in intake between two consecutive days that was always

positive, indicating continual increase in intake throughout the task. Overall, only a

small proportion of the mice (7/35, all non-avoiders) reached a plateau in their

consumption, which somewhat contrasts with the apparent titration observed at the

population level, either here (Fig. 1C) or in previous studies [12,34]. Avoiders and non-

avoiders (phenotypes defined as above throughout the manuscript) showed on

average similar nicotine intake for low concentrations of nicotine (10 and 50 µg/ml, p >

0.05), but while nicotine intake increased throughout the task for non-avoiders (16.9 ±

2.9 mg/kg/day for 200 µg/ml of nicotine) it dropped down to 2.6 ± 0.4 mg/kg/day for

such high nicotine concentration in avoiders (Fig. 1F), and almost reached zero over

the last three days (Fig. 1G). The percent nicotine consumption was fairly constant

throughout the task in non-avoiders, whereas in avoiders, it drastically decreased as

nicotine concentration increased, to approximate zero at the end of the task (Fig. S1B).

We then compared the level of aversion produced by nicotine in avoiders and non-

avoiders, with that produced by quinine, a notoriously bitter molecule. We found that

all of the naive mice actively avoided the quinine-containing solution, and showed a

percent quinine consumption close to zero, as observed with nicotine for avoiders, but

not for non-avoiders (Fig. S1C). Together, these results suggest that avoiders display

strong aversion to nicotine.

<u>The concentration of nicotine triggering aversion differs between mice</u>

Owing to the oral nature of the test, it may be difficult for mice to associate consumption

in a particular bottle with the physiological effects (whether positive or negative) of

nicotine. We thus systematically analyzed the percent nicotine consumption

throughout the task in individual mice, to examine their variability in choice patterns

and behavioral alterations. We found that some non-avoiders actively tracked the side

associated with nicotine when bottles were swapped (e.g. mouse #1 in Fig. 2A),

indicating strong preference for the bottle containing nicotine and active consumption.

Other non-avoiders displayed a strong side preference and never alternated drinking

side (e.g. mouse #2 in Fig. 2A), and hence consumed nicotine in a more passive

fashion. In contrast, all avoider mice (n = 17) displayed avoidance of the nicotine-

containing solution, whether they initially tracked the nicotine solution (e.g. mice #3

and 4 in Fig. 2A) or not (e.g. mice #5 and 6 in Fig. 2A). To quantify the evolution of

nicotine consumption at the individual level throughout the task, and to better take into

account the passive consumption behavior of some of the mice, we mapped each

profile in a pseudo-ternary plot where the base represents the nicotine consumption

index (from 0 to 100%), while the top apex represents 100% side bias (Fig. 2B). Such

a ternary representation enables us to graphically distinguish between mice that

actively track the nicotine bottle (bottom right apex, 100% nicotine consumption index),

mice that actively avoid nicotine (bottom left apex, 0% nicotine consumption index),

and mice that have a strong side bias (top apex), and to calculate the shortest distance

for each mouse to each of the three apices. In addition, by representing the trajectory

for each individual from the water condition to the 200 µg/ml nicotine condition, this

plot can be used to reveal and quantify behavioral adaptations (or lack thereof) in each

individual. Overall, we found that the behavior of non-avoiders was on average fairly

consistent throughout the task, i.e. their distance from the three apices was not really

impacted by the modifications in nicotine concentration (Fig. 2C and Fig. S2), whether

they consumed nicotine actively or passively. In contrast, the behavior of avoiders was

highly nicotine concentration-dependent, with mice going further away from the 100%

nicotine apex as nicotine concentration increases (Fig. 2C and Fig. S2A-B). Aversion

to nicotine in avoider mice mainly occurred at the transition from 100 to 200 µg/ml, but

some mice displayed aversion at concentrations as low as 10 µg/ml (Fig. 1E, 2A and

2C). Together, these results show that most mice can actively track or avoid nicotine,

indicating that they can discriminate nicotine from the control solution. Avoiders started

actively avoiding the nicotine-containing bottle once a specific drug concentration was

reached, suggesting the existence of a threshold at which nicotine aversion is

triggered, which apparently differs between mice.

Do avoiders learn to keep away from the nicotine-containing solution, or do they just

rapidly react to the nicotine concentration in the bottle to adjust their daily intake? To

answer this question, we added at the end of the two-bottle choice task, i.e. after the

200 µg/ml nicotine concentration, a condition with a low concentration of nicotine (50

µg/ml) for four days. We chose 50 µg/ml of nicotine because avoiders and non-

avoiders initially displayed comparable nicotine intake (Fig. 1F) and percent

consumption (Fig. S1B) at this concentration. We hypothesized that if avoiders

increased their percent nicotine consumption at the 200 to 50 µg/ml transition, it would

indicate a rapid adjustment to the concentration proposed, so to maintain their level of

intake constant. In contrast, if avoiders maintained a steady, low percent nicotine

consumption, it would indicate that nicotine aversion persists, independently of the

dose. We indeed found that lowering nicotine concentration from 200 to 50 µg/ml did

not increase percent nicotine consumption in avoiders (Fig. 2D), at least for the four

days mice were subjected to this concentration, suggesting aversion learning for

nicotine, and a behavioral adaptation that leads to near complete cessation of nicotine

consumption.

<u>Nicotine consumption negatively correlated with the amplitude of nicotine-evoked</u>

<u>currents in the IPN</u>

We then investigated the neural correlates of such aversion to nicotine. We

hypothesized that the IPN, which is involved in nicotine aversion and in negative

affective states [3,38,39], might be differently activated by nicotine in avoiders and non-

avoiders. Therefore, we used whole-cell patch-clamp recordings in brain slices to

assess, at completion of the two-bottle choice task, the functional expression level of

nAChRs in IPN neurons. We recorded neurons from the dorsal and rostral IPN

because these neurons have high nAChR density [14,15,40,41]. To record nicotine-evoked

currents, we used a local puff application of nicotine at a concentration (30 µM) close

to the EC50 for heteromeric nAChRs [42]. We found that the amplitude of nicotine-

evoked currents was higher in IPN neurons of avoider mice than in non-avoiders (Fig.

2E). Moreover, we found a negative correlation between the average amplitude of

nicotine-evoked current in IPN neurons, and nicotine consumption (measured over the

last 24 hours prior to the patch-clamp recording, Fig. 2F), suggesting that nicotine

consumption in mice is negatively linked to the amplitude of the response to nicotine

in IPN neurons.

<u>Chronic nicotine treatment alters both nicotinic signaling in the IPN and nicotine</u>

<u>consumption.</u>

It is still unclear at this stage whether chronic nicotine exposure progressively alters

the response of IPN neurons to the drug (non-avoiders being further exposed to high

nicotine doses than avoiders) or whether an intrinsic difference pre-exists in avoiders

and non-avoiders. To determine the effect of chronic nicotine exposure on nAChR

current levels in IPN neurons, we passively and continuously exposed mice to nicotine

for 4 weeks, using subcutaneously implanted osmotic minipumps. The concentration

of nicotine in the minipump (10 mg/kg/day) was chosen to match the average voluntary

nicotine intake in the two-bottle choice task (Fig. 1C). We then recorded from acute

brain slices and found that indeed, prolonged exposure to nicotine reduced the

amplitude of nicotine-evoked currents in the IPN of these mice compared to control

mice treated with saline (Fig. 3A). These results are consistent with the reduced current

amplitudes observed in mice that underwent the two-bottle choice task, compared to

naive mice in their home cage (Fig S3A). Because IPN neurons are mostly silent in

brain slices, and in order to preserve the entire circuitry intact, we decided to perform

juxtacellular recordings of IPN neurons *in vivo*, and to characterize their response

(expressed in % of variation from baseline) to an intravenous (i.v.) injection of nicotine

(30 µg/kg). To our knowledge, no description of *in vivo* recordings of IPN neurons, and

thus no criteria for identification, have been reported as yet, hence we solely

considered neurons that were labelled *in vivo* with neurobiotin and confirmed to be

within the IPN for the analysis. We found that nicotine i.v. injections in naive WT mice

induced an increase in the firing rate of IPN neurons compared to an injection of saline,

and that this acute effect of nicotine was reduced after the prolonged (4 weeks) passive

exposure of mice to the drug (Fig.3B). Some of the IPN neurons, however, responded

to nicotine by decreasing their firing rate, and the degree of inhibition was lower in the

group exposed to chronic nicotine (Fig. S3B). Together, these *ex vivo* and *in vivo*

recordings demonstrate that prolonged exposure to nicotine markedly reduces

nicotine-evoked responses in mouse IPN neurons.

To verify the hypothesis that a modification in the cholinergic activity of IPN neurons

impacts nicotine aversion, we did a series of experiments. Firstly, we evaluated the

consequence of prolonged nicotine exposure on nicotine consumption. Mice were

implanted with an osmotic minipump to passively deliver nicotine and, after 20 days,

were subjected to a modified two-bottle task that consisted in a rapid presentation to a

high concentration (100 µg/ml) of nicotine (Fig. 3C). We chose this protocol to avoid

the confounding effects of a gradual exposure to nicotine, and to evoke strong aversion

in mice. We found that indeed, mice pretreated with saline (controls) showed low

percent nicotine consumption (Fig. 3C), which may indicate aversion to such high

nicotine concentration. In contrast, mice pretreated with nicotine did not decrease on

average their percent consumption when nicotine was introduced in the bottle,

suggesting that they may have developed tolerance for the aversive effects of nicotine.

When looking at the data day by day, we observed that control mice abruptly

decreased their percent consumption when nicotine was introduced, while for nicotine-

treated animals the decrease was more gradual over the four days (Fig. S3C). Overall,

this resulted in greater nicotine intake for the group treated with nicotine than for the

group treated with saline (Fig. 3C). When focusing on individuals, we observed that a single saline-pretreated mouse (1/23) increased its percent consumption when nicotine was introduced in the task, while the great majority of the mice actively avoided nicotine (close to the 0% Nicotine apex). In contrast, a marked proportion of the mice treated with nicotine (8/25) increased their percent consumption when nicotine was introduced (Fig. 3D, p-value = 0.037 Chi-squared). The two groups were identical in the water/water session (Fig. 3E, top). However, in the water/nicotine session, mice pretreated with nicotine showed a greater distance from the 0% nicotine consumption apex, and a shorter distance from the 100% nicotine consumption apex than mice pretreated with saline (Fig. 3E, bottom), indicating decreased aversion and increased preference for nicotine in the nicotine-pretreated group. Altogether, these electrophysiological and behavioral data demonstrate that prolonged exposure to nicotine both decreases nicotine efficacy in the IPN, and also decreases aversion to nicotine in individuals, resulting in increased consumption of the drug. Yet, chronic nicotine can produce adaptations in other brain circuits, and whether the neurophysiological changes observed in the IPN are causally related to the variations in aversion sensitivity remains to be demonstrated.

Nicotine avoidance involves β4*nAChRs.

We turned to mutant mice deleted for the gene encoding the nAChR β4 subunit ($β4^{-/-}$ mice), because of the strong and restricted expression of this subunit in the MHb-IPN pathway [32,43,44]. We found that $β4^{-/-}$ mice displayed both greater percent nicotine consumption and greater nicotine intake than WT animals, with minimal concentration-dependent change in percent nicotine consumption (Fig. 4A). When looking at

individuals, we observed both active and passive nicotine-drinking profiles in β4$^{-/-}$ mice,

as already observed in WT mice. Strikingly however, none of the β4$^{-/-}$ mice (0/13)

showed aversion-like behavior at high nicotine concentration, which contrasts with the

high proportion of avoiders in WT animals (17/35, Fig. 4B, Fig. S4A, p =0.04 Pearson's

Chi squared with Yates 'continuity correction). WT mice showed a strong nicotine

concentration-dependent adaptation in their behavior, while β4$^{-/-}$ mice had a more

consistent behavior throughout the task (Fig. 4C and Fig. S4B, C).

To verify that responses to nicotine were affected in IPN neurons of β4$^{-/-}$ mice, we

performed whole-cell patch-clamp recordings. We found that the amplitude of nicotine-

evoked currents in the IPN was on average three-fold lower in β4$^{-/-}$ than in WT mice

(Fig. 4D), confirming that β4*nAChRs are the major receptor subtype in the IPN.

Furthermore, prolonged nicotine treatment had no significant effect on the amplitude

of nAChR currents in these knock-out mice (Fig. 4D), suggesting that the

downregulation observed after chronic nicotine treatment in the IPN of WT mice mainly

affects β4*nAChRs. We then used *in vivo* juxtacellular recordings, and performed

dose-response experiments (7.5 - 30 μg/kg) in order to assess the role of β4*nAChRs

in the response to different doses of nicotine. In WT mice, nicotine i.v. injections

resulted in a dose-dependent increase in IPN neuron activity (Fig. 4E). In β4$^{-/-}$ mice,

responses to nicotine were of smaller amplitude, especially for the highest dose of

nicotine tested (Fig. 4E), further demonstrating the important role of β4*nAChRs in the

response of the IPN to nicotine. In both WT and β4$^{-/-}$ mice, we observed a population

of neurons that decreased their firing rate in a dose-dependent manner, yet with no

difference in the amplitude of the response between the two genotypes (Fig. S4D-F),

suggesting that β4*nAChR are mainly involved in the increase, but not in the decrease,

of neuronal activity in response to nicotine injection. Collectively, our results in β4$^{-/-}$

mice demonstrate the key role of the nAChR β4 subunit in signaling aversion to

nicotine, and its predominant function in the activation of the IPN by nicotine.

<u>β4*nAChRs of the IPN are critically involved in nicotine aversion.</u>

β4 nAChRs are enriched in the IPN, yet they are also expressed to some extent in

other brain regions. Hence, to directly implicate β4*nAChRs of IPN neurons in nicotine

aversion, and more generally in nicotine consumption, we targeted re-expression of β4

in the IPN specifically, using lentiviral vectors in β4$^{-/-}$ mice (KO-β4^{IPN} mice, Fig. 5A).

Mice transduced with eGFP (KO-GFP^{IPN} mice) were used as controls. Proper

transduction in the IPN was verified using immunohistochemistry after completion of

the two-bottle choice task (Fig. 5A), and mice with expression of GFP in the VTA were

excluded from analyses. Transduction of β4, but not of GFP alone, in the IPN increased

the amplitude of nicotine-evoked currents (Fig. 5B) and restored levels found in WT

animals (U test, p = 0.6, Fig. S5A). We compared nicotine intake in the two groups of

mice in the two-bottle choice task. We found that re-expression of β4 in the IPN of β4^{-}

$^{/-}$ mice decreased nicotine intake compared to the group of mice transduced with eGFP

in the IPN (Fig. 5C). Overall, nicotine intake was similar in WT and in KO-β4^{IPN} animals

(p>0.5 for all concentrations), demonstrating the causal role of β4 nAChRs of IPN

neurons in nicotine consumption behaviors. At the individual level, the proportion mice

that avoided nicotine at 200 μg/kg was very low for KO-GFP^{IPN} control mice (2/18), but

greater for KO-β4^{IPN} mice (8/17, Fig. 5D, p = 0.04 Chi squared). The behavior of KO-

GFP^{IPN} was steady throughout the task, whereas it was highly nicotine concentration-

dependent for KO-β4^{IPN} mice (Fig. 5E and Fig. S5B,C), as already observed with WT

mice. Mice with strong percent nicotine consumption were only found in the KO-GFP[IPN]

control group. Collectively, these results show that selective re-expression of β4 in the

IPN of β4[-/-] mice rescued aversion for nicotine, and highlight the specific role of

β4*nAChRs of IPN neurons in signaling aversion to nicotine, and in the control of

nicotine intake.

Discussion

We used a two-bottle choice paradigm to assess inter-individual differences in nicotine

consumption in mice, and to evaluate how pre-exposure to nicotine modifies drug

taking. Oral self-administration is a classical method for chronic nicotine administration

as it provides rodents with *ad libitum* access to nicotine, likely mimicking administration

in human smokers, while minimizing stress from handling [45]. We observed that, on

average, WT mice titrate their intake to achieve a consistent nicotine dose, in

agreement with previous reports [12,34,35], and discovered that mutant mice lacking the

nAChR β4 subunit did not, resulting in greater nicotine intake, notably at high nicotine

concentrations in the drinking solution. These behavioral results are in agreement with

the greater intracranial self-administration observed at high nicotine doses in these

mice [33] and, conversely, with the results obtained with transgenic TABAC mice

overexpressing the β4 subunit at endogenous sites, which avoid nicotine and

consequently consume very little [13,33]. Nevertheless, it should be noted that conflicting

results have also been reported regarding the role of β4 nAChRs in nicotine

consumption. Notably, intravenous self-administration of nicotine is lower in β4[-/-] mice

despite a higher sensitivity of the VTA to nicotine in these mice [32], and self-

administration is higher in TABAC mice despite reduced nicotine-induced activation of

the VTA [46]. The increased consumption at high nicotine concentration reported here

for β4[-/-] mice resembles what was observed in mutant mice either lacking the α5

subunit [12] or with low levels of the α3 subunit [47], likely because the α3, α5, and β4

nAChR subunits, which are encoded by the same gene cluster, co-assemble in brain

structures, notably the MHb-IPN pathway, to produce functional heteromeric nAChRs

that contribute to the control of nicotine intake.

In the two-bottle choice nicotine-drinking test, a percent nicotine consumption below

50% is usually interpreted as a sign of aversion to the drug [13,37]. However, it may rather

indicate a process of regulation of nicotine consumption, through which mice adapt to

an increase in nicotine concentration by reducing the volume of nicotine intake at each

visit. At low nicotine concentrations, percent nicotine consumption never really

exceeded 50% in average, as already seen in previous studies [37], yet very few mice

readily avoided the nicotine-containing bottle, and many mice did track nicotine, which

would rather indicate an actual appetite (as opposed to an aversion) for the drug. As

nicotine concentration increases, the percent nicotine consumption dropped to about

20% on average at the end of the task (Fig. 1D), yet it was associated with an increase

in nicotine intake (in mg/kg/day, Fig. 1C). Hence the nicotine preference score alone

may not be sufficient to determine whether nicotine is aversive or not. We believe that,

instead, aversion to the drug may be better defined by a constantly low consumption,

or a sudden drop in nicotine intake, as seen here with the group of avoider mice (Fig.

1E-G).

One important limitation of population-level analyses is that they greatly limit the ability

to examine inter-individual differences in drug taking behaviors. It is indeed

increasingly acknowledged that in mice, as in humans, there is a substantial variability

in the susceptibility for developing drug use disorders [27-29,48-51]. Yet, it is still unclear

why some individuals are more susceptible than others to become regular users. Here

we inspected drinking profiles in individual isogenic mice, and discovered large inter-

individual differences in nicotine vulnerability: about half of the WT mice, the avoiders,

durably quit nicotine at a certain concentration, whereas the other half, the non-

avoiders, continued consumption even at high concentration of nicotine, classically

described as aversive [3]. Avoiders displayed variable concentration thresholds required

for triggering aversion, and some of them even developed aversion at the beginning of

the two-bottle choice task, when concentrations of nicotine were still low. This finding,

which challenges the popular idea that aversion is only triggered by high doses of

nicotine [3,12-14], is somewhat supported by our electrophysiogical data. Indeed, by

directly measuring the response of IPN neurons to different doses of nicotine *in vivo*,

we show that concentrations of nicotine as low as 7.5 μg/kg can engage the IPN

circuitry, suggesting that nicotine-induced signaling can emerge in this structure even

at low drug concentration. Non-avoiders (WT and $\beta 4^{-/-}$ mice) may retain an aversion

threshold, but which would exceed the highest concentration tested here. Importantly,

very few mice showed what could be considered as titration (plateaued consumption),

emphasizing the needs to consider individual behavior, as opposed to group behavior,

in addiction research.

We also discovered that the functional expression level of $\beta 4$-containing nAChRs in

the IPN underlies these different sensitivities to the aversive properties of nicotine.

Indeed, inter-individual variability for nicotine aversion was nearly eliminated in $\beta 4^{-/-}$

mice, none of which quit drinking nicotine, and was restored after selective re-

expression of the β4 subunit in the IPN. This selective re-expression experiment

discards the possibility that different sensitivities to the bitter taste of nicotine solutions

explain the different drinking profiles of avoiders and non-avoiders. Strikingly, we

observed a negative correlation between nicotine consumption and the response of

the IPN to the drug: mice consuming large amounts of nicotine displayed lower

nicotine-evoked currents in the IPN than mice consuming small amounts of nicotine,

which displayed large currents. The expression level of β4-containing nAChRs in the

IPN may thus determine the level of aversion to the drug, and consequently its intake.

We suggest that β4-containing nAChRs, by engaging the IPN circuitry, initiate a

primary response to nicotine that, if above a certain threshold, will trigger acute

aversion to the drug, thus impacting the balance between drug reward and aversion to

limit drug consumption. In line with this, it was found that pharmacological or

optogenetic stimulation of the MHb-IPN pathway could directly produce aversion

[14,15,34], while pharmacological inactivation of this pathway increases nicotine intake [12].

From our results, we further suggest that the aversion produced by nicotine is not just

an acute response to the dose that has just been ingested. Aversion can also be lasting

for days, as evidenced by the fact that mice persist in avoiding the nicotine-containing

bottle even after the concentration has been reduced to a dose they used to ingest.

This echoes observations made in humans, in which the unpleasant initial responses

to cigarettes is associated with a reduced likelihood of continued smoking [52]. Such a

sustained aversive reaction to nicotine was conditioned, in mice, by nicotine itself, and

required β4-containing nAChRs of the IPN for its onset, but most likely involves other

brain circuits for its persistence in the long-term. Identifying the molecular and cellular

mechanism of long-term aversion to nicotine in mice will be instrumental to progress

in our understanding of human dependence to tobacco.

We did not observe major differences in nicotine intake between avoiders and non-

avoiders at the beginning of the two-bottle choice experiment, for low concentrations

of nicotine (< 100 μg/ml), but cannot completely rule out pre-existing inter-individual

differences that may explain the opposite trajectories taken by the two groups. Indeed,

the mice used in this study were isogenic, yet epigenetic changes during development

or differences in social status, which are known to affect brain circuits and individual

traits [53], may affect the responses of the IPN to nicotine. It is indeed tempting to

speculate that external factors (e.g. stress, social interactions…) that would affect the

expression level of β4-containing nAChRs in the IPN, will have a strong impact on

nicotine consumption. In addition to pre-existing differences, history of nicotine use

could produce long-lasting molecular and cellular adaptations in the IPN circuitry that

may alter nicotine aversion and consumption. In the VTA, chronic nicotine upregulates

the number of β2-containing receptors at the cell surface [54,55]. We discovered that

prolonged nicotine exposure had the opposite effect on β4-containing nAChRs of the

IPN: it downregulated and/or desensitized these receptors, as evidenced by the

decreased response to nicotine both *ex vivo* and *in vivo*. This effect seems to be

specific to β4-containing nAChRs, since chronic nicotine had minimal consequence on

the residual IPN nAChR current in β4$^{-/-}$ mice. These results may appear at odds with

the recent report of increased nAChR currents in IPN slices of nicotine-treated mice [56].

However, differences in the effective agonist concentrations used in the two studies

(30 μM of nicotine with local puff here, *vs.* 50 μM of caged-nicotine, but effective

concentration probably much lower after photo-uncaging in [56]) may explain this

discrepancy. Indeed, by using a lower effective agonist concentration, Arvin et al. [56]

were possibly not recruiting the low-affinity β4-containing receptors, which are the

receptor subtypes we found to be downregulated by chronic nicotine exposure (Fig. 4D). Whatever the reason behind these opposite results is, our behavioral data in nicotine-treated WT and $\beta4^{-/-}$ mice completely match: both displayed reduced responses to nicotine (*ex vivo* and i*n vivo)* in IPN neurons, as well as increased nicotine intake compared to naive, WT animals. We suggest that long-term exposure to nicotine decreases the likelihood to reach the threshold at which mice develop aversion to the drug, which ultimately leads to increased drug consumption. In other words, nicotine intake history weakens the ability of nicotine to induce aversion in mice. In most nicotine replacement therapies, such as gums or patches, nicotine is slowly administered over prolonged periods of time, to supposedly attenuate the negative emotional reactions elicited by nicotine withdrawal [57]. However, our data indicate that mice under prolonged nicotine administration will also develop tolerance to the aversive effects of nicotine, highlighting the necessity to develop alternative medical approaches.

Materials and Methods

Animals

Eight to sixteen-week old wild-type C57BL/6J (Janvier labs, France) and Acnb4 knock-out ($\beta4^{-/-}$) mice (Pasteur Institute, Paris) [58] were used for this study. $\beta4^{-/-}$ mice were backcrossed onto C57BL/6J background for more than twenty generations. Mice were maintained on a 12h light-dark cycle. All experiments were performed in accordance with the recommendations for animal experiments issued by the European Commission directives 219/1990, 220/1990 and 2010/63, and approved by Sorbonne Université.

Two-bottle choice experiment

Mice single-housed in a home cage were presented with two bottles of water (Volvic) for a habituation period of 4 days. After habituation, mice were presented with one bottle of saccharine solution (2%, Sigma Aldrich) and one bottle of nicotine (free base, Sigma Aldrich) plus saccharine (2%) solution diluted in water (adjusted to pH ~7.2 with NaOH). Unless otherwise noted, four different concentrations of nicotine were tested consecutively (10, 50, 100 and 200 µg/ml) with changes in concentration occurring every 4 days. For the two-bottle aversion task, a single nicotine concentration (100 µg/ml) was used after the habituation period. Bottles were swapped every other day to control for side preference. The drinking volume was measured every minute with an automated acquisition system (TSE system, Germany). Nicotine intake was calculated in mg of nicotine per kilogram of mouse body weight per day (mg/kg/d). To minimize stress from handling, mice were weighed every other day, since we found their weigh to be sufficiently stable over two days. Percent nicotine consumption was calculated as the volume of nicotine solution consumed as a percentage of the total fluid

consumed. Mice showing a strong side bias (preference <20% or >80%) in the habituation period were not taken into account for the analyses.

For the pseudo-ternary plot analyses, we determined the percent nicotine consumption on the left-hand side (%c1) and the percent nicotine consumption on the right-hand side (%c2), for each nicotine concentration and each animal. We then calculated the nicotine consumption and side bias indexes, by plotting the minimum min(%c1, %c2) against the maximum max(%c1, %c2), and used a 90° rotation to obtain the pseudo-ternary plot. In this plot, the three apices represent mice that avoid nicotine on both sides (0% nic.), mice that track nicotine on both sides (100% nic.), and finally mice that drink solely one side (side biased).

Prolonged treatment with nicotine

Osmotic minipumps (2004, Alzet minipump) were implanted subcutaneously in 8-week-old mice anesthetized with isoflurane (1%). Minipumps continuously delivered nicotine (10 mg/kg/d) or saline (control) solution with a rate of 0.25 µl/h during 4 weeks.

Brain slice preparation

Mice were weighed and then anaesthetized with an intraperitoneal injection of a mixture of ketamine (150 mg/kg, Imalgene 1000, Merial, Lyon, France) and xylazine (60 mg/kg, Rompun 2%, Bayer France, Lyon, France). Blood was then fluidized by an injection of an anticoagulant (0.1mL, heparin 1000 U/mL, Sigma) into the left ventricle, and an intra-cardiac perfusion of ice-cold (0-4°C), oxygenated (95% O_2/5% CO_2) sucrose-based artificial cerebrospinal fluid (SB-aCSF) was performed. The SB-aCSF solution contained (in mM): 125 NaCl, 2.5 KCl, 1.25 NaH_2PO_4, 5.9 $MgCl_2$, 26 $NaHCO_3$, 25 sucrose, 2.5 glucose, 1 kynurenate (pH 7.2). After rapid brain sampling, slices (250

µm thick) were cut in SB-aCSF at 0-4°C using a Compresstome slicer (VF-200, Precisionary Instruments Inc.). Slices were then transferred to the same solution at 35°C for 10 min, then moved and stored in an oxygenated aCSF solution at room temperature. The aCSF solution contained in mM: 125 NaCl, 2.5 KCl, 1.25 NaH_2PO_4, 2 $CaCl_2$, 1 $MgCl_2$, 26 $NaHCO_3$, 15 sucrose, 10 glucose (pH 7.2). After minimum 1h of rest, slices were placed individually in a recording chamber at room temperature and infused continuously with aCSF recording solution at a constant flow rate of about 2 ml/min.

Ex vivo patch-clamp recordings of IPN neurons

Patch pipettes (5-8 MΩ) were stretched from borosilicate glass capillaries (G150TF-3, Warner instruments) using a pipette puller (Sutter Instruments, P-87, Novato, CA) and filled with a few microliters of an intracellular solution adjusted to pH 7.2, containing (in mM): 116 K-gluconate, 20 HEPES, 0.5 EGTA, 6 KCl, 2 NaCl, 4 ATP, 0.3 GTP and 2 mg/mL biocytin. Biocytin was used to label the recorded neurons. The slice of interest was placed in the recording chamber and viewed using a white light source and a upright microscope coupled to a Dodt contrast lens (Scientifica, Uckfield, UK). Neurons were recorded from the dorsal (IPDL) and rostral (IPR) parts on the IPN. Whole-cell configuration recordings of IPN neurons were performed using an amplifier (Axoclamp 200B, Molecular Devices, Sunnyvale, CA) connected to a digitizer (Digidata 1550 LowNoise acquisition system, Molecular Devices, Sunnyvale, CA). Signal acquisition was performed at 10 kHz, filtered with a lowpass (Bessel, 2 kHz) and collected by the acquisition software pClamp 10.5 (Molecular Devices, Sunnyvale, CA). Nicotine tartrate (30 µM in aCSF) was locally and briefly applied (200 ms puffs) using a puff pipette (glass pipette ~3 µm diameter at the tip) positioned about 20-30 µm from the

soma of the neuron. The pipette was connected to a Picospritzer (PV-800 PicoPump,

World Precision Instruments) controlled with pClamp to generate transient pressure in

the pipette (~2 psi). Nicotine-evoked currents were recorded in voltage-clamp mode at

a membrane potential of -60 mV. All electrophysiology traces were extracted and pre-

processed using Clampfit (Molecular Devices, Sunnyvale, CA) and analyzed with R.

In vivo electrophysiology

Mice were deeply anesthetized with chloral hydrate (8%, 400 mg/kg) and anesthesia

was maintained throughout the experiment with supplements. Catheters were

positioned in the saphenous veins of the mice to perform saline or nicotine intravenous

injections. Nicotine hydrogen tartrate salt (Sigma-Aldrich) was dissolved in 0.9% NaCl

solution and pH was adjusted to 7.4. The nicotine solution was injected at a dose of

7.5, 15 and 30 µg/kg. Borosilicate glass capillaries (1.5 mm O.D. / 1.17 mm I.D.,

Harvard Apparatus) were pulled using a vertical puller (Narishige). Glass pipettes were

broken under a microscope to obtain a ~1 µm diameter at the tip. Electrodes were filled

with a 0.5% NaCl solution containing 1.5% of neurobiotin tracer (AbCys) yielding

impedances of 6-9 MΩ. Electrical signals were amplified by a high-impedance amplifier

(Axon Instruments) and supervised through an audio monitor (A.M. Systems Inc.). The

signal was digitized, sampled at 25 kHz and recorded on a computer using Spike2

(Cambridge Electronic Design) for later analysis. IPN neurons were recorded in an

area corresponding to the following stereotaxic coordinates (4-5° angle): 3.3 - 3.6 mm

posterior to bregma, 0.2 - 0.45 mm from medial to lateral and 4.3 - 5 mm below the

brain surface. A 5 min-baseline was recorded prior to saline or nicotine i.v. injection.

For the dose-response experiments, successive randomized injections of nicotine (or

saline) were performed, interspaced with sufficient amount of time (> 10 min) to allow

the neuron to return to its baseline.

Stereotaxic viral injections

8-week old mice were injected in the IPN with a lentivirus that co-expresses the WT β4

subunit together with eGFP (or only eGFP for control experiments) under the control

of the PGK promoter. Lentiviruses were produced as previously described [7]. For viral

transduction, mice were anaesthetized with a gas mixture containing 1-3% isoflurane

(IsoVet®, Pyramal Healthcare Ltd., Nothumberland, UK) and placed in a stereotactic

apparatus (David Kopf Instruments, Tujunga, CA). Unilateral injections (0.1µl/min) of

1µl of a viral solution (Lenti.pGK.β4.IRES.eGFP, titer 150 ng/µl of p24 protein; or

Lenti.pGK.eGFP, titer 75 ng/µl of p24 protein) were performed using a cannula

(diameter 36G, Phymep, Paris, France). The cannula was connected to a 10 µL

Hamilton syringe (Model 1701, Hamilton Robotics, Bonaduz, Switzerland) placed in a

syringe pump (QSI, Stoelting Co, Chicago, IL, USA). Injections were performed in the

IPN at the following coordinates (5° angle): from bregma ML - 0.4 mm, AP - 3.5 mm,

and DV: - 4.7 mm (according to Paxinos & Franklin). Electrophysiological recordings

were made at least 4 weeks after viral injection, the time required for the expression of

the transgene, and proper expression was subsequently checked using

immunohistochemistry.

Immunocytochemical identification

Immunostaining was performed as described in [10]*, with the following* primary

antibodies: anti-tyrosine hydroxylase 1:500 (anti-TH, Sigma, T1299) and chicken anti-

eYFP 1:500 (Life technologies Molecular Probes, A-6455). Briefly, serial 60 µm-thick

sections of the midbrain were cut with a vibratome. Slices were permeabilized for one

hour in a solution of phosphate-buffered saline (PBS) containing 3% bovine serum

albumin (BSA, Sigma; A4503). Sections were incubated with primary antibodies in a

solution of 1.5 % BSA and 0.2 % Triton X-100 overnight at 4°C, washed with PBS and

then incubated with the secondary antibodies for 1 hour. The secondary antibodies

were Cy3-conjugated anti-mouse (1:500 dilution) and alexa488-conjugated anti-

chicken (1:1000 dilution) (Jackson ImmunoResearch, 715-165-150 and 711-225-152,

respectively). For the juxtacellular immunostaining, the recorded neurons were

identified with the addition of AMCA-conjugated streptavidin (1:200 dilution) in the

solution (Jackson ImmunoResearch). Slices were mounted using Prolong Gold

Antifade Reagent (Invitrogen, P36930). Microscopy was carried out either with a

confocal microscope (Leica) or with an epifluorescence microscope (Leica), and

images were captured using a camera and analyzed with ImageJ.

Statistical analysis.

All statistical analyses were computed using R (The R Project, version 4.0.0). Results

were plotted as a mean ± s.e.m. The total number (n) of observations in each group

and the statistics used are indicated in figure legends. Classical comparisons between

means were performed using parametric tests (Student's T-test, or ANOVA for

comparing more than two groups when parameters followed a normal distribution

(Shapiro test $P > 0.05$)), and non-parametric tests (here, Mann-Whitney or Friedman)

when the distribution was skewed. Multiple comparisons were corrected using a

sequentially rejective multiple test procedure (Holm-Bonferroni correction). All

statistical tests were two-sided. $P > 0.05$ was considered not to be statistically

significant.

Acknowledgments: The authors would like to thank Ines Centeno-Lemaire (Sorbonne Université, Paris, France) for her help with behavioral tests, Jean-Pierre Hardelin (ESPCI, Paris, France) for critical reading of the manuscript, and the animal facility at Institut de Biology Paris Seine (IBPS, Paris, France).

Funding:

Agence Nationale de la Recherche (ANR-21-CE16-0012 CHOLHAB to AM)

Fondation pour la Recherche Médicale (Equipe FRM EQU201903007961 to PF)

Institut National du Cancer Grant TABAC-16-022 and TABAC-19-02 (to PF).

Fundamental research prize from the Fondation Médisite for neuroscience (AM).

Fourth-year PhD fellowship from Fondation pour la Recherche Médicale (FDT201904008060 to RDC and FDT20170437427 to SM).

Fourth-year PhD fellowship from the Biopsy Labex (CN).

Fourth-year PhD fellowship from the Memolife Labex (JJ).

Author contribution:

Conceptualization: SM, PF and AM.

Methodology: SM, PF and AM.

Software: NT and SM.

Validation: SM, PF and AM.

Formal analysis: SM.

Investigation: SM, CN, EV, JJ, RDC, ST and FM.

Resources: SP and UM.

Writing - original draft: AM.

Writing - review and editing: SM, PF and AM.

Visualization: SM.

Supervision: PF and AM.

Funding acquisition: PF and AM.

Competing interests: The authors declare no competing interest.

Data and material availability: All data needed to evaluate the conclusions in the

paper are present in the paper and/or the Supplementary materials.

References

1. *WHO global report on trends in prevalence of tobacco use 2000-2025 third edition*. 1–121 (2019).

2. Wills, L. *et al.* Neurobiological Mechanisms of Nicotine Reward and Aversion. *Pharmacological Reviews* **74,** 271–310 (2022).

3. Fowler, C. D. & Kenny, P. J. Nicotine aversion: Neurobiological mechanisms and relevance to tobacco dependence vulnerability. *Neuropharmacology* **76 Pt B,** 533–544 (2014).

4. Verendeev, A. & Riley, A. L. The role of the aversive effects of drugs in self-administration. *Behavioural Pharmacology* **24,** 363–374 (2013).

5. Taly, A., Corringer, P.-J., Guedin, D., Lestage, P. & Changeux, J.-P. Nicotinic receptors: allosteric transitions and therapeutic targets in the nervous system. *Nat Rev Drug Discov* **8,** 733–750 (2009).

6. Zoli, M., Pistillo, F. & Gotti, C. Diversity of native nicotinic receptor subtypes in mammalian brain. *Neuropharmacology* **96,** 302–311 (2015).

7. Maskos, U. *et al.* Nicotine reinforcement and cognition restored by targeted expression of nicotinic receptors. *Nature* **436,** 103–107 (2005).

8. Tapper, A. R. *et al.* Nicotine activation of alpha4* receptors: sufficient for reward, tolerance, and sensitization. *Science* **306,** 1029–1032 (2004).

9. Tolu, S. *et al.* Co-activation of VTA DA and GABA neurons mediates nicotine reinforcement. *Mol Psychiatry* **18,** 382–393 (2013).

10. Durand-de Cuttoli, R. *et al.* Manipulating midbrain dopamine neurons and reward-related behaviors with light-controllable nicotinic acetylcholine receptors. *eLife* **7,** e37487 (2018).

11. Nguyen, C. *et al.* Nicotine inhibits the VTA-to-amygdala dopamine pathway to promote anxiety. *Neuron* **109,** 2604–2615.e9 (2021).

12. Fowler, C. D., Lu, Q., Johnson, P. M., Marks, M. J. & Kenny, P. J. Habenular α5 nicotinic receptor subunit signalling controls nicotine intake. *Nature* **471,** 597–601 (2011).

13. Frahm, S. *et al.* Aversion to Nicotine Is Regulated by the Balanced Activity of beta4 and alpha5 Nicotinic Receptor Subunits in the Medial Habenula. *Neuron* **70,** 522–535 (2011).

14. Wolfman, S. L. *et al.* Nicotine aversion is mediated by GABAergic
interpeduncular nucleus inputs to laterodorsal tegmentum. *Nature*
Communications **9,** 2710 (2018).

15. Morton, G. *et al.* Chrna5-Expressing Neurons in the Interpeduncular Nucleus
Mediate Aversion Primed by Prior Stimulation or Nicotine Exposure. *J. Neurosci.*
38, 6900–6920 (2018).

16. Otsu, Y. *et al.* Control of aversion by glycine-gated GluN1/GluN3A NMDA
receptors in the adult medial habenula. *Science* **366,** 250–254 (2019).

17. Yamaguchi, T., Danjo, T., Pastan, I., Hikida, T. & Nakanishi, S. Distinct Roles of
Segregated Transmissionof the Septo-Habenular Pathway in Anxiety and Fear.
Neuron **78,** 537–544 (2013).

18. Presynaptic Excitation via GABAB Receptors in Habenula Cholinergic Neurons
Regulates Fear Memory Expression. *Cell* **166,** 716–728 (2016).

19. Salas, R., Sturm, R., Boulter, J. & De Biasi, M. Nicotinic Receptors in the
Habenulo-Interpeduncular System Are Necessary for Nicotine Withdrawal in
Mice. *J. Neurosci.* **29,** 3014–3018 (2009).

20. Zhao-Shea, R. *et al.* Increased CRF signalling in a ventral tegmentalarea-
interpeduncular nucleus-medial habenulacircuit induces anxiety during nicotine
withdrawal. *Nature Communications* **6,** 1–13 (2015).

21. Pang, X. *et al.* Habenula cholinergic neurons regulate anxiety during nicotine
withdrawal via nicotinic acetylcholine receptors. *Neuropharmacology* **107,** 294–
304 (2016).

22. Zhao-Shea, R., Liu, L., Pang, X., Gardner, P. D. & Tapper, A. R. Activation of
GABAergic Neuronsin the Interpeduncular Nucleus Triggers Physical Nicotine
Withdrawal Symptoms. *Current Biology* **23,** 2327–2335 (2013).

23. Forget, B. *et al.* A Human Polymorphism in CHRNA5 Is Linked to Relapse to
Nicotine Seeking in Transgenic Rats. *Curr. Biol.* **28,** 3244–3253.e7 (2018).

24. Bierut, L. J. *et al.* Variants in nicotinic receptors and risk for nicotine dependence.
Am J Psychiatry **165,** 1163–1171 (2008).

25. Lassi, G. *et al.* The CHRNA5–A3–B4 GeneCluster and Smoking: FromDiscovery
to Therapeutics. *Trends in Neurosciences* **39,** 851–861 (2016).

26. George, O. & Koob, G. F. Individual differences in the neuropsychopathology of
addiction. *Dialogues Clin Neurosci* **19,** 217–229 (2017).

27. Deroche-Gamonet, V., Belin, D. & Piazza, P. V. Evidence for addiction-like behavior in the rat. *Science* **305,** 1014–1017 (2004).

28. Siciliano, C. A. *et al.* A cortical-brainstem circuit predicts and governs compulsive alcohol drinking. *Science* **366,** 1008–1012 (2019).

29. Juarez, B. *et al.* Midbrain circuit regulation of individual alcohol drinking behaviors in mice. *Nature Communications* **8,** 2220 (2017).

30. Pascoli, V. *et al.* Stochastic synaptic plasticity underlying compulsion in a model of addiction. *Nature* **564,** 1–22 (2018).

31. Centers for Disease Control and Prevention (US), National Center for Chronic Disease Prevention and Health Promotion (US)Office on Smoking and Health (US). How Tobacco Smoke Causes Disease: The Biology and Behavioral Basis for Smoking-Attributable Disease: A Report of the Surgeon General. (2010).

32. Harrington, L. *et al.* Role of β4* Nicotinic Acetylcholine Receptors in the Habenulo-Interpeduncular Pathway in Nicotine Reinforcement in Mice. *Neuropsychopharmacology* **41,** 1790–1802 (2016).

33. Husson, M. *et al.* β4-Nicotinic Receptors Are Critically Involved in Reward-Related Behaviors and Self-Regulation of Nicotine Reinforcement. *J. Neurosci.* **40,** 3465–3477 (2020).

34. Tuesta, L. M. *et al.* GLP-1 acts on habenular avoidance circuits to control nicotine intake. *Nature Neuroscience* **20,** 708–716 (2017).

35. Antolin-Fontes, B. *et al.* The habenular G-protein-coupled receptor 151 regulates synaptic plasticity and nicotine intake. *Proceedings of the National Academy of Sciences* **117,** 5502–5509 (2020).

36. Bagdas, D. *et al.* Assessing nicotine dependence using an oral nicotine free-choice paradigm in mice. *NP* **157,** 107669 (2019).

37. Matta, S. G. *et al.* Guidelines on nicotine dose selection for in vivo research. *Psychopharmacology* **190,** 269–319 (2006).

38. Molas, S., DeGroot, S. R., Zhao-Shea, R. & Tapper, A. R. Anxiety and Nicotine Dependence: Emerging Role of the Habenulo-Interpeduncular Axis. *Trends in Pharmacological Sciences* **38,** 169–180 (2017).

39. McLaughlin, I., Dani, J. A. & De Biasi, M. The medial habenula and interpeduncular nucleus circuitry is critical in addiction, anxiety, and mood regulation. *J Neurochem* **142,** 130–143 (2017).

786   40.   Hsu, Y.-W. A. *et al.* Medial habenula output circuit mediated by α5 nicotinic
787   receptor-expressing GABAergic neurons in the interpeduncular nucleus. *J.*
788   *Neurosci.* **33,** 18022–18035 (2013).

789   41.   Quina, L. A., Harris, J., Zeng, H. & Turner, E. E. Specific connections of the
790   interpeduncular subnuclei reveal distinct components of the
791   habenulopeduncular pathway. *J. Comp. Neurol.* **525,** 2632–2656 (2017).

792   42.   Fenster, C. P., Rains, M. F., Noerager, B., Quick, M. W. & Lester, R. A. Influence
793   of subunit composition on desensitization of neuronal acetylcholine receptors at
794   low concentrations of nicotine. *J. Neurosci.* **17,** 5747–5759 (1997).

795   43.   Shih, P.-Y. *et al.* Differential expression and function of nicotinic acetylcholine
796   receptors in subdivisions of medial habenula. *J. Neurosci.* **34,** 9789–9802
797   (2014).

798   44.   Grady, S. R. *et al.* Rodent habenulo-interpeduncular pathway expresses a large
799   variety of uncommon nAChR subtypes, but only the alpha3beta4* and
800   alpha3beta3beta4* subtypes mediate acetylcholine release. *J. Neurosci.* **29,**
801   2272–2282 (2009).

802   45.   Collins, A. C., Pogun, S., Nesil, T. & Kanit, L. Oral Nicotine Self-Administration
803   in Rodents. *J Addict Res Ther* **S2,** (2012).

804   46.   Gallego, X. *et al.* Overexpression of the CHRNA5/A3/B4 genomic cluster in mice
805   increases the sensitivity to nicotine and modifies its reinforcing effects. *Amino*
806   *Acids* **43,** 897–909 (2011).

807   47.   Elayouby, K. S. *et al.* α3* Nicotinic Acetylcholine Receptors in the Habenula-
808   Interpeduncular Nucleus Circuit Regulate Nicotine Intake. *J. Neurosci.* **41,** 1779–
809   1787 (2021).

810   48.   Nesil, T., Kanit, L., Collins, A. C. & Pogun, S. Individual differences in oral
811   nicotine intake in rats. *NP* **61,** 189–201 (2011).

812   49.   Garcia-Rivas, V., Cannella, N. & Deroche-Gamonet, V. Individual Variations in
813   the Mechanisms of Nicotine Seeking: A Key for Research on Nicotine
814   Dependence. *Neuropsychopharmacology* **42,** 584–586 (2017).

815   50.   Piazza, P. V., Deminière, J. M., Le Moal, M. & Simon, H. Factors that predict
816   individual vulnerability to amphetamine self-administration. *Science* **245,** 1511–
817   1513 (1989).

818   51.   Dongelmans, M. *et al.* Chronic nicotine increases midbrain dopamine neuron
819   activity and biases individual strategies towards reduced exploration in mice.
820   *Nature Communications* **12,** 6945 (2021).

821   52.   DiFranza, J. R. *et al.* Recollections and repercussions of the first inhaled
822   cigarette. *Addictive Behaviors* **29,** 261–272 (2004).

823   53.   Torquet, N. *et al.* Social interactions impact on the dopaminergic system and
824   drive individuality. *Nature Communications* **9,** 3081 (2018).

825   54.   Sallette, J. *et al.* Nicotine Upregulates Its Own Receptors through Enhanced
826   Intracellular Maturation. *Neuron* **46,** 595–607 (2005).

827   55.   Lester, H. A. *et al.* Nicotine is a Selective Pharmacological Chaperone of
828   Acetylcholine Receptor Number and Stoichiometry. Implications for Drug
829   Discovery. *AAPS J* **11,** 167–177 (2009).

830   56.   Arvin, M. C. *et al.* Chronic Nicotine Exposure Alters the Neurophysiology of
831   Habenulo-Interpeduncular Circuitry. *J. Neurosci.* **39,** 4268–4281 (2019).

832   57.   Hartmann-Boyce, J., Chepkin, S. C., Ye, W., Bullen, C. & Lancaster, T. Nicotine
833   replacement therapy versus control for smoking cessation. *Cochrane Database
834   of Systematic Reviews* **1,** 7–196 (2018).

835   58.   Xu, W. *et al.* Multiorgan autonomic dysfunction in mice lacking the beta2 and the
836   beta4 subunits of neuronal nicotinic acetylcholine receptors. *J. Neurosci.* **19,**
837   9298–9305 (1999).

Figure Legends

FIGURE 1

Figure 1: Two different profiles, avoiders and non-avoiders, emerged in WT mice subjected to a two-bottle choice nicotine-drinking test. A. Continuous access, two-bottle choice setup. **B.** Two-bottle choice paradigm. Each dot represents a bottle and is color-coded according to whether it contains water (blue or light blue), nicotine plus 2% saccharine (red, gradient of color intensities according to the nicotine concentration), or 2% saccharine (white) solutions. The nicotine concentration in the bottle increased progressively from 10 to 50, 100 and 200 µg/ml. Each condition lasted four days, and the bottles were swapped every other day. **C.** Nicotine intake (mg/kg/d), averaged over four days, at different nicotine concentrations (Friedman test, n = 35, df = 3, p < 0.001 and Mann-Whitney post-hoc test with Holm-Bonferroni correction). **D.** Percent nicotine consumption in WT mice for each concentration of nicotine, averaged over four days (Friedman test, n = 35, df = 4, p < 0.001 and Mann-Whitney post-hoc test with Holm-Bonferroni correction) **E.** Left, nicotine intake in individual avoiders (n = 17) and non-avoiders (n = 18). Right, minimum and maximum values of the difference in nicotine intake between two consecutive days, for each individual. **F.** Nicotine intake

in avoiders and non-avoiders for each nicotine concentration, averaged over four days

(Mann-Whitney comparison with a Holm-Bonferroni correction). **G.** Daily nicotine

intake in avoiders and non-avoiders for nicotine 100 and 200 µg/ml (paired Mann-

Whitney). Note the drop in nicotine consumption at day 17 for avoiders. In all figure

panels avoiders are depicted in pinkish-orange while non-avoiders are in blue. *** $p <$

0.001, ** $p < 0.01$, * $p < 0.05$.

FIGURE 2

Figure 2: Nicotine intake was negatively correlated with the amplitude of the response to nicotine in IPN neurons. A. Representative examples of choice behaviors (% consumption on the right vs. left bottle) in WT mice when the right-hand side bottle contains either nicotine + saccharine (red dots, grey stripe) or saccharine only (white dots, white stripes). Mice are same as in Fig. 1. **B.** Pseudo-ternary diagram representing, for each individual (18 non-avoiders and 17 avoiders, see Fig. 1), its nicotine consumption index over its side bias index. Bottom left apex: 0% nicotine consumption (0%); Bottom right apex: 100% nicotine consumption (100%); Top apex: 100% side preference (Side biased, i.e. mice that never switch side). Small dots correspond to the habituation period (water vs. water) while bigger dots correspond to the condition with 200 µg/ml of nicotine in one bottle. Note how all avoider mice end up in the bottom left apex (0% nicotine consumption) at the end of the task. **C.** Average distance from the three apices for each condition in the task (0, 10, 50, 100 and 200 µg/ml of nicotine, color-coded from light to dark), for avoiders and non-avoiders. Only

avoider mice significantly changed their drinking strategy as nicotine concentration increased (paired Mann-Whitney test with Holm-Bonferroni correction). **D.** Average percent nicotine consumption for avoider and non-avoider mice for each concentration of nicotine. The 50 µg/ml nicotine solution was presented a second time to the mice, at the end of the session, for four days. (Mann-Whitney test, Holm-Bonferroni correction, p(50 Nic) = 0.04, p(100 Nic) = 0.005, p(200 Nic) = 0.004). **E.** Representative (left) and average (right) currents recorded in voltage-clamp mode (-60 mV) from IPN neurons of non-avoiders (blue, n = 142 neurons from 18 mice, I = -194 $\pm$ 15 pA) and avoiders (red, n = 65 neurons from 8 mice, I = -297 $\pm$ 30 pA) following a puff application of nicotine (30 µM, 200 ms). Avoiders presented greater nicotine-evoked currents than non-avoiders (Mann-Whitney test, p = 1.4e-11). **G.** Correlation between the dose consumed (log scale, over the last 24 hours prior to the recording) and the averaged nicotine evoked-current per mouse (-pA). In all figure panels avoiders are depicted in pinkish-orange and non-avoiders in blue. *** p<0.001, ** p<0.01, * p<0.05.

Figure 3: Chronic nicotine treatment altered both nAChR expression levels in

the IPN and nicotine intake in WT mice. A. Top, passive nicotine treatment protocol.

Mice were implanted subcutaneously with an osmotic minipump (MP) that continuously

delivers 10 mg/kg/d of nicotine. After 4 weeks of treatment, nicotine-evoked responses

in IPN neurons were recorded in whole-cell voltage-clamp mode (-60 mV) from IPN

slices. Bottom, representative recordings (left) and average current amplitudes (right)

following a puff application of nicotine (30 µM, 200 ms) in IPN neurons of mice treated

with either saline (n = 24 neurons from 3 mice, I = -401 ± 59 pA) or nicotine (n = 34

neurons from 4 mice, I = -202 ± 37 pA). Nicotine treatment reduced the amplitude of

nicotine-evoked currents in IPN neurons (Mann-Whitney test, p = 0.001). **B.** *In vivo*

juxtacellular recordings of nicotine-evoked responses in IPN neurons of saline- and nicotine-treated animals. Top, representative electrophysiological recording of an IPN neuron, during an i.v. injection of nicotine (30 µg/kg). Middle, post-recording identification of neurobiotin-labeled IPN neurons by immunofluorescence. Bottom, average time course and average amplitude of the change in firing frequency from baseline after an i.v. injection of saline and nicotine (30 µg/kg), for IPN neurons from saline- and nicotine-treated mice. Right, firing rate variation from baseline induced by nicotine or saline injection in IPN neurons from saline- (n = 6) or nicotine-treated (n = 13) animals. Responses were decreased by chronic exposure to nicotine (p = 0.035, Mann-Whitney test). All neurons were confirmed to be located within the IPN using juxtacellular labeling with neurobiotin. **C.** Top, modified two-bottle choice protocol used to evaluate the impact of a long-term exposure to nicotine on drug intake. Mice were implanted subcutaneously with a minipump that delivered 10 mg/kg/d of nicotine continuously, for 20 days before performing the modified two-bottle choice task. After four days of water vs. water habituation, mice were directly exposed to a high concentration of nicotine (100 µg/ml). Bottom, percent nicotine consumption and nicotine intake at 0 and 100 µg/ml of nicotine, for mice under a chronic treatment of nicotine or saline. The saline-treated group displayed a decrease in percent nicotine consumption (n = 23, from 47.3 $\pm$ 2.0% to 23.0 $\pm$ 3.4%, p = 1.7e-05, Mann-Whitney paired test), but not the the nicotine-treated group (n = 25, from 48.9 $\pm$ 2.4 to 41.2 $\pm$ 5.5%, p = 0.16, Mann-Whitney paired test). Overall, the saline-treated group displayed a lower percent nicotine consumption (p = 0.003, Mann Whitney) and lower nicotine intake than the nicotine-treated group (p = 0.004, Mann-Whitney). **D.** Pseudo-ternary diagrams representing each saline- and nicotine-treated mouse for its nicotine consumption index over its side bias index. Small dots correspond to the habituation

period (water vs. water) and bigger dots to the condition with 100 µg/ml of nicotine in

one bottle. **E.** Average distance from each apex in the water vs. water (top) and water

vs. nicotine 100 µg/ml conditions (bottom). Saline-treated, but not nicotine-treated mice

developed a strategy to avoid nicotine (p_{Sacc} = 0.013, p_{Side} = 0.27 p_{Nic} = 0.013, Mann-

Whitney test with Holm-Bonferroni correction). In all figure panels nicotine-treated

animals are displayed in red and saline-treated (control) animals in grey.

FIGURE 4

Figure 4: β4-containing nicotinic receptors are essential for triggering nicotine aversion in mice. A. Left, average percent nicotine consumption in WT and β4$^{-/-}$ mice for each concentration of nicotine in the two-bottle choice task. WT mice had lower percent nicotine consumption than β4$^{-/-}$ mice (Mann-Whitney test with Holm-Bonferroni correction). WT mice decreased their percent nicotine consumption throughout the task (Friedman test, n = 35, df = 4, p < 0.001 and Mann-Whitney post-hoc test with Holm-Bonferroni correction) while β4$^{-/-}$ mice displayed a stable percent nicotine

consumption (Friedman test, n = 13, df = 4, p = 0.11). Right, average nicotine intake

(mg/kg/d) in β4$^{-/-}$ and WT mice for the different concentrations of nicotine (Friedman

test, n = 35, df = 3, p < 0.001 and Mann-Whitney post-hoc test with Holm-Bonferroni

correction). β4$^{-/-}$ mice consumed more nicotine than WT mice (Mann-Whitney test). **B.**

Ternary diagram representing each WT and β4$^{-/-}$ individual for its nicotine consumption

index over its side bias index. Small dots correspond to the habituation period (water

vs. water) and bigger dots to the condition with 200 µg/ml of nicotine in one bottle.

Inserts: pie charts illustrating the proportion of avoiders (A, light grey) and non-avoiders

(N, dark grey) for each genotype at the end of the task. Note the absence of avoiders

in β4$^{-/-}$ mice. **C.** Average distance from each apex at 0, 10, 50, 100 and 200 µg/ml of

nicotine (paired Mann-Whitney test with Holm-Bonferroni correction, p(Sacc 0-200) =

0.0002, p(Sacc 0-100) = 0.03; p(Nic 0-200) = 0.0002, p(Sacc 0-100) = 0.03). **D.** Left,

representative currents following a puff application of nicotine (30 µM, 200 ms) in IPN

neurons from naive WT and β4$^{-/-}$ mice, or from saline-treated (orange) and nicotine-

treated (dark orange) β4$^{-/-}$ mice. Right, average nicotine-evoked currents recorded in

IPN neurons from naïve WT (n = 32 neurons from 5 mice, I = -370 $\pm$ 61 pA) and β4$^{-/-}$

(n = 27 neurons from 4 mice, I = -83 $\pm$ 13 pA) mice, and from β4$^{-/-}$ chronically treated

with either saline (Sal, n = 30 neurons from 6 mice, I = -72 $\pm$ 11 pA) or nicotine (Nic, n

= 31 neurons from 5 mice, I = -123 $\pm$ 21 pA). β4$^{-/-}$ mice presented a large decrease in

nicotine-evoked currents (Mann-Whitney test, p = 4e-05). Nicotine treatment did not

alter nicotine-evoked currents in IPN neurons of β4$^{-/-}$ mice (Mann-Whitney test, p =

0.15). **E.** Juxtacellular recordings of nicotine-evoked responses in IPN neurons in naive

WT and β4$^{-/-}$ mice. Top, representative example of the variation in firing frequency of

an IPN neuron, following repeated i.v. injections of nicotine at 7.5, 15 and 30 µg/kg, in

WT and β4$^{-/-}$ mice. Bottom left, dose-dependent change in firing rate from baseline

following i.v. injections of nicotine ($p < 0.05$). All recorded neurons were neurobiotin-

labelled to confirm their location within the IPN. Bottom right, average nicotine-evoked

responses at 30 µg/kg of nicotine in IPN neurons from WT and β4$^{-/-}$ mice. Insert,

average amplitude of the change in firing frequency from baseline after an i.v. injection

of saline and nicotine (30µg/kg), for IPN neurons from WT (n = 18) and β4$^{-/-}$ (n = 7)

mice. Nicotine-induced responses were smaller in β4$^{-/-}$ than in WT mice ($p < 0.001$,

Mann Whitney test). In all figure panels WT animals are depicted in grey and β4$^{-/-}$ mice

in yellow. *** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$.

FIGURE 5

Figure 5: β4-containing nAChRs of the IPN are involved in the control of nicotine consumption and in aversion to nicotine in mice. A. Protocol: stereotaxic transduction of the β4 subunit together with GFP (or GFP alone in control mice) in the IPN of β4$^{-/-}$ mice, and subsequent two-bottle choice task. Right: coronal section highlighting proper viral transduction of lenti-pGK-β4-IRES-GFP in the IPN. **B.** Validation of the re-expression using whole-cell patch-clamp recordings. Representative currents and average responses following a puff application of nicotine (30 μM, 200 ms) on IPN neurons from β4$^{-/-}$ mice transduced in the IPN with either lenti-pGK-β4-IRES-GFP (KO-β4^{IPN}, n = 7 neurons from 2 mice, I = -362 $\pm$ 95 pA) or lenti-pGK-GFP (KO-GFP^{IPN}, n = 7 neurons from 1 mouse, I = -71 $\pm$ 18 pA; Mann-Whitney test, p = 0.004). **C.** Average nicotine intake was lower in KO-β4^{IPN} than in KO-GFP^{IPN} (two-way repeated measure; ANOVA: genotype x dose interaction, $F_{[3, 99]}$ = 6.3, ***p < 0.001; main effect of dose, $F_{[3, 99]}$ = 69.1 ***p < 0.001, effect of genotype, $F_{[1, 33]}$ = 6.637, *p = 0.015). **D.** Ternary diagram representing each β4$^{-/-}$ mouse, transduced with either β4 or GFP, and illustrating its nicotine consumption index over its side bias index. Small dots correspond to the habituation period (water vs. water) and bigger dots to the condition with 200 μg/ml of nicotine in one bottle. Inserts: pie charts

illustrating the proportion of avoiders (A, light grey) and non-avoiders (N, dark grey) for

each condition at the end of the task. **E.** Average distance from each apex during the

two-bottle choice task at 0 and 200 µg/ml of nicotine, for KO-β4^{IPN} and KO-GFP^{IPN}

mice (p_{Sacc} = 0.007, p_{Nic} = 0.008, Mann-Whitney test with Holm-Bonferroni correction).

In all figure panels KO-β4^{IPN} mice are depicted in red and KO-GFP^{IPN} mice (controls)

in green. *** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$.

1002