The costs and benefits of dispersal in small populations

Jitka Polechová[1,*],

1 University of Vienna, Department of Mathematics

∗ E-mail: jitka@univie.ac.at

ABSTRACT

Dispersal has three major effects on adaptation. First, the gene flow mixes alleles adapted to different environments, potentially hindering (swamping) adaptation. Second, it inflates genetic variance: this aids adaptation to spatially (and temporally) varying environments but if selection is hard, it lowers the mean fitness of the population. Third, neighbourhood size, which determines how weak genetic drift is, increases with dispersal – when genetic drift is strong, increase of neighbourhood size with dispersal aids adaptation. In this note I focus on the role of dispersal in environments which change smoothly across space, and when local populations are quite small such that genetic drift has a significant effect. Using individual-based simulations, I show that in small populations, even leptokurtic dispersal benefits adaptation, by reducing the power of genetic drift. This has implications for management of small marginal populations: increased gene flow appears beneficial as long as adaptations involves a quantitative, rather than a discrete, trait. However, heavily leptokurtic dispersal will swamp continuous adaptation along steep environmental gradients so that only patches of locally adapted subpopulations remain.

keywords:

adaptation, environmental gradients, genetic drift, species' range, spatial structure, evolution, migration, gene flow

INTRODUCTION

The role of gene flow in preventing (swamping) local adaptation is well acknowledged (Bürger, 2013; Yeaman, 2015). Dispersal between discrete niches will swamp adaptation when the negative trade-offs in fitness in the different niches are asymmetric, unless selection and/or niche-preference are strong (Barton, 2010; Bulmer, 1972; Haldane, 1930; Lenormand, 2002; Smith, 1970). In the absence of niche preference, often only one type remains unless the selective advantage is symmetric. In his classic study, Haldane (1956) asserted that even in continuous space, gene flow could swamp adaptation, even if the nascent asymmetry in population density was small: the gene flow leads to departure from the optimum, and the maladaptation translates (under hard selection, Christiansen 1975) to a lower density: through this positive (but detrimental) feedback, the asymmetry in population size would grow. The dynamics have been asserted to potentially form a stable species' range margin, an idea also popularised by Mayr (1963). This idea appears supported by studies of adaptation to distinct niches with source-sink dynamics, which may represent marginal habitats (Gomulkiewicz *et al.*, 1999; Kawecki *et al.*, 1997). Yet, swamping of adaption is not generally expected across continuously changing conditions: spatial clines in allele frequencies readily form and are stabilised by density gradients (dips), and also by evolution of linkage disequilibrium with other clines, effectively strengthening selection (Barton, 1986, 1983; Kruuk *et al.*, 1999).

Whether gene flow swamps or benefits adaptation across environments which vary gradually through space depends on further details. It has been shown that when genetic variance is fixed (or constrained to evolve very slowly), dispersal across environments can indeed swamp adaptation (Kirkpatrick & Barton, 1997). As dispersal load (the incurred fitness cost of dispersal across environments on mean fitness) increases, continuous adaptation fails when genetic variance (measured by the variance load) is too small. Yet, when genetic variance increases with gene flow, the increase of variance aids adaptation such that it remains continuous, as long as the local density stays above zero (Barton, 2001). In small populations, the third effect of dispersal becomes important: the increase of the neighbourhood size (Wright, 1931). It has been shown that in two-dimensional habitats, gene flow can thus – perhaps counterintuitively – facilitate adaptation to environmental gradients (Polechová, 2018). This is because neighbourhood size rises with the dispersal distance squared, while the dispersal load only rises with the dispersal distance (as long as the environment varies mainly along one spatial dimension, such as along altitudinal or latitudinal gradients).

Yet, the assumption of Gaussian dispersal kernel (assumed in the theory above) may have been quite

restrictive. It is conceivable that with a long-range component to dispersal (as opposed to just a Gaussian dispersal kernel), the swamping effect would quickly overwhelm the beneficial effect of reducing genetic drift. Yet, there has been very little exploration of the the effect of long-range dispersal on adaptation, even in the absence of temporal change. On the other end of the spectrum, with uniform dispersal within the whole structured population – such as in the island model – differential adaptation is readily prevented when the migration rate and the span of the environments is large enough, aided by asymmetry in selection and/or relative size of the niches (Szép *et al.*, 2021).

MODEL AND RESULTS

I focus on evolution of a species' range in a two-dimensional habitat, assuming stabilising selection towards an environmental gradient b, which is stable through time. The trait z under selection is determined by a large number of additive loci (set to 500), and stabilising selection takes the form of $r_g(z) = -(z - bx)^2/(2V_s)$. Population growth is density-dependent $r_e(N) = r_m(1 - N/K)$, where K gives the carrying capacity, and N the local population density. The mean fitness is $\bar{r}(\bar{z}, N) = r_e(N) + \bar{r}_g(\bar{z}) = r_m(1 - N/K) - (\bar{z} - \theta)^2/(2V_s) - V_P/(2V_s)$. The term $V_P/(2V_s)$ gives the load due to phenotypic variance $V_P = V_G + V_E$, V_s is the variance of stabilising selection. In this model, one can use $V_G \equiv V_P$ without a loss of generality: the loss of fitness due to environmental variance V_E can be included in $r_m^* = r_m - V_E/(2V_s)$, where r_m^*, r_m is the maximum per capita growth rate.

I assume the spatial optimum varies along one dimension (X); but it is constant along the other dimension (Y). The demes form a two-dimensional lattice of 100 demes in both dimensions. Mating is local within demes after migration and selection: the mating pool is given by (haploid) neighbourhood size $\mathcal{N} = 2\pi\sigma^2 N$. The trait is additive, and each bi-allelic haploid locus can contribute to trait value by the effect of $\alpha = 0.2$. The population starts well adapted in the central half of the habitat: the optimum is matched by a series of clines spaced α/b apart, with cline width $w_{pq} = 4\sigma/\sqrt{2s}$, where $s = \alpha^2/(2V_s)$. In contrast to the model in (Polechová, 2018), dispersal is modelled as a mixture of two Gaussians, with two different variances σ_1 and σ_2, one with much higher than the other: $(1 - a)\phi_N(0, \sigma_1) + a\phi_N(0, \sigma_2)$. As a default, $\sigma_2 = 20\sigma_1$. The sizeable asymmetry means that the proportion of long-range dispersal increases with a unless a is very close to 1, when all dispersal comes from the wider Gaussian.

While large populations will maintain clinal variation and gradually expand if the environment varies smoothly (Barton, 2001), adaptation across steep environmental gradients can fail abruptly when genetic

drift is strong. As clinal variation dissipates, continuous adaptation fails and the species range collapses and/or fragments: each subpopulation is only adapted to a distinct optimum (Polechová, 2018). In contrast, continuous adaptation means clines form across the range, facilitating steady range expansion. Figure 1 shows that in small populations, adaptation across steep environmental gradients is rescued by increasing Gaussian dispersal (A). When the proportion of the long-range dispersal is small, adaptation is further facilitated, and range expands faster (B,C). However, as the proportion of long-range dispersal increases (D), gene flow swamps adaptation to steep environmental gradients, and continuous adaptation is no longer possible. This results in a population consisting of locally adapted subpopulations, with large gaps between them. This is demonstrated in Figure 2, which shows the spatial distribution of genetic variance as long-range dispersal increases. In the absence of long-range dispersal, genetic drift is too high for the spatial gradient, very few clines underlying adaptation are maintained, and local genetic variance is (mostly) low. As dispersal increases, genetic drift weakens and adaptation becomes easier: this is quite robust even when dispersal is leptokurtic (B,C). Eventually, though, increasing long-range dispersal swamps continuous adaptation, and the resulting population consists of locally adapted subpopulations spaced-apart by the long-range dispersal (D). Note that even broader dispersal can push the whole population to extinction as the mean fitness drops to zero globally. The parameters here are chosen so that this extreme is avoided; in general, it is dependent on the specific form of density-regulation.

DISCUSSION

The role of dispersal in spatially structured populations is complex as it can both swamp and facilitate adaptation. Additionally, in small population, the effect of reducing genetic drift becomes important Willi *et al.* (2006). While most commonly is discussed the extent of swamping, which can still be balanced by selection (Bürger, 2013; Holt & Barfield, 2011; Yeaman, 2015), number of studies have found that locally favourable alleles can be most readily established with intermediate migration, supplying the new variants (Gomulkiewicz *et al.*, 1999; Uecker *et al.*, 2014). This can be crucial for adaptation to both spatial and temporal variation. It has been hypothesised that for example in trees, variants brought in by long-range pollen dispersal may aid adaptation, notably to temporal change (Kremer *et al.*, 2012). The mixture of two dispersal kernels used this paper can be seen as a model applicable to such a system, with two modes of dispersal: one more local (seeds), and one more global (pollen). It shows that gene flow in continuously heterogeneous environments does not necessarily swamp adaptation even if dispersal is leptokurtic. In small populations, the additional benefit of reducing the power of genetic drift is more important. This can be relevant for management of small marginal populations: the oft-cited result that gene flow swamps adaptation in marginal populations often does not hold when both genetic variance evolves and genetic drift is important (Kirkpatrick & Barton, 1997). Interestingly, this is in line with a recent review which has found little evidence of gene flow swamping adaptation at the range margins (Kottler *et al.*, 2021).

It is to be noted, though, that there is no quantitative meaning of a particular percentage of long-range dispersal in this model: notably, if the Gaussians were more similar, genetic drift would be reduced more effectively. There are several further limitations to the study: adaptation occurs via many loci of uniform and small effect, and hence it assumes weak selection per locus. It is possible that relaxation of this assumption would make pockets of adaptation more robust to the combined effect of genetic drift and swamping by gene flow. Secondly, it simulates small neighbourhoods, where genetic drift is strong and is thus much better suited to addressing adaptation at range margins than at central parts of a species' range. While long-range dispersal may still be beneficial in large populations by bringing in adaptive variants, such an effect is not addressed in this note.

REFERENCES

Barton, N. (1986). The maintenance of polygenic variation through a balance between mutation and stabilizing selection. *Genetical Research*, 47, 209–216.

Barton, N. (2001). Adaptation at the edge of a species' range. In: *Integrating ecology and evolution in a spatial context* (eds. Silvertown, J. & Antonovics, J.). Blackwell, vol. 14, pp. 365–392.

Barton, N. (2010). What role does natural selection play in speciation? *Philosophical Transactions of the Royal Society B: Biological Sciences*, 365, 1825–1840.

Barton, N.H. (1983). Multilocus clines. *Evolution*, pp. 454–471.

Bulmer, M. (1972). Multiple niche polymorphism. *The American Naturalist*, 106, 254–257.

Bürger, R. (2013). A survey on migration-selection models in population genetics. *arXiv preprint arXiv:1309.2576*.

Christiansen, F.B. (1975). Hard and soft selection in a subdivided population. *American Naturalist*, 109, 11–16.

Gomulkiewicz, R., Holt, R.D. & Barfield, M. (1999). The effects of density dependence and immigration on local adaptation and niche evolution in a black-hole sink environment. *Theoretical Population Biology*, 55, 283–296.

Haldane, J.B.S. (1930). A mathematical theory of natural and artificial selection.(part vi, isolation.). In: *Mathematical proceedings of the Cambridge philosophical society*. Cambridge University Press, vol. 26, pp. 220–230.

Haldane, J.B.S. (1956). The relation between density regulation and natural selection. *Proceedings of the Royal Society of London, B: Biological Sciences*, 145, 306–308.

Holt, R.D. & Barfield, M. (2011). Theoretical perspectives on the statics and dynamics of species' borders in patchy environments. *The American Naturalist*, 178, S6–S25.

Kawecki, T., Barton, N. & Fry, J. (1997). Mutational collapse of fitness in marginal habitats and the evolution of ecological specialisation. *Journal of Evolutionary Biology*, 10, 407–429.

Kirkpatrick, M. & Barton, N. (1997). Evolution of a species' range. *The American Naturalist*, 150, 1–23.

Kottler, E.J., Dickman, E.E., Sexton, J.P., Emery, N.C. & Franks, S.J. (2021). Draining the swamping hypothesis: little evidence that gene flow reduces fitness at range edges. *Trends in Ecology & Evolution*.

Kremer, A., Ronce, O., Robledo-Arnuncio, J.J., Guillaume, F., Bohrer, G., Nathan, R., Bridle, J.R., Gomulkiewicz, R., Klein, E.K., Ritland, K. *et al.* (2012). Long-distance gene flow and adaptation of forest trees to rapid climate change. *Ecology letters*, 15, 378–392.

Kruuk, L., Baird, S., Gale, K.S. & Barton, N. (1999). A comparison of multilocus clines maintained by environmental adaptation or by selection against hybrids. *Genetics*, 153, 1959–1971.

Lenormand, T. (2002). Gene flow and the limits to natural selection. *Trends in Ecology & Evolution*, 17, 183–189.

Mayr, E. (1963). *Animal species and evolution*. Harvard University Press.

Polechová, J. (2018). Is the sky the limit? on the expansion threshold of a species' range. *PLoS Biology*.

Smith, J.M. (1970). Genetic polymorphism in a varied environment. *The American Naturalist*, 104, 487–490.

Szép, E., Sachdeva, H. & Barton, N.H. (2021). Polygenic local adaptation in metapopulations: A stochastic eco-evolutionary model. *Evolution*, 75, 1030–1045.

Uecker, H., Otto, S.P. & Hermisson, J. (2014). Evolutionary rescue in structured populations. *The American Naturalist*, 183, E17–E35.

Willi, Y., Van Buskirk, J. & Hoffmann, A.A. (2006). Limits to the adaptive potential of small populations. *Annu. Rev. Ecol. Evol. Syst.*, 37, 433–458.

Wright, S. (1931). Evolution in Mendelian populations. *Genetics*, 16, 97–159.

Yeaman, S. (2015). Local adaptation by alleles of small effect. *The American Naturalist*, 186, S74–S89.

Figure 1: **Increasing long-range dispersal first aids and then swamps adaptation across steep gradients**. (A) In small populations, Gaussian dispersal aids adaptation to spatially varying optimum as the effect on reducing genetic drift is bigger than the cost of swamping: this allows the population to expand continuously along a gradient. Dispersal still aids continuous adaptation when a minor proportion disperses much further, so that the dispersal kernel is leptokurtic (B,C): with 5% resp. 25% of long-range dispersal. However, when the extent of long-range migration increases further (D, 50%), gene flow across steep gradients ($b > 0.5$) starts swamping adaptation. Dashed line (A) evaluates the expansion threshold under Gaussian dispersal $\mathcal{N} \gtrless 6.3\,B + 0.56$, where the effective environmental gradient is $B = b\sigma/(r^* \sqrt{2V_s})$ and the neighbourhood size $\mathcal{N} = 4\pi N_e \sigma^2 = 2\pi N \sigma^2$ (Polechová, 2018). It is shown as a faint dotted line in pictures (B-D) for a reference; recalculation of the approximation of the threshold using the joint variance overestimates the rescue effect for longe-range dispersal (marginally out of the range for (B)). Parameters: local carrying capacity $K = 4r_m/r_*$, rate of return to equilibrium $r_* = r_m - b\sigma/(2\sqrt{V_s}) = 1$, width of stabilising selection $V_s = 1/2$, mutation rate $\mu = 10^{-6}$, habitat width 100 demes along spatial gradient, 100 demes in the neutral direction, along which the optimum is fixed. The colour shows the rate of expansion (blue and purple hues) or indicates past or ongoing contraction (orange and red hues); black triangles denote populations which went extinct within 500 generations.

Figure 2: **Illustration of spatial distribution of genetic variance for steep gradients, with increasing long-range dispersal - as in Fig. 1.** (A) with only local dispersal, neighbourhood size is small and genetic drift overwhelms adaptation to steep environmental gradient: clines only form sparsely. The blue contour line depicts genetic variance of $V_G = b\sigma\sqrt{V_s}$, which would be maintained in the absence of genetic drift with Gaussian dispersal kernel $\phi_N(0, \sigma)$ (Barton, 2001). Increasing long-range dispersal weakens the genetic drift, at first facilitating continuous adaptation across the species' range (B,C). With strongly leptokurtic dispersal, however, continuous adaptation is swamped around singular, locally adapted populations (D). Because the environmental optimum does not change along the Y-dimension, locally adapted sub-populations form stripes: gene flow is only realized along the neutral dimension. Parameters as in Fig. 1, $b = 0.7$; local dispersal $\sim \phi_N(0, \sigma_1 = 0.2)$, long-range dispersal $\sim \phi_N(0, \sigma_2 = 4)$.