The hyperpolarization-activated current shifts the dynamic range of a voltage-dependent electrical synapse

Wolfgang Stein[1*], Margaret L. DeMaegd[1+], Lena Yolanda Braun[1], Andrés Vidal-Gadea[1],
Allison L. Harris[2], Carola Städele[1#]

[1]School of Biological Sciences, Illinois State University. [2]Dept. of Physics, Illinois State University, Normal, IL 61790, USA. [#]present address: Dept. of Neuro-and Sensory Physiology, Univ. Med. Ctr. Göttingen, Univ. of Göttingen, Göttingen, Germany. [+]present address: New York Univ., New York, NY.

* corresponding author: Wolfgang Stein, wstein@neurobiologie.de

Running title: I_h modulates electrical synapse voltage dependence.

Acknowledgments: We would like to thank Alexandra Yarger, Daniel Nuccio, Chris Goldsmith, Doug Schuweiler, Amanda Smith, Bangxia Suo and Dana Tilley for help on early modeling for this project, and their thoughts on voltage-dependent synapses. We would also like to thank Ulm University for their support of LYB's research at Illinois State University. Supported by NSF IOS 1755098, NSF IOS 1354932, NSF 1828136, DFG STE 937/8-1 and STE 937/9-1.

Abstract.

Like their chemical counterparts, electrical synapses show complex dynamics such as rectification and voltage dependence that interact with other electrical processes in neurons. The consequences arising from these interactions for the electrical behavior of the synapse, and the dynamics they create, remain largely unexplored. Using a voltage-dependent electrical synapse between a descending modulatory projection neuron (MCN1) and a motor neuron (LG) in the crustacean stomatogastric ganglion, we find that the influence of the hyperpolarization-activated inward current (I_h) is critical to the function of the electrical synapse. When we blocked I_h with CsCl, the voltage dependence of the electrical synapse shifted by 18.7 mV to more hyperpolarized voltages, placing the dynamic range of the electrical synapse outside of the range of voltages used by the LG motor neuron (-60.2 mV – -44.9 mV). With dual electrode current- and voltage-clamp recordings, we demonstrate that this voltage shift is due to a sustained effect of I_h on the presynaptic MCN1 axon terminal membrane potential. I_h-induced depolarization of the axon terminal membrane potential increased the electrical postsynaptic potentials and currents. With I_h present, the axon terminal resting membrane potential depolarized, shifting the dynamic range of the electrical synapse towards the functional range of the motor neuron. We thus demonstrate that the function of an electrical synapse is critically influenced by a voltage-dependent ionic current (I_h).

Keywords. Central pattern generation, gap junction, electrical synapse, modulation, stomatogastric.

New & Noteworthy.

Electrical synaptic transmission and voltage-gated ionic currents are often studied independently from one another, despite mounting evidence that their interactions can alter synaptic behavior. We show that the hyperpolarization-activated inward ionic current shifts the voltage dependence of an electrical synapse through its depolarizing effect on the membrane potential, enabling it to lie within the functional membrane potential range of a motor neuron. Thus, the electrical synapse's function critically depends on the voltage-gated ionic current.

Introduction

Electrical synapses are ubiquitous throughout the nervous system and have been described in many vertebrate and invertebrate systems. Non-rectifying electrical synapses often support neuronal synchrony between neurons of similar functional and anatomical or biochemical profiles through bilateral current flow [1]. In contrast, rectifying synapses are usually found in feedforward neuronal circuits. Their effects are primarily unilateral and current flows preferentially from the presynaptic to the postsynaptic cell. Electrical synapses show complex dynamics, including facilitation and depression, and electrical synapse conductance is subject to alterations by a host of influences, including temperature and pH, as well as neuromodulators and hormones [1, 2]. Electrical synapses also interact with other electrical processes in neurons, creating emergent properties that can alter or even determine the dynamics of the whole system [3]. These effects also work in the opposite direction, with intrinsic neuronal properties, such as voltage-gated ion channels, altering the dynamics of the electrical coupling.

However, our understanding of the dynamics that such interactions create, and the consequences arising from these interactions for the electrical behavior of the synapse, is still in its infancy.

Included among the currents influencing electrical synaptic transmission are ionic currents, such as persistent Na^+ currents and the hyperpolarization-activated inward current (I_h), which have slow inactivation kinetics. This makes them particularly conducive to changing the properties of electrical synaptic transmission, for example through changes in the extrajunctional membrane resistance or the postsynaptic membrane potential [1, 2]. For instance, the persistent Na^+ current depolarizes postsynaptic cerebellar Golgi neurons, which causes a switch in the sign of the electrical synapse from predominantly hyperpolarizing to predominantly depolarizing [4, 5]. I_h has also been implicated in altering electrical synaptic transmission, although examples are less prevalent. In neurons of the snail *Helisoma*, for example, the modulation of electrical synapses between identical neurons is mediated by an I_h-mediated decrease in extra-junctional membrane resistance [6].

One system that is ideal for studying the effects of I_h on electrical synaptic transmission is the crustacean stomatogastric nervous system. Here, several electrical synapses and their effects on the behavior of the motor circuits in the stomatogastric ganglion (STG) have been characterized. In the crab, *Cancer borealis*, the descending modulatory projection neuron 1 (MCN1) innervates the lateral gastric (LG) motor neuron in the STG's gastric mill central pattern generator via both chemical and electrical synaptic connections [7, 8]. MCN1's chemical synapse provides slow neuropeptidergic excitation and enables LG to burst and produce rhythmic activity. The electrical synapse contributes to the maintenance of the LG burst [9]. Thus, both chemical and electrical synapses are crucial components required for LG's behavior. The electrical synapse shows a peculiar voltage dependence. At rest, each MCN1 action potential elicits a small sub-threshold electrical postsynaptic potential (ePSPs) in LG (Fig. 1A, black arrows). However, when LG depolarizes, these ePSPs increase in amplitude. This voltage dependence appears to be a property of the electrical synapse itself rather than being due to pre- or postsynaptic influences.

While the basic description of the voltage dependence of the MCN1-LG ePSPs has been available for over 25 years now, interactions with (and modulation by) other intrinsic electrical properties of MCN1 and LG are less understood. Here, by quantifying the voltage dependence of the ePSPs and measuring electrical synaptic currents, we show that the ePSPs are not just modulated by the hyperpolarization-activated inward current (I_h), but that this modulation is critical their function. Without I_h modulation, the voltage dependence of the electrical synapse was shifted below physiologically relevant values. Modulation by I_h shifted the dynamic range of the voltage dependence to more positive voltages to match LG's membrane potential. We demonstrate that this voltage shift is due to a sustained depolarization of the MCN1 axon terminal membrane potential that is caused by the presence of I_h.

Methods

Experimental animals. Adult male Jonah crabs (*Cancer borealis*) were obtained from "The Fresh Lobster Company" (Boston, MA, USA) and kept in aerated sea water tanks (~1.025 g / cm^3 salinity, artificial sea salt, AquaMed, USA). The tanks were kept at a temperature of 10 to 12°C with a 12-hour dark/light cycle. All experiments were performed following the "Principles of animal care" publication No. 86-23, revised in 1985 by the National Institute of Health.

Dissection. Animals were cold anesthetized on ice for 30 to 40 minutes before dissection. The dissection was carried out following published protocols [10]. In short, the stomach was removed from the animal and the STNS was dissected under a stereomicroscope. During the dissection, the nervous system was constantly kept at a low temperature by rinsing it with 4°C cold saline as appropriate. During experiments a temperature between 10 and 13°C was maintained by superfusing chilled saline.

Chemicals: Cancer borealis physiological saline consisted of (in mM, pH 7.4 - 76). KCl 11, NaCl 440, CaCl$_2$*2H$_2$O 13, MgCl$_2$*6H$_2$0 26, Trizma base 11.2, Maleic acid 5.1 (all from Sigma Aldrich, USA) dissolved in ultrapure water (18.3MOhm). In some experiments, saline with low Calcium concentration (low Ca^{2+}-saline) was used to block Calcium channels and synaptic release. Low Ca^{2+}- saline consisted of (in mM): KCl 11, NaCl 440, CaCl$_2$*2H$_2$O 0.1, MgCl$_2$*6H$_2$0 26, Trizma base 11.2, Maleic acid 5.1, MnCl$_2$ 12.9. Manganese was added after adjusting the pH to 7.4 – 7.6. To selectively block I$_h$, Cesium chloride (CsCl, Sigma-Aldrich, USA) was dissolved in low Ca^{2+}- saline (5 mM) before every experiment [11].

Extracellular recordings. Extracellular recordings followed established protocols [12]. In short, relevant nerves were electrically isolated with petroleum jelly wells. Subsequently, a recording electrode was placed inside the well while a reference electrode was placed in the grounded outside bath. The signal was differentially amplified using an A-M Systems Model 1700 amplifier (Carlsborg, WA, USA) and digitized with a Micro 1401 mkII (Cambridge Electronics Design, Cambridge, UK). Spike 2 (ver. 7, Cambridge Electronics Design, Cambridge, UK) was used to record and analyze the data. To detect the activity of the lateral gastric motor neuron (LG), the lateral gastric nerve (*lgn*) was recorded. It contains the LG axon.

Intracellular recordings. A dark field condenser (Nikon) was placed in-between LED and Petri dish to facilitate the visual contrast during intracellular recordings. Glass microelectrodes (borosilicate glass capillaries GB100TF-8P, Science Products GmbH, Hofheim, Germany) were pulled with the help of a microelectrode puller (Model P-97, Sutter Instrument, Novato, USA) and filled with a 0.6 M K$_2$SO$_4$ solution. Measurements were not affected by a potential mismatch in osmolarity of electrode solution and cytosol reported elsewhere [13] since the reported effects represent comparisons between precontrol, manipulated condition, and postcontrol. Recordings with obvious changes in membrane voltage or cell input resistance over the course of the experiment were discarded. Electrode resistance ranged between 15-25 MΩ. To impale cells, either a manual micro-manipulator (Leitz, Wetzlar, Germany) or several electrical manipulators (Sutter, Novato, USA) were used to position the electrodes above the STG. LG was identified by comparing its intracellularly recorded action potentials to those on the extracellular *lgn* recording. MCN1 was identified by responding to stimulation of its axons in the inferior esophageal nerve

(*ion*; see stimulation below). Signals were amplified through either Axoclamp-2B, Axoclamp 900B (Molecular Devises, Sunnyvalle, CA, USA) or NPI BA-1s (NPI Electronic GmbH, Tamm, Germany) amplifiers. For two-electrode current or voltage clamp recordings, two electrodes were simultaneously inserted into LG or the MCN1 axon terminal.

Neuronal stainings. To visualize the MCN1 axon terminals and the LG neuropil (Fig. 1B) we iontophoretically injected neurons with a fluorescent dye. Alexa Fluor 568 (MCN1 terminal) and Alexa Fluor 488 (LG) potassium salts (Thermo Fisher Scientific, Waltham, MA, USA) were diluted in 200 mM KCl and loaded in the electrode tip. After dye loading, electrodes were backfilled with 200 mM KCl. Negative current pulses of -10nA were applied for at least 10 minutes (2 seconds on, 0.5 seconds off) to load cells with dye.

Extracellular nerve stimulation. Petroleum jelly wells were placed around each *ion* to stimulate the axon of MCN1, following established protocols [14]. In some experiments the *ions* were cut close to the commissural ganglia that house the somata of the two MCN1 neurons. To activate MCN1, the remaining section of one *ion* still attached to the STG was stimulated. 1 Hz stimulation with a duration of 1 ms were applied elicit individual action potentials [15]. The long inter-pulse interval prevented a temporal summation of chemical PSPs in LG.

Model. We used a model consisting of two electrically coupled point neurons – one presynaptic and one postsynaptic. For each neuron, the membrane potential *Vm* is found from

$$C\frac{dVm}{dt} = -I_{leak} - I_h - I_{syn} + I_{inj}, \tag{1}$$

where C is the capacitance, I_L is the leak current, I_h is the hyperpolarization-activated inward current, I_{syn} is the synaptic current, and I_{inj} is the injected current.

The leak current is given by

$$I_L = g_L(Vm - E_L), \tag{2}$$

where g_L is the leak conductance and E_L is the leak equilibrium potential.

The hyperpolarization-activated inward current is given by

$$I_h = g_h m_{ht}(Vm - E_h), \tag{3}$$

where g_h is the conductance, m_{ht} is the time- and voltage-dependent activation function, and E_h is the equilibrium potential.

The time-dependence of the activation function is found by solving the differential equation

$$\frac{dm_{ht}}{dt} = \frac{(m_h - m_{ht})}{\tau_{mh}}, \tag{4}$$

where the voltage dependence of the activation function is given by

$$m_h = \left[1 + \exp\left(\frac{(Vm - V_{half})}{k_h}\right)\right]^{-1}. \tag{5}$$

The activation function time constant is τ_{mh} and the midpoint and growth rate of the logistic function are V_{half} and $1/k_h$ respectively.

The synaptic current is given by

$$I_{syn} = g_{syn}m_{syn}(Vm_{mpost} - Vm_{pre}),$$ (6)

where g_{syn} is the conductance, m_{syn} is the voltage-dependent activation function, and $Vm_{pre,post}$ are the pre- and postsynaptic membrane potentials, respectively.

The synaptic activation function is given by

$$m_{syn} = \left[1 + \exp\left(Vm_{post} - Vm_{pre} - V_{half}^{syn}\right)/k_{syn}\right],$$ (7)

where V_{half}^{syn} and $1/k_{syn}$ are the midpoint and growth rate respectively.

The injected current is given by a square pulse of duration t_{pulse} and height I_{pulse} for the presynaptic neuron. For the postsynaptic neuron, the injected current is a constant that is varied between -2 nA and +38 nA in the simulation to achieve different membrane potentials.

All model parameters are listed in Table 1

Model Parameter	Presynaptic	Postsynaptic
C	1 nF	1 nF
g_L	100 nS*	100 nS
E_L	- 80 mV*	- 60 mV
g_h	Variable	Variable
E_h	0 V	0 V
g_{syn}	0 S	40 nS
t_{pulse}	0.3 s	--
k_h	6 mV	6 mV
τ_{mh}	3 s	3 s
V_{half}^{syn}	10 mV	10 mV
k_{syn}	- 3 mV	- 3 mV
I_{pulse}	1 nA	--
V_{half}	- 80 mV	- 80 mV

Table 1. Model parameters. *default parameters used unless otherwise stated in the text.

The simulation was performed with Mathematica v12.1 using the NDSolve command for the differential equations. Neuronal behavior was observed for 30 seconds with the presynaptic pulse initiated at $t = 10$ s. The postsynaptic pulse was detected by using the NMaximize function in Mathematica for $10.2 \leq t \leq 11$ with a working precision of 10. Initial conditions for solving the differential equations are listed in Table 2.

Function	Initial Condition
Vm_{pre}	- 70 mV
Vm_{post}	- 60 mV
$m_{ht_{pre}}$	0.5
$m_{ht_{post}}$	0.5

Table 2. Initial conditions used to solve the differential equations in Eqs. (1) and (4).

Analyses and visualizations. Data were recorded and analyzed with Spike 2 script language and built-in functions. Measurements were transferred for spreadsheet analysis to Excel (Version 365 for Windows, Microsoft). Final figures were generated using Corel-Draw (Version X7 for Windows, Corel Corporation, Ottawa, ON, Canada). Statistical tests were carried out with SigmaPlot (Version 11 for Windows, Systat Software GmbH, Erkrath, Germany). Sigmoidal fits were generated using the SigmaPlot Nonlinear Regression - Dynamic Fitting function, using the equation f= a/(1+exp(-(x-x0)/b)) and 200 iterations. Here, x0 represents the midpoint of the sigmoidal fit that was used to compare between low Ca^{2+} saline and low Ca^{2+} saline with added CsCl (5mM). Only data with significant regression ($p<0.05$) were used. Individual results are given in the text or the figure legends. "N" denotes the number of animals tested whereas "n" indicates the number of repetitions used in an experiment.

Results.

The electrical synapse between MCN1 and LG is voltage-dependent.

MCN1 is part of a group of descending projection neurons located in the commissural ganglia (CoGs) [16]. There is one MCN1 neuron in each of the two CoGs and both project several centimeter-long axons to the STG via the inferior esophageal (*ion*s) and stomatogastric nerves (*stn*). After entering the STG neuropil (Fig. 1B), the MCN1 axons innervate pyloric and gastric mill neurons. While the concurrent activity of both MCN1 copies appears important to elicit robust gastric mill activity [17], the two MCN1 axons have identical postsynaptic targets. Each MCN1 axon terminal interacts via several direct synaptic connections with the core gastric mill CPG neuron LG (Fig. 1C). MCN1 excites LG chemically via *Cancer borealis* tachykinin-related peptide Ia (CabTRP Ia) [15]. The slow LG membrane potential depolarization resulting from this chemical excitation eventually activates spike generation in LG, allowing it to escape the inhibition from its antagonist, Interneuron 1 (Int1). LG also has a chemical feedback synapse onto the MCN1 axon terminal, which over time weakens the chemical excitatory input LG obtains from MCN1 [9]. In addition, the MCN1 axon terminal and LG are electrically coupled. MCN1 action potentials elicit subthreshold ePSPs that are readily observable in LG (Supplemental Figure S1A). Fig. 1A highlights several ePSPs during an ongoing gastric mill rhythm (red boxes, arrows). While the interactions between the chemical synapses allow LG to create its burst, the ePSPs contribute to burst maintenance by prolonging burst duration [9, 18]. The electrical synapse between the MCN1 axon terminal and LG appears unaffected by the presynaptic inhibition LG exerts on the axon, but the elicited ePSPs show an obvious voltage dependence: they grow in amplitude when LG depolarizes. Fig. 1A shows this during an ongoing gastric mill rhythm. When LG depolarizes and starts to burst, ePSPs increase in amplitude, prolonging the burst with action potentials riding on top of the ePSPs. This can be seen best by comparing ePSPs before the LG burst (black arrows) to those during and immediately after the burst (outlined arrows). When the LG burst ends due to a reduction of MCN1 neuropeptide release [19], the LG membrane potential hyperpolarizes and the amplitude of the ePSPs decreases again (Fig. 1A, gray arrows after the burst; [9]). The voltage dependence of the ePSPs can also be observed without a gastric mill rhythm, and when chemical release from MCN1 and chemical synapses from other STG neurons (such as Interneuron 1; Fig. 1C) are blocked. Fig. 1D shows averaged ePSPs obtained from an intracellular recording of LG's membrane potential during continuous 1 Hz stimulation of MCN1's axon (see Methods). No gastric mill rhythm was elicited, and chemical synaptic transmission was blocked using low Ca^{2+} saline. When LG was depolarized by current injection, ePSP amplitude increased. In contrast, when LG was hyperpolarized by a similar current injection, ePSP amplitude diminished below control levels. This suggests that the voltage-dependent changes in ePSP amplitude are caused by voltage-dependent alterations of the electrical synapse itself [9]. We noted, however, that the ePSP amplitude and voltage dependence appeared to be altered by I_h. Fig. 1E shows averaged ePSPs obtained from an intracellular recording of LG's membrane potential during continuous 1 Hz stimulation of MCN1's axon (see Methods). For all conditions shown, chemical synaptic transmission was blocked with low Ca^{2+} saline [20]. When we additionally blocked I_h by adding extracellular CsCl (5mM) [21], ePSP amplitudes increased. This suggested that I_h intrinsic to either MCN1's axon terminal or to LG influences the electrical synaptic transmission between them.

Figure 1. MCN1 elicits voltage-dependent electrical postsynaptic potentials in LG. A. Intracellular recording of LG membrane potential during ongoing gastric mill rhythm. In the LG interburst, ePSPs are small (black arrows). At the end of the LG burst, ePSPs are larger (outlined arrows). ePSP amplitudes slowly diminish after the burst (gray arrows). B. Staining of MCN1 axon terminals (red) and LG cell body and neuropil (green) in the STG. C. Schematic of MCN1 and LG synaptic connectivity. MCN1 chemically excites LG and Int1, which reciprocally inhibit each other. LG provides presynaptic inhibitory feedback to the MCN1 axon terminal. In addition, the MCN1 axon terminal is electrically coupled to LG. D. Time-triggered averages of ePSPs in the absence of a gastric mill rhythm, in low Ca^{2+} saline. Left: at resting membrane potential. Middle: during LG depolarization with 1.5 nA. Right: during LG hyperpolarization with -1.5 nA. Individual ePSPs were elicited through MCN1 axon stimulation with 1 Hz. Averages were calculated by aligning individual sweeps of LG membrane potentials to the MCN1 stimulation. E. ePSP amplitude increases when I_h is blocked. Time-triggered averages of MCN1 PSPs in LG. Individual ePSPs were elicited through MCN1 axon stimulation with 1 Hz. Left: in low Calcium saline. Middle: with I_h blocked by bath application of CsCl (5 mM). Right: washout.

The hyperpolarization-activated inward current (I_h) shifts the voltage dependence of the MCN1-LG electrical postsynaptic potentials.

To investigate I_h's influence, we compared the voltage dependence of the MCN1-LG ePSPs before and after blocking I_h. Figure 2A (left) shows original ePSP recordings at different LG membrane potentials. Individual MCN1 action potentials were elicited through 1 Hz stimulation of one *ion*, and chemical synapses were blocked with low Calcium saline. PSPs were recorded with an intracellular electrode in LG's

soma and LG membrane potential was altered with current injection through a second electrode. As expected, depolarizing LG increased ePSP amplitude, and hyperpolarizing LG decreased ePSP amplitude. We noted however that the change in ePSP amplitude was not instantaneous. Instead, it took several hundred ms (477 ± 281 ms, N=7) after a voltage step for the amplitudes to reach their new equilibrium (Supplemental Figure S1B).

Overall, we obtained a mostly sigmoidal relationship between membrane potential and ePSP amplitude (Fig. 2B, magenta), with ePSP amplitudes flattening out towards their smallest values around -70mV, and towards their maximum at around -40mV. The midpoint of the sigmoidal curve was -63.5 mV. Similar results were obtained in all tested animals. Fig. 2C (magenta) shows a summary of all experiments (N= 8 animals), with the averaged sigmoidal of all experiments, and their standard deviation. On average, the midpoint of the ePSP amplitude change was -62.7±3.3 mV (for calculation of the sigmoidal fits see Methods).

When we blocked I_h, ePSP amplitude generally increased, but most dramatically at voltages near the LG resting membrane potential (-75mV in this particular experiment). Fig. 2A (right) shows the resulting ePSP amplitudes for one experiment, and the resulting sigmoidal is shown in Fig. 2B (purple). In this experiment, the midpoint of the sigmoidal had shifted to – 86.7 mV and was thus more hyperpolarized than with I_h present. We found similar results in all experiments (Fig. 2C, purple). On average, the midpoint of ePSP amplitude change was now -81.4±9.0 mV (N=7, Mann-Whitney U-test, P=0.002, significantly different from no CsCl control), and amplitudes already approached their maximum around -70mV. Thus, the dynamic range of the voltage dependence of the electrical synapse appeared no longer within the functional range of the LG membrane potential. Indeed, when we measured the range of subthreshold membrane voltages LG exhibits during a gastric mill rhythm, we found that, on average, the resting membrane potential was -60.2 ± 2.3 mV, the most hyperpolarized membrane potential during an ongoing rhythm was -59.3 ± 2.0 mV, and the spike threshold was -44.9 ± 3.2 mV (N=14). We gathered these data in a different set of experiments, in regular saline without blockers to allow the gastric mill rhythm to occur. Comparing LG's subthreshold voltage range to the voltage dependence of the electrical synapse (Fig. 2C) revealed that 33% of the ePSP amplitude change occurred in that voltage range when I_h was present. In contrast, with I_h blocked, the amplitude changes occurred at more hyperpolarized membrane potentials, and less than 9% occurred within LG's functional voltage range. This suggests that I_h plays a critical role in keeping the dynamics of this synapses in the adequate voltage range.

Figure 2. Blocking I_h shifts the voltage dependence of the MCN1 ePSPS to more hyperpolarized voltages. A. Original traces of MCN1 ePSPs at various LG membrane potentials (-50 to -100 mV). In CsCl (right) ePSPs were larger, and only diminished at more hyperpolarized LG membrane potentials. Individual ePSPs were elicited through MCN1 axon stimulation with 1 Hz. B. Quantification of ePSP amplitudes of the experiment shown in (A). Multiple ePSPs were elicited at each membrane potential. Magenta: in low Calcium saline. Purple: with I_h blocked by CsCl. C. Quantification of all experiments. The voltage dependence of each experiment was fitted with a sigmoidal curve. Mean ± SD for all experiments are shown. Colored areas indicate the range of all observed data. Magenta: in low Calcium saline. Purple: with I_h blocked by CsCl. The gray box indicates the functional range of LG's membrane potential. Light gray indicates SD.

I_h can be detected in both LG and the MCN1 axon terminal.

While many STG neurons possess I_h, it has not been studied in detail in either LG or MCN1. As a first step, we used two-electrode current clamp to test if I_h was present. Fig. 3A shows an example recording in which LG was hyperpolarized through current injection for several second and then released from hyperpolarization. The sag potential (arrows in Fig. 3A) typical for I_h is clearly visible and slowly depolarized LG during the current step. After release from hyperpolarization, LG's membrane potential overshot the

resting membrane potential in a postinhibitory rebound, but only by a few mV. The recording shows small spikelets (2 mV and smaller) that ride on top of the rebound (asterisk in Fig. 3A). These spikelets were not LG action potentials. Instead, these were ePSPs originating from MCN1, which apparently went through its own rebound, elicited by the strong hyperpolarization of LG. In more than 10 LGs in which we caried out similar current injections, only one showed a rebound that elicited LG action potentials. This suggested that I_h was either weak or absent in LG, but present in the MCN1 terminal, and our current injection reached MCN1 through the gap junction. Fig. 3B shows a step protocol where LG was hyperpolarized in multiple steps. I_h was activated, but again only at very hyperpolarized membrane potentials (<-90 mV), using rather large injected currents (>-20 nA). Fig. 3C shows the same step protocol after CsCl application. Here, no sag potential or rebound could be detected anymore, confirming that indeed I_h was recorded. Figures 3D & 3E show the same scenario in the two-electrode voltage clamp with similar results. In control (low Calcium saline), I_h was detectable but absent in CsCl. This was consistent across experiments (N=7 for current clamp, N=11 for voltage clamp). We noted, however, that in all experiments, I_h was only activated at rather negative membrane potentials (see discussion).

To determine if I_h was present presynaptically, we recorded intracellularly from the MCN1 axon terminal. Some of these recordings were done with a single electrode due to the difficulty to find the MCN1 axon and impale it with two electrodes. Nevertheless, we found evidence that I_h is present in the MCN1 axon terminal. Fig. 3F shows an example of a two-electrode recording where we hyperpolarized the MCN1 axon terminal for several seconds. The membrane potential slowly depolarized during the current step. After the end of the current injection, MCN rebounded and fired several action potentials. Notably, these action potentials were generated in the axon terminal and were not reflections of possible postsynaptic action potentials in LG since no action potentials occurred on the extracellular LG recording (not shown). Fig. 3G shows a step protocol where MCN1 was hyperpolarized in a single electrode configuration. As such, currents, rather than membrane potentials, are given for the step protocol. Nevertheless, clear sag potentials were visible, and they occurred despite rather small current injections (4nA and smaller). In the presence of CsCl, these sag potentials as well as the rebound of the membrane potential after release of the current injection were absent (Fig. 3H), demonstrating that they were caused by I_h. This was the case in all preparations (N=6).

Figure 3. The hyperpolarization-activated inward current can be observed in LG and in the MCN1 axon terminal. A. Two-electrode current clamp recording of LG. When LG was hyperpolarized by current injection, the membrane potential showed a slow sag during which the voltage depolarized. After the end of the hyperpolarization, there was a small rebound as the membrane potential surpassed the resting membrane potential. No LG spikes were generated. *ePSPs elicited by MCN1 action potentials during the rebound. Arrows: sag potential. B. Step protocol in two-electrode current clamp in low Calcium saline. A voltage sag was visible below -90 mV. C. Step protocol in two-electrode current clamp in low Calcium saline, but with I_h blocked by CsCl. No voltage sag was detectable. D. Two-electrode voltage clamp recording with step protocol hyperpolarizing the LG membrane potential, in low Calcium saline. A small I_h was visible. E. Same step protocol in CsCl. I_h was no longer detectable.

F. Two-electrode current clamp recording of the MCN1 axon terminal. When the terminal was hyperpolarized by current injection, the membrane potential showed a sag during which the voltage depolarized. After the end of the hyperpolarization, MCN1 rebounded and produced several action potentials (*). G. Current step protocol to hyperpolarize the MCN1 axon terminal in low Calcium saline. A clear sag of the membrane potential was visible. H. Same protocol, but with I_h blocked by CsCl. The voltage sag and the rebound after the end of the voltage steps were absent.

I_h shifts the voltage dependence of an electrical synapse by depolarizing the presynaptic membrane voltage.

To address the question how I_h may affect the voltage dependence of the MCN1-LG synapse, we built a computational model of two neurons connected via a voltage-dependent electrical synapse. In this simplistic model (see Methods for details and default parameters used), each neuron consisted of only one compartment. The electrical synapse only passed current in one direction (from the presynaptic to the postsynaptic neuron). The voltage dependence was described by equation 7 (see Methods), which depends not just on the presynaptic membrane potential, but also on the postsynaptic membrane potential. This sigmoidal activation function resulted in a sigmoidal amplitude increase of the postsynaptic potentials when the postsynaptic membrane potential was depolarized (Fig. 4A, left: model output, Figs. 4B & 4C, red traces of sigmoidal response plot). Both neurons possessed passive membrane properties, but no active sodium or potassium currents (and therefore no action potentials) as those are not necessary to measure synaptic events. A single current pulse into the presynaptic neuron was used to elicit rapid changes in the membrane potential, simulating an action potential. We additionally implemented I_h, either in the presynaptic neuron or the postsynaptic neuron, or in both. We varied the amount of I_h via its maximum conductance (g_h) from 20 to 80 nS. Results were compared to a control condition without I_h (g_h= 0 nS).

Without I_h, amplitudes started to visibly increase at -75 mV and approached a maximum near -65 mV in a sigmoidal curve and a midpoint voltage of -70 mV. Amplitudes decreased slightly at more depolarized membrane voltages. Fig. 4A (left) shows the model output at selected postsynaptic membrane potentials and the resulting sigmoidal change of the ePSP amplitudes is shown in Fig. 4B (red trace). With I_h in the postsynaptic neuron, the obtained voltage dependence of the ePSPs remained unchanged, independently of the amount of I_h (Fig. 4A, middle: model output at selected membrane potentials; Fig. 4B, various colors: sigmoidal changes in ePSP amplitudes), but there was a small overall decrease in ePSP amplitude. The inset in Fig. 4B shows the highest amplitudes achieved without I_h vs. the strongest I_h (g_h= 80 nS). In contrast, when I_h was implemented in the presynaptic neuron, but not the postsynaptic one (Fig. 4A, right: model output), ePSP amplitudes decreased notably with more I_h, and the voltage dependence started to shift toward more depolarized values (Fig. 4C, various colors). At a g_h of 80 nS, the midpoint had reached -62 mV, without dramatic changes to the sigmoidal shape of the curve. These results matched our experimental data where in the presence of I_h, ePSP amplitudes were smaller than when I_h was blocked and the midpoint of the curve was more depolarized with I_h (compare to Fig. 2). We obtained no further changes to the behavior of the ePSPs when we additionally added I_h to the postsynaptic neuron so that it was present in both neurons (Fig. 4C, green), except a small reduction in the maximum amplitude. This further suggested that I_h in the presynaptic, but not the postsynaptic neuron, was responsible for the shift of the ePSP amplitude curve.

How did I_h exert its effects on the ePSPs? There model shows that there are two main effects I_h had on the presynaptic membrane. One is an overall increase in membrane conductance. I_h may thus act to shunt the presynaptic voltage change. Second, I_h is an inward current (reversal potential of 0 mV) and hence depolarizes the presynaptic membrane potential. To separate the impact of these two effects on the ePSP amplitudes, we altered them independently of one another in the presynaptic model neuron. First, we kept the membrane potential constant but increased overall conductance. We chose the resting potential without I_h as our reference potential (-80 mV). Membrane conductance was increased by adding

conductance to the leak current, which has a reversal potential of the reference potential. To estimate the increase in conductance for the leak, we calculated the steady state difference in membrane conductance for each level of I_h we had used in the original model (Supplemental Table T1). The original leak conductance (g_L) was 100 nS. The new values were: g_h=20 nS, g_L =106.25 nS; g_h=40 nS, g_L =109.51 nS; g_h=60 nS, g_L =111.77 nS; g_h=80 nS, g_L =113.51 nS. Adding the additional conductance, but not I_h itself, and keeping the presynaptic membrane potential at -80 mV, we found that ePSP amplitudes generally diminished when more conductance was added (Fig. 4D). However, there was no shift in their voltage dependence.

Figure 4. Modelling predicts that I_h-induced changes to the presynaptic membrane potential shift the voltage dependence of the ePSPs. A. Electrical postsynaptic potentials at various membrane potentials of the postsynaptic neuron. Without I_h (left), the largest increase of ePSP amplitude occurred between -75 and -70 mV. We obtained very similar results with I_h in the postsynaptic neuron (middle). Note, the most hyperpolarized postsynaptic membrane potential was -71 mV because I_h depolarizes the membrane potential. With I_h in the presynaptic neuron (right), the largest increase occurred between -65 and -60 mV. Top: schematic of stimulation and recording

conditions. V_m = membrane potential. Bottom: the resting membrane potential of the presynaptic neuron is given. B. Quantification of ePSP amplitudes with changing postsynaptic membrane potential. The colors indicate different levels of I_h in the postsynaptic cell (g_{max} = 0 – 80 nS). The voltage dependence of the ePSP was not affected by I_h in the postsynaptic cell, but the maximum amplitudes were diminished slightly. Inset: largest ePSPs elicited without I_h and with maximum I_h (g_{max} = 80 nS). C. Quantification of ePSP amplitudes with changing postsynaptic membrane potential. The colors indicate different levels of I_h in the presynaptic cell (g_{max} = 0 – 80 nS). The voltage dependence of the ePSP shifted to more depolarized voltages with more I_h and the maximum amplitudes diminished. Adding maximum I_h (g_{max} = 80 nS) to the postsynaptic cell (green) did not further change the voltage dependence. D. Quantification of ePSP amplitudes with changing postsynaptic membrane potential. The colors indicate different levels of leak current in the presynaptic cell (g_{leak} = 100 – 114 nS). The voltage dependence of the ePSP was unaffected, but the maximum amplitudes diminished with more leak. E. Resting membrane potential changes with different levels of I_h in the presynaptic cell (g_{max} = 0 – 80 nS). More I_h depolarized the presynaptic membrane potential. F. Quantification of ePSP amplitudes with changing postsynaptic membrane potential. The colors indicate different resting membrane potentials (-80 to -70.4 mV). The voltage dependence of the ePSP shifted to more depolarized voltages with more depolarized membrane potentials in the presynaptic cell. No changes to the maximum amplitudes were observed.

In contrast, when we kept the membrane conductance the same, but altered the membrane potential of the presynaptic model neuron, we found that the voltage dependence shifted in the predicted direction. Here, instead of activating I_h, we altered the resting membrane potential of the presynaptic neuron. This was achieved through changing the reversal potential of the general leak current in this neuron. This did not add additional conductance, but readily changed membrane potential. We compared models with resting membrane potentials corresponding to the same I_h conductance used as in previous models (g_h of 0, 20, 40, 60, and 80 nS). We obtained membrane potentials of -80.0, -75.3, -73.0, -71.5, and -70.4 mV, respectively (Fig. 4E; Supplemental Table T1). We found that at the more depolarized membrane potential of the presynaptic cell, the voltage dependence of the ePSP amplitudes shifted to more depolarized membrane voltages as well (Fig. 4F). This occurred without an overall change to the maximum ePSP amplitudes.

Thus, the model predicts that I_h exerts its actions on maximum ePSP amplitude through the addition of conductance, causing a shunt of the presynaptic membrane. Further, the shift of the voltage dependence of the ePSPs requires a shift of the presynaptic membrane potential.

I_h depolarizes the MCN1 axon terminal and shifts the voltage dependence of the MCN1-LG postsynaptic currents.

To test whether the model predictions held true in the biological system, we recorded simultaneously from the MCN1 axon terminal and LG. We first compared changes in axon membrane potential in saline (with I_h) and in CsCl (no I_h). We found that, as predicted, with I_h present, the membrane potential was more depolarized (Fig. 5A). We also noted that ePSP amplitudes changed in the predicted direction when the MCN1 axon membrane potential was manually altered. Fig. 5B shows an original recording of the MCN1 axon where a long hyperpolarizing current pulse was applied. During this pulse, the axon membrane potential slowly depolarized due to I_h. Importantly, the hyperpolarization of the axon caused an immediate increase in ePSP amplitude in LG (outlined arrow in Fig. 5B), which was followed by a slow reduction in ePSP amplitude (red arrow) as the axon membrane potential continued to depolarize (see

also inset of Fig. 5B). Finally, when we blocked I_h with CsCl (Fig. 5C) and repeated the same manipulation, the ePSP amplitude still increased when the axon was hyperpolarized. However, the slow depolarization of the axon membrane potential was gone, and so was the reduction in ePSP amplitude. These results are consistent with I_h causing a depolarization of the axon membrane potential, which in turn changes the voltage dependence of the ePSPs in LG.

Figure 5. Presynaptic I_h shifts the voltage dependence of the MCN1 ePSPs toward the functional range of the LG membrane potential. A. The resting membrane potential of the MCN1 axon terminal hyperpolarized when I_h was blocked by CsCl. On average, the resting potential was -57.3 ± 8.1 mV in low Calcium saline and -61.0 ± 8.2 with CsCl (N = 5; p<0.05, paired t-test). B. Simultaneous original recordings of LG and the MCN1 axon terminal. MCN1 was continuously stimulated at 5 Hz, and MCN1 was hyperpolarized for several seconds. Left: in low Calcium saline, a sag potential slowly depolarized the trough membrane potential (red dotted line). Concurrently, the amplitude of the ePSPs in LG diminished (red dotted line and arrow). Outlined arrow: the hyperpolarization itself caused an increase in MCN ePSP amplitude. Right: magnification of MCN1 trough potential and action potentials (bottom traces), in addition to ePSPs in LG (top). One action potential and one ePSP from the beginning of the hyperpolarization are shown in black. One action potential and one ePSP from the end of the hyperpolarization are shown in red. In the latter, I_h had depolarized the trough membrane potential. C. Same manipulation as in B., but with I_h blocked by CsCl. The slow depolarization of the MCN1 membrane potential in response to hyperpolarization was absent. There was also no more decrease in ePSP amplitude during the MCN1 hyperpolarization. D. Original current traces of two-electrode voltage clamp LG recording, at different clamp potentials (-40 to -90 mV). Individual ePSCs were elicited through MCN1 axon stimulation with 1 Hz. Left: in low Calcium saline. Right: with I_h blocked by CsCl. E. Quantification of synaptic currents obtained in the experiment shown in (A). Magenta: in low Calcium saline. Purple: with I_h blocked by CsCl. F. Quantification of all experiments. The voltage dependence of the synaptic current of each experiment was fitted with a sigmoidal curve. Mean ± SD for all experiments are shown. Colored areas indicate the range of all observed data. Magenta: in low Calcium saline. Purple: with I_h blocked by CsCl. The gray box indicates the functional range of LG's membrane potential. Light gray indicates SD.

If I_h indeed acts presynaptically, i.e., if its effects on the LG ePSPs are mediated by its actions on the MCN1 axon terminal, we would expect to see a change of the synaptic current in LG. We thus performed two-electrode voltage clamp on LG and measured synaptic currents during 1 Hz stimulation of the MCN1 axon in the *ion*. These experiments were carried out in low Ca^{2+} saline to suppress chemical synaptic transmission. We altered LG's membrane potential between -20 and -120 mV. As predicted, the amplitudes of the electrical postsynaptic currents (ePSCs) behaved similarly to the ePSP amplitudes. Specifically, more depolarized membrane potentials caused larger ePSCs. The traces in Fig. 5D show original recordings of LG's ePSCs at selected membrane potentials, with and without I_h. Fig. 5E shows the quantitative analysis for this experiment. Like for the ePSPs, the increase in synaptic current with membrane potential was approximately sigmoidal in shape. Fig. 5F shows the averaged data from all animals tested (N=13). The average midpoint voltage was -60.4±6.0 mV and ePSCs increased notably in amplitude above -80 mV. They approached maximum values around -40 mV. Hence, some of the dynamic range of the synaptic current overlapped with LG's functional range of membrane voltages (gray in Fig. 5F). When we blocked I_h with CsCl (Figs. 5D-F, purple), the midpoint of the ePSC amplitude change shifted to more hyperpolarized voltages, consistent with a presynaptic action of I_h on the voltage dependence of the electrical synapse. On average, the midpoint of the voltage dependence was now -77.6±9.6 mV (N=11, Mann-Whitney U-test, p<0.001, significantly different from no CsCl control). In summary, thus, our data demonstrates that presynaptic I_h modulates the voltage dependence of the electrical synapse between MCN1 and LG by depolarizing the membrane potential of the MCN1 axon terminal.

Discussion

Our data demonstrate that the modulation of the ePSP voltage dependence by I_h is a critical component of the MCN1-LG electrical synapse. It shifts the voltage dependence of the ePSPs to more depolarized voltages, allowing the dynamic range of the voltage dependence to overlap with the functional range of LG membrane potentials.

I_h is critical to the function of the voltage dependence of the MCN1-LG ePSPs.

In the MCN1 version of the gastric mill rhythm, the release of the neuropeptide CabTRP Ia from MCN1 is the main driver behind the slow bursting activity of LG [9, 19]. In contrast, the concurrent electrical PSPs that MCN1 elicits contribute only slightly to LG's ability to burst. When chemical release from the MCN1 axon is blocked, and only ePSPs remain, LG can produce individual action potentials, but no regular bursting can be observed [9]. Instead, MCN1's ePSPs contribute to the maintenance of the burst once it is created by prolonging burst duration and increasing spike number. Their voltage dependence is an important feature that supports this function. At resting membrane potential as well as during the LG interburst, ePSP amplitudes are small. They increase slowly after burst start and reach their largest amplitudes towards the end of the burst. To achieve this, ePSP amplitude must increase in a range of membrane voltages that LG traverses between interburst and burst, and the increase must occur slowly so that only the later portions of the burst are affected. Our data shows that the time course of the ePSP amplitude change is indeed slow, on the order of several hundred ms (Supplemental Figure S1B). We also demonstrate that I_h is a critical factor in allowing the ePSP voltage dependence to fulfill its function. With I_h blocked, ePSP amplitude was already close to maximum at LG's resting membrane potential, and the dynamic range between resting membrane potential and spike threshold was small. In contrast, with I_h present, the dynamic range of the voltage dependence was more depolarized. This facilitated amplitude changes between resting membrane potential and spike threshold.

Is the voltage dependence a gap junction property or caused by factors in the extra-junctional membrane?

When we first measured the time constant with which ePSP amplitudes changed, we were surprised by the extraordinary slow change (several hundred ms, Supplemental Figure S1B). This time constant is roughly on the time course of I_h in gastric mill neurons and initially suggested to us that I_h not just shifts the voltage dependence but is instead involved in creating it. Looking only at the typical voltage range in which LG operates seemed to further support this notion since ePSP amplitudes in this range are large and mostly unchanged when I_h, is blocked. This suggested that blocking I_h also removed the voltage dependence. And indeed, there is evidence that voltage dependence of electrical synapses can arise from intrinsic membrane properties such as spatially close voltage-gated ionic currents. Persistent Na^+ conductances in particular have been implicated in endowing electrical transmission with voltage-dependent properties [5, 22-24]. Their slow inactivation kinetics make them ideal to alter electrical transmission postsynaptically, either through changing the extrajunctional membrane resistance or the membrane potential. In the auditory axon terminals of goldfish, for example, electrical PSPs are amplified by persistent sodium currents and diminished by potassium currents [23]. In cerebellar Golgi neurons, where the de- and hyperpolarizing components of action potentials reach neighboring cells through electrical synapses, the hyperpolarizing components dominate the electrical PSP when the receiving cell

is at resting membrane potential. However, when depolarized, activation of the persistent sodium current amplifies the depolarizing PSP component, switching the PSP from predominantly hyperpolarizing to predominantly depolarizing [4, 5]. I_h has also been implicated in creating voltage dependence at an electrical synapse. Here, the effect was mediated by a decrease in membrane resistance in the extra-junctional membrane, which caused changes in the coupling coefficient of the gap junction [6].

In contrast to these examples, our data shows that the voltage dependence of the MCN1-LG ePSPs is still present and unchanged in its sigmoidal shape when I_h is blocked. The midpoint of the sigmoidal had shifted to more negative values. Hence, I_h could not have been pivotal to endow the electrical synapse with a voltage dependence. Our physiological data are also consistent with previous studies [9] and with our own model data that demonstrate that I_h in a presynaptic neuron can shift the midpoint of a voltage-dependent electrical synapse. Furthermore, when we voltage clamped LG, excluding extra-junctional influences in the postsynaptic cell, the synaptic currents showed a similar voltage dependence as the ePSPs, and their voltage dependence shifted to more negative voltages when I_h was blocked. Finally, the fact that imposed voltage changes in MCN1 as well as those elicited by I_h caused concurrent changes in ePSP amplitude further supports the conclusion that the voltage dependence is a property of the electrical synapse rather than being due to effects on the extra-junctional membrane. The model further supports this conclusion since changing the resistance of the presynaptic membrane had negligible effects on the voltage dependence.

Modulation of ePSP voltage dependence by I_h.

Neurons possess a variety of heterogeneous ionic conductances, creating complex and highly non-linear intrinsic and synaptic dynamics, and there is mounting evidence that diverse sets of ion channels are distributed throughout all neuronal compartments, including axons and their terminals. Three main functions have been attributed to I_h [25]: (1) It contributes to the resting potential. Blocking I_h leads to a shift in the resting potential towards more hyperpolarized voltages. (2) It shapes neuronal responses to hyperpolarizing input. Due to its depolarizing effects, I_h can shape the amplitude and time course of PSPs, and thereby change the response of the cell to synaptic input. (3) It contributes to the pacemaker properties of a cell by causing a postinhibitory rebound that facilitates rhythmic activity.

Our data suggests that I_h's effect on the membrane potential is to cause a shift in the voltage dependence. Our MCN1 recordings show that changing the axon terminal membrane potential has a strong impact on the ePSP amplitude, and that such amplitude changes occur concurrently with membrane potential changes when I_h is activated by hyperpolarizing the terminal. Changes in ePSP amplitude also occur when I_h is blocked and the membrane potential is artificially altered, albeit without associated conductance change. Our model corroborates these conclusions and further suggests that changes in the strength of I_h can impact ePSP voltage dependence only when I_h is in the presynaptic neuron (MCN1). Here, changing the membrane potential was necessary and sufficient to elicit the shift of the voltage dependence.

Signal transmission at electrical synapses is complex and a host of neuronal and synaptic properties affect it, including differences in pre- and postsynaptic membrane resistances, capacitances, cell sizes, and membrane potentials [1]. The key to understanding how changes in membrane potential can affect the voltage dependence of the MCN1-LG ePSPs seems to be the rectification of this synapse. When both the presynaptic MCN1 axon terminal and the postsynaptic LG neuron are at rest, there is little voltage difference across the junction. Hence, junctional conductance is low and there is little driving force across

that small junctional conductance. Incoming MCN1 action potentials thus only elicit small ePSPs in the postsynaptic LG. When LG depolarizes and begins its burst, the voltage difference between LG and the MCN axon terminal increases and so does the junctional conductance. Incoming MCN1 action potentials then produce large voltage differences across the junction. These voltage differences occur rapidly and are too fast to alter the overall synaptic conductance (due to the slow kinetics, see Supplemental Figure S1B), and hence cause the large ePSPs observed during the LG burst. I_h in the presynaptic MCN1 is a critical factor to maintain the appropriate voltage difference between the axon terminal and LG. Without I_h, the MCN1 axon terminal hyperpolarizes, causing a voltage difference between the pre- and postsynaptic sides even when both cells are at rest and LG is not bursting. Consequently, junctional conductance is already high and ePSPs are large, with only small margins left for further amplitude increase when LG bursts.

Where is I_h located?

We noted that while the model suggested that I_h must be presynaptic to exert its effects on ePSP voltage dependence, sag potentials were present in the MCN1 axon terminal and in LG. This suggests that I_h may be present on the pre- and postsynaptic side. However, we recognized that the voltages at which I_h was activated in LG were rather hyperpolarized and large current injections were required to reach these membrane potentials. During gastric mill rhythms, LG's most hyperpolarized membrane potential was -59 mV, which is approximately 30 mV more depolarized than the voltages at which I_h was visibly activated. LG is the largest cell in the STG, with a cell body size of up to 100 μm, and if at all present, I_h is likely to be located in the LG neuropil (similar to other STG neurons). The hyperpolarized activation values could thus indicate a substantial space clamp issue in this large cell, where with increasing distance of the somatic electrode from I_h's location, current measurement degrades and, due to the electrotonic distance, more hyperpolarizing current at the recording site is necessary to cause corresponding voltage changes at the current source. This effect may be exaggerated by impedance changes within the LG neurites [26] that favor current flow from the neuropil toward the soma, but not vice versa. However, I_h has never been recognized in LG before, nor does it seem necessary to create rhythmic gastric mill activity [19]. It also has a rather low expression profile in LG [27]. One of the problems in studies with electrically coupled cells is that it is difficult to tell where observed effects take place. Depending on the coupling strength of the electrical synapse, the two cells may even act as a single physiological unit [28], at least at times. It is thus possible that I_h is exclusively located in the MCN1 axon terminals and our LG recording only picked up the reflection of the MCN1 I_h. This notion is supported by the fact that LG's postinhibitory rebound was typically not accompanied by action potentials. We observed postinhibitory rebound with spike generation predominantly in MCN1, suggesting that I_h was present in the presynaptic terminal. The sag potential was also more pronounced in MCN1 than in LG and was achieved with small current injections. Given the voltage dependence of the electrical synapse and its time constant, it seems unlikely that our current injection into the MCN1 axon hyperpolarized LG far enough to cause a rebound that would be large enough to backpropagate through the gap junction into the MCN1 axon terminal and to an area in MCN's axon where spikes can be initiated. No previous data about I_h in MCN1's axon terminal is available. However, we found a consistent, albeit small, hyperpolarization of the axon resting membrane potential when I_h was blocked, and the sag potential was absent. The small membrane potential change was likely due to the electrotonic distance between recording site and the axon terminal and gap junction. We recorded the MCN1 axon at the anterior entrance to the STG, and thus just outside of the STG neuropil.

It is reasonable to assume that the gap junction to LG lies within the neuropil.

Neuromodulation of the MCN1-LG electrical synapse.

Membrane conductances like I_h are subject to intrinsic and extrinsic plasticity and have been shown to be the target of neuromodulatory transmitters. This suggests that the voltage dependence of the ePSPs at the MCN1-LG synapse could be altered by neuromodulation. The STG is subject to extensive modulation by neuromodulators that reach the STG either in the form of neurohormones [29], or by being released from sensory cells and the axon terminals of the projection neurons [30]. Their presence alters circuit output or robustness [7, 31-33]. With respect to the MCN1-LG electrical synapse, the ePSP voltage dependence could be altered by modulation of I_h in the MCN1 axon terminal. In the STG, I_h is modulated by serotonin and dopamine. The first is released from the gastropyloric receptor (GPR) cells [34, 35] and depolarizes the activation curve of I_h by ~10 mV [36]. Serotonin also inhibits the MCN1 axon terminals [37], suggesting that it is at least in the vicinity of the synaptic connections between MCN1 and LG. It is thus conceivable that the voltage dependence of the MCN1-LG ePSP could be altered through serotonergic modulation of I_h, allowing a quick adaptation of the dynamic range of the electrical synapse. Dopamine is released by projection neurons descending from the CoGs [38, 39] and its effect depends on the type of dopamine receptors [40-42]. Activating D_1 receptors leads to a long-term increase in I_h conductance in STG neurons [43, 44]. For example, dopamine depolarizes the axon of the pyloric PD neuron where it elicits extrasomatic spikes through its excitatory actions on I_h [11, 43, 45]. Other studies in invertebrates corroborate these results, suggesting that activating D_1-receptors leads to an increase in I_h, while the activation of the D_2-receptor may lead to a decrease [46-48].

Besides monoamines, neuropeptides are a major source of modulation in the STG [8, 33, 49, 50]. Proctolin and CabTRP Ia, for example, are even released by MCN1 itself, to activate LG (via CabTRP Ia) and other pyloric and gastric mill neurons (via CabTRP Ia and proctolin, [15]). This opens the fascinating possibility that peptide release from a neuron (MCN1) could adjust the dynamic range of its own electrical synapses. CabTRP Ia and proctolin elicit the same ionic current (called modulator-induced current I_{MI}), and while there are no indications that they affect I_h, I_{MI} is non-inactivating and depolarizes the resting membrane potential [51]. It is thus conceivable that CabTRP Ia or proctolin alter the membrane potential of the MCN1 axon terminal, and thereby shift the voltage dependence of the ePSPs. While we did not test this possibility in our investigation, studying modulatory influences on the MCN1-LG electrical synapse is the next step in unravelling the complex dynamics between intrinsic cell properties, extrinsic influences and electrical synaptic transmission.

Supplemental information.

Supplemental Figure S1. A. Time-triggered averages of MCN1 PSPs in LG. Individual ePSPs were elicited through MCN1 axon stimulation with 1 Hz. Left: in saline. Right: in low Calcium saline to block chemical synaptic transmission. While the amplitude diminished slightly, the major component of the PSP was electrical. B. Changes in ePSP amplitude occur with a slow time course. Two-electrode voltage clamp recording of MCN1 ePSCs in LG, in low Calcium saline. MCN1 was stimulated every 150 ms. (i) The LG membrane potential was hyperpolarized from -50 mV to -80 mV. (ii) The LG membrane potential was depolarized from -80 mV to -50 mV. Red lines show the decrement (i) and increase (ii) or the ePSC amplitudes after the voltage step. On average, the time constant for ePSC amplitude increase when LG was depolarized was 477 ± 281 ms (N=7). The average time constant for ePSC amplitude decrease when LG was hyperpolarized was 385 ±331 ms (N=6).

Calculation for models where the resting membrane potential is changed but conductance stays the same	Maximum conductance of Ih (gh) in S	Membrane potential (in V) of the presynaptic neuron at steady state = new equilibrium potential of the leak current (El)			
The leak conductance was used to alter the resting membrane potential of the presynaptic models	0	-0.08			
This manipulation does not alter the total conductance of the leak	2.00E-08	-0.0752677			
The membrane potential of the presynaptic model at steady state was measured.	4.00E-08	-0.0730134			
This was then used as the new equilibrium potential of the leak current	6.00E-08	-0.071536			
	8.00E-08	-0.0704357			

Calculation for models with additional conductance but same membrane potential						
Here, the membrane potential (Vm) was kept constant at -80, but additional conductance was added. For this, the Ih conductance was calculated for the steady state membrane potential of the presyanptic models (see above).						
Equilibrium potential of Ih	E =	0	0	0	0	0
Maximum conductance of Ih (gh) in S	g =	0.00E+00	2.00E-08	4.00E-08	6.00E-08	8.00E-08
Slope of Ih activation from Table 1	kh1	0.006	0.006	0.006	0.006	0.006
Half maximum potential for Ih from Table 1	nhalf1	-0.08	-0.08	-0.08	-0.08	-0.08
Calculated opening probability of Ih (equation 5)	mh	0.5	0.312444292	0.237862659	0.196128927	0.168814853
Measured membrane potential at steady state with Ih in V	membrane potential of presynaptic model with Ih at steady state	-0.08	-0.0752677	-0.0730134	-0.071536	-0.0704357
Calculated Ih current in A	Ih	0.00E+00	-4.70E-10	-6.95E-10	-8.42E-10	-9.51E-10
Calculated Ih conductance in S	Ih conductance at steady state	0.00E+00	6.25E-09	9.51E-09	1.18E-08	1.35E-08
Original leak conductance (gL) in S	gL original	1.00E-07	1.00E-07	1.00E-07	1.00E-07	1.00E-07
New leak conductance (gL + calculated Ih conductance)	gL new	1.0000E-07	1.0625E-07	1.0951E-07	1.1177E-07	1.1351E-07

Supplemental Table T1. Top: measurement of membrane potential at steady state in models with different amounts of I_h (g_h was varied). Bottom: calculation of changes to leak conductance to mimic the effects of conductance increase by I_h.

References

1. Alcami, P. and A.E. Pereda, *Beyond plasticity: the dynamic impact of electrical synapses on neural circuits.* Nat. Rev. Neurosci., 2019. **20**(5): p. 253-271.
2. Connors, B.W., *Synchrony and so much more: Diverse roles for electrical synapses in neural circuits.* Dev. Neurobiol., 2017. **77**(5): p. 610-624.
3. Gutierrez, G.J., T. O'Leary, and E. Marder, *Multiple mechanisms switch an electrically coupled, synaptically inhibited neuron between competing rhythmic oscillators.* Neuron, 2013. **77**(5): p. 845-58.
4. Pereda, A.E., et al., *Gap junction-mediated electrical transmission: regulatory mechanisms and plasticity.* Biochim. Biophys. Acta, 2013. **1828**(1): p. 134-46.
5. Dugue, G.P., et al., *Electrical coupling mediates tunable low-frequency oscillations and resonance in the cerebellar Golgi cell network.* Neuron, 2009. **61**(1): p. 126-39.
6. Szabo, T.M., J.S. Caplan, and M.J. Zoran, *Serotonin regulates electrical coupling via modulation of extrajunctional conductance: H-current.* Brain Res., 2010. **1349**: p. 21-31.
7. Nusbaum, M.P. and M.P. Beenhakker, *A small-systems approach to motor pattern generation.* Nature, 2002. **417**(6886): p. 343-50.
8. Stein, W., *Stomatogastric Nervous System*, in *Oxford Research Encyclopedia, http://dx.doi.org/10.1093/acrefore/9780190264086.013.153*. 2017, Oxford University Press.
9. Coleman, M.J., P. Meyrand, and M.P. Nusbaum, *A switch between two modes of synaptic transmission mediated by presynaptic inhibition.* Nature, 1995. **378**(6556): p. 502-5.
10. Gutierrez, G.J. and R.G. Grashow, *Cancer borealis stomatogastric nervous system dissection.* J. Vis. Exp., 2009. **25**: p. e1207.
11. Ballo, A.W. and D. Bucher, *Complex intrinsic membrane properties and dopamine shape spiking activity in a motor axon.* J. Neurosci., 2009. **29**(16): p. 5062-74.
12. DeMaegd, M.L., C. Städele, and W. Stein, *Axonal Conduction Velocity Measurement.* Bio-protocol, 2017. **7**(5): p. e2152.
13. Hooper, S.L., et al., *Cell dialysis by sharp electrodes can cause nonphysiological changes in neuron properties.* J. Neurophysiol., 2015. **114**(2): p. 1255-71.
14. Städele, C., M.L. DeMaegd, and W. Stein, *Extracellular Axon Stimulation.* Bio-protocol, 2017. **7**(5): p. e2151.
15. Stein, W., et al., *Divergent co-transmitter actions underlie motor pattern activation by a modulatory projection neuron.* Eur. J. Neurosci., 2007. **26**(5): p. 1148-65.
16. Follmann, R., C.J. Goldsmith, and W. Stein, *Spatial distribution of intermingling pools of projection neurons with distinct targets: A 3D analysis of the commissural ganglia in Cancer borealis.* J. Comp. Neurol., 2017. **525**(8): p. 1827-1843.
17. Colton, G.F., A.P. Cook, and M.P. Nusbaum, *Different microcircuit responses to comparable input from one versus both copies of an identified projection neuron.* J. Exp. Biol., 2020. **223**(Pt 20).
18. Bartos, M., et al., *Coordination of fast and slow rhythmic neuronal circuits.* J. Neurosci., 1999. **19**(15): p. 6650-60.
19. DeLong, N.D., et al., *Parallel regulation of a modulator-activated current via distinct dynamics underlies comodulation of motor circuit output.* J. Neurosci., 2009. **29**(39): p. 12355-67.
20. Beenhakker, M.P., D.M. Blitz, and M.P. Nusbaum, *Long-lasting activation of rhythmic neuronal activity by a novel mechanosensory system in the crustacean stomatogastric nervous system.* J. Neurophysiol., 2004. **91**(1): p. 78-91.
21. Zhu, L., A.I. Selverston, and J. Ayers, *The role of Ih in differentiating the dynamics of the*

gastric mill and pyloric neurons in the stomatogastric ganglion of the lobster, Homarus americanus. J. Neurophysiol., 2016: p. jn 00737 2015.

22. Mann-Metzer, P. and Y. Yarom, *Electrotonic coupling interacts with intrinsic properties to generate synchronized activity in cerebellar networks of inhibitory interneurons.* J. Neurosci., 1999. **19**(9): p. 3298-306.

23. Curti, S. and A.E. Pereda, *Voltage-dependent enhancement of electrical coupling by a subthreshold sodium current.* J. Neurosci., 2004. **24**(16): p. 3999-4010.

24. Haas, J.S. and C.E. Landisman, *State-dependent modulation of gap junction signaling by the persistent sodium current.* Front Cell Neurosci, 2011. **5**: p. 31.

25. Biel, M., et al., *Hyperpolarization-activated cation channels: from genes to function.* Physiol. Rev., 2009. **89**(3): p. 847-85.

26. Otopalik, A.G., J. Pipkin, and E. Marder, *Neuronal morphologies built for reliable physiology in a rhythmic motor circuit.* ELife, 2019. **8**.

27. Northcutt, A.J., et al., *Molecular Profiling to Infer Neuronal Cell Identity: Lessons from small ganglia of the Crab Cancer borealis.* bioRxiv, 2019: p. 690388.

28. Carrow, G.M. and I.B. Levitan, *Selective formation and modulation of electrical synapses between cultured Aplysia neurons.* J. Neurosci., 1989. **9**(10): p. 3657-64.

29. Pulver, S.R. and E. Marder, *Neuromodulatory complement of the pericardial organs in the embryonic lobster, Homarus americanus.* J. Comp. Neurol., 2002. **451**(1): p. 79-90.

30. Nusbaum, M.P., et al., *The roles of co-transmission in neural network modulation.* Trends Neurosci., 2001. **24**(3): p. 146-54.

31. Stein, W., *Modulation of stomatogastric rhythms.* J. Comp. Physiol. A, 2009. **195**(11): p. 989-1009.

32. DeMaegd, M.L. and W. Stein, *Neuropeptide Modulation Increases Dendritic Electrical Spread to Restore Neuronal Activity Disrupted by Temperature.* J. Neurosci., 2021. **41**(36): p. 7607-7622.

33. Marder, E. and D. Bucher, *Understanding circuit dynamics using the stomatogastric nervous system of lobsters and crabs.* Annu. Rev. Physiol., 2007. **69**: p. 291-316.

34. Katz, P.S. and R.M. Harris-Warrick, *Serotonergic/cholinergic muscle receptor cells in the crab stomatogastric nervous system. II. Rapid nicotinic and prolonged modulatory effects on neurons in the stomatogastric ganglion.* J. Neurophysiol., 1989. **62**(2): p. 571-81.

35. Katz, P.S., M.H. Eigg, and R.M. Harris-Warrick, *Serotonergic/cholinergic muscle receptor cells in the crab stomatogastric nervous system. I. Identification and characterization of the gastropyloric receptor cells.* J. Neurophysiol., 1989. **62**(2): p. 558-70.

36. Kiehn, O. and R.M. Harris-Warrick, *5-HT modulation of hyperpolarization-activated inward current and calcium-dependent outward current in a crustacean motor neuron.* J. Neurophysiol., 1992. **68**(2): p. 496-508.

37. DeLong, N.D., M.P. Beenhakker, and M.P. Nusbaum, *Presynaptic inhibition selectively weakens peptidergic cotransmission in a small motor system.* J. Neurophysiol., 2009. **102**(6): p. 3492-504.

38. Barker, D.L., P.D. Kushner, and N.K. Hooper, *Synthesis of dopamine and octopamine in the crustacean stomatogastric nervous system.* Brain Res., 1979. **161**(1): p. 99-113.

39. Tierney, A.J., T. Kim, and R. Abrams, *Dopamine in crayfish and other crustaceans: distribution in the central nervous system and physiological functions.* Microsc. Res. Tech., 2003. **60**(3): p. 325-35.

40. Clark, M.C. and D.J. Baro, *Arthropod D2 receptors positively couple with cAMP through the Gi/o protein family.* Comp. Biochem. Physiol. B Biochem. Mol. Biol., 2007. **146**(1): p. 9-19.

41. Clark, M.C., R. Khan, and D.J. Baro, *Crustacean dopamine receptors: localization and G*

protein coupling in the stomatogastric ganglion. J. Neurochem., 2008. **104**(4): p. 1006-19.

42. Johnson, B.R., J.H. Peck, and R.M. Harris-Warrick, *Amine modulation of electrical coupling in the pyloric network of the lobster stomatogastric ganglion.* J. Comp. Physiol. [A], 1993. **172**(6): p. 715-32.

43. Ballo, A.W., F. Nadim, and D. Bucher, *Dopamine modulation of Ih improves temporal fidelity of spike propagation in an unmyelinated axon.* J. Neurosci., 2012. **32**(15): p. 5106-19.

44. Rodgers, E.W., et al., *Tonic nanomolar dopamine enables an activity-dependent phase recovery mechanism that persistently alters the maximal conductance of the hyperpolarization-activated current in a rhythmically active neuron.* J. Neurosci., 2011. **31**(45): p. 16387-97.

45. Ballo, A.W., et al., *Dopamine Modulates Ih in a Motor Axon.* J. Neurosci., 2010. **30**(25): p. 8425-8434.

46. Gambardella, C., A. Pignatelli, and O. Belluzzi, *The h-current in the substantia Nigra pars compacta neurons: a re-examination.* PLoS ONE, 2012. **7**(12): p. e52329.

47. Kawai, F., M. Horiguchi, and E. Miyachi, *Dopamine modulates the voltage response of human rod photoreceptors by inhibiting the h current.* Invest. Ophthalmol. Vis. Sci., 2011. **52**(7): p. 4113-7.

48. Rosenkranz, J.A. and D. Johnston, *Dopaminergic regulation of neuronal excitability through modulation of Ih in layer V entorhinal cortex.* J. Neurosci., 2006. **26**(12): p. 3229-44.

49. Nusbaum, M.P., *Modulatory Projection Neurons*, in *Encyclopedia of Neuroscience* M.D. Binder, Hirokawa, N., Windhorst, U., Editor. 2013, Springer, Berlin, Heidelberg.

50. Nusbaum, M.P. and D.M. Blitz, *Neuropeptide modulation of microcircuits.* Curr. Opin. Neurobiol., 2012. **22**(4): p. 592-601.

51. Gray, M. and J. Golowasch, *Voltage Dependence of a Neuromodulator-Activated Ionic Current.* eNeuro, 2016. **3**(2).