Tryptophol acetate and tyrosol acetate, metabolites secreted by a probiotic yeast, block cytokine storm

Orit Malka[1†], Ravit Malishev[1†], Marina Bersudsky[2], Manikandan Rajendran[1], Mathumathi Krishnamohan[2], Jakeer Shaik[2], Daniel Chamovitz[1], Evgeni Tikhonov[3], Eliya Sultan[2,4], Omry Koren[3], Ron N. Apte[2], Benyamin Rosental[2,4], Elena Voronov[2*], Raz Jelinek[1,5*].

*Correspondence: elena@bgu.ac.il; razj@bgu.ac.il

[1] Department of Chemistry, Ben-Gurion University of the Negev, 84105 Be'er Sheva, Israel

[2] Department of Microbiology and Immunology, Faculty of Health Sciences, The Cancer Research Center, Ben-Gurion University of the Negev, Beer-Sheva 84105, Israel

[3] The Azrieli Faculty of Medicine, Bar Ilan University, 1311502 Safed, Israel

[4] Regenerative Medicine and Stem Cell Research Center, Ben-Gurion University of the Negev.

[5] Ilse Katz Institute for Nanoscale Science & Technology Ben Gurion University of the Negev, Be'er Sheva 84105, Israel;

† These authors contributed equally to this work

Abstract

Probiotic fermented foods are perceived as contributing to human health and capable of protecting against inflammation processes, however solid mechanistic evidence for the putative therapeutic benefits are generally lacking. Here we report that tryptophol acetate and tyrosol acetate, small molecule metabolites secreted by the probiotic milk-fermented yeast *Kluyveromyces marxianus* exhibit remarkable anti-inflammatory properties. Comprehensive *in vivo* and *in vitro* experimental analyses, employing LPS-induced "cytokine storm" models, reveal dramatic effects of the two molecules, added in tandem, on mice morbidity, laboratory parameters and mortality. In parallel, significant attenuation of pro-inflammatory cytokines including IL-6, IL-1α, IL-1β and TNF-α, and reduction of reactive oxygen species were recorded. Importantly, tryptophol acetate and tyrosol acetate did not completely suppress cytokine generation, but rather brought their concentrations back to baseline levels, enabling maintenance of core immune functions, including phagocytosis. The anti-inflammatory effects of tryptophol acetate and tyrosol acetate were mediated through downregulation of TLR4, IL-1R, and TNFR signaling pathways and increased A20 expression, leading to NF-kB attenuation. Moreover, the two molecules had a significant impact on mice microbiome, increasing the abundance of the genus *Bactericides,* known to exhibit anti-inflammatory properties. Overall, this work illuminates pronounced and broad-based immune modulation properties of probiotic yeast-secreted metabolites, underscoring potential new therapeutic avenues for severe inflammation pathologies.

Importance

Probiotic fermented foods are perceived as contributing to human health and capable of protecting against inflammation, however biological/molecular evidence for the presumptive health benefits are generally lacking. This paper reports the identification of two small molecule metabolites secreted by milk-fermented probiotic yeast, shown to exert remarkable anti-inflammatory effects. In particular, the small molecules - tryptophol acetate and tyrosol acetate – block severe inflammation, commonly known as "cytokine storm", in cell and mice models. Importantly, tryptophol acetate and tyrosol acetate did not completely suppress immune response, but rather brought the immune system back to baseline levels, enabling maintenance of its protective functions. The anti-inflammatory mechanisms of tryptophol acetate and tyrosol acetate were deciphered, shedding light on the immune modulation properties of probiotic yeast. This study underscores a molecular basis for the therapeutic potential of milk-fermented probiotics, and charts new avenues for treatment of severe inflammation.

Introduction

Sepsis is a life-threatening condition manifested by severe inflammation leading to multiple organ dysfunction. The inflammatory host response associated with both innate and adaptive immunity mechanisms play an important role in the development of the clinical and pathological manifestations of sepsis[1]. Recently, it has been found that the prognosis of the disease is dependent not only on the virulence of the microorganisms, but, mainly, on the host response affected by the pathogen-associated molecular patterns (PAMPs) [2,3]. Recruitment of immune cells and secretion of soluble mediators by the cells can exacerbate the severity of the disease [4]. Specifically, the

release of pro-inflammatory molecules, such as interleukin-6 (IL-6), tumor necrosis factor-alpha (TNF-α) and interleukin-1α and 1β (IL-1α and IL-1β), trigger "cytokine storms", systemic inflammation often leading to multi-organ failure and adverse clinical outcomes with high mortality rates [5]. Cytokine storms occur in various disease conditions, including sepsis and septic shock [6] and acute stages in chronic diseases [7]. Severe inflammation and the occurrence of cytokine storms were also shown to be major causes of mortality from COVID-19 [8,9].

Despite the significant progress in inflammation treatment following the discovery of antibiotics, high mortality from sepsis still exists. Thus, new approaches to improve conventional therapies are highly sought. Varied food products have been touted to endow anti-inflammatory properties and as such attract significant interest [10]. Food-extracted substances have been particularly explored as anti-inflammatory agents, including probiotics [11,12], curcumin [13], resveratrol [14], plant extracts [15], and phenolic compounds from natural sources [16]. However, the therapeutic benefits of most such substances against severe inflammation have been limited [17], and, moreover, detailed mechanistic understanding of their perceived anti-inflammatory properties are generally lacking.

Probiotics, particularly milk-fermented microorganism mixtures (yogurt, kefir), have been known to bolster the innate immune system and host-defense mechanisms against pathogens [18,19]. In a recent study, we reported on a yet-unrecognized mechanism for cross-kingdom inhibition of pathogenic bacterial communication and virulence by a small molecule - tryptophol acetate – secreted by the probiotic yeast *Kluyveromyces marxianus* in a milk-fermented probiotic microorganism mixture [20]. Specifically, tryptophol acetate was found to disrupt biofilm formation and reduce virulence of

several human pathogenic bacteria, underscoring a novel mechanism for combating

bacterial colonization and pathogenicity.

Here, we report that tryptophol acetate and tyrosol acetate, another *K. marxianus* –

secreted metabolite, exhibit remarkable anti-inflammation activities, in *in vitro, ex vivo*

and *in vivo* models. Through application of LPS-induced cytokine storm, we observed

that the two molecules had anti-oxidation, anti-inflammatory, clinical, histological and

hematological systemic protective effects against severe inflammation. Importantly,

tryptophol acetate and tyrosol acetate did not give rise to immune system shutdown,

rather reduced pro-inflammatory cytokine production to baseline levels while retaining

core immune processes including phagocytosis and generation of anti-inflammatory

cytokines. Detailed molecular analysis indicates that the anti-inflammatory activities of

tryptophol acetate and tyrosol acetate are mediated through downregulation of *TLR4*,

IL-1R, and *TNFR* signaling pathways and suppression of NF-κB activity. In particular,

we discovered that the molecules enhanced expression of A20, a key modulator of NF-

κB signaling pathways [21]. Overall, this study demonstrates remarkable anti-

inflammatory properties of probiotic yeast-secreted metabolites and furnishes a detailed

mechanistic description of the effects, underscoring their significant therapeutic

potential.

Results and Discussion

Anti-oxidation properties of tryptophol acetate and tyrosol acetate

Scheme 1 depicts the chemical structures of tryptophol acetate (**1**) and tyrosol acetate

(**2**), recently identified as metabolites secreted by the probiotic fungus *Kluyveromyces*

marxianus in milk-fermented microorganism mixture ("kefir"), and shown to exhibit

intriguing antibacterial properties through blocking quorum sensing [22]. Importantly, no

inflammation-modulatory activities have been previously reported for neither

tryptophol acetate nor tyrosol acetate.

Scheme 1. Chemical structures of tryptophol acetate (1) and tyrosol acetate (2).

We first analyzed the anti-oxidation properties of the molecules (Figure 1) as oxidation

processes and generation of reactive oxidative species (ROS) are recognized as

contributing to toxic effects in diverse diseases and pathological conditions [23]. Figure 1

depicts application of a spectrophotometric assay monitoring the visible absorbance of

the 2,2-diphenyl-1-picrylhydrazyl (DPPH) radical, which is widely employed for

evaluating the anti-oxidation activity of biomolecules [24]. In this assay, the DPPH free

radical, stable at room temperature, is reduced in the presence of an antioxidant

molecule, giving rise to a colorless ethanol solution. The bar diagram in Figure 1A

shows the degree of quenching of DPPH radical absorbance (and concomitant anti-

oxidative properties) by tryptophol acetate and tyrosol acetate, when added individually

as well as together, resembling their presence in the kefir. Figure 1A indicates a

dramatic anti-oxidation effect for the mixture of the two molecules. Specifically, while

1 individually gave rise to 10% quenching of DPPH absorbance and **2** alone reduced

the absorbance by ~3%, when the **1** and **2** were added together (at the same 76

concentrations) to the DPPH solution, 30% absorbance attenuation was recorded 77

(Figure 1A). 78

Figure 1. Anti-oxidation effect of tryptophol acetate and tyrosol acetate. Anti- 80
oxidative capacity of the molecules as measured by DPPH (**A**) and ORAC (**B**) assays. 81
A. The results are presented as a percentage of absorbance inhibition (517 nm, the 82
absorbance maximum of DPPH) in the presence of tryptophol acetate (**1**) alone, tyrosol 83
acetate alone (**2**) alone, and **1**+**2** together. **B**. Fluorescence decay of fluorescein induced 84
by AAPH in the absence or presence of **1** and **2**. Reaction mixtures containing 85
fluorescein (60 nM) and AAPH (18.75 mM) in 200 µl of phosphate buffer (75 mM, pH 86
7.4) were incubated at 37°C for 800 min. Changes in fluorescence intensity emitted by 87
fluorescein were monitored. Results are presented as the net area under the curve 88
(AUC). Each value is the mean ± SD of triplicate experiments. ***$p < 0.0001$ 89

The results of the oxygen radical absorbance capacity (ORAC) assay in Figure 1B 91

further demonstrate substantial anti-oxidation activity by tryptophol acetate and tyrosol 92

acetate when added together. The ORAC experiment measures the quenching of 93

fluorescein emission by peroxyl radicals, and inhibition of the oxidation reaction by 94

molecular species added to the solution [25]. As shown in Figure 1B, in the control 95

sample, the fluorescence emission recorded for fluorescein decreased instantaneously 96

because of the significant quenching induced by the peroxyl radicals. However, upon 97

co-addition of **1** or **2**, the fluorescence decays were significantly longer, up to around 98

400 min (in the case of **1,** purple curve) and 800 min (for **2**, beige curve), likely 99

accounting for scavenging of the peroxyl radical by the molecules. Strikingly, 100

significant inhibition of peroxyl-induced fluorescence quenching occurred when 101

tryptophol acetate and tyrosol acetate were added together to the solution (green curve 102

in Figure 1B), underscoring their anti-oxidation effect according to their existence in 103

the kefir. 104

Oral uptake of tryptophol acetate and tyrosol acetate mixture provides protection 106
against LPS-induced cytokine storm in mice 107

As tryptophol acetate and tyrosol acetate exhibit more powerful anti-oxidation activity 108

as a mixture , we next investigated the anti-inflammatory properties of the **1+2** mixture 109

in vivo using a model of lipopolysaccharide (LPS)-induced inflammation, which 110

triggers a massive pro-inflammatory cytokine release identified as a "cytokine storm" 111

[26,27]. In the experiments, C57BL/6 mice were injected with LPS (30 mg/kg), and the 112

effects of the **1** + **2** mixture administrated orally by gavage (each molecule at 150 μg/Kg 113

per mouse) were monitored up to 156 hours after LPS injection (scheme in Figure 2A). 114

LPS and 1 + 2 mixture simultaneous

1 + 2 mixture 28 hrs. after LPS

Figure 2. Tryptophol acetate and tyrosol acetate mixture shield mice from LPS-induced severe inflammation. (**A**) Experimental scheme. After intraperitoneal (IP) injection of LPS (30 mg/kg), mice were randomly divided into two groups, PBS-treated and treated with **1**+**2** immediately after LPS injection (**B, C**). A third group treated with the molecules 28 hours after LPS injection (**D, E**). Survival (**B, D**) and weight loss (**C, E**) of the mice were monitored for 156 hours .(n=6 in each group)

Initially, mice were injected with LPS and at the same time interval were treated with 124
1+2 mixture. The percentage survival and weight loss in the LPS-administered mice, 125
with and without treatment, show that the molecules mixture significantly improved the 126
disease outcome. Remarkably, while all LPS-treated mice died within 72 hours due to 127
the severe inflammation (i.e. the "cytokine storm"), 100% of the mice orally given the 128
mixture of **1** + **2** survived (Figure 2B). Furthermore, the LPS-administered mice 129
experienced substantial weight loss prior to mortality (Figure 2C, black line). In 130
contrast, mice which were orally treated with the molecular mixture, initially lost 131
weight (up to 60 hours), but reverted to their initial values within 156 hours (Figure 2C, 132
green line). 133

To prove the therapeutic effects of the molecules, we carried out another experiments, 134
in which **1+2** mixture was given to mice that were already impacted by the onset of 135
LPS-induced inflammation (Figure 2D-E). In these experiments, the molecular mixture 136
was administered to the mice by gavage 28 hours after LPS injection. At this time- 137
interval, severe signs of the disease were already evident, including decreased motor 138
activities, ruffled fur, diarrhea, substantial eye discharge and respiratory distress. 139
Notably, the results in Figure 2D demonstrate that while all untreated mice died within 140
48 hours after LPS injection, 100% of the mice treated with the **1** + **2** mixture survived. 141
The weight lost patterns in mice treated with the molecules 28 hours after LPS 142
administration when severe inflammation was already developed (Figure 2E) were 143
similar to the case of mice that were orally given the two molecules simultaneously 144
with LPS (e.g., Figure 2C). Overall, the results presented in Figure 2 demonstrate an 145
exceptional therapeutic effect of the tryptophol acetate and tyrosol acetate mixture. 146

Oral uptake of tryptophol acetate and tyrosol acetate decreased tissue damage induced by LPS

Cytokine storms generally account for multisystem failure manifested in considerable damage to the lungs, liver, and the gastrointestinal (GI) tract [28]. To investigate whether the healing effects of tryptophol acetate and tyrosol acetate (e.g., Figure 2) also lead to reduced organ damage, we carried out histology analyses of different tissues obtained from LPS-injected mice treated or non-treated with the **1+2** mixture (Figure 3). The representative hematoxylin and eosin (H&E) staining image of lung tissue from a healthy mouse displays normal alveolar walls and no inflammatory cell infiltration (Figure 3A, top row). In LPS-injected mice, signs of severe lung injury characterized by lung edema, hemorrhage, intensive cellular infiltrate, and accumulation of fibrin were observed already during the first 6 hours (Figure 3A, middle panel).

Figure 3. Tryptophol acetate and tyrosol acetate prevent inflammation-associated 161
tissue damage. A(i), B and C Representative pictures of H&E-stained sections of 162

obtained organs from healthy and LPS-injected mice with or without administration of molecule mixture **(A(i))** lung, **(B)** liver and **(C)** small intestine samples. Original magnification x20 for lung and small intestine, x10 for liver. Quantification of tissue damage was performed, as discussed in Materials and Methods (M&M) and is presented as a bar diagram on top of each panel. **(A (ii))** Lung tissues obtained 6 and 24 h after LPS administration were stained with anti-myeloperoxidase (MPO) antibodies for neutrophils detection. Original magnification x10. of the ammount of neutrophils was counts as discribed in M&M and presented in the graph. **(A(iii))** Monocytes were detected by anti-Ly6C antibodies and macrophages by anti-F4/80 antibodies. Quantification of the positive cells was assessed, as described. Insert in panel B represents an amount of hepatocytes with preserved cell structure. Insert in panel C represents quantification of intestinal crypt damage. Representative pictures are presented in magnification x20. *p < 0.05; **p < 0.01; ***p < 0.0001 ****p<0.00001

Strikingly, lung tissue damage was largely eliminated in mice to which the **1+2** mixture was co-administered with LPS [Figure 3A(i), bottom row]. Indeed, the pulmonary architecture was mostly preserved in those mice, with only small local zones of inflammation. In addition, the alveolar walls were almost not altered, and minimal accumulation of inflammatory cells was observed at the indicated time intervals. Notably, after 72 hours (in which only mice treated with the molecules survived), the lung structure was similar to the structure observed in healthy mice (Figure 3A(i)). Lung tissue damage was assessed by quantification of pathological score [bar diagram in Figure 3A(i), inset]. Indeed, the increase in mean pathological score after LPS administration, was significantly reduced after treatment with **1 + 2** mixture (at 24 and 72 hours). Those findings demonstrate that tryptophol acetate and tyrosol acetate mixture effectively blocked inflammation-induced damage in lungs.

To further examine the protective effects of the molecules on lung damage after LPS-induced inflammation, we assessed the recruitment of myeloid cells into the lungs (Figure 3A,ii-iii). In the experiments, we performed immunohistochemical (IHC) staining with anti-MPO antibodies for neutrophils, anti-Ly6C antibodies for monocytes

190
191
192
193

(Ly6C[positive]) and anti-F4/80 antibodies for macrophages (F4/80[pos]), considered mature macrophages [29]. Figure 3A(ii), top row, shows minimal neutrophil infiltration in lungs from control mice. However, lungs of LPS-treated mice displayed significant recruitment of neutrophils already 6 hours after LPS injection (Figure 3A(ii), middle row). Remarkably, in mice which received the **1**+**2** mixture, the recruitment of neutrophils was significantly decreased and was comparable to healthy (PBS treated) mice (bar diagram in Figure 3A(ii), inset).

Figure 3A(iii) illuminates the recruitment of monocytes and macrophages into the lungs. Both types of cells are strongly associated with inflammation-induced lung pathogenesis [30,31]. Lung-recruited Ly6C[pos] monocytes account for microvascular endothelial cell activation and vascular injury in LPS-induced early endotoxemia, leading to enhanced pulmonary vascular leakage [32,33]. Figure 3A,iii shows that while the abundance of lung-recruited Ly6C[pos] monocytes significantly increased following LPS administration both after 6 and 24 hours, treatment with tryptophol acetate and tyrosol acetate mixture effectively reduced Ly6C[pos] monocytes levels back to baseline (i.e. healthy mice). In contrast, treating mice with **1**+**2** mixture increased F4/80[pos] myeloid cells (Figure 3A,iii, bottom row), as these macrophages are associated with suppression of inflammation in lung tissues [29,34].

Figure 3B indicates that **1**+**2** mixture also prevented tissue damage in the liver. The H&E staining image of healthy liver is presented in Figure 3B, top row. In contrast, the structure of liver tissue was significantly damaged 6 and 24 hours after LPS administration, in which acute inflammatory response was apparent, characterized by decreased cell density, enlarged cell gaps and neutrophil infiltration. Morphological changes in the liver tissue, however, were markedly decreased in mice treated with **1**+**2** mixture (Figure 3B, bottom panels). Indeed, administration of **1** and **2** featured higher

number of normal nuclear cells in comparison to LPS-administered mice that were 219

untreated with the molecules (bar diagram in Figure 3B, inset). 220

Similar prevention of tissue damage by tryptophol acetate and tyrosol acetate was 221

apparent in the small intestine (Figure 3C). Specifically, H&E staining of intestinal 222

tissues extracted from LPS-injected mice displayed an extensive damage of both 223

mucosa and sub-mucosa, loss of intestinal mucosal crypt morphology and recruitment 224

of inflammatory cells (Figure 3C, middle row). In comparison, mice treated with **1** + **2** 225

mixture exhibited markedly lower tissue damage (Figure 3C, bottom row). 226

Quantification of crypt injury (Figure 3C, inset) attests to the significantly lower tissue 227

damage, in all time points, in LPS-injected mice treated with the two molecules. 228

Effects of oral uptake of tryptophol acetate and tyrosol acetate on systemic 230

inflammation. 231

We further assessed the effects of the tryptophol acetate and tyrosol acetate mixture 232

oral uptake on systemic inflammation (Figure 4). In the experiments depicted in Figure 233

4, the peripheral cell blood count (CBC) was evaluated at different time intervals. The 234

results show that in LPS-injected mice that were not treated with **1+2,** significant 235

reductions in red blood cell (RBC) counts, hemoglobin (HGB) and hematocrits (HCT) 236

occurred. Notably, however, addition of **1** + **2** mixture significantly increased these 237

parameters (Figure 4A top row, green lines vs black lines). Treating mice with **1** + **2** 238

mixture also increased thrombocyte counts (PLT), which were reduced after LPS 239

injection (Figure 4A). The CBC data in Figure 4A indicate that tryptophol acetate and 240

tyrosol acetate prevented both inflammation-induced anemia and thrombocytopenia 241

that are usually associated with severe inflammation, especially accompanied by 242

cytokine storms [35]. Notably, Figure 4A further reveals insignificant cell count changes in the white blood cells (WBC) and granulocytes (GRA).

Figure 4. Effects of tryptophol acetate and tyrosol acetate mixture on systemic inflammation. Samples of the peripheral blood were obtained from the tail vein at different time intervals and assessed for CBC. Serums were collected and studied by ELISAs. **A.** The results of RBC, HGB, HCT, PLT, WBC, GRA at different time intervals. LPS-administered mice with PBS treatment (black lines) and treated with **1**+**2 mixture** (green), *p < 0.05. **B.** Cytokine levels in serum, assessed by commercial ELISA kits. Data are presented as mean ± SD in each group (n=3). *p<0.05l, **p<0.005, ***p<0.0001. **(C)** Lungs were obtained from LPS treated mice at indicated time intervals and mRNA was extracted and proceed to RTPCR. Expression of IL-6, IL-1α and IL-1β is shown. Data are shown as mean ± SD in each group (n=4). *p<0.05, **p<0.005.

The clinical hallmark of cytokine storms is the increased secretion of pro-inflammatory   269

cytokines. Accordingly, we determined the effects of tryptophol acetate and tyrosol   270

acetate mixture given orally to the LPS-injected mice on the levels of pro-inflammatory   271

cytokines, including IL-6, IL-1β and TNF-α in serums (Figure 4B) and lungs (Figure   272

4C). Figure 4B demonstrates that while LPS injection increased the levels of cytokines   273

in serum, treatment with **1**+**2** mixture significantly attenuated cytokine concentrations   274

within 6-24 hours (Figure 4B). Specifically, blood concentrations of IL-6 in LPS-   275

administered mice treated with the molecules were lower compared to the untreated   276

LPS-injected mice in all time intervals (note that after 48 hours, LPS-administered mice   277

treated with PBS did not survive, Figure 4B). However, after 48 hours, the IL-6 level   278

of treated mice was almost identical to the healthy mice. Statistically significant   279

reductions of IL-1β and TNF-α levels in LPS-administered mice that received the **1**+**2**   280

mixture was observed 12 hours after LPS injection (Figure 4B).   281

Tryptophol acetate and tyrosol acetate mixture also inhibited the expression of pro-   282

inflammatory cytokines in the lungs (Figure 4C). Specifically, while LPS   283

administration significantly elevated the mRNA levels of IL-6, IL-1α, and IL-1β in lung   284

tissues, oral treatment with **1** + **2** mixture reduced the expression of these cytokines at   285

the examined time intervals (Figure 4C). Thus, oral administration of tryptophol acetate   286

and tyrosol acetate mixture decreased cytokine secretion in serum (Figure 4B) and   287

expression (Figure 4C) thereby inhibiting severe systemic inflammation after LPS   288

injection.   289

Tryptophol acetate and tyrosol acetate reduced the production of ROS and pro-inflammatory cytokines in LPS-activated murine peritoneal macrophages, without reducing phagocytosis function 291 292 293

To investigate the mechanistic basis of the clinical and biological immunomodulation effects of tryptophol acetate and tyrosol acetate (e.g., Figures 1-4), we performed *ex-vivo* experiments using murine peritoneal macrophages. Figure 5A depicts the effects of tryptophol acetate and tyrosol acetate on reactive oxygen species (ROS) generation using the CellRox Deep-Red dye assay. In general, elevated inflammation levels go together with enhanced production of ROS [36]. Oxidative stress conditions were attained in the experiments by stimulating the murine peritoneal macrophages with LPS (100 ng/ml) with or without co-addition of the **1**+**2** mixture. The flow cytometry analysis revealed a significant decrease in intracellular ROS after 16 hours incubation of LPS-stimulated macrophages with **1** + **2** mixture**,** in comparison with the untreated LPS-stimulated macrophages (Figure 5A). 294 295 296 297 298 299 300 301 302 303 304

Figure 5. Anti-inflammatory properties of tryptophol acetate and tyrosol acetate 308
ex vivo. **A**. Intracellular reactive oxygen species (ROS) after LPS stimulation of murine 309
peritoneal macrophages (three replicates flow cytometry analysis), the results are the 310
Geometric Mean Fluorescent Intensity (GMFI) of the CellRox Deep-Red dye. 311
(Unpaired t-test, $p < 0.05$; n=14). **B**. Representative scatter plots and gating conditions 312
for green-fluorescent beads engulfment analysis, depicting bead uptake after 1 hour 313
incubation of the macrophages, with or without co-addition of **1**+**2** mixture. 314
Percentages of bead positive cells are indicated in the bottom right quadrants. **C**. 315
Analysis of phagocytosis function of murine peritoneal macrophages after LPS 316
stimulation incubated for 1 or 3 hours with or without molecules in concentration of 317
100 μM each by using green-fluorescent beads (Unpaired t-test, two-tailed; $p < 0.05$). **D**. 318

Cytokine levels determined by ELISA from supernatants of *ex-vivo* isolated peritoneal murine macrophages. Macrophages were incubated for 16-hour with and without LPS and **1**+**2** mixture. Results presented as mean ± SD; n=3 from two independent experiments. *p<0.05 ***p<0.0001.

Since phagocytosis is a key function of macrophages [37], we examined macrophage phagocytic action in LPS-stimulated murine peritoneal macrophages with and without co-incubation with the **1** + **2** mixture (Figure 5B,C). Phagocytosis was assessed in the experiments by flow cytometry measuring the uptake of green-fluorescent beads by the cells. The representative gating analysis by flow cytometry (1 hour phagocytosis assay, Figure 5B) together with the bar diagram (Figure 5C) demonstrate no difference in the phagocytic activity of LPS-stimulated macrophages in the absence or presence of the **1**+**2** mixture (1 and 3 hours of phagocytosis assays). These results indicate that the tryptophol acetate and tyrosol acetate mixture did not adversely affect the phagocytic ability of macrophages.

To further confirm the immunomodulatory effects of tryptophol acetate and tyrosol acetate *ex vivo*, we quantified production of pro-inflammatory cytokines by the LPS-stimulated murine peritoneal macrophages with and without co-incubation with the molecule mixture (Figure 5D). Indeed, while the production of IL-1α, IL-1β, IL-6, and TNF-α by macrophages increased within 16 hours after LPS stimulation, addition of the **1**+**2** mixture significantly reduced the concentrations of all these pro-inflammatory cytokines, in most cases back to the non-stimulated macrophage levels. In contrast, tryptophol acetate and tyrosol acetate did not affect secretion of the anti-inflammatory cytokines TGF-β and INF-β (Figure 5D). These results again demonstrate that the addition of the molecule mixture did not disrupt core anti-inflammatory mechanisms of the host. Together, the results presented in Figure 5 demonstrate that tryptophol acetate

and tyrosol acetate decreased intracellular ROS and inhibited generation of pro-inflammatory cytokines, while not adversely affecting important immune processes, including macrophage phagocytic function and anti-inflammatory cytokine production.

Effect of tryptophol acetate and tyrosol acetate on NF-κB pathways in LPS-activated macrophages ex vivo and in vitro

To obtain further insight into the molecular contours underlining the immune modulating activities of the tryptophol acetate and tyrosol acetate mixture, we investigated intracellular pathways associated with NF-κB, a key protein in major inflammation gene cascades [38] (Figure 6). Indeed, the representative western blot (WB) image in Figure 6A (**i**) demonstrates that addition of **1** + **2** mixture to the LPS-stimulated macrophages markedly inhibited generation of the phosphorylated form of NF-κB (p65) in comparison to stimulation with LPS (relative band intensities are depicted in the bar diagram in Figure 6A (ii)).

Figure 6. Molecular mechanisms underlining the anti-inflammatory activity of the   361
tryptophol acetate and tyrosol acetate mixture. Intraperitoneal macrophages were   362
obtained 72 hours after thioglycolate injection and cultured with LPS with or without   363
1+2 mixture. (**A**) At different time intervals, cell lysates were assessed by Western blot   364
(WB). (**i**) Representative WB of phosphorylated NF-κB (p65) protein expression, with   365
the effects of the mixture (100 μM each). (**ii**) Relative band intensity histograms. The   366

graph represents an average of three independent experiments. Bars indicate standard 367
errors of the means; **p<0.001, ***p<0.0001. Beta-actin was used to assess equal 368
protein loading. (**B**) RAW 264.7 cell line was used for definition of the NF-kB pathway 369
related receptors. Cells were incubated with LPS with and without molecules at the 370
indicated time intervals. RNA was extracted and cDNA was analyzed by RT-PCR. (**C**) 371
Relative protein and mRNA expression of A20 after stimulation with 1 μg/ml of LPS 372
in the absence and presence of **1**+**2** mixture (100μM), detected by RT-qPCR. Error bars 373
indicate standard deviations of three independent cultures. * P<0.05, **** P< 0.0001 374
ANOVA followed by Tukey's post hoc analysis. (**D**) (**i**) HEK-293T cells were 375
transfected with pCAGGA vector (*HA-A20*) (1.5 μg) and incubated with and without 376
1+**2** mixture (1 or 2 h). Protein expression of A20 in cell lysates is presented in WB. 377
HEK-293T cells were used as a negative control. (**ii**) Relative band intensity 378
histograms. The graph represents an average of three independent experiments. Bars 379
indicate standard error of the means; statistical analysis calculated by Unpaired t-test, 380
p<0.0001. Beta-actin was used to ascertain equal protein loadings. 381

To shed further light on the parameters responsible for NF-κB attenuation incurred by 383

tryptophol acetate and tyrosol acetate in LPS-stimulated macrophages, we evaluated 384

expression of prominent upstream receptors associated with NF-κB cascades, 385

specifically TLR4, IL-1R and TNFR [39] (Figure 6B). The bar diagrams in Figure 6B 386

depicts the levels of the receptors' mRNA recorded in RAW 264.7 macrophage cells, 387

stimulated by LPS in the presence or absence of **1** + **2** mixture. Figure 6B demonstrates 388

that addition of the molecules to the LPS-stimulated cells significantly attenuated 389

expression of the genes of all three receptors. Specifically, LPS stimulation induced 390

expression of *TLR4* during the first hour (black bar, Figure 6B, left), while co-addition 391

of **1**+**2** mixture together with LPS resulted in suppression of *TLR4* expression, reverting 392

to the baseline (pre-inflammation) level. Similarly, tryptophol acetate and tyrosol 393

acetate significantly attenuated expression of both *IL-1R* (grey bars) and *TNFR* (white 394

bars) 4 hours after LPS stimulation, giving rise to baseline expression levels in both 395

receptors. 396

Accounting for the reduction of the upstream receptors in the NF-κB pathways by the tryptophol acetate and tyrosol acetate (e.g., Figure 6B), we further tested the effects of the molecules upon expression of the A20 protein, known to suppress signaling cascades associated with the TLR receptors [40] (Figure 6C-D). The bar diagram in Figure 6C presents experimental data corresponding to RAW 264.7 cells treated with **1+2** mixture and the A20 m-RNA levels measured after 30 min and 60 min. As shown in Figure 6C, both LPS, and the **1+2** mixture co-added with LPS, induced significant expression of A20 gene within 60 minutes after addition to the cells. However, when **1+2** were co-added to the cells together with LPS, significantly higher expression of A20 was found.

To assess the protein level of A20, we further carried out WB analysis (Figure 6D), employing HEK-293T cells transfected with the pCAGGA mammalian expression vector (HA-A20), allowing transient transfection of A20 [41,42]. Specifically, after transfection, the tryptophol acetate and tyrosol acetate mixture was incubated with the cells for different time intervals (HEK-293T cells were used as a negative control, Figure 6D,i). The WB data in Figure 6D demonstrate that **1+2** mixture induced significant expression of A20, both after 1 and 2 h of incubation compared to the control HEK-293T cells comprising the vector. Together, the RT-PCR and WB results in Figure 6C-D suggest that modulation of A20 levels by tryptophol acetate and tyrosol acetate is likely a key factor in attenuation NF-κB expression and concomitant anti-inflammatory effects of the molecules.

Tryptophol acetate and tyrosol acetate affect mouse gut microbiome

In tandem with elucidating the dramatic clinical effects and molecular facets of the

inflammation modulatory activities of tryptophol acetate and tyrosol acetate, we also

examined the effect of the molecules on gut microbiome of the mice (Figure 7). The

relationship between gut microbiota and host immune properties has recently emerged

as a prominent factor in systemic response to inflammation. Previous seminal studies,

for example, linked inflammatory cytokine production and individual variations in

cytokine response to the composition and function of the microbiota [43]. In the

experiments depicted in Figure 7, mice microbiome analysis was carried out through

collecting the stool samples from all participating mice, the day before and at intervals

of 6 h and 24 h after LPS injection. The microbial taxonomic 16S rRNA gene sequences

were subsequently determined.

Figure 7. Effect of tryptophol acetate and tyrosol acetate mixture treatment on mice microbiome. (A, B) Alpha diversity analysis using the Shannon index after (**A**) 6 hours (p=0.46) and (**B**) 24 hours of treatment (p=0.29); quantitative variables were compared between groups with the Wilcoxon U-test, for non-normally distributed data. **(C, D)** Principal coordinate analysis plot representing beta-diversity based on weighted UniFrac distances after **(C)** 6 hours (permutational multivariate analysis of variance, P=0.001) and **(D)** 24 hours of treatment (permutational multivariate analysis of variance, P=0.038). (**E**) Bacterial taxa *Bacteroides ovatus* relative abundance (%) boxplots for untreated and molecules treated groups. The taxa more abundant in treated group 24 h from LPS injection. (Treated and untreated, n=8)

When comparing the microbial richness (i.e., alpha diversity; Figure 7A, B) before and

after treatment with tryptophol acetate and tyrosol acetate (at 6 hours and 24 hours,

respectively), no significant differences were observed (Wilcoxon test, P=0.46 and

P=0.29). However, we found differences in beta diversity (i.e., between sample

diversities; Figure 7C, D) between treated and untreated mice 6 and 24 hours after LPS

injection (PERMANOVA, P=0.001 and P=0.038, respectively). This important

observation indicates that although the species richness was not significantly altered

between the treated and untreated mice, dysbiosis (shift in the microbiome) at the

community level was already observed 6 hours after LPS administration. Dysbiosis 6

hours after LPS treatment was characterized by a profound reduction in the relative

abundance of commensals such as *Blautia*, *Bacteroides*, *Clostridium, Sutterella* and

Parabacteroides and by an increase in the relative abundance of *Ruminococcus* (Table

S1, Supporting information). These changes occurred in both the treated and untreated

groups. Twenty-four hours after LPS administration we observed a decrease in the

relative abundance of *Blautia, Bacteroides, Clostridium and Oscillospira* and increase

in the relative abundance of *Ruminococcus*.

Figure 7E underscores the effect of treating the LPS-administered mice with tryptophol

acetate and tyrosol acetate. Importantly, the abundance of *Bacteroides* increased in the

mice that were treated with **1+2** compared to the mice that were not treated (24 hours

after LPS injection; Figure 7E). This taxon was also abundant in the microbiome of

healthy mice (ANCOM test significance W=68 and W=72 respectively (Table S1)).

This result is important as *Bacteroides* has been linked to varied immune-protective

and anti-pathogenic activities associated with microbiome modifications [44].

Discussion

Current therapy of sepsis is mainly based on anti-bacterial and symptomatic treatments. Important strategies for overcoming sepsis syndromes have been based on blocking cytokines and other pro-inflammatory molecules. While mortality from sepsis significantly decreased in the past century, in many cases of septic shock or cytokine storms, for example associated with the recent SARS-Covid-19 pandemic, mortality is high [2,9]. In most patients experiencing such sepsis syndromes, various inflammatory factors are responsible for the development of multiorgan failure and new therapeutic approaches which target upstream pro-inflammatory molecules are highly sought.

In this work, we identified two metabolites secreted by the probiotic fungus *Kluyveromyces marxianus* in a milk-fermented microorganism mixture (kefir) – tryptophol acetate and tyrosol acetate - which exhibit remarkable anti-inflammatory activities. Comprehensive *in vivo, ex vivo* and *in vitro* experiments demonstrate an effective blocking of the "cytokine storm" by an equimolar mixture of the two molecules. Specifically, we observed that mice inflicted with LPS-induced sepsis and orally treated with tryptophol acetate and tyrosol acetate, had significantly milder patterns of the disease in comparison with LPS-administrated control mice that received only buffer. Importantly, no mice mortality was observed both upon co-administration of the molecules with LPS (i.e., sepsis preventive effect), as well as upon providing the molecules to mice already displaying signs of severe inflammation (Figure 2).

LPS administration in mice yields clinical signs of human sepsis. Thus, we assessed the effects of tryptophol acetate and tyrosol acetate mixture on systemic inflammation and on various vital organs in LPS-injected mice (Figures 3-5). In sepsis, multiorgan destruction due to severe inflammation leads to hepatic and renal failure,

respiratory distress, dysfunction of gastrointestinal system and hematological changes, 494

such as coagulation disorders [45]. The reported experimental data demonstrate that orally 495

administered tryptophol acetate and tyrosol acetate mixture prevented tissue damage 496

and attenuated hematological changes. Specifically, the molecules' mixture attenuated 497

alveolar wall degradation, and in parallel decreased lung congestion (Figure 3A). 498

Tryptophol acetate and tyrosol acetate also significantly affected recruitment of 499

myeloid cells, which are intimately involved in exacerbation of lung inflammation [46]. 500

Indeed, we observed that mixture of tryptophol acetate and tyrosol acetate reduced 501

recruitment of neutrophils and Ly6C-positive monocytes into lungs (Figure 3A, iii). 502

Moreover, histology analyses reveal that, after treatment with the molecules, more 503

abundant mature F4/80-positive macrophages was detected, likely accounting for tissue 504

repair [29,34] (Figure 3A, iii). We also recorded substantial healing effects in the case of 505

LPS-induced liver tissue damage, in which the molecules stemmed the loss of hepatic 506

cells (Figure 3B), and reduced crypt damage in intestinal tissue (Figure 3C). 507

The systemic therapeutic effects of tryptophol acetate and tyrosol acetate were 508

further analyzed *in vivo* and *ex vivo* (Figure 4). We found that LPS-administered mice 509

treated with the molecules' mixture displayed lower anemia and thrombocytopenia that 510

usually accompany severe septic syndromes and cytokine storms (Figure 4A). 511

Furthermore, significant reduction in the expression of pro-inflammatory cytokines in 512

lung tissue and in serum were recorded after treating LPS-injected mice with the 513

tryptophol acetate and tyrosol acetate mixture (Figure 4B and 4C). These results are 514

significant, since release of pro-inflammatory mediators, including cytokines, 515

chemokines and ROS is the hallmark of cytokine storms [47]. 516

Particularly significant in this context were the *in vivo* and *in vitro* data recorded 517

for IL-6, specifically the attenuation of this pro-inflammatory cytokine by orders of 518

magnitude to the baseline level after treatment with tryptophol acetate and tyrosol   519

acetate (Figure 4B, 4C; Figure 5B). Cytokine storm syndromes in COVID-19 patients,   520

for example, have been associated with dramatic increase in blood IL-6 levels, and IL-   521

6 concentrations have been shown as markers for fatality outcomes [48]. As such, the   522

considerable and rapid reduction of IL-6 levels induced by the tryptophol acetate and   523

tyrosol acetate point to potential therapeutic avenue for severe inflammation   524

syndromes, such as affected by COVID-19. Additional *ex vivo* experiments in Figure   525

5D assessing pro-inflammatory cytokine secretion echoed the *in vivo* results,   526

demonstrating attenuated secretion of pro-inflammatory cytokines by LPS-stimulated   527

macrophages in the presence of the tryptophol acetate and tyrosol acetate mixture.   528

Importantly, different from numerous anti-inflammatory strategies reported in the   529

literature and clinically employed, tryptophol acetate and tyrosol acetate did not induce   530

complete immunosuppression, rather the molecules reverted the pro-inflammatory   531

cytokine levels to baseline (pre-inflammation) values.   532

A major question underlying the anti-inflammatory effects of tryptophol acetate   533

and tyrosol acetate concerns their mode of action. *Ex vivo* experiments utilizing LPS-   534

stimulated murine peritoneal macrophages reveal important mechanistic aspects   535

(Figures 5-6). Specifically, in parallel with inhibition of pro-inflammatory cytokines,   536

we recorded a significant reduction in ROS levels. ROS include short-lived bioactive   537

molecules produced by immune competent cells serving as messenger molecules for   538

various physiologic and pathologic processes [49]. ROS particularly activate multiple   539

inflammatory signaling pathways [50]. Notably, we recorded a significant reduction in   540

ROS levels after tryptophol acetate and tyrosol acetate were added to LPS-activated   541

murine peritoneal macrophages (Figure 5A). It should be noted that while tryptophol   542

acetate and tyrosol acetate significantly attenuated the pathophysiological   543

consequences of severe inflammation, the molecules did not adversely affect core 544

immune processes, particularly phagocytosis activity (Figure 5B-C). These results are 545

consistent with previous studies, reporting anti-inflammatory and anti-oxidant effects 546

without impairing macrophage bactericidal activities [51,52]. 547

To account for the broad-based immune modulation effect of tryptophol acetate 548

and tyrosol acetate, particularly the significant reduction in pro-inflammatory cytokine 549

levels, we investigated the impact of the molecules' mixture upon the immune signaling 550

pathway involving NF-κB (Figure 6). NF-κB is central to transcription pathways of 551

many genes encoding pro-inflammatory mediators and plays a crucial role in varied 552

inflammatory disease conditions [37]. Indeed, addition of tryptophol acetate and tyrosol 553

acetate reduced NF-κB level in macrophages stimulated with LPS (Figure 6A). 554

Furthermore, the RT-PCR data in Figure 6 attest to pronounced downregulation of the 555

main upstream receptors involved in NF-κB activation - TLR4, IL-1R and TNFR. 556

These observations are mechanistically significant as it is known that production and 557

secretion of pro-inflammatory cytokines are governed by the TLR/NF-κB signaling 558

pathway, perceived as a major "gateway" cascade in innate immunity, and are 559

associated with the pathogenesis of severe inflammation that promotes lung, liver, and 560

intestinal injuries[44]. For example, binding of LPS to TLR-4 was shown to induce lung 561

parenchymal damage, neutrophil accumulation in the interstitial and alveolar 562

compartments, elevated vascular permeability and pulmonary edema [53,54]. 563

Further molecular insight upon the underlying immunomodulatory mechanism 564

of tryptophol acetate and tyrosol acetate was furnished by analysis of A20 protein levels 565

(Figure 6). A20 is a prominent negative feedback regulator of NF-κB signaling. It has 566

been reported that mice genetically deficient in A20 develop severe inflammation, 567

underscoring the central role of A20 in suppression of NF-kB-dependent inflammation 568

and tissue homeostasis [55]. Importantly, we found that the tryptophol acetate and tyrosol   569

acetate mixtures significantly induced, in both gene and protein levels, the expression   570

A20 (Figure 6 C,D). Thus, our data suggest that the protection furnished against severe   571

inflammation by tryptophol acetate and tyrosol acetate may be mediated by triggering   572

A20 expression and concomitant inhibition of the TLR4/IL-1R/TNFR-NF-κB pathway.   573

Due to our observations that administration of tryptophol acetate and tyrosol   574

acetate orally gave rise to the pronounced anti-inflammatory effects, we assessed the   575

effects of the molecules on the gut microbiota (Figure 7). The relationship between   576

severe inflammation and microbial dysbiosis has been extensively studied in recent   577

years [56,57]. The gut microbiota have been shown to enhance host immunity to pathogens,   578

and dysbiosis has been linked to increased susceptibility of severe inflammation [58].   579

Accordingly, we characterized the gut microbial compositions of mice prior and after   580

LPS injection, and with / without treatment with the mixture of tryptophol acetate and   581

tyrosol acetate. Indeed, oral administration of the molecules had a significant impact on   582

the community composition, manifested by an increase in the abundance of bacteria   583

with anti-inflammatory properties [59]. Notably, the predominant change recorded was   584

the increase in the genus *Bacteroides* in mice that were treated with the molecules 24   585

hours after LPS administration (Figure 7E).   586

Recent studies have reported on the anti-inflammatory properties of *Bacteroides*   587

[60]. The mechanisms of action proposed for these anti-inflammatory activities include   588

inhibition of pathogen colonization [44] and increased mucosal barrier by modifying   589

goblet cells and mucin glycosylation [61]. The higher abundance of *Bacteroides* upon   590

treating the LPS-injected mice with tryptophol acetate and tyrosol acetate may thus   591

account to "cross-talk" between the molecules and host microbiota, promoting   592

intestinal homeostasis. Moreover, the significance of these results are further   593

underlined by a recent study stressing the need for substances that protect gut   594
microbiota such as *Bacteroides* from the collateral damage of widely used antibiotics   595
such as tetracyclines and macrolides [62]. Thereby, consumption of tryptophol acetate and   596
tyrosol acetate in combination with antibiotics may present a new approach for reducing   597
the harmful side effects of antibiotics on host gut microbiome.   598

Overall, varied food-extracted substances have been touted as exhibiting   599
powerful antioxidant and anti-inflammatory properties [63]. However, reports presenting   600
comprehensive clinical, physiological, and mechanistic analyses of the anti-   601
inflammatory activities of probiotic yeast secreted metabolites, such as presented here   602
for tryptophol acetate and tyrosol acetate, have been rare. In particular, to the best of   603
our knowledge, no previous studies have shown an exceptional clinical outcome (i.e.   604
survival and morbidity results, Figure 2), combined with a comprehensive mechanistic   605
analysis. Indeed, our findings uncover, in detail, the immune pathways underlying anti-   606
inflammatory effects of probiotic-sourced molecules.   607

In conclusion, we discovered that metabolites secreted by a probiotic yeast   608
abundant in a milk-fermented microorganism mixture – tryptophol acetate and tyrosol   609
acetate –significantly inhibit systemic severe inflammation effects. Comprehensive *in   610
vivo, ex-vivo,* and *in vitro* data demonstrate the healing and protective effects of the two   611
molecules, including prevention of mortality in mice experiencing LPS-induced severe   612
inflammation, blocking organ damage, reduction of ROS production, and maintaining   613
healthy peripheral blood cell profile. Particularly important, the experiments   614
demonstrate that, unlike many anti-inflammatory treatments, tryptophol acetate and   615
tyrosol acetate did not induce immunosuppression, but rather effectively restored   616
baseline levels of pro-inflammatory cytokines and inhibited activation of NF-κB   617
through elevating A20 expression. Overall, the remarkable systemic anti-inflammatory   618

effects of tryptophol acetate and tyrosol acetate and their presence in a food source that has been consumed by humankind for millennia may open new avenues for anti-inflammatory therapeutics.

Materials and Methods

Materials

Dulbecco's Modified Eagle Medium (DMEM), Roswell Park Memorial Institute (RPMI 1640), heat-inactivated fetal bovine serum (FBS) were purchased from Biological Industries (Beit Haemek, Israel). Penicillin, streptomycin, ECL western blotting substrate, opti-MEM, and Bradford reagent were purchased from Thermo-fisher. Dihydrochloride (AAPH), 2,2-Diphenyl-1-picrylhydrazyl (DPPH), 2,20 –azobis (2-amidinopropane, fluorescein, hydrogen peroxide, sodium citrate, acetonitrile, formic acid, sulfuric acid, ethanol, methanol, sodium carbonate, lipopolysaccharide (from E. coli O55:B5, L2880), propidium iodide (PI) (Solution P4864), NaCl, Triton X-100, BSA, NaF, Na_3VO_4, AEBSF, polyethylenemine (PEI), Leupeptin, and Aprotinin were purchased from Sigma-Aldrich. Xylene was purchased from Epredia (lot 6601). Tris buffer was purchased from Bio-lab (Jerusalem, Israel). Thioglycolate was purchased from life technologies, 2085262 (ultra-pure grade). All reagents and solvents were of analytical grade and were used as received.

DPPH scavenging activity

The effect of tryptophol and tyrosol acetates on DPPH• radical were estimated according to recommendations of Marinova and Batchvarov [24] with some modifications. All of the solutions were prepared in ethanol. The stock solution was prepared by dissolving 13.8 mg DPPH with 20 mL ethanol and stored until needed. The

control (100%) solution was obtained by mixing 225 µL ethanol with 25 µL stock   643

solution to obtain an absorbance of 1.0 ± 0.1 units at 490 nm. 200 µL of tryptophol   644

acetate and tyrosol acetate dissolved in ethanol in a concentration of 200 µM was   645

allowed to react with 25 µL of the DPPH solution for 20 min in the dark at 400 rpm at   646

25ºC. Ethanol (250 µL) was used for the blank control (100%), and 225 µL was used   647

as a blank. The absorbance decrease was recorded at 490 nm. For all evaluated assays,   648

absorbance measurements were performed in triplicate using a Microtiter Plate Reader   649

(Varioskan Flash, Thermo) to calculate radical scavenging activity (% of inhibition)   650

with the formula.   651

Inhibition (%) = 1 – Abs (sample) – Abs (blank) x100   652
                              Abs (control) – Abs (blank)

where Abs (sample) was the absorbance of the reaction in the presence of the sample   654

(sample dilution + DPPH solution), Abs (blank) was the absorbance of the blank for   655

each sample dilution (sample dilution + DPPH solvent), and Abs (control) was the   656

absorbance of the control reaction (sample solvent + DPPH solution).   657

Oxygen radical absorbance capacity (ORAC assay)   658

The ORAC method is based on the oxidative degradation of the fluorescent molecules   659

after mixing with a free radical generator, for example, azo compounds. This method   660

determines the ability of the sample to neutralize short-lived free radicals. The assay   661

was carried out according to Ou et al. [25], with some minor modifications. Prior to the   662

measurements, tryptophol acetate and tyrosol acetate were dissolved in acidified   663

methanol to a concentration of 200 µM. 30 µL of each molecule or 30 µL mixture of   664

both, as well as blank (methanol) were mixed with 180 µL of 112 nM fluorescein   665

solution in a 96-well plate and incubated at 37°C for 15 min. Subsequently, 100 µL of   666

100 mM 2,2' -azobis(2-amidinopropane) dihydrochloride (AAPH) solution was added, 667

and the fluorescence was measured every 70 s for 90 min using a Microtiter Plate 668

Reader (Varioskan Flash, Thermo). The excitation wavelength was 485 nm and the 669

emission recorded at 520 nm. All stock solutions and dilutions of samples were 670

prepared fresh daily in phosphate-buffered saline (PBS, pH 7.4). The experiments were 671

done in six repetitions. Results are presented according to the following calculations : 672

Area under the curve (AUC) was calculated for each sample using the equation: 673

$AUC = 1 + FU_1/FU_0 + FU_2/FU_0 + FU_3/FU_0 \ldots\ldots+$ 674

$FU_0 =$ fluorescence at time zero. 675

$FU_x =$ fluorescence at specific timepoints (e.g., FU_3 is the fluorescence value at three 676

minutes). Net AUC was calculated by subtracting the Blank AUC from the AUC of 677

each sample using the equation: 678

Net AUC = AUC (Antioxidant) – AUC (blank) 679

Animal studies 680

C57BL/6 mice 681

All experiments were performed in accordance with the Institutional Animal Care and 682

Use Committee at Ben-Gurion University (IL-43-07-2020). Eight-week-old C57BL/6 683

mice (Harlen, Rehovot, Israel) weighing approximately 20 gr were held in specific 684

pathogen-free conditions at the University Central Research Facility for at least one 685

week prior to commencing studies. The mice were fed (*ad libitum*) with V1154-703 686

ssniff[TM] (Soest, Germany) and were allowed water continuously. Environmental 687

enrichment was provided to all animals based on the requirements of the particular 688

mouse strain. Temperature was maintained at 21°C, and animals were exposed to a 12 689

h light, 12 h dark cycle with a 15 min ramp-up and ramp-down to simulate dusk and 690

dawn. The mice were divided into two study groups: untreated (mice injected with LPS 691

and administered a placebo), Treated (mice injected with LPS and treated by oral 692

administration of mixture of tryptophol acetate and tyrosol acetate). Mice were housed 693

in groups of six per cage. LPS dose was 30 mg/kg to each mouse (~20 g), which was 694

injected intraperitoneally. 695

Treatment with tryptophol acetate and tyrosol acetate 696

Tryptophol acetate and tyrosol acetate were synthesized as previously described [22] and 697

dissolved in DDW to desired concentrations (final concentration of 150 µg/kg). The 698

molecular mixture was administrated through oral gavage in a volume of 200 µL twice 699

per day, in the morning (08:30–09:30) and evening (19:30–20:30). DDW was used as 700

a placebo and was administrated orally to the control group at the same times. 701

In one set of experiments, the mixture of molecules was administrated concomitantly 702

with LPS injection and treatment was continued two times per day until the end time of 703

the experiment (156 h). In another set of experiments, the first treatment with the 704

molecules started 28 h after LPS injection and was continued two times per day until 705

the end time of the experiment (156 h). 706

Clinical evaluation of inflammation 707

Clinical signs of inflammation were recorded twice daily. Measurement of body weight, 708

signs of diarrhea and appearance of blood traces in stool, as well as signs of dyspnea 709

were assessed. Mean percentage of weight loss was calculated daily as the ratio of 710

measured body weight and weight measured before injection of LPS. Every 6 h after 711

LPS injection till endpoint of the experiments (till 72 h), mice were anesthetized with 712

5% isoflurane (2-chloro-2-(difluoromethoxy)-1,1,1-trifluoro-ethane) and blood 713

samples were got from tail vien. Lungs, livers, colons, and intestinals were obtained at 714

the same time intervals and were cleaned from contents by flushing with 10 ml of sterile 715

PBS. The tissue were taken for extraction of RNA and part of them were embedded in 716

paraffin following a standard protocol. 5-µm sections using a rotary microtome (MRC- 717

Lab, HIS-202A). Paraffin embedded tissues were cut into 5-µm sections using a rotary 718

microtome (MRC-Lab, HIS-202A), rehydrated and stained with hematoxylin. 719

Immunohistochemistry Staining 720

Tissue sections were de-paraffinized in xylene and re-hydrated with decreasing 721

concentrations of alcohol. Subsequently, endogenous peroxide was blocked with 722

hydrogen peroxide, and antigen retrieval was achieved by treating sections with 0.01M 723

sodium citrate, pH 6.0 for 1 min in a pressure cooker. After blocking with universal 724

blocking solution (ZYMED Laboratories, San Francisco, CA), tissue sections were 725

stained with the designated primary antibodies. A Vectastain Elite ABC Peroxidase kit 726

or Universal ImmPRESS kit (Vector Laboratories, Burlingame, CA) were used for 727

secondary antibodies, and visualization was performed using 3-amino-9- 728

ethylcarbazole (AEC) as a substrate (ZYMED Laboratories, San Francisco, CA). 729

Sections were then stained with hematoxylin for counterstaining and mounted using 730

VectaMount AQ Aqueous Mounting Medium (Cat-No. H-5501; Vector Laboratories). 731

Myeloperoxidase (MPO)-positive (Abcam), F4/80-positive (Santa Cruz, 377009), and 732

LY6C-positive cells (Abcam, ab15627) were counted in stained sections in six 733

randomly chosen fields (×200), and bars are indicated standard errors of the means. 734

Blood count

Blood samples were collected from mice tails with EDTA micropipette capillaries
(Exigo) and analyzed directly after collection by a veterinary hematology analyzer
(Exigo H400, Boule Medical AB, Spånga, Sweden).

Cytokine measurements by ELISA

To measure the extent of the inflammatory response in mice, serum obtained from the
blood of LPS-treated and untreated mice, we customized a highly sensitive milliplex®
MAP Kit (Cat # MPXMCYTO-70K; Millipore) with color-coded beads and fluorescent
dyes according to the manufacturer's recommendations.

In vitro macrophage experiments

264.7 cell line

The RAW 264.7 cell lines (third passage) were purchased from American Type Culture
Collection (ATCC1 TIB-71™). Cells were cultured in DMEM with 10% FBS and 1%
penicillin/streptomycin, in an atmosphere of 5% CO_2 and 95% humidity at 37°C. Cells
were regularly tested for mycoplasma contamination.

Peritoneal macrophages

Mouse peritoneal macrophages were obtained from C57BL/6J mice (6–8weeks old) 72
hours after intra-peritoneum injection of Thioglycolate and cultured in complete RPMI
medium with and without LPS (100ng/ml) (O55:B5, L2880, Sigma) and in the presence
or absence of the tryptophol acetate and tyrosol acetate (1+2 mixture).

Cytokine production in peritoneal macrophages

Cells (3×10^5) were seeded in 24-well plates and after 24 h, supernatants were collected
and assessed for cytokine secretion. To measure the extent of the inflammatory
response in the LPS stimulated peritoneal macrophages, we used murine ELISA kits
according to the manufacturers' recommendations for IL-1β (PeproTech,, 900-K47),
IL-6 (Biotest, DY406), TNF-α (Biotest, DY410), TGF-β (Biotest, DY1679), IL-1α
(Biotest, MAB400-500; AB-monoclonal BAF400, AB-polyclonal 400-ML-005), and
INF-β (BioLegend B292426).

ROS analysis

Peritoneal macrophage cells were seeded (1×10^5 cells) into 96-well plates for ROS and
phagocytosis analysis.

After 2 h of incubation in complete medium at 37°C, cells were stimulated with LPS
(100ng\ml) and incubated overnight in the absence or presence of tryptophol acetate
and tyrosol acetate (100 μM each). After 24 hours, the cells were washed two times
with PBS and incubated for 1 h with CellROX Deep Red Reagent (Invitrogen, Carlsbad,
CA) prior to analysis by flow cytometry. Dead cells were gated out using PI. The
presence of ROS was analyzed by the fluorescence geometric mean (GM) of the
CellROX Deep Red fluorescence.

Phagocytosis analysis

Peritoneal macrophage cells were seeded in 96-well plates with RPMI medium. After
2 adherences at 37°C, all the cells were stimulated with LPS (100ng\ml) and incubated
overnight in the absence or presence of the tryptophol acetate and cytosol acetate
mixture (100 μM each). After 24 hours the cells were washed two times with PBS and

incubated for 1 h or 3 h at 37°C in 200 μl RPMI medium, alone (control) or with 783

Fluoresbrite® Yellow-Green carboxylate microspheres (1 μm) (Polysciences Inc., 784

15702) in a ratio of 3:1 beads-to-cells. In the flow cytometry analysis, dead cells were 785

gated out using PI. Free beads were analyzed alone and gated out from the analysis. 786

The fraction of live cells positive for beads was used to estimate phagocytosis 787

functionality. 788

Western blot analysis for NF-κB expression 789

Protein expression was monitored using Western blot analysis. 10×10^6 murine 790

peritoneal macrophages were seeded on petri dishes. After overnight incubation at 37°C 791

the cells were stimulated with LPS (100ng/ml) in the absence or presence of the mixture 792

of tryptophol acetate and tyrosol acetate (100 μM each molecule) and incubated for 30 793

minutes and harvested in lysis buffer (50 mL 3M NaCl, 25 mL 1M Tris pH 7.5, 10 mL 794

Triton X-100, and 10 mL 0.5M EDTA were mixed and added to 905 mL distilled water) 795

in the presence of protease inhibitor cocktail (1:50 complete, Sigma-Aldrich, Israel). 796

Lysates were placed on ice for 30 min and then centrifuged for 30 min (12,000 rpm) at 797

4°C. Supernatants (cytosolic fractions) were collected, and protein concentrations were 798

determined using a Bio-Rad protein assay kit (lot 5000202, Israel). SDS sample buffer 799

was added, and samples were boiled for 5 min and then frozen at -20°C until use. NF- 800

κB (after 30 minutes) was assayed using cell lysates (70μg), separated on SDS-PAGE 801

and analyzed using the following: antibody raised against the anti-NF-κB (p-65) 802

(1:1000; abcam, ab32536), β-actin (1:1000; MP Biomedicals, Santa Ana, CA). 803

Real-time PCR 804

Total mRNA from obtained tissues and RAW 264.7 macrophages was extracted using 805

a RNA extraction kit (ISOLET II RNA Mini Kit, Bioline). cDNA was synthesized from 806

1 μg of RNA using a PrimeScriptTM RT reagent kit (Lifegene, Bio-52073). Subsequent real-time PCR was performed with an iCycler iQ™ Real-Time PCR Detection System (Bio-RAD). PCR results were analyzed with SDS 2.02 software (Applied Biosystems, Thermo). The level of target gene expression was calculated following normalization of the GAPDH gene level in each sample and presented as relative units. Quantitative PCR was performed with Taqman Master Mix (Rhenium, Israel) for: TLR4 (Cat 4453320, Assay ID Mm00445273_m1), Tnfrsf1b (Cat 4453320, Assay ID Mm00441889_m1), GAPDH (Cat 4453320, Assay ID Mm99999915_g1), and IL-1R (Cat 4453320, Assay ID Mm00434237_m1) or with Mix SYBR Green Master Mix (Applied Biosystems).

Gene	Forward sequences (5′–3′)	Reverse sequences (5′–3′)
IL-1α	GTTACAGTGAAACGAAGACTAC	TGCAAGTCTCATGAAGTGAGC
IL-6	TTCCTCTCTGCAAGAGACTTC	GTTGGGAGTGGTATCCTCTGT
IL-1β	TGCCACCTTTTGACAGTGATG	TGTGCTGCTGCGAGATTTGA
HPRT	AGTCCCAGCGTCGTGATTAG	TGGCCTCCCATCTCCTTCAT

Table 1. Primers used for RT-qPCR of target gene

A20 expression in HEK-293 T cells

Cells: HEK-293 T cells were grown and maintained in Dulbecco's Modified Eagle's Medium (DMEM) at 37 °C and 5% CO_2, supplemented with 10% fetal calf serum and antibiotics (50 units/ml penicillin, 50 μg/ml streptomycin).

Antibodies and plasmids

Mouse monoclonal antibodies (mAbs) specific for A20 (Santa Cruz Biotechnology, Inc., Santa Cruz, CA). Mouse anti-β-actin mAbs (Merck Millipore, Darmstadt, Germany). Horseradish peroxidase (HRP)-conjugated goat anti-mouse Abs (Abcam Biotechnology). The full-length human or mouse A20 cDNA with a human influenza hemagglutinin (HA) tag cloned in a pCAGGA mammalian expression vector (HA-A20) was kindly provided by Dr. Shai Cohen (Cancer and Vascular Biology Research Center, The Rappaport Faculty of Medicine and Research Institute, Technion, Israel), previously described [42] .

Transient transfection in HEK-293 T cells

HEK-293 T cells were chosen for transient plasmid transfection. Briefly, 5×10^5 HEK-293 T cells were plated and left to grow for 24 hours in the incubator (37 °C and 5% CO_2). The following day, old DMEM medium was replaced with serum free-fresh DMEM and the cells were left in the incubator for 30 minutes. DNA (plasmids) samples were prepared for transfection; 1.5 μg of purified HA-A20 plasmid DNA was added to 250 μL of Opti-MEM medium, then 5 μL of (final concentration of 5 μg/mL) of polyethylenimine (PEI) was added to the DNA, vortexed and incubated at RT for 20 minutes. The PEI-DNA mix was, added to the cells steadily drop by drop and the cells were left in the incubator (37 °C and 5% CO_2) for 4 hours. After the transfection, the cells were incubated with the Molecule mix (200 μM each) for time range of 1 and 2 h. Cells were harvested at indicated time points and lysed in lysis buffer containing 25mM Tris HCl, pH 7.5, 150mM NaCl, 5mM EDTA, 1mM Na_3VO_4, 50mM NaF, 10μg/ml each of leupeptin and aprotinin, 2mM AEBSF and 1% Triton X-100) followed by 30 min incubation on ice. Lysates were centrifuged at 13,000 rpm for 30 min at 4°C and

the nuclear-free supernatants were used further analysis. 20 µg of protein was resolved on 10% SDS-PAGE and blotted on to nitrocellulose membrane. Membranes were immunoblotted with mouse anti-A20 (1:2000) and mouse anti-β-actin mAbs (1:500) as indicated and visualized following ECL exposure.

Protein concentration

Protein concentration was measured by the method of Bradford using bovine serum albumin as standard.

Electrophoresis and immunoblotting

Whole cell lysates were resolved by electrophoresis on 10 % polyacrylamide gels using Bio-Rad Mini-PROTEAN II cells. Gel proteins were electroblotted onto nitrocellulose membranes (Schleicher and Schuell) at 100 V for 1 h, using BioRad Mini Trans-Blot transfer cells. After 1 h of membrane blocking with 3% BSA in TBS at 37 °C, the membranes were incubated with the indicated primary Abs followed by extensive washings in TBST and incubation with HRP-conjugated goat anti-mouse. Immunoreactive protein bands were visualized using the enhanced chemiluminescent.

Data analysis

All results are expressed as the mean ± SD or mean ±SE, as indicated. Data for all in-vivo and in-vitro experiments were analyzed by a one/two-way ANOVA followed by Tukey post-hoc test using GraphPad Prism (GraphPad Software, San Diego, Ca). A p-value of ≤ 0.05 was considered statistically significant.

Fecal microbiome analysis

Total DNA was extracted from fecal samples using a PureLink microbiome DNA extraction kit (Invitrogen, Carlsbad, CA) followed by a 2-min bead-beating step, as previously described by Shouval, R. et al. [64]. Purified DNA was PCR amplified using PrimeSTAR Max (Takara-Clontech, Shiga, Japan) for the variable V4 region (using 515F-806R barcoded primers) of the 16S rRNA gene. Amplicons were purified using Agencourt AMPure XP magnetic beads (Beckman-Coulter, Brea, CA), and subsequently quantified using a Quant-It Picogreen dsDNA quantitation kit (Invitrogen, Carlsbad, CA). Equimolar DNA amounts from individual samples were pooled and sequenced using the Illumina MiSeq platform at the Genomic Center, Azrieli Faculty of Medicine, Bar-Ilan University, Israel. Sequencing data were processed using QIIME2 version 2019.10. [65]. Single end sequences with a similarity $\geq 99\%$ were assigned as the same feature. Taxonomy was assigned using the GreenGenes database [66]. Chimeric sequences were removed with DADA2 (--p-trunc-len 160) [67]. Samples with <1000 features and features with total frequency <2 were filtered out. Rarefaction was done using 5,000 sequences per sample. Alpha and beta diversity were calculated based on rarefied datasets.

Statistical analysis of microbiome results

The distribution pattern of quantitative variables was examined using a Shapiro-Wilk test. Quantitative variables were compared between groups with a Wilcoxon U-test, for non-normally distributed data, respectively. All tests were 2-tailed, and in all, $p \leq 0.05$ was considered significant . Bacterial alpha diversity was assessed by the Shannon index. Statistical significance of microbial alpha diversity differences was confirmed using a Kruskal–Wallis test, followed by paired Mann-Whitney tests, with Benjamini-

Hochberg correction for the false discovery rate. Beta diversity was calculated using 896

weighted UniFrac distances. Statistical significance was confirmed using a 897

permutational multivariate analysis of variance (PERMANOVA). Differences in 898

relative abundances of bacterial taxa between groups were identified using the ANCOM 899

method [65]. 900

Acknowledgements 902

We thank Dr Shai Cohen (Cancer and Vascular Biology Research Center, The 903

Rappaport Faculty of Medicine and Research Institute, Technion - Israel) for providing 904

the A20-plasmid, and his help with the A20 experiments**.** 905

Author contributions 906

Conceived and designed the experiments O.M., R.M, E.V. and R.J.; experiments and 907

data analysis O.M., R.M., M.B., M.K., J.S., E.T., E.S., and B.R.; writing—original draft 908

O.M., R.M., E.V. and R.J.; writing - reviewed and edited the manuscript B.R., O.K., 909

and E.V.; All authors read and approved the final manuscript. 910

Disclosures 911

A company - Biotic Therapeutics - has been co-founded by O.M. and R.J. based on this 912

research. 913

References 914

1.   Chen, L. *et al.* Inflammatory responses and inflammation-associated diseases in   915
      organs. *Oncotarget* **9**, 7204–7218 (2018).   916
2.   Moriyama, K. & Nishida, O. Targeting cytokines, pathogen-associated molecular   917
      patterns, and damage-associated molecular patterns in sepsis via blood purification.   918
      Int. J. Mol. Sci. **22**, (2021).   919
3.   Akira, S. & Hemmi, H. Recognition of pathogen-associated molecular patterns by   920
      TLR family. *Immunol. Lett.* **85**, 85–95 (2003).   921
4.   Costa, R. T. *et al.* T helper type cytokines in sepsis: time-shared variance and   922

correlation with organ dysfunction and hospital mortality. *Brazilian J. Infect. Dis.* **23**, 79–85 (2019).

5. Melo, A. K. G. *et al.* Biomarkers of cytokine storm as red flags for severe and fatal COVID-19 cases: A living systematic review and meta-analysis. *PLoS One* **16**, 1–21 (2021).

6. Hotchkiss, R. S. *et al.* Sepsis and septic shock Richard. *Physiol. Behav.* **176**, 139–148 (2017).

7. Furman, D. *et al.* Chronic inflammation in the etiology of disease across the life span. *Nat. Med.* **25**, 1822–1832 (2019).

8. Huang, Q. *et al.* Targeting inflammation and cytokine storm in COVID-19. *Pharmacol. Res.* **159**, 105051 (2020).

9. Yang, L. *et al.* The signal pathways and treatment of cytokine storm in COVID-19. *Signal Transduct. Target. Ther.* **6**, 1–20 (2021).

10. Niva, M. 'All foods affect health': Understandings of functional foods and healthy eating among health-oriented Finns. *Appetite* **48**, 384–393 (2007).

11. Dirienzo, D. B. Effect of probiotics on biomarkers of cardiovascular disease: Implications for heart-healthy diets. *Nutr. Rev.* **72**, 18–29 (2014).

12. Luna, R. A. & Foster, J. A. Gut brain axis: Diet microbiota interactions and implications for modulation of anxiety and depression. *Curr. Opin. Biotechnol.* **32**, 35–41 (2015).

13. Hewlings, S. J. & Kalman, D. S. Curcumin: A review of its effects on human health. *Foods* **6**, 1–11 (2017).

14. Meng, X., Zhou, J., Zhao, C. N., Gan, R. Y. & Li, H. Bin. Health benefits and molecular mechanisms of resveratrol: A narrative review. *Foods* **9**, 1–27 (2020).

15. Proestos, C. The Benefits of Plant Extracts for Human Health. *Foods* **9**, 1653 (2020).

16. Giada, M. de L. R. Food Phenolic Compounds: Main Classes, Sources and Their Antioxidant Power, Oxidative Stress and Chronic Degenerative Diseases - A Role for Antioxidants. *Oxidative Stress Chronic Degener. Dis. - A Role Antioxidants* 87–112 (2013).

17. Mennen, L. I., Walker, R., Bennetau-Pelissero, C. & Scalbert, A. Risks and safety of polyphenol consumption. *Am. J. Clin. Nutr.* **81**, 326–329 (2005).

18. Franco, M. C. *et al.* Administration of kefir-fermented milk protects mice against Giardia intestinalis infection. *J. Med. Microbiol.* **62**, 1815–1822 (2013).

19. Vinderola, C. G. *et al.* Immunomodulating capacity of kefir. *J. Dairy Res.* **72**, 195–202 (2005).

20. Malka, O. *et al.* Cross-kingdom inhibition of bacterial virulence and communication by probiotic yeast metabolites. *Microbiome* **9**, 1–15 (2021).

21. Rothschild, D. E., McDaniel, D. K., Ringel-Scaia, V. M. & Allen, I. C. Modulating inflammation through the negative regulation of NF-κB signaling. *J. Leukoc. Biol.* **103**, 1131–1150 (2018).

22. Jelinek R, Malka O, Kalson D, Meijler MM, K. A. Microorganism mixtures, Molecules Derived Therefrom, and Methods of Use Thereof WO 2020/031191. (2020).

23. Bardaweel, S. K. *et al.* Reactive oxygen species: The dual role in physiological and pathological conditions of the human body. *Eurasian J. Med.* **50**, 193–201 (2018).

24. Marinova, G. & Batchvarov, V. methods DPPH. *Bulg. J. Agric. Sci.* **17**, 11–24 (2011).

25. Ou, B., Hampsch-Woodill, M. & Prior, R. L. Development and validation of an improved oxygen radical absorbance capacity assay using fluorescein as the fluorescent probe. *J. Agric. Food Chem.* **49**, 4619–4626 (2001).

26. Anderberg, S. B., Luther, T. & Frithiof, R. Physiological aspects of Toll-like receptor 4 activation in sepsis-induced acute kidney injury. *Acta Physiol.* **219**, 575–590 (2017).

27. Parker, S. J. & Watkins, P. E. Experimental models of Gram-negative sepsis. 22–30 (2001).

28. Shimabukuro-Vornhagen, A. *et al.* Cytokine release syndrome. *J. Immunother. Cancer* **6**, (2018).

29. Arora, M., Poe, S. L., Ray, A. & Ray, P. LPS-induced CD11b+Gr1intF4/80+ regulatory myeloid cells suppress allergen-induced airway inflammation. *Int. Immunopharmacol.* **11**, 827–832 (2011).

30. Hirata, F. *et al.* Insulin enhances leukocyte-endothelial cell adhesion in the retinal microcirculation through surface expression of intercellular adhesion molecule-1. *Microvasc. Res.* **69**, 135–141 (2005).

31. Abraham, E. Neutrophils and acute lung injury. *Crit. Care Med.* **31**, 1–5 (2003).

32. O'Dea, K. P. *et al.* Mobilization and Margination of Bone Marrow Gr-1 high Monocytes during Subclinical Endotoxemia Predisposes the Lungs toward Acute Injury . *J. Immunol.* **182**, 1155–1166 (2009).

33. O'Dea, K. P. *et al.* Lung-marginated monocytes modulate pulmonary microvascular injury during early endotoxemia. *Am. J. Respir. Crit. Care Med.* **172**, 1119–1127 (2005).

34. Uderhardt, S. *et al.* 12/15-Lipoxygenase Orchestrates the Clearance of Apoptotic Cells and Maintains Immunologic Tolerance. *Immunity* **36**, 834–846 (2012).

35. Jabbour, H. N., Sales, K. J., Catalano, R. D. & Norman, J. E. Inflammatory pathways in female reproductive health and disease. *Reproduction* **138**, 903–919 (2009).

36. Grivennikov, S. I., Greten, F. R. & Karin, M. Immunity, Inflammation, and Cancer. *Cell* **140**, 883–899 (2010).

37. Mittal, M., Siddiqui, M. R., Tran, K., Reddy, S. P. & Malik, A. B. Reactive oxygen species in inflammation and tissue injury. *Antioxidants Redox Signal.* **20**, 1126–1167 (2014).

38. Oeckinghaus, A. and & Ghosh, S. The nuclear factor- k B (NF-k B) family of transcription factors. *Cold Spring Harb. Perspect. Biol.* **1**, 1–15 (2009).

39. Verstrepen, L. *et al.* TLR-4, IL-1R and TNF-R signaling to NF-κB: Variations on a common theme. *Cell. Mol. Life Sci.* **65**, 2964–2978 (2008).

40. Kawai, T. & Akira, S. The role of pattern-recognition receptors in innate immunity: Update on toll-like receptors. *Nat. Immunol.* **11**, 373–384 (2010).

41. Hitoshi, N., Ken-ichi, Y. & Jun-ichi, M. Efficient selection for high-expression transfectants with a novel eukaryotic vector. *Gene* **108**, 193–199 (1991).

42. Lapid, D., Lahav-Baratz, S. & Cohen, S. A20 inhibits both the degradation and limited processing of the NF-κB p105 precursor: A novel additional layer to its regulator role. *Biochem. Biophys. Res. Commun.* **493**, 52–57 (2017).

43. Schirmer, M. *et al.* Linking the Human Gut Microbiome to Inflammatory Cytokine Production Capacity. *Cell* **167**, 1125-1136.e8 (2016).

44. Hiippala, K. *et al.* Isolation of anti-inflammatory and epithelium reinforcing bacteroides and parabacteroides spp. From a healthy fecal donor. *Nutrients* **12**, (2020).

45. Huang, M., Cai, S. & Su, J. The pathogenesis of sepsis and potential therapeutic targets. *Int. J. Mol. Sci.* **20**, (2019).

46. Rosseau, S. *et al.* Phenotypic characterization of alveolar monocyte recruitment in acute respiratory distress syndrome. *Am. J. Physiol. - Lung Cell. Mol. Physiol.* **279**, 25–35 (2000).

47. Kumar, V. Toll-like receptors in sepsis-associated cytokine storm and their endogenous negative regulators as future immunomodulatory targets. *Int. Immunopharmacol.* **89**, 107087 (2020).

48. Huang, L. *et al.* Sepsis-associated severe interleukin-6 storm in critical coronavirus disease 2019. *Cell. Mol. Immunol.* **17**, 1092–1094 (2020).

49. Reczek, C. R. & Chandel, N. S. ROS-dependent signal transduction. *Curr. Opin. Cell Biol.* **33**, 8–13 (2015).

50. Forrester, S. J., Kikuchi, D. S., Hernandes, M. S., Xu, Q. & Griendling, K. K. Reactive oxygen species in metabolic and inflammatory signaling. *Circ. Res.* **122**, 877–902 (2018).

51. Hidalgo H., Humberto A., Helmke R., German V. The effect of cyclosporine and dexamethasone on an alveolar macrophage cell line (NR8383). *Transplantation* **53**, 620–623 (1992).

52. Pooladanda, V. *et al.* Nimbolide protects against endotoxin-induced acute respiratory distress syndrome by inhibiting TNF-α mediated NF-κB and HDAC-3 nuclear translocation. *Cell Death Dis.* **10**, (2019).

53. Jeyaseelan, S., Chu, H. W., Young, S. K., Freeman, M. W. & Worthen, G. S. Distinct roles of pattern recognition receptors CD14 and Toll-like receptor 4 in acute lung injury. *Infect. Immun.* **73**, 1754–1763 (2005).

54. Zanoni, I. *et al.* CD14 controls the LPS-induced endocytosis of toll-like receptor 4. *Cell* **147**, 868–880 (2011).

55. Lee, E. G. *et al.* Failure to regulate TNF-induced NF-kappaB and cell death responses in A20-deficient mice. *Science* **289**, 2350–4 (2000).

56. Hakansson, A. & Molin, G. Gut microbiota and inflammation. *Nutrients* **3**, 637–687 (2011).

57. Wang, J., Chen, W. D. & Wang, Y. D. The Relationship Between Gut Microbiota and Inflammatory Diseases: The Role of Macrophages. *Front. Microbiol.* **11**, 1–9 (2020).

58. Belkaid, Y. and T. H. Role of the Microbiota in Immunity and inflammation Yasmine. *Cell* **157**, 121–141 (2015).

59. Krasulova, K. & Illes, P. Intestinal interplay of quorum sensing molecules and human receptors. *Biochimie* **189**, 108–119 (2021).

60. Delday, M., Mulder, I., Logan, E. T. & Grant, G. Bacteroides thetaiotaomicron Ameliorates Colon Inflammation in Preclinical Models of Crohn's Disease. *Inflamm. Bowel Dis.* **25**, 85–96 (2019).

61. Wrzosek, L. *et al.* Bacteroides thetaiotaomicron and Faecalibacterium prausnitziiinfluence the production of mucus glycans and the development of goblet cells in the colonic epithelium of a gnotobiotic model rodent. *BMC Biol.* **11**, 61 (2013).

62. Maier, L. *et al.* Unravelling the collateral damage of antibiotics on gut bacteria. *Nature* **599**, 120–124 (2021).

63. Bakasatae, N., Kunworarath, N., Takahashi Yupanqui, C., Voravuthikunchai, S. P. & Joycharat, N. Bioactive components, antioxidant, and anti-inflammatory activities of the wood of Albizia myriophylla. *Rev. Bras. Farmacogn.* **28**, 444–450 (2018).

64. Shouval, R. *et al.* Patterns of salivary microbiota injury and oral mucositis in recipients of allogeneic hematopoietic stem cell transplantation. *Blood Adv.* **4**, 2912–2917 (2020).

65. Bolyen, E. *et al.* Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. *Nat. Biotechnol.* **37**, 852–857 (2019).

66. DeSantis, T. Z. *et al.* Greengenes, a chimera-checked 16S rRNA gene database and workbench compatible with ARB. *Appl. Environ. Microbiol.* **72**, 5069–5072 (2006).

67. Callahan, B. J. *et al.* DADA2: High-resolution sample inference from Illumina amplicon data. *Nat. Methods* **13**, 581–583 (2016).

Supporting information

Tryptophol acetate and tyrosol acetate, metabolites secreted by a probiotic yeast, display remarkable anti-inflammatory effects

ANCOM results after 6 hours			
ASVs	**Healthy**	**Untreated**	**Treated**
Blautia	High	Low	Low
Bacteroides	High	Low	**Low**
Clostridium	High	Low	Low
Sutterella	High	Low	Low
Ruminococcus	Low	High	High
Parabacteroides	High	Low	Low
ANCOM results after 24 hours			
ASVs	**Healthy**	**Untreated**	**Treated**
Blautia	High	Low	Low
Bacteroides	High	Low	**High**
Clostridium	High	Low	Low
Oscillospira	Low	High	High
Ruminococcus	Low	High	High

Table 1 ANCOM results of mice treated and untreated with molecules. Main bacterial taxa abundance after 6 and 24 hours from LPS administration.