Mapping information-rich genotype-phenotype landscapes with genome-scale Perturb-seq

Joseph M. Replogle[1-5,†], Reuben A. Saunders[2-5,†], Angela N. Pogson[3-5], Jeffrey A. Hussmann[3-5], Alexander Lenail[4,5], Alina Guna[5], Lauren Mascibroda[6], Eric J. Wagner[6,7], Karen Adelman[8], Jessica L. Bonnar[3-5], Marco Jost[3,9], Thomas M. Norman[10,*], & Jonathan S. Weissman[3-5,11,12, *]

[1]Medical Scientist Training Program, University of California, San Francisco; San Francisco, CA 94158, USA.
[2]Tetrad Graduate Program, University of California, San Francisco; San Francisco, CA 94158, USA.
[3]Department of Cellular and Molecular Pharmacology, University of California, San Francisco; San Francisco, CA 94158, USA.
[4]Howard Hughes Medical Institute, Massachusetts Institute of Technology; Cambridge, MA 02142, USA.
[5]Whitehead Institute for Biomedical Research, Massachusetts Institute of Technology; Cambridge, MA 02142, USA.
[6]Department of Biochemistry & Molecular Biology, The University of Texas Medical Branch at Galveston; Galveston, TX 77555, USA.
[7]Department of Biochemistry & Biophysics, University of Rochester School of Medicine and Dentistry; Rochester, NY 14642, USA.
[8]Department of Biological Chemistry and Molecular Pharmacology, Blavatnik Institute, Harvard Medical School; Boston, MA 02115, USA.
[9]Department of Microbiology, Harvard Medical School; Boston, MA 02115, USA.
[10]Program for Computational and Systems Biology, Sloan Kettering Institute, Memorial Sloan Kettering Cancer Center; New York, NY 10065, USA.
[11]David H. Koch Institute for Integrative Cancer Research, Massachusetts Institute of Technology; Cambridge, MA 02142, USA.
[12]Department of Biology, Massachusetts Institute of Technology; Cambridge, MA 02142, USA.

†Co-first authors
*Corresponding authors: J.S.W. (weissman@wi.mit.edu) and T.M.N. (normantm@mskcc.org).

Abstract

A central goal of genetics is to define the relationships between genotypes and phenotypes.

High-content phenotypic screens such as Perturb-seq (pooled CRISPR-based screens with single-

cell RNA-sequencing readouts) enable massively parallel functional genomic mapping but, to date,

have been used at limited scales. Here, we perform genome-scale Perturb-seq targeting all

expressed genes with CRISPR interference (CRISPRi) across >2.5 million human cells and present

a framework to power biological discovery with the resulting genotype-phenotype map. We use

transcriptional phenotypes to predict the function of poorly-characterized genes, uncovering new

regulators of ribosome biogenesis (including *CCDC86*, *ZNF236*, and *SPATA5L1*), transcription

(*C7orf26*), and mitochondrial respiration (*TMEM242*). In addition to assigning gene function,

single-cell transcriptional phenotypes allow for in-depth dissection of complex cellular phenomena

– from RNA processing to differentiation. We leverage this ability to systematically identify the

genetic drivers and consequences of aneuploidy and to discover an unanticipated layer of stress-

specific regulation of the mitochondrial genome. Our information-rich genotype-phenotype map

reveals a multidimensional portrait of gene function and cellular behavior.

Main Text

Mapping the relationship between genetic changes and their phenotypic consequence is critical to understanding gene and cellular function. This mapping is traditionally carried out in either of two ways. A phenotype-centric, "forward genetic" approach reveals the genetic changes that drive a phenotype of interest. Conversely, a gene-centric, "reverse genetic" approach catalogs the diverse phenotypes caused by a defined genetic change.

Recent technological developments have advanced both forward and reverse genetic efforts (*1*). CRISPR-Cas tools now enable the deletion, mutation, repression, or activation of genes with ease (*2*). In forward genetic screens, CRISPR-Cas systems can be used to generate cells with diverse genetic perturbations. This pool of perturbed cells can then be subjected to a selective pressure, with phenotypes assigned to genetic perturbations by sequencing. Forward genetic screens provide powerful tools for the identification of cancer dependencies, essential cellular machinery, differentiation factors, and suppressors of genetic diseases (*3–6*). In parallel, dramatic improvements in molecular phenotyping now allow for single-cell readouts of epigenetic, transcriptomic, proteomic, and imaging information (*7*). Applied to reverse genetics, single-cell profiling can refine the understanding of how select genetic perturbations affect cell types and cell states.

However, both phenotype-centric and gene-centric approaches suffer conceptual and technical limitations. Pooled forward genetic screens typically use low-dimensional phenotypes (e.g., growth, marker gene expression, drug resistance) for selection. The use of simple phenotypes can conflate genes acting via different mechanisms, requiring extensive follow-up studies to disentangle genetic pathways (*8*). Additionally, in forward genetics, serendipitous discovery is constrained by the prerequisite of selecting phenotypes prior to screening. On the other hand, while reverse genetic approaches enable the study of multidimensional and complex phenotypes, they have typically been restricted in scale to rationally chosen targets, limiting the ability to make systematic comparisons.

Single-cell CRISPR screens present a solution to these problems. These screens simultaneously read out the genetic perturbation and high-dimensional phenotype of individual cells in a pooled screening format, thus combining the throughput of forward genetic screens with the rich phenotypes of reverse genetics. While these approaches initially focused on transcriptomic phenotypes (e.g., Perturb-seq, CROP-seq) (*9–13*), technical advances have enabled their application to epigenetic (*14*), imaging (*15*), or multimodal phenotypes (*16–18*). From these rich

data, it is possible to identify genetic perturbations that cause a specific behavior as well as to

catalog the spectrum of phenotypes associated with each genetic perturbation. Despite the promise

of single-cell CRISPR screens, their use has generally been limited to studying at most a few

hundred genetic perturbations, typically chosen with a bias towards predefined biological

questions.

We reasoned that there would be unique value to genome-scale single-cell CRISPR

screens. For example, while the number of perturbations scales linearly with experimental cost,

the number of pairwise comparisons in a screen—and thus its utility for unsupervised classification

of gene function—scales quadratically. Similarly, in large-scale screens, the diversity of

perturbations allows one to explore the range of cell states that can be revealed by rich phenotypes.

Additionally, as many human genes are well-characterized, these genes serve as natural controls

to anchor the interpretation of observations in comprehensive datasets. Finally, genome-scale

experiments could help address fundamental biological questions, such as what fraction of genetic

changes elicit global transcriptional phenotypes and how transcriptional programs are rewired

between cell types, with implications for understanding the organizing principles of cellular

systems (*19*).

Here we perform the first genome-scale Perturb-seq screens. We use a compact,

multiplexed CRISPR interference (CRISPRi) library to assay thousands of loss-of-function genetic

perturbations with single-cell RNA-sequencing (scRNA-seq) in chronic myeloid leukemia (K562)

and retinal pigment epithelial (RPE1) cell lines. Leveraging the scale and diversity of these

perturbations across >2.5 million cells, we show that Perturb-seq can be used to study numerous

complex cellular phenotypes—from RNA splicing to differentiation to chromosomal instability—

in a single screen. We demonstrate how the interpretability of scRNA-seq phenotypes enables the

discovery of gene function and extensively validate our findings with orthogonal experiments.

Finally, we invert our analysis to focus on regulatory networks rather than genetic perturbations

and uncover unanticipated stress-specific regulation of the mitochondrial genome. In sum, we use

Perturb-seq to reveal a multidimensional portrait of cellular behavior, gene function, and

regulatory networks that advances the goal of creating comprehensive genotype-phenotype maps.

Results

A multiplexed CRISPRi strategy for genome-scale Perturb-seq

Perturb-seq uses scRNA-seq to concurrently read out the CRISPR single-guide RNAs (sgRNA) (i.e., genetic perturbation) and transcriptome (i.e., high-dimensional phenotype) of single cells in a pooled format (Fig. 1A). To enable genome-scale Perturb-seq, we considered key parameters that would increase scalability and data quality, such as the genetic perturbation modality and sgRNA library.

Although Perturb-seq is compatible with a range of CRISPR-based perturbations including knockout (*10–12*), knockdown (CRISPRi) (*9*), or activation (CRISPRa) (*20*), we elected to use CRISPRi for several reasons. First, CRISPRi allows the efficacy of the genetic perturbation, knockdown, to be directly measured from scRNA-seq. Exploiting this feature allowed us to target each gene in our library with a single element and empirically exclude unperturbed genes from downstream analysis. Second, CRISPRi tends to yield more homogeneous genetic perturbation than nuclease-based CRISPR knockout, which can generate a subset of cells bearing active in-frame indels (*21*). The relative homogeneity of CRISPRi limits selection for unperturbed cells, especially when studying essential genes. Third, unlike nuclease-based gene knockout, CRISPRi does not lead to activation of the DNA damage response which can alter cell state and transcriptional signatures (*22*).

To improve scalability, we optimized our CRISPRi sgRNA libraries. To maximize CRISPRi efficacy, we used multiplexed CRISPRi libraries in which each construct contains two distinct sgRNAs targeting the same gene (table S1-S3; see *Methods*) (*13*). To avoid low representation of sgRNAs targeting essential genes, we performed preliminary growth screens and, during library synthesis, overrepresented constructs that caused strong growth defects (fig. S1A-D).

Next, we devised a three-pronged Perturb-seq screening approach encompassing multiple timepoints and cell types (Fig. 1A). As a primary cell line, we studied chronic myeloid leukemia (CML) K562 cells engineered to express the CRISPRi effector protein dCas9-KRAB (*23*). In this cell line, we performed two Perturb-seq screens: one targeting all expressed genes sampled at day 8 after transduction (n=9,866 genes; n=10,673 total perturbations; some genes have multiple independent transcripts) and another targeting common essential genes, which was sampled at day

6 after transduction (n=2,057 genes; n=2,176 total perturbations). As a secondary cell line, we used

RPE1 cells engineered to express dCas9 fused to a KRAB domain derived from the gene *ZIM3*,

which was recently shown to yield improved transcriptional repression compared to the *KOX1*-

derived KRAB domain used in previous CRISPRi experiments (*24*). In contrast to K562 cells,

RPE1 cells are a non-cancerous retinal pigment epithelial cell line that are hTERT-immortalized,

near-euploid, adherent, and p53-positive. In RPE1 cells, we performed a screen targeting common

essential genes plus a subset of nonessential genes that produced phenotypes in K562 cells sampled

at day 7 after transduction (n=2,393 genes; n=2,549 total perturbations).

We conducted these three screens with 10x Genomics droplet-based 3' scRNA-seq and

direct sgRNA capture (*13*). After sequencing and read alignment, we performed sgRNA

identification and removed any cells bearing sgRNAs targeting different genes, which are an

expected byproduct of lentiviral recombination between sgRNA cassette or doublet encapsulation

during scRNA-seq. In total, we obtained >2.5 million high-quality cells with a median coverage

of >100 cells per perturbation (fig. S1 E-G; table S4-S6). We observed a median target knockdown

of 85.5% in K562 cells and 91.6% in RPE1 cells (Fig. 1B), confirming both the efficacy of our

CRISPRi libraries and the fidelity of sgRNA assignment (*13*). The difference in performance

between these cell lines was likely due to the use of the optimized *ZIM3*-derived KRAB domain

in the RPE1 cells, suggesting that future efforts would benefit from improved CRISPRi efficacy.

A robust computational framework to detect transcriptional phenotypes

The scale of our experiment provided a unique opportunity to ask what fraction of genetic

perturbations cause a transcriptional phenotype, a preliminary requirement for inferring gene

function. Significant transcriptional phenotypes can take many forms, ranging from altered

occupancy of a given cell state to focused changes in the expression level of a small number of

target genes. To contend with this diversity, we created a robust framework capable of detecting

transcriptional changes between groups of cells in our data. Our experimental design included

many control cells bearing diverse non-targeting sgRNAs. These allow for internal *z*-normalization

of expression measurements, and we found that this procedure corrected for batch effects that

resulted from parallelized scRNA-seq and sequencing (fig. S2). As Perturb-seq captures single-

cell genetic perturbation identities in a pooled format, we can use statistical approaches that treat

each cell as an independent experimental sample. In general, we chose to use conservative, non-

parametric statistical tests to detect transcriptional changes rather than making specific
assumptions about the underlying distribution of gene expression levels.

First, we examined global transcriptional changes using a permuted energy distance test
(see *Methods*). We compared cells bearing each genetic perturbation to non-targeting control cells
at the level of principal components (approximating global transcriptional features like cell state
and gene expression programs). Relative to a permuted null distribution, this test asks whether
cells carrying a given genetic perturbation could have been drawn from the control population. By
this metric, we found that 2,987 of 9,608 genetic perturbations targeting a primary transcript
(31.1%) compared to 11 of 585 non-targeting controls (1.9%) caused a significant transcriptional
phenotype in K562 cells.

While sensitive, the energy distance test assays global shifts in expression without
providing insight into which specific transcripts are altered. To detect individual differentially
expressed genes, we applied the Anderson-Darling (AD) test to compare the distribution of
expression levels for each gene in cells bearing each genetic perturbation against control cells.
Importantly, the AD test is sensitive to transcriptional changes in a subset of cells, enabling us to
find differences even when phenotypes have incomplete penetrance. With the AD test, we found
2,935 of 9,608 genetic perturbations targeting a primary transcript (30.5%) compared to 12 of 585
non-targeting controls (2.1%) caused >10 differentially expressed genes in K562 cells. These
results were well-correlated between time points and cell types (fig. S3A,B; tables S4-S6) and
concordant with the energy distance test (78.7% concordance by Jaccard index).

We then explored features of genetic perturbations that predict a transcriptional phenotype.
We found that the strength of the transcriptional response was correlated with the strength of the
growth defect (Spearman's rho = –0.51) with 86.6% of essential genetic perturbations (gamma <
–0.1) leading to a significant transcriptional response in K562 cells (Fig. 1C; fig. S3C,D). A
substantial number of genetic perturbations that cause a transcriptional phenotype nonetheless
have a negligible growth phenotype (n=771; fig. S3E), indicating that many genetic perturbations
influence cell state but not growth or survival. We also found that highly expressed genes were
more likely to produce transcriptional phenotypes (Spearman's rho = 0.42) (fig. S3C).

Considering that some of our genetic perturbations did not yield strong on-target
knockdown, our estimate of the fraction of genetic perturbations that cause a transcriptional
phenotype is likely to be a lower bound. While a fraction of phenotypes may result from off-target
effects, an advantage of Perturb-seq is the ability to directly detect potential off-target activities

such as the knockdown of neighboring genes. Consistent with earlier studies (*25*), we found that

~7.5% of perturbations caused significant knockdown of a neighboring gene in K562 cells, but

neighbor gene knockdown was not enriched in genetic perturbations with a negligible growth

defect that produced a transcriptional phenotype (fig. S4). Taken together, these results present a

coherent picture where knockdown of a significant fraction of expressed genes causes a

transcriptional response.

Annotating gene function from transcriptional phenotypes

Previous Perturb-seq screens have focused on targeted sets of genetic perturbations that are

often related biologically, such as genes identified in forward genetic screens. Our large-scale

screen targeting all expressed genes in K562 cells presented a unique opportunity to assess how

well transcriptional phenotypes can resolve gene function when used in an unbiased manner.

We focused on a subset of 1,973 perturbations that had strong transcriptional phenotypes

(>50 differentially expressed genes by AD test) (Fig. 2A). Because related perturbations could

have different magnitudes of effect, we used the correlation between mean expression profiles as

a scale-invariant metric of similarity.

To assess the extent to which correlated mean expression profiles between genetic

perturbations indicated common function, we compared our results to two curated sources of

biological relationships. First, among the 1,973 targeted genes, there were 327 protein complexes

from the CORUM 3.0 database with at least two thirds of the complex members present,

representing 14,165 confirmed protein-protein interactions (*26*). The corresponding expression

profile correlations were markedly stronger (median correlation 0.61) than the background

distribution of all possible gene pairs (median correlation 0.10) (Fig. 2B). Second, we compared

the correlation between genetic perturbations to the STRING database of known and predicted

protein-protein interactions, which had scores for 243,558 of the possible gene-gene relationships

within our dataset (*27*). High STRING scores, reflecting high-confidence interacting proteins,

were also strongly associated with high expression correlations (Fig. 2C).

We next performed an unbiased search for global structure to group similar perturbations

within the dataset. We identified 64 discrete clusters based on strong intra-cluster correlations and

annotated their function using CORUM, STRING, and manual searches. To visualize the dataset,

we constructed a minimum distortion embedding that places genes with correlated expression
profiles close to each other in the plane and labeled the location of gene clusters (Fig. 2D).

Both the clusters and the embedding showed clear organization by biological function
spanning a diverse array of different processes including: chromatin modification; transcription;
mRNA splicing, capping, polyadenylation, and turnover; nonsense-mediated decay; translation;
post-translational modification, trafficking, and degradation of proteins; central metabolism;
mitochondrial transcription and translation; DNA replication; cell division; microRNA biogenesis;
and major signaling pathways active in K562 cells such as BCR-ABL and mTOR (table S7). We
further annotated the embedding visualization by labeling CORUM complexes and STRING
clusters whose members were placed in nearby positions, revealing structure at finer resolution
such as identifying the SMN complex, exon junction complex, U6 snRNP, and methylosome
within the spliceosome and the association of ribosome biogenesis factors with the 40S ribosomal
subunit.

In our dataset, we identified many poorly annotated genes whose perturbation led to similar
transcriptional responses to genes of known function, naturally predicting a role for the
uncharacterized genes. To orthogonally test a subset of these predictions, we selected ten poorly
annotated genes whose perturbation response correlated (r>0.6) with subunits and biogenesis
factors of either the large or small subunit of the cytosolic ribosome, which formed distinct clusters
in our data (fig. S5A). This included genes that had no previous association with ribosome
biogenesis (*CCDC86*, *CINP*, *SPATA5L1*, *ZNF236*, *C1orf131*) as well as genes that had not been
associated with functional defects in a particular subunit (*SPOUT1*, *TMA16*, *NOPCHAP1*, *ABCF1*,
and *NEPRO*). We used CRISPRi to target these genes in K562 cells and looked for evidence of
ribosome biogenesis defects by assessing the ratio of 28S to 18S rRNA by Bioanalyzer
electrophoresis. Knockdown of nine of the ten candidate factors led to substantial defects in
ribosome biogenesis, with the exception of *ABCF1* (Fig. 2E). In every case, the affected ribosomal
subunit corresponded to the Perturb-seq clustering across two independent sgRNAs. While this
study was in progress, another group used cryo-EM to identify C1orf131 as a core structural
component of the pre-A1 small subunit processome, complementing our functional evidence (*28*).
This validation suggests that many poorly characterized genes can be assigned functional roles
through Perturb-seq, although a subset of these relationships might be explained by indirect or off-
target effects (fig. S5B,C).

In total, these results show that transcriptional phenotypes revealed by Perturb-seq have
utility beyond studying gene regulation or transcriptional programs, and can serve as valuable tools
for resolving and interrogating many central processes in cell biology.

Delineating functional modules of the Integrator complex

In general, perturbations to members of known protein complexes produced similar
transcriptional phenotypes in our dataset. Therefore, we were surprised by the wide spectrum of
transcriptional responses to knockdown of subunits of Integrator, a metazoan-specific essential
nuclear complex with roles in small nuclear RNA (snRNA) biogenesis and transcription
termination at paused RNA polymerase II (Fig. 3A) (*29*). Each of the fourteen core subunits of
Integrator was targeted in our experiment, allowing us to systematically compare their
transcriptional phenotypes in K562 and RPE1 cells (Fig. 3B; fig. S6A). *INTS1*, *INTS2*, *INTS5*,
INTS7, and *INTS8* formed a tight cluster which weakly correlated with *INTS6* and *INTS12*.
Separately, *INTS3*, *INTS4*, *INTS9*, and *INTS11* clustered together alongside splicing regulators
involved in snRNP assembly and the tri-snRNP. Finally, *INTS10*, *INTS13*, and *INTS14* formed
another discrete cluster together with *C7orf26*, an uncharacterized gene.

These distinct functional modules mirror the architecture of the Integrator complex
observed in recent structures (*30*, *31*). The INTS1-2-5-7-8 functional module contained the
subunits identified as the structural shoulder and backbone of Integrator. The INTS3-4-9-11
functional module contained the subunits identified as the structural cleavage module (as well as
INTS3 which was not resolved). While INTS10, INTS13, and INTS14 were not resolved in the
recent cryo-EM Integrator structures, these subunits have been identified as a stable biochemical
subcomplex (*32*, *33*).

Integrator is an essential and well-studied complex, so we were intrigued by the robust
clustering of the uncharacterized gene *C7orf26* with Integrator subunits 10, 13, and 14. To explore
this, we tested whether loss of C7orf26 impacted the abundance of Integrator subunits. CRISPRi-
based depletion of C7orf26 destabilized INTS10 in K562 cells, confirming either a regulatory or
protein-level relationship (Fig. 3C). Next, we checked for a biochemical interaction between these
proteins. Pulldown of His-INTS10 from cell lysates recovered endogenous C7orf26 alongside
INTS13 and INTS14 (Fig. 3D). Additionally, overexpression of C7orf26 with INTS10, INTS13,
and INTS14 enabled the purification of a stable INTS10-13-14-C7orf26 complex by size-

exclusion chromatography (Fig. 3E; fig. S6B and fig. S7). We also detected a physical interaction between the Drosophila C7orf26 orthologue and fly Integrator in S2 cells and observed co-essentiality between C7orf26 and INTS10, INTS13, INTS14 in the Cancer Dependency Map, suggesting that this relationship is conserved across species and cell types (fig. S8). Together, these results suggest that C7orf26 is a core subunit of a novel INTS10-13-14-C7orf26 Integrator module.

We sought to better understand the distinct transcriptional phenotypes induced by knockdown of INTS10-13-14-C7orf26 compared to the shoulder/backbone and cleavage modules. Comparison of the genes differentially expressed between modules did not reveal function in an obvious way (fig. S6C,D), perhaps owing to the late time point assayed in our experiment. We next explored the canonical role of Integrator in snRNA biogenesis. As mature snRNAs are not captured in 3' scRNAs-seq, we monitored changes in global splicing as a proxy for snRNA biogenesis defects. In our Perturb-seq data, we quantified changes in splicing by comparing the ratio of intronic (unspliced) to exonic (spliced) reads for each gene. Validating our approach, depletion of known splicing factors as well as subunits of the cleavage and shoulder/backbone modules led to gross splicing defects (Fig. 3F). By contrast, depletion of subunits of the INTS10-13-14-C7orf26 module did not cause a substantial splicing defect. To directly test the effect of the INTS10-13-14-C7orf26 module on snRNA biogenesis, we used PRO-seq to probe the positioning of active RNA-polymerase. These data confirmed that extended knockdown of the cleavage and backbone/shoulder modules, but not *INTS10*, *INTS13*, or *C7orf26*, caused a dramatic increase in transcriptional readthrough past the 3' cleavage site of snRNAs (Fig. 3G). In addition, the PRO-seq data confirmed that loss of the INTS10-13-14-C7orf26 module causes a transcriptional phenotype distinct from other modules (fig. S6E).

In sum, our results show that INTS10-13-14-C7orf26 represents a functionally and biochemically distinct module of the Integrator complex, and we propose that *C7orf26* be renamed *INTS15* for future studies (Fig. 3H). Although Integrator has been subjected to extensive structural analyses, it has been difficult to resolve the INTS10-13-14 components in relation to the rest of the complex. Inclusion of C7orf26 may facilitate future structural efforts. Broadly, this example highlights the utility of high-dimensional functional phenotypes for the unsupervised classification of protein complex subunits into functional modules.

Data-driven definition of transcriptional programs

While clustering can organize genetic perturbations into pathways or complexes, it does not reveal the functional consequences of perturbations. To globally summarize the genotype-phenotype relationships in our dataset, we: (i) clustered genes into expression programs based on their co-regulation across perturbations; (ii) clustered perturbations with strong phenotypes based on their transcriptional profiles (as described above); and (iii) computed the average activity of each gene expression program within each perturbation cluster (Fig. 4A,B; fig. S9A; table S7,S8; see *Methods*). This map uncovered many known gene expression programs associated with genetic perturbations, including upregulation of proteasomal subunits due to proteosome dysfunction (*34*), activation of NFkB signaling upon loss of ESCRT proteins (*35*), downregulation of growth-related genes in response to essential genetic perturbations, and upregulation of the cholesterol biosynthesis pathway in response to defects in vesicular trafficking (*36*). Beyond these large-scale relationships, we could also score the effects of individual genetic perturbations on different expression programs. For example, our analysis delineated the canonical branches of the cellular stress response into the independently regulated Unfolded Protein Response (UPR) and Integrated Stress Response (ISR) (Fig. 4C) (*9*). The ISR was highly activated by loss of mitochondrial proteins, aminoacyl-tRNA synthetases, and translation initiation factors, whereas the UPR was activated by loss of ER-resident chaperones and translocation machinery. Collectively, this analysis establishes the ability of Perturb-seq to learn regulatory circuits by leveraging the variability of responses across perturbations.

Interestingly, our unbiased clustering uncovered many perturbations that drove the expression of markers of erythroid or myeloid differentiation, consistent with the known multilineage potential of K562 cells (Fig. 4D) (*37*). The scale of our experiment allowed us to comprehensively search for genes whose modulation promotes cellular differentiation, an application of major interest in both developmental and cancer biology. As expected, loss of central regulators of erythropoiesis (*GATA1*, *LDB1*, *LMO2*, and *KDM1A*) caused myeloid differentiation, whereas knockdown of *BCR-ABL* and its downstream adaptor *GAB2* induced erythroid differentiation (*38*). Surprisingly, loss of a number of common essential genes (i.e., essential across cell lines in the Cancer Dependency Map) also caused expression of either myeloid (e.g., Integrator subcomplex) or erythroid (e.g., NuRD complex, DNA replication machinery) markers. Next, we investigated the differentiation effect of selectively essential genes, which could

be promising targets for differentiation therapy, analogous to ongoing efforts for KDM1A (*39,*

40). We observed that loss of *PTPN1*, a tyrosine phosphatase selectively essential in K562 cells,

drove myeloid differentiation. While inhibitors of PTPN1 have been developed for use in diabetes

and certain cancers (*41*), to our knowledge they have not been tested as a differentiation therapy.

Remarkably, in targeted experiments, we found that knockdown of *PTPN1* and *KDM1A* in

combination caused a substantial increase in differentiation and growth defect compared to either

single genetic perturbation, suggesting that these targets act via different cellular mechanisms (Fig.

4E; fig. S9B). These results highlight the utility of rich phenotypes for understanding

differentiation as well as nominating promising therapeutic targets or combinations.

Hypothesis-driven study of composite phenotypes

We next recognized that our scRNA-seq readout could be used to study phenotypes that

integrate data from across the transcriptome and, therefore, would be difficult to study in

traditional forward genetic screens. Examples of these "composite phenotypes" include total

cellular RNA content and the fraction of RNA derived from transposable elements (TE). We found

numerous composite phenotypes under strong genetic control, with highly reproducible effects

across screen replicates and cell types (Fig. 4F). In the specific case of TE regulation, two major

classes of perturbations increased the fraction of TE RNA by affecting broad classes of elements

including Alu, L1, and MIR (Fig. 4G; fig. S9C). First, loss of subunits of the exosome led to a

substantial increase in the fraction of TE RNA, suggesting that transcripts deriving from TEs might

be preferentially degraded. Second, loss of the CPSF cleavage and polyadenylation complex and

parts of the Integrator complex produced a similar phenotype, suggesting that many of the TE

RNAs observed in K562 cells may be derived from failure of normal transcription termination.

Turning to total RNA content (Fig. 4H), we found that loss of many essential regulators of

S-phase and mitosis increased the RNA content of cells. This is consistent with the observation

that cells tend to increase their size, and thus their RNA content, as they progress through the cell

cycle (fig. S9D), so perturbations that arrest cells in later cell cycle stages yield increased total

RNA content on average. By contrast, loss of essential transcriptional machinery, including

general transcription factors, the Mediator complex, and transcription elongation factors,

decreased total RNA content. In sum, these analyses show that genome-scale Perturb-seq enables

hypothesis-driven exploration of complex cellular features that are challenging to study through
other means.
403
404 ***Exploring genetic drivers and consequences of aneuploidy in single-cells***
405
As Perturb-seq is a single-cell assay, it enables the study of cell-to-cell heterogeneity in
response to genetic perturbations. We reasoned that systematically exploring sources of
heterogeneity could reveal insights into phenotypes that are missed in bulk or averaged
measurements.
To assess the penetrance of perturbation-induced phenotypes, we first applied SVD-based
leverage scores as a metric of single-cell phenotypic magnitude (see *Methods*). In this formulation,
leverage scores quantify how outlying each perturbed cell's transcriptome is relative to non-
targeting control cells without assuming that perturbations drive a single axis of variation.
Supporting this approach, we found that mean leverage scores for each genetic perturbation were
correlated with the number of differentially expressed genes (fig. S10A, Spearman's rho = 0.71),
and reproducible across the day 6 and day 8 K562 experiments (fig. S10B, Spearman's rho = 0.79).
To quantify the degree of heterogeneity in response to genetic perturbations, we then scored
perturbations by the variation in single-cell leverage scores (Fig. 5A; see *Methods*). Comparing
leverage scores across subunits of large essential complexes, we observed evidence for both
biological (e.g., subcomplex function or dosage imbalance) and technical (e.g., selection to escape
toxic perturbations) sources of phenotypic variation in response to genetic perturbations (fig.
S10C-F).
Intriguingly, many genes implicated in chromosome segregation were among the top
drivers of heterogeneity, including *TTK*, *SPC25*, and *DSN1* (Fig. 5B) (*42*). We hypothesized that
the extreme transcriptional variability caused by these genetic perturbations might result from
acute changes in the copy number of individual chromosomes due to mitotic mis-segregation. To
explore this, we used inferCNV (*43*) to estimate single-cell DNA copy number along the genome
by quantifying the change in moving average gene expression compared to control cells.
Consistent with our hypothesis, knockdown of *TTK*, a core component of the spindle assembly
checkpoint (*44*), led to dramatic changes in estimated DNA copy number in both intrinsically
aneuploid K562 and near euploid RPE1 cells (Fig. 5C; fig. S11A). Specifically, in RPE1 cells, we
found that 61/80 (76%) of *TTK* knockdown cells had evidence of karyotypic changes compared to

274/13140 (2%) of unperturbed cells. Notably, *TTK* knockdown cells bore highly variable

karyotypes due to the stochastic gain or loss of chromosomes, accounting for the phenotypic

heterogeneity observed in these cells (Fig. 5C).

An important advantage of the rich data provided by Perturb-seq is the ability to dissect

not only perturbation-phenotype associations but also relationships between cellular phenotypes.

We were curious how chromosomal instability (CIN) would affect cell cycle progression in

euploid, p53-positive RPE1 cells versus constitutively aneuploid, p53-deficient K562 cells.

Expanding our analysis to all cells in our experiment independent of genetic perturbation, we found

that RPE1 cells with abnormal karyotypes tended to arrest in G1 or G0 of the cell cycle (G1 or G0

fraction 0.68 for abnormal karyotype vs. 0.44 for stable karyotype), while K562 cells with altered

karyotypes had less significant shifts in cell cycle occupancy (Fig. 5D,E). Within the population

of RPE1 cells bearing a chromosomal loss, the likelihood of cell cycle arrest directly depended on

the magnitude of karyotypic abnormality (fig. S11B). Additionally, we observed that cells with

the most severe karyotypic changes—those bearing both chromosomal gains and losses—had

marked upregulation of the ISR (Fig. 5F and fig. S11C). These results are consistent with models

in which cell cycle checkpoints are activated by the secondary consequences of aneuploidy (e.g.,

DNA damage or proteostatic stress) rather than changes in chromosome number *per se* (*45*, *46*).

Finally, we looked across all perturbations to systematically identify genetic drivers of

CIN. We assigned a score to each perturbation based on the average magnitude of induced

karyotypic abnormalities. Validating our approach, we found that CIN scores were strongly

correlated across K562 and RPE1 cell lines (*r*=0.69) and identified many known regulators of

chromosomal segregation, including components of the spindle assembly checkpoint, centromere,

and NDC80 complex (Fig. 5G). Remarkably, we uncovered CIN regulators with diverse cellular

roles, from cytoskeletal components to DNA repair machinery (Fig. 5H; table S4-S6). While many

of these genes have previously been associated with chromosomal instability through targeted

studies, the scale and single-cell resolution of Perturb-seq allowed us to identify numerous genetic

drivers of CIN in a single experiment. This analysis also shows the potential of single-cell CRISPR

screens to dissect phenotypes that were not predefined endpoints of the experiment.

Discovery of stress-specific regulation of the mitochondrial genome

Mitochondria arose from the engulfment and endosymbiotic evolution of an ancestral alphaproteobacterium by the precursor to eukaryotic cells (*47*). While the vast majority (~99%) of mitochondrially-localized proteins are encoded in the nuclear genome, mitochondria contain a small (~16.6 kilobase) remnant of their ancestral genome encoding 2 rRNAs, 22 tRNAs, and 13 protein-coding genes in humans. An open question is how the nuclear and mitochondrial genomes coordinate their expression to cope with mitochondrial stress (*48*). The scale of our experiment provided a unique opportunity to investigate this question.

We began by comparing the nuclear transcriptional responses to CRISPRi-based depletion of nuclear-encoded mitochondrial genes (i.e., mitochondrial perturbations). We found that mitochondrial perturbations elicited relatively homogeneous nuclear transcriptional responses, illustrated by well-correlated transcriptional phenotypes across mitochondrial perturbations (Fig. 6A and fig. S12A). While there was some variation in the magnitude of transcriptional responses (e.g., proteostatic injury drove an especially strong ISR activation), nuclear transcriptional responses generally failed to discriminate genetic perturbations by function. Although this result was broadly consistent with recent literature that has highlighted the role of the ISR as response to mitochondrial stress (*49–53*), the lack of functional specificity of the transcriptional response was puzzling in light of: (i) the multifaceted roles of mitochondria in diverse processes such as respiration, intermediary metabolism, iron-sulfur cluster biogenesis, and apoptosis and (ii) the high-resolution separation of cytosolic perturbations by transcriptional response in our data described above.

In contrast to the nuclear transcriptional response, we observed that the expression of mitochondrially encoded genes was highly variable between different mitochondrial perturbations (Fig. 6B; fig. S12B,C,D). When we clustered mitochondrial perturbations based solely on expression levels of the 13 mitochondrially encoded genes, a remarkably intricate and coherent pattern emerged: the clustering separated perturbations to Complex I, Complex IV, Complex III, Complex V (ATP synthase), the mitochondrial large ribosomal subunit, the mitochondrial small ribosomal subunit, chaperones/import machinery, and RNA processing factors (Fig. 6C; fig. S12E). To quantitatively support this observation, we trained a random forest classifier to distinguish cells with perturbations to different mitochondrial complexes and found that the mitochondrial transcriptome was far more predictive than the nuclear transcriptome

(mitochondrial accuracy 0.64; nuclear accuracy: 0.25) (fig. S12F). We then visualized the gene

expression signatures of a subset of representative perturbations (Fig. 6D). The coregulation of

mitochondrial genes tended to reflect function, with the exception of the bicistronic mRNAs

ND4L/ND4 and *ATP8/ATP6* (*54*). However, we did not identify a simple logic to explain the

connection between genetic perturbations to their observed transcriptional consequences. While

previous studies have described distinct regulation of the mitochondrial genome in response to

specific perturbations [notably, related to loss of complex III and complex IV assembly factors

(*55*, *56*)], our data generalize this phenomenon to a comprehensive set of stressors.

Next, we wanted to shed light on the mechanistic basis for this unappreciated complexity

of mitochondrial genome responses. Given its singular origin, the mitochondrial genome is

expressed by unique processes (Fig. 7A) (*57*). Mitochondrially encoded genes are transcribed as

part of three polycistronic transcripts punctuated by tRNAs. These transcripts are then processed

into rRNAs and mRNAs by tRNA excision, and individual mRNAs can be polyadenylated,

expressed, or degraded. This complex system limits the potential for distinct transcriptional control

but presents multiple opportunities for post-transcriptional regulation. To identify modes of

perturbation-elicited differential expression, we examined the distribution of Perturb-seq reads

along the mitochondrial genome (Fig. 7B). As our scRNA-seq used poly-A selection, most reads

aligned to the 3' ends of mRNAs. To validate the utility of this position-based analysis, we

confirmed that knockdown of known regulators of mitochondrial transcription (*TEFM*) and RNA

degradation (*PNPT1*) led to major shifts in the position of reads along the mitochondrial genome.

By contrast, many of the perturbation-specific responses discovered in the present study appeared

to cause shifts in the relative abundance of mRNAs rather than gross shifts in positional

alignments. To determine whether the observed mitochondrial genome responses reflected

regulation of the total level of mitochondrial mRNAs or specific regulation of mRNA

polyadenylation, we performed a bulk RNA-sequencing experiment with no poly-A selection. We

observed perturbation-specific changes in the level of total RNA similar to those measured by

scRNA-seq (cophenetic correlation r=0.79; Fig. 7C). Given the complexity of the observed

responses, we propose that there are likely to be multiple mechanisms that impact the levels of the

various mitochondrially encoded transcripts in response to different stressors.

Finally, we asked whether we could use the detailed clustering produced by the

mitochondrial genome to predict gene function. Knockdown of an unannotated gene, *TMEM242*,

produced a transcriptional signature resembling loss of ATP synthase in both K562 and RPE1 cells

(Fig. 7D; fig. S12G). Supporting this relationship, the top five co-essential genes with *TMEM242*

were components of ATP synthase in the Cancer Dependency Map. Using a Seahorse assay, we

further confirmed that basal respiration was decreased in *TMEM242* knockdown cells (Fig. 7E).

While this work was in progress, another group used a biochemical approach to show that

TMEM242 physically interacts with ATP synthase subunits and regulates ATP synthase complex

assembly in cells (*58*). Together, these experiments discover a novel factor required for ATP

synthase activity and point to the precision of mitochondrial genome regulation.

Discussion

Single-cell CRISPR screens represent an emerging tool to generate rich genotype-

phenotype maps. However, to date, their use has been limited to the study of preselected genes

focused on discrete, predefined biological questions. Here, we perform genome-scale single-cell

CRISPR screens using Perturb-seq and demonstrate how these screens enable data-driven

dissection of a breadth of complex biological phenomena. Reflecting on this study, we highlight

key biological insights and derive principles to guide future discoveries from rich genotype-

phenotype maps.

A primary aim of large-scale functional screens is to organize genes into pathways or

complexes. To this end, we used Perturb-seq to perform high-resolution clustering of genetic

perturbations. From a single assay, we recapitulated thousands of known relationships while also

assigning new, experimentally validated roles to genes involved in ribosome biogenesis or

translation (*CCDC86*, *CINP*, *SPATA5L1*, *ZNF236*, *C1orf131*, *SPOUT1*, *TMA16*, *NOPCHAP1*,

NEPRO), transcription (*C7orf26*), and respiration (*TMEM242*). However, other large-scale

experimental techniques, such as protein-protein interaction mapping, genetic interaction

mapping, and co-essentiality analysis, similarly group genes or proteins by function. How then are

single-cell CRISPR screens distinct?

We argue that these screens are particularly powerful because of the intrinsic

interpretability of comprehensive genotype-phenotype maps, enabling in-depth dissection of the

functional consequences of genetic perturbations that impinge on many distinct aspects of cell

biology. Of particular note is the ability to use the information-rich readouts to study complex,

composite phenotypes, which are difficult to measure by other modalities. These composite

phenotypes can be created in a data-driven (e.g., deriving transcriptional programs) or hypothesis-

driven manner (e.g., measuring intron/exon ratios to study splicing), resulting in an enormous
breadth of measured phenotypes. In the case of scRNA-seq, we show that it measures not only
features such as differential gene expression and the activity of critical transcriptional programs,
but also RNA splicing and processing, expression of transposable elements, differentiation,
transcriptional heterogeneity, cell cycle progression, and chromosomal instability. Once a
phenotype is defined, the genotype-phenotype map can be used to explore its genetic
underpinnings, in a manner analogous to a forward genetic screen, as well as its relationship to
other cellular phenotypes.

An illustrative example of this process is our study of chromosomal instability. Based on
an initial observation of heterogeneous responses to specific perturbations, we suspected that some
cells carried genetically-induced chromosomal gains or losses. In a hypothesis-driven manner, we
then used our rich phenotypic data to discover a large collection of perturbations—which were
only loosely connected by clustering on average transcriptional phenotypes—that promote
chromosomal instability. Importantly, the single-cell nature of our Perturb-seq data also allowed
us to explore the relationship between karyotypic changes and other phenotypes, including cell
cycle progression and stress induction. While aneuploidy is an important hallmark of most cancers,
it has not been easy to study with traditional genetic screens as it requires both a single-cell and
multimodal readout. In future work, this platform could be used to investigate interactions between
genetic perturbations and specific karyotypes, karyotype-dependent stress responses, or the
temporal evolution of karyotypes (*59*).

Genetic perturbations can push cells into extreme states that are not observed in
unperturbed cells. Because composite phenotypes can be generated and explored without being
preregistered at the time of data collection, rich genotype-phenotype maps provide a powerful
resource for the discovery of new cellular behaviors. Using this ability, we discovered a remarkable
degree of stress-specific changes in the expression of mitochondrially encoded transcripts. It was
only possible to appreciate the functional specificity of this regulation by pairing a defined set of
mitochondrial perturbations with a high-dimensional readout. This discovery suggests a
framework to explain how cells cope with diverse insults to mitochondria: a general nuclear
response is layered over perturbation-specific changes in the expression level of mitochondrially
encoded genes (Fig. 7F). Building on this observation, we can ask new questions about the
mitochondrial stress response. The transcriptional changes we observed may reflect adaptive
responses or, alternatively, complex patterns of dysfunction owing to disruption of the intricate

system of mitochondrial gene expression. Understanding how and in what contexts this regulation

is adaptive may have important implications for diseases associated with mitochondrial stress. An

intriguing additional question is whether individual mitochondria are able to regulate their

expression autonomously. Combined with the nuanced responses observed here, this would

support and substantially extend the "co-location for redox regulation" (CoRR) hypothesis which

holds that the endosymbiotically derived mitochondrial genome has been retained through

evolution to enable localized regulation of mitochondrial gene expression (*60*).

A final theme emerging from our work is the flexibility of single-cell CRISPR screens

compared to other functional genomic approaches. Because these screens extract rich information

from each cell in a pooled format, they require only a fraction of the number of cells used by other

approaches and thus are well suited to the study of iPSC-derived cells and *in vivo* samples. As

technologies for single-cell, multimodal phenotyping advance, single-cell screens will continue to

become more powerful. At present, the major limitation of single-cell CRISPR screens is cost.

Careful experimental designs, such as multiplexed libraries or compressed sensing (*61*), together

with advances in single-cell phenotyping (*62*, *63*) and DNA sequencing promise to greatly increase

the scale of these experiments.

In sum, our study presents a blueprint for the construction and analysis of rich genotype-

phenotype maps to serve as a driving force for the systematic exploration of genetic and cellular

function.

Materials and Methods

A complete description of our Material and Methods is found in the Supplementary Material online. This includes methods experimental methods related to Perturb-seq screens and functional experiments, as well as computational methods detailing all data analysis.

References and Notes

1. J. G. Camp, R. Platt, B. Treutlein, Mapping human cell phenotypes to genotypes with single-cell genomics. *Sci New York N Y*. 365, 1401–1405 (2019).

2. J. G. Doench, Am I ready for CRISPR? A user's guide to genetic screens. *Nat Rev Genet*. 19, 67–80 (2018).

3. N. J. Kramer, M. S. Haney, D. W. Morgens, A. Jovičić, J. Couthouis, A. Li, J. Ousey, R. Ma, G. Bieri, C. K. Tsui, Y. Shi, N. T. Hertz, M. Tessier-Lavigne, J. K. Ichida, M. C. Bassik, A. D. Gitler, CRISPR–Cas9 screens in human cells and primary neurons identify modifiers of C9ORF72 dipeptide-repeat-protein toxicity. *Nat Genet*. 50, 603–612 (2018).

4. A. Tsherniak, F. Vazquez, P. G. Montgomery, B. A. Weir, G. Kryukov, G. S. Cowley, S. Gill, W. F. Harrington, S. Pantel, J. M. Krill-Burger, R. M. Meyers, L. Ali, A. Goodale, Y. Lee, G. Jiang, J. Hsiao, W. F. J. Gerath, S. Howell, E. Merkel, M. Ghandi, L. A. Garraway, D. E. Root, T. R. Golub, J. S. Boehm, W. C. Hahn, Defining a Cancer Dependency Map. *Cell*. 170, 564-576.e16 (2017).

5. T. Wang, K. Birsoy, N. W. Hughes, K. M. Krupczak, Y. Post, J. J. Wei, E. S. Lander, D. M. Sabatini, Identification and characterization of essential genes in the human genome. *Science*. 350, 1096–1101 (2015).

6. E. Wang, H. Zhou, B. Nadorp, G. Cayanan, X. Chen, A. H. Yeaton, S. Nomikou, M. T. Witkowski, S. Narang, A. Kloetgen, P. Thandapani, N. Ravn-Boess, A. Tsirigos, I. Aifantis, Surface antigen-guided CRISPR screens identify regulators of myeloid leukemia differentiation. *Cell Stem Cell*. 28, 718-731.e6 (2021).

7. T. Stuart, R. Satija, Integrative single-cell analysis. *Nat Rev Genet*. 20, 257–272 (2019).

8. L. Przybyla, L. A. Gilbert, A new era in functional genomics screens. *Nat Rev Genet*, 1–15

(2021).

9. B. Adamson, T. M. Norman, M. Jost, M. Y. Cho, J. K. Nuñez, Y. Chen, J. E. Villalta, L. A.

Gilbert, M. A. Horlbeck, M. Y. Hein, R. A. Pak, A. N. Gray, C. A. Gross, A. Dixit, O. Parnas, A.

Regev, J. S. Weissman, A Multiplexed Single-Cell CRISPR Screening Platform Enables

Systematic Dissection of the Unfolded Protein Response. *Cell*. 167, 1867-1882.e21 (2016).

10. A. Dixit, O. Parnas, B. Li, J. Chen, C. P. Fulco, L. Jerby-Arnon, N. D. Marjanovic, D.

Dionne, T. Burks, R. Raychowdhury, B. Adamson, T. M. Norman, E. S. Lander, J. S. Weissman,

N. Friedman, A. Regev, Perturb-Seq: Dissecting Molecular Circuits with Scalable Single-Cell

RNA Profiling of Pooled Genetic Screens. *Cell*. 167, 1853-1866.e17 (2016).

11. D. A. Jaitin, A. Weiner, I. Yofe, D. Lara-Astiaso, H. Keren-Shaul, E. David, T. M. Salame,

A. Tanay, A. van Oudenaarden, I. Amit, Dissecting Immune Circuits by Linking CRISPR-

Pooled Screens with Single-Cell RNA-Seq. *Cell*. 167, 1883-1896.e15 (2016).

12. P. Datlinger, A. F. Rendeiro, C. Schmidl, T. Krausgruber, P. Traxler, J. Klughammer, L. C.

Schuster, A. Kuchler, D. Alpar, C. Bock, Pooled CRISPR screening with single-cell

transcriptome readout. *Nat Methods*. 14, 297–301 (2017).

13. J. M. Replogle, T. M. Norman, A. Xu, J. A. Hussmann, J. Chen, J. Z. Cogan, E. J. Meer, J.

M. Terry, D. P. Riordan, N. Srinivas, I. T. Fiddes, J. G. Arthur, L. J. Alvarado, K. A. Pfeiffer, T.

S. Mikkelsen, J. S. Weissman, B. Adamson, Combinatorial single-cell CRISPR screens by direct

guide RNA capture and targeted sequencing. *Nat Biotechnol*. 38, 954–961 (2020).

14. A. J. Rubin, K. R. Parker, A. T. Satpathy, Y. Qi, B. Wu, A. J. Ong, M. R. Mumbach, A. L. Ji,

D. S. Kim, S. W. Cho, B. J. Zarnegar, W. J. Greenleaf, H. Y. Chang, P. A. Khavari, Coupled

Single-Cell CRISPR Screening and Epigenomic Profiling Reveals Causal Gene Regulatory

Networks. *Cell*. 176, 361-376.e17 (2019).

15. D. Feldman, A. Singh, J. L. Schmid-Burgk, R. J. Carlson, A. Mezger, A. J. Garrity, F. Zhang,

P. C. Blainey, Optical Pooled Screens in Human Cells. *Cell*. 179, 787-799.e17 (2019).

16. E. P. Mimitou, A. Cheng, A. Montalbano, S. Hao, M. Stoeckius, M. Legut, T. Roush, A.

Herrera, E. Papalexi, Z. Ouyang, R. Satija, N. E. Sanjana, S. B. Koralov, P. Smibert, Multiplexed

detection of proteins, transcriptomes, clonotypes and CRISPR perturbations in single cells. *Nat*

Methods. 16, 409–412 (2019).

17. C. J. Frangieh, J. C. Melms, P. I. Thakore, K. R. Geiger-Schuller, P. Ho, A. M. Luoma, B.

Cleary, L. Jerby-Arnon, S. Malu, M. S. Cuoco, M. Zhao, C. R. Ager, M. Rogava, L. Hovey, A.

Rotem, C. Bernatchez, K. W. Wucherpfennig, B. E. Johnson, O. Rozenblatt-Rosen, D.

Schadendorf, A. Regev, B. Izar, Multimodal pooled Perturb-CITE-seq screens in patient models

define mechanisms of cancer immune evasion. *Nat Genet*. 53, 332–341 (2021).

18. E. Papalexi, E. P. Mimitou, A. W. Butler, S. Foster, B. Bracken, W. M. Mauck, H.-H.

Wessels, Y. Hao, B. Z. Yeung, P. Smibert, R. Satija, Characterizing the molecular regulation of

inhibitory immune checkpoints with multimodal single-cell screens. *Nat Genet*. 53, 322–331

(2021).

19. A. Tanay, A. Regev, Scaling single-cell genomics from phenomenology to mechanism.

Nature. 541, 331–338 (2017).

20. T. M. Norman, M. A. Horlbeck, J. M. Replogle, A. Y. Ge, A. Xu, M. Jost, L. A. Gilbert, J. S.

Weissman, Exploring genetic interaction manifolds constructed from rich single-cell phenotypes.

Science. 365, 786–793 (2019).

21. A. H. Smits, F. Ziebell, G. Joberty, N. Zinn, W. F. Mueller, S. Clauder-Münster, D.

Eberhard, M. F. Savitski, P. Grandi, P. Jakob, A.-M. Michon, H. Sun, K. Tessmer, T.

Bürckstümmer, M. Bantscheff, L. M. Steinmetz, G. Drewes, W. Huber, Biological plasticity

rescues target activity in CRISPR knock outs. *Nat Methods*. 16, 1087–1093 (2019).

22. E. Haapaniemi, S. Botla, J. Persson, B. Schmierer, J. Taipale, CRISPR–Cas9 genome editing

induces a p53-mediated DNA damage response. *Nat Med*. 24, 927–930 (2018).

23. L. A. Gilbert, M. A. Horlbeck, B. Adamson, J. E. Villalta, Y. Chen, E. H. Whitehead, C.

Guimaraes, B. Panning, H. L. Ploegh, M. C. Bassik, L. S. Qi, M. Kampmann, J. S. Weissman,

Genome-Scale CRISPR-Mediated Control of Gene Repression and Activation. *Cell*. 159, 647–

661 (2014).

24. N. Alerasool, D. Segal, H. Lee, M. Taipale, An efficient KRAB domain for CRISPRi

applications in human cells. *Nat Methods*. 17, 1093–1096 (2020).

25. J. Rosenbluh, H. Xu, W. Harrington, S. Gill, X. Wang, F. Vazquez, D. E. Root, A. Tsherniak,

W. C. Hahn, Complementary information derived from CRISPR Cas9 mediated gene deletion

and suppression. *Nat Commun*. 8, 15403 (2017).

26. M. Giurgiu, J. Reinhard, B. Brauner, I. Dunger-Kaltenbach, G. Fobo, G. Frishman, C.

Montrone, A. Ruepp, CORUM: the comprehensive resource of mammalian protein complexes—

2019. *Nucleic Acids Res*. 47, D559–D563 (2019).

27. D. Szklarczyk, A. L. Gable, D. Lyon, A. Junge, S. Wyder, J. Huerta-Cepas, M. Simonovic,

N. T. Doncheva, J. H. Morris, P. Bork, L. J. Jensen, C. von Mering, STRING v11: protein–

protein association networks with increased coverage, supporting functional discovery in

genome-wide experimental datasets. *Nucleic Acids Res*. 47, D607–D613 (2019).

28. S. Singh, A. V. Broeck, L. Miller, M. Chaker-Margot, S. Klinge, Nucleolar maturation of the

human small subunit processome. *Science*. 373, eabj5338 (2021).

29. N. Kirstein, H. G. D. Santos, E. Blumenthal, R. Shiekhattar, The Integrator complex at the

crossroad of coding and noncoding RNA. *Curr Opin Cell Biol*. 70, 37–43 (2021).

30. I. Fianu, Y. Chen, C. Dienemann, O. Dybkov, A. Linden, H. Urlaub, P. Cramer, Structural

basis of Integrator-mediated transcription regulation. *Science*. 374, 883–887 (2021).

31. H. Zheng, Y. Qi, S. Hu, X. Cao, C. Xu, Z. Yin, X. Chen, Y. Li, W. Liu, J. Li, J. Wang, G.

Wei, K. Liang, F. X. Chen, Y. Xu, Identification of Integrator-PP2A complex (INTAC), an RNA

polymerase II phosphatase. *Sci New York N Y*. 370 (2020), doi:10.1126/science.abb5872.

32. M. M. Pfleiderer, W. P. Galej, Structure of the catalytic core of the Integrator complex. *Mol*

Cell. 81, 1246-1259.e8 (2021).

33. K. Sabath, M. L. Stäubli, S. Marti, A. Leitner, M. Moes, S. Jonas, INTS10–INTS13–INTS14

form a functional module of Integrator that binds nucleic acids and the cleavage module. *Nat*

Commun. 11, 3422 (2020).

34. S. K. Radhakrishnan, C. S. Lee, P. Young, A. Beskow, J. Y. Chan, R. J. Deshaies,

Transcription Factor Nrf1 Mediates the Proteasome Recovery Pathway after Proteasome

Inhibition in Mammalian Cells. *Mol Cell*. 38, 17–28 (2010).

35. A. Mamińska, A. Bartosik, M. Banach-Orłowska, I. Pilecka, K. Jastrzębski, D. Zdżalik-

Bielecka, I. Castanon, M. Poulain, C. Neyen, L. Wolińska-Nizioł, A. Toruń, E. Szymańska, A.

Kowalczyk, K. Piwocka, A. Simonsen, H. Stenmark, M. Fürthauer, M. González-Gaitán, M.

Miaczynska, ESCRT proteins restrict constitutive NF-κB signaling by trafficking cytokine

receptors. *Sci Signal*. 9, ra8–ra8 (2016).

36. J. Luo, H. Yang, B.-L. Song, Mechanisms and regulation of cholesterol homeostasis. *Nat Rev*

Mol Cell Bio. 21, 225–245 (2020).

37. J. F. Leary, B. M. Ohlsson-Wilhelm, R. Giuliano, S. Labella, B. Farley, P. T. Rowley,

Multipotent human hematopoietic cell line K562: Lineage-specific constitutive and inducible

antigens. *Leukemia Res*. 11, 807–815 (1987).

38. S. H. Orkin, L. I. Zon, Hematopoiesis: An Evolving Paradigm for Stem Cell Biology. *Cell*.

132, 631–644 (2008).

39. L. Yu, G. Myers, C.-J. Ku, E. Schneider, Y. Wang, S. A. Singh, N. Jearawiriyapaisarn, A.

White, T. Moriguchi, R. Khoriaty, M. Yamamoto, M. G. Rosenfeld, J. Pedron, J. H. Bushweller,

K.-C. Lim, J. D. Engel, An erythroid-to-myeloid cell fate conversion is elicited by LSD1

inactivation. *Blood*. 138, 1691–1704 (2021).

40. T. Maes, C. Mascaró, I. Tirapu, A. Estiarte, F. Ciceri, S. Lunardi, N. Guibourt, A. Perdones,

M. M. P. Lufino, T. C. P. Somervaille, D. H. Wiseman, C. Duy, A. Melnick, C. Willekens, A.

Ortega, M. Martinell, N. Valls, G. Kurz, M. Fyfe, J. C. Castro-Palomino, C. Buesa, ORY-1001, a

Potent and Selective Covalent KDM1A Inhibitor, for the Treatment of Acute Leukemia. *Cancer*

Cell. 33, 495-511.e12 (2018).

41. B. Sharma, L. Xie, F. Yang, W. Wang, Q. Zhou, M. Xiang, S. Zhou, W. Lv, Y. Jia, L.
Pokhrel, J. Shen, Q. Xiao, L. Gao, W. Deng, Recent advance on PTP1B inhibitors and their
biomedical applications. *Eur J Med Chem*. 199, 112376 (2020).

42. A. Musacchio, E. D. Salmon, The spindle-assembly checkpoint in space and time. *Nat Rev
Mol Cell Bio*. 8, 379–393 (2007).

43. A. P. Patel, I. Tirosh, J. J. Trombetta, A. K. Shalek, S. M. Gillespie, H. Wakimoto, D. P.
Cahill, B. V. Nahed, W. T. Curry, R. L. Martuza, D. N. Louis, O. Rozenblatt-Rosen, M. L. Suvà,
A. Regev, B. E. Bernstein, Single-cell RNA-seq highlights intratumoral heterogeneity in primary
glioblastoma. *Science*. 344, 1396–1401 (2014).

44. N. Jelluma, A. B. Brenkman, N. J. F. van den Broek, C. W. A. Cruijsen, M. H. J. van Osch,
S. M. A. Lens, R. H. Medema, G. J. P. L. Kops, Mps1 Phosphorylates Borealin to Control
Aurora B Activity and Chromosome Alignment. *Cell*. 132, 233–246 (2008).

45. S. Santaguida, A. Richardson, D. R. Iyer, O. M'Saad, L. Zasadil, K. A. Knouse, Y. L. Wong,
N. Rhind, A. Desai, A. Amon, Chromosome Mis-segregation Generates Cell-Cycle-Arrested
Cells with Complex Karyotypes that Are Eliminated by the Immune System. *Dev Cell*. 41, 638-
651.e5 (2017).

46. S. Santaguida, A. Amon, Short- and long-term effects of chromosome mis-segregation and
aneuploidy. *Nat Rev Mol Cell Bio*. 16, 473–485 (2015).

47. J. R. Friedman, J. Nunnari, Mitochondrial form and function. *Nature*. 505, 335–343 (2014).

48. P. M. Quirós, A. Mottis, J. Auwerx, Mitonuclear communication in homeostasis and stress.
Nat Rev Mol Cell Biology. 17, 213–26 (2016).

49. E. Mick, D. V. Titov, O. S. Skinner, R. Sharma, A. A. Jourdain, V. K. Mootha, Distinct
mitochondrial defects trigger the integrated stress response depending on the metabolic state of
the cell. *Elife*. 9, e49178 (2020).

50. P. M. Quirós, M. A. Prado, N. Zamboni, D. D'Amico, R. W. Williams, D. Finley, S. P. Gygi,
J. Auwerx, Multi-omics analysis identifies ATF4 as a key regulator of the mitochondrial stress
response in mammals. *J Cell Biol*. 216, 2027–2045 (2017).

51. C. Münch, J. W. Harper, Mitochondrial unfolded protein response controls matrix pre-RNA
processing and translation. *Nature*. 534, 710–713 (2016).

52. E. Fessler, E.-M. Eckl, S. Schmitt, I. A. Mancilla, M. F. Meyer-Bender, M. Hanf, J.
Philippou-Massier, S. Krebs, H. Zischka, L. T. Jae, A pathway coordinated by DELE1 relays
mitochondrial stress to the cytosol. *Nature*. 579, 433–437 (2020).

53. X. Guo, G. Aviles, Y. Liu, R. Tian, B. A. Unger, Y.-H. T. Lin, A. P. Wiita, K. Xu, M. A.
Correia, M. Kampmann, Mitochondrial stress is relayed to the cytosol by an OMA1-DELE1-HRI
pathway. *Nature*. 579, 427–432 (2020).

54. T. R. Mercer, S. Neph, M. E. Dinger, J. Crawford, M. A. Smith, A.-M. J. Shearwood, E.
Haugen, C. P. Bracken, O. Rackham, J. A. Stamatoyannopoulos, A. Filipovska, J. S. Mattick,
The Human Mitochondrial Transcriptome. *Cell*. 146, 645–658 (2011).

55. R. Richter-Dennerlein, S. Oeljeklaus, I. Lorenzi, C. Ronsör, B. Bareth, A. B. Schendzielorz,
C. Wang, B. Warscheid, P. Rehling, S. Dennerlein, Mitochondrial Protein Synthesis Adapts to
Influx of Nuclear-Encoded Protein. *Cell*. 167, 471-483.e10 (2016).

56. R. Salvatori, K. Kehrein, A. P. Singh, W. Aftab, B. V. Möller-Hergt, I. Forne, A. Imhof, M.
Ott, Molecular Wiring of a Mitochondrial Translational Feedback Loop. *Mol Cell*. 77, 887-
900.e5 (2020).

57. E. Kummer, N. Ban, Mechanisms and regulation of protein synthesis in mitochondria. *Nat
Rev Mol Cell Bio*. 22, 307–325 (2021).

58. J. Carroll, J. He, S. Ding, I. M. Fearnley, J. E. Walker, TMEM70 and TMEM242 help to
assemble the rotor ring of human ATP synthase and interact with assembly factors for complex I.
P Natl Acad Sci Usa. 118, e2100558118 (2021).

59. U. Ben-David, A. Amon, Context is everything: aneuploidy in cancer. *Nat Rev Genet*. 21,
44–62 (2020).

60. J. F. Allen, The CoRR hypothesis for genes in organelles. *J Theor Biol*. 434, 50–57 (2017).

61. B. Cleary, L. Cong, A. Cheung, E. S. Lander, A. Regev, Efficient Generation of
Transcriptomic Profiles by Random Composite Measurements. *Cell*. 171, 1424-1436.e18 (2017).

62. P. Datlinger, A. F. Rendeiro, T. Boenke, M. Senekowitsch, T. Krausgruber, D. Barreca, C.
Bock, Ultra-high-throughput single-cell RNA sequencing and perturbation screening with
combinatorial fluidic indexing. *Nat Methods*. 18, 635–642 (2021).

63. B. K. Martin, C. Qiu, E. Nichols, M. Phung, R. Green-Gladden, S. Srivatsan, R. Blecher-
Gonen, B. J. Beliveau, C. Trapnell, J. Cao, J. Shendure, An optimized protocol for single cell
transcriptional profiling by combinatorial indexing. *Arxiv* (2021).

64. M. Jost, Y. Chen, L. A. Gilbert, M. A. Horlbeck, L. Krenning, G. Menchon, A. Rai, M. Y.
Cho, J. J. Stern, A. E. Prota, M. Kampmann, A. Akhmanova, M. O. Steinmetz, M. E.
Tanenbaum, J. S. Weissman, Combined CRISPRi/a-Based Chemical Genetic Screens Reveal
that Rigosertib Is a Microtubule-Destabilizing Agent. *Mol Cell*. 68, 210-223.e6 (2017).

65. S. A. Lambert, A. Jolma, L. F. Campitelli, P. K. Das, Y. Yin, M. Albu, X. Chen, J. Taipale,
T. R. Hughes, M. T. Weirauch, The Human Transcription Factors. *Cell*. 172, 650–665 (2018).

66. M. A. Horlbeck, L. A. Gilbert, J. E. Villalta, B. Adamson, R. A. Pak, Y. Chen, A. P. Fields,
C. Y. Park, J. E. Corn, M. Kampmann, J. S. Weissman, Compact and highly active next-
generation libraries for CRISPR-mediated gene repression and activation. *Elife*. 5, e19760
(2016).

67. N. D. Elrod, T. Henriques, K.-L. Huang, D. C. Tatomer, J. E. Wilusz, E. J. Wagner, K.
Adelman, The Integrator Complex Attenuates Promoter-Proximal Transcription at Protein-
Coding Genes. *Mol Cell*. 76, 738-752.e7 (2019).

68. K.-L. Huang, D. Jee, C. B. Stein, N. D. Elrod, T. Henriques, L. G. Mascibroda, D. Baillat, W.
K. Russell, K. Adelman, E. J. Wagner, Integrator Recruits Protein Phosphatase 2A to Prevent
Pause Release and Facilitate Transcription Termination. *Mol Cell*. 80, 345-358.e9 (2020).

69. A. P. Anderson, X. Luo, W. Russell, Y. W. Yin, Oxidative damage diminishes mitochondrial
DNA polymerase replication fidelity. *Nucleic Acids Res*. 48, 817–829 (2020).

70. K. A. Reimer, C. A. Mimoso, K. Adelman, K. M. Neugebauer, Co-transcriptional splicing
regulates 3′ end cleavage during mammalian erythropoiesis. *Mol Cell*. 81, 998-1012.e7 (2021).

71. Z. Xie, A. Bailey, M. V. Kuleshov, D. J. B. Clarke, J. E. Evangelista, S. L. Jenkins, A.
Lachmann, M. L. Wojciechowicz, E. Kropiwnicki, K. M. Jagodnik, M. Jeon, A. Ma'ayan, Gene
Set Knowledge Discovery with Enrichr. *Curr Protoc*. 1, e90 (2021).

72. J. He, I. A. Babarinde, L. Sun, S. Xu, R. Chen, J. Shi, Y. Wei, Y. Li, G. Ma, Q. Zhuang, A.
P. Hutchins, J. Chen, Identifying transposable element expression dynamics and heterogeneity
during development at the single-cell level with a processing pipeline scTE. *Nat Commun*. 12,
1456 (2021).

73. P. Ma, M. W. Mahoney, B. Yu, A Statistical Perspective on Algorithmic Leveraging. *Arxiv*
(2013).

74. R. Satija, J. A. Farrell, D. Gennert, A. F. Schier, A. Regev, Spatial reconstruction of single-
cell gene expression data. *Nat Biotechnol*. 33, 495–502 (2015).

Acknowledgements: We thank S. Vazquez, L. Gilbert, F. Urnov, C. Cotta-Ramusino, V.
Sankaran, K. Loh, W. Allen, B. Do, P. Hsu, F. Diehl, C. Jan, J. Replogle, D. Phizicky, and all
members of the Weissman and Norman labs for helpful discussions. We thank Jorge Dinis and
the Innovative Genomics Institute for critical administrative and scientific support. We also
thank: the UCSF Center for Advanced Technology, Eric Chow, and Delsy Martinez for
sequencing and access to FACS machines; the Whitehead Institute Flow Cytometry Core and
Kathy Daniels for access to FACS machines; the Whitehead Institute Genome Technology Core
for support with bulk RNA-seq library preparation and rRNA bioanalyzers; and the Harvard
Nascent Transcriptomics Core for support with PRO-seq library preparation.

Author contributions: JMR, RAS, TMN, and JSW were responsible for the conception, design,
and interpretation of the experiments and wrote the manuscript. JMR led the Perturb-seq screens.
JMR and TMN led the Perturb-seq data analysis. RAS led the functional studies. ANP generated
and validated the ZIM3 CRISPRi RPE1 cell line and optimized Perturb-seq protocols across cell
lines. JAH assisted with data interpretation and analysis. AL designed the interactive web
visualization. AG produced preliminary data for Perturb-seq across cell lines. EJW and LM
performed and supervised Drosophila Integrator biochemistry. KA supervised PRO-seq library
preparation. JLB and MJ cloned pJB108 and validated the ZIM3 KRAB domain. JMR and RAS
helped obtain funding for experiments. All authors provided feedback on the manuscript.

Funding: Research reported in this publication was supported by:
Defense Advanced Research Projects Agency (DARPA) grant HR0011-19-2-0007 (JSW)
National Institutes of Health (NIH) Centers of Excellence in Genomic Science (CEGS) (JSW)
Howard Hughes Medical Institute (JSW)
Chan Zuckerberg Initiative (JSW)
NIH grant 1DP2 GM140925-01 (TMN)
NIH grant R00-GM130964 (MJ)
NIH grant R01-GM134539 (EJW, KA)
Fannie and John Hertz Foundation Fellowship (RAS)
NSF Graduate Research Fellowship (RAS)
NIH grant F31-NS115380 (JMR)

Competing interests: JMR consults for Maze Therapeutics and is a consultant for and equity

holder in Waypoint Bio. RAS consults for Maze Therapeutics. KA is a consultant for Syros

Pharmaceuticals, is on the SAB of CAMP4 Therapeutics, and received research funding from

Novartis not related to this work. MJ consults for Maze Therapeutics and Gate Bioscience. TMN

consults for Maze Therapeutics. JSW declares outside interest in 5 AM Venture, Amgen,

Chroma Medicine, KSQ Therapeutics, Maze Therapeutics, Tenaya Therapeutics, Tessera

Therapeutics and Third Rock Ventures. The Regents of the University of California with RAS,

TMN, MJ, and JSW as inventors have filed patent applications related to CRISPRi/a screening

and Perturb-seq.

Data and materials availability: Raw sequencing data will be deposited into SRA. An interactive

data browser including processed, downloadable single-cell and pseudobulk populations will be

made available upon publication. Our previously published analytic framework for Perturb-seq

analysis is available at https://github.com/thomasmaxwellnorman/Perturbseq_GI. Scripts for

guide assignment are available at https://github.com/josephreplogle/guide_calling. Additional

code related to specific analyses will be made available on github upon publication.

Supplementary Materials:

Materials and Methods

Figs. S1 to S12

Tables S1 to S9

Figure 1: Genome-scale Perturb-seq via multiplexed CRISPRi. (A) Schematic experimental strategy. A multiplexed CRISPRi sgRNA library was used to knock down all expressed genes (in K562 cells) or all common essential genes (in RPE1 and K562 cells). Cells were transcriptionally profiled using droplet-based single-cell RNA-sequencing, with genetic perturbations assigned to cells by direct capture and sequencing of sgRNAs. **(B)** On-target knockdown statistics. Cumulative density plot of on-target knockdown, for n=9,464 target genes in K562 cells (red) and n=2,333 target genes in RPE1 cells (blue). **(C)** Comparing growth phenotype versus the number of differentially expressed genes (DEGs) for each multiplexed guide pair in K562 cells. Growth phenotypes are reported as the $\log_2$ guide enrichment per cell doubling (gamma). DEGs were determined using a two-sample Anderson-Darling test compared against non-targeting guides, and a pseudocount of 1 was added to the number of DEGs before $\log_{10}$ transformation. Dots are colored by Energy distance as either permutation significant (purple) or not significant (grey). The growth phenotype and number of DEGs are anticorrelated (Spearman's rho=-0.51).

Figure 2: Data-driven inference of gene function from transcriptional phenotypes. (A) Schematic of analysis. To examine the ability of transcriptional phenotypes to assign gene function, we analyzed 1973 genetic perturbations that elicited strong responses. Perturbations were compared and clustered using the correlation of gene expression across 2319 highly variable genes. **(B)** Expression profile correlations among genes in curated complexes. 327 protein complexes from the CORUM3.0 database have at least two thirds of complex subunits within the dataset. Plot compares the distribution of pairwise expression profile correlations among genes in complexes vs. all possible gene-gene pairs. **(C)** Comparing expression profile correlations to predicted protein-protein interactions from STRING. 243,558 gene-gene relationships within the dataset are scored within STRING. The relationships were sorted into 6 equally spaced bins based on expression profile correlation. Plot shows kernel density estimates of STRING scores within each bin. **(D)** Minimum distortion embedding of dataset. Each dot represents a genetic perturbation, arranged so that perturbations with correlated expression profiles are nearby in the two dimensional embedding. Manual annotations (black labels) of

cluster function are placed near the median location of genes within the cluster. CORUM complexes or STRING clusters (green labels) are annotated when involved genes are nearby within the embedding. **(E)** Quantification of 28s to 18s rRNA ratio. Poorly characterized genes with Perturb-seq predicted roles in ribosome biogenesis were targeted by CRISPRi. The 28s/18s rRNA ratio was measured by Bioanalyzer electrophoresis in biological duplicate with two distinct sgRNAs per gene (green and blue; solid grey lines represent mean). Dotted grey lines represent two standard deviations above and below the mean of non-targeting controls.

Figure 3. (legend on next page)

Figure 3: Perturb-seq discovers a novel gene member and functional submodules of the Integrator complex. (A) Location of known Integrator complex members in the minimum distortion embedding. **(B)** Relationship between Integrator complex members and *C7orf26* in K562 cells (top) and RPE1 cells (bottom). The heatmap displays the Pearson correlation between pseudobulk *z*-normalized gene expression profiles of Integrator complex members. Genetic perturbations are ordered by average linkage hierarchical clustering based on correlation in K562 cells. Functional modules suggested by the clustering are highlighted. **(C)** Co-depletion of Integrator complex members. Individual Integrator complex members were depleted in CRISPRi K562 cells. Lysates were then probed for other module members by western blot. **(D)** Co-immunoprecipitation of endogenous C7orf26 with His-INTS10. HEK293T were transfected with His-INTS10 or INTS10. Cell lysates were affinity purified and select Integrator proteins were probed by western blot. **(E)** Purification of a INTS10-INTS13-INTS14-C7orf26 complex. His-INTS10, INTS13, INTS14, and C7orf26 were overexpressed in Expi293 cells, affinity purified, and separated via SEC. The INTS10-INTS13-INTS14-C7orf26 proteins co-fractionated as visualized by Western blotting. **(F)** Effects of Integrator modules on splicing from Perturb-seq data. Histogram (kernel density estimate) compares gene-level splicing scores. Splicing scores represent the change in the $\log_2$ ratio of total to unspliced reads for each gene relative to non-targeting control guides. Representative genetic perturbations from Integrator modules as well as the spliceosome are shown colored by module. **(G)** Density of PRO-seq reads at the snRNA *RNU1-1* locus mapping actively engaged RNA polymerase II. For each perturbation, densities are shown relative to the maximum read count in the locus. **(H)** Structure of the Integrator complex colored by functional modules revealed by Perturb-seq. The endonuclease (blue) and shoulder/backbone (orange) modules were obtained from the cryo-EM structure (*31*). The model of the newly discovered 10-13-14-C7orf26 module was built by docking the crystal structure of INTS13-INTS14 (*33*) with an AlphaFold multimeric model of INTS10 and C7orf26.

Figure 4: Summarizing genotype-phenotype relationships with Perturb-seq. (A) Schematic of analysis. To produce a high-level summary of genotype-phenotype relationships in K562 cells, 1973 genetic perturbations that elicited strong responses and 2319 highly variable genes were clustered using HDBSCAN after nonlinear embedding. Alternatively, composite phenotypes were derived from global metrics in a hypothesis-driven manner. **(B)** Heatmap of the high-level genotype-phenotype map. The heatmap represents the mean z-scored expression for gene expression and perturbation clusters. For a subset of clusters, clustered are labelled with manual annotations (black labels) of cluster function along with example genes within the cluster (light gray labels). **(C)** Comparison of ISR and UPR scores for genetic perturbations. Scores were recovered from unbiased clustering of genes by genetic dependency, and manually annotated. **(D)** Comparison of erythroid and myeloid differentiation scores for genetic perturbations. Scores were recovered from unbiased clustering of genes by genetic dependency, and manually annotated. Genetic perturbations are colored to reflect cluster identity. **(E)** Expression of CD11b/ITGAM in

K562 cells upon knockdown of *PTPN1* or *KDM1A*. CD11b was labelled by cell surface staining with anti-CD11b antibody and measured by flow cytometry. **(F)** Correlation of composite phenotypes across time points and cell types. Composite phenotypes were defined in a hypothesis-driven manner. Fraction TE (repetitive and transposable element) represents the number of non-intronic reads mapped to TEs over total, averaged over all cells bearing each perturbation (both collapsed on UMIs). Fraction mtRNA represents the mean number of reads mapped to mitochondrial genome protein-coding genes over total. Total RNA represents the mean total RNA content (number of UMIs). **(G)** Comparison of TE expression across time points. The mean fraction TE reads per perturbation is highly correlated across time points in K562 cells (r=0.79). Genetic perturbations are colored to reflect cluster identity. **(H)** Comparison of total RNA content across time points. Total RNA content per perturbation is highly correlated across time points in K562 cells (r=0.74). Genetic perturbations are colored to reflect cluster identity.

Figure 5: Exploring acute consequences and genetic drivers of aneuploidy in single-cells. (A) Schematic of heterogeneity statistic. Single-cell leverage scores quantify how outlying each cell is relative to non-targeting control cells by PCA. For each perturbation, heterogeneity of single-cell phenotypes is quantified as the standard deviation of leverage scores. **(B)** Identifying heterogeneous perturbations. Known regulators of chromosome segregation were among the perturbations with the highest single-cell heterogeneity (high variability of leverage scores), especially compared to their number of differentially expressed genes (based on Anderson-Darling test). **(C)** Heatmap of chromosomal copy number inference from Perturb-seq data. For all genes (expressed >0.05 UMI per cell), the log-fold change in expression is calculated with respect to the average of non-targeting control cells, and genes are ordered along the genome. A weighted moving average of 100 genes is used infer copy number changes (columns) in single-cells (rows) with noise and median filtering. 80 *TTK* knockdown RPE1 cells and 80 randomly sampled non-targeting control RPE1 cells are shown. Cells are ordered by average linkage hierarchical clustering based on correlation of chromosomal copy number profiles. **(D, E)** Comparison of cell

cycle occupancy upon acute karyotypic changes. Abnormal karyotypic cells were defined as having ≥1 chromosome with evidence of changes in chromosomal copy number for >80% of the chromosomal length. For single-cells, cell-cycle positioning was inferred by UMAP dimension reduction on differential expression profiles of 199 selected cell-cycle regulated genes. Cell cycle occupancy is shown as a 2D kernel density estimate of a random subset of 1000 cells per karyotypic status. Approximate gates between cell cycle phases (G1 or G0; S; G2 or M) are shown as dotted lines, and the fraction of cells in each cell cycle phase are indicated. **(F)** Effect of chromosomal instability (CIN) on activation of the Integrated Stress Response (ISR). Histogram (kernel density estimate) compares the ISR score versus CIN status in RPE1 cells. CIN status is defined as evidence of gain or loss of chromosomal copy number for >80% of the chromosomal length, with 240,768 stable cells, 5,522 cells bearing chromosomal loss, 1987 cells bearing chromosomal gain, and 904 cells bearing gain and loss of chromosomes. ISR score is defined as the sum of z-normalized expression of ISR marker genes where increased values indicate stronger ISR activation. **(G)** Comparison of the effect of genetic perturbations on the CIN score across cell types. For each genetic perturbation, the CIN score is calculated as the mean single-cell sum of squared CIN values, *z*-normalized relative to non-targeting control perturbations. The CIN score is correlated across cell types (r=0.69). **(H)** Schematic of a subset of genetic perturbations that drive CIN. CIN drivers play diverse roles in mitosis, cell cycle regulation, and DNA repair.

Figure 6: Global organization of the transcriptional response to mitochondrial stress. **(A)** Clustering mitochondrial perturbations by nuclear transcriptional response. CRISPRi enables knockdown of nuclear-encoded genes whose protein products are targeted to mitochondria (mitochondrial perturbations). Mitochondrial perturbations were annotated by MitoCarta3.0 and subset to those with a strong transcriptional phenotype (n=268 mitochondrial perturbations). Gene expression profiles were restricted to nuclear encoded genes which includes 99% of mitochondrial proteins. The heatmap displays the Pearson correlation between pseudobulk *z*-normalized gene expression profiles of mitochondrial perturbations in K562 cells. Genetic perturbations were clustered by HDBSCAN with a correlation metric. **(B)** Comparing variability in the mitochondrial transcriptome by perturbation localization. The mitochondrial genome encodes 13 protein-coding

genes. Genetic perturbations were grouped based on localization of their protein products as determined by the Human Protein Atlas. For each of these 13 mitochondrially encoded genes, the variance in pseudobulk z-normalized expression profiles was calculated between all perturbations with the same localization. Barplots represent the average across genes with 95% confidence interval obtained by bootstrapping. **(C)** Clustering mitochondrial perturbations by mitochondrial transcriptional response. Mitochondrial perturbations were annotated by MitoCarta3.0 and subset to those with a strong transcriptional phenotype as above (n=268 mitochondrial perturbations). Gene expression profiles were restricted to the 13 mitochondrial-encoded genes. The heatmap displays the Pearson correlation between pseudobulk *z*-normalized gene expression profiles of mitochondrial perturbations in K562 cells. Genetic perturbations are clustered by HDBSCAN with a correlation metric. Clusters were manually annotated. **(D)** Heatmap visualizing the mitochondrial genome transcriptional response to diverse mitochondrial stressors. The expression (log$_2$ fold-change relative to non-targeting controls) of the 13 mitochondrially encoded genes is shown for a subset of perturbations representative of different mitochondrial complexes or function. Neither the ISR score nor mean fraction of mitochondrial RNA (mtRNA) would allow for high-resolution clustering by function as provided by the mitochondrial genome response. Genetic perturbations and genes are ordered by average linkage hierarchical clustering with a correlation metric.

Figure 7: Investigating regulation of the mitochondrial genome in stress. (A) Schematic of the mitochondrial transcriptome. Each human cell contains many copies of the circular 16.6 kb mitochondrial genome distributed throughout the mitochondrial network. The human mitochondrial genome encodes 2 rRNAs, 22 tRNAs, and 13 protein-coding genes. Both the heavy (H) and light (L) strand of the genome are transcribed as polycistronic transcripts punctuated by tRNAs. Excision of tRNAs from transcripts generates nascent mRNA precursors (colored by complex membership). mRNA precursors can then be polyadenylated, stabilized, or

degraded. **(B)** Density of Perturb-seq reads along the mitochondrial genome from select genetic perturbations. Reads are aligned to both the H-strand (dark grey) and L-strand (light grey). For each perturbation, densities are shown relative to the maximum read count in the locus. **(C)** Comparison of mitochondrial gene expression profiles between Perturb-seq and bulk RNA-seq. Heatmap displays $\log_2$-fold changes in expression of the 13 mitochondrial encoded genes (columns) for genetic perturbations (rows) in Perturb-seq and bulk RNA-seq data collected from K562 cells. Bulk RNA-seq was conducted to analyze total RNA (including non-polyadenylated RNA), with data representing the average of biological replicates. Genetic perturbations and genes are ordered by average linkage hierarchical clustering with a Euclidean distance metric. The profiles are strongly correlated ($r=0.79$, $p<10^{-39}$). **(D)** Clustering of *TMEM242* genetic perturbation based on the mitochondrial transcriptome. Genetic perturbations to members of ATP synthase and Complex I of the respiratory chain were compared to knockdown of *TMEM242*, a mitochondrial gene of unknown function. Gene expression profiles were restricted to the 13 mitochondrially encoded genes. The heatmap displays the Pearson correlation between pseudobulk *z*-normalized gene expression profiles of mitochondrial perturbations in K562 cells. Genetic perturbations are ordered by HDBSCAN with a correlation metric. **(E)** Effect of *TMEM242* knockdown on mitochondrial respiration. A Seahorse analyzer was used to monitor oxygen consumption rate (OCR). The Mito Stress Test consists of sequential addition of oligomycin (an ATP synthase inhibitor that enables measurement of ATP-productive respiration), FCCP (an uncoupling agent that enables measurement of maximal respiratory capacity), and a mixture of rotenone and antimycin A (inhibitors of Complex I and Complex III, respectively, that enable measurement of non-mitochondrial respiration). Data is presented as average ± SEM, n=6. **(F)** Schematic diagram of mitochondrial stress response.