Effects of footshock stress on social behavior and neuronal activation in the medial

prefrontal cortex and amygdala of male and female mice

Short title: *Stress, social behavior, and limbic system activation*

Mariia Dorofeikova[1,2], Chandrashekhar D Borkar[1,2], Katherine Weissmuller[3], Lydia Smith-

Osborne[1,4], Samhita Basavanhalli[3], Erin Bean[3], Avery Smith[3], Anh Duong[1,3], Alexis Resendez[1,2],

Jonathan P Fadok[1,2*]

Affiliations:

[1]Department of Psychology, Tulane University, New Orleans, LA 70118, USA.

[2]Tulane Brain Institute, Tulane University, New Orleans, LA 70118, USA.

[3]Neuroscience Program, Tulane University, New Orleans, LA 70118, USA.

[4]Tulane National Primate Research Center, Covington, LA 70433, USA.

*Corresponding author: Jonathan P. Fadok, jfadok@tulane.edu

Abstract

Social behavior is complex and fundamental, and its deficits are common pathological features for several psychiatric disorders including anxiety, depression, and posttraumatic stress disorder. Acute stress may have a negative impact on social behavior, and these effects can vary based on sex. The aim of this study was to explore the effect of two days of footshock stress on the sociability of male and female C57Bl/6J mice measured using a standard social interaction test. Animals were divided into two main groups of footshock exposure or context exposure control. Each group had mice that were treated with either the benzodiazepine alprazolam, or vehicle. In all groups, neuronal activation during social interaction was assessed using immunohistochemistry against the immediate early gene product cFos. Although footshock stress did not significantly alter sociability or latency to approach a social interaction counterpart, it did increase defensive tail-rattling behavior specifically in males. This stress-induced increase in tail-rattling was alleviated by alprazolam, yet alprazolam had no effect on female tail-rattling behavior in the stress group. Alprazolam lowered cFos expression in the medial prefrontal cortex, and social interaction induced sex-dependent differences in cFos activation in the ventromedial intercalated cell clusters. Overall, our results suggest that acute footshock stress induces sex-dependent alterations in defensiveness and patterns of cFos activation during social interaction tests.

Introduction

Social behavior is important for survival, and social deficits are common pathological features for a variety of mental illnesses including social anxiety disorder, depression, and posttraumatic stress disorder [1, 2]. More women than men suffer from these disorders, yet there is a paucity of data on sex differences in social behavior after stress [3]. There are data suggesting that childhood trauma leads to more aggressive behavior in men and more social withdrawal and avoidance in women [4], and women with social anxiety disorder report greater clinical severity, which may be associated with stressful life experiences [5]. Therefore, understanding sex differences in the effects of traumatic stress on social behavior, as well as the underlying neural substrates that potentially control this behavior, has important translational relevance.

Animal models of stress and trauma show alterations in many aspects of behavior [6]. In general, both acute and chronic stress have been found to lead to social fear and withdrawal in rodents [7]. One common model of traumatic stress in rodents is footshock exposure [8]. However, there is a lack of data investigating whether footshock stress influences mouse social behavior in a sex-dependent manner.

It is also unclear whether there are sex differences in the activation of brain regions responsive to social encounters following acute stress. Several lines of evidence suggest that neural activity in the medial prefrontal cortex (mPFC) is important for social behavior and subregions of the mPFC are differentially activated following social interaction. Neurons in the infralimbic cortex (IL) are preferentially activated in response to social cues compared to neurons in the prelimbic cortex (PL) [9], and a social neural ensemble within the IL may contribute to the social buffering of fear after fear conditioning [10].

Several subnuclei of the amygdala have also been implicated in social behavior. The role of the basolateral amygdala in social interaction following different stress paradigms has been well-established [12-14]. For example, activation of the basolateral amygdala leads to

reduced social interaction in a social interaction test [11]. The medial nucleus of the amygdala

(MeA) is also involved in both social behaviors and responses to stressors [15]. cFos

expression in the intercalated nucleus of the amygdala (ITC) is increased during social buffering

in rats [16], and social interaction is impaired in mice with altered migration and differentiation of

ITCs [17, 18]. Among the amygdala regions implicated in social behavior, the central nucleus of

the amygdala (CeA) is relatively unexplored. There is recent evidence, however, that CeA

circuits may be linked to sociability, and some manipulations of CeA activity impact social

behavior [18-21].

Thus, multiple brain regions are involved in social cognition [15, 16], but the differences

in social approach or avoidance behavior following acute stress are still poorly understood. In

the current study, our goal was to assess the extent to which there are sex differences in

sociability using a social interaction test that eliminates the possibility of direct physical

aggression. We hypothesized that footshock stress would negatively affect sociability, and that

those changes might depend on sex and be reversible with the fast-acting benzodiazepine

alprazolam. Therefore, we performed social interaction tests 24 hours after two consecutive

days of footshock stress. Additionally, we aimed to determine patterns of neuronal activation in

the mPFC, CeA, MeA, and ventral ITC associated with social behaviors using expression of the

immediate early gene cFos.

Materials and methods

Animals

2–4-month-old male and female C57BL/6J mice were obtained from the Jackson Laboratory

(Bar Harbor, ME, Stock No: 000664) and housed on a 12 h light/dark cycle with *ad libitum*

access to water and chow under standard laboratory conditions. Mice were individually housed

for 7 days before the start of and all throughout the experiments. Experiments were performed

during the light phase. All animal procedures were performed in accordance with institutional

guidelines and were approved by the Institutional Animal Care & Use Committee of Tulane

University. Unfamiliar strain-, sex- and age-matched mice were used as the passively

interacting counterparts (stimulus mice) during social interaction tests.

Footshock stress exposure

Footshock exposure or control context exposure was conducted in standard mouse operant

conditioning chambers (ENV-307W, Med Associates, Inc., St. Albans, VT) enclosed within

sound- and light-attenuating cubicles (ENV-022MD, Med Associates, Inc., St. Albans, VT). The

chambers were connected to a computer through an interface and controlled by MED-PC

software. The chamber was equipped with a grid floor and a house light, which was cleaned

using 70% ethanol.

Seven days after single housing, mice underwent footshock exposure for two

consecutive days. Each of the shock sessions included five 1 s, 0.9 mA footshocks presented

with a 120 s average pseudorandom intertrial interval (range 90–150 s), totaling 800 s in the

chamber. Mice in the control group were exposed to the same chambers for the same period

but did not experience footshock.

Social interaction test

The day after footshock exposure, the mice underwent the social interaction test in a square 46

x 46 X 38 cm arena constructed from sheets of white plexiglass. Behavioral videos were

recorded using a digital camera (Allied Vision "Pike" camera, Germany) and Plexon Studio

tracking software (Plexon, Dallas, TX). Tests were conducted under dim (10.6 lux) white

fluorescent lighting. Stimulus mice were single housed for 3 days before tests. Each of the

stimulus mice interacted with three experimental mice with at least 30 min between tests.

Experimental mice were perfused 90 min after the test to assess cFos expression.

An indirect social interaction method was chosen to avoid physical aggression between

male mice. For the first 3 min, mice were allowed to explore the open arena with two rectangular

(15 X 5 X 6 cm) or circular (8 cm diameter, 10 cm high) metallic mesh boxes located in

opposing corners 5 cm away from the walls. After the initial exploration, an unfamiliar, untreated

stimulus mouse was put underneath one of the boxes. Behavior was recorded for an additional

5 minutes, and sociability was scored using time spent sniffing the mesh box containing the

stimulus mouse as a percentage of total box interaction time (mouse preference, %), the latency

to approach the stimulus mouse, and the number of defensive tail rattles. Total exploration of

the mesh boxes was also scored in seconds to assess general activity. All behavioral

measurements were scored by an observer blinded to condition. Consistent with other social

interaction scoring protocols [22], sniffing directed to the upper and top part of the mesh boxes,

sniffing of feces, bar biting and circulating around the corral without sniffing, were not scored as

social interaction.

Alprazolam treatment

Alprazolam (Sigma-Aldrich, St. Louis, MO) was dissolved in a drop of Tween 80 (Merck,

Germany) and saline was added to make a final dose of 0.25 mg/kg. This dose was shown to

have anxiolytic effects [23] with minimal motor impairment in C57BL/6J mice [24]. Tween 80 +

saline solution was used for vehicle injections. Solutions were administered at 10 ml/kg volume,

intraperitoneally, 30 min before social interaction tests.

Histology

Following testing, mice were anesthetized with 2,2,2-tribromoethanol (240 mg/kg, ip, Sigma)

and subsequently transcardially perfused with 4% paraformaldehyde in phosphate-buffered

saline (PBS). cFos expression was assessed in mice that were perfused 90 min after the social

interaction test. Fixed brains were cut on a compresstome vibrating microtome (Precisionary,

Greenville, NC) in 80 μm coronal slices.

Antibody staining was performed on free-floating tissue sections. After 3 x 10 min

washes with 0.5% PBST slices were put in 5% donkey serum for 2 hours. Sections were then

incubated overnight in primary rabbit anti-cFos antibody (dilution 1:1500; #226 003, Synaptic

Systems, Germany) at 4°C. On the next day sections were washed in 0.5% PBST (3 X 10 min),

and then went through a 2 hr incubation with secondary donkey anti-rabbit antibody AlexaFluor

488 (dilution 1:500; #A-21206, Thermo Fisher Scientific, Waltham, MA) at 4°C. After 3 x 10 min

washes in PBS slices were mounted with mounting medium with DAPI (Biotium, Fremont, CA).

Images were obtained using an AxioScan.Z1 slide-scanning microscope (Zeiss,

Germany) and a Nikon A1 Confocal microscope (Nikon, Japan). cFos-positive nuclei were

quantified using Fiji ImageJ software (NIH, Bethesda, USA), and averaged for each animal. A

blinded observer quantified cFos expression in 2-5 slices per structure per mouse.

Statistical analysis

Data were analyzed using Prism 9 (GraphPad Software, San Diego, CA). The definition of

statistical significance was P ≤ 0.05. Two male mice that showed no interaction with either mesh

box during the social interaction test were excluded from analysis. One female outlier that

exhibited tail rattling behavior 7 times was also excluded. Because the data were

noncontinuous, tail-rattling was analyzed using Fisher's exact test. To assess the interaction of

factors, a 2- or 3-way ANOVA was used. If a significant effect was detected, Sidak's multiple

comparisons test was used, because it assumes independent comparisons and has more

power than the Bonferroni method. Correlations between cFos expression and behavior were

analyzed for control and stressed mice using pooled data from alprazolam and vehicle

treatment groups of both sexes. Using the Kolmogorov-Smirnov test, the data were determined

to be non-parametric, so Spearman's correlation coefficient was used for this analysis. Because

there was a Drug x Sex interaction and a main effect of Drug in cFos expression in the IL, we

did not include these data in the correlation analysis. For the sake of clarity, we report the

results of the interaction tests, the significant simple main effects, and the significant post-hoc

tests in the main text. The results of all tests are reported in **Table 1**. All statistical tests were

two-tailed.

Table 1: Results of Statistical Analyses

Effects of alprazolam x footshock stress x sex: 3-way ANOVA					
Source	Sum of squares	DFn	DFd	F	p
Fig. 1A Sociability (Mouse preference, %)					
Alprazolam*Stress*Sex	0.560	1	73	0.001	0.97
Alprazolam*Stress	2348	1	73	5.71	**0.02***
Alprazolam*Sex	182.4	1	73	0.44	0.51
Stress*Sex	347.5	1	73	0.85	0.36
Alprazolam	490.2	1	73	1.19	0.28
Stress	259.2	1	73	0.63	0.43
Sex	16.89	1	73	0.04	0.83
Fig. 1B Latency to approach					
Alprazolam*Stress*Sex	44.45	1	73	0.006	0.94
Alprazolam*Stress	13492	1	73	1.70	0.20
Alprazolam*Sex	12303	1	73	1.55	0.22
Stress*Sex	4270	1	73	0.54	0.47
Alprazolam	478.7	1	73	0.06	0.81
Stress	2646	1	73	0.33	0.57
Sex	11364	1	73	1.43	0.24
Fig. 1D Exploration					
Alprazolam*Stress*Sex	47.43	1	73	0.28	0.60
Alprazolam*Stress	257.0	1	73	1.51	0.22
Alprazolam*Sex	190.5	1	73	1.13	0.29
Stress*Sex	0.266	1	73	0.002	0.97
Alprazolam	540.9	1	73	3.20	0.08
Stress	180.2	1	73	1.06	0.31
Sex	3.060	1	73	0.02	0.89
Effects of alprazolam x footshock stress x sex on cFos: 3-way ANOVA					
Fig. 2C IL PFC					
Alprazolam*Stress*Sex	579981	1	41	2.43	0.13
Alprazolam*Stress	172328	1	41	0.72	0.40
Alprazolam*Sex	1378855	1	41	5.78	**0.02***
Stress*Sex	169279	1	41	0.71	0.40
Alprazolam	3014936	1	41	12.6	**0.001***
Stress	2231	1	41	0.01	0.92
Sex	538912	1	41	2.26	0.14
Fig. 2D PL PFC					

Alprazolam*Stress*Sex	657.3	1	33	0.008	0.93
Alprazolam*Stress	322417	1	33	3.86	0.06
Alprazolam*Sex	169139	1	33	2.03	0.16
Stress*Sex	7228	1	33	0.09	0.77
Alprazolam	546812	1	33	6.55	**0.02***
Stress	3566	1	33	0.04	0.84
Sex	29411	1	33	0.35	0.56
Fig. 2E Capsular CeA					
Alprazolam*Stress*Sex	24040	1	34	0.55	0.46
Alprazolam*Stress	19319	1	34	0.45	0.51
Alprazolam*Sex	54202	1	34	1.25	0.27
Stress*Sex	45335	1	34	1.05	0.31
Alprazolam	38512	1	34	0.89	0.35
Stress	36099	1	34	0.83	0.37
Sex	82985	1	34	1.91	0.18
Fig. 2F Medial CeA					
Alprazolam*Stress*Sex	5538	1	34	0.92	0.34
Alprazolam*Stress	3419	1	34	0.57	0.46
Alprazolam*Sex	8969	1	34	1.49	0.23
Stress*Sex	8175	1	34	1.35	0.25
Alprazolam	15566	1	34	2.59	0.12
Stress	307.6	1	34	0.05	0.83
Sex	24836	1	34	4.11	0.05
Fig. 2G Lateral CeA					
Alprazolam*Stress*Sex	6744	1	34	0.38	0.54
Alprazolam*Stress	10219	1	34	0.58	0.45
Alprazolam*Sex	762.9	1	34	0.04	0.84
Stress*Sex	11.84	1	34	0.0005	0.98
Alprazolam	11499	1	34	0.65	0.42
Stress	3797	1	34	0.22	0.64
Sex	69608	1	34	3.97	0.05
Fig. 2H ITC					
Alprazolam*Stress*Sex	56271	1	34	0.12	0.73
Alprazolam*Stress	46481	1	34	0.10	0.75
Alprazolam*Sex	153940	1	34	0.33	0.57
Stress*Sex	14920	1	34	0.03	0.86
Alprazolam	19359	1	34	0.04	0.84
Stress	9053	1	34	0.02	0.89
Sex	2104455	1	34	4.53	0.04*
Fig. 2I MeA					
Alprazolam*Stress*Sex	291674	1	34	1.87	0.18
Alprazolam*Stress	4035	1	34	0.03	0.87
Alprazolam*Sex	278969	1	34	1.79	0.19
Stress*Sex	103189	1	34	0.66	0.42
Alprazolam	38692	1	34	0.25	0.62
Stress	6305	1	34	0.04	0.84

163 **Figure 3A: Control group correlations (Spearman's coefficient)**

		PL	CeC	CeL	CeM	ITC	MeA
Sociability	r	-.236	-.627	-.320	.118	-.339	.313
	p	.347	.015*	.244	.674	.216	.255
Exploration	r	.118	.105	-.149	.294	.375	.501
	p	.641	.706	.593	.284	.168	.059
Tail rattling	r	.017	.115	-.184	.465	-.205	.130
	p	.947	.686	.519	.086	.486	.648
Latency to approach	r	.065	.557	.044	.171	-.016	.005
	p	.797	.033*	.876	.538	.957	.986

164 **Figure 3B: Footshock stress group correlations (Spearman's coefficient)**

		PL	CeC	CeL	CeM	ITC	MeA
Sociability	r	-.663	-.746	-.557	-.307	-.593	-.690
	p	.001**	<.001****	.007**	.165	.004**	<.001***
Exploration	r	-.057	-.310	-.104	.034	-.185	-.292
	p	.808	.161	.647	.882	.410	.187
Tail rattling	r	-.016	-.072	-.209	-.0149	-.208	-.053
	p	.944	.750	.351	.948	.353	.815
Latency to approach	r	.646	.385	.245	-.209	.399	.468
	p	.002**	.077	.271	.350	.066	.028*

Results

Effects of footshock stress and alprazolam treatment on social behavior

Following two days of footshock stress or context exposure, male (N = 18 control, N = 22 stress)

and female (N = 17 control, N = 24 stress) mice were allotted to the vehicle (N = 49, 24 males

and 25 females) and alprazolam (N = 32, 16 males and 16 females) treatment groups. These

mice were then subjected to a social interaction test designed to measure sociability (**Fig. 1A,**

B). A three-way ANOVA revealed that there was statistically significant interaction between the

effects of alprazolam and stress on sociability (**Fig. 1C**; drug X stress, $F_{(1,73)}$ = 5.7, *p* = 0.02).

Simple main effects analysis showed that there was no statistically significant effect of sex,

alprazolam, or stress on sociability (**Fig. 1C; Table 1**). There was no significant interaction

between stress, alprazolam, and sex on the latency to approach the social stimulus (**Fig. 1D**; 2-

way ANOVA, sex X stress, $F_{(1, 73)} = 0.0056$, $p = 0.94$). In addition, simple main effects analysis

showed that neither stress, alprazolam, nor sex had a significant effect. Further, we applied

Fisher's exact test to analyze noncontinuous tail-rattling data. Interestingly, stressed males

displayed significantly more tail-rattling behavior than control males during the social interaction

test ($p = 0.002$), while females in both groups displayed equivalent levels of tail rattling (**Fig. 1E**;

$p > 0.99$). On the other hand, alprazolam treatment significantly reduced tail-rattling in males (p

$= 0.03$) but did not affect tail-rattling in females. In the control vehicle condition, females show

higher tail-rattling than males, although this difference did not reach statistical significance ($p =$

0.07). Exploratory behavior, measured as total interaction time with both mesh boxes, was

unaffected by footshock stress or alprazolam treatment (**Fig. 1F**; 3-way ANOVA, sex X stress X

drug, $F_{(1, 73)} = 1.5$, $p = 0.22$).

Figure 1. Effects of acute stress and alprazolam on behavior in the social interaction test.

A. Experimental timeline. **B.** Schematic of the social interaction arena. **C.** There was a

significant interaction between the drug and stress condition, but there were no significant

results from post hoc multiple comparison tests. **D.** Latency to approach was not significantly

affected by stress or alprazolam treatment. **E.** Tail rattling was increased by stress and was

significantly reduced by alprazolam in males. **F.** Exploratory behavior was not significantly

altered by stress or drug treatment. Data are presented as means $\pm$ SEM. *$p<0.05$ post hoc

tests.

cFos expression analysis

We next quantified expression of cFos in several brain regions involved in the regulation of

social behavior (**Fig. 2A, B**). A three-way ANOVA was performed to test for the effects of sex,

stress, and alprazolam treatment on cFos expression in the IL, PL, CeA, ventromedial ITC, and

MeA (N = 13 control vehicle (5 male, 8 female), N = 10 control alprazolam (5 male, 5 female), N

= 14 stress vehicle (5 male, 9 female), N = 12 stress alprazolam (6 male, 6 female)). There was

no significant three-way interaction between the effects of these variables on cFos expression in

any of the brain areas analyzed (see **Table 1**). There was a significant sex by alprazolam

interaction effect on cFos expression in IL (**Fig. 2C**; $F_{(1,41)} = 5.78$, $p = 0.02$) and a main effects

analysis showed that alprazolam treatment significantly reduced cFos expression in the IL (drug

effect, $F_{(1,41)} = 12.6$, $p = 0.001$). Post hoc analysis showed that control females injected with

alprazolam had significantly fewer cFos+ cells in the IL compared to vehicle-injected female

controls (Sidak's multiple comparisons test, $p = 0.03$). There was a main effect of drug on cFos

expression in the PL (**Fig. 2D**; drug effect, $F_{(1,33)} = 6.55$, $p = 0.02$). There were no significant

simple main effects of sex, stress, or drug on cFos expression in the capsular subdivision of the

CeA (**Fig. 2E**), the medial (**Fig. 2F**, sex, $F_{(1, 34)} = 4.11$, $p = 0.0505$) or lateral subdivision of the

CeA (**Fig. 2G**, sex, $F_{(1, 34)} = 3.96$, $p = 0.0546$), or in the medial amygdala (**Fig. 2I**). There was a

significant effect of sex on cFos expression in the ventromedial intercalated nucleus of

amygdala, with greater expression levels in males (**Fig. 2H**; sex effect, $F_{(1,34)} = 4.53$, $p = 0.04$).

Figure 2. cFos expression patterns following social interaction. A. Schematic of the

experiment. **B.** Representative images of cFos staining results in mPFC (*left*) and amygdala

(*right*). IL = infralimbic mPFC, CeC = capsular CeA, CeL = lateral CeA, CeM = medial CeA, ITC

= ventromedial intercalated nucleus of amygdala, MEAd = dorsal MEA, MEAv = ventral MEA.

Scale: 1000 µm. **C-I.** Average number of cFos+ cells / mouse. **C.** Alprazolam reduced cFos

expression in the IL in control females. **D**. Alprazolam lowered cFos+ cells in the PL. **E-G.** cFos

expression in the capsular, medial, or lateral CeA was not significantly affected by stress, sex,

or drug. **H.** cFos expression in the ventromedial ITCs were greater in males. **I.** cFos expression

in the MeA was unaffected by stress, sex, or drug. Data are presented as means □±□SEM.

*p<0.05 post hoc tests. +p<0.05, main effect of sex, #p < 0.05, main effect of drug.

Correlations between cFos levels and behavior following the social interaction test

Spearman's correlation coefficient was used to assess the relationship between behavioral variables and cFos expression levels (**Fig. 3** and **Table 1**). These correlation analyses were performed for the control and stressed groups. For the brain areas in which we did not find any significant effect of drug or sex, we pooled the data from their respective vehicle or alprazolam treatment groups together for analysis. In controls (**Fig. 3A**), there was a significant negative correlation between CeC cFos expression and sociability (r = -0.63, p = 0.01), and a positive correlation with latency to approach (r = 0.56, p = 0.03). In the stress group (**Fig. 3B**), there was a significant negative correlation between sociability and cFos expression in numerous areas, with more socially active mice demonstrating less neuronal activation in the PL (r = -0.66, p = 0.001), capsular (CeC, r = -0.75, p = 0.0001) and lateral (r = -0.56, p = 0.007) subdivisions of CeA, as well as in the ventromedial ITC (r = -0.59, p = 0.003) and medial amygdala (MEA, r = -0.69, p = 0.0004).

Figure 3. Correlations between behavioral variables and cFos expression. Spearman's correlation coefficients (r) between the number of cFos+ cells and behavioral parameters in PL, CeA subdivisions, ITC and MeA of vehicle or alprazolam treated mice pooled together in the control (**A**) and stressed (**B**) groups. **C-D**, the distribution of correlation points for sociability and latency to approach against cFos+ cells/mm^2. Spearman's correlation coefficient represented with * indicating significant correlations (p<0.05). PL = prelimbic cortex, CeC = capsular CeA, CeM = medial CeA, CeL = lateral CeA, ITC = ventromedial intercalated nucleus of amygdala.

Latency to approach the social stimulus positively correlated with the number of cFos+ cells in the PL (r = 0.64, p = 0.001), and MEA (r = 0.47, p = 0.02), with a trend to significance in

CeC (r = 0.39, *p* = 0.07) and ITC (r = 0.40, *p* = 0.06; **Fig. 3B**). Distribution of data points and

regression lines for the respective brain area for sociability and latency is shown in **Fig. 3C-D**.

Discussion

Exposure to footshock is a commonly used model of acute traumatic stress to assay core

features of stress disorders such as social withdrawal [25]; however, sociability after footshock

stress has been investigated predominantly in male rodents. In the current study, several

measurements of social interaction behavior were taken in male and female mice to investigate

potential sex differences in the impact of footshock stress on sociability (**Fig. 1**) and neuronal

activation in the mPFC, CeA, and MEA (**Fig. 2**).

Our findings suggest that tail rattling behavior during social interaction is sex-dependent,

which is affected by stress specifically in males and diminished by alprazolam (**Fig. 1**). Tail

rattling is elicited due to territorial aggression in male mice [26, 27], and it has been suggested

that it can be a measurement of threat-induced defensive aggression [27]. Our results are

therefore comparable to other mouse sociability studies in which footshock stress produced an

enhancement of aggression in male mice [10]. Interestingly, female mice exhibit more tail

rattling than males during fear conditioning [28], suggesting that sex differences in tail rattling

behavior are both stress- and context-dependent. The significant reduction of tail rattling by

alprazolam further supports the link between this behavior and negative valence states. Overall,

the observed sex differences reinforce the idea that sex is a crucial factor that should be

considered in stress-related studies.

We found a significant interaction between the effects of alprazolam and stress on

preference for the social stimulus with alprazolam treatment leading to a decrease in sociability

in the control group, while there were no such trends in the stress group. There is evidence that

social interaction itself is anxiogenic, even under control conditions [29]. This is further

supported by the negative correlation between sociability and cFos expression in the CeA (**Fig.**

3). Acute stress by itself did not significantly lower sociability in contrast to other studies that

were focused on the effects of stress on social behavior, which mostly employed chronic stress

models [7]. Because the mice in our experiment underwent 9 days of single housing before the

social interaction test, it could be hypothesized that social isolation could enhance social

interest, thus leveling out social withdrawal [30].

Our results show that cFos expression in the ITC is higher in males compared to

females after social interaction. Given the known role of amygdala regions in mediating threat

responses [17, 31], these data suggest that social interaction may induce a higher level of

defensiveness in male mice. Consistent with this hypothesis, previous work has shown that CeA

neuronal activation is associated with male mouse aggression during social interaction in the

resident intruder assay [12, 32]. These findings emphasize the need to study sex differences

while deciphering the relationship between stress and social behaviors. Further investigations

should be conducted into sex differences in different neuronal populations of CeA, expressing

molecular markers such as somatostatin or corticotropin-releasing factor. This is especially

relevant given that there is a greater number of corticotropin-releasing factor receptor 1-

containing neurons in the male compared to female CeA, and they have different neuronal

excitability in response to corticotropin-releasing factor [33].

IL and PL cFos levels were decreased due to alprazolam treatment both in stressed and

control mice. Similarly, in humans, benzodiazepines decrease cerebral blood flow in PFC [34].

cFos expression in PL and IL mPFC in our study was not significantly affected by stress. Thus,

although the IL has been shown to be activated in response to a social cue [9], the mPFC does

not seem to modulate the interaction between traumatic stress and sociability in mice. Acute

restraint stress has been shown to hyperactivate PL, but not IL or CeA, and social interaction

following stress further activated PL PFC and did not affect total CeA activation [35]. A similar

lack of differences in cFos expression was observed in our study, which may occur due to the

delay between stress exposure and sacrifice of the animals.

Among control vehicle-injected mice, females had more cFos+ cells in IL. Similar results

were obtained by Tan et al. who observed that pyramidal neurons in the mPFC of females after

chronic adolescent social isolation stress showed a blunted increase of discharge rates during

sociability tests [36]. Sex differences in the modulation of social behavior by the mPFC were

previously explored regarding oxytocin-receptor-expressing interneurons which have differential

responses to oxytocin in male and female mice and regulate the social motivation of female

mice to interact with male mice during estrus [37]. Alprazolam significantly lowered the number

of cFos+ cells in the IL compared to vehicle-injected female controls, consistent with the

benzodiazepine effects observed in humans: women demonstrate decreased activity in frontal

regions after treatment, while an opposite effect is present in males [38]. This effect could be

related to females having significantly higher GABA-A benzodiazepine receptor availability [39].

The MEA has been linked to a wide variety of social behaviors, such as aggression,

mating, and parenting [40]. Here, we find that MEA cFos expression induced by the social

interaction test is not affected by sex, stress, or alprazolam. Prior studies have demonstrated

that socially defeated females housed with aggressive male residents exhibit increased cFos

activation in the medial amygdala [41], and the MEA is more responsive to aggressive than to

benign social interaction [42]. We designed our experiments to exclude the possibility of

aggressive interactions, therefore free interactions between the experimental and stimulus mice

were not possible. Future studies could investigate the effects of stress and sex on MEA

activation in free interaction paradigms.

In stressed animals, sociability was significantly negatively correlated with cFos

expression in the PL, CeC and CeL, as well as in the ventromedial ITC and MEA. In non-

stressed animals, MEA activation has been demonstrated previously as a result of social

interaction [43], and an opposite relationship has been shown between CeA cFos expression

and sociability after anxiogenic synthetic amphetamine treatment [44]. Because mPFC

activation has been demonstrated to suppress social behaviors [45], the negative correlation

observed in the current study could be expected, although enhanced activity of a subset of

mPFC neurons was correlated with social approach behavior previously [46]. In controls, there

was a significant negative correlation between sociability and CeC cFos expression, and an

opposite correlation between the latter and the latency to approach. Latency to approach the

social stimulus also positively correlated with the number of cFos+ cells in the PL and MEA.

The social consequences of stress have been extensively studied using the social defeat

stress model that entails exposure to emotional / psychological stress and leads to depression-

related outcomes [7]. After the social defeat procedure, most mice develop a decreased drive to

approach and interact with the social target [47]; however, social defeat stress is difficult to

achieve in female mice. Acute stress paradigms utilizing footshock facilitate investigations into

sex differences in social behavior following trauma. Future work should define the optimal

conditions, such as footshock intensity or lighting conditions, that influence sociability after

stress. It could be especially valuable to develop paradigms that stratify mice as resilient and

susceptible to further validate acute footshock stress as a tool for PTSD research.

Funding and Disclosure

The authors declare no competing interests.

This work was supported by the Louisiana Board of Regents through the Board of Regents

Support Fund (LEQSF(2018-21)-RD-A-17) and the National Institute of Mental Health of the

National Institutes of Health under award number R01MH122561 to JPF. The content is solely

the responsibility of the authors and does not necessarily represent the official views of the

National Institutes of Health.

Author contributions

Conceptualization—MD, CDB, JPF

Formal analysis—MD, CDB, JPF

Funding acquisition—JPF

Investigation—MD, CDB, LSO, KW, AR, SB, EB, AD, AS

Methodology—MD, CDB, JPF

Project Administration / Supervision—MD, JPF

Resources—JPF

Visualization—MD, CDB, JPF

Writing, original draft—MD, JPF

Writing, review & editing—MD, JPF, CDB, LSO, KW

References

1. Hames JL, Hagan CR, Joiner TE. Interpersonal processes in depression. Annu Rev Clin Psycho. 2013;9;355–377.

2. Dutton CE, Rojas SM, Badour CL, Wanklyn SG, Feldner MT. Posttraumatic stress disorder and suicidal behavior: Indirect effects of impaired social functioning. Arch Suicide Res. 2016;20(4);567–579.

3. Masi G, Berloffa S, Milone A, Brovedani P. Social withdrawal and gender differences: Clinical phenotypes and biological bases. J Neurosci Res. 2021. 10.1002/jnr.24802.

4. Ullman SE, Filipas HH. Gender differences in social reactions to abuse disclosures, post-abuse coping, and PTSD of child sexual abuse survivors. Child Abuse Negl. 2005;29(7); 767-782.

5. Asher M, Asnaani A, Aderka IM. Gender differences in social anxiety disorder: A review. Clin
Psychol Rev. 2007;56;1-12.

6.   Rincón-Cortés M, Herman JP, Lupien S, Maguire J, Shansky RM. Stress: Influence of sex,
reproductive status and gender. Neurobiol Stress. 2019;10;100155.

7. Toth I, Neumann ID. Animal models of social avoidance and social fear. Cell Tissue Res.
2013;354(1);107–118.

8. Zhang L, Hu XZ, Li H, Li X, Yu T, Dohl J, et al. Updates in PTSD animal models
characterization, methods. Mol Biol. 2019;2011;331-344.

9. Huang WC, Zucca A, Levy J, Page DT. Social behavior is modulated by valence-encoding
mPFC-amygdala sub-circuitry. Cell Rep. 2020;32(2);107899.

10. Gutzeit VA, Ahuna K, Santos TL, Cunningham AM, Sadsad Rooney M, Muñoz Zamora A, et
al. Optogenetic reactivation of prefrontal social neural ensembles mimics social buffering of
fear. Neuropsychopharmacol. 2020;45(6);1068-1077.

11. Paine TA, Swedlow N, Swetschinski L. Decreasing GABA function within the medial
prefrontal cortex or basolateral amygdala decreases sociability. Behav Brain Res.
2017;317;542-552.

12. Felix-Ortiz AC, Burgos-Robles A, Bhagat ND, Leppla CA, Tye KM. Bidirectional modulation
of anxiety-related and social behaviors by amygdala projections to the medial prefrontal
cortex. Neurosci. 2016;321;197-209.

13. Munshi S, Loh MK, Ferrara N, DeJoseph MR, Ritger A, Padival M. Repeated stress induces
a pro-inflammatory state, increases amygdala neuronal and microglial activation, and
causes anxiety in adult male rats. Brain Behav Immun. 2020;84;180-199.

14. Varlinskaya EI, Johnson JM, Przybysz KR, Deak T, Diaz MR. Adolescent forced swim stress
increases social anxiety-like behaviors and alters kappa opioid receptor function in the
basolateral amygdala of male rats. Prog Neuropsychopharmacol Biol Psychiatry.
2020;98;109812.

15. Nordman J, Ma X, Li Z. Traumatic stress induces prolonged aggression increase through

synaptic potentiation in the medial amygdala circuits. ENeuro. 2020;7(4);1-14.

16. Minami S, Kiyokawa Y, Takeuchi Y. The lateral intercalated cell mass of the amygdala is

activated during social buffering of conditioned fear responses in male rats. Behav Brain

Res. 2019;372;112065.

17. Kuerbitz J, Arnett M, Ehrman S, Williams MT, Vorhees CV, Fisher SE, et al. Loss of

intercalated cells (ITCs) in the mouse amygdala of Tshz1 mutants correlates with fear,

depression, and social interaction phenotypes. J Neurosci. 2018;38(5);1160-1177.

18. Wellman LL, Forcelli PA, Aguilar BL, Malkova L. Bidirectional control of social behavior by

activity within basolateral and central amygdala of primates. J Neurosci. 2016;36(33);8746-

8756.

19. Venniro M, Zhang M, Caprioli D, Hoots JK, Golden SA, Heins C, et al. Volitional social

interaction prevents drug addiction in rat models. Nat Neurosci. 2018;21(11);1520-1529.

20. Dong N, Du P, Hao X, He Z, Hou W, Wang L, et al. Involvement of GABAA receptors in the

regulation of social preference and emotional behaviors by oxytocin in the central amygdala

of female mandarin voles, Neuropeptides 66 (2017) 8-17.

21. Jones PW, Thornton AE, Jones AA, Knerich VM, Lang DJ, Woodward ML, et al. Amygdala

nuclei volumes are selectively associated with social network size in homeless and

precariously housed persons. Front Behav Neurosci. 2020;14;97.

22. Macbeth AH, Edds JS, Young WS 3[rd]. Housing conditions and stimulus females: a robust

social discrimination task for studying male rodent social recognition. Nature protoc.

2009;4(11);1574-1581.

23. Goyal R, Anil K. Protective effect of alprazolam in acute immobilization stress-induced

certain behavioral and biochemical alterations in mice. Pharmacol Rep. 2007;59(3);284-90.

24. Weizman R, Paz L, Backer MM, Amiri Z, Modai I, Pick CG. Mouse strains differ in their

sensitivity to alprazolam effect in the staircase test. Brain Res. 1999;839(1);58-65.

25. Schöner J, Heinz A, Endres M, Gertz K, Kronenberg G. Post-traumatic stress disorder and

beyond: an overview of rodent stress models. J Cell Mol Med.2017;21(10);2248-2256.

26. Aleyasin H, Flanigan ME, Golden SA, Takahashi A, Menard C, Pfau ML, et al. Cell-type-

specific role of ΔFosB in nucleus accumbens in modulating intermale aggression. J

Neurosci. 2018;38(26);5913-5924.

27. Nelson RJ, Chiavegatto S. Aggression in knockout mice. ILAR Journal 2000;41(3);153-162.

28. Borkar CD, Dorofeikova M, Le QE, Vutukuri RV, Vo C, Hereford D, et al. Sex differences in

behavioral responses during a conditioned flight paradigm. Behav Brain Res

2020;389;112623.

29. Hart PC, Bergne CL, Smolinsky AN, Dufour BD, Egan RJ, Laporte JL, et al. Experimental

models of anxiety for drug discovery and brain research. Methods Mol Biol. 2010;602;299-

321.

30. van den Berg CL, Pijlman FT, Koning HA, Diergaarde L, van Ree JM, Spruijt BM. Isolation

changes the incentive value of sucrose and social behaviour in juvenile and adult rats.

Behavioural Brain Research 1999;106;133–142.

31. Shackman AJ, Fox AS. Contributions of the central extended amygdala to fear and anxiety.

J Neurosci. 2016;36(31);8050-8063.

32. Veenema AH, Neumann ID. Neurobiological mechanisms of aggression and stress coping:

a comparative study in mouse and rat selection lines. Brain Behav Evol. 2007;70(4);274-85.

33. Agoglia AE, Tella J, Herman MA. Sex differences in corticotropin releasing factor peptide

regulation of inhibitory control and excitability in central amygdala corticotropin releasing

factor receptor 1-neurons. Neuropharmacol. 2020;180;108296.

34. Bagary M, Fluck E, File SE, Joyce E, Lockwood G, Grasby P. Is benzodiazepine-induced

amnesia due to deactivation of the left prefrontal cortex? Psychopharmacology (Berl).

2000;150(3);292-9.

35. Nosjean A, de Chaumont F, Olivo-Marin JC, Granon S. Stress-induced brain activation:

buffering role of social behavior and neuronal nicotinic receptors. Brain Struct Funct.

2018;223(9);4259-4274.

36. Tan T, Wang W, Liu T, Zhong P, Conrow-Graham M, Tian X, Yan Z. Neural circuits and

activity dynamics underlying sex-specific effects of chronic social isolation stress. Cell Rep.

2021;34(12);108874.

37. Nakajima M, Görlich A, Heintz N. Oxytocin modulates female sociosexual behavior through

a specific class of prefrontal cortical interneurons. Cell 2014;159(2);295-305.

38. Muñoz-Torres Z, Armony JL, Trejo-Martínez D, Conde R, Corsi-Cabrera M. Prefrontal

activity decline in women under a single dose of diazepam during rule-guided responses: an

fMRI study. Exp Brain Res. 2016;234(12);3483-3495.

39. Esterlis I, McKee SA, Kirk K, Lee D, Bois F, Stiklus SM, Seibyl JP, Krishnan-Sarin S,

O'Malley SS, Staley JK, Cosgrove KP. Sex-specific differences in GABA(A) -benzodiazepine

receptor availability: relationship with sensitivity to pain and tobacco smoking craving. Addict

Biol. 2013;18(2);370-8.

40. Raam T, Hong W. Organization of neural circuits underlying social behavior: A consideration

of the medial amygdala. Curr Opin Neurobiol. 2021;68;124-136.

41. Newman EL, Covington HE 3rd, Suh J, Bicakci MB, Ressler KJ, DeBold JF, et al. Fighting

females: neural and behavioral consequences of social defeat stress in female mice. Biol

Psychiatry. 2019;86(9);657-668.

42. Weathington JM, Strahan JA, Cooke BM. Social experience induces sex-specific fos

expression in the amygdala of the juvenile rat. Horm Behav. 2012;62(2);154-61.

43. Ferguson JN, Aldag JM, Insel TR, Young LJ. Oxytocin in the medial amygdala is essential

for social recognition in the mouse. J Neurosci. 2001;21(20);8278-85.

44. Navarro JF, Rivera A, Maldonado E, Cavas M, de la Calle A. Anxiogenic-like activity of 3,4-

methylenedioxy-methamphetamine ("Ecstasy") in the social interaction test is accompanied

by an increase of c-fos expression in mice amygdala. Prog Neuropsychopharmacol Biol

Psychiatry. 2004;28(2);249-54.

45. Benekareddy M, Stachniak TJ, Bruns A, Knoflach F, von Kienlin M, Künnecke B, Ghosh A.

Identification of a Corticohabenular Circuit Regulating Socially Directed Behavior. Biol

Psychiatry 2018;83(7);607-617.

46. Lee E, Rhim I, Lee JW, Ghim JW, Lee S, Kim E, Jung MW. Enhanced Neuronal Activity in

the Medial Prefrontal Cortex during Social Approach Behavior. J Neurosci.

2016;36(26);6926-36.

47. Milic M, Schmitt U, Lutz B, Müller MB. Individual baseline behavioral traits predict the

resilience phenotype after chronic social defeat. Neurobiol Stress. 2020;14;100290.

Figure legends

Figure 1. Effects of acute stress and alprazolam on behavior in the social interaction test. **A.** Experimental timeline. **B.** Schematic of the social interaction arena. **C.** There was a significant interaction between the drug and stress condition, but there were no significant results from post hoc multiple comparison tests. **D.** Latency to approach was not significantly affected by stress or alprazolam treatment. **E.** Tail rattling was increased by stress and was significantly reduced by alprazolam in males. **F.** Exploratory behavior was not significantly altered by stress or drug treatment. Data are presented as means $\pm$ SEM. *p<0.05 post hoc tests.

Figure 2. cFos expression patterns following social interaction. A. Schematic of the experiment. **B.** Representative images of cFos staining results in mPFC (*left*) and amygdala (*right*). IL = infralimbic mPFC, CeC = capsular CeA, CeL = lateral CeA, CeM = medial CeA, ITC = ventromedial intercalated nucleus of amygdala, MEAd = dorsal MEA, MEAv = ventral MEA. Scale: 1000 μm. **C-I.** Average number of cFos+ cells / mouse. **C.** Alprazolam reduced cFos expression in the IL in control females. **D**. Alprazolam lowered cFos+ cells in the PL. **E-G.** cFos expression in the capsular, medial or lateral CeA was not significantly affected by stress, sex, or drug. **H.** cFos expression in the ventromedial ITCs were greater in males. **I.** cFos expression in the MeA was unaffected by stress, sex, or drug. Data are presented as means $\pm$ SEM. *p<0.05 post hoc tests. +p<0.05, main effect of sex, #p < 0.05, main effect of drug.

Figure 3. Correlations between behavioral variables and cFos expression. Spearman's correlation coefficients (r) between the number of cFos+ cells and behavioral parameters in PL, CeA subdivisions, ITC and MeA of vehicle or alprazolam treated mice pooled together in the control (**A**) and stressed (**B**) groups. **C-D**, the distribution of correlation points for sociability and

516   latency to approach against cFos+ cells/mm^2. Spearman's correlation coefficient represented

517   with * indicating significant correlations (p<0.05). PL = prelimbic cortex, CeC = capsular CeA,

518   CeM = medial CeA, CeL = lateral CeA, ITC = ventromedial intercalated nucleus of amygdala.

A

| Days 1 & 2 | Foot shock exposure or conditioning chamber exposure |

| Day 3 | Alprazolam / vehicle Social interaction test Perfusion |

B

C

Control   Stress

Sociability, %

Males Females   Males Females

D

Control   Stress

Latency to approach, s

Vehicle
Alprazolam

Males Females   Males Females

E

Control   Stress

Tail-rattling

**

*

Males Females   Males Females

F

Control   Stress

Total exploration, s

Males Females   Males Females