ACE2-independent SARS-CoV-2 infection and mouse adaption emerge after passage in cells

expressing human and mouse ACE2

Kexin Yan[1], Troy Dumenil[1], Bing Tang[1], Thuy T. Le[1], Cameron Bishop[1], Andreas Suhrbier[1,2],

Daniel J. Rawle[1*].

[1] Immunology Department, QIMR Berghofer Medical Research Institute, Brisbane, Queensland.

4029, Australia.

[2] Australian Infectious Disease Research Centre, GVN Center of Excellence, Brisbane,

Queensland, 4029 and 4072, Australia.

[*]Corresponding author. Email: Daniel.Rawle@qimrberghofer.edu.au

SUMMARY

Human ACE2 (hACE2) is the key cell attachment and entry receptor for SARS-CoV-2, with the original SARS-CoV-2 isolates unable to use mouse ACE2 (mACE2). Herein we describe a new system for generating mouse-adapted SARS-CoV-2 *in vitro* by serial passaging virus in co-cultures of cell lines expressing hACE2 and mACE2. Mouse-adapted viruses emerged with up to five amino acid changes in the spike protein, all of which have been seen in human isolates. Mouse-adapted viruses replicated to high titers in C57BL/6J mouse lungs and nasal turbinates, and caused severe lung histopathology. One mouse-adapted virus was also able to replicate efficiently in ACE2-negative cell lines, with ACE2-independent entry by SARS-CoV-2 representing a new biology for SARS-CoV-2 that has potential widespread implications for disease and intervention development.

INTRODUCTION

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) emerged in 2019, causing a global pandemic of coronavirus disease 2019 (COVID-19)[1]. The human ACE2 (hACE2) receptor is generally held to be the key cell attachment and entry receptor for SARS-CoV-2, with the receptor binding domain (RBD) of the spike protein binding to hACE2[2]. ACE2 binding has deep ancestral origins within the sarbecovirus lineage of coronaviruses[3], with SARS-CoV, SARS-CoV-2, and SARS-related bat coronaviruses all using ACE2 as their entry receptor[4]. Genetically diverse spike proteins are able to bind ACE2[3,4], with all the SARS-CoV-2 variants of concern (Alpha, Beta, Gamma, Delta and Omicron) reported to require ACE2 binding for efficient infection[5,6].

The original isolates of SARS-CoV-2 (Wuhan strain) are unable to bind mouse ACE2 (mACE2) for infection[2], therefore, to study these viruses a series of mouse models were developed that express hACE2[2,7-10]. Some subsequent SARS-CoV-2 variants of concern emerged to be able to use mACE2, including Alpha, Beta, Gamma, and Omicron, but not Delta[6,11]. In addition, mouse-adapted SARS-CoV-2 have been generated by serial passage of original SARS-CoV-2 isolates in mouse lungs[12-16] or by reverse genetics[17] to introduce amino acid changes that allow binding to mACE2.

As infection of wild animals by SARS-CoV-2 is increasingly being reported[18-23], with omicron also potentially arising from rodents[24,25], we further sought to characterize the process of mouse adaptation. We have previously reported the use of HEK293T cells that express hACE2 (HEK293T-hACE2) or mACE2 (HEK293T-mACE2) by virtue of lentiviral transduction, with the former, but not the latter, able to support efficient replication of an original SARS-CoV-2 isolate, hCoV-19/Australia/QLD02/2020 (SARS-CoV-2$_{QLD02}$)[2]. We also showed that

transduction with mACE2 containing the N31K and H353K mouse-to-human amino acid

changes (to generate HEK293T-mACE2$^{N31K/H353K}$ cells), represented the minimum changes

required to support SARS-CoV-2 $_{QLD02}$ replication[2]. Herein we serially passaged SARS-CoV-

2$_{QLD02}$ in HEK293T-hACE2 or HEK293T-mACE2$^{N31K/H353K}$ cells co-cultured with HEK293T-

mACE2 cells, followed by passaging in HEK293T-mACE2 cells to generate five different

mouse-adapted (MA) viruses. These MA viruses were able to replicate in both HEK293T-

hACE2 and HEK293T-mACE2 cells. These viruses were also able to replicate in C57BL/6J

mice, leading to characteristic COVID-19 histopathological lesions and inflammatory responses.

Remarkably, one of these viruses (MA1) was also able to replicate in HEK293T cells in the

absence of lentiviral ACE2 expression. Furthermore, MA1 replicated in a number of other cell

lines that do not express significant levels of ACE2 and do not support SARS-CoV-2$_{QLD02}$

infection. This represents just the second report of productive infection in unmodified ACE2-

negative cell lines[26], with potential implications for the biology of SARS-CoV-2 infections,

evolution, tropism, disease and perhaps also intervention development.

RESULTS

SARS-CoV-2 $_{QLD02}$ co-culture passaging to select for MA viruses

To investigate the process of mouse adaptation, SARS-CoV-2$_{QLD02}$ was passaged in co-cultures

of HEK293T-hACE2 and HEK293T-mACE2, or HEK293T-mACE2$^{N31K/H353K}$ and HEK293T-

mACE2 cells, followed by passage in HEK293T-mACE2 cells (Fig. 1a. After 4 co-culture

passages, no viruses emerged that were able to replicate efficiently in HEK293T-mACE2 cells

(Fig. 1b, Passage 4). However, after 9 co-culture passages, 5 viruses (MA1-5) demonstrated a

clear ability to replicate in HEK293T-mACE2 cells (Fig. 1b, Passage 9). As expected, SARS-

CoV-2$_{QLD02}$ did not replicate in HEK293T-mACE2 cells (Fig. 1b). All 5 MA viruses showed

overt cytopathic effects (CPE) in HEK293T-mACE2 cells 2 days after infection (Fig. 1c).

Importantly, these viruses were also able to replicate in lungs of wild-type C57BL/6J mice after

intrapulmonary inoculation (via the intranasal route) with 1.5×10^4 CCID$_{50}$ per mouse (Fig. 1d).

MA1 and MA2 showed the highest lung titers, replicating to above 10^6 CCID$_{50}$/g (Fig. 1d). As

expected SARS-CoV-2$_{QLD02}$ was unable to replicate in C57BL/6J mice (Fig. 1d). The co-culture

and *in vitro* passaging of SARS-CoV-2$_{QLD02}$ thus generated five viruses able to replicate in wild-

type mice.

Amino acid changes in MA viruses have previously been reported in human isolates

MA1-5 viruses were sequenced before and after infection of C57BL/6J mice (Fig. 1a,

Sequencing) by RNA-Seq. Sequences for MA viruses before and after infection of C57BL/6J

mice (Fig. 1a, Sequencing) were almost identical (Supplementary File 1), illustrating that the

single passage in mice did not generate further significant changes. There were 13 different

amino acid changes and two deletions in the spike protein across the five MA viruses (Fig. 2a,

b). All these changes have previously been identified in human isolates (Fig. 2c). The changes

are described in detail, with structural information in Supplementary Figs. 1-3. The Q498H and

Q493R substitutions (Fig. 2a) have been identified previously for mouse adapted viruses and

increase affinity for mACE2[12-17]. The latter contention is supported by our modeling

(Supplementary Fig. 2). The non-spike changes comprise nine different amino acid substitutions

in Orf1ab (Fig. 2a); with all these changes also identified in human isolates (Supplementary Fig.

4).

MA1 replicates efficiently in several ACE2-negative cell lines

In vitro growth kinetics of MA1 and MA2 were studied in more detail using stock virus derived from infected mouse lungs and prepared in HEK293T-mACE2 cells (Fig. 1a). As expected, MA1 and MA2, as well as SARS-CoV-2$_{QLD02,}$ and Alpha and Beta strain viruses, all replicated in HEK293T-hACE2 cells (Fig. 3a). MA1, MA2 and Alpha, Beta, and Omicron strain viruses, but not SARS-CoV-2$_{QLD02}$, replicated in HEK293T-mACE2 cells (Fig. 3b); consistent with Fig. 1b-d and the reported ability of Alpha, Beta and Omicron variants to replicate in mice[6]. Perhaps of interest, given the postulated rodent origins of Omicron[24,25], our Omicron isolate replicated more efficiently in HEK293T-mACE2 than in HEK293T-hACE2 cells (Fig. 3a, b).

Remarkably, when replication was examined in control untransduced HEK293T cells, both MA1 and MA2 showed significant replication, with MA1 replicating to ≈8 log$_{10}$CCID$_{50}$/ml in 2 days (Fig. 3c). None of the other viruses were able to replicate efficiently in untransduced HEK293T cells (Fig. 3c), consistent with previous reports[2,27,28]. Replication was also tested in a series of cell lines (Fig. 3d-i) in which (i) publically available RNA-Seq data showed negligible levels of ACE2 mRNA expression (Supplementary Fig. 5a) or (ii) surface expression of ACE2 is absent (LLC-PK1)[29]. MA1 was able to replicate in all the aforementioned cell lines (Fig. 3d-i). In contrast, MA2 and SARS-CoV-2$_{QLD02}$ were unable to replicate in any of these cell lines (Fig. 3d-i); the SARS-CoV-2$_{QLD02}$ results are consistent with previous reports[2,29-32]. Overall, the data suggests ACE2-independent infection by MA1 is robust across multiple cell lines and is functional, at least *in vitro*, across 4 different species (human, mouse, hamster and pig).

None of these cell lines expressed TMPRSS2 mRNA (Supplementary Fig. 5b), with virus

fusion instead likely to rely on cathepsin L[33], with cathepsin L mRNA abundantly expressed in

all the cell lines (except BHK-21 which does not have an annotated orthologue) (Supplementary

Fig. 5c).

MA1 and MA2 infection of C57BL/6J mice

To characterize the behavior of MA1 and MA2 during mouse infections, C57BL/6J mice

were infected by intrapulmonary inoculation (via the intranasal route) with 10^5 CCID$_{50}$ MA1 or

MA2 per mouse. Virus titers in lungs reached 6-8 log$_{10}$ CCID$_{50}$/g on day 2, with no significant

differences seen between MA1 and MA2 (Fig. 4a). These results are broadly similar to those

reported previously for mouse adapted viruses[12,17]. Nasal turbinate titers on day 2 were

significantly higher for MA1 ($\approx$7 log$_{10}$ CCID$_{50}$/g), when compared with MA2 (Fig. 4b). MA1

and MA2 virus titers were below the level of detection for all other mice and tissues tested

(similar to other reports[14]), with the exception of heart where MA1 infection resulted in

detectable heart infection on day 2 in 2/4 mice (Fig. 4c). (Infectious virus can also be detected in

the heart of K18-hACE mice – Supplementary Fig. 6a). There was no significant weight loss

observed for MA1 or MA2 infected mice (Supplementary Fig. 6b).

Immunohistochemistry (IHC) staining with anti-SARS-CoV-2 spike monoclonal

antibody showed that MA1 virus infection was localized primarily to the epithelium of bronchi

(Fig. 4d-e) and bronchioles (Fig. 4f) in C57BL/6J mouse lungs, with occasional staining of cells

in the alveoli (Fig. 4g). A similar picture emerged for MA2 (Supplementary Fig. 7). There was

also occasionally prodigious staining of bronchial epithelium in MA1 infected lungs

(Supplementary Fig. 8a). Other studies using mouse-adapted SARS-CoV-2 viruses also showed

viral staining in bronchial epithelium[14,17,34], and such staining is also seen after infection with

non-mouse adapted viruses of mice expressing hACE2 from the mACE2 promoter[35]. Staining of

alveoli was not a prominent feature in MA1 (Fig. 4d, g) or MA2 infected (Supplementary Fig. 7)

C57BL/6J mice, with similar results reported for SARS-CoV-2 infection of mice expressing

hACE2 from the mACE2 promoter[35]. This contrasts markedly with SARS-CoV-2 infection of

K18-hACE2 mice, which show widespread infection of alveoli[35]. Occasional staining of MA1

was also observed in tracheal columnar epithelial cells (Supplementary Fig. 8b), indicating upper

respiratory tract infection. Omicron viruses similarly show a propensity to replicate in the upper

respiratory tract, rather than in lung parenchyma[36-38].

Mouse-to-mouse (C57BL/6J to K18-hACE2) transmission was not seen for either MA1

or MA2 (Supplementary Fig. 9a,b). Such transmission has been reported for K18-hACE2 to

K18-hACE2 mice[39] and for deer mice[23]. The complete loss of an intact furin cleavage site in

MA1 and MA2 may, at least in part, be responsible (Supplementary Fig. 9c), with this site

required for transmission in ferrets[40], as well as pathogenicity in humans[41]. Loss of this site

commonly arises after *in vitro* passage of SARS-CoV-2[42] and provides a built-in safety feature

for our MA viruses.

Lung pathology after MA1 and MA2 infections of C57BL/6J mice

C57BL/6J mice were infected with MA1 and MA2 and lungs harvested on days 0, 2, 4 and 7

post infection and analyzed by hematoxylin and eosin (H&E) staining. Examples of low

magnification images of whole lung sections illustrate the loss of alveolar airspace (Fig. 5a),

which reached significance by image analysis on day 7 for MA1 infected mice (Fig. 5b). The

ratio of blue (nuclear) to red (cytoplasmic) pixels in H&E stained sections is a measure of

cellular infiltration[43], with lungs from MA1 or MA2 infected mice showing significantly higher

cellular infiltrates compared to uninfected mice if all days post infection are taken together (Fig.

5c).

H&E staining revealed clear evidence of bronchiolar sloughing and smooth muscle

hyperplasia in MA1-infected mice (Fig. 5d). The collapse of alveolar spaces, with some

pulmonary edema is also evident (Fig. 5e) in MA1 infected lungs, consistent with Fig. 5b. Areas

with dense cellular infiltrates were occasionally pronounced in MA1-infected lungs (Fig. 5e,

dotted circle), consistent with Fig. 5c. Overall similar lung pathology was seen in MA2-infected

mice (Supplementary Fig. 10). MA infection of C57BL/6J mice results in robust lung pathology,

with no overtly unique features when compared, for instance, with the infection of K18-hACE2

mice with SARS-CoV-2$_{QLD02}$ [9].

Inflammatory responses in MA1 infected C57BL/6J mouse lungs are similar to, but less

severe than, SARS-CoV-2$_{QLD02}$ infection of K18-hACE2 mouse lungs

RNA-Seq was used to determine the inflammatory responses in C57BL/6J mouse lungs on day 4

post MA1 infection (full gene list in Supplementary File 2a). Differentially expressed genes

(DEGs) (n = 1027, Supplementary File 2b) were analyzed for 'UpStream Regulators' (USRs)

(Supplementary File 2c) and 'Diseases and Functions' (Supplementary File 2d) tools of the

Ingenuity Pathway Analysis (IPA) software. The same analyses were performed for day 4 lungs

from K18-hACE2 mice infected with SARS-CoV-2$_{QLD02}$[44] (n=1349 DEGs; Supplementary File

2e-h, respectively). A highly significant correlation emerged for cytokine USRs (Fig. 6a;

Supplementary File 2i). A number of USR annotations (Fig. 6a, red) indicated that inflammation

in K18-hACE2 is more severe (i) with higher z scores for the pro-inflammatory cytokine TNF,

and osteopontin (SPP1), a proposed severity marker of COVID-19 [45], and (ii) lower z-scores for

anti-inflammatory cytokines IL10 and IL37, and Secretoglobin family 1a member 1

(SCGB1A1), a pulmonary surfactant protein that blunts alveolar macrophage responses and

mitigates against cytokine surges[46].

The Disease and Functions annotation z scores also showed a highly significant correlation,

although K18-hACE2 annotations again indicated a more severe inflammatory response, with

increased cell movement, cell death, and weight loss (Fig. 6b; Supplementary File 2j), consistent

with histology[9] (Fig. 5) and body weight measurements[44] (Supplementary Fig. 6b). Thus

although MA1 can infect cells lacking ACE2, the inflammatory responses in C57BL/6J mice

were very similar to those seen in the K18-hACE2/SARS-CoV-2$_{QLD02}$ model, which is

dependent on hACE2.

DISCUSSION

Herein we describe the rapid *in vitro* adaptation of an original isolate of SARS-CoV-2 to

produce viruses capable of infecting mice, with one virus (MA1) also capable of ACE2-

independent infection. The latter was robust across multiple cell lines and was functional in at

least 4 different species (human, mouse hamster and pig). Our study again highlights the ability

of SARS-CoV-2 to evolve rapidly in the face of selection pressure[47], a feature that has also been

documented in individual patients[48]. Our study again highlights the relatively small number of

changes (that include Q498H, Q493R/K, and N501T; Supplementary Fig. 1) that allow SARS-

CoV-2 to jump species[2,12-14,16,17].

The ability to infect cells in an ACE2-independent manner, would appear to require the

E484D substitution, identified herein and previously[26]. This substitution has been identified in

multiple human isolates, in all variants of concern and in a large range of geographical locations

around the world (Supplementary Fig. 11). Omicron has an E484A substitution in this position,

with Omicron requiring ACE2 expression for infection (Fig. 3)[36]. Modeling suggests the E484D

substitution retracts the carboxyl group located between Y489 and F490 of the RBD

(Supplementary Fig. 12), perhaps thereby providing a novel binding site. A series of alternative

or additional receptors have been reported for SARS-CoV-2 Wuhan strain and include DC-

SIGN/L-SIGN[49], Kidney Injury Molecule-1/T cell immunoglobulin mucin domain 1 (KIM-

1/TIM-1)[29], AXL Receptor Tyrosine Kinase[28], neuropilin 1 (NRP1)[27] and CD147[50]. Analysis of

publically available RNA-Seq data indicated that only CD147 was expressed at high levels in all

cell lines tested herein (Supplementary Fig. 13). However, Wang *et al*. reported replication of an

original isolate of SARS-CoV-2 in BHK-21 cells only when human CD147 was over-

expressed[50]. The MA viruses are likely to enter via the endosomal route, given the loss of the

furin cleavage site[37,51]. This loss improves cleavage by Cathepsin L, which substitutes for

TMPRSS2 by cleaving S2' in endosomes and releasing viral RNA into the cytoplasm[33].

Omicron, for different unknown reasons, shows reduced reliance on TMPRSS2, and relies more

on endosomal entry[36,52,53]. The D614G change, which is present in all SARS-CoV-2 variants

(except the original Wuhan strain), also increases entry via the cathepsin L route[33]. Cathepsin L

is widely expressed in human nasal and lung epithelial cells[33] and in the cell lines tested herein

(Supplementary Fig. 5c).

The concept that Omicron might have arisen via rodents[24,54] is perhaps supported by the

observation that Omicron replicated more efficiently in mACE2-expressing cells when compared

with hACE2-expressing cells (Fig. 3a,b). Omicron binding to mACE2 is significantly higher

than Alpha and Beta variants, which are also able to use mACE2 as a receptor[55]. Increased

mACE2-binding by Omicron may involve the Q493R, Q498R and N501Y substitutions (Fig. 2c,

Omicron), given these changes are selected in mouse-adapted viruses[12-15,17]. Our MA viruses

have Q493R, Q498H and N501T substitutions (Fig. 2c; Supplementary Fig. 1), with modeling

suggesting Q493R and Q498H increase mACE2 binding (Supplementary Fig. 2). Finally, like

Omicron[36], MA1, MA2 and other mouse adapted viruses[14,17], show a preference for replication in

the upper respiratory tract.

The pathological consequences of ACE-2 independent infection remain to be characterized,

and one might speculate that ACE-independent infection may play a role in long-COVID[56,57].

Our studies of acute MA1 infections in C57BL/6J mice argue that when ACE2-dependent

infection remains possible, the ability also to infect via an ACE2-independent mechanism does

not result in overtly unique pathology or immunopathology. Both the development of mouse

models of long COVID, evaluation of infection and disease in ACE2$^{-/-}$ mice and the

identification of the alternate receptor used by MA1 are thus the focus of our future studies.

ACKNOWLEDGEMENTS

From QIMR Berghofer Medical Research Institute we thank Dr I Anraku for managing the PC3

(BSL3) facility, animal house staff for mouse breeding and agistment, Paul Collins for library

preparation and RNA-Seq, and Dr Clay Winterford, Sang-Hee Park and Crystal Chang for the

histology and immunohistochemistry. We thank Dr Alyssa Pyke and Mr Fredrick Moore

(Queensland Health, Brisbane) for providing the SARS-CoV-2 isolates. We thank Monash

Genome Modification Platform for providing the plasmid containing the mouse-codon optimized

human ACE2 gene. We thank Clive Berghofer and the Brazil Family Foundation (and many

others) for their generous philanthropic donations to support SARS-CoV-2 research at QIMR

Berghofer MRI. The project was also partly funded by an intramural QIMRB seed grant

awarded to D.J.R. A.S. holds an Investigator grant from the National Health and Medical

Research Council (NHMRC) of Australia (APP1173880).

AUTHOR CONTRIBUTIONS

275　Conceptualization, D.J.R.; Methodology, D.J.R. and A.S.; Formal analysis, D.J.R., A.S., T.D.,

276　C.B.; Investigation, D.J.R., K.Y., T.T.L., and B.T.; Resources, A.S.; Data curation, D.J.R., A.S.,

277　and T.D.; Writing – D.J.R., AS; Visualization, D.J.R., A.S., and T.D.; Supervision, D.J.R. and

278　A.S., Project administration, D.J.R. and A.S.; Funding acquisition, A.S. and D.J.R.

DECLARATION OF INTERESTS

281　The authors declare no competing interests.

DATA AVAILABILITY

284　All raw sequencing data (fastq files) are available from the Sequence Read Archive (SRA),

285　BioProject accession: PRJNA804321. All other data is available within the paper and supporting

286　information files.

MATERIALS and METHODS

Ethics statement and regulatory compliance

All mouse work was conducted in accordance with the "Australian code for the care and use of

animals for scientific purposes" as defined by the National Health and Medical Research Council

of Australia. Mouse work was approved by the QIMR Berghofer Medical Research Institute

Animal Ethics Committee (P3600, A2003-607). For intrapulmonary inoculations, mice were

anesthetized using isoflurane. Mice were euthanized using CO_2 or cervical dislocation. Obtaining

samples from patients and using human SARS-CoV-2 isolates was approved by the QIMR

Berghofer MRI Human Research Ethics Committee (P3600).

Breeding and use of GM mice was approved under a Notifiable Low Risk Dealing (NLRD)

Identifier: NLRD_Suhrbier_Oct2020: NLRD 1.1(a). Cloning and use of lentiviral vectors for

transduction of ACE2 into cell lines was approved under an NLRD (OGTR identifier:

NLRD_Suhrbier_Feb2021: NLRD 2.1(l), NLRD 2.1(m)).

All infectious SARS-CoV-2 work was conducted in a dedicated suite in a biosafety level-3

(PC3) facility at the QIMR Berghofer MRI (Australian Department of Agriculture, Water and the

Environment certification Q2326 and Office of the Gene Technology Regulator certification

3445). All work was approved by the QIMR Berghofer Medical Research Institute Safety

Committee (P3600).

Cell lines and SARS-CoV-2 culture

Vero E6 (C1008, ECACC, Wiltshire, England; obtained via Sigma Aldrich, St. Louis, MO,

USA), Lenti-X 293T (Takara Bio), AE17 (a gift from Dr Delia Nelson, Faculty of Health

Sciences, Curtin Medical School), NIH-3T3 (American Type Culture Collection, ATCC, CRL-

1658), LLC-PK1 (a gift from Prof. Roy Hall, University of Queensland), A549 (ATCC CCL-

185), BHK-21 (ATCC# CCL-10) and HeLa (ATCC-CLL 2) cells were cultured in medium

comprising DMEM for Lenti-X 293T and A549 cells, M199 for LLC-PK1 cells or RPMI1640

for all the others (Gibco), supplemented with (endotoxin free[58]) 10% fetal bovine serum (FBS),

penicillin (100 IU/ml)/streptomycin (100 µg/ml) (Gibco/Life Technologies) and L-glutamine (2

mM) (Life Technologies). Cells were cultured at 37°C and 5% CO_2. Cells were routinely

checked for mycoplasma (MycoAlert Mycoplasma Detection Kit MycoAlert, Lonza) and FBS

was assayed for endotoxin contamination before purchase[58].

The SARS-CoV-2 original, Alpha and Beta isolates were kindly provided by Dr Alyssa

Pyke (Queensland Health Forensic & Scientific Services, Queensland Department of Health,

Brisbane, Australia). The viruses (hCoV-19/Australia/QLD02/2020, Alpha variant hCoV-

19/Australia/QLD1517/2021 and Beta variant hCoV-19/Australia/QLD1520/2020) were isolated

from patients and sequences deposited at GISAID (https://www.gisaid.org/). Virus stocks were

generated by infection of Vero E6 cells at multiplicity of infection (MOI)≈0.01, with supernatant

collected after 2-3 days, cell debris removed by centrifugation at 3000 x g for 15 min at 4°C, and

virus aliquoted and stored at -80°C. Virus titers were determined using standard $CCID_{50}$ assays

(see below). The virus was determined to be mycoplasma free using co-culture with a non-

permissive cell line (i.e. HeLa) and Hoechst staining as described[59].

An Omicron isolate was isolated from a SARS-CoV-2 infected individual in Brisbane

Australia. A nasal swab was acquired, submerged and agitated in 1 ml RPMI + 10% FBS +

penicillin (100 IU/ml)/streptomycin (100 µg/ml) (Gibco/Life Technologies), and was transported

to the laboratory on ice. After centrifugation at 20,000 x g for 10 min the supernatant was

cultured on $2x10^6$ Vero E6-TMPRSS2 cells[9] in a T25 flask for 3 days at 37°C and 5% CO_2; 1 ml

was passaged on to a confluent T75 flask of Vero E6-TMPRSS2 cells, and supernatant harvested

and aliquoted after 3 days. Virus titers were determined using standard $CCID_{50}$ assays (see

below). For sequencing, viral RNA was isolated from infected Vero E6-TMPRSS2 cells by

TRIzol extraction as per manufacturer's instructions. Reverse transcription was performed using

ProtoScript® II First Strand cDNA Synthesis Kit (New England Biolabs) as per manufacturer's

instructions. PCR of a spike gene fragment was performed using Q5 High-Fidelity 2X Master

Mix (New England Biolabs) and the following primers: Forward 5' –

TTGAACTTCTACATGCACCAGC – 3' and Reverse 5' – CCAGAAGTGATTGTACCCGC –

3'. The fragment was gel purified using Monarch DNA Gel Extraction Kit (New England

Biolabs), and Sanger sequencing was performed using BigDye Terminator v3.1 and either the

forward or the reverse primer. The virus was confirmed as the Omicron variant due to

containing the following amino acid changes compared to SARS-CoV-2$_{QLD02}$; N547K, D614G,

H655Y, N679K, P681H, N764K, D796Y, N856K.

CCID$_{50}$ assays

Vero E6 cells were plated into 96 well flat bottom plates at 2×10^4 cells per well in 100 µl of

medium. For tissue titer, tissue was homogenized in tubes each containing 4 ceramic beads

twice at 6000 x g for 15 seconds, followed by centrifugation twice at 21000 x g for 5 min before

5 fold serial dilutions in 100 µl RPMI1640 supplemented with 2% FBS. For cell culture

supernatant, 10 fold serial dilutions were performed in 100 µl RPMI1640 supplemented with 2%

FBS. 100 µl of serially diluted samples were added to Vero E6 cells and the plates cultured for 5

days at 37°C and 5% CO_2. The virus titer was determined by the method of Spearman and

Karber.

SARS-CoV-2 passaging in hACE2 and mACE2 co-cultures

Lentivirus encoding hACE2, mACE2 or mACE2$^{N31K/H353K}$ were produced in HEK293T cells by

plasmid transfection and were used to transduce HEK293T cells, as described previously[2].

HEK293T-hACE2 or HEK293T-mACE2$^{N31K/H353K}$ were mixed with equal cell numbers of

HEK293T-mACE2, and 500,000 cells (i.e. 250,000 HEK293T-hACE2 or mACE2$^{N31K/H353K}$ +

250,000 HEK293T-mACE2) were seeded in 6 well plates overnight. Co-cultures were infected

with SARS-CoV-2 at MOI≈0.1. Every 3-4 days, supernatant was collected and centrifuged at

2000 x g at 4°C for 5 min, and 200 µl was added to fresh co-cultures and the remaining was

stored at -80°C. This was performed a total of 9 times before 1 ml was passaged on to

HEK293T-mACE2 cells. After one passage in HEK293T-mACE2 cells, supernatant was used to

infect new HEK293T-mACE2 cells for 2 hrs, then inoculum was removed and cells were washed

3 times with PBS and media replaced. Supernatant was harvested at 0 hr and 72 hrs post

infection for virus titration by CCID$_{50}$ to confirm virus replication in mACE2 expressing cells.

Virus was then passaged one additional time in HEK293T-mACE2 cells and supernatant stored

at -80°C for use in sequencing and mouse infections.

For growth kinetics experiments, HEK293T, HEK293T-hACE2 and HEK293T-mACE2,

NIH-3T3, AE17, BHK-21, A549, HeLa or LLC-PK1 cells were infected with SARS-CoV-2

(QLD02, MA1, MA2, Alpha, Beta or Omicron) at MOI≈0.1 for 1 hr at 37°C, cells were washed

with PBS and media replaced. Culture supernatant was harvested at the indicated time points

and titered by CCID$_{50}$ assay as described above.

Mouse intrapulmonary SARS-CoV-2 infection

Female C57BL/6J mice (~ 6 months old at the time of infection) were purchased from Animal

Resources Centre (Canning Vale, WA, Australia). The conditions the mice were kept are as

follows: light = 12:12 h dark/light cycle, 7:45 a.m. sunrise and 7:45 p.m. sunset, 15 min light

dark and dark light ramping time. Enclosures: M.I.C.E cage (Animal Care Systems, Colorado,

USA). Ventilation: 100% fresh air, eight complete air exchange/h/rooms. In-house enrichment:

paper cups (Impact-Australia); tissue paper, cardboard rolls. Bedding: PuraChips (Able

scientific) (aspen fine). Food: Double bagged norco rat and mouse pellet (AIRR, Darra, QLD).

Water: deionized water acidified with HCl (pH = 3.2). Mice were anesthetized using isoflurane

and given an intrapulmonary inoculation $\approx 5 \times 10^4$ $CCID_{50}$ SARS-CoV-2 delivered via the

intranasal route in 50 µl. Mice were sacrificed by cervical dislocation at day 2, 4 or 7 and lungs,

nasal turbinates, brain, small intestine, colon, liver, kidney and spleen were collected. Right lung

and all other organs were immediately homogenized in tubes each containing 4 beads twice at

6000 x g for 15 seconds, and used in tissue titration as described above. Left lungs were fixed in

10% formalin for histology.

K18-hACE2 mice (strain B6.Cg-Tg(K18-ACE2)2Prlmn/J, JAX Stock No: 034860)[60]

were purchased from The Jackson Laboratory, USA, and bred and maintained in-house at

QIMRB as heterozygotes by crossing with C57BL/6J mice[10]. Mice were genotyped using

Extract-N-Amp Tissue PCR Kit (Sigma Aldrich) according to manufacturers' instructions with

the following primers; Forward 5'-CTTGGTGATATGTGGGGTAGA-3' and Reverse 5'-

CGCTTCATCTCCCACCACTT-3' (recommended by NIOBIOHN, Osaka, Japan).

Thermocycling conditions were as follows; 94°C 3 min, 35 cycles of 94°C 30 s, 55.8 °C 30 s,

72°C 1 min, and final extension of 72°C 10 min.

RNA sequencing

For viral RNA purification, cell culture supernatants were processed using the NucleoSpin RNA

Virus kit (Machery-Nagel) as per manufacturers' instructions. For mouse lung RNA isolation,

lungs were transferred from RNAlater to TRIzol (Life Technologies) and were homogenized

twice at 6000 x g for 15 sec. Homogenates were centrifuged at 14,000 × g for 10 min and RNA

was isolated as per manufacturers' instructions.

RNA concentration and quality was measured using TapeStation D1K TapeScreen assay

(Agilent). cDNA libraries were prepared using the Illumina TruSeq Stranded mRNA library

prep kit and the sequencing performed on the Illumina Nextseq 550 platform generating 75 bp

paired end reads.

For virus sequences, per base sequence quality for >90% bases was above Q30 for all

samples. The quality of raw sequencing reads was assessed using FastQC[61] (v0.11.80), and

trimmed using Cutadapt[62] (v2.3) to remove adapter sequences and low-quality bases. Trimmed

reads were aligned using STAR[63] (v2.7.1a) to a SARS-CoV-2 isolate Wuhan-Hu-1

(NC_045512.2; 29903 bp). Aligned reads were viewed using Integrative Genome Viewer

(IGV)[64], and any position with >20% change compared to the reference genome was manually

curated. SAMtools mpileup was used to produce a consensus sequence from mapped reads[65].

For mouse lungs, per base sequence quality for >90% bases was above Q30 for all

samples. The quality of raw sequencing reads was assessed using FastQC[61] (v0.11.80), and

trimmed using Cutadapt[62] (v2.3) to remove adapter sequences and low-quality bases. Trimmed

reads were aligned using STAR[63] (v2.7.1a) to a combined reference that included the mouse

GRCm39 primary assembly and the GENCODE M27 gene model[66], SARS-CoV-2 isolate

Wuhan-Hu-1 (NC_045512.2; 29903 bp). Mouse gene expression was estimated using RSEM[67]

(v1.3.0). Reads aligned to SARS-CoV-2 were counted using SAMtools[65] (v1.9). Differential

gene expression in the mouse was analyzed using EdgeR (3.22.3) and modelled using the quasi-

likelihood F-test, glmQLFTest().

Pathway analysis

431 Up-Stream Regulators (USRs) and Diseases and Functions enriched in differentially expressed

432 genes in direct and indirect interactions were investigated using Ingenuity Pathway Analysis

433 (IPA) (QIAGEN).

SARS-CoV-2 amino acid change analyses and modelling

The GISAID (https://www.gisaid.org/phylodynamics/global/nextstrain/)[68] EpiCoV 'search'

function (https://www.epicov.org/epi3/frontend#577a0f) was used to identify the number of

GISAID SARS-CoV-2 sequence submissions containing each amino acid change in the MA

viruses. Data was filtered for submissions that selected "Human" as the host. Data was accessed

on 21st February 2021 when the number of total submissions was 8,601,773, and this was used to

calculate the proportion of each amino acid change in the MA viruses among total GISAID

sequence submissions.

PyMOL v4.60 (Schrodinger) was used for mutagenesis of the crystal structure of SARS-

CoV-2 spike bound with ACE2 from the protein data bank (7DF4)[69].

Lung histopathology and immunohistochemistry

Lungs and trachea (via pluck necropsy. i.e. removal of tongue, larynx, trachea, lungs, heart, and

part of the esophagus in one piece) were fixed in 10% formalin, embedded in paraffin, and

sections stained with H&E (Sigma Aldrich). Slides were scanned using Aperio AT Turbo

(Aperio, Vista, CA USA) and analyzed using Aperio ImageScope software (LeicaBiosystems,

Mt Waverley, Australia) (v10) and the Positive Pixel Count v9 algorithm. Automatic

quantitation of white space was undertaken using QuPath v0.2.3[70]. Immunohistochemistry for

SARS-CoV-2 antigen was undertaken using mouse anti-SARS-CoV-2 spike monoclonal

antibody 1E8 (Hobson-Peters *et al*. in preparation) as described previously[2].

Statistics

Statistical analyses of experimental data were performed using IBM SPSS Statistics for

Windows, Version 19.0 (IBM Corp., Armonk, NY, USA). The t-test was used when the

difference in variances was <4, skewness was > -2 and kurtosis was <2. Otherwise, the non-

parametric Kolmogorov-Smirnov test was used.

REFERENCES

1. Wu, F. *et al.* A new coronavirus associated with human respiratory disease in China. *Nature* **579**, 265-269, doi:10.1038/s41586-020-2008-3 (2020).

2. Rawle, D. J. *et al.* ACE2-lentiviral transduction enables mouse SARS-CoV-2 infection and mapping of receptor interactions. *PLOS Pathogens* **17**, e1009723, doi:10.1371/journal.ppat.1009723 (2021).

3. Starr, T. N. *et al.* ACE2 binding is an ancestral and evolvable trait of sarbecoviruses. *Nature*, https://doi.org/10.1038/s41586-41022-04464-z, doi:10.1101/2021.07.17.452804 (2022).

4. Hu, B. *et al.* Discovery of a rich gene pool of bat SARS-related coronaviruses provides new insights into the origin of SARS coronavirus. *PLOS Pathogens* **13**, e1006698, doi:10.1371/journal.ppat.1006698 (2017).

5. Cele, S. *et al.* SARS-CoV-2 Omicron has extensive but incomplete escape of Pfizer BNT162b2 elicited neutralization and requires ACE2 for infection. *medRxiv*, 2021.2012.2008.21267417, doi:10.1101/2021.12.08.21267417 (2021).

6. Shuai, H. *et al.* Emerging SARS-CoV-2 variants expand species tropism to murines. *EBioMedicine* **73**, 103643, doi:10.1016/j.ebiom.2021.103643 (2021).

7. Muñoz-Fontela, C. *et al.* Animal models for COVID-19. *Nature* **586**, 509-515, doi:10.1038/s41586-020-2787-6 (2020).

8. Shou, S. *et al.* Animal Models for COVID-19: Hamsters, Mouse, Ferret, Mink, Tree Shrew, and Non-human Primates. *Frontiers in Microbiology* **12**, 626553, doi:https://doi.org/10.3389/fmicb.2021.626553 (2021).

9. Amarilla, A. A. *et al.* A versatile reverse genetics platform for SARS-CoV-2 and other positive-strand RNA viruses. *Nat Commun* **12**, 3431, doi:10.1038/s41467-021-23779-5 (2021).

10. Bishop, C. R. *et al.* Mouse models of COVID-19 recapitulate inflammatory pathways rather than gene expression. *bioRxiv*, 2022.2002.2024.481866, doi:10.1101/2022.02.24.481866 (2022).

11. Halfmann, P. J. *et al.* SARS-CoV-2 Omicron virus causes attenuated disease in mice and hamsters. *Nature* **603**, 687–692, doi:10.1038/s41586-022-04441-6 (2022).

12. Leist, S. R. *et al.* A Mouse-Adapted SARS-CoV-2 Induces Acute Lung Injury and Mortality in Standard Laboratory Mice. *Cell* **183**, 1070-1085.e1012, doi:https://doi.org/10.1016/j.cell.2020.09.050 (2020).

13. Zhang, Y. *et al.* SARS-CoV-2 Rapidly Adapts in Aged BALB/c Mice and Induces Typical Pneumonia. *Journal of Virology* **95**, e02477-02420, doi:10.1128/JVI.02477-20 (2021).

14. Huang, K. *et al.* Q493K and Q498H substitutions in Spike promote adaptation of SARS-CoV-2 in mice. *EBioMedicine* **67**, 103381, doi:https://doi.org/10.1016/j.ebiom.2021.103381 (2021).

15. Gu, H. *et al.* Adaptation of SARS-CoV-2 in BALB/c mice for testing vaccine efficacy. *Science* **369**, 1603, doi:10.1126/science.abc4730 (2020).

16. Wang, J. *et al.* Mouse-adapted SARS-CoV-2 replicates efficiently in the upper and lower respiratory tract of BALB/c and C57BL/6J mice. *Protein Cell* **11**, 776-782, doi:10.1007/s13238-020-00767-x (2020).

17. Dinnon, K. H. *et al.* A mouse-adapted model of SARS-CoV-2 to test COVID-19 countermeasures. *Nature* **586**, 560-566, doi:10.1038/s41586-020-2708-8 (2020).

18. Palermo, P. M., Orbegozo, J., Watts, D. M. & Morrill, J. C. SARS-CoV-2 Neutralizing Antibodies in White-Tailed Deer from Texas. *Vector Borne Zoonotic Dis* **22**, 62-64, doi:http://doi.org/10.1089/vbz.2021.0094 (2021).

19. Diaz, A. V., Walker, M. & Webster, J. P. Surveillance and control of SARS-CoV-2 in mustelids: An evolutionary perspective. *Evol Appl*, 2715-2725, doi:10.1111/eva.13310 (2021).

20 Langereis, M. A. *et al.* An alphavirus replicon-based vaccine expressing a stabilized Spike antigen
induces protective immunity and prevents transmission of SARS-CoV-2 between cats. *npj*
Vaccines **6**, 122, doi:10.1038/s41541-021-00390-9 (2021).
21 Welkers, M. R. A., Han, A. X., Reusken, C. B. E. M. & Eggink, D. Possible host-adaptation of SARS-
CoV-2 due to improved ACE2 receptor binding in mink. *Virus Evolution* **7**, veaa094,
doi:10.1093/ve/veaa094 (2021).
22 Chandler, J. C. *et al.* SARS-CoV-2 exposure in wild white-tailed deer (Odocoileus
virginianus). *Proceedings of the National Academy of Sciences* **118**, e2114828118,
doi:10.1073/pnas.2114828118 (2021).
23 Griffin, B. D. *et al.* SARS-CoV-2 infection and transmission in the North American deer mouse.
Nat Commun **12**, 3612, doi:10.1038/s41467-021-23848-9 (2021).
24 Mallapaty, S. Where did Omicron come from? Three key theories. The hunt for the origins of
omicron. . *Nature* **602**, 26-28 (2022).
25 Wei, C. *et al.* Evidence for a mouse origin of the SARS-CoV-2 Omicron variant. *J Genet Genomics*
48, 1111-1121, doi:https://doi.org/10.1016/j.jgg.2021.12.003 (2021).
26 Puray-Chavez, M. *et al.* Systematic analysis of SARS-CoV-2 infection of an ACE2-negative human
airway cell. *Cell Reports* **36**, 109364, doi:https://doi.org/10.1016/j.celrep.2021.109364 (2021).
27 Cantuti-Castelvetri, L. *et al.* Neuropilin-1 facilitates SARS-CoV-2 cell entry and infectivity. *Science*
370, 856-860, doi:10.1126/science.abd2985 (2020).
28 Wang, S. *et al.* AXL is a candidate receptor for SARS-CoV-2 that promotes infection of pulmonary
and bronchial epithelial cells. *Cell Res* **31**, 126-140, doi:10.1038/s41422-020-00460-y (2021).
29 Mori, Y. *et al.* KIM-1/TIM-1 is a Receptor for SARS-CoV-2 in Lung and Kidney. *medRxiv*,
2020.2009.2016.20190694, doi:10.1101/2020.09.16.20190694 (2022).
30 Hoffmann, M. *et al.* SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a
Clinically Proven Protease Inhibitor. *Cell* **181**, 271-280.e278, doi:10.1016/j.cell.2020.02.052
(2020).
31 Chu, H. *et al.* Comparative tropism, replication kinetics, and cell damage profiling of SARS-CoV-2
and SARS-CoV with implications for clinical manifestations, transmissibility, and laboratory
studies of COVID-19: an observational study. *The Lancet Microbe* **1**, e14-e23,
doi:https://doi.org/10.1016/S2666-5247(20)30004-5 (2020).
32 Trimarco, J. D. *et al.* TMEM41B is a host factor required for the replication of diverse
coronaviruses including SARS-CoV-2. *PLoS Pathog* **17**, e1009599,
doi:10.1371/journal.ppat.1009599 (2021).
33 Laporte, M. *et al.* The SARS-CoV-2 and other human coronavirus spike proteins are fine-tuned
towards temperature and proteases of the human airways. *PLOS Pathogens* **17**, e1009500,
doi:10.1371/journal.ppat.1009500 (2021).
34 DiPiazza, A. T. *et al.* COVID-19 vaccine mRNA-1273 elicits a protective immune profile in mice
that is not associated with vaccine-enhanced disease upon SARS-CoV-2 challenge. *Immunity* **54**,
1869-1882.e1866, doi:https://doi.org/10.1016/j.immuni.2021.06.018 (2021).
35 Winkler Emma, S. *et al.* SARS-CoV-2 causes lung infection without severe disease in human ACE2
knock-in mice. *Journal of Virology* **96**, e0151121, doi:10.1128/JVI.01511-21 (2021).
36 Hui, K. P. Y. *et al.* SARS-CoV-2 Omicron variant replication in human bronchus and lung ex vivo.
Nature, doi:10.1038/s41586-022-04479-6 (2022).
37 Peacock, T. P. *et al.* The SARS-CoV-2 variant, Omicron, shows rapid replication in human primary
nasal epithelial cultures and efficiently uses the endosomal route of entry. *bioRxiv*,
2021.2012.2031.474653, doi:10.1101/2021.12.31.474653 (2022).
38 McMahan, K. *et al.* Reduced Pathogenicity of the SARS-CoV-2 Omicron Variant in Hamsters.
bioRxiv, 2022.2001.2002.474743, doi:10.1101/2022.01.02.474743 (2022).

39 Bao, L. *et al.* Transmission of Severe Acute Respiratory Syndrome Coronavirus 2 via Close
Contact and Respiratory Droplets Among Human Angiotensin-Converting Enzyme 2 Mice. *J*
Infect Dis **222**, 551-555, doi:10.1093/infdis/jiaa281 (2020).
40 Peacock, T. P. *et al.* The furin cleavage site in the SARS-CoV-2 spike protein is required for
transmission in ferrets. *Nature Microbiology* **6**, 899-909, doi:10.1038/s41564-021-00908-w
(2021).
41 Johnson, B. A. *et al.* Loss of furin cleavage site attenuates SARS-CoV-2 pathogenesis. *Nature* **591**,
293-299, doi:10.1038/s41586-021-03237-4 (2021).
42 Liu, Z. *et al.* Identification of Common Deletions in the Spike Protein of Severe Acute Respiratory
Syndrome Coronavirus 2. *Journal of Virology* **94**, e00790-00720, doi:10.1128/JVI.00790-20
(2020).
43 Prow, N. A. *et al.* Exacerbation of Chikungunya Virus Rheumatic Immunopathology by a High
Fiber Diet and Butyrate. *Front Immunol* **10**, 2736-2736, doi:10.3389/fimmu.2019.02736 (2019).
44 Mills, R. J. *et al.* BET inhibition blocks inflammation-induced cardiac dysfunction and SARS-CoV-2
infection. *Cell* **184**, 2167-2182.e2122, doi:https://doi.org/10.1016/j.cell.2021.03.026 (2021).
45 MacDonald, L. *et al.* COVID-19 and RA share an SPP1 myeloid pathway that drives PD-L1+
neutrophils and CD14+ monocytes. *JCI Insight* **6**, e147413, doi:10.1172/jci.insight.147413 (2021).
46 Xu, M., Yang, W., Wang, X. & Nayak, D. K. Lung Secretoglobin Scgb1a1 Influences Alveolar
Macrophage-Mediated Inflammation and Immunity. *Front Immunol* **11**, 584310,
doi:10.3389/fimmu.2020.584310 (2020).
47 Ramirez, S. *et al.* Overcoming Culture Restriction for SARS-CoV-2 in Human Cells Facilitates the
Screening of Compounds Inhibiting Viral Replication. *Antimicrob Agents Chemother* **65**,
e0009721, doi:10.1128/AAC.00097-21 (2021).
48 Jary, A. *et al.* Spike Gene Evolution and Immune Escape Mutations in Patients with Mild or
Moderate Forms of COVID-19 and Treated with Monoclonal Antibodies Therapies. *Viruses* **14**,
226, doi:10.3390/v14020226 (2022).
49 Amraei, R. *et al.* CD209L/L-SIGN and CD209/DC-SIGN Act as Receptors for SARS-CoV-2. *ACS Cent*
Sci **7**, 1156-1165, doi:10.1021/acscentsci.0c01537 (2021).
50 Wang, K. *et al.* CD147-spike protein is a novel route for SARS-CoV-2 infection to host cells. *Signal*
Transduction and Targeted Therapy **5**, 283, doi:10.1038/s41392-020-00426-x (2020).
51 Bestle, D. *et al.* TMPRSS2 and furin are both essential for proteolytic activation of SARS-CoV-2 in
human airway cells. *Life Science Alliance* **3**, e202000786, doi:10.26508/lsa.202000786 (2020).
52 Meng, B. *et al.* Altered TMPRSS2 usage by SARS-CoV-2 Omicron impacts tropism and
fusogenicity. *Nature* **603**, 706-714, doi:10.1038/s41586-022-04474-x (2022).
53 Shuai, H. *et al.* Attenuated replication and pathogenicity of SARS-CoV-2 B.1.1.529 Omicron.
Nature, 693-699, doi:10.1038/s41586-022-04442-5 (2022).
54 Wei, C. *et al.* Evidence for a mouse origin of the SARS-CoV-2 Omicron variant. *Journal of*
Genetics and Genomics **48**, 1111-1121, doi:https://doi.org/10.1016/j.jgg.2021.12.003 (2021).
55 Cameroni, E. *et al.* Broadly neutralizing antibodies overcome SARS-CoV-2 Omicron antigenic
shift. *bioRxiv*, 2021.2012.2012.472269, doi:10.1101/2021.12.12.472269 (2021).
56 Lopez-Leon, S. *et al.* More than 50 long-term effects of COVID-19: a systematic review and meta-
analysis. *Scientific reports* **11**, 16144-16144, doi:10.1038/s41598-021-95565-8 (2021).
57 Mehandru, S. & Merad, M. Pathological sequelae of long-haul COVID. *Nat Immunol* **23**, 194-202,
doi:10.1038/s41590-021-01104-y (2022).
58 Johnson, B. J. *et al.* Heat shock protein 10 inhibits lipopolysaccharide-induced inflammatory
mediator production. *J Biol Chem* **280**, 4037-4047, doi:10.1074/jbc.M411569200 (2005).

59   La Linn, M., Bellett, A. J., Parsons, P. G. & Suhrbier, A. Complete removal of mycoplasma from
viral preparations using solvent extraction. *J Virol Methods* **52**, 51-54, doi:10.1016/0166-
0934(94)00136-5 (1995).
60   McCray, P. B., Jr. *et al.* Lethal infection of K18-hACE2 mice infected with severe acute respiratory
syndrome coronavirus. *J Virol* **81**, 813-821, doi:10.1128/jvi.02012-06 (2007).
61   Andrews, S. A quality control tool for high throughput sequence data. (2010).
62   Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet.journal; Vol 17, No 1: Next Generation Sequencing Data AnalysisDO -
10.14806/ej.17.1.200 (2011).
63   Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21,
doi:10.1093/bioinformatics/bts635 (2013).
64   Robinson, J. T. *et al.* Integrative genomics viewer. *Nat Biotechnol* **29**, 24-26,
doi:10.1038/nbt.1754 (2011).
65   Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078-2079,
doi:10.1093/bioinformatics/btp352 (2009).
66   Harrow, J. *et al.* GENCODE: the reference human genome annotation for The ENCODE Project.
Genome Res **22**, 1760-1774, doi:10.1101/gr.135350.111 (2012).
67   Li, B. & Dewey, C. N. RSEM: accurate transcript quantification from RNA-Seq data with or
without a reference genome. *BMC Bioinformatics* **12**, 323, doi:10.1186/1471-2105-12-323
(2011).
68   Elbe, S. & Buckland-Merrett, G. Data, disease and diplomacy: GISAID's innovative contribution to
global health. *Glob Chall* **1**, 33-46, doi:10.1002/gch2.1018 (2017).
69   Xu, C. *et al.* Conformational dynamics of SARS-CoV-2 trimeric spike glycoprotein in complex with
receptor ACE2 revealed by cryo-EM. *Sci Adv* **7**, eabe5575, doi:10.1126/sciadv.abe5575 (2021).
70   Bankhead, P. *et al.* QuPath: Open source software for digital pathology image analysis. *Scientific*
Reports **7**, 16878, doi:10.1038/s41598-017-17204-5 (2017).

Figures

Figure 1. *In vitro* adaptation of SARS-CoV-2$_{QLD02}$ to mACE2 utilization. a Schematic of SARS-CoV-2$_{QLD02}$ passaging in HEK293T-hACE2 or HEK293T-mACE2$^{N31K/H353K}$ cells, co-cultured with HEK293T-mACE2 cells (9 passages), followed by passaging in HEK293T-mACE2 cells (3 passages) and mouse infections. Viruses used to infect mice and viruses derived from mice lungs (2 dpi) were sequenced. Stock virus was prepared for MA1 and MA2 from mouse lungs 2 dpi for use in subsequent experiments. **b** Supernatants from passage 4 and 9 (from the co-cultures – blue arrows) were used to infect HEK93T-mACE2 cells and viral growth over 72 hours determined by CCID$_{50}$ assays. Dotted line – limit of detection. MA1 and MA3 (underlined) were derived from HEK293T-hACE2/HEK293T-mACE2 co-cultures, MA2, 4, 5 and 6 from HEK293T-mACE2$^{N31K/H353K}$/HEK293T-mACE2 co-cultures. **c** Inverted light microscopy images of CPE in HEK293T-mACE2 cells infected with passage 9 supernatants at 72 hours post-infection. Images are representative of at least three replicates. **d** MA viruses obtained after 3 passages in HEK293T-mACE2 cells were used to infect C57BL/6J mice. Lung tissue titers are shown for 2 dpi for MA1-5 (n=1 for each) and for SARS-CoV-2$_{QLD02}$ (n=6).

Figure 2. Sequencing of MA viruses. a Amino acid changes and deletions (Δ) in MA1, MA2, MA3, MA4 and MA5 (full dataset in Supplementary File 1). Amino acid changes that only appear in one MA virus are in red, with other colors used to show amino acid changes common

between at least two MA viruses. **b** Amino acids changes and deletions in any of the MA viruses

shown on the structure of SARS-CoV-2$_{QLD02}$ spike bound to hACE2 (PDB: 7DF4). Underlining

indicates non-conservative amino changes. **c** Spike changes in the MA viruses. * - amino acid

changes located within the RBD. Underlining indicates non-conservative amino acid changes.

GISAID n – number of GISAID submissions that contain this change; GISAID % - percentage

of all GISAID submissions with this change. Alpha, Beta, Gamma, Delta, Omicron: black text -

hallmark changes in variants of concern; grey text –not a hallmark change in variants of concern.

Match of amino acid changes in MA viruses with hallmark changes in variants of concern (green

- exact match, blue - conservative, brown - non-conservative change).

Figure 3. *In vitro* growth kinetics of mACE2-adapted viruses reveal an ACE2-independent entry mechanism. Growth kinetics of the indicated viruses in HEK293T-hACE2 cells **(a)**, HEK293T-mACE2 cells **(b)**, untransduced HEK293T cells **(c)**, 3T3 cells **(d)**, AE17 cells **(e)**, BHK-21 cells **(f)**, A549 cells **(g)**, HeLa cells **(h)** and LLC-PK1 **(i)** cells after infection at MOI = 0.1. n = 3-6 replicates per virus strain per cell line. Dotted line – limit of detection. * p<0.05; statistics by Kolmogorov–Smirnov test for 3 dpi versus 0 dpi.

Figure 4. MA1 and MA2 infection in C57BL/6J mice. a C57BL/6J mice were infected with
MA1 or MA2 and lungs collected at day 2, 4 or 7 post-infection and tissue titers determined by
$CCID_{50}$ assays. **b** As for (**a**) for nasal turbinates. Statistics by t-test. **c** As for (**a**) for the
indicated tissues. **d-g** IHC using an anti-SARS-CoV-2 spike monoclonal antibody and lungs
taken on day 2 after infection of C57BL/6J mice with MA1. Images are representative of lung
sections from 4 mice. Dark brown staining (Nova Red) indicates infected cells, often with the
expected, clearly discernable, cytoplasmic staining pattern. The large unstained areas in (**d**) and
(**g**) are included to illustrate the specificity of the IHC staining.

702
703 **Figure 5. H&E staining of lungs after infection of C57BL/6J mice with MA1 and MA2. a**
704 Representative low magnification images of H&E stained lung sections for uninfected and MA1
705 infected C57BL/6J mice taken 4 days post infection. **b** Image analysis to measure lung
706 consolidation quantitated as area of white space (unstained air spaces within the lung) per μm^2
707 (n=4 mice per group, with 3 sections scanned per lung and values averaged to produce one value

for each lung). Statistics by t-test. **c** Image analysis to quantitate leukocyte infiltration. The ratios of nuclear (blue/dark purple) to cytoplasmic (red) staining of H&E stained lung n=4 mice per group, with 3 sections scanned per lung and values averaged to produce one value for each lung. Statistics by t-tests, with data from the 3 time points for infected mice combined. **d** High magnification image showing bronchus from uninfected mice (left) and MA1 infected mice (right) on day 4 post infection. The latter shows sloughing of the bronchial epithelium (S) and smooth muscle hyperplasia (M). **e** High magnification image showing uninfected lungs (top) and MA1 infected lungs (4 dpi) (bottom). The latter shows lung consolidation and loss of alveolar air spaces, as well as pulmonary edema (dotted ovals, bottom right). **f** High magnification images of uninfected lungs (top) and in MA1-infected lungs on 4 dpi (bottom). The latter shows dense cellular infiltrate (dotted circle).

Figure 6. RNA-Seq comparison of MA1 infected C57BL/6J versus SARS-CoV-2$_{QLD02}$
infected K18-hACE2 mouse lungs. RNA-Seq was performed on mouse lungs 4 dpi for MA1
infection of C57BL/6J mice, and SARS-CoV-2$_{QLD02}$ infection of K18-hACE2 mice. DEGs (n =
1027 for C57BL/6J, n = 1349 for K18-hACE2) were analyzed by Ingenuity Pathway Analysis,
and z-scores for significant UpStream Regulators **(a)** or Diseases and Functions **(b)** were plotted
with C57BL/6J+MA1 on x-axis and K18-hACE2+QLD02 on the y-axis. Statistics by Pearson
correlations. Dotted black lines - 95% confidence limits. Solid black line - line of best fit.
Annotations outside the 95% confidence limits are colored in red. Selected annotations are
labelled (red text). Full data sets are provided in Supplemental File 2.