Resistance to miltefosine results from amplification of the *RTA3* floppase or inactivation of flippases in *Candida parapsilosis*

Sean A. Bergin[1*], Fang Zhao[1*], Adam P. Ryan[1], Carolin A. Müller[2], Conrad A. Nieduszynski[3], Bing Zhai[4,5,6], Thierry Rolling[4,5], Tobias M. Hohl[4,5,7], Florent Morio[8,9], Jillian Scully[1], Kenneth H. Wolfe[10] and Geraldine Butler[1].

* Equal contribution

[1] School of Biomolecular and Biomedical Science, Conway Institute, University College Dublin, Belfield, Dublin 4, Ireland

[2] Sir William Dunn School of Pathology, University of Oxford, Oxford, UK

[3] Earlham Institute, Norwich Research Park, Norwich NR4 7UZ, UK

[4] Infectious Disease Service, Department of Medicine, Memorial Sloan Kettering Cancer Center, New York, NY, USA

[5] Immunology Program, Sloan Kettering Institute, Memorial Sloan Kettering Cancer Center, New York, NY, USA

[6] Laboratory of Quantitative Engineering Biology, Shenzhen Institute of Synthetic Biology, Shenzhen Institutes of Advanced Technology, Chinese Academy of Sciences, Shenzhen, China

[7] Department of Medicine, Weill Cornell Medical College, New York, NY, USA

[8] Laboratoire de Parasitologie et Mycologie, CHU de Nantes, Nantes, France

[9] Cibles et Médicaments des Infections et du Cancer, IICiMed UPRES EA 1155, UFR de Sciences Pharmaceutiques et Biologiques, 1 rue Gaston Veil, 44035 Nantes, France

[10] School of Medicine, Conway Institute, University College Dublin, Belfield, Dublin 4, Ireland.

Running title: Miltefosine resistance in *Candida parapsilosis*

corresponding author: Geraldine Butler, School of Biomolecular and Biomedical Science, Conway Institute, University College Dublin, Belfield, Dublin 4, Ireland

email: gbutler@ucd.ie

Abstract

Flippases and floppases are two classes of proteins that have opposing functions in the
maintenance of lipid asymmetry of the plasma membrane. Flippases translocate lipids from
the exoplasmic leaflet to the cytosolic leaflet, and floppases act in the opposite direction.
Phosphatidylcholine (PC) is a major component of the eukaryotic plasma membrane and is
asymmetrically distributed, being more abundant in the exoplasmic leaflet. Here we show
that gene amplification of a putative PC floppase or double disruption of two PC flippases in
the pathogenic yeast *Candida parapsilosis* results in resistance to miltefosine, an
alkylphosphocholine drug that affects PC metabolism that has recently been granted orphan
drug designation approval by the US FDA for treatment of invasive candidiasis. We analysed
the genomes of 170 *C. parapsilosis* isolates and found that 107 of them have copy number
variations (CNVs) at the *RTA3* gene. *RTA3* encodes a putative PC floppase whose deletion
is known to increase the inward translocation of PC in *Candida albicans*. *RTA3* copy number
ranges from 2 to >40 across the *C. parapsilosis* isolates. Interestingly, 16 distinct CNVs with
unique endpoints were identified, and phylogenetic analysis shows that almost all of them
have originated only once. We found that increased copy number of *RTA3* correlates with
miltefosine resistance. Additionally, we conducted an adaptive laboratory evolution
experiment in which two *C. parapsilosis* isolates were cultured in increasing concentrations
of miltefosine over 26 days. Two genes, *CPAR2_303950* and *CPAR2_102700*, gained
homozygous protein-disrupting mutations in the evolved strains and code for putative PC
flippases homologous to *S. cerevisiae DNF1.* Our results indicate that alteration of lipid
asymmetry across the plasma membrane is a key mechanism of miltefosine resistance. We
also find that *C. parapsilosis* is likely to gain resistance to miltefosine rapidly, because many
isolates carry loss-of-function alleles in one of the flippase genes.

Author summary

Miltefosine was developed as an anticancer drug but is commonly used to treat infections
with the protozoan parasites *Leishmania* and *Trypanosoma cruzi*. More recently, it has been
used to treat fungal infections, and in 2021 it was designated as an orphan drug by the US
Food and Drug Administration for treatment of invasive candidiasis. Miltefosine is a
derivative of phosphatidylcholine (PC), a major constituent of the cell membrane. PC and
other phospholipids are asymmetrically distributed across the cell membrane. The
mechanism of action of miltefosine is unknown. Here, we show that either increasing the
activity of a putative floppase, which controls outward "flop" movement of phospholipids, or
decreasing the activity of flippases, which control inward "flip" movement, results in

increased resistance of the fungal pathogen *Candida parapsilosis* to miltefosine. This result

suggests that miltefosine acts by controlling the localisation of PC or other phospholipids in

the membrane. Importantly, we find that many *C. parapsilosis* isolates carry mutations in one

flippase gene, which renders them partially resistant to miltefosine, and prone to easily

acquiring increased resistance.

Introduction

Lipids in yeast plasma membranes belong to three main families: sterols, sphingolipids, and phospholipids (or glycerophospholipids) [1]. Phospholipids (including phosphatidylcholine (PC), phosphatidylethanolamine (PE), phosphatidylinositol (PI), and phosphatidylserine (PS)) and sphingolipids (including the long chain bases (LCBs) dihydrosphingosine (DHS) and phytosphingosine (PHS)) are distributed asymmetrically in the plasma membrane bilayer [2, 3]. Plasma asymmetry is determined by the action of lipid translocators via rapid inward (flip) and outward (flop) movements [4]. Most flippases are members of the P4-ATPase family [5], and most floppases are ABC-type transporters [6]. The Rta1 family of 7 transmembrane domain proteins are also predicted lipid transporters [7].

Several antimicrobial drugs are targeted against lipid synthesis, such as azole drugs, which target ergosterol synthesis in fungi [8-10]. Miltefosine (hexadecylphosphocholine) is an alkylphosphocholine derivative that inhibits PC biosynthesis or localization [11]. Miltefosine was initially investigated as an antitumor agent, but it was found to have a very limited anticancer effect [12]. However, it has been used successfully to treat infections with *Leishmania* and *Trypanosoma cruz*i [13-15]. Recently, miltefosine has also been shown to have antifungal activity, including against *Aspergillus* [16], *Cryptococcus* [17] and *Paracoccidioides* [18]. Anti-*Candida* activity has also been reported, including against *Candida auris* [19, 20], *Candida krusei* [21] and *Candida albicans*, where it inhibits growth of both planktonic and biofilm cells [22]. In 2021, miltefosine was granted orphan drug designation approval by the US Food and Drug Administration (FDA) for treatment of invasive candidiasis (https://www.accessdata.fda.gov/scripts/opdlisting/oopd/detailedIndex.cfm?cfgridkey=843921).

The mechanism of action of miltefosine is not completely understood. In liver cells, miltefosine is proposed to reduce the biosynthesis of PC by inhibiting the activity of CTP:phosphocholine cytidylyltransferase (CT) and PE N-methyltransferase activity [23]. Treatment of *Leishmania* with miltefosine decreases the amount of PC and increases PE in the membrane, because of a reduction of intracellular choline [24]. In addition, cytochrome c activity in the mitochondria is inhibited, and the plasma membrane Ca^{2+} channel is disrupted [25]. Studies in *Saccharomyces cerevisiae* suggest that related drugs, such as edelfosine (lysophosphatidylcholine) also disrupt lipid rafts in the cell membrane, perhaps by directly interacting with sterols [26]. In *Aspergillus fumigatus*, miltefosine affects metabolism of lipids and fatty acids, as well as by increasing mitochondrial fragmentation [16].

Most flippases belonging to the P4-ATPase family, which consist of two subunits, a catalytic alpha-subunit with 10 transmembrane alpha-helices, and a beta-subunit with two transmembrane alpha-helices belonging to the Cdc50 family [5]. Flipping of PE, PS and PC is specifically associated with Class 3 P4-ATPases, represented by Dnf1 and Dnf2 in *S. cerevisiae* [27]. Disruption of *DNF1* or *DNF2*, or of *LEM3* (*CDC50*), results in miltefosine resistance in *S. cerevisiae* [28]. Similarly, mutations in P4-ATPases in *Leishmania* species confer resistance to miltefosine [29-32].

In *Candida* species, flippases have not yet been associated with miltefosine resistance. However, deletion of an unrelated gene, *RTA3*, results in increased miltefosine sensitivity in *C. albicans* [33]. Rta3 is a member of the Rta1/Rsb1-like family in *S. cerevisiae*, which encode putative transporters with seven transmembrane domains (TMs). ScRsb1 controls the localization of sphingoid bases in *S. cerevisiae*, including phytosphingosine and dihydrosphingosine [34, 35]. Rsb1 is localized to the membrane and is likely to act as a membrane transporter (a floppase) [7], or possibly as a regulator of transporters [36]. Overexpression of another family member, *ScRTA1*, correlates with increased resistance to 7-amino cholesterol, an inhibitor of 8-7 sterol isomerase and 14-sterol reductase [37]. The Rsb1 family members have not been fully characterised in *Candida* species. As well as showing that deleting *C. albicans RTA3* (*CaRTA3*) increases sensitivity to miltefosine, Srivastava et al [33] also showed that CaRta3 localizes to the plasma membrane, and that a fluorophore-labeled PC accumulates in the inner leaflet of the membrane in an *CaRTA3* deletion. They proposed that CaRta3 is either a floppase, or that it inhibits the activity of a flippase.

Some previous reports suggest that *CaRTA3* is also associated with resistance to azoles. Rogers and Barker [38, 39] found that transcription of *CaRTA3* is up-regulated in fluconazole-resistant isolates from AIDS patents, and Coste et al. [40] showed that Tac1p, a transcription factor regulating expression of the MDR transporters *CDR1* and *CDR2*, also regulates expression of *CaRTA3* via binding to the drug-response element. Whaley et al [41] reported that deleting *CaRTA3* reduced fluconazole resistance of a clinical isolate of *C. albicans*, and that overexpressing *CaRTA3* increased resistance. However, Srivastava et al. [33] showed that deleting *CaRTA3* in a reference isolate had no effect on fluconazole resistance. The role of Rta3 in azole resistance is therefore unclear.

In 2020, we found that the *RTA3* gene has undergone amplification in 23 clinical isolates of
Candida parapsilosis, resulting in increased copy number of 24-33x [42]. Variable copy
numbers of *RTA3* were also found by West et al [43] in four isolates. Here, by genome
sequencing, we report that *RTA3* amplification is very common in *C. parapsilosis*. We find 16
distinct amplifications of *RTA3* with unique endpoints, indicative of independent and parallel
amplification events. We also find a fusion to a related neighboring gene, *RTA2*. Copy
number of *RTA3* is correlated with resistance to miltefosine, but not to azoles. In addition, we
find that resistance to miltefosine induced by exposure to the drug is associated with loss-of-
function mutations in two P4-ATPase flippase genes.
123
124   **Results**
125   ***C. parapsilosis* phylogeny shows independent origins of *RTA3* CNVs**
126
We explored the relationships of 170 *Candida parapsilosis* isolates using a genome-wide
SNP alignment approach, generating the largest phylogeny of *C. parapsilosis* isolates to
date (Fig 1). Isolates were obtained from several locations, including some previously
published, and 127 that are sequenced and described here for the first time (S1 Table). Most
isolates originated from the Memorial Sloan Kettering Cancer in New York (designated by
MSK) and the Centre Hospitalier Universitaire de Nantes, France (designated by FM) (Fig
1).
134

Figure 1. SNP-based unrooted phylogeny of 170 *C. parapsilosis* isolates.

Isolates harboring a CNV at *RTA3* are marked with a thick gray bar and a letter (A-P) corresponding

to one of the 16 CNVs found. Isolates from Memorial Sloan Kettering Cancer Center (MSK) are

named in blue (the prefix MSK is omitted from these strain names), and isolates from CHU de Nantes

(FM) are named in red. Green dashed lines label each of five apparent clades. Fifty similar isolates in

Clade 1, and 19 similar isolates in Clade 4, were grouped and are shown as blue triangles. The

phylogeny was constructed by calling SNPs for each sample using the GATK HaplotypeCaller tool

[44]. Filtered heterozygous sites were resolved using 100 iterations of Random Repeated Haplotype

Sampling to provide haploid inputs for tree construction [45]. The tree was then constructed using

RAxML with the GTRGAMMA model of nucleotide substitution.

Isolates fall into five major clades, with almost half (84/170), belonging to clade I. Most clade

I isolates were isolated at MSK. Half (8/16) of the FM isolates are found in Clade 4. There

are 49 highly similar isolates in Clade 1, and 19 highly similar isolates in Clade 4, all from

patients at MSK. These may have recently originated from two single isolates, and they are

indicated with blue triangles in Fig 1. Some other isolates (e.g. MSK2233 and MSK2234 in

Clade 2) may also share a recent origin. However, overall there is little evidence for

geographical clustering. Each clade includes at least one isolate from both MSK and FM (Fig

1). Three isolates from a clinical setting in Kuwait [46] are each located in a different clade

(designated by Kw in Fig 1). Interestingly, an environmental sample isolated from Irish soil,

UCD321, groups with clinical isolates from both MSK and FM (in Clade 5). The diversity of

isolates obtained from the same clinical setting, and the close relationship between isolates

from different geographical settings, highlight the global nature of *C. parapsilosis* as a

human pathogen.

We previously showed that the *RTA3* gene, encoding a putative floppase, had undergone

extensive copy number amplification in 23 *C. parapsilosis* isolates [42]. This was also

observed in a small number of isolates by West et al [43]. We therefore characterized copy

number variants (CNVs) at the *RTA3* locus in all sequenced isolates. We identified

increased *RTA3* copy numbers in 104 of 170 isolates. Sixteen CNVs with unique endpoints

were observed (assigned letters A-P, Figs. 1, 2). Nine different CNVs were observed in

isolates from Clade 1, whereas only CNV-L is found in isolates in Clade 3, and there are no

CNVs in Clade 2 isolates (Fig 1). Most (15/16) of the CNVs have a single evolutionary origin,

and some are present in only a single isolate (Fig 1). However, CNV-A has three separate

origins on the tree (one in Clade 1, and two in Clade 5).

16 unique CNVs amplify *RTA3*

We found that *RTA3* has been amplified in 16 different types of CNV, each with unique

endpoints. Thirteen CNV patterns result from tandem duplications that amplify the entire

RTA3 coding sequence (Fig 2A), and these repeat units range in size from 2.3 to 4.5 kb.

Some of these CNVs extend into the upstream neighbouring gene *MAK16,* and one (CNV-N)

includes the first 17 bp in the coding sequence of the downstream gene *RTA2*. The copy

number of *RTA3* varies both in isolates that share the same CNV, and in isolates with

different CNVs (Fig 2B). Isolate MSK807 (CNV-G) has the lowest estimated copy number of

RTA3 among isolates with CNVs, at four copies, whereas isolate FM16 (CNV-J) has the

highest copy number, at 50 copies.

Figure 2. CNVs amplify different sequences at the *RTA3* locus

A.   Span of 16 different CNVs at the *RTA3* locus in *C. parapsilosis*. The coding sequence of *RTA3*
is shown by a dark purple box, with the flanking UTRs in a lighter color. The neighbouring genes
MAK16 and *RTA2* are also shown. The extents of the tandemly repeating units in each of 16 CNVs
are shown by black boxes, labelled (A-P) on the left, and their lengths are indicated in white. Exact
breakpoints were identified by interrogating split reads from the Illumina data, and the array structure
of CNVs D and K was verified by MinION sequencing of isolates UCD321, MSK478, and MSK812.

B.   Copy number of *RTA3* in each *C. parapsilosis* isolate. Each dot represents a single isolate:
blue isolates from MSK, red from CHU de Nantes, and dark grey from other sources. Median and
interquartile ranges are shown for CNVs present in more than one strain. Dots are jittered for clarity.

C.   *RTA2/3* fusion gene in strain 434 containing CNV-P. A coverage plot created by
PyGenomeTracks [47] is shown at the top. The schematic shows the position of CNV breakpoints in
relation to the CDS of both genes, and the structure of the inferred array of fusion genes.

D.   Amplifications of *RTA3* in *C. orthopsilosis*. Orientation of the chromosome has been flipped to
highlight the similarities to *RTA3* CNVs in *C. parapsilosis*.

E.   A deletion in *C. orthopsilosis* generates a *CoRTA3/CoRTA2* fusion. A coverage plot of the

locus is shown at the top. The schematic shows CNV breakpoints in both *CoRTA3* and *CoRTA2* and

how the resulting fusion gene is formed at one allele, leaving the other allele intact.

For CNV-K, three isolates have roughly half the *RTA3* copy number of other isolates with the

same CNV (~13x instead of ~26x; Fig 2B; S1 Table). We considered the possibility that only

one allele of *RTA3* was amplified in these isolates (*C. parapsilosis* is diploid), and that both

alleles were amplified in other isolates. We explored this issue by using long read

sequencing (Oxford Nanopore) of *C. parapsilosis* MSK812 which is one of the CNV-K

isolates with fewer copies. We found that it has 8 copies of *RTA3* at one allele and 6 copies

at the other (S1A Fig). Similarly, sequencing of isolate MSK478 (which also has CNV-K, with

higher copy number of *RTA3*) showed that it has at least 11 copies at both alleles (S1B Fig).

Therefore the variation in copy number among isolates with CNV-K is due to expansion or

contraction of the array in both alleles, and not due to hemizygosity for the amplification.

Two of the *RTA3* CNVs alter the structure of the protein it encodes. The amplification in

CNV-P starts within the *RTA3* open reading frame, and ends within the related adjacent

gene *RTA2* (Fig 2A,C). This repeat structure generates an array of in-frame *RTA2/RTA3*

gene fusions, with the N-terminus derived from *RTA2* and the C-terminus from *RTA3* (Fig

2C). *RTA3* and *RTA2* are very similar genes, which likely facilitated the fusion event. In

CNV-H, the Rta3 protein is slightly truncated because this CNV consists of a tandem

duplication with an endpoint 31 bp upstream of the stop codon of *RTA3*, resulting in a protein

that is 10 amino acids shorter than the wild type.

CNV-B is substantially different from the others because it is an amplification of only a small

region (269 bp) that is upstream and completely outside the *RTA3* coding sequence (Fig

2A). Sequencing read coverage analysis suggests that the repeat has a complex

organization in which direct repeats are interspersed with inverted copies of a central

segment flanked by inverted-repeats, resulting in an N:(2N-1):N copy number pattern (S2A

Fig). This repeat structure was confirmed by MinION sequencing of isolates MSK802 and

MSK803. The structure of the CNV-B repeat is similar to the DUP-TRP/INV-DUP structure

seen in some human CNVs [48] and is reminiscent of amplifications formed via Origin-

Dependent Inverted Repeat Amplification (ODIRA) [49]. The ODIRA model proposes that

replication errors at origins that are surrounded by short inverted repeats generate

extrachromosomal intermediates, which are then integrated at the original site [49]. This

results in complex CNVs with repeat units in head-to-head and tail-to-tail arrangements that

are similar to the CNV-B structure. To date, ODIRA has only been observed in *S. cerevisiae*,

and it is unclear whether it occurs in other yeasts [49]. If CNV-B were caused by ODIRA, we

would anticipate that the central region of the repeat would contain an origin of replication

(S2B Fig). However, determined the temporal order of replication in *C. parapsilosis* using

Sort-seq (S2C Fig), and we found no evidence that there is an origin near *RTA3* [50].

We also looked for evidence of *RTA3* amplification in other *Candida* species*. RTA3* is not

amplified in 200 sequenced *C. albicans* genomes [51, 52]. However, we identified four

different *RTA3* CNVs among 36 strains of *Candida orthopsilosis*, a close relative of *C.*

parapsilosis (Fig 2D). Only one of these (CNV-Co1) consists of a simple tandem

amplification of the entire *CoRTA3* ORF. Two others, CNV-Co2 and CNV-Co3, amplify a

fusion of the *CoRTA2* and *CoRTA3* genes, similar to CNV-P in *C. parapsilosis* (Fig 2D).

Interestingly, the final *C. orthopsilosis* CNV (CNV-Co4) involves a deletion. In the two strains

containing CNV-Co4, the 3' end of *CoRTA3*, the 5' end of *CoRTA2,* and the intergenic space

between them are deleted at one allele (Fig 2E). This results in a new fusion gene, with the

N-terminus derived from *CoRTA3* and the C-terminus derived from *CoRTA2*. Single copies

of *CoRTA2* and *CoRTA3* are intact on the other allele (Fig 2E). This is the only example we

have seen of an *RTA3* deletion in *C. parapsilosis, C. orthopsilosis* or *C. albicans*.

Copy number of *RTA3* is correlated with miltefosine, but not fluconazole, resistance

Deletion of *RTA3* in *C. albicans* has been shown to increase susceptibility to miltefosine [33]

and possibly to fluconazole [41]. We therefore investigated the effect of copy number on

resistance of *C. parapsilosis*. Fig 3A shows that amplification of *RTA3* is associated with

miltefosine resistance. For example, *C. parapsilosis* strains with only two copies of *RTA3*

(one at each allele, e.g. the reference strain CLIB214) fail to grow at miltefosine

concentrations of 10 µg/ml, whereas all the strains with *RTA3* amplifications can grow at this

concentration. Strains with CNVs A, H, I, J and L can tolerate miltefosine concentrations up

to at least 30 µg/ml, as can isolates with CNV-B (which amplifies only the region upstream of

RTA3). The CNV with the weakest effect on MF resistance is CNV-G, which can tolerate 10-

15 µg/ml, which is still higher than the reference strain CLIB214.

When different isolates carrying the same CNV are compared, miltefosine resistance is seen

to be correlated with the copy number of *RTA3* (Fig 3B). For example, *C. parapsilosis*

isolates MSK812 and MSK1129, which have 13-14 copies of CNV-K, can tolerate miltefosine

concentration up to ~20 µg/ml, whereas *C. parapsilosis* MSK2086 which has 27 copies

survives up to 30 µg/ml. Similarly, *C. parapsilosis* MSK815 with 17 copies of CNV-D and

MSK794 with 5 copies of CNV-F tolerate miltefosine concentrations up to ~20-25 µg/ml

 whereas UCD321 and FM43 with 27 and 15 copies of CNV-D and CNV- F respectively

273   survive up to ~30 µg/ml (Fig 3B).

Figure 3. CNVs of *RTA3* correlate with resistance to miltefosine

A.   A representative isolate of most CNV patterns (Fig 2) was grown on YPD with increasing

concentrations of miltefosine (MF) or fluconazole (FLC) as shown. Cells were serially diluted (1/5).

The CNV pattern is indicated on the left, with the strain name shown on the right. MF and FLC plates

were incubated for 48 hr. "Ref" indicates the reference strain *C. parapsilosis* CLIB214.

B.   Copy number is directly correlated with miltefosine resistance. Isolates with different copy

numbers of CNV-K, CNV-D and CNV-F were grown as in (A). The *RTA3* copy number of each isolate

is shown on the left.

C.   Deleting *RTA3* reduces resistance to miltefosine. *RTA3* was deleted in *C. parapsilosis*

CLIB214 using CRISPR-Cas9. The growth of two biological replicates is shown, as in (A), except that

incubations on FLC were for 72 h.

The isolates have variable levels of sensitivity to fluconazole (Fig 3), but the copy number of

RTA3 does not correlate with susceptibility to this drug. For example, isolates MSK815 and

UCD321 which have 17 and 27 copies of CNV-D differ in their susceptibility to miltefosine,

but they both tolerate fluconazole levels of 6 µg/ml.

We also found that deleting *RTA3* in the reference isolate of *C. parapsilosis* (CLIB214) by

CRISPR-Cas9 editing [53] results in increased sensitivity to miltefosine (Fig 3C). Very low

levels of miltefosine were used (up to 4 µg/ml), because the parental strain, without any

RTA3 amplifications, is very sensitive. However, susceptibility to fluconazole was not

affected by deleting *RTA3* (Fig 3C).

Increasing the *RTA3* copy number correlates with increased expression, as shown by West

et al [43]. However we wondered whether amplifying the upstream region (CNV-B) has the

same effect as amplifying the entire open reading frame. To explore this further, we

measured *RTA3* expression by RT-PCR in one isolate of *C. parapsilosis*, MSK808, with

approximately 42 copies of *RTA3* (CNV-I), and two isolates (MSK802 and MSK803) in which

only the promoter region is amplified (CNV-B). We found that *RTA3* expression is

approximately 22-fold higher in *C. parapsilosis* MSK808, and 2.8- to 6.6-fold higher in *C.*

parapsilosis MSK802 and MSK803 than in the reference strain CLIB214 (Table 1). There are

28 copies of the direct repeat in CNV-B in *C. parapsilosis* MSK802 and 24 in *C. parapsilosis*

MSK803. Thus, *RTA3* promoter region amplification (CNV-B) can lead to increased

expression.

Generation of miltefosine-resistant strains by experimental evolution

The prevalence of *RTA3* amplification in *C. parapsilosis* isolates suggests that there may be

some strong selective pressure inducing amplification. To determine if miltefosine is the

driving force, we characterized the effect of exposing isolates to increasing concentrations of

miltefosine, in an adaptive laboratory evolution approach. We started with two isolates with

only one copy of *RTA3* at each allele that were in the same clade as other isolates that had

undergone amplification - *C. parapsilosis* MSK795 from Clade 1, which is related to isolates

that have undergone 4 different CNVs (B, C, E, and F), and *C. parapsilosis* MSK247 from

Clade 5, which is related to isolates with CNVs A, D, J and M (Fig 1; Fig 4). Five lineages

were evolved from MSK247 (247A to 247E) and one from MSK795 (795B). Isolates were

cultured in YPD with increasing concentrations of miltefosine, up to a maximum of 32 µg/ml,

over a 26-day period (Fig 4A). Sixteen randomly picked evolved colonies from each lineage

tolerated miltefosine levels of 40 µg/ml (Fig 4B). The genomes of ten isolates derived from

MSK247 and two from MSK795 were sequenced, together with the parental strains. The

sequenced strains are listed in Methods.

Figure 4. Generation of miltefosine-resistant *C. parapsilosis* isolates by adaptive laboratory evolution

A.   Six colonies (5 derived from *C. parapsilosis* MSK247 and one from *C. parapsilosis* MSK795) were inoculated in YPD at A_{600}=0.1,and incubated at 30℃ for 10 h. Miltefosine (indicated with a blue drop) was added to a final concentration of 1 µg/ml, the cultures were incubated for a further 14 h, and then re-inoculated into the same media using a 1/100 dilution and incubated for 24 h. The dilution was repeated every 24 h for 3 days. Cells were then inoculated in fresh media, grown for 10 h, and the miltefosine concentration was doubled. The process was repeated until the concentration of miltefosine reached 32 µg/ml (24 days). On day 25, the overnight cultures were plated on YPD with 16 µg/ml miltefosine. Sixteen colonies picked from each lineage for further analysis, and the genomes of 10 were sequenced.

B.   Growth of representative isolates evolved from *C. parapsilosis* MSK247 and *C. parapsilosis* MSK795 on 40 µg/ml miltefosine. WT = parental strain.

Miltefosine-resistant isolates acquired homozygous disruptions in two flippase genes

We did not find any evidence of amplification of the *RTA3* locus in any of the evolved miltefosine-resistant strains. However, by comparing the sequences of the evolved strains to those of the parental strains, we identified two genes with homozygous loss-of-function variants in all 10 resistant isolates: *CPAR2_102700* and *CPAR2_303950*. These variants include frameshifts, nonsense mutations, and missense mutations that are predicted to be deleterious (Fig 5A). *CPAR2_102700* and *CPAR2_303950* encode putative Class 3 P4-ATPases and are homologs of the PC flippase genes *DNF1* and *DNF2* in *S. cerevisiae* (Fig 5B).

The two sequenced isolates derived from *C. parapsilosis* MSK795 (795B1 and 795B16) acquired mutations in both *CPAR2_102700* and *CPAR2_303950*, whereas lineages derived from *C. parapsilosis* MSK247 acquired mutations only in *CPAR2_303950* (Fig 5A). However, subsequent analysis revealed that *C. parapsilosis* MSK247 contains a homozygous natural variant in *CPAR2_102700*, resulting in a Trp-to-Stop nonsense mutation (W1280X; Fig 5A). *C. parapsilosis* MSK247 tolerates miltefosine concentrations of 5 µg/ml, whereas *C. parapsilosis* MSK795 fails to grow (Fig 5C). Derivatives of both parents that carry homozygous inactivating mutations in both *CPAR2_30395* and *CPAR2_102700* can grow up to 30 µg/ml.

Figure 5. Protein-disrupting mutations in two flippase genes confer resistance to miltefosine in laboratory evolved strains

A. Schematic showing homozygous mutations in P4-ATPases *CPAR2_102700* and *CPAR2_303950*. Mutations named in black were acquired during the laboratory evolution experiment, in the evolved strains named below the mutation name. The mutation named in red is a natural homozygous variant found in the *C. parapsilosis* MSK247 isolate and all strains derived from it. The predicted transmembrane topology of each protein (from TMHMM [54]) is shown, where red peaks show predicted transmembrane domains.

B. Phylogeny of P4-ATPases in *C. parapsilosis*, *C. albicans*, and *S. cerevisiae*. The P4-ATPase genes in three yeast species were identified by BLAST, aligned and a tree was constructed using SeaView [55]. Monophyletic groups of genes were assigned to established classes (1-4) [56]. Both *CPAR2_102700* and *CPAR2_303950* (in bold) belong to Class 3 P4-ATPases.

C. Deleting *CPAR2_102700* and *CPAR2_303950* increases resistance to miltefosine, but not to fluconazole. *CPAR2_102700* and *CPAR2_303950* were deleted singly or together in the *C. parapsilosis* CLIB214 background, and growth was compared to isolates evolved from *C. parapsilosis* MSK247 (247C1) and *C. parapsilosis* MSK795 (795B1), as in Figure 3.

This result suggests that maximum levels of miltefosine resistance requires inactivation of both genes, *CPAR2_30395* and *CPAR2_102700*. To test this hypothesis, we deleted them, both separately and together, in the *C. parapsilosis* CLIB214 genetic background using

CRISPR-Cas9 editing (Fig 5C). Deleting *CPAR2_102700* alone in *C. parapsilosis* CLIB214

allows growth up to 5 µg/ml miltefosine, whereas deleting *CPAR2_303950* alone has no

effect (Fig 5C). However, strains in which both *CPAR2_303950* and *CPAR2_102700* are

deleted tolerate at least 30 µg/ml miltefosine, a similar concentration to strains derived from

the experimentally evolved isolates (MSK795B1, MSK247C1) (Fig 5C). Deleting

CPAR2_102700 alone, or in combination with *CPAR2_303950*, slightly increases sensitivity

to fluconazole (Fig 5C).

391 Discussion

There is some debate over whether Rta3-like proteins are transporters or signaling

receptors. Rta3 is a member of the Rta1-family, which encodes proteins with 7

transmembrane domains, similar to the structure of G-protein-coupled receptors (GPCRs)

[7]. Within this family, Rta3 is more closely related to *S. cerevisiae* Rsb1 than to other

members. Most of the early evidence suggested that Rsb1 directly flips LCBs in the plasma

membrane [35]. However, Johnson et al [36] suggested that rather than acting as a

transporter, Rsb1 determines resistance to the LCB phytosphingosine by regulating

endocytosis of the tryptophan transporter Tat2, either by signalling through a G-protein (as a

GPCR), or by an arrestin-mediated effect on targeting of ubiquitin ligases. In addition,

Srivistava et al (2017) found that deleting *RTA3* in *C. albicans* resulted in increased flipping

of a fluorescently labelled PC, possibly by controlling activity of an unknown flippase [33].

However, several pieces of evidence suggest that Rsb1 is a direct lipid transporter,

specifically a floppase. Firstly, Rsb1 shares no sequence similarity with other yeast GPCRs

[7]. Secondly, Makuta et al [57] showed that Rsb1 (and not other Rta1-family members)

regulates LCB transport, and that this activity is dependent on a loop region following TMS5,

and not the C-terminus as expected for a GPCR. Rsb1 in *S. cerevisiae* and Rta3 in *C.*

albicans are located in the plasma membrane [33, 34, 41]. The increased flipping observed

by Srivistava et al (2017) [33] may be due to crosstalk between sphingolipids and

glycerophospholipids (e.g. PC), as shown in *S. cerevisiae* by Kihara and Igarashi (2004)

[35]. Changes in the distribution of one may be compensated by changing the distribution of

the other [7]. The association of *RTA3* copy number amplification with miltefosine resistance

in *C. parapsilosis* is also consistent with a role as a transporter. It is therefore likely that Rta3

is a transporter of LCBs, similar to Rsb1, and that its overexpression and deletion cause

changes in the localization of both PC and LCBs. Our results also suggest that *C.*

parapsilosis CPAR2_102700 and *CPAR2_303950*, members of the class 3 family of P4-

ATPases, are the direct flippases of PC. Our data however cannot rule out the alternative

model of Johnson et al [36] whereby Rta3 regulates the function of *CPAR2_102700* and

CPAR2_303950, rather than directly acting as a transporter.

RTA3 is adjacent to a paralogous gene (called *RTA2*) in many species in the *Candida* (CUG-

Ser1) clade [58, 59]. In fact, *C. parapsilosis* CNV-P results from amplification of an

RTA3/RTA2 fusion, with the N-terminus derived from Rta2 and the C-terminus from Rta3

(Fig 2C). Deleting *RTA2* in *C. albicans* increases sensitivity to azole drugs, and

overexpression increases resistance [60]. Whaley et al [41] showed that deleting *RTA3*

increases sensitivity of an azole-resistant isolate of *C. albicans*, but does not have any effect

on a reference isolate. Our results show that *C. parapsilosis RTA3* is unlikely to play any role

in azole resistance: deleting the gene does not increase sensitivity, and levels of resistance

are not associated with *RTA3* copy number in several different genetic backgrounds.

However, the sequence similarity between *RTA2* and *RTA3* in *C. parapsilosis* (and in the

related species *C. orthopsilosis* and *C. metapsilosis*) is much higher than between *C.*

albicans RTA2 and *RTA3* (S3A Fig). This similarity, and the phylogenetic relationship among

these sequences (S3A Fig), indicates that in the ancestor of *C. parapsilosis*/*C.*

orthopsilosis/*C. metapsilosis*, a gene conversion event occurred in which *RTA3* overwrote

RTA2. The biggest difference between Rta3 and Rta2 in *C. parapsilosis* is that the former

has a longer C-terminus (S3B Fig). Because we have never observed amplifications of

RTA2 in miltefosine-resistant *C. parapsilosis* isolates, we suggest that the two genes may

have different functions, and that the difference may be related to the C terminus. In *C.*

albicans, *RTA2* (and possibly *RTA3*) also contribute to tunicamycin-induced ER stress,

which has not been tested in *C. parapsilosis* [61, 62].

Of the 14 CNVs resulting in tandem array amplification of *RTA3,* only CNV-A and CNV-E

have obvious repeat sequences at the CNV endpoints, direct repeats of 19 bp and 10 bp

respectively. The other 12 either have no repeats, or the repeat is <4 bp. The longer repeat

sequence at the breakpoints of CNV-A may explain why it has originated independently

three times (Fig 1). CNV-A and CNV-E may result from non-allelic homologous

recombination between the direct repeats [48]. It is likely that CNV-P also originated by non-

allelic homologous recombination due to the high sequence similarity between *RTA3* and

RTA2. CNVs with shorter direct repeats at their endpoints may result from microhomology-

mediated break induced replication [63]. However, there is no obvious mechanism for CNVs

with no repeats at the breakpoints.

CNV-B (upstream from the coding sequence) is particularly intriguing, because it results in

inversions and triplications of part of the repeated sequence (S2A Fig). This structure is

reminiscent of the origin-dependent ODIRA model for CNV generation [49]. We did not find

any evidence of a replication origin in the amplified sequence (S2A Fig), but it remains

possible that a rarely used origin is present. It is likely that the amplified region contains a

promoter because expression of *RTA3* is increased (Table 1).

Our results strongly suggest that there is selective pressure driving amplification of *RTA3* in

C. parapsilosis. This conclusion is based on our observation that amplification occurred in

isolates from diverse genetic and environmental backgrounds, distributed across the *C.*

parapsilosis phylogeny (Fig 1). We also found that there have been multiple independent

events of amplification, with at least 16 unique endpoints. West et al (2021) previously

described *RTA3* amplifications in 4 *C. parapsilosis* isolates, including one from the New York

subway [43]. Those amplifications also have different endpoints, but because the data

comes from metagenomics analyses, we cannot determine if their endpoints differ from the

16 CNVs we describe.

The CNVs sometimes encompass parts of the adjacent genes, but *RTA3* is the only

complete gene amplified in all of them (except for CNV-B, where its likely promoter is

amplified, and CNV-H where its 3' end is slightly truncated in the amplified copies). The

majority of the isolates in this study originated from clinical sources, although one with an

RTA3 amplification (*C. parapsilosis* UCD321) has an environmental origin (it was isolated

from soil). There are several known examples of environmental conditions that induce or

selecting for CNVs, such as copper resistance resulting from amplification of *CUP1* in *S.*

cerevisiae [64, 65], and CNVs associated with growth of *C. albicans* in the presence of

fluconazole [66]. It is highly unlikely that the *RTA3* amplifications we discovered were driven

by exposure to miltefosine because it is not a commonly used antifungal drug and, as we

have shown, exposure to miltefosine is more likely to result in loss-of-function mutations in

flippase genes. At present, we do not know what kind of selective pressure led to the

widespread amplifications of *RTA3* in *C. parapsilosis*.

Miltefosine has potential as an antifungal drug [20-22], and it has recently been designated

as an orphan drug for the treatment of invasive candidiasis

(https://www.accessdata.fda.gov/scripts/opdlisting/oopd/detailedIndex.cfm?cfgridkey=843921

). However, our analyses suggest that it would be a poor choice, at least for invasive

Candida parapsilosis infections. Many isolates are naturally resistant because of

amplifications of *RTA3*, and in others, resistance rapidly arises due to loss-of-function

mutations in the flippase genes. Although mutations in two flippase genes are required for

resistance, we note that many of the *C. parapsilosis* samples we examined (56 of 170) have

predicted loss-of-function variants in *CPAR2_102700*, meaning that acquiring a mutation in

CPAR2_303950 would be sufficient to render them highly resistant to miltefosine.

Materials and methods

Strains and growth. Isolates used are listed in S1 Table. Isolates were maintained on YPD

agar (1% Bacto Yeast Extract (212750, Sigma), 2% Bacto Peptone (211677, Sigma), 2%

Bacto Agar (214010, Sigma), 2% D-(+)-Glucose (G8270, Sigma)), and liquid cultures were

grown in 5 ml YPD broth without agar at 30ºC and 200 rpm shaking overnight. For serial

dilutions, 0.5 ml of the overnight cultures were harvested at 13,000 rpm at room temperature

for 1 min, washed twice in 0.5 ml of phosphate-buffered saline (PBS) buffer (BR0014G,

Thermofisher), resuspended in 0.5 ml PBS, and diluted to A_{600}= 0.0625 (approximately 6.25 x

10^5 cells/ml) in PBS. Five-fold serial dilutions were made in PBS and transferred with a

pinner to YPD agar plates containing miltefosine (M5571, Sigma Aldrich) or fluconazole

(F8929, Sigma Aldrich) at the indicated concentrations. Pinned plates were incubated at

30ºC for the indicated times and photographed using a Singer PhenoBooth.

Gene deletions/disruptions. The entire open reading frames of *CPAR2_104610* (*RTA3*)

and the flippases *CPAR2_303950* and *CPAR2_102700* were deleted using CRISPR-Cas9

with the pCP-tRNA system, as described in Lombardi et al [53]. All primers used for gRNA

and repair template synthesis are listed in S2 Table.

Microevolution of miltefosine resistant isolates. The microevolution method was

modified from Papp et al [67] and Ene et al. [68]. Three colonies from *C. parapsilosis*

MSK247 and *C. parapsilosis* MSK795 were originally chosen. Subsequent analysis showed

that 5 evolved lineages were derived from *C. parapsilosis* MSK247 (247A to E), and one

from *C. parapsilosis* MSK795 (795B). Colonies were incubated for 10 h in 5 ml YPD at 200

rpm and 30ºC. Miltefosine was added to a final concentration of 1.0 μg/ml, and the cultures

were incubated for a further 14 h. The cultures were diluted (1:100) into fresh media

containing the same concentration of miltefosine every 24 h for 3 days. Overnight cultures

were then diluted to an A_{600} = 0.1, incubated for 10 h, and the miltefosine concentration was

doubled. The 5-day cycle was repeated until a concentration of 32 μg/ml of miltefosine was

reached (see Fig 4A). Cultures were diluted and plated on YPD agar plates containing 16

μg/ml miltefosine, yielding 100-200 colonies each plate, and were incubated at 30ºC for 48

h. The genomes of ten resistant isolates (247A1, 247B1, 247C1, 247D1, 247D2,

247D16,247E1, 247E16, 797B1 and 795B16), together with the parental strains were

sequenced by Beijing Genomic Institute via DNBseq.

Illumina sequencing. Illumina sequencing of MSK isolates was carried out as described in

Zhai et al [42]. 45 *C. parapsilosis* isolates from Centre Hospitalier Universitaire de Nantes,

France were screened for resistance to miltefosine, and 16 isolates were chosen for

sequencing. Genomic DNA was isolated from 5 ml overnight cultures in YPD at 30℃ using

phenol-chloroform-isoamyl alcohol extraction. Cell pellets were resuspended in 200 µl of

extraction buffer (Triton X 100 2% m/v, NaCl 100mM, Tris 10mM pH 7.4, EDTA 1mM, SDS

1% m/v), transferred to screw cap tubes, and approximately 0.3 g of acid-washed beads and

200 µl of Phenol-chloroform-isoamyl alcohol (25:24:1) was added. Cells were lysed using a

1600 MiniG bead beater from Spex SamplePrep for 6 x 30 seconds with 30 s pauses in

between, and then centrifuged at 14000 rpm for 10 min at room temperature. The

supernatant was extracted twice more with 200 µl of TE and 200 µl of phenol-chloroform-

isoamyl alcohol and one 30 s agitation in the bead beater. DNA was precipitated using 80 µl

of ammonium acetate (7.5M) and 1 ml of 100% isopropanol, washed with 1 ml of 70%

ethanol and air-dried. Pellets were resuspended in 400 µl of TE with 1 µl of RNAse A (100

mg/ml) incubated overnight at 37℃, DNA was precipitated again, and resuspended in 150 µl

water. Illumina sequencing was performed by the UCD Conway Genomics Core using a

NextSeq 500. 1 ng of gDNA was tagmented (fragmented and tagged with adapter

sequences) using the Nextera kit transposome, Dual-indexed paired-end libraries were

prepared using Nextera XT DNA Library Prep kit. An Illumina NextSeq500 mid output 300

cycle sequencing kit was used to prepare and run the flowcell (HVGWJAFX2).

Oxford Nanopore sequencing. Strains were grown overnight in 50 ml YPD broth and

genomic DNA was extracted from approximately $4x10^9$ cells using a QIAGEN Genomic-tip

100/G kit (10223, QIAGEN) with minor modifications. The lyticase incubation was extended

to 2 h, and the proteinase K incubation was extended to overnight (~15 h). DNA libraries

were prepared using three different kits as per manufacturer's instructions. Libraries from *C.*

parapsilosis MSK812 and UCD321 were prepared using a Ligation Sequencing Kit (SQK-

LSK109, Oxford Nanopore) using 1 µg of DNA per strain. DNA was repaired using a

NEBNext FFPE DNA Repair Mix (M6630, New England Biolabs) and NEBNext Ultra II End

repair/dA-tailing Module (E75460, New England Biolabs). Adapters were ligated using

NEBNext Quick Ligation Module (E60560, New England Biolabs). Libraries were sequenced

on a MinION Mk1C device. Priming, loading and washing was carried using the EXP-

FLP002, and SQK-LSK109 and EXP-WSH002 Oxford Nanopore kits respectively as per

manufacturer's instructions. The genomes were sequenced in parallel on one flow cell; the

first isolate (*C. parapsilosis* MSK812) was sequenced for 24 h, and the flow cell was

washed, re-primed and loaded with the *C. parapsilosis* UCD321 isolate for an additional ~48

h. Libraries from *C. parapsilosis* MSK802 and MSK803 were prepared using a Rapid

Barcode Sequencing Kit (SQK-RBK004, Oxford Nanopore) using 400 ng of DNA per

sample. Samples were multiplexed using barcodes R04 and R05 respectively. Libraries

were sequenced on an original MinION device for 72 h. Priming and loading was performed

using the EXP-FLP002 and SQK-RBK004 kits respectively. The *C. parapsilosis* MSK478

library was prepared using a Rapid Sequencing Kit (SQK-RAD004, Oxford Nanopore) using

400 ng of DNA, and sequenced for 72 h using an original MinION device. Basecalling was

performed for all samples using guppy_basecaller with the following parameters: "--

input_path fast5 --save_path fastq -c dna_r9.4.1_450bps_fast.cfg --verbose_logs --

cpu_threads_per_caller 5 --num_callers 10". Guppy v.4.2.2+effbaf8 was used for samples

MSK812 and UCD321 and Guppy v.3.6 was used for samples MSK802 and MSK803. For

multiplexed samples MSK802 and MSK803, demultiplexing was performed using

guppy_barcoder with the following parameters: "--barcode_kits SQK-RBK004 -t 30 --

verbose_logs --trim_barcodes". Reads were filtered using NanoFilt with the following

parameters: "-l 1000 -q 7" [69].

Sequence analysis. Illumina reads were trimmed with Skewer version 0.2.2 using tags "-m

pe -t 4 -l 35 -q 30 -Q 30" [70]. The trimmed reads were aligned to the *C. parapsilosis*

reference genome using bwa mem version 0.7.12 . The resulting BAM files were sorted and

duplicate reads were marked using GenomeAnalysisToolkit (GATK version 4.0.1.2) SortSam

and MarkDuplicates tools respectively [44]. Variants were called with GATK HaplotypeCaller

using the tag "--genotyping_mode DISCOVERY", combined using GATK CombineGVCFs

and joint-genotyped using GATK GenotypeGVCFs. Variant files were filtered for read depth

(<15) using and genotype quality (<40) using GATK VariantFiltration. Additionally, clusters of

SNPs (5 SNPs in a 100bp window) were filtered using GATK VariantFiltration. A custom

script was used to remove variants that were flanked on either side by a long string of mono-

or dinucleotide repeats, and variants that were called as heterozygous but had an allele

depth ratio <0.25 or >0.75 (https://github.com/CMOTsean/milt_variant_filtration).

Additionally, for tree construction, indels were excluded using GATK SelectVariants with the

tag "--select-type-to-include SNP". For analysis of the evolved strains, a custom script was

used to filter out variants in the evolved strains that were also present in the respective

parent strain (https://github.com/CMOTsean/milt_variant_filtration).

SIFT4G analysis. A SIFT prediction database was created for *C. parapsilosis* using the

SIFT4G algorithm [71], and the recommended Uniref90 database as a reference for protein

sequences [72]. The *C. parapsilosis* prediction database was used to annotate variants from

the evolved strains with whether they are likely to be deleterious to protein function.

For each annotated gene in *C. parapsilosis*, the number of evolved strains which carried a

variant predicted to be protein function-affecting by SIFT in that gene was tallied. Variants

were also visualised using Integrative Genomics Viewer (IGV) to manually check results

[73].

Phylogeny construction. Called SNPS were concatenated and heterozygous sites were

resolved randomly to either allele by 1000 iterations of Random Repeated Haplotype

Sampling [45]. SNP trees were then constructed from each of the 1000 haploid inputs, using

RAxML (v8.2.12) with the GTRGAMMA model of nucleotide substitution and the random

number seed "-p 12345" [74]. The tree with the highest maximum likelihood score was

chosen, and the remaining 999 trees were used to generate branch support values.

Estimating copy number. For each strain with a CNV at *RTA3*, the average coverage

across the *RTA3* ORF was found using BEDTools coverage (v2.29.2) [75]. This value was

divided by the average genome coverage (found with BEDTools genomecov) and multiplied

by two to adjust for ploidy, to calculate an estimate for the copy number of *RTA3*. For CNVs

that did not cover the entire *RTA3* ORF, the average coverage of a representative section of

the CNV was used instead.

MinION read sequences were used to identify the exact CNV copy number for a set of

strains. The respective CNV sequences plus 1 kb flanking sequence on either side, were

searched against the set of MinION reads for strains MSK478, MSK802, MSK803, MSK812,

and UCD321 using BLASTN (v2.10.0) [76]. The search outputs were parsed with RECON-

EBB (https://github.com/CMOTsean/recon-ebb) to estimate and visualise copy number from

reads which include hits for multiple copies of the CNV sequence and both regions of

flanking sequence, i.e the reads which covered the entirety of the repeat region.

Replication timing profiles. Relative replication time was determined by sort-seq as

described previously [77]. Briefly, replicating (S phase) and non-replicating (G2 phase) cells

were enriched from an asynchronously growing culture by FACS based on DNA content. In

each sample, genomic DNA was extracted and subjected to Illumina sequencing to measure

relative DNA copy number. Replication timing profiles were generated by normalizing the

replicating (S phase) sample read count to the nonreplicating (G2) sample read count in 1-kb

windows.

Quantitative RT-PCR. Cell harvesting and RNA extraction methods were adapted from Cravener and Mitchell, 2020 [78]. Cells were inoculated from overnight cultures to an A_{600} of 0.2 in 25 ml of pre-warmed YPD broth and incubated at 30°C for 6 h at 200 RPM using an orbital shaker. Cells were then harvested via vacuum filtration using MicroPlus-21 Sterile 0.45µm filters (10407713, Whatman) and stored at -80°C for at least 24 h prior to RNA extraction.Cells were lysed by mechanical disruption as recommended in the QIAGEN RNeasy Mini Kit (74104, QIAGEN) with some modification [78]. Cells were lysed in RLT lysis buffer (QIAGEN) and Phenol:Chloroform:Isoamyl Alcohol (25:24:1)(P3803, Sigma) in a 1:1 ratio. Lysis was performed using a 1600 MiniG from Spex SamplePrep using 30 s lysis followed by 30 s chilling on ice for a total of 6 min. RNA was extracted using a QIAGEN RNeasy Mini Kit followed by two rounds of DNase digestion: one on-column using the QIAGEN RNase-Free DNase Set (79254, QIAGEN); and one off-column using Invitrogen's TURBO DNA-free Kit (AM1907, ThermoFisher). cDNA was synthesized from 1 µg RNA using M-MLV reverse transcriptase (9PIM170, Promega) and Oligo(dT)15 primers (C110A, Promega), following manufacturer's instructions. qPCR was performed in 20 µl reactions using 50 ng cDNA using FastStart Universal SYBR Green Master (Rox) (4913850001, Sigma) as per manufacturer's instructions on an Agilent Technologies Stratagene Mx3005p machine using default "two-step" settings. All primers are listed in S2 Table. Relative quantification was performed using the 2(-Delta Delta C(T)) method by comparing expression to *ACT1*, and using *C. parapsilosis* CLIB214 as the calibrator strain. Calculations and statistics were performed in R using the pcr package [79].

Data availability. All sequencing data is deposited at NCBI under BioProject number XXXX (awaiting accession numbers).

Acknowledgements
For Open Access, the authors have applied a CC BY public copyright license to any Author Accepted Manuscript version arising from this submission. This work was supported by Science Foundation Ireland (grant numbers 19/FFP/6668 and 18/CRT/6214 to G.B. and 20/FFP-A/8795 to K.H.W), the Irish Research Council (A.R.), the Chinese Scholarship Scheme (FZ), Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) grant RO-5328/2 (T.R.), National Institutes of Health (NIH) grants R01 AI093808 (T.M.H.), R21 AI156157 (T.M.H.), the Ludwig Center for Cancer Immunotherapy (T.M.H.), the Susan and Peter Solomon Divisional Genomics Program (T.M.H.), and by NIH P30 CA008748 (Cancer Center Core Grant to MSKCC). CAN acknowledges support from the Biotechnology and Biological Sciences Research Council (BBSRC), part of UK Research and Innovation, Core Capability Grant BB/CCG1720/1 and the National Capability (BBS/E/T/000PR9814).

667   Thanks to the genomics core facility in the Conway Institute UCD for help with Illumina

668   sequencing, to Eoin Ó Cinnéide and Letal Salzberg (UCD) for help with MinION sequencing,

669   and to Lisa Lombardi for help with the adaptive evolution experiment.

672   Table 1. Expression of *RTA3* in strains with different amplifications

Isolate	Relative Expression	Range	P Value
MSK802 (CNV-B)	6.57	2.12 - 20.35	7.37E-03
MSK803 (CNV-B)	2.79	1.24 - 6.25	3.39E-02
MSK808 (CNV-I)	22.50	6.52 - 77.62	6.92E-05
CLIB214 (Reference)	1.00	0.19 - 5.27	N/A

References

1.   Renne MF, de Kroon AI. The role of phospholipid molecular species in determining

the physical properties of yeast membranes. FEBS Lett. 2018;592(8):1330-45.

2.   Sharma SC. Implications of sterol structure for membrane lipid composition, fluidity

and phospholipid asymmetry in *Saccharomyces cerevisiae*. FEMS Yeast Res.

2006;6(7):1047-51.

3.   Grillitsch K, Tarazona P, Klug L, Wriessnegger T, Zellnig G, Leitner E, et al. Isolation

and characterization of the plasma membrane from the yeast *Pichia pastoris*. BBA-

Biomemb. 2014;1838(7):1889-97.

4.   Devaux PF, Herrmann A, Ohlwein N, Kozlov MM. How lipid flippases can modulate

membrane structure. BBA-Biomemb. 2008;1778(7-8):1591-600.

5.   Andersen JP, Vestergaard AL, Mikkelsen SA, Mogensen LS, Chalat M, Molday RS.

P4-ATPases as phospholipid flippases—structure, function, and enigmas. Front Physiol.

2016;7:275.

6.   Daleke DL. Regulation of transbilayer plasma membrane phospholipid asymmetry. J

Lipid Res. 2003;44(2):233-42.

7.   Manente M, Ghislain M. The lipid-translocating exporter family and membrane

phospholipid homeostasis in yeast. FEMS Yeast Res. 2009;9(5):673-87.

8.   Heimark L, Shipkova P, Greene J, Munayyer H, Yarosh-Tomaine T, DiDomenico B,

et al. Mechanism of azole antifungal activity as determined by liquid chromatographic/mass

spectrometric monitoring of ergosterol biosynthesis. J Mass Spectrom. 2002;37(3):265-9.

9.   Joseph-Horne T, Hollomon DW. Molecular mechanisms of azole resistance in fungi.

FEMS Microbiol Lett. 1997;149(2):141-9.

10.   Van den Bossche H, Willemsens G, Cools W, Marichal P, Lauwers W. Hypothesis on

the molecular basis of the antifungal activity of N-substituted imidazoles and triazoles:

Portland Press Ltd.; 1983.

11.   Iacano AJ, Lewis H, Hazen JE, Andro H, Smith JD, Gulshan K. Miltefosine increases

macrophage cholesterol release and inhibits NLRP3-inflammasome assembly and IL-1β

release. Sci Rep. 2019;9(1):1-12.

12.   Kaleağasıoğlu F, Zaharieva MM, Konstantinov SM, Berger MR. Alkylphospholipids

are signal transduction modulators with potential for anticancer therapy. Anticancer Agents

Med Chem. 2019;19(1):66-91.

13.   Benaim G, Paniz-Mondolfi AE, Sordillo EM, Martinez-Sotillo N. Disruption of

intracellular calcium homeostasis as a therapeutic target against *Trypanosoma cruzi*. Front

Cell Infect Microbiol. 2020;10:46.

14.   Roatt BM, de Oliveira Cardoso JM, De Brito RCF, Coura-Vital W, de Oliveira Aguiar-

Soares RD, Reis AB. Recent advances and new strategies on leishmaniasis treatment. Appl

Microbiol Biotechnol. 2020:1-13.

15.   Papadakis MA, McPhee S, Rabow M. Current medical diagnosis & treatment New

York, NY: McGraw-Hill Education LLC; 2018.

16.   Dos Reis TF, Horta MAC, Colabardini AC, Fernandes CM, Silva LP, Bastos RW, et

al. Screening of chemical libraries for new antifungal drugs against *Aspergillus fumigatus*

reveals sphingolipids are involved in the mechanism of action of miltefosine. mBIO.

2021;12(4):e0145821. Epub 20210810. doi: 10.1128/mBio.01458-21. PubMed PMID:

34372704; PubMed Central PMCID: PMCPMC8406317.

17.   Widmer F, Wright LC, Obando D, Handke R, Ganendren R, Ellis DH, et al.

Hexadecylphosphocholine (miltefosine) has broad-spectrum fungicidal activity and is

efficacious in a mouse model of cryptococcosis. Antimicrob Agents Chemother.

2006;50(2):414-21.

18.   Rossi DCP, de Castro Spadari C, Nosanchuk JD, Taborda CP, Ishida K. Miltefosine

is fungicidal to *Paracoccidioides* spp. yeast cells but subinhibitory concentrations induce

melanisation. Int J Antimicrob Agents. 2017;49(4):465-71.

19.   Wu Y, Totten M, Memon W, Ying C, Zhang SX. In vitro antifungal susceptibility of

miltefosine alone and in combination with amphotericin B against emerging multidrug

resistant *Candida auris*. Antimicrob Agents Chemother. 2019.

20.   Barreto TL, Rossato L, de Freitas ALD, Meis JF, Lopes LB, Colombo AL, et al.

Miltefosine as an alternative strategy in the treatment of the emerging fungus *Candida auris*.

Int J Antimicrob Agents. 2020;56(2):106049.

21.   Wu Y, Wu M, Gao J, Ying C. Antifungal activity and mode of action of miltefosine

against clinical isolates of *Candida krusei*. Front Microbiol. 2020;11:854.

22.   Vila T, Ishida K, Seabra SH, Rozental S. Miltefosine inhibits *Candida albicans* and
non-*albicans Candida* spp. biofilms and impairs the dispersion of infectious cells. Int J
Antimicrob Agents. 2016;48(5):512-20.

23.   Jiménez-López JM, Carrasco MP, Marco C, Segovia JL. Hexadecylphosphocholine
disrupts cholesterol homeostasis and induces the accumulation of free cholesterol in HepG2
tumour cells. Biochem Pharmacol. 2006;71(8):1114-21. Epub 20060208. doi:
10.1016/j.bcp.2005.08.001. PubMed PMID: 16466701.

24.   Rakotomanga M, Blanc S, Gaudin K, Chaminade P, Loiseau P. Miltefosine affects
lipid metabolism in *Leishmania donovani* promastigotes. Antimicrob Agents Chemother.
2007;51(4):1425-30.

25.   Pinto-Martinez AK, Rodriguez-Durán J, Serrano-Martin X, Hernandez-Rodriguez V,
Benaim G. Mechanism of action of miltefosine on *Leishmania donovani* involves the
impairment of acidocalcisome function and the activation of the sphingosine-dependent
plasma membrane Ca2+ channel. Antimicrob Agents Chemother. 2018;62(1):e01614-17.

26.   Cuesta-Marbán Á, Botet J, Czyz O, Cacharro LM, Gajate C, Hornillos V, et al. Drug
uptake, lipid rafts, and vesicle trafficking modulate resistance to an anticancer
lysophosphatidylcholine analogue in yeast. J Biol Chem. 2013;288(12):8405-18.

27.   Pomorski T, Lombardi R, Riezman H, Devaux PF, van Meer G, Holthuis JC. Drs2p-
related P-type ATPases Dnf1p and Dnf2p are required for phospholipid translocation across
the yeast plasma membrane and serve a role in endocytosis. Mol Biol Cell. 2003;14(3):1240-
54.

28.   Hanson PK, Malone L, Birchmore JL, Nichols JW. Lem3p is essential for the uptake
and potency of alkylphosphocholine drugs, edelfosine and miltefosine. J Biol Chem.
2003;278(38):36041-50.

29.   Pérez-Victoria FJ, Castanys S, Gamarro F. *Leishmania donovani* resistance to
miltefosine involves a defective inward translocation of the drug. Antimicrob Agents
Chemother. 2003;47(8):2397-403.

30.   Pérez-Victoria FJ, Sánchez-Cañete MP, Seifert K, Croft SL, Sundar S, Castanys S,
et al. Mechanisms of experimental resistance of *Leishmania* to miltefosine: implications for
clinical use. Drug Resistance Updates. 2006;9(1-2):26-39.

31.   Seifert K, Matu S, Perez-Victoria FJ, Castanys S, Gamarro F, Croft SL.
Characterisation of *Leishmania donovani* promastigotes resistant to
hexadecylphosphocholine (miltefosine). Int J Antimicrob Agents. 2003;22(4):380-7.

32.   Mondelaers A, Sanchez-Cañete MP, Hendrickx S, Eberhardt E, Garcia-Hernandez
R, Lachaud L, et al. Genomic and molecular characterization of miltefosine resistance in
Leishmania infantum strains with either natural or acquired resistance through experimental
selection of intracellular amastigotes. PLoS One. 2016;11(4):e0154101.

33.   Srivastava A, Sircaik S, Husain F, Thomas E, Ror S, Rastogi S, et al. Distinct roles of
the 7-transmembrane receptor protein Rta3 in regulating the asymmetric distribution of
phosphatidylcholine across the plasma membrane and biofilm formation in *Candida*
albicans. Cell Microbiol. 2017;19(12). Epub 20171004. doi: 10.1111/cmi.12767. PubMed
PMID: 28745020; PubMed Central PMCID: PMCPMC5720375.
34.   Kihara A, Igarashi Y. Identification and characterization of a *Saccharomyces*
cerevisiae gene, *RSB1*, involved in sphingoid Long-chain Base release. J Biol Chem.
2002;277(33):30048-54.
35.   Kihara A, Igarashi Y. Cross talk between sphingolipids and glycerophospholipids in
the establishment of plasma membrane asymmetry. Mol Biol Cell. 2004;15(11):4949-59.
36.   Johnson SS, Hanson PK, Manoharlal R, Brice SE, Cowart LA, Moye-Rowley WS.
Regulation of yeast nutrient permease endocytosis by ATP-binding cassette transporters
and a seven-transmembrane protein, RSB1. J Biol Chem. 2010;285(46):35792-802.
37.   Soustre I, Letourneux Y, Karst F. Characterization of the *Saccharomyces cerevisiae*
RTA1 gene involved in 7-aminocholesterol resistance. Curr Genet. 1996;30(2):121-5.
38.   Rogers PD, Barker KS. Evaluation of differential gene expression in fluconazole-
susceptible and-resistant isolates of *Candida albicans* by cDNA microarray analysis.
Antimicrob Agents Chemother. 2002;46(11):3412-7.
39.   Rogers PD, Barker KS. Genome-wide expression profile analysis reveals
coordinately regulated genes associated with stepwise acquisition of azole resistance in
Candida albicans clinical isolates. Antimicrob Agents Chemother. 2003;47(4):1220-7.
40.   Coste AT, Karababa M, Ischer F, Bille J, Sanglard D. *TAC1*, transcriptional activator
of *CDR* genes, is a new transcription factor involved in the regulation of *Candida albicans*
ABC transporters *CDR1* and *CDR2*. Eukaryot Cell. 2004;3(6):1639-52.
41.   Whaley SG, Tsao S, Weber S, Zhang Q, Barker KS, Raymond M, et al. The *RTA3*
gene, encoding a putative lipid translocase, influences the susceptibility of *Candida albicans*
to fluconazole. Antimicrob Agents Chemother. 2016;60(10):6060-6.
42.   Zhai B, Ola M, Rolling T, Tosini NL, Joshowitz S, Littmann ER, et al. High-resolution
mycobiota analysis reveals dynamic intestinal translocation preceding invasive candidiasis.
Nat Med. 2020;26(1):59-64.
43.   West PT, Peters SL, Olm MR, Feiqiao BY, Gause H, Lou YC, et al. Genetic and
behavioral adaptation of *Candida parapsilosis* to the microbiome of hospitalized infants
revealed by *in situ* genomics, transcriptomics, and proteomics. Microbiome. 2021;9(1):1-17.
44.   McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, et al. The
Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA
sequencing data. Genome Res. 2010;20(9):1297-303.

45.   Lischer HE, Excoffier L, Heckel G. Ignoring heterozygous sites biases phylogenomic

estimates of divergence times: implications for the evolutionary history of Microtus voles. Mol

Biol Evol. 2014;31(4):817-31.

46.   Asadzadeh M, Dashti M, Ahmad S, Alfouzan W, Alameer A. Whole-Genome and

Targeted-Amplicon sequencing of fluconazole-susceptible and-resistant *Candida*

parapsilosis isolates from Kuwait reveals a previously undescribed N1132D polymorphism in

CDR1. Antimicrob Agents Chemother. 2021;65(2):e01633-20.

47.   Lopez-Delisle L, Rabbani L, Wolff J, Bhardwaj V, Backofen R, Grüning B, et al.

pyGenomeTracks: reproducible plots for multivariate genomic datasets. Bioinformatics.

2021;37(3):422.

48.   Carvalho CM, Zhang F, Liu P, Patel A, Sahoo T, Bacino CA, et al. Complex

rearrangements in patients with duplications of MECP2 can occur by fork stalling and

template switching. Hum Mol Genet. 2009;18(12):2188-203.

49.   Brewer BJ, Payen C, Di Rienzi SC, Higgins MM, Ong G, Dunham MJ, et al. Origin-

dependent inverted-repeat amplification: tests of a model for inverted DNA amplification.

PLoS Genet. 2015;11(12):e1005699.

50.   Müller CA, Hawkins M, Retkute R, Malla S, Wilson R, Blythe MJ, et al. The dynamics

of genome replication using deep sequencing. Nucleic Acids Res. 2014;42(1):e3-e.

51.   Ropars J, Maufrais C, Diogo D, Marcet-Houben M, Perin A, Sertour N, et al. Gene

flow contributes to diversification of the major fungal pathogen *Candida albicans*. Nat

Commun. 2018;9(1):2253. Epub 20180608. doi: 10.1038/s41467-018-04787-4. PubMed

PMID: 29884848; PubMed Central PMCID: PMCPMC5993739.

52.   Sitterlé E, Maufrais C, Sertour N, Palayret M, d'Enfert C, Bougnoux ME. Within-host

genomic diversity of *Candida albicans* in healthy carriers. Sci Rep. 2019;9(1):2563. Epub

20190222. doi: 10.1038/s41598-019-38768-4. PubMed PMID: 30796326; PubMed Central

PMCID: PMCPMC6385308.

53.   Lombardi L, Oliveira-Pacheco J, Butler G. Plasmid-based CRISPR-Cas9 gene

editing in multiple *Candida* species. mSphere. 2019;4(2):e00125-19.

54.   Krogh A, Larsson B, Von Heijne G, Sonnhammer EL. Predicting transmembrane

protein topology with a hidden Markov model: application to complete genomes. J Mol Biol.

2001;305(3):567-80.

55.   Gouy M, Guindon S, Gascuel O. SeaView version 4: a multiplatform graphical user

interface for sequence alignment and phylogenetic tree building. Mol Biol Evol.

2010;27(2):221-4.

56.   Van der Mark VA, Elferink RP, Paulusma CC. P4 ATPases: flippases in health and

disease. Int J Mol Sci. 2013;14(4):7897-922.

57.   Makuta H, Obara K, Kihara A. Loop 5 region is important for the activity of the long-
chain base transporter Rsb1. The journal of biochemistry. 2017;161(2):207-13.
58.   Fitzpatrick DA, O'Gaora P, Byrne KP, Butler G. Analysis of gene evolution and
metabolic pathways using the *Candida* Gene Order Browser. BMC Genomics. 2010;11(1):1-
14.
59.   Maguire SL, ÓhÉigeartaigh SS, Byrne KP, Schröder MS, O'Gaora P, Wolfe KH, et al.
Comparative genome analysis and gene finding in *Candida* species using CGOB. Mol Biol
Evol. 2013;30(6):1281-91.
60.   Jia X, Wang Y, Jia Y, Gao P, Xu Y, Wang L, et al. *RTA2* is involved in calcineurin-
mediated azole resistance and sphingoid long-chain base release in *Candida albicans*. Cell
Mol Life Sci. 2009;66(1):122-34.
61.   Thomas E, Sircaik S, Roman E, Brunel J-M, Johri AK, Pla J, et al. The activity of
RTA2, a downstream effector of the calcineurin pathway, is required during tunicamycin-
induced ER stress response in *Candida albicans*. FEMS Yeast Res. 2015;15(8).
62.   Yang F, Gritsenko V, Slor Futterman Y, Gao L, Zhen C, Lu H, et al. Tunicamycin
potentiates antifungal drug tolerance via aneuploidy in *Candida albicans*. mBIO.
2021;12(4):e02272-21.
63.   Sakofsky CJ, Ayyar S, Deem AK, Chung W-H, Ira G, Malkova A. Translesion
polymerases drive microhomology-mediated break-induced replication leading to complex
chromosomal rearrangements. Mol Cell. 2015;60(6):860-72.
64.   Fogel S, Welch JW. Tandem gene amplification mediates copper resistance in yeast.
Proc Natl Acad Sci (USA). 1982;79(17):5342-6.
65.   Hull RM, Cruz C, Jack CV, Houseley J. Environmental change drives accelerated
adaptation through stimulated copy number variation. PLoS Biol. 2017;15(6):e2001333.
66.   Todd RT, Selmecki A. Expandable and reversible copy number amplification drives
rapid adaptation to antifungal drugs. Elife. 2020;9:e58349.
67.   Papp C, Kocsis K, Tóth R, Bodai L, Willis JR, Ksiezopolska E, et al. Echinocandin-
induced microevolution of *Candida parapsilosis* influences virulence and abiotic stress
tolerance. mSphere. 2018;3(6):e00547-18.
68.   Ene IV, Farrer RA, Hirakawa MP, Agwamba K, Cuomo CA, Bennett RJ. Global
analysis of mutations driving microevolution of a heterozygous diploid fungal pathogen. Proc
Natl Acad Sci (USA). 2018;115(37):E8688-E97.
69.   De Coster W, D'Hert S, Schultz DT, Cruts M, Van Broeckhoven C. NanoPack:
visualizing and processing long-read sequencing data. Bioinformatics. 2018;34(15):2666-9.
70.   Jiang H, Lei R, Ding S-W, Zhu S. Skewer: a fast and accurate adapter trimmer for
next-generation sequencing paired-end reads. BMC Bioinformatics. 2014;15(1):1-12.

71.   Vaser R, Adusumalli S, Leng SN, Sikic M, Ng PC. SIFT missense predictions for

genomes. Nat Protoc. 2016;11(1):1-9.

72.   Suzek BE, Wang Y, Huang H, McGarvey PB, Wu CH, Consortium U. UniRef

clusters: a comprehensive and scalable alternative for improving sequence similarity

searches. Bioinformatics. 2015;31(6):926-32.

73.   Thorvaldsdóttir H, Robinson JT, Mesirov JP. Integrative Genomics Viewer (IGV):

high-performance genomics data visualization and exploration. Brief Bioinform.

2013;14(2):178-92.

74.   Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of

large phylogenies. Bioinformatics. 2014;30(9):1312-3.

75.   Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic

features. Bioinformatics. 2010;26(6):841-2.

76.   Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, et al. Gapped

BLAST and PSI-BLAST: a new generation of protein database search programs. Nucleic

Acids Res. 1997;25(17):3389-402.

77.   Batrakou DG, Müller CA, Wilson RHC, Nieduszynski CA. DNA copy-number

measurement of genome replication dynamics by high-throughput sequencing: the sort-seq,

sync-seq and MFA-seq family. Nat Protoc. 2020;15(3):1255-84. Epub 20200212. doi:

10.1038/s41596-019-0287-7. PubMed PMID: 32051615.

78.   Cravener MV, Mitchell AP. *Candida albicans* culture, cell Harvesting, and total RNA

extraction. Bio-protocol. 2020;10(21):e3803-e.

79.   Ahmed M, Kim DR. pcr: an R package for quality assessment, analysis and testing of

qPCR data. PeerJ. 2018;6:e4473.

Supplementary Table 1 List of strains used.

Supplementary Table 2. List of primers used

Supplementary Figure 1. Copy number determination of CNV-K repeat
Visualisations of BLASTN results using the CNV-K repeat unit plus 1 kb flanking sequence
as query against MinION reads for isolates MSK478 and MSK812 where each plot
represents hits against a single read. Each line represents a hit, and adjacent hits are
separated vertically for clarity. Read identifiers are shown on the y-axis.
A.   The exact copy number of CNV-K at both alleles was identified in isolate MSK812. (i)
Seven reads in the MSK812 MinION dataset have 8 copies of the CNV-K repeat unit, one is

shown as an example. (ii) Eighteen reads in the MSK812 dataset have 6 copies of the CNV-

K repeat unit, one is shown as an example.

B.   MSK478 has at least 11 copies on both alleles. No reads in the MSK478 dataset

covered the entirety of the repeat array, i.e. no reads had sequence matching both sides of

the query flanking DNA. The read with the highest number of copies of the CNV-K repeat

contains 11 copies, establishing a likely lower bound for copy number at both alleles.

Supplementary Figure 2. Structure of CNV-B

A.   (i) CNV-B consists of a central region (B) flanked by two regions (A and C) bounded

by inverted repeat pairs (inward-facing triangles). The CNV occurs upstream of the *RTA3*

coding sequence. (ii) CNV-B resolves as a repeat array of regions ABC interspersed with

inverted copies of region B.

B.   The ODIRA model of complex CNV generation, adapted from Brewer et al (2015)

[49] under the terms of the Creative Commons Attribution Licence. The topmost diagram has

been labelled to demonstrate relationship to the observed CNV in (A).

C.   Replication profile of strain MSK802 mapped to the *C. parapsilosis* reference

genome. Relative DNA copy number, as a proxy for replication time, is on the Y-axis, where

high values denote earlier replication. The region containing *RTA3* on chromosome 1 is

denoted by a red bar.

Supplementary Figure 3. Comparison of Rta2 and Rta3 proteins.

A.   A phylogenetic tree was generated from Muscle alignments of Rta2 and Rta3

sequences from CUG-Ser species (CGOB) using PhyML, implemented in SeaView

[55]. Bootstrap values are shown. The pink box highlights the Rta2 and Rta3 sequences

from the *Candida parapsilosis* complex. The gene names are taken from the Candida Gene

Order Browser.

B.   Alignment of *C. parapsilosis* Rta2 and Rta3 generated using Muscle implemented in

SeaView.

940