OPEN leaf: **an open-source cloud-based phenotyping system**

for tracking dynamic changes at leaf-specific resolution in

Arabidopsis.

Landon G. Swartz[1], Suxing Liu[3], Drew Dahlquist[1], Skyler Kramer[4],

Emily S. Walter[2], Sam McInturf[2], Alexander Bucksch[3], David G.

Mendoza-Cozatl[1,2,*]

[1]Department of Electrical Engineering and Computer Science, University

of Missouri-Columbia, USA

[2]Department of Plant Science and Technology, College of Agriculture,

Food, and Natural Resources, University of Missouri, Columbia, USA

[3]Department of Plant Biology, University of Georgia, Athens, GA 30602,

Warnell School of Forestry and Natural Resources, University of

Georgia, Athens, and Institute of Bioinformatics, University of Georgia,

Athens, GA 30602

[4]MU Institute of Data Science and Informatics, University of Missouri-

Columbia, USA

Short title: Cloud-based phenotyping.

Keywords: CyVerse, High-throughput phenotyping, hydroponics,

phenomics, Plant nutrition, *Arabidopsis thaliana,* Internet of

Phenotyping

Corresponding author:

D. G. Mendoza-Cozatl

Division of Plant Sciences,

University of Missouri-Columbia

1201 Rollins St, Bond-Life Sciences Center, Room 271F

Columbia, MO 65211, U.S.A.

Phone: (+1) 573-882 1892

Email: mendozad@missouri.edu

Abstract:

The first draft of the *Arabidopsis* genome was released more than 20 years

ago and despite intensive molecular research, more than 30% of

Arabidopsis genes remained uncharacterized or without an assigned

function. This is in part due to gene redundancy within gene families or

the essential nature of genes, where their deletion results in lethality (*i.e.,*

the *dark genome*).

High-throughput plant phenotyping (HTPP) offers an automated and

unbiased approach to characterize subtle or transient phenotypes resulting

from gene redundancy or inducible gene silencing; however, commercial

HTPP platforms remain unaffordable. Here we describe the design and

implementation of *OPEN leaf*, an open-source HTPP system with cloud

connectivity and remote bilateral communication to facilitate data

collection, sharing and processing.

OPEN leaf, coupled with the SMART imaging processing package was

able to consistently document and quantify dynamic morphological

changes over time at the whole rosette level and also at leaf-specific

resolution when plants experienced changes in nutrient availability.

The modular design of *OPEN leaf* allows for additional sensor integration.

Notably, our data demonstrate that VIS sensors remain underutilized and

can be used in high-throughput screens to identify characterize previously

unidentified phenotypes in a leaf-specific manner.

Significance Statement

Many bottlenecks exist in high-throughput phenotyping involving

computing power for processing and a lack of focus on abiotic stresses that

has prevented an advancement in phenotyping on par with genotyping.

Therefore, we create an automated HTP system that performs nutrient

studies on *Arabidopsis thaliana* with cloud-based image processing that

quantifies plant traits at a whole and leaf-level.

Introduction

Understanding genome-to-phenome relationships is at the core of *predictive biology*, whose aim is to predict a biological outcome from a known input (Lopatkin & Collins, 2020). The phenome, however, is also the result of specific environmental conditions and the ability of organisms to sense and adapt to changes in their environment (Des Marais *et al.*, 2013; Xu, 2016; Buckner *et al.*, 2019). Thus, *predictive biology* requires deep mechanistic understanding of the genetic makeup of a given organism but also its plastic response to environmental cues. In the case of plants, these cues may include changes in light intensity, nutrient, and water availability, and even the presence of other organisms, including microbes or additional plant species (Lundberg *et al.*, 2012; Legris *et al.*, 2016). While significant advances have been made to understand plant behavior at the molecular, physiological, and ecological level, the prediction of plant responses during changing environments remains a formidable challenge that requires consistent collection of reproducible data at the "-*omic*" and environmental levels.

Collecting large amounts of genomic and environmental data have become reasonably affordable in recent years; however, integrating, sharing, and analyzing such data in real-time is currently a bottleneck in biology. Moreover, the methods for characterizing genomes and phenomes have not advanced at the same rate (Moore *et al.*, 2013; Yang *et al.*, 2020). While genomic techniques have benefited from new developments in DNA sequencing, including higher resolution and lower costs, plant phenotyping has improved at a slower rate and commercial plant phenotyping platforms are still costly and practically inaccessible for the majority of plant laboratories (Shendure & Ji, 2008; Jackson *et al.*, 2011; White *et al.*, 2012; Reynolds *et al.*, 2019). More importantly, this gap between genomic and phenomic technologies are preventing the full use of resources available in several plant species such as mutant collections or diversity panels to pursue genome-wide studies or the characterization of complex traits at a higher resolution (Xiao *et al.*, 2017). At the same time, the technology needed for high-throughput plant phenotyping studies including sensors, computer vision, and information

technology (IT), have become more accessible, allowing the development
of diverse platforms for plant phenotyping experiments including roots
(Le Marié *et al.*, 2014; Mathieu *et al.*, 2015), shoots (Awlia *et al.*, 2016;
Flood *et al.*, 2016), plant growth within laboratory or greenhouse spaces
(van der Heijden *et al.*, 2012; Yang *et al.*, 2014) and large field crops
(Sadeghi-Tehran *et al.*, 2017; Maes & Steppe, 2019). Aside from
biological data, these platforms have also identified three major limitations
in high-throughput plant phenotyping (HTPP) platforms, that is: (i) the
ability to consistently reproduce abiotic stresses such as water and nutrient
limitation (ii) data management that is standardized and comprehensive
for collaborative analysis and re-analysis (iii) and the high cost of
commercial high-throughput phenotyping machines (Billiau *et al.*, 2012;
Tsaftaris & Scharr, 2019; Reynolds *et al.*, 2019; Yang *et al.*, 2020).

Here we describe *OPEN leaf* [Open PhENotyper for *leaf*
tracking], an open-source plant phenotyping platform designed to track
growth and development of rosette leaves in a leaf-specific manner. Our
system was built considering accessible materials worldwide and it is
primarily based on a high-resolution camera hovering over user-defined
positions to track the morphology (size, shape, color) of the whole rosette
and specific leaves. As a semi-autonomous platform, *OPEN leaf* is enabled
to communicate remotely with users on a user-defined basis to track data
in real-time, thus allowing adjustments throughout the experiments. *OPEN*
leaf was developed to be modular, scalable, and includes cloud-based
capabilities for data sharing and processing. As a proof-of-concept, here
we describe the use of *OPEN leaf* to characterize *Arabidopsis* plants grown
hydroponically with different nutrient levels. Reproducibility in the field
of plant nutrition is a major roadblock, particularly in micronutrient
deficiency studies, where the absolute concentration of elements in the
nutrient solution and their ratio with other micronutrients have profound
effects on day-to-day phenotypes (Nguyen *et al.*, 2016). With its cloud-
based capabilities enabled, *OPEN leaf* was able to resolve dynamic
phenotypes in a leaf-specific resolution during changes iron/zinc ratios.
Our results demonstrate that within the *Arabidopsis* rosette, multiple

phenotypes remain to be identified and that clear developmental changes

occur in parallel to the standard chlorosis induced by iron deficiency.

Materials and Methods

Hardware

OPEN leaf was built using the open-source T-slot aluminum

beams from 80/20 (https://8020.net) and a C-Beam Linear Actuator

Bundle from OpenBuilds, which contains the track, a gantry plate, and a

NEMA23 stepper motor (https://openbuildspartstore.com/c-beam-linear-

actuator-bundle/). A high-resolution camera (Allied Vision Mako G-503)

was mounted on the gantry plate using a bracket 3D printed on a Prusa i3

MK3 (STL file provided in Supp data; https://www.prusa3d.com). The

track system moves to pre-determined positions using GRBL, an open-

source, embedded, and high-performance g-code parser controlled by an

Arduino microcontroller (https://www.arduino.cc). For a detailed bill of

materials, 3D files, schematics and assembly manual see *Supplementary*

data.

Software

OPEN *leaf* is controlled by an open-source desktop application

written in C# (*OPEN Controller*, github.com/DMendozaLab/OPEN-

Controller), and allows multi-machine control. From the desktop

application, the user can identify and select the proper controller (machine)

and its unique camera, or other sensors associated with it. The user also

selects pre-defined camera settings including gain and aperture, the folder

where the pictures will be saved and the GRBL commands to move the

gantry plate with the camera attached to specified locations on the track.

At each specific location, the camera will take a picture at one time point

or a time series for a period of time defined by the user. Installation of

OPEN Controller is streamlined by a package (*i.e.,* installer), which

downloads and compiles all the required dependencies.

Cloud Integration

Integration for CyVerse (https://CyVerse.org) is an optional
module of the OPEN series of devices that will send data to CyVerse for
cloud storage, access, management, sharing, and analysis. CyVerse
integration is done through the desktop application Cyberduck and a
command-line interface (CLI) that interacts with CyVerse. Using CyVerse
configuration profiles, one can log into a desired CyVerse account to have
remote access to a respective CyVerse Data Store. For automated syncing
(uploading), we created a python script titled *cyverse.py* that runs a duck -
synchronize command to synchronize all data in the local directory to the
remote directory located on CyVerse.

Slack Integration

We used the Slack tool called "bot" account to relay messages
regarding the operational status of our OPEN machines. The "bot" account
is setup onto a desired channel within a Slack teamspace with the proper
permissions. Instructions to integrate Slack with OPEN machines can be
found on (github.com/DMendozaLab/OPEN-leaf-cloud). For messages,
we first query whether the HTP machine is currently running or not. If the
machine is running then a message of "Status: Running" is sent with the
current experiment name, and the latest single cycle images are sent. If the
machine is not running, then a message of "Status: Dormant" is sent. After
seven dormant messages are sent, no more dormant messages are sent until
a new experiment is started and the status is updated to "Running". The
status of the machine can be scheduled at specific times per day or "on
demand" using slash commands (*e.g.,* /srp-get-status).

Image processing using *SMART*
SMART (Speedy Measurement of Arabidopsis Rosette Traits) is a
complete set of parameter-free pipeline to analyze plant images in the
visible (VIS) and near-infrared spectrum (NIR). The SMART pipeline use
top-view images of *Arabidopsis* to compute geometrical traits of the whole
rosette and if desired, SMART can also extract information from
individual leaves in within the rosette (Table 1). For plant object

segmentation and unlike other plant image pipelines that use a threshold
value in color space, SMART uses a dominant color clustering method to
segment the plant object from the background. The pipeline used are
available as source code on GitHub (https://github.com/Computational-
Plant-Science/SMART) and are also available as a prepackaged Docker
container (https://hub.docker.com/r/computationalplantscience/smart).
For the identification and characterization of individual leaves and their
corresponding parameters, SMART adopted a watershed segmentation
algorithm. This algorithm first uses Euclidean Distance Transform to
compute the Euclidean distance map to the closest zero (*i.e.,* background
pixel) for each of the foreground pixels. Then each peak in the Euclidean
distance map was fed as the markers for watershed segmentation to
segment and count each of the plant leaves (van der Walt *et al.*, 2014). In
addition, SMART was able to compute the color distribution and
recognize distinct shades of color on the plant leaf surface.
221
Plant growth conditions

Arabidopsis Col-0 seeds were surfaced sterilized and geminated on 1/4
Murashige and Skoog (MS) medium after stratification for 2-days at 4℃
in the dark. Then, seeds were allowed to germinate under 16 h light/ 8h
dark cycles, 23°C day/ 21°C night (60 % humidity) as previously
described (Nguyen *et al.*, 2016). After the emerging of the first true leaves
(10-12 days after seed plating), plants were transferred to black magenta
boxes using 3D printed floaters (See Supp data for STL files). Unless
otherwise stated, plants were maintained in standard Arabidopsis
hydroponic media as described in Nguyen et al. (2016) and imaged every
four hours using the *OPEN leaf* system described here. Hydroponic media
were saturated with air using standard aquarium air pumps and bubbling
stones and the nutrient solution was changed every three days. After 4-6
days of continuous image capturing, the nutrient solution of some plants
was replaced with hydroponic without Fe or with hydroponic media
containing 10 μM Zn with and without Fe (the standard Zn concentration
was 1 μM Zn).

Results

OPEN leaf design

High-throughput plant phenotyping (HTPP) has revolutionized the way plant scientists gather, process and extract information from plants, including model and crop species.. However, commercial HTPP stations remain exceptionally expensive for laboratories around the world. Alternatively, several open-source HTPP devices are now available, but their assembly and operation still require a high degree of proficiency in mechanical engineering or computer sciences. To address this issue, we designed a top-view single row phenotyper (*OPEN leaf*; Figure 1A). We streamlined the assembly of *OPEN leaf* by taking advantage of affordable pre-assembled tracks available worldwide combined with controllers and open-source software (Figure 1B). At the core of *OPEN leaf* is a high-resolution camera that captures images suitable to be processed with current computer vision pipelines. In addition, we took advantage of CyVerse, a publicly accessible cyberinfrastructure, and Slack, a team communication platform, to enable cloud-storage, data sharing and processing, and a two-way communication protocol to remotely request operational status and data from *OPEN leaf* on a regularly basis or on-demand (Figure 1C).

Figure 1 *OPEN leaf* design and capabilities. (a) The main components of
the OPEN *leaf* imaging system are aluminum extrusion beans, a linear
actuator and a high-resolution VIS camera to capture top-view images of
Arabidopsis plants grown hydroponically. (b) The software and hardware
required to operate OPEN *leaf* is based on publicly available packages and
materials available worldwide. (c) To facilitate data management, we have
included optional cloud-based connectivity for remote data sharing
processing and real-time operational status updates initiated on-demand
by the end user.

OPEN leaf user interface and remote connectivity

The *OPEN leaf* user interface (*OPEN controller*) was designed so
users can easily identify the critical parts of the system (*i.e.,* camera,
storage path, and coordinates to stop along the track; Figure 2A) and have
a one-stop access to modify each of them as needed. *OPEN controller* also
allows users to capture images at one time point (single cycle) or over a
user-defined period of time (time lapse). Images are initially stored locally
in a hard drive, but we have also taken advantage of the publicly available
cloud cyberinfrastructure CyVerse and its *Discovery Environment* to
upload images regularly and facilitate data sharing and processing (scripts
available in Supplementary Data). To further engage remote access and
communication with *OPEN leaf,* we implemented a two-way
communication using Slack (Figure 2B, scripts also available in
Supplementary Data). This remote communication allows regularly
updates on the operational status of the machine together with the last set
of images taken when an experiment is ongoing. Alternatively, if the
machine has finished the required cycles or the machine has stopped
working due to a power interruption or computer malfunction the user will
receive "dormant" status at the specified time (Figure 2B-C). Operational
status can also be requested on demand with a similar outcome: "running
+ experiment + data" or "dormant" (Figure 2D-E).

Figure 2 *OPEN* leaf user interface and remote communication. (a) The

OPEN *leaf* user-interface was designed to be linear and intuitive with one-

to-one relationships to control the different settings of the imaging system.

(b-c) By enabling the communication platform Slack, users can receive

automatic updates of ongoing experiments indicating the status of the

machine and the latest acquired data. (c) In addition, users can request on-

demand updates for independent machines running simultaneously.

Feature extraction of rosette leaves using SMART

To begin testing the capabilities of *OPEN leaf* and the SMART

image processing package, we first tracked the growth of *Arabidopsis*

rosettes grown hydroponically in replete media or lacking Fe for 7 days.
In these experiments, plants were first germinated on agar plates for 7-10
days before being transferred to magenta boxes with replete nutrient
solution. After two days of acclimation in the hydroponic system, plants
were placed under the *OPEN leaf* track and imaged continuously for two
weeks. During the first week plants were grown in replete media to capture
the baseline growth and after a week, half of the plants were grown on
hydroponic media lacking Fe. While the experiment was running, images
were uploaded to CyVerse and processed using the *Speedy Measurement*
of Arabidopsis Rosette Traits (SMART) package (Figure 3A). SMART
uses the top view of the *Arabidopsis* rosette to extract several geometrical
traits of the whole rosette (Table 1). Figure 3B shows an example of the
automated rosette segmentation of plants grown in replete media or Fe
deficiency. Only 9 representative images throughout the entire experiment
are shown but all traits were calculated at each experimental time point.
Table 2 shows additional traits calculated for these 9 samples and the full
table containing all the data points and traits, including color clustering, is
available as Supplementary Table 1. Figure 3C shows the area of the entire
rosette of plants grown in the presence and absence of Fe and illustrates
how Fe deficiency has a severe negative impact on growth and
development, here measured as full rosette area. Note that the sensor used
in these experiments was a visible image sensor (*i.e.*, a RGB color
camera), As such, data collection was only possible during the light cycle
of the imposed photoperiod (shown as open blocks on the *x* axis of Figure
3C). The selection of a visible image sensor was selected to maintain an
affordable cost with the highest resolution possible and allowed us to
uncover novel phenotypes (see below); however, other sensors such as
NIR cameras are compatible with the *OPEN leaf* track system and the
SMART imaging software.

Figure 3 SMART imaging capabilities. (a) The *Speedy Measurement of Arabidopsis Rosette Traits* (SMART) is an automated pipeline to extract morphological information from Arabidopsis plants grown on the OPEN *leaf* imaging system, (b) the SMART pipeline can segment the Arabidopsis rosette and quantify several traits including leaf number, area and color [for a full list of traits see Table 1]. (c) OPEN *leaf* and SMART are capable of detecting visual changes of Arabidopsis plants grown in the presence or absence of iron in the growth media. The figure shows a representative experiment and similar trends were observed in three independent experiments.

Next, we tested whether our phenotyping platform was sensitive enough to detect changes when plants experience a combination of micronutrient availability. In these experiments, plants were grown hydroponically as described before but the nutrient challenge consisted in Fe deficiency experiments in the presence of standard zinc (Zn) concentrations (1 μM) or a mild elevated yet non-lethal concentration of Zn (10 μM) in the hydroponic solution. Figure 4A shows representative rosette images of each condition extracted using the SMART package and Figure 4B shows the area of the whole rosette plotted against time. Both, the images and the corresponding rosette, were able to capture differences in growth and development in each of the growth conditions. Notably, the addition of 10 μM Zn to the hydroponic media resulted in a smaller rosette area compared to plants grown in 1 μM Zn grown with replete Fe (1 μM Zn+Fe) or exposed to Fe deficiency (1 μM Zn-Fe). Moreover, the color clustering algorithm revealed clear differences in color distribution across treatments, a trait that was found to be independent of rosette size and therefore should be extracted and analyzed in parallel. That is, the rosette area of plants grown in 1 μM Zn-Fe was larger than plants grown in 10 μM Zn+Fe, but the color distribution reported as hexadecimal values was consistent with chlorosis only in plants grown without Fe (Figure 4B).

Figure 4 Arabidopsis phenotyping during contrasting changes in nutrient
availability. (a) Whole rosette area and color are affected when the ratio
of iron (Fe) and zinc (Zn) are modified in the hydroponic media. (b) the
OPEN leaf-SMART pipeline was able to detect subtle changes in rosette
area and color when the Fe/Zn ratio was modified. (c) Changes in Fe and
Zn lead to unique phenotypes in individual leaves within the entire rosette
in a pattern consistent with sink-source relationships. These experiments
were repeated in triplicate with similar results.

SMART leaf-specific phenotyping

During the course of these experiments, we noticed that within
treatments, the color of individual leaves within the *Arabidopsis* rosette
changed substantially, in a manner consistent with source-sink

relationships (Figure 4C). While these Fe-associated leaf-specific
phenotypes have been described at the molecular level (Schuler *et al.*,
2012), segmentation and quantification of visible traits in a leaf-specific
manner is not standard in current phenotyping pipelines. Hence, we
implemented a watershed segmentation algorithm to segment individual
leaves within the *Arabidopsis* rosette (Figure 5). This watershed
segmentation algorithm first uses *Euclidean Distance Transform* to
calculate the Euclidean distance map to the closest zero (*i.e.*, background
pixel) for each of the foreground pixels. In the following step, each peak
in the Euclidean distance map was used as a marker for the watershed
segmentation to identify and count each of the leaves within the rosette
(Figure 5B). Once individual leaves were isolated, all different traits
described for the full rosette were extracted including color using the
dominant clustering method described before (Table 1; Supplementary
Table 2, and Figure 5C-D).

Figure 5 Leaf-specific segmentation of Arabidopsis rosette using
SMART. (a-b) SMART is capable of segmenting individual leaves
through watershed segmentation. (c-d) color is described as the top four
dominant colors using hexadecimal values, and once the individual leaves
have been identified and extracted, all the morphological traits described
in Table 1 can be quantified in a leaf-specific manner.

To test the capabilities of our leaf-specific phenotyping approach,

we compared plants grown in standard Zn levels (1 μM) plus Fe (1 μM

Zn+Fe) with plants grown in 10 μM Zn without Fe (10 μM Zn-Fe). We

chose these two conditions as starting point based on the data obtained

using whole rosette phenotyping (Figure 4), where color changes in a leaf-

specific manner were consistently observed. In turn, our individual leaf

segmentation approach allowed us to classify individual rosette leaves in

three main clusters (Figure 6). Cluster 1 represents the first true leaves

within the rosette, whose area remain constant through the progression of

the experiment; however, and as excepted for a *source* leaf, these leaves

receive and retain the majority of micronutrients coming through the

xylem stream and dictate the distribution of micronutrient across the

rosette in a pattern consistent with their orthostichy (Khan *et al.*, 2018).

Despite minimal changes in individual leaf area, Cluster 1 leaves show

clear differences in color, and under the experimental conditions imposed,

leaves from plants grown with 10 μM Zn-Fe display hexadecimal color

values consistent with darker green compared to the same leaves from

plants grown in 1 μM Zn+Fe (Figure 6B and C). In contrast, Cluster 2

leaves represent individual leaves transitioning from sink (developing

leaves) to source leaves (fully photosynthetic). In this cluster, the leaf area

and color became more dependent on the environmental conditions (*i.e.*,

nutrient availability) and their unique and autonomous developmental

program. For instance, in 1 μM Zn+Fe, the leaf area of Cluster 2 leaves

increased over time and their color remain constant through the entire

experiment. However, growth of Cluster 2 leaves from plants grown in 10

μM Zn-Fe was variable, with some leaves being larger than plants grown

with 1 μM Zn+Fe, while others remain smaller (Figure 6B). Moreover,

capturing color throughout the experiment allowed us to observe and

quantify the transition of Cluster 2 leaves from dark green to yellow and

this transition was consistent with the availability of high Zn and Fe at the

beginning of the experiment to lack of Fe in the later days of the

experiment. (Figure 6C). Lastly, Cluster 3 leaves represent mostly *sink*

leaves; thus, their appearance, color and size were more dependent on the

nutrient availability. For instance, Cluster 3 leaves appearance was

delayed in plants grown in 10 µM Zn-Fe compared to plants grown in 1

µM Zn+Fe (Figure 6A-B). Moreover, Cluster 3 leaves from plants grown

in 10 µM Zn+Fe were more chlorotic compared to leaves from plants

grown in 1 µM Zn+Fe (Figure 6C). Altogether, these results demonstrate

that *OPEN leaf* and SMART were able to capture the dynamic behavior of

individual leaves within the *Arabidopsis* rosette when plants were exposed

to different levels of nutrient availability.

Figure 6 Nutrient availability dictates the fate of individual leaves within

the Arabidopsis rosette. (a) the *OPEN leaf* – SMART pipeline is able to

track individual leaves during plant growth and development. (b) Changes

in the Fe/Zn ratio have a severe impact on individual leaf size and color

and leaf clustering was consistent with the leaf developmental stage

(source or sink). (c) Individual color clustering of the 4 dominant colors
allowed the dynamic tracking of color transitioning during the entire
experiment a at leaf-specific resolution. Each bar plot represents the
proportional distribution of each dominant color in individual leaves. The
plots are representative of one experiment and similar results were
obtained in three independent experiments.

Discussion

The Arabidopsis genome was released in the year 2000 and after
more than two decades of intensive molecular research, 30% of its 25,000+
genes remain without an assigned function and a greater number have been
annotated based on homology without experimental evidence of their
putative function (Wang *et al.*, 2019). There are many factors behind this
gap in knowledge, including gene lethality or redundancy within gene
families, which is not uncommon in higher plants. In addition, many
mutations lead to subtle or transient phenotypes which are difficult to
reproduce and capture with standard end-point documentation methods
(*i.e.,* photos at the end of an experiment) or subject to experimental bias
from environmental conditions or image processing protocols such as
manual extraction of leaf traits including shape, size and color (Dobrescu
et al., 2017; Zhang *et al.*, 2020; Hernández-Herrera *et al.*, 2021).

High throughput plant phenotyping (HTPP) has emerged as a
technology to solve many of these issues including consistency during data
acquisition and processing (Acosta-Gamboa *et al.*, 2016; Vasseur *et al.*,
2018) HTPP has facilitated the discovery of novel genes and gene function
through the use of large mutant collections and diversity panels or
ecotypes (Campos *et al.*, 2021; Xie *et al.*, 2021). Unfortunately,
commercial HTPP devices remain prohibitively expensive and thus are
limited to the very few labs that can afford them. Many open-source HTPP
devices have also been described but their assembly and operation require
significant expertise in mechanical engineering and computer science,
which is not standard nor common in plant biology departments. However,
a full and comprehensive characterization of the *dark genome* will require
a collective effort within the plant community and at the core of this efforts

is accessibility to HTPP technologies and the use of standard practices to
conduct experiments under well-defined environmental conditions. This
Internet of Phenotyping (IoP) in which creating an integrated network of
HTPP technologies, user-interactions, and image processing can greatly
accelerate development in HTPP.

Here we described *OPEN leaf*, an open-source HTPP device with cloud
connectivity and remote bilateral communication with users (Figures 1
and 2). *OPEN leaf* design is user-centered and takes advantage of
affordable materials available worldwide, with an approximate cost of
$1000 USD (as of Dec 2021; Manual assembly available in supplementary
data). *OPEN leaf* together with the SMART processing suite offers a one-
stop solution for HTPP of *Arabidopsis* rosettes, from continuous and
automatic data collection to data storage, sharing and processing. The
cloud integration solves one of the main issues in HTP approaches known
as the FAIR principle for data management, where HTP data should be
ideally findable, available, identifiable, and reusable for reasonable
scientific advancement (Wilkinson *et al.*, 2016). The modular structure of
OPEN leaf allows users to capture images at a user-defined frequency
without the need for cloud or internet connectivity. This modularity also
allows users to incorporate additional sensors such as NIR, fluorescence,
multi- and hyperspectral cameras; however, each of these sensors will
proportionally increase the cost of the HTPP device and data output.

In this work, we also aimed to demonstrate that despite being one
of the most basic sensors, visible cameras and RGB images, are still
underutilized and several novel phenotypes occurring within the rosette of
Arabidopsis are yet to be identified, quantified, and characterized. For
instance, rosette growth and development is known to be controlled by the
intrinsic genetic program of the plant but can also be modified by
environmental factors such as nutrient availability or the presence of
pathogens (Hindt & Guerinot, 2012; Kim *et al.*, 2014). In turn, our whole-
rosette phenotyping pipeline was able to document the growth delay of
plants experiencing Fe deficiency (Figure 3). Growth delay in Arabidopsis
roots has been found to be mediated by the gibberellin pathway and the
corresponding DELLA repressor proteins (Wild *et al.*, 2016). For instance,

Arabidopsis plants lacking all DELLA proteins have longer roots during
Fe deficiency compared to wildtype plants, suggesting that growth arrest
during Fe deficiency is the result of proper Fe sensing followed by
downstream hormone-developmental events but not the result of energy
shortages due to impaired photosynthesis and respiration. Notably, Fe
sensing at the whole plant level is sensed in source leaves, which
communicate the Fe status of the plant to roots *via* the phloem to either
increase or repressed Fe uptake at the root level (Khan *et al.*, 2018). It is
therefore expected that a similar hormone-developmental signaling
pathway associated with Fe sensing in leaves takes place; however, the
molecular mechanisms behind Fe sensing in leaves remain largely
unknown. In turn, our phenotyping system was capable of providing
detailed dynamic information about the timing of Fe-deficiency responses
in leaves thus offering researchers the opportunity to select ideal time
windows for further in-depth molecular analyses (Figure 3).

Hydroponic experiments have provided a unique opportunity to
explore plant responses to nutrient availability at a very high resolution
and our *OPEN leaf*-SMART pipeline captured subtle differences in rosette
size and color during experiments were Fe and Zn availabilities were
modified (Figure 4). Moreover, our phenotyping pipeline demonstrates
that significant changes occur within the *Arabidopsis* rosette, in a leaf-
specific manner, when the ratio of Fe/Zn availability changes (Figure 5).
Leaf responses to nutrient availability are expected to be striking different
from roots as the *Arabidopsis* rosette includes both sink and source tissues
(*i.e.,* young and mature leaves), while roots are by definition a sink tissue.
In fact, unique leaf-specific transcriptional responses have been described
during Fe deficiency experiments in *Arabidopsis* where prolonged Fe
deprivation leads to a restricted distribution of Fe to developing (sink)
leaves to preserve optimal photosynthesis in mature (source) leaves
(Hantzis *et al.*, 2018; McInturf *et al.*, 2021).

While the precise molecular mechanisms behind Fe economy
reprograming is ongoing, our *OPEN leaf*-SMART phenotyping pipeline
was able to capture the timing of such leaf-specific changes, thus offering
the sought-after time window to pursue additional molecular analyses to

unravel the mechanisms behind plant responses and adaptation to nutrient

availability in a leaf-specific manner.

Acknowledgments

This research was supported in part by the US National Science Foundation (MCB-1818312 and IOS- 1734145 to DMC), the MU Honors College and Peggy and Andrew Cherng Charitable Foundation (LS), the USDOE ARPA-E ROOTS (DE-AR0000821), the NSF CAREER Award No. 1845760 to A.B, and the US National Institute of Health BD2K Training Grant T32HG009060 grant (STK). We would also like to thank Wesley Paul Bonelli for his contribution to design the script collection data Any opinions findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect those of ARPA-E or the National Science Foundation.

Author Contribution

LGS, SM and DM conceived the rpoject, LGD and SM designed HTPP device. EW performed plant experiments on the system. SL and AB developed the SMART imaging pipeline. LGS, SK and DM worked on data interpretation and visualization. LS and DM wrote the manuscript.

Data Availability

The data that supports the findings of this study are available in the supplementary material of this article

575 **Table 1** Summary of plant traits measurement and units

Measurement	Units	Trait type*	Description
Whole plant area	pixels	WP	Sum of the whole plant area
Whole plant width/height	pixels	WP	The width/height of the plant object, computed from major/minor axis values of the fitted bounding ellipse.
Whole plant solidity	None	WP	The ratio of fitted whole plant contour area to its convex hull area, describe the shape of the whole plant, A value of 1 signifies a solid object and a value less than 1 will signify an object having an irregular boundary.
Whole plant curvature	None	WP	The derivative of the inclination of the tangent with respect to arc length. The mean curvature value of all leaves, describing the boundary shape change of individual leaf.
Number of leaves	None	WP	Count of the total number of leaves
Color distribution	None	LS	The distribution and percentage of distinct shades of color on the plant leaf surface.
Leaf area	pixels	LS	Individual leaf area
Leaf width/height	pixels	LS	The width/height of the individual leaf object, computed from major/minor axis values of the fitted bounding ellipse.
Leaf solidity	None	LS	The ratio of fitted leaf contour area to its convex hull area.

576　　　* WP, whole plant; LS, Leaf specific.

579 **Table 2** Main traits extracted from Arabidopsis plants grown

580 hydroponically shown in Figure 3B.

Time (min)	Area		Solidity		Max_width		Max_height		Curvature		Leaf number	
	Full	-Fe	Full	-Fe	Full	-Fe	Full	-Fe	Full	-Fe	Full	-Fe
72	29.41	33.10	0.571	0.611	211	288	302	269	0.026	0.022	6	5
102	43.19	51.08	0.547	0.588	285	371	370	324	0.022	0.020	7	6
126	58.25	72.87	0.520	0.552	360	463	416	426	0.017	0.018	7	7
168	104.2	109.1	0.478	0.488	626	539	544	636	0.017	0.019	9	10
189	123.6	129.1	0.484	0.512	718	538	576	658	0.016	0.017	10	11
213	143.7	156.0	0.482	0.493	808	666	596	731	0.017	0.016	11	12
237	161.9	170.2	0.542	0.432	651	818	620	792	0.012	0.014	8	11
261	237.9	200.8	0.509	0.436	963	891	782	816	0.013	0.014	11	12
295	290.8	257.3	0.454	0.431	1089	982	1001	936	0.014	0.013	11	10

*See Table 1 for units.

Supplementary data

Manual assembly

STL files (floaters and camera assembly)

Supplementary Table 1 Rosette traits extracted using SMART.

Supplementary Table 2 Leaf-specific traits extracted using SMART.

References

Acosta-Gamboa L, Liu S, Langley E, Campbell Z, Guerrero N, Mendoza Cozatl D, Lorence A. **2016**. Moderate to severe water limitation differentially affects the phenome and ionome of Arabidopsis. *Functional Plant Biology* **44**.

Awlia M, Nigro A, Fajkus J, Schmoeckel SM, Negrão S, Santelia D, Trtílek M, Tester M, Julkowska MM, Panzarová K. **2016**. High-Throughput Non-destructive Phenotyping of Traits that Contribute to Salinity Tolerance in Arabidopsis thaliana. *Frontiers in Plant Science* **7**: 1414.

Billiau K, Sprenger H, Schudoma C, Walther D, Köhl KI. **2012**. Data management pipeline for plant phenotyping in a multisite project. *Functional Plant Biology* **39**: 948–957.

Buckner E, Madison I, Chou H, Matthiadis A, Melvin CE, Sozzani R, Williams C, Long TA. **2019**. Automated Imaging, Tracking, and Analytics Pipeline for Differentiating Environmental Effects on Root Meristematic Cell Division. *Frontiers in Plant Science* **10**: 1487.

Campos ACAL, van Dijk WFA, Ramakrishna P, Giles T, Korte P, Douglas A, Smith P, Salt DE. **2021**. 1,135 ionomes reveal the global pattern of leaf and seed mineral nutrient and trace element diversity in Arabidopsis thaliana. *The Plant Journal* **106**: 536–554.

Des Marais DL, Hernandez KM, Juenger TE. **2013**. Genotype-by-Environment Interaction and Plasticity: Exploring Genomic Responses of Plants to the Abiotic Environment. *Annual Review of Ecology, Evolution, and Systematics* **44**: 5–29.

Dobrescu A, Scorza LCT, Tsaftaris SA, McCormick AJ. **2017**. A "Do-It-Yourself" phenotyping system: measuring growth and morphology throughout the diel cycle in rosette shaped plants. *Plant Methods* **13**: 95.

Flood PJ, Kruijer W, Schnabel SK, van der Schoor R, Jalink H, Snel JFH, Harbinson J, Aarts MGM. **2016**. Phenomics for photosynthesis, growth and reflectance in Arabidopsis thaliana reveals circadian and long-term fluctuations in heritability. *Plant methods* **12**: 14–14.

Hantzis LJ, Kroh GE, Jahn CE, Cantrell M, Peers G, Pilon M, Ravet K. **2018**. A Program for Iron Economy during Deficiency Targets Specific Fe Proteins. *Plant Physiology* **176**: 596–610.

van der Heijden G, Song Y, Horgan G, Polder G, Dieleman A, Bink M, Palloix A, van Eeuwijk F, Glasbey C. **2012**. SPICY: towards automated phenotyping of large pepper plants in the greenhouse. *Functional Plant Biology* **39**: 870–877.

Hernández-Herrera P, Ugartechea-Chirino Y, Torres-Martínez HH, Arzola AV, Chairez-Veloz JE, García-Ponce B, Sánchez M de la P, Garay-Arroyo A, Álvarez-Buylla ER, Dubrovsky JG, *et al.* 2021. Live Plant Cell Tracking: Fiji plugin to analyze cell proliferation dynamics and understand morphogenesis. *Plant Physiology*: kiab530.

Hindt MN, Guerinot ML. **2012**. Getting a sense for signals: regulation of the plant iron deficiency response. *Biochimica et biophysica acta* **1823**: 1521–1530.

Jackson SA, Iwata A, Lee S-H, Schmutz J, Shoemaker R. **2011**. Sequencing crop genomes: approaches and applications. *The New phytologist* **191**: 915–925.

Khan MA, Castro-Guerrero NA, McInturf SA, Nguyen NT, Dame AN, Wang J, Bindbeutel RK, Joshi T, Jurisson SS, Nusinow DA, *et al.* 2018. Changes in iron availability in Arabidopsis are rapidly sensed in the leaf vasculature and impaired sensing leads to opposite transcriptional programs in leaves and roots. *Plant, Cell & Environment* **41**: 2263–2276.

Kim SH, Son GH, Bhattacharjee S, Kim HJ, Nam JC, Nguyen PDT, Hong JC, Gassmann W. **2014**. The Arabidopsis immune adaptor SRFR1 interacts with TCP transcription factors that redundantly contribute to effector-triggered immunity. *The Plant Journal* **78**: 978–989.

Le Marié C, Kirchgessner N, Marschall D, Walter A, Hund A. **2014**. Rhizoslides: paper-based growth system for non-destructive, high throughput phenotyping of root development by means of image analysis. *Plant Methods* **10**: 13.

Legris M, Klose C, Burgie ES, Rojas CCR, Neme M, Hiltbrunner A, Wigge PA, Schäfer E, Vierstra RD, Casal JJ. **2016**. Phytochrome B integrates light and temperature signals in Arabidopsis. *Science* **354**: 897.

Lopatkin AJ, Collins JJ. **2020**. Predictive biology: modelling, understanding and harnessing microbial complexity. *Nature Reviews Microbiology* **18**: 507–520.

Lundberg DS, Lebeis SL, Paredes SH, Yourstone S, Gehring J, Malfatti S, Tremblay J, Engelbrektson A, Kunin V, Del Rio TG, *et al.* 2012. Defining the core Arabidopsis thaliana root microbiome. *Nature* **488**: 86–90.

Maes WH, Steppe K. **2019**. Perspectives for Remote Sensing with Unmanned Aerial Vehicles in Precision Agriculture. *Trends in Plant Science* **24**: 152–164.

Mathieu L, Lobet G, Tocquin P, Périlleux C. **2015**. "Rhizoponics": a novel hydroponic rhizotron for root system analyses on mature Arabidopsis thaliana plants. *Plant Methods* **11**: 3.

McInturf SA, Khan MA, Gokul A, Castro-Guerrero NA, Höhner R, Li J, Marjault H-B, Fichman Y, Kunz H-H, Goggin FL, *et al.* 2021. Cadmium interference with iron sensing reveals transcriptional programs sensitive and insensitive to reactive oxygen species. *Journal of Experimental Botany*: erab393.

Moore C, Johnson L, Kwak I-Y, Livny M, Broman K, Spalding E. **2013**. High-Throughput Computer Vision Introduces the Time Axis to a Quantitative Trait Map of a Plant Growth Response. *Genetics* **195**.

Nguyen NT, McInturf SA, Mendoza-Cózatl DG. **2016**. Hydroponics: A Versatile System to Study Nutrient Allocation and Plant Responses to Nutrient Availability and Exposure to Toxic Elements. *Journal of visualized experiments : JoVE*: 54317.

Reynolds D, Baret F, Welcker C, Bostrom A, Ball J, Cellini F, Lorence A, Chawade A, Khafif M, Noshita K, *et al.* 2019. What is cost-efficient

phenotyping? Optimizing costs for different scenarios. *The 4th International*
Plant Phenotyping Symposium **282**: 14–22.

Sadeghi-Tehran P, Sabermanesh K, Virlet N, Hawkesford MJ. **2017**.
Automated Method to Determine Two Critical Growth Stages of Wheat:
Heading and Flowering. *Frontiers in Plant Science* **8**: 252.

Schuler M, Rellán-Álvarez R, Fink-Straube C, Abadía J, Bauer P. **2012**.
Nicotianamine functions in the Phloem-based transport of iron to sink organs, in
pollen development and pollen tube growth in Arabidopsis. *The Plant cell* **24**:
2380–2400.

Shendure J, Ji H. **2008**. Next-generation DNA sequencing. *Nature*
Biotechnology **26**: 1135–1145.

Tsaftaris SA, Scharr H. **2019**. Sharing the Right Data Right: A Symbiosis with
Machine Learning. *Trends in Plant Science* **24**: 99–102.

Vasseur F, Bresson J, Wang G, Schwab R, Weigel D. **2018**. Image-based
methods for phenotyping growth dynamics and fitness components in
Arabidopsis thaliana. *Plant Methods* **14**: 63.

van der Walt S, Schönberger JL, Nunez-Iglesias J, Boulogne F, Warner JD,
Yager N, Gouillart E, Yu T, scikit-image contributors. **2014**. scikit-image:
image processing in Python. *PeerJ* **2**: e453–e453.

Wang W, Zhang X, Niittylä T. **2019**. OPENER Is a Nuclear Envelope and
Mitochondria Localized Protein Required for Cell Cycle Progression in
Arabidopsis. *The Plant cell* **31**: 1446–1465.

White JW, Andrade-Sanchez P, Gore MA, Bronson KF, Coffelt TA, Conley
MM, Feldmann KA, French AN, Heun JT, Hunsaker DJ, *et al.* **2012**. Field-
based phenomics for plant genetics research. *Field Crops Research* **133**: 101–
112.

Wild M, Davière J-M, Regnault T, Sakvarelidze-Achard L, Carrera E,
Lopez Diaz I, Cayrel A, Dubeaux G, Vert G, Achard P. **2016**. Tissue-
Specific Regulation of Gibberellin Signaling Fine-Tunes Arabidopsis Iron-
Deficiency Responses. *Developmental Cell* **37**: 190–200.

Wilkinson MD, Dumontier M, Aalbersberg IjJ, Appleton G, Axton M,
Baak A, Blomberg N, Boiten J-W, da Silva Santos LB, Bourne PE, *et al.*
2016. The FAIR Guiding Principles for scientific data management and
stewardship. *Scientific Data* **3**: 160018.

Xiao Y, Liu H, Wu L, Warburton M, Yan J. **2017**. Genome-wide Association
Studies in Maize: Praise and Stargaze. *Molecular plant* **10**: 359–374.

Xie J, Fernandes SB, Mayfield-Jones D, Erice G, Choi M, E Lipka A,
Leakey ADB. **2021**. Optical topometry and machine learning to rapidly
phenotype stomatal patterning traits for maize QTL mapping. *Plant Physiology*
187: 1462–1480.

Xu Y. **2016**. Envirotyping for deciphering environmental impacts on crop
plants. *Theoretical and Applied Genetics* **129**: 653–673.

Yang W, Feng H, Zhang X, Zhang J, Doonan JH, Batchelor WD, Xiong L,
Yan J. **2020**. Crop Phenomics and High-Throughput Phenotyping: Past
Decades, Current Challenges, and Future Perspectives. *Molecular Plant* **13**:
187–214.

Yang W, Guo Z, Huang C, Duan L, Chen G, Jiang N, Fang W, Feng H, Xie
W, Lian X, *et al.* **2014**. Combining high-throughput phenotyping and genome-
wide association studies to reveal natural genetic variation in rice. *Nature*
Communications **5**: 5087.

Zhang X, Hu Z, Guo Y, Shan X, Li X, Lin J. **2020**. High-efficiency procedure
to characterize, segment, and quantify complex multicellularity in raw
micrographs in plants. *Plant Methods* **16**: 100.

728