1

Antibiotic Resistance Genes and Taxa Analysis from Mat and Planktonic Microbiomes of

Antarctic Perennial Ice-covered Lake Fryxell and Lake Bonney

By Sheetal Tallada,[1][†] Grant Hall,[1][†] Daniel Barich,[1] Rachael M. Morgan-Kiss,[2] and Joan L.

Slonczewski[1][*]

Address:

[1]Department of Biology, Kenyon College, Gambier, Ohio, 43022, USA

[2]Department of Microbiology, Miami University, Oxford, Ohio, 45056, USA

[†]These two authors contributed equally.

*Corresponding Author:

Joan L. Slonczewski, slonczewski@kenyon.edu

ABSTRACT

The perennial ice-covered lakes of the Antarctic McMurdo Dry Valleys harbor oligotrophic

microbial communities that are separated geographically from other aquatic systems. Their

microbiomes include planktonic microbes as well as lift-off mat communities that emerge from

the ice. We used ShortBRED to quantify the antibiotic resistance genes (ARGs) from

metagenomes of lift-off mats emerging from ice, and from filtered water samples of Lake Fryxell

and Lake Bonney. The overall proportion of ARG hits was similar to that found in temperate-

zone rural water bodies with moderate human inputs. Specific ARGs showed distinct

distributions for the two lakes, and for mat versus planktonic sources. Metagenomic taxa

distributions showed that mat phototrophs consisted mainly of cyanobacteria or

betaproteobacteria, whereas the water column phototrophs were mainly protists. An enrichment

culture of the betaproteobacterium *Rhodoferax antarcticus* from a Lake Fryxell mat sample

showed an unusual mat-forming phenotype not previously reported for this species. Its genome

showed no ARGs associated with betaproteobacteria, but had ARGs consistent with a minor

Pseudomonas component. The Antarctic lake mats and water showed specific ARGs distinctive

to the mat and water sources, but overall ARG levels were similar to those of temperate water

bodies with moderate human inputs.

INTRODUCTION

One of Earth's coldest dry deserts, the McMurdo Dry Valleys in Antarctica possess a string of stratified lakes with unique geochemistry and microbial communities (Spigel & Priscu 1998, Roberts *et al.* 2004, Cavicchioli 2015, Sohm *et al.* 2020). The microbial genomic analysis of these lakes remains limited (Dillon *et al.* 2020, Koo *et al.* 2018, Wei Li *et al.* 2020). There are a number of reports on antibiotic resistance genes (ARGs) in Antarctic soil or marine bacteria (Antelo *et al.* 2021, Tam *et al.* 2015, Wang *et al.* 2016, Na *et al.* 2019, van Goethem *et al.* 2018) but few studies of ARGs in Antarctic lake water or their mat communities (Jara *et al.* 2020). The prevalence of ARGs in relatively pristine habitats is of compelling importance for human health and environment, as it provides a baseline for assessment of human-associated ARG contamination (Allen *et al.* 2010, D'Costa *et al.* 2011). Antimicrobial resistance is a leading cause of death worldwide (Murray *et al.* 2022).

We report on the taxonomic composition and ARG prevalence in the water of Lake Fryxell and Lake Bonney, and in the phototrophic mat communities of Lake Fryxell (**Fig. 1A**). Lake Fryxell extends along the Taylor Valley, 10 km downstream of Lake Bonney, 80 km across the Ross Sea from McMurdo Station. No permanent stream connects the two lakes; their main hydrology involves glacial meltwater, and each lake has an underground brine aquifer (Mikucki *et al.* 2015). Their water ecosystems are entirely microscopic, with no fauna larger than nematodes. The lakes and their ecosystems are studied as models for ancient Mars (Head & Marchant 2014) and as indicators of climate change (Hall *et al.* 2017).

Lakes Fryxell and Bonney have permanent stratified layers in the water column that give rise to oligotrophic microbial communities (Roberts *et al.* 2004, Li *et al.* 2016, Kwon *et al.*

2017). The lack of turnover allows for the growth of benthic microbial mats that form flame-

shaped towers several centimeters tall (Jungblut *et al.* 2016, Hawes *et al.* 2013). The microbial mats

have been studied with primary emphasis on the role of cyanobacteria such as *Microcoleus* and

Oscillatoria (Taton *et al.* 2003, Jungblut *et al.* 2016) although metagenomes reveal a high

abundance of proteobacteria (Dillon *et al.* 2020) and protists (Li *et al.* 2016, Bielewicz *et al.* 2011).

Known as "lift-off" mats, the benthic biofilms reach upward, buoyed by oxygen bubbles, until

pieces break off and float upward to the undersurface of the ice cover (Parker *et al.* 1982,

Moorehead *et al.* 1999). During winter, water freezes beneath the mat fragments, adding to the ice

layer, but above the lake the valley's dry winds ablate the ice. Ice sections reveal 6-8 annual

layers of freezing below followed by ablation above (Parker *et al.* 1982) leading to exposure of the

dessicated mats (**Fig. 1B**). Surviving several years trapped in ice, the mat material remains

viable, and psychrophiles may continue growing (Boetius *et al.* 2015). When mat organisms reach

the surface of ablated ice, they can blow off and enter "moats" of melted ice that occur in

summer, surrounding the permanent ice in the main part of the lake. The mat material carries

communities that include microscopic arthropods and nematodes; thus, the mats can act as

vectors for transport of entire ecosystems (Parker *et al.* 1982, Brambilla *et al.* 2001, Dillon *et al.* 2020).

Molecular genetic data on McMurdo Dry Valley microbes remained limited until recently

(Taton *et al.* 2003, Vick-Majors *et al.* 2014, Kwon *et al.* 2017). In December, 2014, the austral

summer, we obtained water samples from Fryxell and Bonney as well as desiccated mat samples

emerging from the ice cover of Fryxell. For the present study, we applied tools of metagenome

analysis to compare the taxonomic diversity and ARG abundance of mat and lake samples. We

also sequenced the genome of a novel ecotype of *Rhoderax antarcticus* from a mat-forming

enrichment culture, whose phenotype differs markedly from planktonic isolates of this species

(Madigan *et al.* 2000, Baker *et al.* 2017, Jung *et al.* 2004).

Our study addressed the important question of ARG prevalence in relatively pristine

Antarctic lakes. A modest level of antibiotic resistance is an ancient, widespread phenomenon

naturally and historically occurring in all environments (Allen *et al.* 2010, D'Costa *et al.* 2011). Non-

anthropogenic processes can select for ARGs in pristine habitats; for example, cyanobacterial

blooms drive increases in bacterial ARG prevalence and diversity (Zhang *et al.* 2020). But inputs

of human origin can add substantially to the native ARG pool as well as add additional types of

ARGs (Antelo *et al.* 2021, Tam *et al.* 2015, Jara *et al.* 2020). We surveyed our mat and water

metagenomes for the presence of ARGs, referenced to the Comprehensive Antibiotic Resistance

Database (CARD) (Alcock *et al.* 2020).

METHODS

Sample collection and culture.

Microbial communities were sampled in December, 2014 from two meromictic lakes of the

Taylor Valley, Victoria Land, Antarctica. Lake Fryxell has a maximum depth of 20 m (Kwon *et*

al. 2017, Lawrence & Hendy 1985), while Lake Bonney has a depth of 40 m (Priscu & Spigel 1996).

Both lakes are covered by a perpetual ice layer, approximately 4 m thick (Priscu 2018), although

summer melting occurs near the shoreline.

Microbial lift-off mat samples were collected from independent mat tufts emerging

separately from the Lake Fryxell ice surface, within the GPS area of: -77.60491, 163.16315; -

77.60473, 163.16290; -77.60463, 163.16405; -77.60495, 163.16495. Each sample consisted of a

separate tuft of dessicated microbial mat, collected with alcohol-sterilized forceps and stored at

-20°C (4 weeks) then at -80°C (indefinitely).

Lake water was sampled from the permanent chemoclines of Lake Fryxell (-77.605,

163.163, 9 m depth) and Lake Bonney, east lobe (-77.719, 162.283, 15 m depth). Samples were

obtained using ice holes established by the McMurdo Dry Valleys Long-Term Ecological

Research (LTER) program (Priscu 2022b, Priscu 2022a, Priscu 2021b, Priscu 2021a). The LTER

geochemical data are presented in Table 1. Water samples were collected in 1-liter cubitainers

pre-washed in 10% HCl. Each water sample was concentrated by filtration through a 0.45 µm

filter.

Enrichment culture for anaerobic phototrophs was performed using Harwood PM

medium supplemented with 10 mM succinate (Fixen *et al.* 2019, Kim & Harwood 1991, Rey *et al.*

2006). Screwcap pyrex tubes were filled with medium and inoculated with approximately 0.05 g

dessicated material from sample Mat-04. Sealed tubes were incubated under approximately 10%

PAR at 10°C for five weeks. Portions of biofilm were serially subcultured for two-week periods,

then frozen at -80°C. Gram stain was performed by standard methods (Remel kit).

Testing for growth range of pH and NaCl amendment was performed using a *Rhodoferax*

medium modified from references (Tayeh & Madigan 1987, Madigan *et al.* 2000). The medium

contained: yeast extract (0.5 g/l), EDTA (20 mg/l), malic acid (4 g/l), $(NH_4)_2SO_4$ (1 g/l),

$MgSO_4 \cdot 7H_2O$ (200 mg/l), $FeSO_4 \cdot 7 H_2O$ (12 mg/l), K_2HPO_4 (0.9 g/l), KH_2PO_4 (0.6 g/l), 10 mM

sodium succinate, 100 mM MOPS (3-(N-morpholino)propanesulfonic acid) adjusted to pH 7.0

with KOH. In addition, 1ml/l of a trace elements solution was added [H_3BO_3 (2.8 g/l), $MnSO_4$

$\cdot H_2O$ (1.6 g/l), $Na_2MoO_4 \cdot 2H_2O$ (0.76 g/l), $ZnSO_4 \cdot 7 H_2O$ (240 mg/l), $Cu(NO_3)_2 \cdot 3 H_2O$ (40

mg/l), $CoCl_2 \cdot 6 H_2O$ (200 mg/l)]. For pH 6.0, the MOPS buffer was replaced with 100 mM MES

(2-(*N*-morpholino)ethanesulfonic acid) and for pH 8.0, the buffer was 100 mM TAPS

([tris(hydroxymethyl)methylamino]propanesulfonic acid). All buffered media were adjusted for

pH with KOH. Growth of the enriched culture of *Rhodoferax antarcticus* JLS was compared

with culture of the type strain *Rhodoferax antarcticus* ANT.BR obtained from the American

Type Culture Collection (ATCC 700587).

DNA isolation and sequencing.

Mat DNA was extracted by the PowerBiofilm kit (MO BIO). From each sample, approximately

20 mg material was extracted, and 0.75-1.50 µg DNA (measured by Qubit) was sent for

sequencing. From lake water filters, 300-400 ng DNA was obtained by extraction using an MP

FastDNA SPIN DNA kit (MP Biomedicals, CA) (Bielewicz *et al.* 2011). All metagenomic DNA

was sequenced at DOE JGI (Joint Genome Institute Community Science Program award 1936).

Shotgun metagenomic library construction and sequencing were carried out at JGI using standard

protocols on the Illumina HiSeq 2500 platform. The raw reads were quality filtered using JGI

standard protocols, and 1.2 Tb total sequences were obtained. Illumina reads were further

processed using Trimmomatic (Bolger *et al.* 2014) to remove adapters and low-quality sequences.

Trimmomatic counts the number of reads per FASTQ file, and it was used to determine the total

reads per sample metagenome.

Cultured *Rhodoferax* biofilm DNA was isolated by the PowerBiofilm kit (MO BIO).

Sequencing was performed by the Michigan State University Genomics Core. Libraries were

prepared using the Illumina TruSeq Nano DNA Library Preparation Kit. Sequencing was

performed on an Illumina MiSeq done in a 2x250bp format using an Illumina 500 cycle v2

reagent cartridge. The raw reads were quality filtered and 1 Tb total sequences were obtained.

Identification of ARGs by ShortBRED.

ShortBRED (Kaminski *et al.* 2015) is a pipeline that identifies protein family sequences by

generating specific peptide markers from a database of interest, then screening the markers for

specificity against the UniProt universal database of protein sequences (Bateman *et al.* 2021). The

shortbred_identify command scans a reference database for protein family-specific peptide

sequences. It was used to scan UniRef (the UniProt protein reference database) for sequences of

the ARGs in version 3.0.7 of the Comprehensive Antibiotic Resistance Database (CARD) (Alcock

et al. 2020). The minimum marker length accepted was 8 amino-acid residues, and the maximum

length for combined markers for a given protein was 200. Markers that matched nonspecific

genes in UniProt were discarded from the marker dataset.

The ShortBRED marker set was then used by the shortbred_quantify command to assign

ARGs in the 150-kb reads from lake water and mat samples. From each target read, six reading

frames were translated using tblastn. A marker hit required 95% match to the target. Sample

class differences for ARG abundance were tested for significance (P ≤ 0.05) using the Kruskal-

Wallis test.

Taxa classification of metagenomic reads.

The 150-kb sequence reads from mat and water metagenomes were assigned to taxonomic

classes using the Kraken 2 classifier pipeline (Wood & Salzberg 2014, Wood *et al.* 2019). Kraken

associates genomic *k*-mers (short sequence strings) in the reads with lowest common ancestor

taxa. The read classifications were then used to calculate taxa percentages via the Bracken

abundance estimator (Lu *et al.* 2017). The *k*-mer length was set at 150. For the Kraken 2-Bracken

pipeline, the reference database used was the 5/17/2021 Standard Collection accessed at:

https://benlangmead.github.io/aws-indexes/k2.

Eukaryotic taxa were further characterized using the pipeline EukDetect (Lind & Pollard

2021). This pipeline assigns reads to a database of 521,824 marker genes from 241 gene families,

out of 3713 genomes and transcriptomes of fungi, protists and invertebrates.

For an alternative classifier, mat and water metagenomes were mapped to core taxonomic

markers using the MetaPhlAn2 pipeline (Segata *et al.* 2012, Truong *et al.* 2015). MetaPhlAn2

assigns short reads to taxa using a set of marker genes identified from approximately 17,000

microbial reference genomes, primarily bacteria and archaea.

Genome assembly by *breseq*.

The pipeline *breseq* 0.35.6 (Deatherage & Barrick 2014) was used to assemble reads from two

samples of the enrichment culture obtained from Mat-04. The reads were mapped to the *R.*

antarcticus ANT.BRT (DSMZ24876) reference genome (Baker *et al.* 2017).

RESULTS

Taxa abundance of mat and water samples.

Metagenomes were sequenced from three microbial sources: the water columns of Lake Fryxell

(FRY-01, 02, 03) and Lake Bonney (BON-01, 02, 03), and the lift-off mat from the ice surface of

Lake Fryxell (Mat 01 – 06). A total of 4 billion reads were sequenced, with a range of between

210 – 420 million reads per metagenome. The average read length was 147 bp, SD = 26.

From the three groups of samples, we characterized the taxa abundance using

Kraken2/Bracken pipelines (Wood *et al.* 2019, Lu *et al.* 2017). The ice-surface mat samples (Mat-01

through Mat-06) showed substantial amounts of cyanobacteria, ranging from 9-60% of the total

(**Fig. 2A**). The predominant orders of cyanobacteria were Oscillatoriales and Nostocales (**Fig.**

2B). High abundance of Oscillatoriales and Nostocales is consistent with previous reports for

Fryxell mats obtained from benthic samples (Dillon *et al.* 2020). For Mat-01, 02, 03, 06 the

cyanobacterial assignments were primarily Oscillatoriales and Nostocales. Mat-04 and Mat-05

however showed depletion of Oscillatoriales, with predominant abundance of

Betaproteobacteria. Throughout the six mat samples, other taxa with significant abundance

included Actinobacteria, Bacteroidetes, and Alphaproteobacteria, with smaller abundance of

Planctomycetes, Bacteroidetes and Firmicutes. Thus the ice-trapped, air-exposed community

showed a composition remarkably similar to that of the benthic mat samples from which lift-off

mats originate (Dillon *et al.* 2020).

The filtered water from both lakes Fryxell and Bonney contained abundant bacterial taxa

of Actinobacteria, Bacteroides, Betaproteobacteria, and Alphaproteobacteria, similar to previous

reports of Fryxell water (Kwon *et al.* 2017) and other glacier-associated water bodies (Boetius *et al.*

2015). The water column showed almost no cyanobacteria but a high proportion of eukaryotes, in

some cases more than 50% of the reads classified by Kraken2/Bracken. By contrast, the mat

samples showed virtually no detectable eukaryotic DNA. The one-sided Mann-Whitney U test

confirms that water from each lake contains more eukaryotic DNA than the mat samples (p =

0.01) and that the mat samples contain more cyanobacteria than the water microbiomes of either

lake (p = 0.01).

To classify the eukaryotes, we used EukDetect (Lind & Pollard 2021) a pipeline that

matches short read data to marker genes from fungal, protist and invertebrate genomes (**Fig. 3**;

full output provided in **Table S1**). EukDetect identified the protist species *Geminigera cryophila*

(van den Hoff *et al.* 2020), *Mesodinium rubrum* (Yih *et al.* 2004), *Nannochloropsis limnetica* (Kong *et*

al. 2012), and *Chlamydomonas* sp. ICE-L (Lizotte *et al.* 1996). The two lakes showed similar taxa

profiles, except that *Chlamydomonas* was found only in Lake Bonney. Mat samples had too few

eukaryotic read counts to classify.

The validity of taxa classifier pipelines is highly dependent on their algorithm and taxa

database. For comparison with Kraken2/Braken output, our mat and water reads were mapped to

core taxonomic markers using an alternative pipeline, MetaPhlAn2 (Segata *et al.* 2012, Truong *et*

al. 2015) (**Fig. 4**). Four of the six mat samples showed mainly cyanobacteria, consistent with

longstanding reports of cyanobacterial mats (Taton *et al.* 2003, Hawes *et al.* 2013, Jungblut *et al.*

2016). Consistent with the Kraken analysis, Mat-04 and Mat-05 samples showed major amounts

of Betaproteobacteria and Alphaproteobacteria, with depletion of cyanobacteria. For the filtered

water samples, MetaPhlAn2 markers include limited content of Eukarya, and the pipeline did not

find the eukaryotic taxa predicted by Kraken2 (**Fig. 2**). The filtered water from both lakes

showed mainly Actinobacteria, with some Betaproteobacteria. Overall MetaPhlAn2 appeared

less effective than Kraken 2/Bracken at classifying our samples.

ARG composition characterized using ShortBRED.

Gene sequences encoding various forms of antibiotic resistance, including genomic loci as well

as mobile elements, are collected in the Comprehensive Antibiotic Resistance Database (CARD)

(Alcock *et al.* 2020). We used the ShortBRED pipeline to identify ARGs in our samples, based on

a set of marker peptides matching CARD sequences. The percentage of reads from each sample

that matched ARGs ranged from 0.0001-0.0006% (**Fig. 5**). The overall ARG abundances were

similar amongst the three sample classes (Fryxell mat, Fryxell water, Bonney water).

The specific ARGs identified by ShortBRED were sorted by abundance across samples

(**Table 2;** Supplementary file **Table S2**). Eight of the top 20 sorted ARGs showed a difference

among the three sample classes (Kruskal-Wallis test, P ≤ 0.05); of these results, 7 of 8 are

significant. Of the top-ranked ARGs, BUT-1 and vancomycin-resistance genes *vanYA*, *vanTG*

and *vanI* appeared mainly in the water column. BUT-1 is class-C beta-lactamase reported in

Buttiauxella agrestis, an environmental Gammaproteobacterium found in environmental water

sources (Fihman *et al.* 2002, Nakamichi *et al.* 2021). The vancomycin resistance genes occur in

Enterococcus and other Gram-positives (Courvalin 2006). By contrast, the beta-lactamase

AAC(3)-Ia (Riccio *et al.* 2003) occurred only in the mat samples. AAC(3)-Ia is encoded on

Pseudomonas aeruginosa integrons as well as in other Proteobacteria. Other ARGs showed

possible differences in prevalence between Lake Bonney and Lake Fryxell water; for example,

the Streptomyces-associated beta-lactamase AAC(3)-VIIa (López-Cabrera *et al.* 1989) was more

abundant in Bonney water whereas *vanYA* was more abundant in Fryxell water.

Overall, the three sample classes each had distinctive ARG-abundance signatures, as

indicated by Kruskal-Wallis test. Ten of the top 20 ranked ARGs were associated with

Proteobacteria, which roughly corresponds with the proportion of bacterial taxa abundances

predicted by Kraken2/Bracken (**Fig. 2**).

Enrichment of Mat-forming *Rhodoferax antarcticus*

The relatively high proportions of Betaproteobacteria and Alphaproteobacteria in MAT-04

differed from that of other samples, and the color of the sample material was more red than

green. We cultured the various samples anaerobically (in closed tubes) at 10°C with illumination.

From MAT-04 red spots of biofilm were obtained. There was no evidence of planktonic growth

(cloudiness) or cell motility within the culture medium, but the organism grew as a mat, forming

red blobs with a "red nose" appearance. In addition the culture extended as a red film-like mat

growing along the inside of the culture tube (**Fig. 6A**). It was not possible to obtain isolated

colonies, but serial culture of the "red noses" showed consistent biofilm formation and Gram-

stain morphology (**Fig. 6B**). This finding is consistent with an earlier report that the Fryxell

sedimentary biofilm includes a layer of phototrophic purple bacteria beneath the cyanobacterial

layer (Buffan-Dubau *et al.* 2001). Ours is the first report of such biofilm cultured from material at

the ice surface, rather than from its inferred origin in the sediment at the bottom of the lake.

The MAT-04 biofilm culture was subjected to genome analysis. DNA was sequenced and

short reads were used to BLAST the NCBI database. The predominant hits were to the genome

of *Rhodoferax antarcticus* ANT.BRT, a Betaproteobacterium originally isolated from a saltwater

pond at Cape Royds, Ross Island, Antarctica (Madigan *et al.* 2000, Baker *et al.* 2017). Kraken

2/Bracken analysis of sequenced reads agreed with this classification, assigning the samples as

95% *R. antarcticus* with about 1% *Pseudomonas.*

We used the *breseq* pipeline (Deatherage & Barrick 2014) to assemble reads from two non-

axenic samples (7-RN1 and 8-RN2) whose DNA sequence reads were assembled to the *R.*

antarcticus ANT.BRT reference genome (Supplementary file, **Table S3** and **S4**). These tables

present "mutations," that is, all differences from the ANT.BR reference sequence that show

100% read coverage. The two samples, 7-RN1 and 8-RN2, showed 276 and 271 sequence

differences respectively, in their main chromosome compared with that of the reference genome,

length 3,809,266 bp (Baker *et al.* 2017). The sample genomes showed > 99.99% identity with the

main chromosome and > 99% identity with one plasmid in the reference. A large part of the

observed sequence differences from reference were silent mutations. These observations confirm

the close relatedness of our culture to the published strain.

The *R. antarcticus* enrichment culture (designated *R. antarcticus* JLS) was tested for

ARG abundance using ShortBRED (**Table 3**). More than thirty ARGs showed hits, but nearly all

were associated with *Pseudomonas aeruginosa*. By contrast, no ARGs matched those from

CARD database that were associated with *Rhodoferax.*

Characterization of the mat-forming *R. antarcticus* JLS

The *R. antarcticus* JLS biofilm was further tested for growth range of pH and salinity.

The biofilm was subcultured anaerobically with illumination, in a malate-succinate *Rhodoferax*

medium modified from that of references (Madigan *et al.* 2000, Tayeh & Madigan 1987) as described

under Methods (**Fig. 7A, B, C**). After 45 days, at pH 7, the bacteria formed globules as well as a

red coating along the glass. At pH 6, spots of biofilm grew slowly, and at pH 8 little growth was

seen. Growth was also tested for cultures buffered at pH 7 with NaCl amendment (**Fig. 7D, E,**

F). The fullest growth was seen in the absence of NaCl amendment (the core medium contains

approximately 10 mM Na$^+$ ions). Less growth was seen with added NaCl (23 mM or 46 mM).

Despite the high sequence identity, the *R. antarcticus* JLS enrichment culture differed

from the *R. antarcticus* ANT.BR strain (Madigan *et al.* 2000) in its growth phenotype, under all

conditions of pH and NaCl concentration that were tested for both strains. Unlike the motile,

planktonic *R. antarcticus* ANT.BR, our culture showed no motility and little sign of planktonic

single-celled growth. Instead, the subcultured material grew entirely as a biofilm, in globular

spots and as a film-like growth along the interior surface of the glass tube.

The type strain ANT.BR was cultured at the same time, under all conditions. The original

strain never formed a biofilm; it appeared planktonic and motile under all conditions of pH and

NaCl concentration.

DISCUSSION

Identification of antibiotic resistance genes from relatively pristine environments is of

interest for several reasons. Long before human introduction of high-dose antibiotics,

environmental bacteria evolved multidrug pumps to efflux toxic products of their own

metabolism, as well as antimicrobial substances produced by their competitors (Allen *et al.* 2010,

Wright 2019). Many antibiotics possess signalling capabilities and other unknown functions.

Phylogeny dates the origin of beta-lactamases to hundreds of millions of years ago. Vancomycin

resistance genes are found in 30,000-year-old permafrost (Dcosta *et al.* 2011). But the specific

kinds of ARGs found in environmental sources may differ from those prevalent in human

microbiomes, those conferring resistance to the drugs we depend on for therapy (Zeng *et al.* 2019).

In our Taylor Valley lake metagenomes, the top-scoring ARG was BUT-1, a

cephalosporinase previously found in a clinical isolate of *Buttauxiella* (Fihman *et al.* 2002). Two

other ARGs of clinical origin (*vanYA*, *vanTG*) encode vancomycin resistance components in

Enterococcus (Courvalin 2006, Boyd *et al.* 2006). The finding of clinical ARGs in Antarctic water

bodies is concerning. The rest of the top 20 ARGs we found appear common in environmental

organisms. Eight were beta-lactamases, which are commonly found in environmental organisms

but can also be readily transferred between environmental and pathogenic strains (Hooban *et al.*

2020).

In polar regions, previous metagenomic studies reveal a range of naturally occurring

ARGs. Surface soils of the Antarctic Mackay Glacier region show ARGs encoding multidrug

pumps, beta-lactamases and aminoglycoside inactivators, largely associated with Gram-negative

bacteria (van Goethem *et al.* 2018). A study of Tibetan soils showed a number of abundant ARGs,

most notably encoding vancomycin resistance (Bo Li *et al.* 2020). Our study of ARGs from

Antarctic lakes adds to this picture, showing that both water and microbial mat sources contain

familiar ARGs, and likely contain others not yet discovered in Antarctic strains.

It was interesting to compare the total ARG abundance of our Antarctic lake samples

with those of temperate-zone Ohio rural water bodies with moderate human inputs, in a study

where samples were prepared by the same method and analyzed using the same ShortBRED

marker set (Murphy *et al.* 2021). The overall ARG abundance is about the same for the two

environmental sources, except for river samples obtained just downstream of a wastewater

effluent pipe, where the total ARG prevalence is increased approximately five-fold. This result is

consistent with a model that natural microbial communities generally harbor a small balanced

prevalence of ARGs, which gets amplified by concentrated human input. Note however that a

database such as CARD can only represent a fraction of the actual ARGs out there; new drug

resistance families and related mobility agents are continually discovered.

The mat lift-off samples we obtained from the ice surface showed a range of DNA taxa

consistent with those reported for samples obtained from benthic mats (Dillon *et al.* 2020). Given

that our collected mat organisms had survived several years within ice, followed by air

dessication and prolonged UV radiation exposure, it is impressive how much of the Antarctic

mat microbiomes retain viability with intact DNA. Even ciliated protists and invertebrate worms

were obtained alive from samples cultured after months of storage at -80°C (not shown). It is

likely that some of these organisms are psychrophiles that continue to grow within the ice

(Boetius *et al.* 2015).

Previous studies emphasize the cyanobacterial content of lift-off mats, primarily

Oscillatoriales genera such as *Microcoleus*, as well as *Nostoc* (Taton *et al.* 2003, Jungblut *et al.*

2016). While most of our lift-off samples showed abundance of cyanobacteria, one sample

yielded cultures from which the majority of reads matched *R. antarcticus*. The finding of mat

samples enriched for *R. antarcticus* indicates that portions of the lower layer of *Rhodoferax* mat

(Buffan-Dubau *et al.* 2001), along with the cyanobacterial upper layer, can break off and form lift-

off patches that emerge from the ice. Our culture of *R. antarcticus* (*R. antarcticus* JLS) obtained

from Lake Fryxell showed mat-forming morphology very different from the motile single cells

of *R. antarcticus* ANT.BR isolated from Cape Royds (Madigan *et al.* 2000, Baker *et al.* 2017).

Despite the high genetic similarity, our cultured organism appears to represent a novel ecotype of

R. antarcticus.

370   Our genomic reads from *R. antarcticus* JLS showed no matches to our ShortBRED

371   antibiotic resistance markers, although the reference genome does indeed include various

372   resistance genes including numerous RND and MFS transporters as well as multidrug efflux

373   components. Thus, many naturally-occurring ARGs are likely to be missed by standard marker

374   searches.

375   The prevalence of eukaryotic sequences in the lake water metagenomes is consistent with

previous reports that protists play important roles in the Taylor Valley lake communities (Li *et al.*

2016, Bielewicz *et al.* 2011, Glatz *et al.* 2006, Lizotte *et al.* 1996). Microbial eukaryotes including

phototrophs and mixotrophs provide prominent functions in the lake ecology (Li *et al.* 2016). In

our data, the eukaryotic community of Lakes Fryxell and Bonney showed three major taxa in

common: *Gemingera cryophila*, *Mesodinium rubrum*, and *Nannochloropsis limnetica*. *G.*

cryophila is a mixotrophic cryptophyte that feeds on bacteria but also conducts photosynthesis as

a secondary endosymbiont alga (van den Hoff *et al.* 2020). *M. rubrum* is a ciliate that consumes

cryptophytes but also uses the prey chloroplasts to conduct photosynthesis (kleptoplasty) (Yih *et*

al. 2004). *N. limnetica* is a heterokont alga, with red alga-derived chloroplasts, primarily a

phototroph (Kong *et al.* 2012). Our Lake Bonney samples also showed sequences from

Chlamydomonas, a green alga that dominates some parts of the Lake Bonney water column

(Lizotte *et al.* 1996).

We note that in the water samples, smaller aquatic phototrophs were likely missed by the

0.45-µm filter; 0.2-µm filters would have been preferable but were not available in the field at

the time. Even 0.2-µm filters miss important microbial community members (Brown *et al.* 2015).

The mat samples however had no filtration, so a broader spectrum of cell sizes was captured.

It is interesting that the Lake Fryxell water shows mainly eukaryotic phototrophs whereas

the mat shows mainly cyanobacteria and proteobacterial phototrophs. The mat bacteria are likely

to survive a wider range of light and temperature conditions than the eukaryotes. From the

standpoint of drug resistance, cyanobacteria are more likely than eukaryotes to harbor and

transfer ARGs of potential bacterial pathogens. Nevertheless, protists can regulate bacterial ARG

composition in terrestrial communities (Nguyen *et al.* 2020), so this factor may be of interest when

assessing the lake water ARG pools.

Acknowledgments and Funding

Fieldwork was supported by the National Science Foundation award OPP-1637708 to RM.

For metagenome sequencing we thank the U.S. Department of Energy Joint Genome Institute, a

DOE Office of Science User Facility, supported by the Office of Science of the U.S. Department

of Energy Contract No. DE-AC02-05CH11231, JGI Metagenome award 1936 to JLS.

Bioinformatic analysis and biofilm culture and sequencing were supported by the National

Science Foundation award MCB-1923077 to JLS. We thank Caroline Harwood for guidance

regarding culture of the mat biofilm, and Emma Stuart-Bates for the biofilm culture micrograph.

We especially thank Darcy Blankenhorn for expert technical support.

Author Contributions Statement

ST designed the bioinformatic analysis of ARGs and drafted the manuscript. GH characterized

the *Rhodoferax* enrichment culture and contributed to the manuscript. DB ran the bioinformatic

pipelines and performed statistics. RM led the field expedition and collected water samples. JLS

conceived the central concept, collected the mat samples and isolated DNA, mentored students,

and completed the manuscript.

Availability of data and materials

Sequence read files from lake water and mat samples are deposited at NCBI under SRA

accession numbers SRP104818, SRP104821, SRP098041, SRP098040, SRP098042,

SRP104817, SRP098044, SRP104822, SRP098050, SRP104819, SRP104820, SRP104823.

Sequence read files from cultured *Rhodoferax antarcticus* JLS are deposited at NCBI under SRA

accession number PRJNA736311.

REFERENCES

ALCOCK, B.P., RAPHENYA, A.R., LAU, T.T.Y., TSANG, K.K., BOUCHARD, M., EDALATMAND, A., HUYNH, W., et al.
2020. CARD 2020: antibiotic resistome surveillance with the comprehensive antibiotic resistance
database. *Nucleic acids research*, **48**, D517–D525, 10.1093/nar/gkz935.

ALLEN, H.K., DONATO, J., WANG, H.H., CLOUD-HANSEN, K.A., DAVIES, J. & HANDELSMAN, J. 2010. Call of the wild:
Antibiotic resistance genes in natural environments. *Nature Reviews Microbiology*, **8**, 251–259,
10.1038/nrmicro2312.

ANTELO, V., GIMÉNEZ, M., AZZIZ, G., VALDESPINO-CASTILLO, P., FALCÓN, L.I., RUBERTO, L.A.M., MAC CORMACK,
W.P., MAZEL, D. & BATISTA, S. 2021. Metagenomic strategies identify diverse integron-integrase and
antibiotic resistance genes in the Antarctic environment. *MicrobiologyOpen*, **10**, 1–14,
10.1002/mbo3.1219.

BAKER, J., RIESTER, C., SKINNER, B., NEWELL, A., SWINGLEY, W., MADIGAN, M., JUNG, D., et al. 2017. Genome
Sequence of Rhodoferax antarcticus ANT.BRT; A Psychrophilic Purple Nonsulfur Bacterium from an
Antarctic Microbial Mat. *Microorganisms*, **5**, 8, 10.3390/microorganisms5010008.

BATEMAN, A., MARTIN, M.J., ORCHARD, S., MAGRANE, M., AGIVETOVA, R., AHMAD, S., ALPI, E., et al. 2021.
UniProt: The universal protein knowledgebase in 2021. *Nucleic Acids Research*, **49**, D480–D489,
10.1093/nar/gkaa1100.

BIELEWICZ, S., BELL, E., KONG, W., FRIEDBERG, I., PRISCU, J.C. & MORGAN-KISS, R.M. 2011. Protist diversity in a
permanently ice-covered Antarctic Lake during the polar night transition. *ISME Journal*, **5**, 1559–
1564, 10.1038/ismej.2011.23.

BOETIUS, A., ANESIO, A.M., DEMING, J.W., MIKUCKI, J.A. & RAPP, J.Z. 2015. Microbial ecology of the
cryosphere: Sea ice and glacial habitats. *Nature Reviews Microbiology*, **13**, 677–690,
10.1038/nrmicro3522.

BOLGER, A.M., LOHSE, M. & USADEL, B. 2014. Trimmomatic: A flexible trimmer for Illumina sequence data.
Bioinformatics, **30**, 2114–2120, 10.1093/bioinformatics/btu170.

BOYD, D.A., DU, T., HIZON, R., KAPLEN, B., MURPHY, T., TYLER, S., BROWN, S., JAMIESON, F., WEISS, K. & MULVEY,
M.R. 2006. VanG-type vancomycin-resistant Enterococcus faecalis strains isolated in Canada.
Antimicrobial Agents and Chemotherapy, **50**, 2217–2221, 10.1128/AAC.01541-05.

BRAMBILLA, E., HIPPE, H., HAGELSTEIN, A., TINDALL, B.J., STACKEBRANDT, E., FRYXELL, L. & VALLEYS, M.D. 2001.
16S rDNA diversity of cultured and uncultured prokaryotes of a mat sample. *International Journal*,
23–33.

BROWN, C.T., HUG, L.A., THOMAS, B.C., SHARON, I., CASTELLE, C.J., SINGH, A., WILKINS, M.J., WRIGHTON, K.C.,
WILLIAMS, K.H. & BANFIELD, J.F. 2015. Unusual biology across a group comprising more than 15% of
domain Bacteria. *Nature*, **523**, 208–211, 10.1038/nature14486.

BUFFAN-DUBAU, E., PRINGAULT, O. & DE WIT, R. 2001. Artificial cold-adapted microbial mats cultured from
Antarctic lake samples. 1. Formation and structure. *Aquatic Microbial Ecology*, **26**, 115–125.

CAVICCHIOLI, R. 2015. Microbial ecology of Antarctic aquatic systems. *Nature Reviews Microbiology*, **13**,
691–706, 10.1038/nrmicro3549.

COURVALIN, P. 2006. Vancomycin resistance in Gram-positive cocci. *Clinical Infectious Diseases*, **42**, S25–
S34.

D'COSTA, V.M., KING, C.E., KALAN, L., MORAR, M., SUNG, W.W.L., SCHWARZ, C., FROESE, D., et al. 2011.
Antibiotic resistance is ancient. *Nature*, **477**, 457–461, 10.1038/nature10388.

DCOSTA, V.M., KING, C.E., KALAN, L., MORAR, M., SUNG, W.W.L., SCHWARZ, C., FROESE, D., et al. 2011.
Antibiotic resistance is ancient. *Nature*, **477**, 457–461, 10.1038/nature10388.

DEATHERAGE, D.E. & BARRICK, J.E. 2014. Identification of mutations in laboratory-evolved microbes from
next-generation sequencing data using *breseq* Sun, L. & Shou, W., *eds. Methods in Molecular*
Biology, **1151**, 165–188, 10.1007/978-1-4939-0554-6.

DILLON, M.L., HAWES, I., JUNGBLUT, A.D., MACKEY, T.J., EISEN, J.A., DORAN, P.T. & SUMNER, D.Y. 2020.
Energetic and Environmental Constraints on the Community Structure of Benthic Microbial Mats in
Lake Fryxell, Antarctica. *FEMS Microbiology Ecology*, **96**, 1–10, 10.1093/femsec/fiz207.

FIHMAN, V., ROTTMAN, M., BENZERARA, Y., DELISLE, F., LABIA, R., PHILIPPON, A. & ARLET, G. 2002. BUT-1: A new
member in the chromosomal inducible class C β-lactamases family from a clinical isolate of
Buttiauxella sp. *FEMS Microbiology Letters*, **213**, 103–111, 10.1016/S0378-1097(02)00792-9.

FIXEN, K.R., ODA, Y. & HARWOOD, C.S. 2019. Redox regulation of a light-harvesting antenna complex in an
anoxygenic phototroph. *mBio*, **10**, 10.1128/mBio.02838-19.

GLATZ, R.E., LEPP, P.W., WARD, B.B. & FRANCIS, C.A. 2006. Planktonic microbial community composition
across steep physical/chemical gradients in permanently ice-covered Lake Bonney, Antarctica.
Geobiology, **4**, 53–67, 10.1111/j.1472-4669.2006.00057.x.

HALL, C.M., CASTRO, M.C., KENIG, F. & DORAN, P.T. 2017. Constraining the recent history of the perennially
ice-covered Lake Bonney, East Antarctica using He, Kr and Xe concentrations. *Geochimica et*
Cosmochimica Acta, **209**, 233–253, 10.1016/j.gca.2017.04.023.

HAWES, I., SUMNER, D., ANDERSEN, D., JUNGBLUT, A. & MACKEY, T. 2013. Timescales of growth response of
microbial mats to environmental change in an ice-covered Antarctic lake. *Biology*, **2**, 151–176,
10.3390/biology2010151.

HEAD, J.W. & MARCHANT, D.R. 2014. The climate history of early Mars: insights from the Antarctic
McMurdo Dry Valleys hydrologic system. *Antarctic Science*, **26**, 774–800,
10.1017/S0954102014000686.

HOOBAN, B., JOYCE, A., FITZHENRY, K., CHIQUE, C. & MORRIS, D. 2020. The role of the natural aquatic
environment in the dissemination of extended spectrum beta-lactamase and carbapenemase
encoding genes: A scoping review. *Water Research*, **180**, 10.1016/j.watres.2020.115880.

JARA, D., BELLO-TOLEDO, H., DOMÍNGUEZ, M., CIGARROA, C., FERNÁNDEZ, P., VERGARA, L., QUEZADA-AGUILUZ, M.,
OPAZO-CAPURRO, A., LIMA, C.A. & GONZÁLEZ-ROCHA, G. 2020. Antibiotic resistance in bacterial isolates
from freshwater samples in Fildes Peninsula, King George Island, Antarctica. *Scientific reports*, **10**,
3145, 10.1038/s41598-020-60035-0.

JUNG, D.O., ACHENBACH, L.A., KARR, E.A., TAKAICHI, S. & MADIGAN, M.T. 2004. A gas vesiculate planktonic
strain of the purple non-sulfur bacterium Rhodoferax antarcticus isolated from lake Fryxell, Dry
Valleys, Antarctica. *Archives of Microbiology*, **182**, 236–243, 10.1007/s00203-004-0719-8.

JUNGBLUT, A.D., HAWES, I., MACKEY, T.J., KRUSOR, M., DORAN, P.T., SUMNER, D.Y., EISEN, J.A., HILLMAN, C. &
GORONCY, A.K. 2016. Microbial mat communities along an oxygen gradient in a perennially ice-
covered Antarctic lake. *Applied and Environmental Microbiology*, **82**, 620–630,
10.1128/AEM.02699-15.

Kaminski, J., Gibson, M.K., Franzosa, E.A., Segata, N., Dantas, G. & Huttenhower, C. 2015. High-specificity
targeted functional profiling in microbial communities with ShortBRED. *PLoS computational*
biology, **11**.

Kim, M.K. & Harwood, C.S. 1991. Regulation of benzoate-CoA ligase in Rhodopseudomonas palustris.
FEMS Microbiology Letters, **83**, 199–203, 10.1111/j.1574-6968.1991.tb04440.x-i1.

Kong, W., Ream, D.C., Priscu, J.C. & Morgan-Kiss, R.M. 2012. Diversity and expression of RubisCO genes
in a perennially ice-covered antarctic lake during the polar night transition. *Applied and*
Environmental Microbiology, **78**, 4358–4366, 10.1128/AEM.00029-12.

Koo, H., Hakim, J.A., Morrow, C.D., Crowley, M.R., Andersen, D.T. & Bej, A.K. 2018. Metagenomic
analysis of microbial community compositions and cold-responsive stress genes in selected
antarctic lacustrine and soil ecosystems. *Life*, **8**, 10.3390/life8030029.

Kwon, M., Kim, M., Takacs-Vesbach, C., Lee, J., Hong, S.G., Kim, S.J., Priscu, J.C. & Kim, O.S. 2017. Niche
specialization of bacteria in permanently ice-covered lakes of the McMurdo Dry Valleys, Antarctica.
Environmental Microbiology, **19**, 2258–2271, 10.1111/1462-2920.13721.

Lawrence, M.J.F. & Hendy, C.H. 1985. Water column and sediment characteristics of lake fryxell, taylor
valley, Antarctica. *New Zealand Journal of Geology and Geophysics*, **28**, 543–552,
10.1080/00288306.1985.10421206.

Li, Bo, Chen, Z., Zhang, F., Liu, Y. & Yan, T. 2020. Abundance, diversity and mobility potential of antibiotic
resistance genes in pristine Tibetan Plateau soil as revealed by soil metagenomics. *FEMS*
Microbiology Ecology, **96**, 1–10, 10.1093/femsec/fiaa172.

Li, Wei, Dore, J.E., Steigmeyer, A.J., Cho, Y.J., Kim, O.S., Liu, Y., Morgan-Kiss, R.M., Skidmore, M.L. & Priscu,
J.C. 2020. Methane production in the oxygenated water column of a perennially ice-covered
Antarctic lake. *Limnology and Oceanography*, **65**, 143–156, 10.1002/lno.11257.

Li, W., Podar, M. & Morgan-Kiss, R.M. 2016. Ultrastructural and single-cell-level characterization reveals
metabolic versatility in a microbial eukaryote community from an ice-covered Antarctic lake.
Applied and Environmental Microbiology, **82**, 3659–3670, 10.1128/AEM.00478-16.

Lind, A.L. & Pollard, K.S. 2021. Accurate and sensitive detection of microbial eukaryotes from whole
metagenome shotgun sequencing. *Microbiome*, **9**, 1–18, 10.1186/s40168-021-01015-y.

Lizotte, M.P., Sharp, T.R. & Priscu, J.C. 1996. Phytoplankton dynamics in the stratified water column of
Lake Bonney, Antarctica: I. Biomass and productivity during the winter-spring transition. *Polar*
Biology, **16**, 155–162, 10.1007/BF02329203.

López-Cabrera, M., Pérez-González, J.A., Heinzel, P., Piepersberg, W. & Jiménez, A. 1989. Isolation and
nucleotide sequencing of an aminocyclitol acetyltransferase gene from Streptomyces rimosus
forma paromomycinus. *Journal of bacteriology*, **171**, 321–328, 10.1128/jb.171.1.321-328.1989.

Lu, J., Breitwieser, F.P., Thielen, P. & Salzberg, S.L. 2017. Bracken: Estimating species abundance in
metagenomics data. *PeerJ Computer Science*, **2017**, 1–17, 10.7717/peerj-cs.104.

MADIGAN, M.T., JUNG, D.O., WOESE, C.R. & ACHENBACH, L.A. 2000. Rhodoferax antarcticus sp. nov., a
moderately psychrophilic purple nonsulfur bacterium isolated from an Antarctic microbial mat.
Archives of Microbiology, **173**, 269–277, 10.1007/s002030000140.

MIKUCKI, J.A., AUKEN, E., TULACZYK, S., VIRGINIA, R.A., SCHAMPER, C., SØRENSEN, K.I., DORAN, P.T., DUGAN, H. &
FOLEY, N. 2015. Deep groundwater and potential subsurface habitats beneath an Antarctic dry
valley. *Nature Communications*, **6**, 6831, 10.1038/ncomms7831.

MOOREHEAD, D.L., DORAN, P.T., FOUNTAIN, A.G., LYONS, W.B., MCNIGHT, D.M., PRISCU, J.C., VIRGINIA, R.A. &
WALL, D.H. 1999. Ecological legacies: Impacts on ecosystems of the McMurdo Dry Valleys.
Bioscience, **49**, 1009–1019.

MURPHY, A., BARICH, D., FENNESSY, M.S. & SLONCZEWSKI, J.L. 2021. An Ohio State Scenic River Shows
Elevated Antibiotic Resistance Genes, Including Acinetobacter Tetracycline and Macrolide
Resistance, Downstream of Wastewater Treatment Plant Effluent. *Microbiology Spectrum*, **9**,
e00941.

MURRAY, C.J., IKUTA, K.S., SHARARA, F., SWETSCHINSKI, L., ROBLES AGUILAR, G., GRAY, A., HAN, C., et al. 2022.
Global burden of bacterial antimicrobial resistance in 2019: a systematic analysis. *The Lancet*, **6736**,
10.1016/s0140-6736(21)02724-0.

NA, G., WANG, C., GAO, H., LI, R., JIN, S., ZHANG, W. & ZONG, H. 2019. The occurrence of sulfonamide and
quinolone resistance genes at the Fildes Peninsula in Antarctica. *Marine Pollution Bulletin*, **149**,
110503, 10.1016/j.marpolbul.2019.110503.

NAKAMICHI, N., MORIUCHI, R., DOHRA, H., FUTAMATA, H. & TASHIRO, Y. 2021. Complete Genome Sequence of
Buttiauxella agrestis DSM 9389. *Microbiology Resource Announcements*, **10**, 20–22,
10.1128/mra.00301-21.

NGUYEN, B.A.T., CHEN, Q.L., HE, J.Z. & HU, H.W. 2020. Microbial regulation of natural antibiotic resistance:
Understanding the protist-bacteria interactions for evolution of soil resistome. *Science of the Total
Environment*, **705**, 135882, 10.1016/j.scitotenv.2019.135882.

PARKER, B.C., SIMMONS, G.M., WHARTON, R.A., SEABURG, K.G. & LOVE, F.G. 1982. Removal of organic and
inorganic matter from Anarctic lakes by aerial escape of bluegreen algal mats. **18**, 72–78.

PRISCU, J. 2021a. Chlorophyll-a concentrations in discrete water column samples collected from lakes in
the McMurdo Dry Valleys, Antarctica (1993-2020, ongoing). *Environmental Data Initiative*.

PRISCU, J. 2022a. CTD profiles in lakes, McMurdo Dry Valleys, Antarctica (1993-2019, ongoing).
Environmental Data Initiative.

PRISCU, J. 2018. McMurdo Dry Valleys Ice Thickness for Taylor Valley Lakes, Antarctica ver 12.
Environmental Data Initiative, 12.

PRISCU, J. 2022b. Nitrogen and phosphorus concentrations in discrete water column samples collected
from lakes in the McMurdo Dry Valleys, Antarctica (1993-2020, ongoing). *Environmental Data
Initiative*.

PRISCU, J. 2021b. Underwater photosynthetically active radiation (PAR) vertical profiles collected from
lakes in the McMurdo Dry Valleys, Antarctica (1993-2020, ongoing). *Environmental Data Initiative*.

PRISCU, J.C. & SPIGEL, R.H. 1996. Evolution of temperature and salt structure of Lake Bonney, a chemically
stratified Antarctic lake. *Hydrobiologia*, **321**, 177–190.

REY, F.E., ODA, Y. & HARWOOD, C.S. 2006. Regulation of uptake hydrogenase and effects of hydrogen
utilization on gene expression in Rhodopseudomonas palustris. *Journal of Bacteriology*, **188**, 6143–
6152, 10.1128/JB.00381-06.

RICCIO, M.L., DOCQUIER, J.D., DELL'AMICO, E., LUZZARO, F., AMICOSANTE, G. & ROSSOLINI, G.M. 2003. Novel 3-N-
aminoglycoside acetyltransferase gene, aac(3)-Ic, from a Pseudomonas aeruginosa integron.
Antimicrobial Agents and Chemotherapy, **47**, 1746–1748, 10.1128/AAC.47.5.1746-1748.2003.

ROBERTS, E.C., PRISCU, J.C., WOLF, C., LYONS, W.B. & LAYBOURN-PARRY, J. 2004. The distribution of
microplankton in the McMurdo Dry Valley Lakes, Antarctica: response to ecosystem legacy or
present-day climatic controls? *Polar Biology*, **27**, 238–249, 10.1007/s00300-003-0582-0.

SEGATA, N., WALDRON, L., BALLARINI, A., NARASIMHAN, V., JOUSSON, O. & HUTTENHOWER, C. 2012.
Metagenomic microbial community profiling using unique clade-specific marker genes. *Nature
Methods*, **9**, 811–814, 10.1038/nmeth.2066.

SOHM, J.A., NIEDERBERGER, T.D., PARKER, A.E., TIRINDELLI, J., GUNDERSON, T., CARY, S.C., CAPONE, D.G. &
CARPENTER, E.J. 2020. Microbial Mats of the McMurdo Dry Valleys, Antarctica: Oases of Biological
Activity in a Very Cold Desert. *Frontiers in Microbiology*, **11**, 1–12, 10.3389/fmicb.2020.537960.

SPIGEL, R.H. & PRISCU, J.C. 1998. Physical limnology of the McMurdo Dry Valleys lakes. *In Ecosystem
Dynamics in a Polar Desert: the McMurdo Dry Valleys, Antartica*. 153–187., 10.1029/AR072p0153.

TAM, H.K., WONG, C.M.V.L., YONG, S.T., BLAMEY, J. & GONZÁLEZ, M. 2015. Multiple-antibiotic-resistant
bacteria from the maritime Antarctic. *Polar Biology*, **38**, 1129–1141, 10.1007/s00300-015-1671-6.

TATON, A., GRUBISIC, S., BRAMBILLA, E., DE WIT, R. & WILMOTTE, A. 2003. Cyanobacterial diversity in natural
and artificial microbial mats of Lake Fryxell (McMurdo Dry Valleys, Antarctica): a morphological and
molecular approach. *Applied and Environmental Microbiology*, **69**, 5157–5169.

TAYEH, M.A. & MADIGAN, M.T. 1987. Malate dehydrogenase in phototrophic purple bacteria: purification,
molecular weight, and quaternary structure. *Journal of bacteriology*, **169**, 4196–4202,
10.1128/jb.169.9.4196-4202.1987.

TRUONG, D.T., FRANZOSA, E.A., TICKLE, T.L., SCHOLZ, M., WEINGART, G., PASOLLI, E., TETT, A., HUTTENHOWER, C. &
SEGATA, N. 2015. MetaPhlAn2 for enhanced metagenomic taxonomic profiling. *Nature Methods*, **12**,
902–903, 10.1038/nmeth.3589.

VAN DEN HOFF, J., BELL, E. & WHITTOCK, L. 2020. Dimorphism in the Antarctic cryptophyte Geminigera
cryophila (Cryptophyceae). *Journal of Phycology*, **56**, 1028–1038, 10.1111/jpy.13004.

VAN GOETHEM, M.W., PIERNEEF, R., BEZUIDT, O.K.I., VAN DE PEER, Y., COWAN, D.A. & MAKHALANYANE, T.P. 2018.
A reservoir of "historical" antibiotic resistance genes in remote pristine Antarctic soils. *Microbiome*,
6, 1–12, 10.1186/s40168-018-0424-5.

VICK-MAJORS, T.J., PRISCU, J.C. & AMARAL-ZETTLER, L.A. 2014. Modular community structure suggests
metabolic plasticity during the transition to polar night in ice-covered Antarctic lakes. *ISME Journal*,
8, 778–789, 10.1038/ismej.2013.190.

WANG, F., STEDTFELD, R.D., KIM, O.S., CHAI, B., YANG, L., STEDTFELD, T.M., HONG, S.G., et al. 2016. Influence of
soil characteristics and proximity to antarctic research stations on abundance of antibiotic
resistance genes in soils. *Environmental Science and Technology*, **50**, 12621–12629,
10.1021/acs.est.6b02863.

WOOD, D.E., LU, J. & LANGMEAD, B. 2019. Improved metagenomic analysis with Kraken 2. *bioRxiv*, 1–13,
10.1101/762302.

WOOD, D.E. & SALZBERG, S.L. 2014. Kraken: Ultrafast metagenomic sequence classification using exact
alignments. *Genome Biology*, **15**, 10.1186/gb-2014-15-3-r46.

WRIGHT, G.D. 2019. Environmental and clinical antibiotic resistomes, same only different. *Current
Opinion in Microbiology*, **51**, 57–63, 10.1016/j.mib.2019.06.005.

YIH, W., HYUNG, S.K., HAE, J.J., MYUNG, G. & YOUNG, G.K. 2004. Ingestion of cryptophyte cells by the
marine photosynthetic ciliate Mesodinium rubrum. *Aquatic Microbial Ecology*, **36**, 165–170,
10.3354/ame036165.

ZENG, J., PAN, Y., YANG, J., HOU, M., ZENG, Z. & XIONG, W. 2019. Metagenomic insights into the distribution
of antibiotic resistome between the gut-associated environments and the pristine environments.
Environment International, **126**, 346–354, 10.1016/j.envint.2019.02.052.

ZHANG, Q., ZHANG, Z., LU, T., PEIJNENBURG, W.J.G.M., GILLINGS, M., YANG, X., CHEN, J., et al. 2020.
Cyanobacterial blooms contribute to the diversity of antibiotic-resistance genes in aquatic
ecosystems. *Communications Biology*, **3**, 1–10, 10.1038/s42003-020-01468-1.

TABLES

Table 1. Physical, chemical and biological parameters from Lakes Fryxell and Bonney (east

lobe).*

Site	Conductivity	Temperature	PAR	NH$_4^+$	SRP	Chlorophyll
	mS cm^{-1}	^{o}C	µmol m^{-2}s^{-1}	µM	µM	
Lake Fryxell, 9m depth	3.23	2.67	1.67	1.04	008	6.85
Lake Bonney, 15 m depth	17.21	4.58	8.46	6.07	0.04	0.72

644 *Data were retrieved from the McMurdo LTER long term record (Priscu 2022b, Priscu 2022a,

645 Priscu 2021b, Priscu 2021a). PAR, photosynthetically active radiation; SRP, signal recognition

646 protein.

Table 2. Mat and water antibiotic resistance genes (ARGs) identified by ShortBRED

marker peptides*

ARG Family	MAT-01	MAT-02	MAT-03	MAT-04	MAT-05	MAT-06	FRY-01	FRY-02	FRY-03	BON-01	BON-02	BON-03	TOTAL	P
ARO_3004294|BUT-1 *Buttiauxella agrestis*	0	1	2	0	0	0	554	1720	967	266	262	323	4095	0.027
ARO_3002541|AAC(3)-VIIa *Streptomyces rimosus*	89	235	114	173	152	54	44	31	31	323	313	231	1790	0.038
ARO_3002528|AAC(3)-Ia *Pseudomonas aeruginosa*	302	189	131	492	238	196	0	0	0	0	0	0	1548	0.022
ARO_3003167|CTX-M-159 *Proteobacteria*	36	9	25	1032	111	18	1	0	0	0	0	3	1235	0.052
ARO_3001461|OXA-156 *Pandoraea pulmonicola*	43	18	75	146	70	36	14	15	19	70	60	53	619	0.099
ARO_3002894|otrC *Streptomyces rimosus*	40	122	16	27	13	23	162	73	81	11	20	10	598	0.079
ARO_3003599|OXA-443 *Ralstonia mannitolilytica*	41	260	13	42	33	16	64	23	12	2	3	4	513	0.061
ARO_3001338|SHV-100 *Klebsiella pneumoniae*	58	1	49	144	56	44	4	1	5	10	6	11	389	0.241
ARO_3002955|vanYA *Enterococcus faecium*	0	3	2	1	1	0	101	59	47	19	29	31	293	0.027
ARO_3001855|ACT-35 *Enterobacter cloacae*	4	7	26	32	37	5	30	22	22	40	22	30	277	0.214
ARO_3000410|sul1 *Proteobacteria*	72	0	32	19	34	62	1	0	0	0	0	0	220	0.199
ARO_3003022|dfrB3 *Klebsiella oxytoca*	13	4	9	19	23	8	54	21	18	9	12	18	208	0.069
ARO_3002972|vanTG *Enterococcus faecalis*	1	0	1	0	2	0	10	9	13	50	53	55	194	0.027
ARO_3001517|OXA-329 *Acinetobacter calcoaceticus*	2	2	1	2	0	1	76	52	24	1	15	5	181	0.054
ARO_3002701|cmr *Rhodococcus fascians*	6	5	17	22	19	2	39	47	16	3	2	2	180	0.038
ARO_3003723|van *Desulfitobacterium hafniense*	0	0	0	0	0	0	53	50	40	9	16	10	178	0.024
ARO_3003942|abcA *Staphylococcus aureus*	31	75	2	9	17	20	0	0	0	7	4	5	170	0.055
ARO_3003583|basS *Pseudomonas aeruginosa*	12	1	30	8	7	12	1	0	0	47	32	19	169	0.046
ARO_3004652|Erm(O)-lrm *Streptomyces lividans*	28	8	16	37	35	10	8	7	4	5	2	5	165	0.061
ARO_3001299|tlrB_tylosin *Streptomyces fradiae*	18	8	10	30	26	8	2	2	0	18	18	22	162	0.034

*Sample class differences for ARG abundance were tested for significance (P ≤ 0.05) using the

Kruskal-Wallis test.

656 **Table 3.** *Rhodoferax* **enrichment culture ARGs identified by ShortBRED***

A. *Rhodoferax* (95%), *Pseudomonas* (1.5%)	Hits	Marker Length	B. *Rhodoferax* (95%), *Pseudomonas* (0.7%)	Hits	Marker Length
ARO_3003679|TriA__Pseudomonas_aeruginosa_PAO1	8	242	ARO_3003680|TriB__Pseudomonas_aeruginosa_PAO1	3	144
ARO_3001796|OXA-50__Pseudomonas_aeruginosa	4	116	ARO_3003031|mexW__Pseudomonas_aeruginosa_PAO1	1	56
ARO_3003681|TriC__Pseudomonas_aeruginosa_PAO1	3	113	ARO_3002320|KPC-10__Acinetobacter_baumannii	1	66
ARO_3003710|mexL__Pseudomonas_aeruginosa_PAO1	3	134	ARO_3003583|basS__Pseudomonas_aeruginosa	1	290
ARO_3003030|mexV__Pseudomonas_aeruginosa_PAO1	3	176	ARO_3004777|CMH-1__Enterobacter_cloacae	1	25
ARO_3002507|PDC-8__Pseudomonas_aeruginosa	3	155	ARO_3003705|mexN__Pseudomonas_aeruginosa	1	63
ARO_3000802|OprJ__Pseudomonas_aeruginosa	2	36	ARO_3000808|mexI__Pseudomonas_aeruginosa_PAO1	1	118
ARO_3003680|TriB__Pseudomonas_aeruginosa_PAO1	2	144	ARO_3004054|CpxR__Pseudomonas_aeruginosa	1	26
ARO_3000379|OprM__Pseudomonas_aeruginosa_PAO1	2	35			
ARO_3000805|OprN__Pseudomonas_aeruginosa_PAO1	2	102			
ARO_3002985|arnA__Pseudomonas_aeruginosa_PAO1	2	102			
ARO_3003692|mexJ__Pseudomonas_aeruginosa_PAO1	2	120			
ARO_3003031|mexW__Pseudomonas_aeruginosa_PAO1	2	56			
ARO_3004038|emrE__Pseudomonas_aeruginosa_PAO1	2	51			
ARO_3003705|mexN__Pseudomonas_aeruginosa	2	63			
ARO_3003698|mexP__Pseudomonas_aeruginosa	2	85			
ARO_3002645|APH(3)-IIb__Pseudomonas_aeruginosa	2	153			
ARO_3000377|MexA__Pseudomonas_aeruginosa_PAO1	2	106			
ARO_3001214|mdtM__Escherichia_coli_str__K-12	1	71			
ARO_3004077|PmpM__Pseudomonas_aeruginosa_PAO	1	109			
ARO_3000804|MexF__Pseudomonas_aeruginosa_PAO1	1	102			
ARO_3004072|OpmB__Pseudomonas_aeruginosa_PAO	1	111			
ARO_3004075|MuxC__Pseudomonas_aeruginosa_PAO1	1	177			
ARO_3003693|mexK__Pseudomonas_aeruginosa_PAO1	1	44			
ARO_3000809|opmD__Pseudomonas_aeruginosa_PAO	1	116			
ARO_3003682|OpmH__Pseudomonas_aeruginosa_PAO	1	79			
ARO_3002695|cmlA5__uncultured_bacterium	1	99			
ARO_3003583|basS__Pseudomonas_aeruginosa	1	290			
ARO_3000795|mdtE__Escherichia_coli_str__K-12	1	54			
ARO_3004612|ampH__Escherichia_coli_DH1	1	68			

658 *Enrichment cultures A and B, taxa proportions were estimated by Kraken2/Bracken pipeline.

662 **FIGURES**

664 **Figure 1. Lake Fryxell uplift mats.**

665 (**A**) Lake Fryxell with permanent ice cover, weathered by katabatic winds; December 10, 2014.

666 (**B**) Source of MAT-4 DNA. Uplift mat emerges from the ice; approximately 40 cm across.

Figure 2. Microbial community composition of mat and water samples. Relative abundance

of phyla was determined by read alignment to marker genes using Kraken2/Bracken pipeline.

(**A**) Phyla and classes. (**B**) Orders of cyanobacteria in mat samples.

MAT (Lake Fryxell, mat samples); FRY (Lake Fryxell, water samples); BON (Lake Bonney,

water samples). MAT samples were from ice surface, unfiltered. The water samples were

collected on a 0.45-µm filter. PVC=Planctomycetes-Verrucomicrobia-Chlamydiae

677 **Figure 3. Eukaryotic taxa classified by EukDetect.** Relative abundance of protist taxa were

678 determined by read alignment to marker genes using EukDetect (Lind & Pollard 2021).

Figure 4. Microbial community composition via MetaPhlAn2. Relative abundance of class-

level taxa was determined by read alignment to marker genes using MetaPhlAn2 (Segata *et al.*

2012, Truong *et al.* 2015). Samples were as indicated for Figure 2.

687 **Figure 5. ARG abundance in lake samples.** Percentage of reads matched to ShortBRED

688 markers. Total reads were counted using Trimmomatic. Bars indicate total ARG hits per sample

689 (blue) and top 60 ranked ARG hits (orange).

A

B

Figure 6. Mat-forming enrichment of Mat-04 *R. antarcticus* JLS.

(**A**) Biofilm of cultured *R. antarcticus* within closed tube of medium, showing globular form of

growth. Initial Mat-04 sample was subcultured serially three times in PM medium, in horizontal

closed tubes at 10°C with 30 umoles/m^{2}s illumination.

(**B**) Gram stain of culture, 1000X with oil immersion. Courtesy of Emma Stuart-Bates.

Figure 7. pH and NaCl dependence of *R. antarcticus* JLS.

(A, B, C) Medium PM contained **(A)** 100 mM MES pH 6; **(B)** 100 mM MOPS pH 7; **(C)** 100

mM TAPS pH 8. Cultures were incubated 45 days.

(D, E, F) Medium PM was amended with **(D)** no NaCl added; **(E)** 23 mM NaCl; **(F)** 46 mM

NaCl. Cultures were incubated 45 days.