Lipid hydroperoxides promote muscle atrophy through lysosomal amplification

Hiroaki Eshima,[1-3] Piyarat Siripoksup,[1,4] Justin L. Shahtout,[1,4] MacKenzie J. Pearson,[5] Ziad

S. Mahmassani,[1,2,4] Patrick J. Ferrara,[1,2,6] Alexis W. Lyons,[1] J. Alan Maschek,[1,6,7] Alek D.

Peterlin,[1,6] Anthony R. P. Verkerke,[1,6] Jordan M. Johnson,[1,6] Anahy Salcedo,[1] Jonathan J.

Petrocelli,[1,4] Ethan J. Anderson,[8] Sihem Boudina,[1,2,6] Qitao Ran,[9] James E. Cox,[1,7,10] Micah

J. Drummond,[1,2,4] Katsuhiko Funai,[1,2,4,6]*

[1]Diabetes & Metabolism Research Center, University of Utah, Salt Lake City, Utah, USA

[2]Molecular Medicine Program, University of Utah, Salt Lake City, Utah, USA

[3]Department of International Tourism, Nagasaki International University, Sasebo, Nagasaki,

Japan

[4]Department of Physical Therapy & Athletic Training, University of Utah, Salt Lake City,

Utah, USA

[5]Sciex, Framingham, MA, USA

[6]Department of Nutrition & Integrative Physiology, University of Utah, Salt Lake City, Utah,

USA

[7]Metabolomics Core Research Facility, University of Utah, Salt Lake City, Utah, USA

[8]Fraternal Order of Eagles Diabetes Research Center, University of Iowa, Iowa City, Iowa,

USA

[9]Department of Cell Systems and Anatomy, University of Texas Health Science Center at

San Antonio, San Antonio, Texas, USA

[10]Department of Biochemistry, University of Utah, Salt Lake City, Utah, USA

*Corresponding Author:

Katsuhiko Funai, Ph.D.

Diabetes & Metabolism Research Center

15 N 2030 E, Salt Lake City, UT 84112

29   Phone: (801) 585-1781

30   Fax: (801) 585-0701

31   Email: kfunai@health.utah.edu

Abstract

Reactive oxygen species (ROS) is a cardinal feature of skeletal muscle atrophy. However,

ROS refers to a collection of radical molecules whose cellular signals are vast, and it is

unclear which downstream consequences of ROS are responsible for the loss of muscle

mass and strength.[1,2] Here we show that lipid hydroperoxides (LOOH) are increased with

age and disuse, and the accumulation of LOOH by suppression of glutathione peroxidase 4

(GPx4) is sufficient to augment muscle atrophy. Strikingly, genetic and pharmacologic

suppression of muscle LOOH robustly prevented the reduction of both muscle mass and

force-generating capacity. LOOH promoted atrophy in a lysosomal-dependent,

proteasomal-independent manner, and the suppression of autophagic machinery was

sufficient to prevent muscle atrophy and weakness. Indeed, the lysosome is essential for

the amplification of LOOH induced by oxidative stress. Our findings provide novel insights

for the role of LOOH in muscle atrophy including a therapeutic implication by

pharmacologic suppression.

Main

Loss of muscle mass and function with age is detrimental to health and quality of life.[3,4] A

pharmacologic therapy for muscle loss does not exist, and current diet or exercise

therapeutic approaches are often ineffective or unfeasible. Oxidative stress has been

implicated in muscle atrophy by accelerating proteolysis,[2] but the exact mechanism by

which ROS contributes to the decrease in muscle mass and strength is not well

understood. Lipid hydroperoxide (LOOH) is a class of lipid ROS molecules that has been

implicated in cell damage, particularly as a trigger to induce ferroptosis, a non-apoptotic

form of regulated cell death.[5,6] LOOH propagation is endogenously suppressed by GPx4

that catalyzes the reaction by which LOOH is reduced to its nonreactive hydroxyl

metabolite.[7] However, the role of LOOH in skeletal muscle is not well understood.[8,9]

We first evaluated the changes in skeletal muscle LOOH with aging. In humans and

in mice, aging promoted a reduction in the expression of GPx4 in skeletal muscle (Fig.

1a&b). To examine the changes in skeletal muscle LOOH landscape with age, we

performed a comprehensive oxidolipidomic analysis in muscle samples from young (4

months) and old (20 months) mice (Fig. 1c&d). We detected over 300 species of oxidized

lipids with an effect distribution that was highly class-dependent. Among these, age had the

most robust effect on oxidized phosphatidylethanolamine (Fig. 1c, red), a class of lipids that

have been implicated as a potential lipid signal to induce ferroptosis.[10] Among the top ten

oxidized lipid species whose abundance was most robustly increased with age, six of them

were oxidized phosphatidylethanolamine (Figure 1d), and they were substantially more

highly abundant compared to other oxidized lipids. LOOH can be indirectly assessed by

quantifying LOOH adducts such as 4-hydroxynonenal (4-HNE) and malondialdehyde

(MDA). We confirmed increased muscle 4-HNE and MDA with age (Extended Data Fig. 1a-

c).

Disuse also promotes skeletal muscle atrophy that is likely contributed by ROS.[2] To

model disuse atrophy, mice underwent a hindlimb unloading (HU) procedure as previously

described.[11,12] As expected, HU induced muscle atrophy and weakness (Fig. 1e&f)

concomitant to reduction in body and lean mass (Extended Data Fig. 1d-f). Disuse robustly

elevated muscle LOOH levels (Fig. 1g, Extended Data Fig. 1g) without significant changes

in mitochondrial bioenergetics (Extended Data Fig. 1h-j). An increase in muscle LOOH

preceded atrophy, consistent with the notion that LOOH may trigger mechanisms to

promote loss of muscle mass.

Next, we tested our hypothesis that LOOH contributes to muscle atrophy using

C2C12 myotubes (Fig. 1h). Lentivirus-mediated knockdown (KD) of GPx4 increased LOOH

and markers of ferroptosis (Extended Data Fig. 2a-f) concomitant with a decrease in

myotube diameter (Fig. 1i&j). We also recapitulated these findings with erastin (system X_c^-

inhibitor that suppresses glutathione synthesis) (Fig. 1i&j and Extended Data Fig. 2g-k) and

RSL3 (GPx4 inhibitor) (Fig. 1i&j and Extended Data Fig. 2l-o), commonly used acute

pharmacological interventions to elevate intracellular LOOH. We then translated these

findings in vivo with global heterozygous GPx4 knockout mice (GPx4$^{+/-}$). Germline deletion

of GPx4 is embryonically lethal,[13] but GPx4$^{+/-}$ mice appear normal and do not have an

observable muscle phenotype at baseline.[7,14] We studied 4 months (young) and 20 months

(old) GPx4$^{+/-}$ and wildtype littermates with or without HU (Extended Data Fig. 3a-d). In

young mice, GPx4 haploinsufficiency augmented the loss in muscle mass induced by HU

(Fig. 2a and Extended Data Fig. 3e). However, muscle masses between old GPx4$^{+/-}$ and

wildtype mice were not different. We interpret these findings to mean that disuse in old

mice promotes an increase in LOOH that has already reached a maximally effective

threshold with age such that GPx4 deletion had no further effect. In support of this, we saw

no differences in 4-HNE or MDA levels between old GPx4$^{+/-}$ and wildtype mice (Fig. 2b and

Extended Data Fig. 3f&g). GPx4 haploinsufficiency did not alter force-generating capacity

(Fig. 2c and Extended Data Fig. 3h-j).

Because GPx4 is expressed globally, we also studied mice with skeletal muscle-

specific tamoxifen-inducible GPx4 knockout (GPx4-MKO) (Fig. 2d and Extended Fig. 4a).[15]

Consistent with GPx4+/- mice, GPx4-MKO mice were also more prone to developing

disuse-induced skeletal muscle atrophy (Fig. 2e&f and Extended Data Fig. 4b-f)

concomitant to elevated LOOH (Fig. 2g and Extended Data Fig. 4g&h), suggesting that loss

of GPx4 in muscle augments atrophy in a cell-autonomous manner. Histological analyses

revealed that reduced muscle mass was consistent with reduced cross-sectional area of

myofibers regardless of fiber-type compositions (Fig. 2h&i and Extended Data Fig. 4i&j).

These data implicate that LOOH directly reduces muscle cell size in vivo.

GPx4 primarily neutralizes LOOH but it also exhibits some activity towards other

peroxides.[16] To confirm that the effects of GPx4 deletion to promote atrophy is specific to

LOOH, we diminished the ability of cells to incorporate polyunsaturated fatty acids (PUFAs)

into phospholipids by deleting lysophosphatidylcholine acyltransferase 3 (LPCAT3).[17-19]

LPCAT3 is an enzyme of Lands cycle that preferentially acylates lysophospholipids with

PUFAs, and thus an essential component of ferroptosis[10]. Indeed, LPCAT3 KD rescued the

increase in 4-HNE induced by GPx4 KD (Fig. 2j and Extended Data Fig. 5a-c).

Remarkably, deletion of LPCAT3 KD completely restored the reduction in myotube

diameter induced by GPx4 KD (Fig. 2k&l). Similarly, LPCAT3 deletion also prevented

LOOH and cell death induced by erastin (Extended Data Fig. 5d-h). These findings indicate

that muscle atrophy induced by loss of GPx4 or erastin treatment is due to the

accumulation of LOOH and not other peroxides.

Would suppression of LOOH be sufficient to ameliorate skeletal muscle atrophy? We

studied young (4 months) and old (20 months) global GPx4-overexpressing (GPx4Tg)

mice[20] with or without HU (Extended Data Fig. 6a-e). Strikingly, both young and old

GPx4Tg mice were resistant to disuse-induced muscle atrophy (Fig. 3a and Extended Data

Fig. 6f&g). Perhaps even more impactful was the effect of GPx4 overexpression on skeletal

muscle force-generating capacity such that, in both young and old, GPx4 overexpression

robustly protected mice from muscle weakness induced by HU (Fig. 3b and Extended Data

Fig. 6h-j). These findings are in contrast to our experiments in GPx4$^{+/-}$ mice where muscle

mass phenotype was only present in the young mice (Fig. 2a) and no phenotype on muscle

strength (Fig. 2b). Consistent with the notion that GPx4 overexpression acts on LOOH, HU-

induced increase in 4-HNE was completely suppressed in GPx4Tg mice (Fig. 3c and

Extended Data Fig. 6k). We also found that the protection from muscle atrophy was

explained by greater myofiber cross-sectional area regardless of fiber-type (Fig. 3d&e and

Extended Data Fig. 6l&m).

Next, we explored opportunities to pharmacologically suppress LOOH to prevent

muscle atrophy. Ferrostatin-1 is an inhibitor of ferroptosis that is widely used to study

LOOH.[21,22] Indeed, incubation of cells with ferrostatin-1 was sufficient to suppress LOOH

induced by GPx4 KD (Fig. 3f and Extended Data Fig. 7a) concomitant with protection from

myotube atrophy (Fig. 3g&h and Extended Data Fig. 7b&c). Nevertheless, ferrostatin-1 is

currently not an FDA approved drug with questions surrounding safety. Thus, we tested L-

carnosine, a dipeptide composed of beta-alanine and L-histidine that has the ability to

scavenge LOOH.[23,24] Similar to ferrostatin-1, L-carnosine was sufficient to rescue LOOH

and cell death induced by GPx4 KD (Extended Data Fig. 7d-f) or erastin (Extended Data

Fig. 7g-i). Leveraging these data, we performed a preclinical trial for L-carnosine provided

in drinking water ad lib (80 mM) in young wildtype C57BL6/J mice. L-carnosine treatment

did not alter body mass, body composition, food intake, and water intake (Extended Data

Fig. 7j-m), and successfully suppressed muscle LOOH induced by HU (Extended Data Fig.

7n&o). Remarkably, mice provided with L-carnosine were partly protected from disuse-

induced muscle atrophy (Extended Data Fig. 7p).

In humans, L-carnosine is rapidly degraded by a circulating carnosinase[25] that may

render oral carnosine treatment ineffective. In contrast, N-acetylcarnosine has a longer

half-life and may be a more effective reagent in humans thus improving its translational

potential. Similar to ferrostatin-1 and L-carnosine, N-acetylcarnosine also prevented LOOH

and cell death induced by GPx4 KD (Extended Data Fig. 8a-c) or erastin (Extended Data

Fig. 8d-f). Leveraging these findings, we proceeded with a preclinical trial for N-

acetylcarnosine in drinking water (80 mM, Fig. 3i) in young (4 months, C57BL6/J; Jax

colony) and old (20 months, C57BL/6; NIA rodent colony) wildtype mice. Similar to L-

carnosine treatment, N-acetylcarnosine did not alter body mass, body composition, food

intake, or water intake (Extended Data Fig. 8g-k), and successfully suppressed muscle

LOOH (Fig. 3j and Extended Data Fig. 8l&m). Strikingly, similar to our findings in GPx4Tg

mice, N-acetylcarnosine ameliorated muscle atrophy (Fig. 3k and Extended Data Fig. 8n)

and weakness (Fig. 3l and Extended Data Fig. 8o-q) in both young and old mice. Protection

from muscle atrophy was similarly explained by greater myofiber cross-sectional area

regardless of fiber-type (Fig. 3m&n and Extended Data Fig. 8r&s). These promising

observations inform a potential clinical trial to test the efficacy of N-acetylcarnosine

treatment in ameliorating muscle atrophy in humans.

What is the mechanism by which LOOH promotes muscle atrophy? C2C12 myotubes

were pretreated with Bafilomycin A1 (BafA1) or MG132 prior to erastin incubation to

determine whether LOOH increases protein degradation in a lysosomal- or proteasomal-

dependent manner, respectively. Erastin-induced reduction in myotube diameter was

suppressed with BafA1, but not with MG132 (Fig. 4a&b and Extended Data Fig. 9a),

suggesting that the lysosome mediates protein degradation by LOOH.[26] We also

reproduced these findings with RSL3 treatment (Extended Data Fig. 9b&c). How does

LOOH, a lipid molecule, promote lysosomal-degradation? Upstream of the lysosome,

autophagosome formation is mediated by a lipidation of LC3 by ATG3.[27] Thus, we

hypothesized that LOOH may affect the lipidation of LC3. Indeed, GPx4 KD drastically

reduced the protein content of p62, LC3-I and LC3-II (Fig. 4c and Extended Data Fig. 9d-h),

potentially suggesting that LOOH may accelerate lysosomal degradation by affecting LC3

lipidation. To test this possibility, we performed a targeted deletion of ATG3 in vitro. Indeed,

ATG3 KD completely rescued the reduction in myotube diameter induced by GPx4 KD (Fig.

4d&e and Extended Data Fig. 9i-l). Leveraging these findings, we generated mice with

skeletal muscle-specific tamoxifen-inducible ATG3 knockout (ATG3-MKO) (Fig. 4f and

Extended Data Fig. 10a-c) and studied them with or without HU (Extended Data Fig. 10d-f).

Loss of muscle ATG3 was protective from disuse-induced atrophy (Fig. 4g and Extended

Data Fig. 10g) and weakness (Fig. 4h and Extended Data Fig. 10h) which can be explained

by greater myofiber cross-sectional area (Fig. 4i&j and Extended Data Fig. 10i).

We initially hypothesized that lysosomal-degradation mediates LOOH-induced

protein degradation to contribute to muscle atrophy. However, further assessment of

muscle LOOH illuminated a more complex interaction between the lysosome and

LOOH.[28,29] Unexpectedly, quantification of 4-HNE revealed that inhibition of the autophagy-

lysosome axis by ATG3 deletion or BafA1 was sufficient to inhibit LOOH induced by GPx4

deletion, erastin, or RSL3 (Fig. 4k and Extended Data Fig. 10j-p). These findings suggest

that the autophagy-lysosome axis is essential for LOOH amplification, in addition to its

potential role in mediating protein degradation downstream. Indeed, immunofluorescence

experiments revealed that 4-HNE is highly co-localized to LAMP2 (Fig. 4l and Extended

Data Fig. 10q), consistent with the notion that the lysosome is necessary for LOOH

propagation. To support this idea, an increase in LOOH by hydrogen peroxide or

carmustine (agents that increase pan oxidative stress without acting on GPx4 directly) was

completely inhibited by lysosomal inhibition (Fig. 4n and Extended Fig. 10r-t). Together,

these observations suggests that the propagation of LOOH may be mediated by the

lysosome (Fig. 4o).

In summary, the current findings demonstrate a novel mechanism that indicate LOOH

as the key downstream molecule by which oxidative stress promotes muscle atrophy and

weakness. Skeletal muscle LOOH was robustly upregulated with aging and/or disuse, and

genetic or pharmacologic suppression of LOOH was sufficient to rescue muscle atrophy. In

particular, N-acetylcarnosine treatment shows a potent effect in preserving muscle mass

and strength with disuse in both young and old mice, informing the potential trial to utilize

this compound to ameliorate loss of muscle function in humans. We further provide

evidence that the autophagy-lysosome axis is essential for the propagation of LOOH,

suggesting a novel interaction between these cellular processes that could be interrogated

to ameliorate LOOH-induced damage.

Methods

Animal models. GPx4+/- and GPx4Tg mice were generated previously.[13,20] Conditional GPx4 knockout (GPx4cKO+/+) mice were acquired from Jackson Laboratory (Stock No: 027964).[15] Conditional ATG3 knockout (ATG3cKO+/+) mice were previously described.[30] GPx4cKO+/+ mice or ATG3cKO+/+ mice were then crossed with tamoxifen-inducible, skeletal muscle–specific Cre recombinase (HSA-MerCreMer+/-) mice[31] to generate GPx4cKO+/+; HSAMerCreMer-/- (control) and GPx4cKO+/+; HSA-MerCreMer+/- (skeletal muscle–specific GPx4 knockout; GPx4-MKO) mice or ATG3cKO+/+; HSAMerCreMer-/- (control) and ATG3cKO+/+; HSA-MerCreMer+/- (ATG3-MKO) mice. Tamoxifen-injected (7.5 µg/g body mass, 5 consecutive days) littermates were used. Mice were maintained on a 12-hour light/12-hour dark cycle in a temperature-controlled room. Body composition measurements were taken immediately before terminal experiments with a Bruker Minispec MQ20 nuclear magnetic resonance (NMR) analyzer (Bruker, Rheinstetten, Germany). All mice were bred onto C57BL/6J background and were born at normal Mendelian ratios. Body mass were measured every day during HU. All protocols were approved by Institutional Animal Care and Use Committees at the University of Utah.

Hindlimb unloading. Mice underwent 1, 7, or 14 days of HU (2 mice/cage) using a previously described protocol[11,12] based on the traditional Morey-Holton design to study disuse atrophy in rodents. Along with daily monitoring of body mass, food intake was monitored every other day to ensure that the mice did not experience excessive weight loss due to malnutrition or dehydration. Following 1, 7, or 14 days of HU, mice were fasted for 4 h and given an intraperitoneal injection of 80 mg/kg ketamine and 10 mg/kg xylazine, after which tissues were harvested. Extensor digitorum longus (EDL), and soleus (SOL) were carefully dissected for weight measurements.

Muscle force generation. Force-generating properties of soleus and EDL muscles were measured as previously described.[32,33] Briefly, soleus/EDL muscles were sutured at each tendon, and muscles were suspended at optimal length (Lo), which was determined by pulse stimulation. After Lo was identified, muscles were stimulated (0.35 seconds, pulse width 0.2 milliseconds) at frequencies ranging from 10 to 200 Hz. Muscle length and mass were measured to quantify cross-sectional area for force normalization.

Quantitative reverse transcription PCR. Samples were homogenized in TRIzol reagent (Life Technologies) to extract total RNA. One microgram RNA was reverse-transcribed using an IScript cDNA synthesis kit (Bio-Rad). Reverse transcription PCR (RT-PCR) was performed with the Viia 7 Real-Time PCR System (Life Technologies) using SYBR Green

reagent (Life Technologies). All data were normalized to ribosomal L32 gene expression and
were normalized to the mean of the control group. Primers were based on sequences in
public databases.

Western blot. Whole muscle or cells were homogenized, and western blots were performed
as previously described.[11] Protein homogenates were analyzed for abundance of
phosphorylated 4-hydroxynonenal (4-HNE; ab48506; Abcam), GPx4 (ab125066, Abcam),
actin (A2066, MilliporeSigma), GAPDH (14C10, Cell Signaling Technology), p62 (ab56416,
Abcam), LC3B (83506, Cell Signaling Technology).

Mass spectrometry. Oxidolipidomics samples were analyzed on the SCIEX 7500 system
coupled with ExionLC (SCIEX, Concord, Canada) using multiple reaction monitoring (MRM)
analysis. Mobile phase A is composed of 93:7 acetonitrile:dichloromethane containing 2mM
ammonium acetate and mobile phase B is composed of 50:50 acetonitrile:water containing
2mM ammonium acetate. A Phenomenex Luna® NH2 3 µm particle size (4.6x150mm) was
used for separation and column temperature was kept at 40°C. The total flow rate is 0.7
mL/min with a total run time of 17-minutes. Samples were extracted using the Bligh & Dyer
method. Lower layer was collected, dried down and resuspended in mobile phase A.

Cell culture. C2C12 myoblasts were grown and maintained in high-glucose Dulbecco's
modified Eagle's medium (DMEM), with 10% fetal bovine serum (FBS), and 0.1 %
penicillin/streptomycin. Once 90 to 100% confluent, C2C12 cells were differentiated into
myotubes with low-glucose DMEM, with l-glutamine and 110 mg/L sodium pyruvate;
supplemented with 2% horse serum, and 0.1% penicillin-streptomycin. For experiments with
erastin (E7781, MilliporeSigma), Ferrostatin-1 (SML0583, MilliporeSigma), and RSL3
(SML2234, MilliporeSigma), C2C12 myotubes were incubated with either 10 µM erastin/10
µM Ferrostatin-1/5 µM RSL3/ or equal-volume DMSO directly dissolved into medium. For
experiments with L-carnosine (C9625, MilliporeSigma), and N-acetylcarnosine (18817,
Cayman), C2C12 myotubes were incubated with 10 mM of L-carnosine/N-acetylcarnosine
directly dissolved into medium.

Lentivirus-mediated knockdown of GPx4/LPCAT3/ATG3 Lentivirus-mediated knockdown
of experiments were performed as previously described.[12,18,34] Vectors were decreased using
pLKO.1 lentiviral-RNAi system. Plasmids encoding short hairpin RNA (shRNA) for mouse
GPx4 (shGPx4: TRCN0000076552), mouse LPCAT3 (shLPCAT3: TRCN0000121437), and
mouse ATG3 (shATG3: TRCN0000247442) were obtained from MilliporeSigma. Packaging
vector psPAX2 (ID 12260), envelope vector pMD2.G (ID 12259), and scrambled shRNA
plasmid (SC: ID 1864) were obtained from Addgene. HEK293T cells in 10 cm dishes were

transfected using 50 μL 0.1% polyethylenimine, 200 μL 0.15 M sodium chloride, and 500 μL

Opti-MEM (with HEPES, 2.4 g/L sodium bicarbonate, and l-glutamine; Gibco 31985) with

2.66 μg of psPAX2, 0.75 μg of pMD2.G, and 3 μg of either scrambled or GPx4/LPCAT3/ATG3

shRNA plasmids. After 48 hours, growth medium was collected, filtered using 0.22 μm

vacuum filters, and used to treat undifferentiated C2C12 cells for 48 hours. To ensure that

only cells infected with shRNA vectors were viable, cells were selected with puromycin

throughout differentiation.

Measurements of myotube diameter. Images of myotubes were visualized at ×20

magnification using an inverted light microscope and captured with a camera (DP74,

Olympus). Myotube diameter was measured for at least 100 myotubes from 5 random fields

in each group using ImageJ software. The average diameter per myotube was calculated as

the mean of ten short-axis measurements taken along the length of the myotube.

Assessment of cell death. Cell death levels were examined by counting the numbers of

cells with trypan blue staining. The cells were trypsinized and stained with 0.2% trypan blue

for 5 min. Stained and non-stained cells were counted under a microscope using a

hemocytometer.

Immunofluorescence. C2C12 myotubes were fixed with 4% paraformaldehyde for 10 min

and permeabilized with 0.2% Triton X-100 for 15 min. After blocking with bovine serum

albumin, immunocytochemistry was performed with anti-HNE (ab48506, Abcam), anti-

lysosome associated membrane protein 2 (Lamp-2) (NB300-591, Novus), and Alexa Fluor-

conjugated secondary antibodies Alexa Fluor® 568 (ab175472, abcam), Alexa Fluor® 488

(ab150073, abcam), and DAPI (D1306, Invitrogen). Images were captured using a 63× 1.4

NA oil immersion objective on a Leica SP5 confocal system (Leica). For an experiment,

C2C12 myotubes incubated with erastin with or without pretreatment of BaFA1 (SML1661,

MilliporeSigma). Soleus muscles were embedded in optimal cutting temperature (OCT) gel

and sectioned at 10 μm with a cryostat (Microtome Plus). The sections underwent blocking

for 1 hr with M.O.M. mouse IgG Blocking Reagent (Vector Laboratories, MKB-2213), 1 hr

with primary antibodies (BA.D5, SC.71, BF.F3 all at 1:100 from DSHB). Sections were then

probed with the following secondary antibodies: Alexa Fluor 647 (1:250; Invitrogen, A21242),

Alexa Fluor 488 (1:500; Invitrogen, A21121), and Alexa Fluor 555 (1:500; Invitrogen, A21426).

Negative stained fibers were considered to be IIx. Slides were imaged with an automated

wide-field light microscope (Nikon Corp.) using a 10x objective lens. Cross-sectional area

and fiber type composition was then quantified utilizing ImageJ software.

Mitochondrial respiration measurements. Mitochondrial O_2 utilization was measured

using the Oroboros O$_2$K Oxygraphs, as previously described.[11,12] Isolated mitochondria were

added to the oxygraph chambers containing buffer Z. Respiration was measured in response

to the following substrate concentrations: 0.5 mM malate, 5 mM pyruvate, 2 mM ADP, 10 mM

succinate, and 1.5 μM FCCP.

Mitochondrial H$_2$O$_2$ measurements. Mitochondrial H$_2$O$_2$ production was measured using

the Horiba Fluoromax-4, as previously described.[11,12] Briefly, skeletal muscle was minced in

mitochondria isolation medium (300 mM sucrose, 10 mM HEPES, 1 mM EGTA) and

subsequently homogenized using a Teflon glass system. Homogenates were then

centrifuged at 800 g for 10 min, after which the supernatant was taken and centrifuged at

12,000 g for 10 min. The resulting pellet was carefully resuspended in mitochondria isolation

medium. JH$_2$O$_2$ was measured in buffer Z (MES potassium salt; 105 mM, KCl 30 mM,

KH$_2$PO$_4$ 10 mM, MgCl$_2$ 5 mM, and BSA 0.5 mg/ml) supplemented with 10 μM Amplex

UltraRed (Invitrogen) and 20 U/mL CuZnSOD in the presence of the following substrates: 10

mM succinate, 100 μM 1,3-bis(2-chloroethyl)-1-nitrosourea (BCNU/carmustine), and 1 μM

auranofin. The appearance of the fluorescent product was measured with excitation/emission

at 565/600 nm.

Administration of L-carnosine/N-acetylcarnosine in vivo. Carnosine was administered

as previously described (75). Briefly, young (4-month-old) or old (20-month-old) C57BL/6J

mice were supplemented with 80 mM carnosine dissolved in drinking water (pH 7.5) for 2

weeks (1 week of pretreatment and 1 week during HU). Bottles were refreshed two times a

week (L-carnosine, C9625, MilliporeSigma), and or everyday (N-acetyl carnosine, 18817,

Cayman).

MDA quantification. MDA content was quantified in fresh gastrocnemius muscles using a

lipid peroxidation assay kit (ab118970, Abcam) according to the manufacturer's instruction.

Rates of appearance of MDA-thiobarbituric acid adduct were quantified colorimetrically at

532 nm using a spectrophotometer.

Statistical analyses. Data are presented as means ± s.e.m. Statistical analyses were

performed using GraphPad Prism 7.03. Independent sample t-tests (two-sided) were used

to compare two groups. For multiple comparisons, one- or two-way analysis of variance

(ANOVA) were performed followed by appropriate post-hoc tests corrected for multiple

comparisons. For all tests $P < 0.05$ was considered statistically significant.

References

Scicchitano, B. M., Pelosi, L., Sica, G. & Musaro, A. The physiopathologic role of oxidative stress in skeletal muscle. *Mech Ageing Dev* **170**, 37-44, doi:10.1016/j.mad.2017.08.009 (2018).

Powers, S. K., Smuder, A. J. & Criswell, D. S. Mechanistic links between oxidative stress and disuse muscle atrophy. *Antioxid Redox Signal* **15**, 2519-2528, doi:10.1089/ars.2011.3973 (2011).

Evans, W. J. Skeletal muscle loss: cachexia, sarcopenia, and inactivity. *Am J Clin Nutr* **91**, 1123S-1127S, doi:10.3945/ajcn.2010.28608A (2010).

Larsson, L. *et al.* Sarcopenia: Aging-Related Loss of Muscle Mass and Function. *Physiol Rev* **99**, 427-511, doi:10.1152/physrev.00061.2017 (2019).

Wiernicki, B. *et al.* Excessive phospholipid peroxidation distinguishes ferroptosis from other cell death modes including pyroptosis. *Cell Death Dis* **11**, 922, doi:10.1038/s41419-020-03118-0 (2020).

Yang, W. S. *et al.* Regulation of ferroptotic cancer cell death by GPX4. *Cell* **156**, 317-331, doi:10.1016/j.cell.2013.12.010 (2014).

Anderson, E. J. *et al.* A carnosine analog mitigates metabolic disorders of obesity by reducing carbonyl stress. *J Clin Invest* **128**, 5280-5293, doi:10.1172/JCI94307 (2018).

Bhattacharya, A. *et al.* Denervation induces cytosolic phospholipase A2-mediated fatty acid hydroperoxide generation by muscle mitochondria. *J Biol Chem* **284**, 46-55, doi:10.1074/jbc.M806311200 (2009).

Pharaoh, G. *et al.* Targeting cPLA2 derived lipid hydroperoxides as a potential intervention for sarcopenia. *Sci Rep* **10**, 13968, doi:10.1038/s41598-020-70792-7 (2020).

Kagan, V. E. *et al.* Oxidized arachidonic and adrenic PEs navigate cells to ferroptosis. *Nat Chem Biol* **13**, 81-90, doi:10.1038/nchembio.2238 (2017).

Eshima, H. *et al.* Neutralizing mitochondrial ROS does not rescue muscle atrophy induced by hindlimb unloading in female mice. *J Appl Physiol (1985)* **129**, 124-132, doi:10.1152/japplphysiol.00456.2019 (2020).

Heden, T. D. *et al.* Mitochondrial PE potentiates respiratory enzymes to amplify skeletal muscle aerobic capacity. *Sci Adv* **5**, eaax8352, doi:10.1126/sciadv.aax8352 (2019).

Yant, L. J. *et al.* The selenoprotein GPX4 is essential for mouse development and protects from radiation and oxidative damage insults. *Free Radic Biol Med* **34**, 496-502, doi:10.1016/s0891-5849(02)01360-6 (2003).

Katunga, L. A. *et al.* Obesity in a model of gpx4 haploinsufficiency uncovers a causal role for lipid-derived aldehydes in human metabolic disease and cardiomyopathy. *Mol Metab* **4**, 493-506, doi:10.1016/j.molmet.2015.04.001 (2015).

Yoo, S. E. *et al.* Gpx4 ablation in adult mice results in a lethal phenotype accompanied by neuronal loss in brain. *Free Radic Biol Med* **52**, 1820-1827, doi:10.1016/j.freeradbiomed.2012.02.043 (2012).

Imai, H. & Nakagawa, Y. Biological significance of phospholipid hydroperoxide glutathione peroxidase (PHGPx, GPx4) in mammalian cells. *Free Radic Biol Med* **34**, 145-169, doi:10.1016/s0891-5849(02)01197-8 (2003).

Lee, J. Y., Kim, W. K., Bae, K. H., Lee, S. C. & Lee, E. W. Lipid Metabolism and Ferroptosis. *Biology (Basel)* **10**, doi:10.3390/biology10030184 (2021).

Ferrara, P. J. *et al.* Lysophospholipid acylation modulates plasma membrane lipid organization and insulin sensitivity in skeletal muscle. *J Clin Invest* **131**, doi:10.1172/JCI135963 (2021).

Ferrara, P. J. *et al.* Low lysophosphatidylcholine induces skeletal muscle myopathy that is aggravated by high-fat diet feeding. *FASEB J* **35**, e21867, doi:10.1096/fj.202101104R (2021).

20   Ran, Q. *et al.* Transgenic mice overexpressing glutathione peroxidase 4 are
protected against oxidative stress-induced apoptosis. *J Biol Chem* **279**, 55137-
55146, doi:10.1074/jbc.M410387200 (2004).
21   Dixon, S. J. *et al.* Ferroptosis: an iron-dependent form of nonapoptotic cell death.
Cell **149**, 1060-1072, doi:10.1016/j.cell.2012.03.042 (2012).
22   Codenotti, S. *et al.* Cell growth potential drives ferroptosis susceptibility in
rhabdomyosarcoma and myoblast cell lines. *J Cancer Res Clin Oncol* **144**, 1717-
1730, doi:10.1007/s00432-018-2699-0 (2018).
23   Cripps, M. J. *et al.* Carnosine scavenging of glucolipotoxic free radicals enhances
insulin secretion and glucose uptake. *Sci Rep* **7**, 13313, doi:10.1038/s41598-017-
13649-w (2017).
24   Everaert, I., De Naeyer, H., Taes, Y. & Derave, W. Gene expression of carnosine-
related enzymes and transporters in skeletal muscle. *Eur J Appl Physiol* **113**, 1169-
1179, doi:10.1007/s00421-012-2540-4 (2013).
25   Boldyrev, A. A., Aldini, G. & Derave, W. Physiology and pathophysiology of
carnosine. *Physiol Rev* **93**, 1803-1845, doi:10.1152/physrev.00039.2012 (2013).
26   Gao, H. *et al.* Ferroptosis is a lysosomal cell death process. *Biochem Biophys Res
Commun* **503**, 1550-1556, doi:10.1016/j.bbrc.2018.07.078 (2018).
27   Ichimura, Y. *et al.* A ubiquitin-like system mediates protein lipidation. *Nature* **408**,
488-492, doi:10.1038/35044114 (2000).
28   Chen, X., Yu, C., Kang, R., Kroemer, G. & Tang, D. Cellular degradation systems in
ferroptosis. *Cell Death Differ* **28**, 1135-1148, doi:10.1038/s41418-020-00728-1
(2021).
29   Gao, M. *et al.* Ferroptosis is an autophagic cell death process. *Cell Res* **26**, 1021-
1032, doi:10.1038/cr.2016.95 (2016).
30   Cai, J. *et al.* Autophagy Ablation in Adipocytes Induces Insulin Resistance and
Reveals Roles for Lipid Peroxide and Nrf2 Signaling in Adipose-Liver Crosstalk.
Cell Rep **25**, 1708-1717 e1705, doi:10.1016/j.celrep.2018.10.040 (2018).
31   McCarthy, J. J., Srikuea, R., Kirby, T. J., Peterson, C. A. & Esser, K. A. Inducible
Cre transgenic mouse strain for skeletal muscle-specific gene targeting. *Skelet
Muscle* **2**, 8, doi:10.1186/2044-5040-2-8 (2012).
32   Ferrara, P. J., Verkerke, A. R. P., Brault, J. J. & Funai, K. Hypothermia Decreases
O2 Cost for Ex Vivo Contraction in Mouse Skeletal Muscle. *Med Sci Sports Exerc*
50, 2015-2023, doi:10.1249/MSS.0000000000001673 (2018).
33   Verkerke, A. R. P. *et al.* Phospholipid methylation regulates muscle metabolic rate
through Ca(2+) transport efficiency. *Nat Metab* **1**, 876-885, doi:10.1038/s42255-
019-0111-2 (2019).
34   Johnson, J. M. *et al.* Alternative splicing of UCP1 by non-cell-autonomous action of
PEMT. *Mol Metab* **31**, 55-66, doi:10.1016/j.molmet.2019.10.007 (2020).

Acknowledgements

This research is supported by NIH DK107397, GM144613, AG063077, University of

Utah Center on Aging (to K.F.), AG050781 (to M.J.D.), HL122863, AG057006 (E.J.A),

AG064078 (to Q.R.), HL149870 (S.B.), HL139451 (to Z.S.M), DK130555 (to A.D.P.),

AG073493 (to J.J.P.), American Heart Association 18PRE33960491 (to A.R.P.V.) and

19PRE34380991 (to J.M.J.), Larry H. & Gail Miller Family Foundation (to P.J.F.), and

Uehara Memorial Foundation (to H.E.). We would like to thank Diana Lim from the

University of Utah Molecular Medicine Program for assistance with figures.

Author contributions

H.E. and K.F. contributed to study concept design and wrote the manuscript. P.S. and J.L.S.

contributed to study design and data analysis. M.J.P., J.A.M., and J.E.C. performed mass

spectrometry analyses. M.J.D. and Z.S.M. performed human muscle biopsies. H.E., P.S.,

J.L.S., A.W.L., J.M.J., J.J.P., and A.S., performed mouse experiments. H.E. performed all

biochemical assays, body composition measurements and analysis of muscle force

production. A.R.P.V. and P.J.F. assisted in muscle functional measurements. P.J.F. assisted

in cell culture experiments. E.J.A assisted in L-carnosine and N-acetylcarnosine experiments.

S.B. and Q.R. designed and generated mouse models. All contributed to editing the

manuscript.

Competing interests

No competing interests to disclose.

Materials & Correspondence

Correspondence and material requests should be addressed to K. Funai.

Figure legends

Fig. 1 | LOOH increases with aging and disuse in skeletal muscle. a,b GPx4 mRNA levels in skeletal muscle biopsy samples from young and old humans (n = 9 for young, n = 18 for old) (**a**) or skeletal muscles from young and old mice (n = 8 for young, n = 7 for old) (**b**). **c,d**, Oxidized phospholipid content in skeletal muscle from young and old mice (n = 8 per group). **e,f**, Soleus muscle mass (**e**), force-frequency curve (**f**) in soleus muscle following 1 and 7 days of HU from young WT C57BL/6J mice (n = 4 for sham, n = 6 for 1 day HU, n = 6 for 7 day HU). **g**, Immunoblotting of 4-HNE following 1 and 7 days of HU. **h**. A schematic on how pathways that regulate LOOH may promote muscle atrophy. PUFA: phospholipids containing polyunsaturated fatty acids. **i,j**, Representative images (**i**), and quantification (**j**) of myotube diameter (n = 104 for scrambled: SC, n = 107 for GPx4 KD, n = 117 for Vehicle, n = 120 for erastin, n = 104 for Vehicle, n = 110 for RSL3). Scale bar, 100 µm. Data are shown as the mean ± s.e.m. Statistical analyses in **a,b,c,d** and **j** were performed with an unpaired two-tailed t-test. Statistical analyses in **e** were performed with a one-way ANOVA with Dunnett's multiple comparisons test. Statistical analyses in **f** were performed with a two-way ANOVA and multiple comparisons were performed using Tukey's multiple comparisons tests.

Fig. 2 | Elevated LOOH is sufficient to augment disuse-induced muscle atrophy in young and old mice. a, Soleus muscle mass from young or old WT or GPx4+/- mice with or without HU (n = 8-11 per young group, n = 4-8 per old mice group). **b**, Force-frequency curve from old WT or GPx4+/- mice (n = 4-7 per group). **c**, Immunoblotting of 4-HNE from old WT or GPx4+/- mice. **d**, mRNA levels of GPx4 from young control or GPx4-MKO mice (n = 4 per group). **e,f**, Soleus muscle mass (**e**) (n = 7-8 per group) or force-frequency curve (**f**) from young control or GPx4-MKO mice (n = 4-7 per group). **g**, Immunoblotting of 4-HNE and GPx4 from young GPx4-MKO. **h,i**, Representative images of MHC immunofluorescence (**h**) and muscle fiber CSA by fiber type (**i**) for soleus muscles in young control or GPx4-MKO mice with HU (n = 4 per group). **j**, Immunoblotting of 4-HNE, GPx4 and Actin protein in C2C12 myotubes with or without GPx4 KD and/or LPCAT3 KD. **k,l**, Representative images (**k**), and quantification of myotube diameter (**l**) from C2C12 myotubes with or without GPx4 KD and/or without LPCAT3 KD (n = 104-114 per group). Scale bar, 100 µm. Data are shown as the mean ± s.e.m. Statistical analyses in **d** were performed with an unpaired two-tailed t-test. Statistical analyses in **l** were performed with a one-way ANOVA with Dunnett's multiple comparisons test. Statistical analyses in **a,c,e,f**, and **i** were performed with a two-way ANOVA and multiple comparisons were performed using Tukey's (**c,e,f,i**) or Sidak's (**a**) multiple comparisons tests.

Fig. 3 | Genetic and pharmacologic suppression of LOOH ameliorates muscle atrophy and weakness in young and old mice. a, Soleus muscle mass from young or old WT or GPx4Tg mice with or without HU (n = 6-11 per young group, n = 7 per old group). **b**, Force-frequency curve from old WT or GPx4Tg mice (n = 5-7 per group). **c**, Immunoblotting of muscle 4-HNE from old WT or GPx4Tg mice. **d,e**, Representative images of MHC immunofluorescence (**d**) and muscle fiber CSA by fiber type (**e**) for soleus muscles in old WT or GPx4Tg mice with HU (n = 4-6 per group). **f**, Immunoblotting of 4-HNE, and actin in C2C12 myotubes with or without GPx4 KD and/or Ferrostatin-1. **g,h**, Representative images (**g**), and quantification of myotube diameter (**h**) from C2C12 myotubes with GPx4 KD and/or Ferrostatin-1 treatments (n = 102-110 per group). Scale bar, 100 μm. **i**, Schematic illustration of the protocol for administration of N-acetylcarnosine in vivo. **j**, Immunoblotting of muscle 4-HNE from N-acetylcarnosine treatment in old mice. **k**, Soleus muscle mass from young or old mice with or without N-acetylcarnosine treatment (n = 6-8 per young group, n = 7 per old group). **l**, Force-frequency curve from N-acetylcarnosine study in old mice (n = 4-5 per group). **m,n**, Representative images of MHC immunofluorescence (**m**) and muscle fiber CSA by fiber type (**n**) for soleus muscles (n = 3-4 per group) in old mice from the N-acetylcarnosine study. Data are shown as the mean ± s.e.m. Statistical analyses in **h** was performed with a one-way ANOVA with Dunnett's multiple comparisons test, and Statistical analyses in **a,b,e,k,l**, and **n** were performed with a two-way ANOVA and were performed using Tukey's (**b,e,l,n**) or Sidak's (**a,k**) multiple comparisons tests.

Fig. 4 | Autophagy-lysosomal system amplifies LOOH to promote muscle atrophy. a,b, Representative images (**a**), and quantification of myotube diameter (**b**) from erastin-stimulated C2C12 myotubes with proteasomal inhibitor MG132 or lysosomal inhibitor BaFA1. (n = 21-120 per group). Scale bar, 100 μm. **c**, Autophagic flux analyses immunoblotting for LC3-I, LC3-II, p62, and actin in SC or GPx4 KD C2C12 myotuebs with or without BafA1. **d,e**, Representative images (**d**) and quantification of myotube diameter (**e**) from C2C12 myotubes with or without GPx4 KD and/or ATG3 KD) (n = 104-121 per group). **f**, mRNA levels of ATG3 (n = 8-32 per group) from young control or ATG3-MKO mice. **g**, Soleus muscle mass from control or ATG3-MKO mice (n = 10-12 per group). **h**, Force-frequency curve from young control or ATG3-MKO mice (n = 9-12 per group). **i,j**, Representative images of MHC immunofluorescence (**i**) and muscle fiber CSA by fiber type (**j**) of soleus muscles in young control or ATG3-MKO mice with HU (n = 6 per group). **k**, Immunoblotting of 4-HNE, GPx4, and actin in C2C12 myotubes with or without GPx4 KD and/or ATG3 KD. **l**, Confocal

immunofluorescence of LAMP2 (lysosome) and 4-HNE in erastin-stimulated C2C12

myotubes with or without BaFA1. **m**, Representative images of H_2O_2 or BCNU-stimulated

C2C12 myotubes with or without BaFA1. Scale bar, 100 µm. **n**, Immunoblotting of 4-HNE

and actin in H_2O_2-stimulated C2C12 myotubes with or without BafA1. **o**, A novel role of

lysosome in amplifying LOOH. Data are shown as the mean ± s.e.m. Statistical analyses in

f were performed with an unpaired two-tailed t-test. Statistical analyses in **b,e** was performed

with a one-way ANOVA with Dunnett's multiple comparisons test, and **g,h**, and **j** were

performed with a two-way ANOVA and multiple comparisons were performed using Tukey's

multiple comparisons tests.

Extended Data Fig. 1 (Related to age or disuse experiments in Figure 1). (**a**) Immunoblotting
of 4-HNE and GPx4 in skeletal muscle from young and old C57BL6/J mice. (**b**,**c**)
Quantification of 4- HNE (**b**) and MDA levels (**c**) (n = 5 per group) in young and old C57BL6/J
mice. (**d-f**) body mass (**d**), body composition (**e**), food intake (**f**), following 1, and 7 days of
HU from young WT C57BL/6J mice (n = 4 for sham, n = 6 for 1day HU, n = 6 for 7day HU).
(**g**) Quantification of 4- HNE (n = 3 per group). (**h**) Rates of oxygen consumption measured
in isolated skeletal muscle mitochondria with Krebs cycle substrates (n = 4 for sham, n = 5
for 1 day HU, n = 5 for 7 day HU). ADP, adenosine diphosphate; FCCP, carbonyl cyanide-p-
trifluoromethoxyphenylhydrazone; Mal, malate; Pyr, pyruvate. (**i**) Rate of mitochondrial H_2O_2
production in response to succinate or Aurorafin/BCNU (n = 4 for sham, n = 5 for 1 day HU,
n = 5 for 7 day HU). BCNU, 1,3-bis[2-chloroethyl]-1-nitrosourea. (**j**) Percentage of electron
leak in muscle mitochondria (n = 4 for sham, n = 5 for 1 day HU, n = 5 for 7 day HU). Data
are shown as the mean ± s.e.m. Statistical analyses in **b** and **c** were performed with an
unpaired two-tailed t-test. Statistical analyses in **d,e,f,g**, and **j** were performed with an one-
way ANOVA and Dunnett's multiple comparisons tests. Statistical analyses in **h**, and **i** were
performed with a two-way ANOVA and Tukey's multiple comparison test.

Extended Data Fig. 2 (Related to in vitro experiments in Figure 1). (**a**) GPx4 mRNA levels
in C2C12 myotubes with or without GPx4 knockdown (GPx4 KD) (n = 6 per group). (**b-d**)
Immunoblotting of 4-HNE, GPx4, and actin (**b**), quantification of 4-HNE (**c**) proteins and MDA
levels (**d**) (n = 3 per group). (**e**) mRNA levels for CHAC1 and PTGS2 (n = 6 per group). (**f**)
cell death levels in GPx4 KD myoblast or myotubes (n = 3 independent repeats). (**g-k**)
Quantification of 4-HNE (**g**), immunoblotting of 4-HNE (**h**) proteins, MDA levels (**i**), mRNA
levels for CHAC1 and PTGS2 (**j**), and cell death levels (**k**) in erastin-stimulated C2C12
myotubes (n = 3 independent repeats). (**l-o**) mRNA levels of GPx4 (**l**), immunoblotting of
GPx4 and actin (**m**), quantification of GPx4 (**n**), CHAC1 and PTGS2 (**o**) in RSL3-stimulated
C2C12 myotubes (n = 6 per group). Data are shown as the mean ± s.e.m. Statistical analyses
in **a,c,d,f,i,j,k,l,n** and **o** were performed with an unpaired two-tailed t-test. Statistical analyses
in **g** were performed with a one-way ANOVA and Dunnett's multiple comparisons tests.
Statistical analyses in **e** were performed with a two-way ANOVA and Tukey's multiple
comparison test.

Extended Data Fig. 3 (Related to GPx4+/- mice in Figure 2). (**a**) Body mass from young and
old WT or GPx4+/- mice. (**b**,**c**) body composition from young (n = 9 for WT sham, n = 8 for

GPx4 +/- sham, n = 7 for WT HU, n = 6 for GPx4 +/- HU) (**b**) and old (n = 4 for WT sham, n

= 5 for GPx4 +/- sham, n = 8 for WT HU, n = 6 for GPx4 +/- HU) (**c**) WT or GPx4+/- mice. (**d**)

Food intake during hindlimb unloading (HU) in young (n = 7 for WT, n = 6 for GPx4 +/-) and

old (n = 8 for WT, n = 6 for GPx4 +/-) WT or GPx4+/- mice. (**e**) EDL muscle mass from young

(n = 9 for WT sham, n = 8 for GPx4 +/- sham, n = 7 for WT HU, n = 6 for GPx4 +/- HU) and

old (n = 9 for WT sham, n = 8 for GPx4 +/- sham, n = 7 for WT HU, n = 6 for GPx4 +/- HU)

WT or GPx4+/- mice. (**f**) Quantification of muscle 4-HNE from old mice (n = 3 per group). (**g**)

MDA levels from old mice (n = 3 per group). (**h**) Force-frequency curve from soleus muscle

from young WT or GPx4+/- mice (n = 5 for WT sham, n = 5 for GPx4 +/- sham, n = 7 for WT

HU, n = 5 for GPx4 +/- HU). (**i,j**) Force-frequency curve in EDL muscle from young (n = 5 for

WT sham, n = 5 for GPx4 +/- sham, n = 7 for WT HU, n = 5 for GPx4 +/- HU) (**i**) and old (n =

3 for WT sham, n = 4 for GPx4 +/- sham, n = 6 for WT HU, n = 6 for GPx4 +/- HU) (**j**) WT or

GPx4+/- mice. Data are shown as the mean ± s.e.m. Statistical analyses in **a-j** were

performed with a two-way ANOVA and using Tukey's multiple comparisons tests.

Extended Data Fig. 4 (Related to GPx4-MKO mice in Figure 2). (**a**) Genotyping gel for

GPx4cKO+/+ mice crossed with HSA-MCM+/- mice to yield GPx4cKO+/+; HSA-MCM+/-

(GPx4-MKO) mice. Control littermates (GPx4cKO +/+; HSA-MCM -/-) were used for

experiments. Body mass (**b**), and body composition (**c**) (n = 7 for Ctrl sham, n = 4 for GPx4-

MKO sham, n = 7 for Ctrl HU, n = 8 for GPx4-MKO HU). (**d**) Food intake during HU (n = 7 for

Ctrl, n = 8 for GPx4-MKO). (**e,f**) EDL muscle mass (n = 6 for Ctrl sham, n = 4 for GPx4-MKO

sham, n = 7 for Ctrl HU, n = 8 for GPx4-MKO HU) (**e**) and force-frequency curve from young

mice (n = 6 for Ctrl sham, n = 4 for GPx4-MKO sham, n = 6 for Ctrl HU, n = 7 for GPx4-MKO

HU) (**f**). (**g**) Quantification of western blotting for 4-HNE from young mice (n = 3 per group).

(**h**) mRNA levels of CHAC1 and PTGS2 in skeletal muscles (n = 4 per group). (**i,j**) Mean

muscle fiber CSA (**i**) and fiber type composition (**j**) for soleus muscles following HU (*n* = 4

per group). Data are shown as the mean ± s.e.m. Statistical analyses in **d**, and **i**, were

performed with an unpaired two-tailed t-test. Statistical analyses in **b,c,e,f,g,h** and **j** were

performed with a two-way ANOVA and Tukey's multiple comparison test.

Extended Data Fig. 5 (Related to LPCAT3 KD studies in Figure 2). (**a**) mRNA levels of

LPCAT3 in C2C12 myotubes with or without LPCAT3 KD (n = 3 per group). (**b**) mRNA levels

of GPx4 and LPCAT3 in C2C12 myotubes with double knockdown for GPx4 and LPCAT3 (n

= 3 per group). (**c**) Quantification of 4-HNE proteins in C2C12 myotubes with or without

LPCAT3 KD and/or GPx4 KD (n=3 per group). (**d,e**) immunoblotting (**d**) and quantification (**e**) of 4-HNE from C2C12 myotubes with or without LPCAT 3 KD and/or erastin (n = 3 per group). (**f**) mRNA levels of LPCAT3 from erastin-stimulated myotubes (n = 3 per group). (**g,h**) Representative images of myotubes (**g**) and cell death levels (**h**) from C2C12 myotubes with or without LPCAT3 KD and/or erastin (n = 3 independent repeats). Scale bar, 100 μm. Data are shown as the mean ± s.e.m. Statistical analyses in **a**,**b**, and **f** were performed with an unpaired two-tailed t-test. Statistical analyses in **c**,**e**, and **h** were performed with a two-way ANOVA and Tukey's multiple comparison test.

Extended Data Fig. 6 (Related to GPx4Tg mice in Figure 3). (**a**) Quantification of GPx4 proteins from WT or GPx4Tg with or without HU (n = 3 per group). (**b**) Body mass from young (n = 9 for WT, n = 8 for GPx4Tg) and old (n = 7 for WT, n = 7 for GPx4Tg) mice. (**c,d**) body composition from young mice (**c**) and old mice (**d**). (**e**) Food intake during HU from young and old mice. (**f,g**) EDL muscle mass from young (n = 6 for WT sham, n = 8 for GPx4Tg sham, n = 9 for WT HU, n = 8 for GPx4Tg HU)(**f**) and old (n = 7 per group) (**g**) mice. (**h**) Force-frequency curve in soleus muscle from young mice (n = 5 for WT sham, n = 6 for GPx4Tg sham, n = 9 for WT HU, n = 8 for GPx4Tg HU). (**i,j**) Force-frequency curve in EDL muscle from young mice (n = 5 for WT sham, n = 7 for GPx4Tg sham, n = 9 for WT HU, n = 8 for GPx4Tg HU) (**i**) and old (n = 5 for WT sham, n = 5 for GPx4Tg sham, n = 5 for WT HU, n = 6 for GPx4Tg HU) (**j**) mice. (**k**) Quantification of muscle 4-HNE proteins from old mice (n = 3 per group). (**l,m**) Mean muscle fiber CSA (**l**) and fiber type composition (**m**) for soleus muscles from old mice following HU (n = 6 for WT, n = 4 for GPx4Tg). Data are shown as the mean ± s.e.m. Statistical analyses in **l** were performed with an unpaired two-tailed t-test. Statistical analyses in **a-m** were performed with a two-way ANOVA and Tukey's multiple comparison test.

Extended Data Fig. 7 (A pilot L-carnosine study leading to N-acetylcarnosine study in Figure 3). (**a-c**) Quantification of 4-HNE proteins (n = 2 per group) (**a**), cell death level (n = 3 independent repeats) (**b**), and mRNA levels of CHAC1 and PTGS2 with or without GPx4 KD and/or Ferrostatin-1 (n = 6 for SC, n = 5 for GPx4 KD) (**c**) in C2C12 myotubes. (**d-f**) Immunoblotting (**d**) and quantification (**e**) of 4-HNE, and actin (n = 3 per group) and cell death level (n = 3 independent repeats) (**f**) with or without erastin and/or L-carnosine in C2C12 myotubes. (**g-i**) Immunoblotting (**g**) and quantification (**h**) of 4-HNE, and actin (n = 3 per group), and (**i**) cell death level (n = 3 independent repeats) in C2C12 myotubes with or without erastin and/or L-carnosine. (**j-p**) Administration of L-carnosine in young (4-month-old)

C57BL/6J mice with or without HU. Body mass (**j**), body composition (n = 6 for Vehicle sham,

n = 7 for L-carnosine sham, n = 12 for vehicle HU, n = 13 for L-carnosine HU) (**k**), food intake

(**l**), water intake (n = 6 for Vehicle sham, n = 6 for L-carnosine sham, n = 6 for vehicle HU, n

= 7 for L-carnosine HU) (**m**), immunoblotting (**n**) and quantification (**o**) of 4-HNE (n = 3 per

group), and soleus muscle mass (n = 6 for Vehicle sham, n = 7 for L-carnosine sham, n = 12

for vehicle HU, n = 13 for L-carnosine HU) (**p**). Data are shown as the mean ± s.e.m.

Statistical analyses in **e**, and **f** were performed with a one-way ANOVA and Tukey's multiple

comparison test. Statistical analyses in **a**,**b**,**c**,**h**,**i**,**j**,**k**,**l**,**m**,**o** and **p** were performed with a two-

way ANOVA and Tukey's multiple comparison test.

Extended Data Fig. 8 (Related to N-acetylcarnosinestudies in Figure 3). (**a-c**)

Immunoblotting (**a**) and quantification (**b**) for 4-HNE, GPx4 and actin (n = 3 per group) and

cell death level (n = 3 independent repeats) (**c**) from C2C12 myotubes with or without GPx4

KD and/or N-acetylcarnosine. (**d-f**) Immunoblotting (**d**) and quantification (**e**) of 4-HNE and

actin (n = 3 per group) and cell death level (n = 3 independent repeats) (**f**) in C2C12 myotubes

with or without erastin and/or N-acetylcarnosine. (**g-s**) Administration of N-acetylcarnosine

from young (4-month-old) and old (20-month-old) C57BL/6J mice. Body mass (**g**), body

composition from young mice (**h**) and old mice (**i**). Food intake (**j**), water intake (**k**),

immunoblotting (**l**) and quantification (**m**) of 4-HNE (n = 3 per group). (**n**) EDL muscle mass

from young mice and old mice. (**o**) Force-frequency curve in soleus muscle from young mice

(n = 5 for vehicle sham, n = 6 for N-acetylcarnosine sham, n = 5 for vehicle HU, n = 5 for N-

acetylcarnosine HU). (**p**,**q**) Force-frequency curve in EDL muscle from young mice (n = 7 for

vehicle sham, n = 7 for N-acetylcarnosine sham, n = 6 for vehicle HU, n = 7 for N-

acetylcarnosine HU) (**p**) and old mice (n = 6 for vehicle sham, n = 7 for N-acetylcarnosine

sham, n = 6 for vehicle HU, n = 6 for N-acetylcarnosine HU) (**q**). (**r**,**s**) Average of muscle fiber

CSA (**r**) and fiber type composition (**s**) for soleus muscles from old mice following HU (*n* = 3

for vehicle, n = 4 for N-acetylcarnosine). Data are shown as the mean ± s.e.m. Statistical

analyses in **r** were performed with an unpaired two-tailed t-test. Statistical analyses in **e**, and

f were performed with a one-way ANOVA and Tukey's multiple comparison test. Statistical

analyses in **b**,**c**,**g**,**h**,**i**,**j**,**k**,**m**,**n**,**o**,**p**,**q**, and **s** were performed with a two-way ANOVA and

Tukey's multiple comparison test.

Extended Data Fig. 9 (Related to in vitro experiments in Figure 4). (**a**) Cell death levels in

erastin-stimulated myotubes with MG132 or BaFA1 (n = 3 independent repeats). (**b**,**c**)

Representative images (**b**) and cell death level (**c**) in RSL3-stimulated myotubes with BaFA1

(n = 3 independent repeats). Immunoblotting (**d**) and quantification (**e**) of LC3-I, LC3-II, p62

(all static measurements without BafA1), and actin protein from C2C12 myotubes with or

without GPX4 KD (n = 4 per group). (**f-h**) quantification of western blotting for LC3-I (**f**), LC3-

II (**g**), and p62 (**h**) protein from GPx4 KD myotubes with or without BaFA1 (n = 3 per group).

(**i**) mRNA levels of ATG3 in C2C12 myotubes with or without (n = 3 per group). (**j**) mRNA

levels of GPx4 and ATG3 from C2C12 myotubes with double knockdown for ATG3 and GPx4

(n = 3 per group). (**k**,**l**) Immunoblotting (**k**) and quantification (**l**) of LC3, p62, and GPx4 protein

from C2C12 myotubes with or without ATG3 KD (n = 3 per group). Data are shown as the

mean ± s.e.m. Statistical analyses in **c**,**i** and **l** were performed with an unpaired two-tailed t-

test. Statistical analyses in **a** were performed with a one-way ANOVA and Tukey's multiple

comparison test. Statistical analyses in **e**,**f**,**g**,**h**,**i**, and **j** were performed with a two-way

ANOVA and Tukey's multiple comparison test.

Extended Data Fig. 10 (Related to in vivo and in vitro experiments in Figure 4). (**a**)

Genotyping of ATG3cKO+/+ mice crossed with HSA-MCM+/- mice to yield ATG3cKO+/+;

HSA-MCM+/- (ATG3-MKO) mice. Control littermates (ATG3cKO +/+; HSA-MCM -/-) were

used for experiments. (**b**) Immunoblotting of p62, LC3, and GAPDH proteins from skeletal

muscles from control and ATG3-MKO mice. (**c**) Quantification of LC3-II/I ratio in muscles

from control and ATG3-MKO mice (n = 8 per group). (**d**) Body weight, (**e**) lean mass, and (**f**)

fat mass from control and ATG3-MKO mice (control sham n = 9, ATG3-MKO sham n = 10,

control HU n = 10, ATG3-MKO HU n = 10). (**g**) EDL muscle mass (control sham n = 10,

ATG3-MKO sham n = 12, control HU n = 10, ATG3-MKO HU n = 10). (**h**) Force frequency

curve for EDL muscle (control sham n = 9, ATG3-MKO sham n = 8, control HU n = 10, ATG3-

MKO HU n = 9). (**i**) Fiber type composition for SOL muscle (control HU n = 6, ATG3MKO HU

n = 6). (**j**) quantification of 4-HNE protein from C2C12 myotubes with or without GPx4 KD

and/or ATG3 KD (n = 3 per group). (**k**,**l**) Immunoblotting (**k**) and quantification (**l**) of 4-HNE

protein from C2C12 myotubes with or without erastin and/or ATG3 KD (n = 3 per group).

(**m**,**n**) Immunoblotting (**m**) and quantification (**n**) of 4-HNE protein from C2C12 myotubes with

or without erastin and/or BaFA1 (n = 3 per group). (**o**,**p**) Immunoblotting (**o**) and quantification

(**p**) of 4-HNE protein from C2C12 myotubes with or without RSL3 and/or BaFA1 (n = 3 per

group). (**q**) Confocal fluorescence microscope images of erastin-stimulated myotubes with or

without BaFA1. Boxed regions are shown enlarged at far right. 4-HNE protein from H_2O_2-

stimulated (**r**), or BCNU-stimulated (**s**,**t**) C2C12 myotubes with or without BafA1. Data are

shown as the mean ± s.e.m. Statistical analyses in **c**,**d**,**e**,**f**,**g**,**h**,**i**,**j**,**l**,**n**,**p**,**r** and **t** were performed

with a two-way ANOVA and Tukey's multiple comparison test.

Figure 1

K. Funai, et al.

Figure 2

K. Funai, et al.

Figure 3

Figure 4

K. Funai, et al.

K. Funai, et al.

ED09

K. Funai, et al.