1

Possible Cross Reactivity of Feline and White-tailed Deer Antibodies Against the SARS-CoV-2

Receptor Binding Domain

Trevor J. Hancock[1], Peyton Hickman[1], Niloo Kazerooni[2], Melissa Kennedy[2], Stephen A. Kania[2],

Michelle Dennis[2], Nicole Szafranski[2], Richard Gerhold[2], Chunlei Su[1], Tom Masi[3], Stephen

Smith[4], Tim E. Sparer[1]*

[1] Department of Microbiology, University of Tennessee, Knoxville, Tennessee, USA

[2] Department of Biomedical and Diagnostic Sciences, University of Tennessee, Knoxville,

Tennessee, USA

[3] Graduate School of Medicine, University of Tennessee Medical Center, Knoxville, Tennessee,

USA

[4] MEDIC Regional Blood Center, Knoxville, Tennessee, USA

* Corresponding author

Email: tsparer@utk.edu (TES)

Abstract

In late 2019, a novel coronavirus began circulating within humans in central China. It was designated SARS-CoV-2 because of its genetic similarities to the 2003 SARS coronavirus (SARS-CoV). Now that SARS-CoV-2 has spread worldwide, there is a risk of it establishing new animal reservoirs and recombination with native circulating coronaviruses. To screen local animal populations in the United States for exposure to SARS-like coronaviruses, we developed a serological assay using the receptor binding domain (RBD) from SARS-CoV-2. SARS-CoV-2's RBD is antigenically distinct from common human and animal coronaviruses allowing us to identify animals previously infected with SARS-CoV or SARS-CoV-2. Using an indirect ELISA for SARS-CoV-2's RBD, we screened serum from wild and domestic animals for the presence of antibodies against SARS-CoV-2's RBD. Surprisingly pre-pandemic feline serum samples submitted to the University of Tennessee Veterinary Hospital were ~50% positive for anti-SARS RBD antibodies. Some of these samples were serologically negative for feline coronavirus (FCoV), raising the question of the etiological agent generating anti-SARS-CoV-2 RBD cross-reactivity. We also identified several white-tailed deer from South Carolina with anti-SARS-CoV-2 antibodies. These results are intriguing as cross-reactive antibodies towards SARS-CoV-2 RBD have not been reported to date. The etiological agent responsible for seropositivity was not readily apparent, but finding seropositive cats prior to the current SARS-CoV-2 pandemic highlights our lack of information about circulating coronaviruses in other species.

Importance

We report cross-reactive antibodies from pre-pandemic cats and post-pandemic South Carolina white-tailed deer that are specific for that SARS-CoV RBD. There are several potential explanations for this cross-reactivity, each with important implications to coronavirus disease surveillance. Perhaps the most intriguing possibility is the existence and transmission of an etiological agent (such as another coronavirus) with similarity to SARS-CoV-2's RBD region. However, we lack conclusive evidence of pre-pandemic transmission of a SARS-like virus. Our findings provide impetus for the adoption of a One Health Initiative focusing on infectious disease surveillance of multiple animal species to predict the next zoonotic transmission to humans and future pandemics.

Introduction

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is an emergent zoonotic
beta-coronavirus initially identified in late 2019 after human-to-human transmission within
central China [1]. By early 2020, the virus had caused a pandemic infecting millions of people
and continues to circulate throughout the world. Like other human coronaviruses it is spread *via*
aerosolized particles, leading to respiratory infections [1, 2]. Infected individuals develop a range
of symptoms from mild/asymptomatic infection to severe pneumonia-like disease (i.e.,
coronavirus disease (COVID)) [2]. Sequence analysis of known SARS coronaviruses points to a
bat origin with probable intermediate hosts prior to human adaptation [3-5]. However, the exact
intermediate host and factors that led to its zoonosis and establishment within humans are under
investigation.

Secretion of SARS-CoV-2 is thought to be primarily *via* aerosolized particles with high
viral loads in the lungs and nasopharyngeal secretions of infected individuals [6, 7]. However,
both viral RNA and infectious particles have been detected in fecal samples of acutely infected
individuals. In the original SARS-CoV outbreak, there was documented fecal-oral transmission
of infection [6, 8-12]. Fecal to oral spread and shedding is a common route of transmission of
other animal coronaviruses. Oropharyngeal viral RNA shedding of SARS-CoV-2 in humans lasts
for ~17 days on average but persists up to 60-120 days in the respiratory tract and stool [13].
Similarly, oropharyngeal secretion of infectious SARS-CoV-2 in cats appears to cease by 5-10
days post infection (dpi) [14]. Infected felids shed SARS-CoV-2 viral RNA in their feces for at
least 5 dpi, but whether that represents infectious virus, or a potential route of transmission is yet
to be demonstrated [15].

Due to the multiple routes of spread and close contact with other species, transmission of
SARS-CoV-2 from humans to animals is plausible [16]. Human-to-animal and animal-to-animal
transmission of SARS-CoV-2 has been documented or experimentally demonstrated. Companion
animals such as cats and dogs are susceptible to experimental as well as natural infection from
COVID-positive owners [14, 17-21]. In addition, susceptible animals are capable of transmitting
infection to cohoused animals [14, 22]. In the case of minks, transmission from humans-to-minks
and back to humans was demonstrated [23]. This is similar to a situation at the Amoy Garden
complex during the original SARS-CoV outbreak where animal-to-human transmission occurred
when an animal vector potentially contracted and spread SARS-CoV throughout the complex
[24-26]. Human transmission of SARS-CoV-2 to companion animals opens up the potential for

spillover into wild animal populations. Indeed, transmission from humans to deer within North
America has been proposed, as post-pandemic deer have been demonstrated seropositive in
multiple states and SARS-CoV-2 genome sequenced from lymph nodes [27-29]. Human to
animal transmission could contribute to the spread of SARS-like coronaviruses and the
establishment of new reservoirs for recombination and the generation of future novel coronavirus
outbreaks.

Infected humans and animals mount humoral responses to SARS-CoV-2 [13, 14, 30-32].
In humans, SARS-CoV-2 antibodies arise within 5-14 days post-infection/symptom onset and
peak around 17-20 dpi [13, 31, 33]. For cats experimentally inoculated or naturally exposed to
SARS-CoV-2, detectable antibody titers appeared by 7-14 days post-infection peaking ~21 dpi
[14]. This matches anti-FCoV responses where high antibody levels can arise within ~9 dpi [34-
36]. Immunity to coronaviruses in cats is typically short-lived, with the average FCoV humoral
responses lasting several months to 2 years [37]. Anti-SARS-CoV-2 RBD responses in
seropositive cats had similar declines in antibody titers only lasting around 4-5 months [38].
However, humans infected with the initial SARS-CoV mounted robust responses detectable 1-2
years post exposure [39-41]. The duration of anti-SARS-CoV-2 antibody responses is the subject
ongoing research, but natural exposure is unlikely to induce long-term or lifelong
immunity/seropositivity [42].

Major antigenic targets for SARS-CoV-2 infected individuals are the nucleocapsid,
which is one of the most abundantly produced viral proteins [43], and spike protein, which is
responsible for viral entry [44]. The spike has high immunogenicity and diverges from other
coronaviruses [32, 44, 45]. Spike is composed of two subunits (S1/S2). The S1 subunit contains
the receptor binding domain (RBD) responsible for binding to host ACE-2 and determining
tropism/entry, while the S2 domain contains the fusogenic region of the spike [44, 45]. SARS
coronaviruses share very low similarity to other coronaviruses within the spike protein [32], but
antibodies against the S2 subunit can cross-react with common human coronaviruses [46-48].
Cross-reactivity of the S1 subunit occurs at very low rates. Within the S1 region, the RBD is
highly immunogenic and unique to SARS-CoV-2 [32, 49]. Serum from humans infected with
common human coronaviruses such as OC43, NL63, and 229E failed to recognize the RBD from
SARS-CoV-2 [32, 46, 49]. Animals infected or immunized with other coronaviruses similarly
fail to generate cross-reactive antibodies against SARS-CoV-2's RBD [32]. For infected cats,

SARS-CoV-2 seroconversion was not impacted by pre-existing immunity against feline
coronavirus (FCoV), an alpha-coronavirus with limited similarity to SARS-CoV-2 [38].
Collectively, seropositivity against the RBD of SARS-CoV-2 is a specific marker of SARS-
CoV-2 exposure and has led several groups to create highly specific indirect ELISAs against
SARS-CoV-2's RBD to screen for SARS-CoV-2 exposure [30, 32, 33]. A final consideration of
antibodies targeting the RBD is they could be either neutralizing or non-neutralizing [33, 50-53].
This may explain why serum from humans and animals exposed to the original SARS-CoV were
able to recognize the spike and RBD of SARS-CoV-2 while their cross-neutralization potential
was variable [54, 55].

Despite limited similarity in the spike protein of SARS-CoV-2 vs common circulating
coronaviruses, there are reports of pre-pandemic, pre-existing SARS-CoV-2 reactive serum in
humans [48, 49, 54]. These cross-reactive antibodies represent a rare response to common human
coronaviruses within conserved epitopes of SARS-CoV-2's spike protein (usually in the S2
region) with reports of ~0.6% prevalence of pre-existing anti-RBD responses [46, 48, 49].
Although there is increasing evidence for earlier timelines of SARS-CoV-2 spread among
humans, pre-existing seropositivity among other species has not been reported [38, 56-58].
Indeed, even within central China, researchers failed to find evidence of SARS-CoV-2 exposure
prior to the pandemic [38, 56, 58].

As SARS-CoV-2 spreads and encounter's new species, there is a need for monitoring
local populations for SARS-CoV-2 transmission and the potential establishment of local
reservoirs. Currently, we have a limited understanding of coronavirus reservoirs, spread, and
recombination among diverse species. The original SARS outbreak in 2003 was a harbinger of
the potential risk of crossover coronaviruses. At that time, animal coronavirus surveillance was a
high priority. Unfortunately, this investment was not sustained. Our aim was to address whether
SARS-CoV-2 is being introduced into companion animals of North America by tracking
seroconversion using an in-house indirect ELISA against the RBD of SARS-CoV-2. We chose to
focus on companion animals (i.e., cats and dogs) as they represent a significant source of human-
animal interactions with potential for contact and further spillover into wild animal populations.
Surprisingly, we found evidence of anti-RBD seropositive animals pre-dating the pandemic by
several months to years. Our study provides evidence for the existence and prevalence of SARS-
CoV-2 serum reactivity prior to the current pandemic.

Materials and methods

Recombinant RBD production and purification

Recombinant RBD production has been previously published [56]. Our lab deviated from the prior published method to utilize equipment readily available. Briefly, the plasmid containing the RBD of SARS-CoV-2 was produced under federal contract HHSN272201400008C and obtained through BEI Resources, NIAID, NIH. Vector pCAGGS contains the SARS-related coronavirus 2, Wuhan-Hu-1 spike glycoprotein RBD, NR-52309. To produce recombinant RBD, the pCAGGS-RBD plasmid was transfected into ~$5x10^7$ adherent HEK-293/T17 cells (ATCC CRL-11268) in a T-175 using PEI (Polyethylenimine, linear 25,000 m.w.(Polysciences, Warrington, PA, USA). Plasmid was mixed at a 1:3 ratio with PEI (20ug of plasmid : 60ug PEI for a T-175 transfection) in 1 mL serum-free DMEM for 30 minutes at room temperature. Media was aspirated and the transfection mixture was added to 14mL fresh growth media and placed onto cells. Three to four hours post-transfection, media was changed and replaced with DMEM containing either 2% or 5% Fetal Bovine Serum (Hyclone FetalClone III, Cytiva Life Sciences, USA). Maintenance in a lower serum prevents overgrowth. However, we found higher protein yields when supplemented with 5% FBS. Transfection efficiency was nearly 100% as assessed by GFP-positive transfected cells in a control flask.

Supernatants from transfected HEK-293 T17 cells were collected into 50mL conical tubes and frozen at days 3 and 6 post-transfection. Pooled supernatants were thawed and incubated with Ni-NTA (Ni-NTA Agarose, Qiagen, Germany) resin with gentle rocking overnight. The resin was spun down at >3400x g in a swing-bucket Sorvall RT centrifuge for 10 minutes at 4^0C. Ni-NTA resin was resuspended in 1mL wash buffer (20mM imidazole, 5mM $NaH_2PO_4.H_2O$, 0.3M NaCl in H_20), transferred to a 2mL microcentrifuge tube, gently rocked for 10 minutes at room temperature, spun, and resuspended in fresh buffer. Resin was washed between 3-7 times until OD_{230} was ≤ wash buffer. Once the supernatant OD dropped sufficiently, 1mL elution buffer (235mM Imidazole, 5mM $NaH_2PO_4.H_2O$, 0.3M NaCl in H_20) was added to elute the RBD from the nickel resin. Eluate was rocked for 10 minutes at room temp and then centrifuged. Two elution steps were performed with a third final elution using 0.5M imidazole. Protein concentration was determined by standard curve analysis of a silver-stained (Pierce

Silver Stain Kit, Thermo Scientific, USA) 12% SDS-PAGE gel using a standard curve of BSA (bovine serum albumin). Analysis was performed using Image Studio Lite ver. 5.3 (Li-Cor Biosciences, Lincoln, NE, USA).

Serum and plasma samples

Previously collected pre-SARS-CoV-2 de-identified human serum samples were kindly donated by Dr. Jon Wall and Steve Foster (University of Tennessee Medical Center, Knoxville, Tennessee, USA). De-identified COVID-positive plasma samples were donated from: MEDIC Regional Blood Center (Knoxville, Tennessee, USA) and Dr. Mark Slifka (Oregon Health Sciences University, Portland, Oregon, USA). The following reagents were obtained through BEI Resources, NIAID, NIH: Human Plasma, Sample ID WU353-073, NR-53643; WU353-074, NR-53644; WU353-075, NR-53645; WU353-076, NR-53646; WU353-076, NR-53647, were contributed by Ali Ellebedy, Ph.D., Washington University School of Medicine, St. Louis, Missouri, USA. The following reagents were obtained through BEI Resources, NIAID, NIH: polyclonal anti-feline infectious peritonitis virus, 79-1146 (antiserum, guinea pig), NR-2518; polyclonal anti-canine coronavirus, UCD1 (antiserum, guinea pig), NR-2727; polyclonal anti-bovine coronavirus, Mebus (antiserum, guinea pig), NR-455; polyclonal anti-porcine respiratory coronavirus, ISU-1 (antiserum, guinea pig), NR-459; polyclonal anti-turkey coronavirus, Indiana (antiserum, guinea pig), NR-9465. Client-owned canine and feline serum samples were submitted to the University of Tennessee Veterinary Hospital for routine animal testing. Canine samples (n=36) were collected post-pandemic from local, East Tennessee animals. All feline samples were pre-pandemic. Samples from East Tennessee feral cats (n=36) were collected from 2007-2012. Client-owned feline samples (n=92) were collected nationwide as part of clinical diagnostic testing (see Table 1). Twenty cat samples were grouped into FCoV positive and negative groups based on feline infectious peritonitis (FIP) serology using an immunofluorescence assay (IFA) against FIP serotypes I and II, as well as TGEV (transmissible gastroenteritis virus) (VMRD, Pullman, WA, USA). Normal cat serum was purchased from Jackson ImmunoResearch (West Grove, PA, USA). Tennessee-resident cows (n=33) and tigers (n=9) were collected pre-pandemic for routine diagnostic testing. Post-mortem, post-pandemic samples were collected from East Tennessee elk (n=12) and South Carolina white-tailed deer (n=22).

Anti-RBD ELISA

Anti-RBD ELISA was based on the published protocol by Amanat et al. and Stadlbauer
et al. [30, 59]. Purified RBD was diluted to 2ug/mL in PBS and 50uL was placed into each well
of a 96 well plate (Immulon 4HBX, Thermo Fisher, USA) and allowed to incubate overnight at
4^0C. Unbound RBD was removed and wells were washed 3x with PBS-T (PBS with 0.1%
Tween-20). Rinsed wells were blocked with 5% milk in PBS for 2 hours at room temp. Block
was removed and serum or plasma samples were added at 1:50 dilution for the initial screen in
PBS with 1% milk and incubated at room temp for 1 hour. After 1 hour, wells were washed 3x
with PBS-T and a secondary antibody for that species was added (i.e., HRP goat-anti-human
IgG, Rockland Immunochemicals, Pottstown, PA, USA; HRP goat-anti-dog IgG, Bethyl
Laboratories, Montgomery, TX, USA; HRP goat-anti-cat IgG, Invitrogen, Waltham, MA, USA;
HRP goat-anti-guinea pig IgG, Life Technologies Corp, Carlsbad, CA, USA; HRP rabbit-anti-
deer IgG, KPL, Gaithersburg, MD, USA; HRP sheep-anti-cow IgG, Bethyl Laboratories,
Montgomery, TX, USA) at dilutions of 1:10,000 (anti-human, cat, dog, tiger) or 1:250 (anti-cow,
deer, elk) in PBS with 1% milk. Optimal secondary antibody concentrations were determined by
titration on either 5% milk (negative control) or 1:50 dilution of that species serum (positive
control). Secondary antibodies were allowed to incubate for 1 hour at room temperature before
being washed 3x with PBS-T. ELISA was developed with 50uL TMB (1-Step Ultra TMB-
ELISA, Thermo Fisher, Waltham, MA, USA) for 10 minutes. Reactions were stopped by the
addition of 2M sulfuric acid and read using a BioTek Synergy 2 or Synergy H1 plate reader set at
450nm (BioTek, Winooski, VT, USA). Receiver operator curve (ROC) analysis was performed
to the determine the appropriate threshold to yield 100% specificity of ELISAs performed at a
1:50 dilution. For titrations of seropositive and seronegative samples, threshold values for each
dilution were calculated as the average of negative samples plus 3 times the standard deviation.
Titrated samples were initially diluted 1:100 and then serially diluted 1:3 (final dilution of
1:8100). Serum dilutions were made in PBS with 1% milk and added to RBD-coated and
blocked wells. OD_{450} values for each species and titrations were graphed in GraphPad Prism ver.
8 (GraphPad Software, San Diego, CA, USA).

Western and dot blots

For dot blots, 5-10uL of sample was applied directly onto nitrocellulose membranes and
allowed to dry. Western blots were loaded with 30uL (~3ug) of purified recombinant RBD,
resolved in a 12% SDS-PAGE gel and transferred to a nitrocellulose membrane. Blots were

blocked overnight at 4^0C with 5% milk in PBS. Mouse anti-6His-HRP conjugated antibody

(1:5,000) (Proteintech, Rosemont, IL, USA) or polyclonal serum samples (1:20) were incubated

with the blots at room temperature for 2 hours and subsequently washed 2x with TBS-T (tris

buffered saline with 0.1% Tween-20). For polyclonal serum, species specific HRP anti-IgG

antibodies (1:5,000 dilution) were incubated for an additional 2 hours and washed 2x as above.

Chemiluminescent substrate (Pierce SuperSignal West Pico PLUS, Thermo Fisher, USA) was

added and luminescence was detected using BioRad ChemiDoc (Bio-Rad, Hercules, CA, USA).

Neutralization assays

Serum samples were screened for neutralization using LEGENDplex SARS-CoV-2

neutralizing antibody assay (BioLegend, San Diego, CA, USA) following manufacturer

recommendations. Briefly, serum was diluted 1:100 and incubated with biotinylated SARS-CoV-

2 S1 subunit containing the RBD and human ACE-2 (hACE-2) conjugated to fluorescent beads.

Streptavidin-PE (phycoerythrin) was added to detect SARS-CoV-2 S1 subunit bound to

beads/hACE-2. Binding/PE levels were detected *via* a BD LSR-II equipped with 488 and 633nm

lasers (Becton Dickinson, Franklin Lakes, NJ, USA). Data was analyzed using BioLegend

LEGENDplex Data Analysis Software. Mean fluorescent intensity (MFI) was normalized and

graphed in GraphPad prism v9 (GraphPad Software, San Diego, CA, USA).

Fecal coronavirus PCR screen and sequence alignments

De-identified fecal samples from thirty healthy East Tennessee cats were collected and

stored at -80^{o}C. Samples were resuspended in PBS to yield a 10% solution and centrifuged to

clarify. Fecal RNA was extracted using a Qiagen viral RNA extraction kit (Qiagen, Hilden,

Germany) and RNA was reverse transcribed using Verso cDNA kit with random hexamers and

RT enhancer (Thermo Fisher, Waltham, MA, USA). PCR amplification of conserved

coronavirus regions using previously reported primer pairs was used to screen the cDNA [60].

PCR amplicons were visualized on a 1% agarose gel and positive PCR samples were Sanger

dideoxy sequenced. Sequences were viewed using 4Peaks software (Nucleobytes, Amsterdam,

Netherlands). Sequences from fecal samples were Clustal W aligned to common coronaviruses,

trimmed and phylogenetic trees were generated using Maximum-Likelihood method for each

positive loci using MEGA X [61, 62]. For phylogenetic trees using multiple loci, aligned and

trimmed sequences for each loci were concatenated together prior to Maximum-Likelihood tree

construction. Phylogenetic trees were tested by bootstrap testing with 1000 iterations. Common
coronavirus sequences for ORF1ab (RdRp and helicase loci) and spike were obtained from the
following: porcine coronavirus HKU15 (NC039208), SARS-CoV-2 (MN988668), SARS-CoV
(NC004718), porcine respiratory coronavirus/PRCoV (KY406735), human coronavirus OC43
(NC006213), MERS-CoV (NC038294), feline coronavirus/FCoV (NC002306), canine
respiratory coronavirus/CRCoV (KX432213), canine coronavirus/CCoV (JQ404410), bovine
coronavirus/BCoV (NC003045), avian coronavirus (NC048214), human coronavirus 229E
(NC002645), transmissible gastroenteritis virus/TGEV (NC038861), murine hepatitis
virus/MHV (NC048217), human coronavirus NL63 (NC005831), feline coronavirus strain UU8
(FJ938055), feline coronavirus strain UU19 (HQ392470), feline coronavirus strain Black
(EU186072), feline coronavirus strain RM (FJ938051), feline coronavirus strain Felix
(MG893511). All generated sequences were deposited in GenBank under accession numbers:
MZ220722 through MZ220762 (sample ID, positive loci, animal location, and accession
numbers are shown in Table 2).

Statistics

All graphs and statistical analysis were performed in GraphPad Prism ver. 9 (GraphPad
Software, San Diego, CA, USA). ELISA OD_{450} results were normalized to an inter-plate
replicate run with all assays. Student's one-tailed t-tests with Welch's correction and one-way
ANOVA with multiple comparisons tests were performed on ELISA results and documented in
the respective figure legends. Descriptive statistics were provided for each ELISA group (mean,
median, and quartiles). Receiver operator curve (ROC) analysis was performed to determine
appropriate threshold values for human, cat, and deer serum samples. Area under the curve was
calculated for each titrated ELISA sample and graphed. Neutralization data was normalized with
negative control group (normal cat serum) representing 100% MFI.

Data will be made publicly available upon publication and upon request for peer review.
294

Results

We developed an in-house ELISA to serologically screen companion animals based on a
protocol developed at Mt. Sinai [30, 59]. To examine cross-reactivity of our in-house anti-SARS-
CoV-2 RBD indirect ELISA, we used polyclonal guinea pig serum raised against different

animal coronaviruses (Fig 1A). Consistent with previous reports, no cross-reactive antibodies for

any of the common coronaviruses were found [18, 32, 38, 56]. Only antibodies from SARS-CoV

or SARS-CoV-2 infected individuals reacted (Fig 1A, 1B). Human serum collected from

individuals prior to the SARS-CoV-2 pandemic or plasma from recovered SARS-CoV-2 donors

were used to validate our ELISA screen (Fig 1B). ROC analysis determined the positive cutoff

threshold, using a value that gave highest specificity and sensitivity with pre-pandemic human

serum and serum from confirmed SARS-CoV-2 infected individuals. ROC analysis was in

agreement with the commonly used threshold determination method of three standard deviations

above the mean negative value. Our assay based on RBD screening showed high sensitivity

(96.96%) and specificity (95.45%) with 66 SARS-CoV-2 samples and 22 pre-SARS-CoV-2

samples (Fig 1B). While Stadlbauer et al. performed two diagnostic ELISAs, one with RBD and

the other with full-length spike, our results using only the RBD-based screen are in good

agreement with their published data. Others have also demonstrated the accuracy of an RBD-

only based ELISA [33, 38]. A western blot using an anti-6His antibody (Fig 1C) shows the

expected size of purified RBD with a single band ~32kDa. This shows that our isolated RBD is

the correct size and runs as a monomer. Silver stain of the same affinity-purified SARS-CoV-2

RBD demonstrates relative purity (Fig 1D). However, there are co-purified proteins present at

lower levels.

To establish a baseline for future SARS-CoV-2 screening of companion animals, 128

pre-pandemic feline serum samples collected prior to December 2019 were retrospectively

screened using our in-house ELISA. Nineteen samples were of a known FCoV serological status,

with the remaining 109 of unknown FCoV status. Following the same protocol used for

screening human serum samples (Fig. 1), feline samples were tested for antibodies against

SARS-CoV-2's RBD (Fig 2A). There were two batches tested. Serum samples from feral cats in

East Tennessee collected from 2007 to 2012 (2007-2012) (n=36) and convenience samples from

client-owned cats undergoing routine blood work (listed as Pre-pandemic) (n=92) (Fig 2A). As

expected, SARS-CoV-2 experimentally infected cats [14] tested positive with high relative

OD_{450}, and normal cat serum (i.e., negative control) with very low relative OD_{450} (Fig 2A).

Despite pre-dating the pandemic, 52% (67/128) of the cat samples tested positive for antibodies

against SARS-CoV-2 RBD. This is surprising as there was a lack of high cross-reactivity in

guinea pigs immunized with FCoV in Fig 1A. Several reports also showed a lack of similarity

and cross reactivity between alpha coronaviruses and SARS-CoVs [18, 32, 38, 56]. Indeed, two

other groups found that pre-existing immunity to FCoV had no impact on seropositivity of feline

samples [38, 58]. To ensure that the positive ELISA results were specific to the RBD and not to a

co-purified protein, a western blot was carried out using serum from a positive sample (Fig 2B).

Positive cat serum bound a ~32 kDa protein, the size of the RBD protein (Fig 1C). Notably,

normal cat serum did not react with any other protein despite the presence of co-purified

proteins. To further show the specificity of the anti-RBD response, we titrated seropositive and

seronegative samples. Starting with serum from cats experimentally infected with SARS-CoV-2

(Fig 2A) and normal cat serum, saw a normalized OD_{450} >3 standard deviations above the

negative control (i.e., normal cat serum) at all dilutions. This gives a titer >8100 (Fig 2C). 17

seropositive and 10 seronegative pre-2020 cat samples were titrated and assayed in our ELISA

(Fig 2D). Titers ranged from 900 to 8100, with a median titer of 2700 demonstrating both a high

anti-SARS RBD prevalence and titer. Titrations of these earlier seropositive and seronegative

feral cat samples have a similarly high titer (Fig 2E) (median titer 8100) which is on par with the

pre-pandemic samples. AUC for all groups is shown to the right side of their respective titration.

AUC analysis of the titrated samples showed a significant difference between all positive and

negative samples.

Following the surprising presence and prevalence of anti-RBD responses in pre-pandemic

cats, we explored the epidemiological characteristics of our samples. Pre-pandemic convenience

samples were submitted to the University of Tennessee for diagnostic testing of feline

herpesvirus, feline calicivirus, and FCoV. Age, sex, and location of seropositive and

seronegative samples are shown in Table 1. Both seropositive and seronegative samples had a

mean age of >3 years with no difference between the groups and contained similar ratios of

male: female animals (Table 1). Seropositive samples were found in disparate geographic

locations from opposite coasts of the United States (i.e., New York to California (Table 1)). This

observation indicates that seropositivity is not confined to a single geographic region (e.g., East

Tennessee). Based on our limited sampling, we were unable to identify any unique characteristic

or identifier for seropositive vs seronegative samples.

With our discovery of pre-existing antibodies against SARS-CoV-2's RBD, it was

pertinent to examine samples from dogs, another companion animal with high human contact.

Serum samples from dogs (n=36) were collected and retrospectively screened as part of a tick

study during a 7-month period beginning in Jan 2020 and extending into July 2020. These

samples are considered post-pandemic because the timeframe straddles the arrival of SARS-

CoV-2 in East Tennessee (~March 2020). The initial ELISA screen identified 97% seropositivity

in the dog samples (Fig 3A) with only 1 sample falling below the cutoff established on human

serum. Surprisingly, serum from purpose-bred research animals housed at the University of

Tennessee also showed high levels of reactivity (Fig 3A). This raised suspicion about the

specificity of the response. To address this, western blot analysis with canine serum (Fig 3B)

identified a protein other than the RBD (see the ~32 kDa protein in Fig 1C and 2B). The canine

serum recognized a ~60 kDa protein which is likely a co-purified protein present after RBD

purification and is faintly seen in the silver-stained gel in Fig 1D. This co-purified protein was

not detected in the blots performed for Fig. 1C and 2B using anti-6His monoclonal antibody and

cat serum, respectively. Although there is a possibility that canine serum recognizes an oligomer

of RBD [63], based on Fig 1C the anti-6His antibody does not detect any protein >32kDa,

eliminating this possibility.

Following our observation of high levels of anti-SARS-CoV-2 RBD antibodies in North

American cats, we began examining other regional animals. Serum from Tennessee resident, pre-

pandemic cows (n=33) and tigers (n=9), post-pandemic East Tennessee elk (n=12), and post-

pandemic South Carolina white-tailed deer (n=22) were tested for anti-SARS-CoV-2 RBD

antibodies (Fig 4A). Of the four species tested, only the deer from South Carolina showed any

seropositive samples (2/22). Serum titrations show the two seropositive samples have a high titer

>8100 (Fig 4B), and AUC of the titrations show a significant difference between seropositive

and negative deer samples (Fig 4C). Unfortunately, due to limited sample volume, we were

unable to run western blots to demonstrate the specificity for the RBD protein. The deer are post-

pandemic and could represent recent transmission of SARS-CoV-2 into the deer population.

Although these animals probably have had limited contact with humans, white-tailed deer are

susceptible to and capable of transmitting SARS-CoV-2 [64]. Another possibility is that this

species was exposed to the same etiological agent as our pre-pandemic seropositive cats.

To address whether our ELISA-positive animal samples can neutralize SARS-CoV-2

infections, we measured the ability of cat serum to block the interaction of the spike protein with

the human ACE-2 (hACE-2) receptor using a commercially available flow cytometry-based bead

assay. In this assay, neutralization is characterized as the decrease in fluorescence when

antibodies block the fluorescently labeled SARS-CoV-2 S1 subunit from binding to hACE-2

conjugated beads (Fig 5A). Because this assay is not species specific or immunoglobulin type

dependent, it is applicable for assessing both human and feline serum. The internal antibody

control shows a decrease in fluorescence corresponding to levels of neutralizing monoclonal

antibody against SARS-CoV-2. Serum from experimentally infected cats showed potent

neutralization at 1:100 dilution. However, only one ELISA-positive, pre-pandemic cat sample

showed neutralization (Fig 5B). One of the seropositive white-tailed deer samples, and a single

serum sample from mice immunized with PRCoV also showed slight neutralization, clearing the

determined ROC threshold/cutoff value (Fig 5B). Notably, we were unable to detect high levels

of neutralization/neutralizing antibodies even in several of the human convalescent serum

samples (Fig 5B).

Because cross-reactivity of antibodies to SARS-CoV-2 RBD independent of SARS-CoV-

2 infection has not been previously reported in felines, we suspected that the etiological agent

could be another coronavirus [38, 58]. Fecal samples were collected from healthy East

Tennessee cats and screened for coronaviruses using pan-coronavirus primers amplifying

conserved regions of the RNA-dependent RNA polymerase (RdRp), helicase (Hel), and spike (S)

genes [60]. Coronavirus viral RNA, whether common animal coronavirus or SARS-like

coronavirus, is potentially shed in feces [8, 9, 11, 13]. Collection of fecal samples represented a

non-invasive collection method, and SARS-CoV-2 has been reported to have prolonged shedding

within fecal samples of humans [8, 9, 11, 13]. Fifteen out of thirty samples (50%) tested positive

for at least one loci, with most yielding positive results for multiple loci (Table 2). Not

surprisingly, sequences cluster within the alpha-coronavirus group and with high similarity to

previously identified FCoV strains. When all five loci were aligned and concatenated together,

the Maximum-Likelihood phylogenetic tree places the concatenated coronavirus sequences

within the alpha-coronavirus lineage, closely related to FCoV (Fig 6A). We were unable to

amplify or identify any sequences which resemble SARS-like coronaviruses or beta-

coronaviruses. Partial sequencing of the S1 region was able to amplify the RBD from several

coronavirus RNA positive samples. The sequenced RBDs were again highly similar to FCoV

based on Maximum-Likelihood phylogenetic tree (Fig 7A). Along with the phylogenetic tree, a

similarity matrix demonstrates high RBD similarity between previous FCoV strains and those

sequenced here (~80%) (Fig 7B). The RBD from these fecal samples displays low similarity to

betacoronaviruses such as MERS, SARS-CoV, and SARS-CoV-2 (~30%) as previously reported

(Fig 7B).

Discussion

We have employed serological screening as a method to detect potential SARS-CoV-2

exposure in animal populations. Tracking active viral spread in wild and domesticated animals in

real-time *via* sequencing and RT-qPCR is expensive and low throughput. In addition, the

unknown transient nature of viral shedding from different secretions/locations makes this type of

surveillance prohibitively expensive with no guarantee of identifying a virus. Serological

detection of antibodies against SARS-CoV-2 is comparatively high-throughput and inexpensive

while still maintaining sensitivity. A downside of this methodology is the lack of an up-to-date

picture of cross-species transmission, as serology trails initial infections by several days to weeks

[22, 33]. On the other hand, due to the lowered cost of serological testing there is a compensatory

increase in testing capability allowing a broader swath of animals and regions to be sampled with

more frequent re-sampling to track spillover into new species. Our adapted protocol yields

recombinant SARS-CoV-2 RBD protein allowing production of a low-cost indirect anti-RBD

ELISA. The recombinant RBD was sufficient for serological screening *via* ELISA and is

amenable to most labs with prior tissue culture capabilities and does not require large initial

investments in cell lines, culture media, or specialized incubators. We validated our method

demonstrating low cross-reactivity with other common animal coronaviruses (Fig 1A) and >95%

sensitivity and specificity on human serum samples (Fig 1B).

For the seropositive samples identified in our study, mean titers for positive cat samples

were relatively high at ~2700 (Fig 2E, G), which based on reported rapid declines in anti-RBD

responses for SARS-CoV and FCoV points to exposure within the past few years [37, 38, 40].

Further, based on FCoV studies, animals with high titers typically correlate with active viral

shedding and spread within a household, which highlights a potential overlap between

seropositivity and viral shedding [37]. This is in stark contrast to SARS-CoV-2 serosurveys on

pre- and post-pandemic feline samples from Central China. They found no evidence of exposure

before the outbreak, but also positivity levels post-pandemic were significantly lower than shown

here (~12% Central China vs >50% USA) (Fig 2A) [38, 58]. OD_{450} and titers of pre-pandemic

seropositive cat samples, while high, were lower than SARS-CoV-2 experimentally inoculated

cats (6 weeks post infection) (Fig 2A, C, E, G). This likely represents a natural decline in titer
over time for the environmental samples but could also represent lower titers of cross-reactive
antibodies from another coronavirus.

Unfortunately, dog serum was shown to bind to a co-purified protein (Fig 3B), leaving us
unable to utilize our assay for examining cross-species transmission of SARS-like coronaviruses
to canines. We can show that recombinant RBDs produced and purified by groups at both Mt.
Sinai and Emory contain co-purified proteins at approximately the same size as shown in Fig 1A
[30, 33]. As such, screens for SARS-CoV-2 exposure in canines would likely require producing
and purifying the RBD using a different strategy that eliminates non-RBD protein contamination.
Recently both SARS-CoV-2's RBD and soluble full-length spike have been produced and
purified in plants [65]. This alternative method may prove useful for animal SARS-CoV-2
screening by reducing or eliminating false positives due to co-purified proteins.

After the discovery of pre-pandemic seropositive cats, we examined other commercial
(cow) and convenience samples from local wild species (deer, elk, tiger) (Fig 4A). Two out of
twenty-two (9%) white-tailed deer from South Carolina were positive for antibodies against the
RBD (Fig 4B). Unlike our cat samples, the two seropositive deer could represent transmission of
SARS-CoV-2 into the local deer population because these samples were collected post-
pandemic. Interestingly, a recent report showed that ~40% of white-tailed deer from 4 states
(Illinois, Michigan, New York, and Pennsylvania) were positive for SARS-CoV-2 antibodies
[29]. Seropositive animals were only observed from 2019 onward, with pre-pandemic deer
testing negative on their SARS-CoV-2 neutralization assay. This information supports our
finding in South Carolina deer (Fig 4). SARS-CoV-2 sequences were also recently isolated from
the retropharyngeal lymph nodes of wild and captive deer [27, 28]. The dominant genotype of
deer-isolated SARS-CoV-2 genotypes closely corresponded to those circulating within humans
at the time, pointing to potential rapid transmission from humans to animals [28]. This highlights
the importance of the One Health Initiative to provide information on the potential exposure and
spillover into other species and whether there is recombination with native coronaviruses
occurring to generate new variants or establishment of new reservoirs in North America. Further
work is needed to determine the prevalence, spread, and identity of SARS and other
coronaviruses circulating within North American deer and associated species.

That samples from cats experimentally infected with SARS-CoV-2 displayed potent
neutralization (Fig 5B) is unsurprising because of their high ELISA titers (Fig 2C). These
samples were collected at ~8 weeks post infection and likely represent peak titer and
neutralization capacity [14]. Neutralization of SARS-CoV-2 S1 subunit was variable for
environmental feline ELISA-positive samples (Fig 5B). There was no significant difference in
MFI/neutralization between the anti-RBD seropositive and seronegative feline samples (Fig 5B).
Even sera from convalescent, COVID-recovered individuals showed none to minimal
neutralization (Fig 5B). Several groups have found that not all anti-RBD responses generate
neutralizing antibodies [50-53, 66]. Indeed, even in convalescent serum, high levels of RBD
recognition does not guarantee high neutralizing titers, consistent with our own observations (Fig
5B) [50]. Based on ELISA and neutralization results (Fig 2A, Fig 5B), we suspect that these
animals contain antibodies recognizing SARS-CoV-2's RBD, but likely bind to non-neutralizing
epitopes of the RBD domain.

Because ~50% of cats surveyed were seropositive, we reasoned that isolation of the
suspected infectious agent or coronavirus might be possible. Based on fecal viral RNA shedding
following animal coronaviruses infections, PCR amplification with universal coronavirus
primers was used to screen for potential causative agents of anti-RBD seropositivity. This
allowed for non-invasive testing and isolation of coronavirus RNA from infected cats. In line
with previous studies on other wild animals, we did not identify any non-alphacoronaviruses
circulating in felines [67-70]. The coronavirus sequences that were isolated and sequenced likely
represent normal circulating FCoVs (Fig 6). Due to the opportunistic nature of our sampling, we
were unable to obtain any paired blood and fecal samples from the same animal. As such, we are
unable to conclude whether the cats with identified FCoVs would produce cross-reactive
antibodies against SARS-CoV-2's RBD. However, based on our ELISA results in Fig 1A and 2A
showing no correlation with SARS RBD antibodies and FCoV infection, we would suspect not.
Furthermore, not knowing when the seropositive cats were exposed (i.e., cats could have been
infected potentially years prior to any fecal sampling) fecal sampling and sequencing would not
detect a novel coronavirus if it had been cleared. Following the identification of coronavirus
positive fecal samples, we attempted to amplify and sequence the entire RBD region from
positive samples to look for similarity to SARS-CoV-2 or SARS-like viruses. Large portions of

the S1 region spanning the RBD were sequenced and contained RBD regions similar to
previously isolated FCoV strains (Fig. 7), with no similarity to SARS or betacoronaviruses.

The current study demonstrates cross-reactivity of pre-pandemic feline samples with the
RBD of SARS-CoV-2. Our indirect ELISA screen has provided evidence for seropositivity of
serum from North American cats and deer to a SARS-CoV-2 protein previously shown to be
highly specific to SARS coronaviruses [32, 38, 46, 49]. This is the first study to demonstrate
seropositivity of animal samples pre-pandemic. What induces this cross-reactive response was
not readily apparent. However, we propose several possibilities: exposure to another infectious
agent generating cross-reactive antibodies, infection with multiple common coronavirus strains
(Feline coronavirus or otherwise) generating cross-reactive antibodies, or exposure of animals to
a SARS-like coronavirus pre-pandemic.

There is evidence both for and against these explanations for the seropositivity observed.
While we cannot discount a non-coronavirus infection generating cross-reactive antibodies, the
RBD of SARS-like viruses is thought to be unique with no previous evidence of RBD cross-
reactivity [32, 38, 46, 49]. A plausible explanation for seropositivity against SARS-CoV-2's
RBD is infections with coronaviruses generating an atypical response. To-date, cross-reactivity
against the RBD of SARS-CoV-2 has only been reported for SARS and SARS-like
coronaviruses [32, 46]. The common human and animal coronaviruses (both alpha and beta
coronavirus families) individually do not generate cross-reactive antibodies against this protein,
presumably making it a SARS-specific response [30, 32, 33, 38, 46, 49]. For example, prior
FCoV exposure did not impact SARS-CoV-2 RBD recognition [38]. Our own observations
further demonstrated that FCoV (both serotype I and II) and TGEV did not correlate with
serostatus for SARS-CoV-2's RBD (Fig 2A) (FCoV+, FCoV-). Additionally, humans and
animals are exposed to multiple coronaviruses throughout their lifetime, generating measurable
immune responses to antigenic proteins [71, 72]. With humans, CoV infections occur on average
once a year, with protective immunity appearing to wane a few months after initial exposure [71,
72]. Despite reports of cross-reactivity towards portions of SARS-CoV-2's spike protein induced
by other human coronaviruses, there have been no reports of cross-reaction against the distinct
RBD region [32, 46, 49]. While it does not disprove multiple CoV infections generating cross
reactive antibodies, it does cast doubt on this possibility. This leads us to propose prior exposure
to a SARS-like coronavirus for North American cats.

There is limited evidence to support prior transmission of a SARS-like virus within
felines. However, cats are susceptible to infection and transmission of SARS-CoV-2 [14]. SARS
coronaviruses evolved from bat coronaviruses and maintain a high degree of sequence similarity,
even in the RBD region [32]. Conceivably, our definition of SARS-like coronavirus could also
encompass bat coronaviruses due to amino acid similarity and potential cross-reactivity in the
RBD region. At least within Tennessee there are numerous cave systems with several native
species of bats potentially leading to an inter-species transmission. Feline and bat interactions
could lead to direct transfer of a SARS-like agent, or an intermediary species could be involved.
North American deer mice were recently shown to be susceptible to human SARS-CoV-2 and
capable of mouse-to-mouse transmission representing another potential point of introduction into
the feline population [73]. The pre-pandemic exposure of cats to a SARS-like agent does have
detractions. Other groups examining pre-pandemic samples have failed to find evidence of RBD-
reactive serum even within Central China [38, 56, 58]. The ELISA-positive samples from our
environmental samples (i.e., feline and deer) did not neutralize to the same degree as
experimentally inoculated felines (Fig 5B) [14]. However, significant neutralization was not
observed for most samples, even convalescent serum from COVID-positive individuals (Fig 5B).
Despite ~50% seroprevalence in our samples, we were unable to identify any coronavirus
sequences capable of eliciting an anti-SARS RBD response (Fig 6 and 7). Additionally, there is
currently no evidence of a circulating SARS-like or bat betacoronavirus in North America [67-
70]. While we cannot conclusively demonstrate the origin of the anti-RBD responses shown
here, this study indicates that there could be a virus (or another infectious agent) that can
generate cross reactive antibodies to the SARS-CoV-2 RBD.

Limitations of this study

A major limitation of this study is the number of serum samples surveyed for all species.
Due to the small sample size, we have refrained from making extrapolations from our data to
larger animal populations or geographic regions. Similarly, a relatively small number of feline
fecal samples were screened for coronavirus viral RNA (n=30), limiting our ability to detect
coronaviruses. Our choice of sampling source may have also limited our ability to detect novel
coronaviruses or SARS-like coronaviruses. We chose to utilize fecal samples for coronavirus
screening because it is non-invasive, animal coronavirus shedding within feces is common, and
many are transmitted fecal to oral. Additionally, SARS-CoV and SARS-CoV-2 viral RNA has

been detected in stool samples of infected humans [8, 9, 11, 13]. This fecally shed RNA persists longer than nasal or oral sources [13]. However, sampling nasal/oral sources rather than fecal samples may allow for better detection of SARS-CoV-2. Indeed, isolation of SARS-CoV-2 RNA and whole genome sequencing was possible from deer retropharyngeal lymph nodes [28]. Along with sampling site limitations, infection and shedding of viral RNA is a transient process with a relatively small window to detect and sequence the causative agent. While ~20 feline samples were tested for anti-FCoV antibodies, the majority of our samples were of an unknown FCoV status. Additionally, medical histories were not available for these animals and were not screened for prior exposure to other infectious agents. As such, we cannot definitively rule out some other infectious agent conferring the anti-RBD responses.

Conclusion

Our initial goal was to develop an ELISA assay for tracking the reverse zoonosis of SARS-CoV-2. However, when establishing our baseline on pre-pandemic cat samples we discovered seropositive serum for an antigen previously reported to be SARS-specific (i.e., RBD). Seropositivity was ~50% in feline samples and could be found several years prior to the genesis of the current SARS-CoV-2 pandemic. What generated the RBD-reactive antibodies is unknown, but we proposed three possibilities: cross-reaction caused by another infectious agent, multiple coronavirus infections creating a rare cross-reactive antibody, or existence and circulation of a SARS-related virus containing the RBD sequence. Regardless, the high rate of SARS-CoV-2 RBD seropositivity within a common companion animal further highlights our need for a better understanding of the prevalence and crossover potential of wild coronaviruses. Further investigations should address shedding of viral RNA from the seropositive species (i.e., cats and deer) identified here to isolate, sequence, and identify the agent enabling cross-reactivity against the RBD of SARS-CoV-2.

Acknowledgments

We would like to thank the following for their generous contributions making this work possible:

MEDIC Regional Blood Center

Mark Slifka – Oregon Health Sciences University

Jon Wall and Steve Foster – University of Tennessee Medical Center Knoxville

Krammer Lab – Icahn School of Medicine at Mount Sinai

Wrammert Lab – Emory University

Angela Bosco-Lauth – Colorado State University

Jennifer Weisent, Rebekah DeBolt, Sarah Englebert, Jessica Thompson – University of

Tennessee College of Veterinary Medicine Shelter Medicine Service

References

1.   Poon, L.L.M. and M. Peiris, *Emergence of a novel human coronavirus threatening*
613   *human health.* Nat Med, 2020. **26**(3): p. 317-319.
614   2.   Guan, W.J., et al., *Clinical Characteristics of Coronavirus Disease 2019 in China.* N
615   Engl J Med, 2020. **382**(18): p. 1708-1720.
616   3.   Latinne, A., et al., *Origin and cross-species transmission of bat coronaviruses in China.*
617   Nat Commun, 2020. **11**(1): p. 4235.
618   4.   Li, X., et al., *Emergence of SARS-CoV-2 through recombination and strong purifying*
619   *selection.* Sci Adv, 2020. **6**(27).
620   5.   Andersen, K.G., et al., *The proximal origin of SARS-CoV-2.* Nat Med, 2020. **26**(4): p.
621   450-452.
622   6.   Sun, J., et al., *Prolonged Persistence of SARS-CoV-2 RNA in Body Fluids.* Emerg Infect
623   Dis, 2020. **26**(8): p. 1834-1838.
624   7.   Meyerowitz, E.A., et al., *Transmission of SARS-CoV-2: A Review of Viral, Host, and*
625   *Environmental Factors.* Ann Intern Med, 2021. **174**(1): p. 69-79.
626   8.   Wang, W., et al., *Detection of SARS-CoV-2 in Different Types of Clinical Specimens.*
627   JAMA, 2020. **323**(18): p. 1843-1844.
628   9.   Kim, J.M., et al., *Detection and Isolation of SARS-CoV-2 in Serum, Urine, and Stool*
629   *Specimens of COVID-19 Patients from the Republic of Korea.* Osong Public Health Res
630   Perspect, 2020. **11**(3): p. 112-117.
631   10.   Kang, M., et al., *Probable Evidence of Fecal Aerosol Transmission of SARS-CoV-2 in a*
632   *High-Rise Building.* Ann Intern Med, 2020. **173**(12): p. 974-980.
633   11.   Xiao, F., et al., *Infectious SARS-CoV-2 in Feces of Patient with Severe COVID-19.*
634   Emerg Infect Dis, 2020. **26**(8): p. 1920-1922.
635   12.   Yu, I.T., et al., *Evidence of airborne transmission of the severe acute respiratory*
636   *syndrome virus.* N Engl J Med, 2004. **350**(17): p. 1731-9.
637   13.   Roltgen, K., et al., *Defining the features and duration of antibody responses to SARS-*
638   *CoV-2 infection associated with disease severity and outcome.* Sci Immunol, 2020. **5**(54).
639   14.   Bosco-Lauth, A.M., et al., *Experimental infection of domestic dogs and cats with SARS-*
640   *CoV-2: Pathogenesis, transmission, and response to reexposure in cats.* Proc Natl Acad
641   Sci U S A, 2020. **117**(42): p. 26382-26388.
642   15.   Shi, J., et al., *Susceptibility of ferrets, cats, dogs, and other domesticated animals to*
643   *SARS-coronavirus 2.* Science, 2020. **368**(6494): p. 1016-1020.
644   16.   Wu, L., et al., *Broad host range of SARS-CoV-2 and the molecular basis for SARS-CoV-2*
645   *binding to cat ACE2.* Cell Discov, 2020. **6**: p. 68.
646   17.   Segales, J., et al., *Detection of SARS-CoV-2 in a cat owned by a COVID-19-affected*
647   *patient in Spain.* Proc Natl Acad Sci U S A, 2020. **117**(40): p. 24790-24793.

18.   Fritz, M., et al., *High prevalence of SARS-CoV-2 antibodies in pets from COVID-19+*
households. One Health, 2021. **11**: p. 100192.
19.   Sit, T.H.C., et al., *Infection of dogs with SARS-CoV-2.* Nature, 2020. **586**(7831): p. 776-
778.
20.   Patterson, E.I., et al., *Evidence of exposure to SARS-CoV-2 in cats and dogs from*
households in Italy. Nat Commun, 2020. **11**(1): p. 6231.
21.   Barrs, V.R., et al., *SARS-CoV-2 in Quarantined Domestic Cats from COVID-19*
Households or Close Contacts, Hong Kong, China. Emerg Infect Dis, 2020. **26**(12): p.
3071-3074.
22.   Halfmann, P.J., et al., *Transmission of SARS-CoV-2 in Domestic Cats.* N Engl J Med,
2020. **383**(6): p. 592-594.
23.   Oude Munnink, B.B., et al., *Transmission of SARS-CoV-2 on mink farms between*
humans and mink and back to humans. Science, 2021. **371**(6525): p. 172-177.
24.   Martina, B.E., et al., *Virology: SARS virus infection of cats and ferrets.* Nature, 2003.
425(6961): p. 915.
25.   Ng, S.K., *Possible role of an animal vector in the SARS outbreak at Amoy Gardens.*
Lancet, 2003. **362**(9383): p. 570-2.
26.   Lun, Z.R. and L.H. Qu, *Animal-to-human SARS-associated coronavirus transmission?*
Emerg Infect Dis, 2004. **10**(5): p. 959.
27.   Hale, V.L., et al., *SARS-CoV-2 infection in free-ranging white-tailed deer.* Nature, 2021.
28.   Kuchipudi, S.V., et al., *Multiple spillovers from humans and onward transmission of*
SARS-CoV-2 in white-tailed deer. Proc Natl Acad Sci U S A, 2022. **119**(6).
29.   Chandler, J.C., et al., *SARS-CoV-2 exposure in wild white-tailed deer (Odocoileus*
virginianus). Proc Natl Acad Sci U S A, 2021. **118**(47).
30.   Amanat, F., et al., *A serological assay to detect SARS-CoV-2 seroconversion in humans.*
Nat Med, 2020. **26**(7): p. 1033-1036.
31.   Long, Q.X., et al., *Antibody responses to SARS-CoV-2 in patients with COVID-19.* Nat
Med, 2020. **26**(6): p. 845-848.
32.   Premkumar, L., et al., *The receptor binding domain of the viral spike protein is an*
immunodominant and highly specific target of antibodies in SARS-CoV-2 patients. Sci
Immunol, 2020. **5**(48).
33.   Suthar, M.S., et al., *Rapid Generation of Neutralizing Antibody Responses in COVID-19*
Patients. Cell Rep Med, 2020. **1**(3): p. 100040.
34.   Desmarets, L.M., et al., *Experimental feline enteric coronavirus infection reveals an*
aberrant infection pattern and shedding of mutants with impaired infectivity in enterocyte
cultures. Sci Rep, 2016. **6**: p. 20022.
35.   Vogel, L., et al., *Pathogenic characteristics of persistent feline enteric coronavirus*
infection in cats. Vet Res, 2010. **41**(5): p. 71.
36.   Cevik, M., et al., *SARS-CoV-2, SARS-CoV, and MERS-CoV viral load dynamics,*
duration of viral shedding, and infectiousness: a systematic review and meta-analysis.
The Lancet Microbe.
37.   Addie, D.D. and O. Jarrett, *Use of a reverse-transcriptase polymerase chain reaction for*
monitoring the shedding of feline coronavirus by healthy cats. Vet Rec, 2001. **148**(21): p.
649-53.
38.   Zhang, Q., et al., *A serological survey of SARS-CoV-2 in cat in Wuhan.* Emerg Microbes
Infect, 2020. **9**(1): p. 2013-2019.

39.   Liu, W., et al., *Two-year prospective study of the humoral immune response of patients*
695   *with severe acute respiratory syndrome.* J Infect Dis, 2006. **193**(6): p. 792-5.
696   40.   Cao, W.C., et al., *Disappearance of antibodies to SARS-associated coronavirus after*
697   *recovery.* N Engl J Med, 2007. **357**(11): p. 1162-3.
698   41.   Lin, Q., et al., *Duration of serum neutralizing antibodies for SARS-CoV-2: Lessons from*
699   *SARS-CoV infection.* J Microbiol Immunol Infect, 2020. **53**(5): p. 821-822.
700   42.   Tang, F., et al., *Lack of peripheral memory B cell responses in recovered patients with*
701   *severe acute respiratory syndrome: a six-year follow-up study.* J Immunol, 2011.
702   **186**(12): p. 7264-8.
703   43.   Zeng, W., et al., *Biochemical characterization of SARS-CoV-2 nucleocapsid protein.*
704   Biochem Biophys Res Commun, 2020. **527**(3): p. 618-623.
705   44.   Hoffmann, M., et al., *SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is*
706   *Blocked by a Clinically Proven Protease Inhibitor.* Cell, 2020. **181**(2): p. 271-280 e8.
707   45.   Lan, J., et al., *Structure of the SARS-CoV-2 spike receptor-binding domain bound to the*
708   *ACE2 receptor.* Nature, 2020. **581**(7807): p. 215-220.
709   46.   Ng, K.W., et al., *Preexisting and de novo humoral immunity to SARS-CoV-2 in humans.*
710   Science, 2020. **370**(6522): p. 1339-1343.
711   47.   Chi, X., et al., *A neutralizing human antibody binds to the N-terminal domain of the Spike*
712   *protein of SARS-CoV-2.* Science, 2020. **369**(6504): p. 650-655.
713   48.   Song, G., et al., *Cross-reactive serum and memory B-cell responses to spike protein in*
714   *SARS-CoV-2 and endemic coronavirus infection.* Nat Commun, 2021. **12**(1): p. 2938.
715   49.   Anderson, E.M., et al., *Seasonal human coronavirus antibodies are boosted upon SARS-*
716   *CoV-2 infection but not associated with protection.* Cell, 2021. **184**(7): p. 1858-1864 e10.
717   50.   Lee, W.T., et al., *Neutralizing Antibody Responses in COVID-19 Convalescent Sera.* J
718   Infect Dis, 2021. **223**(1): p. 47-55.
719   51.   Rogers, T.F., et al., *Isolation of potent SARS-CoV-2 neutralizing antibodies and*
720   *protection from disease in a small animal model.* Science, 2020. **369**(6506): p. 956-963.
721   52.   Niu, L., et al., *A Structural Landscape of Neutralizing Antibodies Against SARS-CoV-2*
722   *Receptor Binding Domain.* Front Immunol, 2021. **12**: p. 647934.
723   53.   Zhou, Y., et al., *Enhancement versus neutralization by SARS-CoV-2 antibodies from a*
724   *convalescent donor associates with distinct epitopes on the RBD.* Cell Rep, 2021. **34**(5):
725   p. 108699.
726   54.   Zhu, Y., et al., *Cross-reactive neutralization of SARS-CoV-2 by serum antibodies from*
727   *recovered SARS patients and immunized animals.* Sci Adv, 2020. **6**(45).
728   55.   Lv, H., et al., *Cross-reactive Antibody Response between SARS-CoV-2 and SARS-CoV*
729   *Infections.* Cell Rep, 2020. **31**(9): p. 107725.
730   56.   Zhao, S., et al., *Serological Screening for Coronavirus Infections in Cats.* Viruses, 2019.
731   **11**(8).
732   57.   Deng, J., et al., *Serological survey of SARS-CoV-2 for experimental, domestic,*
733   *companion and wild animals excludes intermediate hosts of 35 different species of*
734   *animals.* Transbound Emerg Dis, 2020. **67**(4): p. 1745-1749.
735   58.   Deng, J., et al., *SARS-CoV-2 Serological Survey of Cats in China before and after the*
736   *Pandemic.* Virol Sin, 2020.
737   59.   Stadlbauer, D., et al., *SARS-CoV-2 Seroconversion in Humans: A Detailed Protocol for a*
738   *Serological Assay, Antigen Production, and Test Setup.* Curr Protoc Microbiol, 2020.
739   **57**(1): p. e100.

60. Poon, L.L., et al., *Identification of a novel coronavirus in bats.* J Virol, 2005. **79**(4): p.
2001-9.
61. Thompson, J.D., D.G. Higgins, and T.J. Gibson, *CLUSTAL W: improving the sensitivity*
of progressive multiple sequence alignment through sequence weighting, position-
specific gap penalties and weight matrix choice. Nucleic Acids Res, 1994. **22**(22): p.
4673-80.
62. Kumar, S., et al., *MEGA X: Molecular Evolutionary Genetics Analysis across Computing*
Platforms. Mol Biol Evol, 2018. **35**(6): p. 1547-1549.
63. Johari, Y.B., et al., *Production of trimeric SARS-CoV-2 spike protein by CHO cells for*
serological COVID-19 testing. Biotechnol Bioeng, 2021. **118**(2): p. 1013-1021.
64. Palmer, M.V., et al., *Susceptibility of white-tailed deer (Odocoileus virginianus) to SARS-*
CoV-2. J Virol, 2021.
65. Rattanapisit, K., et al., *Rapid production of SARS-CoV-2 receptor binding domain (RBD)*
and spike specific monoclonal antibody CR3022 in Nicotiana benthamiana. Sci Rep,
2020. **10**(1): p. 17698.
66. Xiaojie, S., et al., *Neutralizing antibodies targeting SARS-CoV-2 spike protein.* Stem Cell
Res, 2020. **50**: p. 102125.
67. Olival, K.J., et al., *Possibility for reverse zoonotic transmission of SARS-CoV-2 to free-*
ranging wildlife: A case study of bats. PLoS Pathog, 2020. **16**(9): p. e1008758.
68. Dominguez, S.R., et al., *Detection of group 1 coronaviruses in bats in North America.*
Emerg Infect Dis, 2007. **13**(9): p. 1295-300.
69. Misra, V., et al., *Detection of polyoma and corona viruses in bats of Canada.* J Gen
Virol, 2009. **90**(Pt 8): p. 2015-2022.
70. Osborne, C., et al., *Alphacoronaviruses in New World bats: prevalence, persistence,*
phylogeny, and potential for interaction with humans. PLoS One, 2011. **6**(5): p. e19156.
71. Macnaughton, M.R., *Occurrence and frequency of coronavirus infections in humans as*
determined by enzyme-linked immunosorbent assay. Infect Immun, 1982. **38**(2): p. 419-
23.
72. Schmidt, O.W., et al., *Rises in titers of antibody to human coronaviruses OC43 and 229E*
in Seattle families during 1975-1979. Am J Epidemiol, 1986. **123**(5): p. 862-8.
73. Griffin, B.D., et al., *SARS-CoV-2 infection and transmission in the North American deer*
mouse. Nat Commun, 2021. **12**(1): p. 3612.

Figure 1: Anti-SARS-CoV-2 ELISA Sensitivity and Specificity. (A) Cross reactivity of anti-

CoV antibodies against SARS-CoV-2 RBD. Polyclonal sera from guinea pigs immunized with

common animal coronaviruses (turkey coronavirus, TCoV; porcine respiratory coronavirus,

PRCoV; canine coronavirus, CCoV; feline coronavirus, FCoV; bovine coronavirus, BCoV) was

used in a SARS-CoV-2 RBD indirect ELISA. Positive samples consisted of polyclonal serum

from a SARS-CoV-2 infected patient and a monoclonal antibody to SARS-CoV (CR3022). The

negative control group was comprised of pre-pandemic human serum. Secondary antibodies

were either anti-human IgG (1:10,000) (Rockland Immunochemicals, USA) or anti-guinea pig

IgG (1:10,000) (Life Technologies Corp, USA). Bars represent mean and standard deviation

(n>3 for all samples). (B) ELISA validation using 66 human Covid-positive plasma and 22

negative serum samples. Human antibodies against the SARS-CoV-2 RBD were detected with

an indirect RBD-specific ELISA. Secondary antibody was the anti-human IgG (1:10,000)

(Rockland Immunochemicals, USA). ROC analysis determined the positive OD_{450} cutoff value

(dashed line). Positive plasma samples were donated COVID recovered patients and pre-

pandemic serum samples were the negative controls. Based on the experimentally determined

cutoff value, 64 of the 66 positive samples were anti-RBD positive, giving a sensitivity value of

96.96%. All but one of the negative samples were below the cutoff value for a specificity of

95.45%. Adjacent tables list first and third quartiles along with mean and median OD_{450} values

of COVID-positive and negative human samples. Bars represent mean and standard deviation

(n>3). (C) Anti-6His western blot on HEK-293T17 purified RBD from 5% serum conditions.

Samples were run on a 12% denaturing SDS-PAGE gel. Protein was transferred to nitrocellulose

and probed with anti-6His antibody at 1:10,000 (Proteintech, USA). White light and

chemiluminescent images were overlaid and from left to right, ladder (lane 1) and purified RBD

(lane 2). (D) Silver stain of purified recombinant SARS-Cov-2 RBD produced in HEK-293T17.

From left to right: Ladder, HEK-RBD under 5% serum conditions, HEK-RBD from 2% serum

conditions. Samples were denatured and run on a 12% SDS-PAGE gel and Silver Stained

(Thermo Fisher Scientific, USA). For B and C, representative data shown.

Figure 2: Pre-Pandemic Feline Antibodies Cross-React with SARS-CoV-2 RBD. (A) ELISA

results of cat serum RBD reactivity. 93 pre-pandemic feline serum samples were tested for

reactivity in our anti-RBD ELISA with anti-felid IgG secondary (1:10,000) (Invitrogen, USA).

Cutoff values were determined by receiver operator curve (ROC) analysis. OD_{450} for samples in

each group were plotted with the dotted line representing the positive threshold. Two sets of pre-

pandemic cat samples were collected. Pre-pandemic cat convenience samples (n=73) were

collected in local clinics and sent to the University of Tennessee for diagnostic testing or during

feral cat studies (2007-2012) (n=36). Pre-pandemic convenience samples were subdivided into

feline coronavirus positive (FCoV+) and negative (FCoV-) subgroups. Normal cat serum

(Jackson ImmunoResearch Laboratories, USA) serves as the negative control and SARS-CoV-

2+ serum from two cats experimentally inoculated with SARS-CoV-2 as positive controls. Side

table lists first and third quartiles and mean and median OD_{450} values for all samples. Bars
represent mean +/- standard deviation (n>3 for all samples). (B) Western blot of purified RBD
using serum from a single positive cat sample. Purified RBD was run under denaturing
conditions and blotted onto nitrocellulose. The RBD blot was first probed with cat serum from an
ELISA positive sample (1:20 dilution) followed by anti-felid IgG-HRP conjugated (1:10,000
dilution) (Invitrogen, USA). White light and chemiluminescent images were overlaid. Lane 1 is
the molecular weight ladder and lane 2 is purified RBD. (C, D, E) Titration of seropositive and
seronegative serums assessed *via* RBD ELISA. OD_{450} values were plotted against the reciprocal
dilution. Samples were considered positive if they were 3 standard deviations above the negative
average for each dilution. Anti-RBD titer was designated as the last dilution above the negative
cutoff. Positive controls were human COVID-positive serum and negative controls were normal
human and cat serum (Jackson ImmunoResearch Laboratories, USA). Statistics for the positive
sample titrations are included in the table along with AUC analysis. (C) Serum from two SARS-
CoV-2 infected cats (●) and normal cat serum (■) were titrated in an anti-RBD ELISA. (D)
Titration of 17 seropositive and 10 seronegative, pre-pandemic cat samples. (E) Titration of four
seropositive and seronegative cat samples collected from 2007-12. For A and B, representative
data shown. For A, Tukey's one-way ANOVA with multiple comparisons was performed. For C,
D, E AUC analysis and Student's one-tailed t-test with Welch's correction was performed. $p<$
$0.05 = *$, $p< 0.01 = **$, $p< 0.001 = ***$.

Figure 3: Dog Serum Cross-reacts to a Co-Purified Protein. (A) Anti-SARS-CoV-2 RBD
ELISA with dog serum. Serum from thirty-six client-owned and two purpose-bred research dogs
were tested in an anti-RBD ELISA with anti-canine IgG secondary HRP (1:10,000) (Bethyl
Laboratories, USA). Table to the right lists the first and third quartiles, median, and mean OD_{450}
values for all samples. Bars represent mean +/- standard deviation (n>3). (B) Western blot of
purified RBD using serum from a positive dog sample. Purified RBD was probed with dog
serum from an ELISA positive sample (1:20 dilution) followed by anti-canine IgG-HRP
(1:10,000 dilution) (Bethyl Laboratories, USA). White light and chemiluminescent images were
overlaid. Lane 1 (from left to right) ladder and lane 2: purified RBD. For all figures,
representative data shown.

Figure 4: Serological Testing of Other Regional Animals. (A). Anti-SARS-CoV-2 RBD

ELISA with bovine, elk, tiger, and deer serum. Thirty-three pre-pandemic East Tennessee cows,

twelve post-pandemic East Tennessee elk, nine pre-pandemic East Tennessee tigers, and twenty-

two post-pandemic South Carolina deer serum samples were tested for anti-RBD antibodies.

Species-specific secondary antibodies were used at the following dilutions: anti-bovine 1:250

(Bethyl Laboratories, USA), anti-elk/deer 1:250 (KPL, USA), anti-tiger/cat 1:10,000 (Invitrogen,

USA), and anti-deer 1:250 (KPL, USA). Bars represent mean +/- standard deviation (n>3 for all

samples). (B) Titration of two seropositive (●) and four seronegative (■) deer samples. OD_{450}

values are plotted against the reciprocal dilution of each sample. Samples were considered

positive if they were 3 standard deviations above the negative average for each dilution. Positive

and negative controls were human COVID-positive and negative samples, respectively. Statistics

for the positive sample titrations are included in the table. The AUC analysis for titrations of deer

ELISA positive and negative samples is shown to the right. For all figures, representative data

shown. For AUC analysis Student's one-tailed t-test with Welch's correction was performed. $p<$

$0.05 = *$, $p< 0.01 = **$, $p< 0.001 = ***$.

Figure 5: Neutralization Assays. (A) Schematic of the neutralization assay. Neutralization is

measured as the decrease in binding of phycoerythrin (PE)-labeled SARS-CoV-2 S1 subunit to

human ACE-2 conjugated beads. Addition of neutralizing antibodies results in a decreased mean

fluorescent intensity (MFI) as measured by flow cytometry. (B) Neutralization of SARS-CoV-2

S1 subunit interaction with hACE2. Serum from several ELISA positive and negative cats

(ELISA+ and ELISA-, respectively), serum from South Carolina white-tailed deer (SCWD, 2

ELISA-positive and 1 negative), mice immunized with other common coronaviruses

(BCoV=bovine coronavirus, TCoV= Turkey coronavirus, PRCoV=porcine respiratory

coronavirus, FCoV=feline coronavirus, CCoV=canine coronavirus), and human serum samples

(Covid+ = convalescent plasma from Covid+ humans, Covid- = pre-SARS-CoV-2 serum

samples, Vaccine = serum post vaccination against SARS-CoV-2 Spike protein). SARS-CoV-2

infected cats and normal cat serum served as positive and negative controls, respectively. Data

was normalized to normal cat serum representing 100% binding of SARS-CoV-2 S1 subunit to

hACE2 beads. ROC analysis was used to generate a positive reduction threshold (dotted line).

Each point is an average of 2 replicates. A Tukey's one-way ANOVA with multiple comparisons
was used to analyze experimental groups. $p < 0.05 = *$, $p < 0.0001 = ****$.

Figure 6: Pan Coronavirus Screen of East Tennessee Felines. Fecal samples from healthy cats
were collected and screened for conserved coronavirus sequences. Phylogenetic tree were
generated consisting of common human and animal coronaviruses: CRCoV (canine respiratory
coronavirus), BCoV (bovine coronavirus), OC43 (human beta-coronavirus), MHV (murine
hepatitis virus), MERS-CoV (Middle East respiratory coronavirus), SARS-CoV-2 (severe acute
respiratory coronavirus-2), SARS-CoV (severe acute respiratory coronavirus), AvianCoV (duck
coronavirus), NL63 (human alpha-coronavirus), 229E (human alpha-coronavirus), TGEV
(transmissible gastroenteritis virus), PRCoV (porcine respiratory coronavirus), FCoV (feline
coronavirus strains UU19, Felix, RM, Black, UU8), CCov (canine coronavirus), HKU15
(porcine delta-coronavirus), as well as a locally identified coronavirus (CP-20-26). Sequences
from five coronavirus loci were independently aligned, trimmed, and concatenated together.
Concatenated sequences were aligned, and phylogenetic trees generated with the Maximum-
Likelihood method with bootstrap analysis in MEGA X. Bootstrap values for each branch are
shown with lengths to scale. Coronavirus lineages are annotated on the tree.

Figure 7: RBD Coronavirus Screen of East Tennessee Felines. (A) Fecal samples from
healthy cats were collected and screened for S1/RBD coronavirus sequences. Phylogenetic tree
consisting of common human and animal coronaviruses: CRCoV (canine respiratory
coronavirus), BCoV (bovine coronavirus), OC43 (human beta-coronavirus), MHV (murine
hepatitis virus), MERS-CoV (Middle East respiratory coronavirus), SARS-CoV-2 (severe acute
respiratory coronavirus-2), SARS-CoV (severe acute respiratory coronavirus), AvianCoV (duck
coronavirus), NL63 (human alpha-coronavirus), 229E (human alpha-coronavirus), TGEV
(transmissible gastroenteritis virus), PRCoV (porcine respiratory coronavirus), FCoV (feline
coronavirus strains UU19, Felix, RM, Black, UU8), CCoV (canine coronavirus), HKU15
(porcine delta-coronavirus), as well as locally identified coronaviruses (CP-20-13, CP-20-19,
CP-20-23). Sequences from the S1 region were aligned and trimmed. Maximum-Likelihood
phylogenetic trees were generated with bootstrap analysis in MEGA X. Bootstrap values for each
branch are shown with lengths to scale. (B) Percent similarity matrix of select coronaviruses.

Aligned and truncated RBD regions from the shown coronaviruses were analyzed *via* Clustal

Omega to determine percent identity matrix.

Table 1: Characteristics of Feline Samples.

	Seropositive	%	Seronegative	%
Age (n =)	58		15	
Average Age (years +/- S.D.)	3.56 +/- 3.67		3.87 +/- 4.37	
< 1 year	15	25.86	5	33.33
1-3 years	22	37.93	4	26.67
4-6 years	10	17.24	2	13.33
> 6 years	11	18.97	4	26.67
Sex (n =)	56		19	
Male	30	53.57	12	63.16
Female	26	46.43	7	36.84
Location (n =)	62		21	
TN	10	16.12	7	33.33
NY	19	30.64	9	42.85
CA	27	43.54	5	23.8
MA	4	6.45	0	0
SC	1	1.61	0	0
WI	1	1.61	0	0

Cats were grouped based on the seronegative/positive status from the ELISA from Fig 2A. Student's t-test was used to determine significance (ns = not significant).

A.

OD$_{450}$	α-TCoV	α-PRCoV	α-CCoV	α-FCoV	α-BCoV	α-SARS-CoV	α-SARS-CoV-2	Pre-Pandemic
25%	0.038	0.227	0.161	0.109	0.084	0.935	0.996	0.078
Median	0.048	0.253	0.193	0.159	0.102	1.008	1.000	0.088
Mean	0.058	0.258	0.199	0.173	0.127	1.051	1.000	0.095
75%	0.054	0.280	0.249	0.246	0.180	1.209	1.004	0.115

B.

Sensitivity = 96.96% (64/66)

Specificity = 95.45% (21/22)

OD$_{450}$	Positive	Negative	SARS-CoV
25%	0.427	0.050	0.821
Median	0.679	0.054	0.981
Mean	0.669	0.068	0.985
75%	0.862	0.077	1.151

A.

Normalized OD$_{450}$

All = 97.22% (35/36)

OD$_{450}$	Client	SPF-free
25%	0.397	0.981
Median	0.813	1.104
Mean	0.731	1.065
75%	1.062	1.126

B.

75+ kDa
50 kDa
37 kDa
25 kDa

A.

OD$_{450}$	Cow	Elk	Tiger	White Tail Deer
25%	0.041	0.039	0.065	0.041
Median	0.044	0.041	0.108	0.049
Mean	0.047	0.043	0.103	0.119
75%	0.052	0.048	0.142	0.058

B.

● Positive Control
■ Negative Control

	Titer
Mean	1:8100
Median	1:8100
Minimum	1:8100
Maximum	1:8100

Table 2: Fecal Coronavirus Screening Information.

Sample	RdRp1	RdRp2	Spike1	Spike2	Spike3	Helicase	Location	Accession #'s
CP-20-01		■		■		■	Jefferson County, TN, USA	MZ220722-MZ220724
CP-20-02		■		■		■	Washington County, TN, USA	MZ220725-MZ220727
CP-20-03							TN, USA	N/A
CP-20-04	■	■				■	Monroe County, TN, USA	MZ220728-MZ220730
CP-20-05	■						TN, USA	N/A
CP-20-06							Sullivan County, TN, USA	N/A
CP-20-07	■	■		■		■	Grainger County, TN, USA	MZ220731-MZ220734
CP-20-08	■						Monroe County, TN, USA	MZ220735
CP-20-09		■		■		■	Sullivan County, TN, USA	MZ220736-MZ220738
CP-20-10							Washington County, TN, USA	N/A
CP-20-11							Jefferson County, TN, USA	N/A
CP-20-12	■	■	■	■			Monroe County, TN, USA	MZ220739-MZ220742
CP-20-13	■	■		■			Grainger County, TN, USA	MZ220743-MZ220745
CP-20-14							Jefferson County, TN, USA	N/A
CP-20-15							Sullivan County, TN, USA	N/A
CP-20-16							Sevier County, TN, USA	N/A
CP-20-17							Cocke County, TN, USA	N/A
CP-20-18							Hamblin County, TN, USA	N/A
CP-20-19			■	■			Washington/Sullivan County, TN, USA	MZ220746-MZ220747
CP-20-20				■			Hamblin County, TN, USA	MZ220748
CP-20-21							Washington/Sullivan County, TN, USA	N/A
CP-20-22							Grainger County, TN, USA	N/A
CP-20-23		■		■			Jefferson County, TN, USA	MZ220749-MZ220750
CP-20-24							Jefferson County, TN, USA	N/A
CP-20-25							Grainger County, TN, USA	N/A
CP-20-26	■	■	■	■		■	Grainger County, TN, USA	MZ220751-MZ220755
CP-20-27	■	■		■		■	Grainger County, TN, USA	MZ220756-MZ220759
CP-20-28	■						Washington/Sullivan County, TN, USA	MZ220760
CP-20-29							TN, USA	N/A
CP-20-30	■					■	Grainger County, TN, USA	MZ220761-MZ220762

Sample IDs were assigned to de-identified fecal samples from healthy cats. RNA was extracted from each and PCR screened for amplification of 6 coronavirus loci across 3 conserved genes (RdRp, spike, and helicase). Positive amplification and sequencing is shown by a black shaded box. 15/30 samples were positive for at least one coronavirus loci. Animals originated from multiple East Tennessee counties, and, when known, exact county of origin is listed. Accession numbers for each sample are shown to the right and consist of each sequenced loci.

Percent Similarity	MERS-CoV	SARS-CoV	SARS-CoV-2	CP-20-23	CP-20-19	CP-20-13	FCoV (Black)	CCoV	FCoV	PRCoV	TGEV	229E	NL63
MERS-CoV	100												
SARS-CoV	32.2	100											
SARS-CoV-2	38.05	69.86	100										
CP-20-23	32.29	26.27	29.9	100									
CP-20-19	33.76	26.75	30.14	87.07	100								
CP-20-13	33.76	26.75	30.14	87.07	100	100							
FCoV (Black)	32.29	25.54	28.23	86.56	87.76	87.76	100						
CCoV	29.91	33.41	29.81	45.27	45.96	45.96	44.06	100					
FCoV	29.17	33.41	29.81	46.3	45.96	45.96	44.75	96.94	100				
PRCoV	30.34	34.14	30.53	45.61	46.64	46.64	44.92	88.1	88.78	100			
TGEV	31.01	34.14	31.49	45.96	46.81	46.81	45.09	91.67	91.67	94.9	100		
229E	32.02	31.52	28.47	45.17	44.05	44.05	42.19	45.54	44.98	45.54	44.98	100	
NL63	33.27	31.02	31.03	52.41	51.48	51.48	50	49.91	49.17	48.8	48.61	57.55	100