Parental provisioning drives brain size in birds

Griesser, Michael 1-4

Drobniak, Szymon M. 5-6

Graber, Sereina M. 4

van Schaik, Carel P. 4,7

1) Department of Biology, University of Konstanz, Konstanz, Germany

2) Center for the Advanced Study of Collective Behavior, University of Konstanz, Konstanz, Germany.

3) Department of Collective Behavior, Max Planck Institute of Animal Behavior, Konstanz, Germany

4) Department of Anthropology, University of Zurich, Zurich, Switzerland

5) Evolution & Ecology Research Centre, School of Biological, Environmental & Earth Sciences, University of New South Wales, Sydney, Australia

6) Institute of Environmental Sciences; Jagiellonian University, Krakow, Poland

7) Centre for Interdisciplinary Study of Language Evolution, University of Zurich, Zürich, Switzerland

Contact information

Michael Griesser: michael.griesser@uni-konstanz.de

24   **Abstract**

Larger brains should be adaptive because they support numerous eco- and socio-cognitive benefits, but these benefits explain only a modest part of the interspecific variation in brain

27   size. Notably underexplored are the high energetic costs of developing brains, and thus the possible role of parental provisioning in the evolution of adult brain size. We explore this idea in birds, which show considerable variation in both socio-ecological traits and the

30   energy transfer from parents to offspring. Comparative analyses of 1,176 bird species show that the combination of adult body mass, mode of development at hatching, relative egg mass, and the time spent provisioning the young in combination strongly predict relative

33   brain size across species. Adding adult eco- and socio-cognitive predictors only marginally adds explanatory value. We therefore conclude that parental provisioning enabled bird species to evolve into skill-intensive niches, reducing interspecific competition and

36   consequently promoting survival prospects and population stability. Critically, parental provisioning also explains why precocial bird species have smaller brains than altricial ones. Finally, these results suggest that the cognitive adaptations that provide the behavioral

39   flexibility to improve reproductive success and survival are intrinsically linked to successful parental provisioning. Our findings also suggest that the traditionally assessed cognitive abilities may not predict relative brain size.

Significance statement

Species vary substantially in their brain size, and the idea that larger-brained species are

45   those with a complex social life or enhanced technological skills has generated much interest. Here we explore another explanation. Because the cognitive benefits provided by brains materialize after a long delay, and constructing them independently involves a major

48   bootstrapping problem for the young, parental inputs may be critical for a species' brain size. Comparative analyses of a large sample of bird species show that parental provisioning strongly predicts relative brain size across species, swamping the effects of previously

51   considered social or technological cognitive abilities. Thus, cognitive adaptations in birds that enable successful parental provisioning also provide the behavioral flexibility to improve survival and reproductive success.

Introduction

Brain size varies extensively within and across lineages, and has increased considerably over evolutionary time (1). This intriguing variability has raised questions about the adaptive benefits of increased relative brain size (i.e., relative to adult body mass). An obvious benefit is improved cognitive abilities (2-4), and numerous comparative studies explored this idea. In mammals, work has prominently focused on the social brain hypothesis, which emphasizes the social benefits of relatively larger brains, but eco-cognitive benefits have also been explored (5-7). In birds, work focused on both social drivers, especially pair bond maintenance (5, 7, 8), and ecological ones, especially environmental variability (9-12). Whereas comparative studies have difficulty assessing whether larger brains improve reproductive success, they can more easily show whether they improve survival. Indeed, comparative studies support a positive association between relative brain size and longevity in diverse lineages (13-15).

Other research independently focused on the high energy costs of brain tissue (16, 17). This aspect is pivotal for the expensive tissue hypothesis, which focuses on the trade-off between brain size and other energetically expensive tissues, such as guts (18). Subsequent work highlighted a general link between relative brain size and the energy available to sustaining the brain (19). These costs imply that natural selection cannot always respond to opportunities to evolve specific cognitive adaptations via an increase in brain size, even if this would in principle lead to higher fitness. Indeed, recent comparative work has found that species living in fluctuating environments that cause inevitable reductions of energy intake show reduced relative brain size (20, 21). These trade-offs also have a developmental aspect: the maternal energy hypothesis (22) argues that the ability of mammalian mothers to transfer energy to their offspring constrains brain size (23). Various studies reported a link between maternal allocation and relative brain size in mammals (23, 24), birds (8), cichlid fish (25), and sharks and rays (26). However, they failed to provide an adaptive explanation for this association or did not evaluate alternative or complementary hypotheses.

Here we propose that paying systematic attention to the developmental costs of brains should improve our understanding of brain size evolution via its correlated evolution with

87   parental provisioning. Offspring face a seemingly insurmountable bootstrapping problem

posed by growing and differentiating large brains. How can offspring develop large brains

when they would need them to provide the energy necessary for this process? This catch-22

90   leads to the prediction that adult brain size across species depends on the parents' capacity

to provision their young. Thus, we examine the effect of parental provisioning on relative

brain size in birds, while simultaneously assessing hypotheses proposing eco- or socio-

93   cognitive benefits. This lineage is well suited for an integrated test of these different

hypotheses: birds show appreciable variation in many relevant traits, and the energy

transfer from parents to offspring can readily be assessed via egg volumes and the time

96   offspring are provisioned. Moreover, birds have a deeply rooted split in the developmental

state of the hatchlings, ranging from precocial (no or little provisioning of the young) to

altricial (extended provisioning), henceforth referred to as developmental mode (27).

99   Previous work noted that precocial species have smaller adult brain size than altricial ones

(28, 29), but the evolutionary causes of this pattern remain unclear. Birds also vary in the

number of providers (30). These differences allow us to distinguish the effects of investment

102   in eggs from those of provisioning of young. Moreover, birds show major variation in

ecology (aspects of their ecological niche, the climate in their geographic range) (31) and in

sociality (32).

This high variability in all relevant variables in birds allowed us to simultaneously test all

these effects on adult brain size. We assembled information on adult brain size, adult body

108   mass, predictors assessing parental provisioning (egg mass, clutch size, duration of food

provisioning, number of providers), ecology (fiber and energy content of food, complexity of

food acquisition, foraging substrate, sedentariness, insularity, climate in the geographic

111   range), and sociality (strength and stability of pair bonds, breeding alone or in colonies,

group sizes outside the breeding season) for 1,176 bird species. For the comparative tests,

we used phylogenetically controlled Bayesian mixed models (33) to assess the effects on

114   relative brain size across species of i) parental provisioning, ii) eco-social predictors, and iii)

all predictors combined. Next, to understand how the different predictors may have

affected each other's evolution, we used phylogenetic d-separation path analysis (34).

Results

Because body mass is tightly correlated with brain size, any model that incorporates body mass as an independent predictor of brain size will tend to mask the impact of the other, more informative predictors. We therefore report the results using relative brain size scaled by body mass as the response variable, but note that analyses including absolute brain size as a response variable produce qualitatively the same results (see S1a-c Table). Moreover, body mass scales many aspects of bird physiology and life-history in a non-linear manner. Thus, we include body mass and its interaction with key predictors in all models to capture these nonlinearities.

Phylogenetically controlled mixed models showed that larger relative brain sizes in birds were positively associated with all components of parental provisioning (Fig 1, S2a-d Tables). Larger relative brains were found in species that are altricial, have larger relative egg mass, smaller clutches, and feed their young for a longer time. The effect of these predictors was modified by interactions among them, further highlighting the importance of parental provisioning (Fig 1). Accordingly, only precocial species showed a negative correlation between relative brain size and clutch size (Fig 2a), while only altricial species showed a positive correlation between relative brain size and the time offspring are fed (Fig 2b). The remaining interaction effects reflected fundamental allometries or differences in body mass between precocial and altricial species (Fig 1). In sum, the total amount of energy invested into each individual young strongly affected adult brain size.

Fig 1. Estimated coefficients and effect sizes of phylogenetically controlled mixed models of the parental provisioning model, the eco-social model, and the combined model on relative brain size in birds. Color-filled circles denote estimated effects, lines denote the 95% lower and upper confidence limits generated in the R package MCMCglmm (33) based on a consensus phylogeny. Symbols below the circles denote the proportion of random tree models that reach statistical significance (p<0.05) for each predictor based on model averaging of 200 models, using a set of random trees from http://birdtree.org (35). The corresponding full models are shown in S2a-c Table.

150   When only examining the effect of eco-social predictors on relative brain size,
      phylogenetically controlled mixed models primarily showed that sedentary species and
      those that forage arboreally, had larger relative brain size than those that are migratory or

153   use other foraging substrates (Fig 2c,d). In terms of interaction effects, we found that
      relative brain size increases with body mass only in arboreal foragers, and that higher
      energy content of food translated into larger relative brain size only in larger species (Fig 1).

156   Social factors had a much weaker impact on relative brain size: species with enduring pair

bonds tended to have a larger relative brain size than species without pair bonds, whereas no correlations were found for the other social predictors (Fig 1).

Fig 2. The effect of parental provisioning (a, b) and ecological parameters (c, d) on relative brain size in birds (N=1,176 species). a) Clutch size vs developmental mode (precocial vs altricial). **b)** Time provisioned vs developmental mode (precocial vs altricial). **c)** Movement pattern: sedentary vs migratory. **d)** Foraging substrate: others vs arboreal. In **a-b**, solid lines designate linear trend estimates, grey shading designate standard error bands. In **c-d**, thick lines designate medians, the grey boxes designate inter-quantile ranges (IQR, 0.25-0.75 quantile), whiskers designate ±1.5 IQR, circles designate outliers.

The eco-social model explained markedly less variance in relative brain sizes than the

171   parental provisioning model ($R^2 = 0.38$ vs $R^2 = 0.50$). However, a combined model containing

both parental provisioning and eco-social predictors performed best ($R^2 = 0.57$). Although

confirming the patterns of the separate models (Fig 1), it revealed that the posterior

174   estimates of effect sizes of parental provisioning were far larger than those of the eco-social

factors. Among the latter, the ecological predictors that remain significant in the combined

model were largely linked to the adult foraging niche and movement patterns, whereas

177   social predictors no longer played any role.

Additional models showed that data limitations are not responsible for the observed

180   patterns. Models that excluded parameters with limited data availability (time fed, social

grouping; S3 a-b Tables) confirmed a key role of parental provisioning for relative brain size.

Similarly, using a more detailed categorization of the development mode spectrum instead

183   of binary categorization recovered the same patterns (S3c Table), and so did a model that

controlled for the possible confounding effect of longevity on parental provisioning

parameters (S2d Table).

The effect of parental provisioning on relative brain size became obvious when

superimposed on a phylogenetic tree (Fig 3). Starting at 3 o'clock and moving counter-

189   clockwise, early emerging, precocial lineages had relatively small brains, while inter-

dispersed altricial lineages (seabirds, pigeons, hummingbirds, swifts) had slightly larger

brains. With the origin of Inopinaves (core land birds) (36) some 65 million years ago,

192   systematic altriciality and extensive parental provisioning arose, resulting in several clusters

of notably large-brained lineages, including owls, parrots, and corvids, while one lineage

(nightjars) reverted to precociality and small brains. The most recently evolved and highly

195   diverse other passerines were small-bodied and did not have brains of exceptionally large

relative size, although they are still relatively larger than those of precocial lineages.

198   **Fig 3. Phylogenetic distribution of relative brain size, developmental mode (precocial vs
altricial), and time parent(s) feed their offspring (N=1,176 bird species).** Size of bars in
outer circle represent body mass (ln(grams)); size of bars in middle circle represents relative

201   brain size (ln(mL)) while their color represents the time parents feed their offspring (ln(days

+ 1)); inner circle represents developmental mode (precocial, altricial). Phylogeny follows birdrtree.org (35), whereas taxonomy largely follows Prum et al. (36), explaining why some clades appear polyphyletic.

207 Because the predictors analyzed above may not only have affected relative brain size directly, but may also have selected for evolutionary changes in values of other predictors and thus affected brain size indirectly, we performed phylogenetic path analyses (34, 37) to

210 gain better insights into the causal relationships between traits. We included well-established developmental and eco-social relationships in all models, but also added biologically plausible ones involving brain size that were varied across the different models.

The model assessing the relationships among the various parental provisioning predictors of relative brain size showed that adult body mass affects all other traits, including

216 developmental mode (Fig 4a). Both a large adult body mass and altriciality had a negative effect on clutch size and residual egg mass. Altriciality had a strong positive effect on the time offspring are fed, and subsequently on brain size. These patterns reflect established

219 life-history trade-offs and their cascading effects on the total amount of energy transferred from parents to their offspring at the different developmental stages. The remaining effect of body mass on relative brain size depended on developmental mode, reflecting that larger

222 altricial species had larger relative brain sizes than precocial species, relative to what would be predicted from body size alone (see Fig 1). This pattern therefore fully supports the pervasive effect of parental provisioning on a species' brain size.

Turning to the eco-social predictors, the path model assessing the evolutionary relationships among them showed that some of these predictors have strong evolutionary links among

228 themselves, but rather weak selective effects on relative brain size (Fig 4b). A combined path analysis (S1 Fig) including both parental provisioning and eco-social predictors retained the relationships revealed by the separate and combined phylogenetically controlled mixed

231 models. Parental provisioning continued to have a direct and strong effect on relative brain size, whereas that of the eco-social variables remains weak.

234 **Fig 4. Phylogenetic associations among predictors and relative brain size, assessed by d-separation path analyses** (34) **(N=1,176 bird species).** Arrows thickness and numeric values show the strength of the association, arrow color shows its direction (positive: orange,

237 negative: blue). **a,** Associations among parental provisioning predictors. **b,** Associations

among eco-social predictors. The plot showing the model including both the parental provisioning and eco-social parameters is shown in S1 Fig.

243 **Discussion**

These findings have three major implications, which we now discuss in turn. First, parental provisioning is the main predictor of adult relative brain size in birds: in the combined model, adult eco- and socio-cognitive demands hardly play a role. Thus, large brains can only be sustained in species with extensive energy inputs during development. This association is supported by previous work in diverse lineages (8, 24-26). However, these studies either did not provide an underlying mechanism or did not consider the relationship to reflect adaptation (23, 24). The parental provisioning hypothesis offers a novel theoretical underpinning to this pattern. Parental provisioning enables growing young to construct a large brain during a life stage when they themselves have not yet learned the skills needed to acquire the energy necessary to sustain such a growing and learning brain. Accordingly, parental provisioning removed this bottleneck for the evolution of larger brain size, and enabled bird species to evolve into skill-intensive niches, which reduce interspecific competition and thereby promoting survival prospects and population stability (see (12)). Its importance is also highlighted by the phylogenetic path analyses, which confirm that its components form a tightly coevolved set.

Second, the parental provisioning hypothesis can explain the well-established pattern that altricial species have relatively larger brains than precocial (28, 29). Precocial young are only provisioned through egg mass, which is subject to obvious physical limitations. As a result, they experience serious constraints on relative brain size and therefore niche complexity. The evolutionary transitions to altriciality allowed species to evolve into cognitively more demanding niches, which require more elaborate food-handling or anti-predator skills (29). This was made possible by parents donating the requisite large brain through extended parental provisioning. However, these transitions were actually quite rare (29). Although the underlying reasons for this rarity remain unclear, we speculate that precocial species remain stuck in a simple ecological niche unless they find a way to breed in safer places, which opens the path to provisioning and therefore altriciality.

Finally, our findings imply a modest role at best for socio- and eco-cognitive abilities in the evolution of brain size in birds. We found almost no detectable effect of the cognitive demands of adult social life. We did, however, find weak evidence for effects of eco-cognitive parameters independent of parental provisioning that affect the energy balance of

276   adults. First, migratory species have smaller brains than resident species (38), indicating that the high energy cost of migration reduces the ability to sustain a large adult brain. Second, the effect of arboreal foraging may be more related to adult survival, as in mammals (39),

279   thus removing a constraint on brain enlargement.

The absence of an effect of the traditionally measured socio- and eco-cognitive variables on

relative brain size is puzzling. It may actually reflect a more general problem in comparative cognition, as popular tests often fail to produce a differentiation between species generally thought to show major differences (40). Thus, these tests may not assess those cognitive

adaptations that produce fitness benefits by scaffolding critical foraging and predation-avoidance skills and improving the ability to provision the young and/or improve survival. Thus, our results suggest we should pay closer attention to cognitive adaptations such as

general behavioral flexibility (as enabled by executive functions and domain-general intelligence (41)), including the coordination of parental duties (essentially linked to planning, decision-making, and time management). These cognitive abilities may be

expressed behaviorally in avoidance of predation on eggs, nestlings and adults (42, 43), in maximization of food intake under everchanging food and weather conditions, or in the socio-cognitive ability to coordinate offspring provisioning (5). For example, some birds

flexibly adjust parenting in response to predation risk to themselves and their nestlings (44), as confirmed experimentally (45). However, because these cognitive abilities are not readily estimated in a way that can be compared across species, they are so far not included in

comparative analyses of brain size evolution.

To conclude, the amount and duration of parental provisioning is the key to understand

300   brain size evolution in birds. It could also explain the major differences in relative brain size between endotherms and ectotherms (1), which is broadly correlated with the amount of parental provisioning. Moreover, it may also apply to human evolution because we humans

303   have the largest relative brain size among all animals, and parents and other caregivers engage in provisioning children well into adolescence (46).

306   **Methods**

Data collection

We gathered information quantifying parental provisioning and the socio-ecological
correlates identified by previous work as influencing brain size in animals. We collected
these data from the online version of the Handbooks of the Birds of the World (47), region
specific handbooks (48, 49), existing published datasets (32, 50-52), as well as species
specific studies listed in the dataset. Data on brain size were collected from three sources
(9, 10, 53).

The following parental provisioning predictors were included: adult body mass,
developmental mode, egg mass, clutch size, the time offspring are fed by their parents until
nutritional independence, and the number of caretakers. We did not use imputed data
based on mean values of related species, which limited our sample size compared to other
comparative studies. Adult body mass data were calculated as means if mass of males and
females were listed separately. The developmental mode was assessed dichotomously.
Precocial species are those which hatch with open eyes and abandon the nest within 2 days
after hatching, and here also encompass semi-precocial species (young hatch with open
eyes and are soon capable of walking, but they remain in the nest to be fed by parents).
Altricial species are those in which young hatch with closed eyes and largely lack feathers
and are fed by parents in the nest, and here also encompass semi-altricial species (young
hatch with open eyes and are feathered, but remain in the nest to be fed by their parents)
(29). Given that this dichotomous split could obscure patterns, we did re-run the main
analyses using all four categories (fully precocial, semi-precocial, semi-altricial, fully altricial).
Egg mass was either assessed through direct measurements of egg mass, or was calculated
based on egg measurements, using the formula by Hoyt (54). For those species where
multiple egg dimension measurements were available in the literature, the mean values of
all provided measurements were used. For those species where egg dimension or mass data
were not available in handbooks, we used information in published data compilations listed
in the dataset (52, 55-57). Clutch size data corresponded to the median clutch size (43, 57).
The time young are fed by their parents included both the time parents provision their
young in the nest as well as thereafter until young are nutritionally independent from their
parents, as reported in handbooks (see above), and published datasets (28). We also
included the mean number of caretakers, where mound-nesting species had zero
caretakers, species with uniparental care one, species with bi-parental care two, and

cooperatively breeding species the mean value of caretakers. These data were obtained from handbooks (47), existing datasets (30, 53), and species-specific studies listed in the dataset.

Based on previous studies on the drivers of brain size in vertebrates, the following ecological predictors were included: food energy, fiber content of food, number of processing steps required to extract food, foraging substrate, sedentariness, insularity, and climatic predictors. The predictors relating to food were calculated based on an existing database (50). We calculated the mean energy and fiber content of foods per 100g from previous studies (19, 58), as well as the number of processing steps required to extract food (59). Given that these three predictors are highly correlated (that is, low caloric food usually has a high fiber content and requires only a single processing step), we used a PCA approach and extracted two independent PCs from these predictors (see below). The foraging substrate data were obtained from a published database (51). We categorized species into those that forage arboreally, compared to all other foraging substrates. In mammals, arboreal foraging has been shown to be associated with increased longevity, which in turn allows the evolution of larger brains (39). Sedentariness was assessed based on the maximum movement pattern of a species, separating sedentary species (including local movements) from migratory species (short- and long-distance migrants, altitudinal migrants), combining data from three sources (9, 32, 51). In addition, we also included data on the insularity of species, distinguishing continental from insular species (11, 51). Missing data were directly obtained from the primary sources listed above. Finally, we included data on the climate in the species' geographic range (60, 61), using a PCA approach (see below). Values of species with contradicting values in different published datasets were confirmed based on the literature listed above, using the above definitions.

The social predictors were also based on previous studies and included social bond strength, group size, and coloniality. Social bond data were based on published data (62), separating species with short bonds during mating only, seasonal bonds, and long-term bonds that extend beyond a single breeding season. The grouping patterns of species outside the breeding season distinguished between asocial species (usually on their own), pair living species (usually in pairs), species that live in small groups (usually 3-30 individuals), and

372   large groups (usually more than 30 individuals). The split between small and large groups

should capture the difference between personalized groups where group members know

each other individually, including family groups (32), and ephemeral and/or anonymous

375   groups. We also included whether species breed singly or whether they breed in colonies.

Analyses

378   All statistical analyses were carried out in the R 4.0.2 environment (63). The analyses were

structured as a series of linear models with increasing complexity, all including phylogenetic

relatedness across species (in the form of a phylogenetic correlation matrix based on well-

381   resolved avian phylogenetic trees (35)). For predictors that are expected to be allometrically

linked to body mass (brain size, egg mass), the models included their residuals from

regressions against body mass. Brain size (the response variable) and adult body mass were

384   log-transformed prior to extracting relative brain sizes. All continuous predictors were

mean-centered and scaled to a standard deviation of one, to aid in the interpretation of

intercepts and the comparison of effect sizes. The three predictors relating to food (food

387   energy, fiber content, food handling levels) were strongly collinear (S4a Table). Thus, we

used a Principal Component analysis in the package psych (64) (using varimax rotation) to

reduce these predictors to two uncorrelated components. Based on their loadings, the

390   resulting two components describe the energy content of food and the food handling levels

(including the absence of fibers) (S4b Table). Using the same approach, we extracted three

uncorrelated components from 14 climatic parameters. PC1 reflects a latitudinal gradient

393   from the tropics to high latitudes. PC2 reflects inter-annual variability. PC3 reflects a

gradient from dry, open and highly seasonal environments to stable and closed

environments.

We fitted three phylogenetically controlled mixed models that assessed the dependence of

relative brain size on provisioning predictors (model 1), eco-social predictors (model 2), and

399   a combination of both (model 3). Each model included a set of main effects and first-order

interactions with meaningful biological interpretation (see S1-3 Tables for a full list of

predictors and interactions in each of the models). Since the size of models (both in terms of

402   data and numbers of predictors) prevented a full search of all predictor combinations via

selection based on information criteria (IC), we conducted backward simplification of

models, removing non-significant interactions (starting with the ones having the largest

prediction error). To remain conservative, we did not remove non-significant main effects. Backward selection has been shown to perform well in comparison to IC-based model selection in similar contexts (65), and is robust in cases where the number of individual

observations exceeds several fold the number of estimated predictors, as is the case here. We report final outputs of the most parsimonious models (Fig 1, S1-3 Tables). All models were fitted using an MCMC-based mixed modelling approach using the package

MCMCglmm (33). We used chains of 50'000 iterations, with the first 10'000 iterations discarded as burn-ins and thinned every 40 iterations. Inspection of the final MCMC samples did not show any sign of autocorrelation. All effective sample sizes were close to the actual

numbers of sampled draws from predictors' posteriors. In all models, inverse-gamma priors were used for residual variances (parametrized as inverse-Wishart with V = 1 and v = 0.002). The prior for phylogenetic effect was formed as a weakly informative half-Cauchy density

(parameter expanded priors with V = 1, v = 1, α_μ = 0 and α_V = 10'000). Priors for fixed effects were left as default (Gaussian densities with μ = 0 and large variance). Using a modification of the variance inflation factor (VIF) analysis adjusted to the phylogenetic comparative

mixed model used, we verified that none of the other predictors were significantly collinear, as all VIF values were <2 (S5 Table).

The necessity of complete information on all predictors reduced the sample size of the main models to 1,176 species. Thus, we fitted additional models excluding those predictors that most strongly limited our sample size: excluding duration of parental provisioning (N=1,458

species); excluding duration of parental provisioning and social grouping (N=1,594 species). Both models recaptured the patterns observed in the full model (S2 Table), confirming that our sample was not biased by limited data (S3a-b Tables).

429
Subsequent to the phylogenetically controlled mixed models, we conducted phylogenetic path analyses to assess the causal links among the predictors included in parental

432 provisioning, the eco-social and the combined model. Briefly, for each model we defined a set of d-separation statements (34, 37, 66), describing a hypothesis represented by a directed acyclic graph, linking all involved predictors with causal links. We included well-

435 established developmental and eco-social patterns in all models, but also added biologically

plausible relationships that were varied across the different models. The sequence of models considered reflects hypotheses related to brain size predictors, as outlined in S2-4

Figs. The statements were then represented as sets of linear mixed models and fitted using a phylogenetic comparative framework implemented in the ASReml-R package (67). In models involving binary responses (e.g., precocial/altricial), we used generalized linear

mixed models with probit link function to fit models, which makes their regression coefficients directly comparable with standardized regression coefficients from general linear models for continuous predictors. Since all continuous predictors were scaled to unit

standard deviation, results for categorical predictors can be interpreted as standardized Cohen's d effect sizes, and thus, can be directly compared with continuous standardized regression coefficients. The best-fitting hypothesis in each model set was determined via

comparisons of CICc (corrected Fisher's C Information Criterion, a path analysis analogue of AICc (37)). In the parental provisioning and eco-social model, the best model was clearly supported over the alternatives (ΔCICc > 2 for top models). In the combined model, two

models were clearly supported over the alternatives, but were conceptually identical (S6 Table).

Code availability

Code used for the analyses and plotting the figures are available from S.M.D. upon request.

456 **Data availability**

The data are provided in a supplement.

459 **References**

1. H. J. Jerison, *Evolution of the Brain and Intelligence* (Academic Press, New York, 1973).

462 2. S. M. Reader, Y. Hager, K. N. Laland, The evolution of primate general and cultural intelligence. *Philosophical Transactions of the Royal Society B: Biological Sciences* **366**, 1017-1027 (2011).

465 3. R. O. Deaner, K. Isler, J. Burkart, C. Van Schaik, Overall brain size, and not encephalization quotient, best predicts cognitive ability across non-human primates. *Brain, behavior and evolution* **70**, 115-124 (2007).

468 4. S. E. Overington, J. Morand-Ferron, N. J. Boogert, L. Lefebvre, Technical innovations drive the relationship between innovativeness and residual brain size in birds. *Animal Behaviour* **78**, 1001-1010 (2009).

471 5. R. I. Dunbar, S. Shultz, Evolution in the social brain. *Science* **317**, 1344-1347 (2007).

6.   S. Shultz, R. Dunbar, Encephalization is not a universal macroevolutionary phenomenon in mammals but is associated with sociality. *Proceedings of the National Academy of Sciences* **107**, 21582-21586 (2010).

7.   R. I. Dunbar, The social brain hypothesis. *Evolutionary Anthropology: Issues, News, and Reviews: Issues, News, and Reviews* **6**, 178-190 (1998).

8.   S. Shultz, R. I. Dunbar, Social bonds in birds are associated with brain size and contingent on the correlated evolution of life-history and increased parental investment. *Biological Journal of the Linnean Society* **100**, 111-123 (2010).

9.   F. Sayol *et al.*, Environmental variation and the evolution of large brains in birds. *Nature Communications* **7**, 13971 (2016).

10.   T. S. Fristoe, C. A. Botero, Alternative ecological strategies lead to avian brain size bimodality in variable habitats. *Nature Communications* **10**, 1-9 (2019).

11.   F. Sayol, P. A. Downing, A. N. Iwaniuk, J. Maspons, D. Sol, Predictable evolution towards larger brains in birds colonizing oceanic islands. *Nature Communications* **9**, 2820 (2018).

12.   T. S. Fristoe, A. N. Iwaniuk, C. A. Botero, Big brains stabilize populations and facilitate colonization of variable habitats in birds. *Nature ecology & evolution* **1**, 1706-1715 (2017).

13.   P. Minias, P. Podlaszczuk, Longevity is associated with relative brain size in birds. *Ecology and evolution* **7**, 3558-3566 (2017).

14.   X. Yu *et al.*, Large-brained frogs mature later and live longer. *Evolution* **72**, 1174-1183 (2018).

15.   N. L. Barrickman, M. L. Bastian, K. Isler, C. P. van Schaik, Life history costs and benefits of encephalization: a comparative test using data from long-term studies of primates in the wild. *Journal of human evolution* **54**, 568-590 (2008).

16.   A. L. Bauernfeind *et al.*, Aerobic glycolysis in the primate brain: reconsidering the implications for growth and maintenance. *Brain Structure and Function* **219**, 1149-1167 (2014).

17.   J. W. Mink, R. J. Blumenschine, D. B. Adams, Ratio of central nervous system to body metabolism in vertebrates: its constancy and functional basis. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology* **241**, R203-R212 (1981).

18.   L. C. Aiello, P. Wheeler, The expensive-tissue hypothesis: the brain and the digestive system in human and primate evolution. *Current anthropology* **36**, 199-221 (1995).

19.   K. Isler, C. van Schaik, Costs of encephalization: the energy trade-off hypothesis tested on birds. *Journal of Human Evolution* **51**, 228-243 (2006).

20.   J. T. van Woerden, E. P. Willems, C. P. van Schaik, K. Isler, Large brains buffer energetic effects of seasonal habitats in catarrhine primates. *Evolution: International Journal of Organic Evolution* **66**, 191-199 (2012).

21.   S. A. Heldstab, K. Isler, C. P. van Schaik, Hibernation constrains brain size evolution in mammals. *J. Evol. Biol.* **31**, 1582-1588 (2018).

22.   R. D. Martin, Relative brain size and basal metabolic rate in terrestrial vertebrates. *Nature* **293**, 57-60 (1981).

23.   R. D. Martin, Scaling of the mammalian brain: the maternal energy hypothesis. *Physiology* **11**, 149-156 (1996).

24.   R. A. Barton, I. Capellini, Maternal investment, life histories, and the costs of brain growth in mammals. *Proceedings of the National Academy of Sciences* **108**, 6169-6174 (2011).

25. M. Tsuboi *et al.*, Comparative support for the expensive tissue hypothesis: big brains are correlated with smaller gut and greater parental investment in Lake Tanganyika cichlids. *Evolution* **69**, 190-200 (2015).

26. C. G. Mull, K. E. Yopak, N. K. Dulvy, Maternal investment, ecological lifestyle, and brain evolution in sharks and rays. *The American Naturalist* **195**, 1056-1069 (2020).

27. A. Portmann, Etudes sur la cérébralisation chez les oiseaux. III Cérébralisation et mode ontogénétique. *Alauda* **15**, 1-171 (1947).

28. A. N. Iwaniuk, J. E. Nelson, Developmental differences are correlated with relative brain size in birds: a comparative analysis. *Canadian Journal of Zoology* **81**, 1913-1928 (2003).

29. J. M. Starck, R. E. Ricklefs, *Avian growth and development: evolution within the altricial-precocial spectrum* (Oxford University Press, 1998).

30. A. Cockburn, Prevalence of different modes of parental care in birds. *Proceedings of the Royal Society B-Biological Sciences* **273**, 1375-1383 (2006).

31. A. L. Pigot *et al.*, Macroevolutionary convergence connects morphological form to ecological function in birds. *Nature Ecology & Evolution* **4**, 230-239 (2020).

32. M. Griesser, S. M. Drobniak, S. Nakagawa, C. A. Botero, Family living sets the stage for cooperative breeding and ecological resilience in birds. *PLoS Biology* **15**, e2000483 (2017).

33. J. D. Hadfield, MCMC methods for multi-response generalized linear mixed models: the MCMCglmm R package. *Journal of Statistical Software* **33**, 1-22 (2010).

34. A. v. Hardenberg, A. Gonzalez-Voyer, Disentangling evolutionary cause-effect relationships with phylogenetic confirmatory path analysis. *Evolution: International Journal of Organic Evolution* **67**, 378-387 (2013).

35. W. Jetz, G. H. Thomas, J. B. Joy, K. Hartmann, A. O. Mooers, The global diversity of birds in space and time. *Nature* **491**, 444-448 (2012).

36. R. O. Prum *et al.*, A comprehensive phylogeny of birds (Aves) using targeted next-generation DNA sequencing. *Nature* **526**, 569 (2015).

37. B. Shipley, The AIC model selection method applied to path analytic models compared using ad-separation test. *Ecology* **94**, 560-564 (2013).

38. O. Vincze, Light enough to travel or wise enough to stay? Brain size evolution and migratory behavior in birds. *Evolution* **70**, 2123-2133 (2016).

39. C. P. van Schaik, R. O. Deaner, Life history and brain evolution. *Animal social complexity: intelligence, culture, and individualized societies (eds FBM de Waal & PL Tyack)*, 1-25 (2003).

40. C. Fichtel, K. Dinter, P. M. Kappeler, The lemur baseline: how lemurs compare to monkeys and apes in the Primate Cognition Test Battery. *PeerJ* **8**, e10025 (2020).

41. J. M. Burkart, M. N. Schubiger, C. P. van Schaik, The evolution of general intelligence. *Behavioral and Brain Sciences* **40** (2017).

42. T. E. Martin, J. V. Briskie, "Predation on Dependent Offspring A Review of the Consequences for Mean Expression and Phenotypic Plasticity in Avian Life History Traits" in Year in Evolutionary Biology 2009. (Blackwell Publishing, Oxford, 2009), vol. 1168, pp. 201-217.

43. M. Valcu, J. Dale, M. Griesser, S. Nakagawa, B. Kempenaers, Global gradients of avian longevity support the classic evolutionary theory of ageing. *Ecography* **37**, 930-938 (2014).

44. S. L. Lima, Predators and the breeding bird: behavioral and reproductive flexibility under the risk of predation. *Biol. Rev.* **84**, 485-513 (2009).

45. S. Eggers, M. Griesser, J. Ekman, Predator-induced plasticity in nest visitation rates in the Siberian jay (Perisoreus infaustus). *Behavioral Ecology* **16**, 309-315 (2005).

46. H. Kaplan, K. Hill, J. Lancaster, A. M. Hurtado, A theory of human life history evolution: Diet, intelligence, and longevity. *Evolutionary Anthropology: Issues, News, and Reviews: Issues, News, and Reviews* **9**, 156-185 (2000).

47. J. Del Hoyo, A. Elliot, J. Sargatal, D. A. Christie, *Handbook of the Birds of the World* (Lynx Editions, Barcelona, 2011).

48. P. A. R. Hockey, W. J. J. Dean, P. G. Ryan, *Roberts Birds of Southern Africa* (Trustees of the John Voelcker Bird Book Fund, Cape Town, ed. 7th edition, 2005).

49. P. J. Higgins, S. Marchant, J. M. Peter, S. Cowling, J. Davies, *Handbook of Australian, New Zealand & Antarctic Birds* (Oxford University Press, 2007).

50. H. Wilman *et al.*, EltonTraits 1.0: Species-level foraging attributes of the world's birds and mammals. *Ecology* **95**, 2027-2027 (2014).

51. J. A. Tobias, A. L. Pigot, Integrating behaviour and ecology into global biodiversity conservation strategies. *Philosophical Transactions of the Royal Society B: Biological Sciences* **374**, 20190012 (2019).

52. T. Lislevand, J. Figuerola, T. Székely, Avian body sizes in relation to fecundity, mating system, display behavior, and resource sharing. *Ecology* **88**, 1605-1605 (2007).

53. S. Graber (2017) Social and ecological aspects of brain size evolution: a comparative approach. in *Department of Anthropology* (University of Zurich, Zurich), p 182.

54. D. F. Hoyt, Practical methods of estimating volume and fresh weight of bird eggs. *The Auk* **96**, 73-77 (1979).

55. K. Schläpfer, *The colorful eggs of the tinamous* (Published by author, Engelburg, Switzerland, 2017).

56. J.-Y. Juang *et al.*, The avian egg exhibits general allometric invariances in mechanical design. *Scientific Reports* **7**, 1-11 (2017).

57. C. Brown, P. Bridgeford, S. Braine, M. Paxton, W. Versfeld, Breeding data on the birds of Namibia: laying months, colony and clutch sizes and egg measurements. *Biodiversity Observations*, 92-196 (2015).

58. P. Langer, Lactation, weaning period, food quality, and digestive tract differentiations in eutheria. *Evolution* **57**, 1196-1215 (2003).

59. C. Schuppli, S. M. Graber, K. Isler, C. P. van Schaik, Life history, cognition and the evolution of complex foraging niches. *Journal of Human Evolution* **92**, 91-100 (2016).

60. U. Roll *et al.*, The global distribution of tetrapods reveals a need for targeted reptile conservation. *Nature Ecology & Evolution* **1**, 1677-1682 (2017).

61. G. Murali, R. Gumbs, S. Meiri, U. Roll, Global determinants and conservation of evolutionary and geographic rarity in land vertebrates. *Science advances* **7**, eabe5582 (2021).

62. J. A. Tobias *et al.*, Territoriality, social bonds, and the evolution of communal signaling in birds. *Frontiers in Ecology and Evolution* **4**, 74 (2016).

63. R Core Team (2016) R: A language and environment for statistical computing. (R Foundation for Statistical Computing, Vienna, Austria.).

64. W. Revelle (2012) Procedures for psychological, psychometric, and personality Research. R Package Version 1.9.12. (Evanston, Illinois).

65. P. A. Murtaugh, Performance of several variable-selection methods applied to real ecological data. *Ecology letters* **12**, 1061-1068 (2009).

66. B. Shipley, Confirmatory path analysis in a generalized multilevel context. *Ecology* **90**, 363-368 (2009).

67. D. Butler (2019) Asreml: Fits the linear mixed model. R package version 4.1.0.110. (VSNi, Hemel Hempstead, UK).

Acknowledgements

We thank all field biologists who collected all the natural history data underlying this study.

621   We thank Uri Roll for sharing climatic data.

Funding

624   M.G.: Heisenberg Grant nr. GR 4650/2-1 by the German Research Foundation DFG.

S.M.D: Polish National Science Centre, grant no. UMO-2015/18/E/NZ8/00505 and Australian Research Council, DECRA award no. DE180100202.

627   C.P.S: University of Zurich.

Author contributions

630   M.G. contributed to acquisition of funding, conceptualizing, data collection, data analyses, interpretation of the data, drafting the manuscript.

S.M.D. contributed the data analyses, interpretation of the data.

633   S.M.G. contributed to data collection, data analyses, revising the manuscript.

C.P. S. contributed to acquisition of funding, conceptualizing, interpretation of the data, drafting the manuscript.

Competing interests

None.

Materials & Correspondence

M.G.

Supporting Information

S1 Fig. Phylogenetic associations among parental provisioning (orange), ecological and social predictors (green) and relative brain size, assessed by d-separation path analyses (34) **(N=1,176 bird species).** Arrow thickness and numeric values show the strength of the association, arrow color its direction (orange: positive, blue: negative).

651 **S2 Fig. Network topologies of the top 15 parental provisioning models of the phylogenetic path analyses, arranged by increasing cCIC.** Grey arrows indicate relationships retained in all models, black arrows indicate relationships that varied between different models.

S3 Fig. Network topologies of the top 15 eco-social models of the phylogenetic path analyses,
arranged by increasing cCIC. Grey arrows indicate relationships retained in all models, black arrows indicate relationships that varied between different models.

S4 Fig. Network topologies of all 10 combined models of the phylogenetic path analyses, arranged by increasing cCIC. Grey arrows indicate relationships retained in all models, black arrows indicate relationships that varied between different models.

666 **S1a-c Table. Phylogenetically controlled mixed models in MCMCglmm(33) assessing the effect on *absolute* brain size in birds (N=1,176 species) of a) parental provisioning predictors, b) eco-social predictors, and c) both combined.**

	Predictor	Post. mean	lower ▪ upper 95% CI	N	p MCMC
a) Parental investment model	**Intercept**	**-0.30**	**-0.46 ▪ -0.11**	**1000**	**<0.001**
	Mass	-0.03	-0.07 ▪ 0.01	893	0.11
	Development mode (precocial vs altricial)	0.12	-0.02 ▪ 0.25	1000	0.10
	Clutch size	**-0.07**	**-0.09 ▪ -0.04**	**1000**	**<0.001**
	Residual egg mass	**0.09**	**0.04 ▪ 0.14**	**1000**	**0.002**
	Time fed	**0.09**	**0.04 ▪ 0.14**	**1000**	**<0.001**
	Number of caretakers	0.00	-0.01 ▪ 0.01	1000	0.51
	Mass x development mode	**0.18**	**0.13 ▪ 0.24**	**1000**	**<0.001**
	Mass x clutch size	**0.02**	**0.01 ▪ 0.04**	**1000**	**0.008**
	Mass x residual egg mass	**0.07**	**0.03 ▪ 0.11**	**1000**	**<0.001**
	Development mode x clutch size	**0.09**	**0.06 ▪ 0.13**	**1000**	**<0.001**
	Development mode x residual egg mass	**0.19**	**0.12 ▪ 0.28**	**1087**	**<0.001**
	Development mode x time fed	0.07	0.00 ▪ 0.16	1000	0.051
	Residual egg mass x clutch size	**0.09**	**0.07 ▪ 0.11**	**1000**	**<0.001**
b) Eco-social model	**Intercept**	**-0.4**	**-0.57 ▪ -0.22**	**1155**	**<0.001**
	Mass	**0.98**	**0.95 ▪ 1**	**1000**	**<0.001**
	Foraging substrate (others vs arboreal)	**0.11**	**0.06 ▪ 0.15**	**954**	**<0.001**
	Migratory behavior (sedentary vs migratory)	**0.05**	**0.03 ▪ 0.07**	**1190**	**<0.001**
	Energy content of food (PC)	0.00	-0.01 ▪ 0.01	1000	0.92
	Food handling level (PC)	0.00	-0.02 ▪ 0.02	1000	0.93
	Insularity (continental vs insular)	0.00	-0.04 ▪ 0.06	1000	0.89
	Mean diurnal temperature range (PC)	0.00	0.00 ▪ 0.01	1000	0.33
	Annual temperature and rainfall (PC)	-0.01	-0.03 ▪ 0.00	1000	0.064
	Climatic seasonality (PC)	0.00	-0.01 ▪ 0.01	1000	0.88
	Short vs seasonal social bonds	0.03	-0.01 ▪ 0.09	1000	0.18
	Short vs long-term social bonds	**0.07**	**0.02 ▪ 0.13**	**1000**	**0.002**
	Colonial vs solitary breeding	0.00	-0.02 ▪ 0.02	1131	0.82
	Asocial vs pair living	-0.03	-0.06 ▪ 0.01	1000	0.11
	Asocial vs small groups	-0.02	-0.05 ▪ 0.02	1000	0.35
	Asocial vs large groups	0.00	-0.03 ▪ 0.03	1288	0.96
	Mass x foraging substrate	**0.1**	**0.06 ▪ 0.14**	**861**	**<0.001**
	Mass x sedentariness	**-0.03**	**-0.05 ▪ -0.01**	**1000**	**0.01**
	Mass x energy content of food (PC)	**0.02**	**0.01 ▪ 0.03**	**1000**	**0.004**
c) Combined model	**Intercept**	**-0.36**	**-0.55 ▪ -0.18**	**1000**	**<0.001**
	Mass	**0.94**	**0.90 ▪ 0.97**	**901**	**<0.001**
	Development mode (precocial vs altricial)	**0.17**	**0.05 ▪ 0.29**	**1000**	**0.01**
	Clutch size	**-0.06**	**-0.09 ▪ -0.04**	**1000**	**<0.001**
	Residual egg mass	**0.1**	**0.05 ▪ 0.14**	**1000**	**<0.001**
	Time fed	**0.06**	**0.01 ▪ 0.11**	**1000**	**0.018**
	Number of caretakers	0.00	-0.01 ▪ 0.01	1000	0.67
	Mass x development mode	**0.16**	**0.11 ▪ 0.21**	**891**	**<0.001**
	Mass x clutch size	**0.02**	**0.00 ▪ 0.03**	**1000**	**0.016**
	Mass x residual egg mass	**0.06**	**0.03 ▪ 0.09**	**1000**	**<0.001**
	Development mode x clutch size	**0.11**	**0.08 ▪ 0.15**	**1000**	**<0.001**
	Development mode x residual egg mass	**0.17**	**0.10 ▪ 0.24**	**1000**	**<0.001**
	Development mode x time fed	0.07	0.00 ▪ 0.15	1000	0.078
	Residual egg mass x clutch size	**0.08**	**0.06 ▪ 0.10**	**1108**	**<0.001**
	Foraging substrate (others vs arboreal)	**0.09**	**0.05 ▪ 0.14**	**1000**	**<0.001**
	Migratory behavior (sedentary vs migratory)	**0.04**	**0.02 ▪ 0.06**	**1079**	**<0.001**
	Energy content of food (PC)	0.01	0.00 ▪ 0.03	1179	0.056
	Food handling level (PC)	0.02	-0.01 ▪ 0.04	972	0.22
	Insularity (continental vs insular)	0.01	-0.03 ▪ 0.06	1000	0.68
	Mean diurnal temperature range (PC)	0.00	-0.01 ▪ 0.01	1000	0.64
	Annual temperature and rainfall (PC)	-0.01	-0.02 ▪ 0.00	1000	0.23
	Climatic seasonality (PC)	0.00	-0.01 ▪ 0.01	1000	0.53
	Short vs seasonal social bonds	-0.01	-0.06 ▪ 0.04	1000	0.77
	Short vs long-term social bonds	0.02	-0.03 ▪ 0.07	1000	0.39
	Colonial vs solitary breeding	-0.01	-0.03 ▪ 0.02	1000	0.59
	Asocial vs pair living	-0.03	-0.06 ▪ 0.00	1098	0.10
	Asocial vs small groups	-0.02	-0.05 ▪ 0.01	1000	0.27
	Asocial vs large groups	0.00	-0.03 ▪ 0.02	1215	0.78
	Mass x foraging substrate	**0.06**	**0.02 ▪ 0.10**	**1000**	**0.008**
	Mass x sedentariness	**-0.03**	**-0.05 ▪ -0.01**	**1000**	**<0.001**
	Mass x energy content of food (PC)	0.01	0.00 ▪ 0.02	1000	0.13

	Post. mean	CI	N	p MCMC
Developmental mode x food handling level (PC)	-0.05	-0.08 ▪ -0.01	900	0.018
Clutch size x energy content of food (PC)	-0.01	-0.02 ▪ 0.00	1000	0.04
Clutch size x food handling level (PC)	-0.01	-0.03 ▪ 0.00	1000	0.014

Significant predictors are highlighted in bold. Post. mean = mean of the posterior distribution of a

672   parameter, CI = posterior credibility interval, N = effective sample size (corrected for MCMC

timeseries autocorrelation), p MCMC = P-value. The Principal Component analyses (PCAs) of climatic

and food parameters are shown in S4 Table.

S2a-d Table. Phylogenetically controlled mixed models in MCMCglmm(33) assessing the effect on *relative* brain size in birds (model a-c: N=1,176 species, model d: N= 886 species) of a) parental provisioning predictors, b) eco-social predictors, c) both combined, and d) both combined and including longevity.

	Predictor	Post. mean	lower ▪ upper 95% CI	p MCMC
a) Parental investment model	**Intercept**	**-0.29**	**-0.49 ▪ -0.10**	**1**
	Mass	-0.03	-0.07 ▪ 0.01	0.02
	Development mode (precocial vs altricial)	0.11	-0.02 ▪ 0.24	0.04
	Clutch size	**-0.07**	**-0.10 ▪ -0.04**	**1**
	Residual egg mass	**0.10**	**0.04 ▪ 0.15**	**1**
	Time fed	**0.09**	**0.04 ▪ 0.14**	**1**
	Number of caretakers	0.00	-0.01 ▪ 0.01	0
	Mass x development mode	**0.19**	**0.13 ▪ 0.24**	**1**
	Mass x clutch size	**0.02**	**0.01 ▪ 0.04**	**1**
	Mass x residual egg mass	**0.07**	**0.03 ▪ 0.10**	**1**
	Development mode x clutch size	**0.09**	**0.05 ▪ 0.13**	**1**
	Development mode x residual egg mass	**0.19**	**0.11 ▪ 0.27**	**1**
	Development mode x time fed	0.07	-0.01 ▪ 0.15	0
	Residual egg mass x clutch size	**0.09**	**0.06 ▪ 0.11**	**1**
b) Eco-social model	**Intercept**	**-0.44**	**-0.64 ▪ -0.24**	**1**
	Mass	**0.05**	**0.02 ▪ 0.07**	**1**
	Foraging substrate (others vs arboreal)	**0.12**	**0.07 ▪ 0.17**	**1**
	Migratory behavior (sedentary vs migratory)	**0.05**	**0.02 ▪ 0.07**	**1**
	Energy content of food (PC)	0.00	-0.01 ▪ 0.01	0
	Food handling level (PC)	0.00	-0.02 ▪ 0.02	0
	Insularity (continental vs insular)	0.00	-0.05 ▪ 0.06	0
	Mean diurnal temperature range (PC)	0.01	0.00 ▪ 0.02	0
	Annual temperature and rainfall (PC)	-0.01	-0.03 ▪ 0.00	0.34
	Climatic seasonality (PC)	0.00	-0.01 ▪ 0.01	0
	Short vs seasonal social bonds	0.04	-0.02 ▪ 0.09	0
	Short vs long-term social bonds	**0.08**	**0.02 ▪ 0.14**	**1**
	Colonial vs solitary breeding	0.00	-0.02 ▪ 0.03	0
	Asocial vs pair living	-0.03	-0.06 ▪ 0.01	0
	Asocial vs small groups	-0.02	-0.06 ▪ 0.02	0
	Asocial vs large groups	0.00	-0.04 ▪ 0.03	0
	Mass x foraging substrate	**0.10**	**0.06 ▪ 0.15**	**1**
	Mass x sedentariness	**-0.03**	**-0.05 ▪ -0.01**	**1**
	Mass x energy content of food (PC)	**0.02**	**0.01 ▪ 0.03**	**1**
c) Combined model	**Intercept**	**-0.37**	**-0.57 ▪ -0.17**	**1**
	Mass	-0.01	-0.05 ▪ 0.03	0
	Development mode (precocial vs altricial)	**0.17**	**0.04 ▪ 0.30**	**0.99**
	Clutch size	**-0.07**	**-0.10 ▪ -0.04**	**1**
	Residual egg mass	**0.10**	**0.05 ▪ 0.16**	**1**
	Time fed	**0.07**	**0.02 ▪ 0.12**	**1**
	Number of caretakers	0.00	-0.01 ▪ 0.01	0
	Mass x development mode	**0.18**	**0.13 ▪ 0.23**	**1**
	Mass x clutch size	**0.02**	**0.00 ▪ 0.04**	**1**
	Mass x residual egg mass	**0.07**	**0.03 ▪ 0.10**	**1**
	Development mode x clutch size	**0.12**	**0.08 ▪ 0.16**	**1**
	Development mode x residual egg mass	**0.19**	**0.11 ▪ 0.27**	**1**
	Development mode x time fed	0.07	-0.01 ▪ 0.15	0.02
	Residual egg mass x clutch size	**0.09**	**0.07 ▪ 0.11**	**1**
	Foraging substrate (others vs arboreal)	**0.10**	**0.05 ▪ 0.14**	**1**
	Migratory behavior (sedentary vs migratory)	**0.05**	**0.02 ▪ 0.07**	**1**
	Energy content of food (PC)	0.01	0.00 ▪ 0.03	0.05
	Food handling level (PC)	0.02	-0.01 ▪ 0.05	0
	Insularity (continental vs insular)	0.01	-0.04 ▪ 0.05	0
	Mean diurnal temperature range (PC)	0.00	-0.01 ▪ 0.01	0
	Annual temperature and rainfall (PC)	-0.01	-0.02 ▪ 0.00	0
	Climatic seasonality (PC)	0.00	0.00 ▪ 0.01	0
	Short vs seasonal social bonds	-0.01	-0.06 ▪ 0.04	0
	Short vs long-term social bonds	0.02	-0.03 ▪ 0.08	0
	Colonial vs solitary breeding	-0.01	-0.03 ▪ 0.02	0
	Asocial vs pair living	-0.03	-0.06 ▪ 0.01	0.01
	Asocial vs small groups	-0.02	-0.06 ▪ 0.02	0
	Asocial vs large groups	-0.01	-0.04 ▪ 0.03	0
	Mass x foraging substrate	**0.06**	**0.02 ▪ 0.10**	**1**

	Post. mean	CI	p MCMC
Mass x sedentariness	**-0.03**	**-0.05 ▪ -0.01**	**1**
Developmental mode x food handling level (PC)	**-0.05**	**-0.09 ▪ -0.02**	**1**
Clutch size x energy content of food (PC)	-0.01	-0.02 ▪ 0.00	0.36
Clutch size x food handling level (PC)	**-0.01**	**-0.03 ▪ 0.00**	**0.96**

		Post. mean	CI	p MCMC
d) Combined model including longevity	**Intercept**	**-0.36**	**-0.58 ▪ -0.14**	**1**
	Mass	0.01	-0.04 ▪ 0.06	0
	Development mode (precocial vs altricial)	**0.18**	**0.04 ▪ 0.33**	**0.99**
	Clutch size	**-0.08**	**-0.12 ▪ -0.04**	**1**
	Residual egg mass	**0.12**	**0.05 ▪ 0.18**	**1**
	Time fed	0.05	0.00 ▪ 0.11	0.1
	Number of caretakers	0.00	-0.01 ▪ 0.01	0
	Mass x development mode	**0.15**	**0.09 ▪ 0.22**	**1**
	Mass x clutch size	0.01	-0.01 ▪ 0.03	0
	Mass x residual egg mass	0.04	0.00 ▪ 0.08	0.26
	Development mode x clutch size	**0.13**	**0.09 ▪ 0.18**	**1**
	Development mode x residual egg mass	**0.17**	**0.08 ▪ 0.26**	**1**
	Development mode x time fed	**0.13**	**0.04 ▪ 0.23**	**1**
	Residual egg mass x clutch size	**0.08**	**0.06 ▪ 0.11**	**1**
	Foraging substrate (others vs arboreal)	**0.09**	**0.04 ▪ 0.14**	**1**
	Migratory behavior (sedentary vs migratory)	**0.04**	**0.01 ▪ 0.07**	**1**
	Energy content of food (PC)	0.01	0.00 ▪ 0.03	0
	Food handling level (PC)	0.01	-0.03 ▪ 0.04	0
	Insularity (continental vs insular)	-0.01	-0.07 ▪ 0.05	0
	Mean diurnal temperature range (PC)	-0.01	-0.02 ▪ 0.01	0
	Annual temperature and rainfall (PC)	0.00	-0.02 ▪ 0.01	0
	Climatic seasonality (PC)	0.00	-0.01 ▪ 0.01	0
	Short vs seasonal social bonds	0.00	-0.07 ▪ 0.06	0
	Short vs long-term social bonds	0.04	-0.03 ▪ 0.11	0
	Colonial vs solitary breeding	-0.01	-0.04 ▪ 0.01	0
	Asocial vs pair living	-0.02	-0.06 ▪ 0.02	0
	Asocial vs small groups	-0.04	-0.08 ▪ 0.01	0
	Asocial vs large groups	0.00	-0.04 ▪ 0.04	0
	Mass x foraging substrate	**0.06**	**0.01 ▪ 0.11**	**0.98**
	Mass x sedentariness	**-0.03**	**-0.05 ▪ -0.01**	**1**
	Mass x energy content of food (PC)	0.01	-0.01 ▪ 0.02	0
	Developmental mode x food handling level (PC)	-0.05	-0.09 ▪ 0.00	0.71
	Clutch size x energy content of food (PC)	-0.01	-0.03 ▪ 0.00	0.12
	Clutch size x food handling level (PC)	-0.01	-0.03 ▪ 0.00	0.02
	Longevity	0.01	-0.01 ▪ 0.02	0
	Mass x longevity	**-0.01**	**-0.03 ▪ 0.00**	**0.98**

681 Significant predictors that are supported by at least 95% of different models are highlighted in bold.

Post. mean = mean of the posterior distribution of a parameter, CI = posterior credibility interval, p

MCMC = averaged P-values over all 200 models with different phylogenetic trees. The Principal

684 Component analyses (PCAs) of climatic and food parameters are shown in S4 Table.

S3 Table. Additional phylogenetically controlled mixed models in MCMCglmm(33) assessing the effect of parental provisioning and eco-social parameters on the evolution of relative brain size in birds.

	Predictor	Post. mean	lower ▪ upper 95% CI	N	pMCMC
a) excluding duration parental provisioning	**Intercept**	**-0.44**	**-0.64 ▪ -0.26**	**1000**	**<0.001**
	Mass	-0.01	-0.04 ▪ 0.03	1000	0.77
	Development mode (precocial vs altricial)	**0.24**	**0.13 ▪ 0.36**	**977**	**<0.001**
	Clutch size	**-0.07**	**-0.10 ▪ -0.04**	**1722**	**<0.001**
	Residual egg mass	**0.05**	**0.00 ▪ 0.10**	**895**	**0.034**
	Number of caretakers	0.00	-0.01 ▪ 0.01	1000	0.42
	Mass x development mode	**0.19**	**0.14 ▪ 0.24**	**1000**	**<0.001**
	Mass x clutch size	**0.02**	**0.01 ▪ 0.03**	**1000**	**<0.001**
	Mass x residual egg mass	**0.11**	**0.08 ▪ 0.14**	**1117**	**<0.001**
	Development mode x clutch size	**0.11**	**0.07 ▪ 0.15**	**1446**	**<0.001**
	Development mode x residual egg mass	**0.27**	**0.21 ▪ 0.34**	**1000**	**<0.001**
	Residual egg mass x clutch size	**0.08**	**0.06 ▪ 0.10**	**1000**	**<0.001**
	Foraging substrate (others vs arboreal)	**0.11**	**0.08 ▪ 0.15**	**1000**	**<0.001**
	Migratory behavior (sedentary vs migratory)	**0.05**	**0.03 ▪ 0.07**	**980**	**<0.001**
	Energy content of food (PC)	**0.02**	**0.00 ▪ 0.03**	**1000**	**0.008**
	Food handling level (PC)	0.02	-0.01 ▪ 0.04	1151	0.16
	Insularity (continental vs insular)	0.03	-0.01 ▪ 0.08	1000	0.13
	Mean diurnal temperature range (PC)	0.00	-0.01 ▪ 0.01	1220	1.00
	Annual temperature and rainfall (PC)	**-0.01**	**-0.02 ▪ 0.00**	**1000**	**0.018**
	Climatic seasonality (PC)	0.00	-0.01 ▪ 0.01	1000	0.90
	Short vs seasonal social bonds	-0.02	-0.07 ▪ 0.03	844	0.48
	Short vs long-term social bonds	0.01	-0.04 ▪ 0.06	876	0.77
	Colonial vs solitary breeding	0.00	-0.02 ▪ 0.02	1000	0.73
	Asocial vs pair living	-0.01	-0.05 ▪ 0.01	978	0.36
	Asocial vs small groups	-0.01	-0.05 ▪ 0.02	1000	0.61
	Asocial vs large groups	0.01	-0.02 ▪ 0.04	1000	0.38
	Mass x foraging substrate	**0.07**	**0.03 ▪ 0.11**	**1000**	**0.002**
	Mass x sedentariness	**-0.03**	**-0.05 ▪ -0.01**	**1000**	**<0.001**
	Developmental mode x food handling level (PC)	**-0.04**	**-0.08 ▪ -0.01**	**1115**	**0.016**
	Clutch size x energy content of food (PC)	**-0.01**	**-0.02 ▪ 0.00**	**1000**	**0.044**
	Clutch size x food handling level (PC)	**-0.01**	**-0.02 ▪ 0.00**	**1189**	**0.02**
b) excluding parental provisioning and social grouping	**Intercept**	**-0.45**	**-0.62 ▪ -0.27**	**1000**	**<0.001**
	Mass	0.00	-0.04 ▪ 0.04	1099	0.99
	Development mode (precocial vs altricial)	**0.23**	**0.12 ▪ 0.34**	**1000**	**<0.001**
	Clutch size	**-0.06**	**-0.09 ▪ -0.04**	**1000**	**<0.001**
	Residual egg mass	**0.08**	**0.03 ▪ 0.13**	**1000**	**0.006**
	Number of caretakers	0.01	0.00 ▪ 0.01	1020	0.18
	Mass x development mode	**0.16**	**0.11 ▪ 0.21**	**1000**	**<0.001**
	Mass x clutch size	**0.02**	**0.00 ▪ 0.03**	**1108**	**0.008**
	Mass x residual egg mass	**0.09**	**0.06 ▪ 0.12**	**1097**	**<0.001**
	Development mode x clutch size	**0.09**	**0.05 ▪ 0.12**	**1000**	**<0.001**
	Development mode x residual egg mass	**0.19**	**0.14 ▪ 0.26**	**1000**	**<0.001**
	Residual egg mass x clutch size	**0.08**	**0.06 ▪ 0.10**	**1000**	**<0.001**
	Foraging substrate (others vs arboreal)	**0.11**	**0.08 ▪ 0.15**	**1000**	**<0.001**
	Migratory behavior (sedentary vs migratory)	**0.06**	**0.04 ▪ 0.07**	**1000**	**<0.001**
	Energy content of food (PC)	0.01	0.00 ▪ 0.02	1093	0.062
	Food handling level (PC)	0.01	-0.01 ▪ 0.04	1000	0.34
	Insularity (continental vs insular)	0.02	-0.03 ▪ 0.06	1000	0.41
	Mean diurnal temperature range (PC)	0.00	-0.01 ▪ 0.01	1000	0.60
	Annual temperature and rainfall (PC)	**-0.01**	**-0.02 ▪ 0.00**	**1359**	**0.046**
	Climatic seasonality (PC)	0.00	-0.01 ▪ 0.01	1506	0.79
	Short vs seasonal social bonds	-0.02	-0.07 ▪ 0.03	1100	0.48
	Short vs long-term social bonds	0.01	-0.04 ▪ 0.06	1000	0.79
	Colonial vs solitary breeding	-0.01	-0.03 ▪ 0.02	1000	0.61
	Mass x foraging substrate	**0.07**	**0.04 ▪ 0.11**	**1000**	**<0.001**
	Mass x sedentariness	**-0.03**	**-0.05 ▪ -0.01**	**1000**	**0.002**
	Developmental mode x food handling level (PC)	-0.03	-0.06 ▪ 0.00	1000	0.052
	Clutch size x food handling level (PC)	**-0.01**	**-0.02 ▪ 0.00**	**1000**	**0.008**
c) using 4 developmental mode categories	Intercept	-0.11	-0.32 ▪ 0.06	1000	0.25
	Mass	**0.17**	**0.14 ▪ 0.21**	**1134**	**<0.001**
	Development mode (altricial vs precocial)	-0.23	-0.53 ▪ 0.11	1000	0.16
	Development mode (altricial vs semi-altricial)	-0.13	-0.32 ▪ 0.05	985	0.17
	Development mode (altricial vs semi-precocial)	**-0.26**	**-0.41 ▪ -0.11**	**1000**	**<0.001**
	Clutch size	0.02	-0.01 ▪ 0.05	1000	0.11

Residual egg mass	**0.29**	**0.22 ▪ 0.36**	**1000**	**<0.001**
Time fed	**0.20**	**0.13 ▪ 0.27**	**874**	**<0.001**
Number of caretakers	0.00	-0.01 ▪ 0.01	1000	0.71
Mass x development mode (altricial vs precocial)	**-0.22**	**-0.28 ▪ -0.16**	**1120**	**<0.001**
Mass x development mode (altricial vs semi-altricial)	-0.03	-0.12 ▪ 0.07	1000	0.54
Mass x development mode (alt. vs semi-precocial)	**-0.13**	**-0.21 ▪ -0.06**	**1000**	**<0.001**
Mass x clutch size	0.02	0.00 ▪ 0.04	1000	0.05
Mass x residual egg mass	**0.06**	**0.02 ▪ 0.10**	**1000**	**0.002**
Development mode (alt. vs precocial) x clutch size	**-0.06**	**-0.11 ▪ -0.01**	**1000**	**0.012**
Development mode (alt. vs semi-altricial) x clutch size	-0.01	-0.07 ▪ 0.07	1000	**0.89**
Development mode (alt. vs semi-prec.) x clutch size	**-0.13**	**-0.19 ▪ -0.07**	**1000**	**<0.001**
Develop. mode (alt. vs prec.) x residual egg mass	**-0.19**	**-0.33 ▪ -0.08**	**1095**	**0.002**
Develop. mode (alt. vs semi-alt.) x residual egg mass	0.06	-0.14 ▪ 0.25	1000	**0.484**
Develop. mode (alt. vs semi-prec.) x res. egg mass	**-0.19**	**-0.29 ▪ -0.08**	**1095**	**<0.001**
Development mode (altricial vs precocial) x time fed	-0.12	-0.30 ▪ 0.06	1000	0.164
Development mode (alt. vs semi-altricial) x time fed	**-0.14**	**-0.3 ▪ -0.01**	**764**	**0.048**
Development mode (alt. vs semi-prec.) x time fed	**-0.18**	**-0.27 ▪ -0.07**	**1000**	**<0.001**
Residual egg mass x clutch size	**0.09**	**0.05 ▪ 0.12**	**1000**	**<0.001**

690   Model **a)** combined model excluding the duration of parental provisioning: N = 1,458 species; model

b) combined model excluding parental provisioning and social grouping parameters: N = 1,594

species; model **c)** using four developmental mode categories: N=1,176 species. Significant predictors

693   are highlighted in bold. Post. mean = mean of the posterior distribution of a parameter, CI =

posterior credibility interval, N = effective sample size (corrected for MCMC timeseries

autocorrelation). The Principal Component analyses (PCAs) of climatic and food parameters are

696   shown in S4 Table.

S4 Table. Principal Component Analyses (PCA) loadings and variance explained. Loadings of the
699   **main contributors to the different component are highlighted in bold.**

a) Food PCA	PC F1	PC F2
Food energy content / 100g	0.12	**0.99**
Food fiber content / 100g	**-0.96**	-0.04
Food handling levels	**0.95**	0.17
SS loadings	1.84	1.01
Proportion Variance	0.61	0.34
Cumulative Variance	0.61	0.95
Proportion Explained	0.64	0.36
Cumulative Proportion	0.64	1.00

b) Climatic PCA	PC C1	PC C2	PC C3
Bio1_mean Annual Mean Temperature	**0.89**	0.06	0.04
Bio1_stdev	**-0.62**	**0.53**	0.04
Bio2_mean Mean Diurnal Range	-0.14	0.19	**0.81**
Bio2_stdev	-0.30	**0.71**	0.43
Bio3_mean Isothermality	**0.89**	-0.02	-0.22
Bio3_stdev	-0.02	0.87	0.10
Bio4_mean Temperature Seasonality	**-0.96**	0.02	0.13
Bio4_stdev	**-0.78**	0.50	0.14
Bio7_mean Temperature Annual Range	**-0.90**	0.08	0.34
Bio7_stdev	**-0.69**	**0.61**	0.22
Bio12_mean Annual Precipitation	**0.56**	-0.01	**-0.73**
Bio12_stdev	0.28	0.65	-0.44
Bio15_mean Precipitation Seasonality	0.01	0.17	**0.80**
Bio15_stdev	-0.11	0.74	0.39
SS loadings	5.32	3.22	2.63
Proportion Variance	0.38	0.23	0.19
Cumulative Variance	0.38	0.61	0.80
Proportion Explained	0.48	0.29	0.24
Cumulative Proportion	0.48	0.76	1.00

702   **a)** Food items PCA. **b)** Climatic PCA.

705   **S5 Table. Variance inflation factor (VIF) analyses of all parameters included in the combined model.**

Predictor	VIF
Mass	1.188
Development mode (precocial vs altricial)	1.162
Clutch size	1.219
Residual egg mass	1.273
Time fed	1.906
Number of caretakers	1.167
Foraging substrate (others vs arboreal)	1.075
Migratory behavior (sedentary vs migratory)	1.352
Energy content of food (PC)	1.095
Food handling level (PC)	1.135
Insularity (continental vs insular)	1.564
Mean diurnal temperature range (PC)	1.600
Annual temperature and rainfall (PC)	2.076
Climatic seasonality (PC)	1.108
Social bonds	1.567
Colonial vs solitary breeding	1.031
Grouping pattern	1.073

708

Variance inflation factors (VIF) were calculated from a Bayesian phylogenetically controlled mixed model including parental provisioning and eco-social predictors. VIF is assumed to be the term

711   inflating the regression coefficient's variance estimate $\widehat{var}(\beta_i) = \left(\frac{s^2}{(n-1)\widehat{var}(X_i)}\right)\frac{1}{1-R_i^2} =$

$\left(\frac{s^2}{(n-1)\widehat{var}(X_i)}\right)VIF_i$, where s^2 is the root mean squared error of regression, n is the sample size, X_i is

the i-th column of the design matrix. R_i^2 then becomes the GLMM R^2 analogue for the model: $X_i =$

714 $\alpha_0 + \alpha_1 X_1 + \cdots + \alpha_{i-1} X_{i-1} + \alpha_{i+1} X_{i+1} + \cdots + \alpha_k X_k.$

S6 Table. Model comparisons of the models included in the phylogenetic d-separation path analyses.

	Model	C	CICc	ΔCICc		Model	C	CICc	ΔCICc
a, Parental	Model_A1	54.22	92.88	0.00	b, Eco-social	Model_E1	222.73	282.24	0.00
provisioning	Model_A2	62.77	99.36	6.48	model	Model_E2	238.15	291.37	9.13
model	Model_A3	62.07	100.72	7.85		Model_E3	241.22	296.54	14.29
	Model_A4	72.04	110.70	17.82		Model_E4	242.74	298.06	15.82
	Model_A5	79.89	118.54	25.67		Model_E5	242.16	299.57	17.33
	Model_A6	87.46	124.05	31.17		Model_E6	249.66	302.88	20.63
	Model_A7	91.65	128.24	35.36		Model_E7	254.88	303.92	21.67
	Model_A8	99.22	133.75	40.87		Model_E8	249.07	304.39	22.14
	Model_A9	116.06	150.59	57.71		Model_E9	258.17	307.21	24.96
	Model_A10	119.17	151.64	58.77		Model_E10	257.58	308.71	26.46
	Model_A11	115.36	151.95	59.07		Model_E11	255.57	308.79	26.54
	Model_A12	118.47	153.00	60.12		Model_E12	260.31	309.35	27.10
	Model_A13	134.86	169.39	76.51		Model_E13	265.08	312.04	29.79
	Model_A14	133.18	169.77	76.89		Model_E14	261.24	312.37	30.12
	Model_A15	134.15	170.74	77.87		Model_E15	264.49	313.54	31.29
	Model_A16	136.29	170.82	77.94		Model_E16	260.65	313.87	31.62
	Model_A17	141.97	174.44	81.56		Model_E17	268.15	317.19	34.95
	Model_A18	141.26	175.79	82.92		Model_E18	267.56	318.69	36.45
	Model_A19	141.72	176.25	83.37		Model_E19	274.89	319.77	37.53
	Model_A20	144.94	179.47	86.59		Model_E20	274.31	321.26	39.02
	Model_A21	148.05	180.52	87.64		Model_E21	272.62	323.75	41.50
	Model_A22	152.56	185.03	92.15		Model_E22	281.81	324.61	42.36
	Model_A23	151.97	188.57	95.69		Model_E23	275.81	324.85	42.60
	Model_A24	154.53	189.06	96.18		Model_E24	280.54	325.42	43.18
	Model_A25	160.02	192.49	99.62		Model_E25	281.22	326.10	43.85
	Model_A26	159.09	193.61	100.74		Model_E26	275.00	326.13	43.88
	Model_A27	159.54	194.07	101.20		Model_E27	279.73	326.69	44.45
	Model_A28	160.00	194.53	101.65		Model_E28	282.72	329.68	47.43
	Model_A29	163.73	198.26	105.39		Model_E29	287.46	330.26	48.01
	Model_A30	170.85	203.32	110.44		Model_E30	281.91	330.95	48.71
	Model_A31	171.30	203.77	110.90		Model_E31	286.65	331.53	49.28
	Model_A32	172.36	206.88	114.01		Model_E32	292.64	339.59	57.35
	Model_A33	177.82	212.35	119.48		Model_E33	292.05	341.09	58.84
	Model_A34	181.43	213.90	121.03		Model_E34	299.55	344.43	62.18
	Model_A35	187.43	217.84	124.96		Model_E35	298.96	345.92	63.67
	Model_A36	187.41	219.87	127.00		Model_E36	308.51	351.31	69.06
	Model_A37	187.41	219.87	127.00		Model_E37	316.24	354.90	72.65
	Model_A38	188.15	220.62	127.74	c, Combined	Model_C1	422.63	544.76	0.00
	Model_A39	188.90	221.37	128.49	model	Model_C2	420.92	545.26	0.50
	Model_A40	191.26	221.67	128.80		Model_C3	430.92	548.62	3.87
	Model_A41	187.44	221.97	129.09		Model_C4	429.21	549.12	4.36
	Model_A42	190.56	223.02	130.15		Model_C5	429.68	549.59	4.83
	Model_A43	205.23	237.70	144.82		Model_C6	433.82	551.52	6.76
	Model_A44	205.23	237.70	144.82		Model_C7	433.20	553.11	8.35
	Model_A45	205.26	239.79	146.92		Model_C8	444.99	560.49	15.73
	Model_A46	208.38	240.85	147.97		Model_C9	471.68	574.03	29.27
						Model_C10	470.11	576.82	32.06
						Model_C11	933.65	991.06	446.31

720   Models are sorted by their CICc values. Parental provisioning models with a ΔCICc >150 are not

       shown (N=21 models). C: Fisher's C-statistics. The topologies of the top models (N=15 for parental

       provisioning and eco-social model, N=10 for combined model) are illustrated in S2-4 Figs.