Depletion of *ALMS1* affects TGF-β signalling pathway and

downstream processes such as cell migration and

adhesion capacity.

Brais Bea-Mascato[1,2], Elena Neira-Goyanes[1,2], Antía Iglesias-Rodríguez[1,2], Diana

Valverde[1,2]*

1 CINBIO, Universidad de Vigo, 36310, Spain.

2 Grupo de Investigación en Enfermedades Raras y Medicina Pediátrica, Instituto de

Investigación Sanitaria Galicia Sur (IIS Galicia Sur), SERGAS-UVIGO, Vigo, Spain

*Correspondence: Diana Valverde

CINBIO Facultad de Biología, Universidad de Vigo, Campus As Lagoas-Marcosende s/n,

36310 Vigo, Spain

Tel +34 986 811 953

Email dianaval@uvigo.es

ABSTRACT:

Background:

ALMS1 is a ubiquitous gene associated with Alström syndrome (ALMS). The main symptoms of ALMS affect multiple organs and tissues, generating at last, multi-organic fibrosis in the lungs, kidneys and liver. TGF-β is one of the main pathways implicated in fibrosis, controlling the cell cycle, apoptosis, cell migration, cell adhesion and epithelial-mesenchymal transition (EMT). Nevertheless, the role of *ALMS1* gene in fibrosis generation and other implicated processes such as cell migration or cell adhesion via the TGF- β pathway has not been elucidated yet.

Methods:

Initially, we evaluated how depletion of *ALMS1* affects different processes like apoptosis, cell cycle and mitochondrial activity in HeLa cells. Then, we performed proteomic profiling with TGF-β stimuli in HeLa ALMS1 -/- cells and validated the results by examining different EMT biomarkers using qPCR. The expression of these EMT biomarkers were also studied in hTERT-BJ-5ta ALMS1 -/-. Finally, we also evaluated the SMAD3 and SMAD2 phosphorylation and cell migration capacity in both models.

Results:

Depletion of *ALMS1* generated apoptosis resistance to thapsigargin (THAP) and C2-Ceramide (C2-C), and G2/M cell cycle arrest in HeLa cells. For mitochondrial activity, results did not show significant differences between *ALMS1 +/+* and *ALMS1 -/-*. Proteomic results showed inhibition of downstream pathways regulated by TGF-β. The protein-coding genes (PCG) were associated with processes like focal adhesion or cell-substrate adherens junction in HeLa. *SNAI1* showed an opposite pattern to what would be expected when activating the EMT in HeLa and BJ-5ta. Finally, in BJ-5ta model a reduced activation of SMAD3 but not SMAD2 were also observed. In HeLa models no alterations in the canonical TGF-β pathway were observed but both cell lines showed a reduction in migration capacity.

Conclusion:

ALMS1 has a role in controlling the cell cycle and the apoptosis processes. Moreover, the depletion of *ALMS1* affects the signal transduction through the TGF-β and other processes like the cell migration and adhesion capacity.

Keywords: ALMS1, Alström syndrome, ciliopathies, primary cilia, TGF-β, epithelial-mesenchymal transition (EMT), fibrosis, cell cycle arrest, cell migration, apoptosis.

INTRODUCTION:

ALMS1 is a ubiquitous gene with 23 exons located on chromosome 2p13. Mutations in this gene are related to Alström syndrome (ALMS; OMIM #203800), a rare, autosomal and recessive disease with a prevalence of 1-9 cases per 1.000.000 inhabitants and approximately 950 cases reported worldwide (Orphanet). The main symptoms of ALMS affect multiple organs and tissues [1], such as cone-rod retinal dystrophy, dilated cardiomyopathy (DCM) and hypertension [2], hypertriglyceridemia, type II diabetes mellitus (T2D) [3], and hepatic, renal and pulmonary dysfunction caused by multi-organic fibrosis [1], [4], which is the main cause of death in these patients.

The protein encoded by *ALMS1* plays a structural role in the eukaryotic cell centrosomes [5], a microtubule (MT)-nucleating organelle strongly related to primary cilia. The ALMS1 basal body location [5], [6] has been related to different functions controlled by the primary cilium such as centrosome cohesion, cell cycle control and endosomal trafficking [6]–[9].

Although numerous roles of ALMS1 have been described at both ciliary and extraciliary levels, the regulation of the pathways underlying these cellular processes remain unknown. Recently, our group has established the relationship between the *ALMS1* gene and the TGF-β signaling pathway [10], which could be related to the development of fibrosis in these patients [11], [12]. TGF-β is a signalling pathway regulated by clathrin-dependent endocytosis (CDE) in the pocket region of primary cilium [13], and coordinates cell cycle, apoptosis and migration processes such as epithelium-mesenchyme transition (EMT) [14]. This pathway acts in close coordination with other signalling cascades such as Wnt, Hippo, Notch, Hedgehog (Hh), mitogen-activated protein kinase (MAPK), and phosphoinositide 3-kinase (PI3K)-Akt, resulting in many cases in complex regulatory networks dependent on the primary cilium [15].

Stimulation of the TGF-β pathway can activate two main cascades: the canonical (SMAD-dependent) pathway and the non-canonical (SMAD-independent) pathway [16]. The canonical pathway is coordinated by transcription factors called R-SMAD (SMAD2 and SMAD3) which join co-SMAD (SMAD4) to activate the expression of several genes such as *SNAI1* [17], [18], a key modulator of EMT [19], [20]. This pathway is inhibited by an I-SMAD, SMAD7, which inhibits R-SMAD signalling in a variety of ways, binding to TGF-β receptors or directly in the nucleus, avoiding the binding of the R-Smad-Smad4 complex to DNA [21]. On the other hand, the non-canonical activation pathway includes a wide group of pathways such as p38 MAPK, PI3K/AKT, or Rho GTPases [16]. Recently, studies have linked the abolition of several centrosomal proteins such as ALMS1 and CEP128 to the inhibition of the SMAD-dependent canonical pathway in

zebrafish and various human cell types such as hTERT-RPE1 and ciliated human foreskin
fibroblasts (hFF) [10], [22]. However, deciphering how centrosomal proteins coordinate and
regulate the TGF-β signalling pathway and their downstream processes is a challenge.

Understanding how the absence of *ALMS1* influence cell migration and adhesion processes such
as EMT through the TGF-β pathway is key to understanding how the multi-organ fibrosis occurs
in these patients.

In this article, we analysed how the lack of *ALMS1* could affect different cell processes such as
mitochondrial activity, cell cycle, apoptosis, and signal transduction through the TGF-β pathway.
Moreover, we evaluated the expression pattern for different EMT biomarkers after TGF-β
pathway stimulation in two independent CRISPR-KO cell models (HeLa and BJ-5TA) for *ALMS1*.
Finally, we measured p-SMAD3 and p-SMAD2 activation and migration capacity in in the
presence and absence of the TGF-β1 ligand.

MATERIALS AND METHODS:

Cell culture:

HeLa human cervical cancer cell line from American Type Culture Collection (ATCC), hTERT BJ-
5ta human dermal fibroblasts (ATCC) and 293T human embryonic kidney cell line from ATCC
were used in this study. HeLa and 293T cells were maintained in Dulbecco's minimum essential
medium (DMEM, Gibco, Invitrogen, NY, USA), supplemented with 10% fetal bovine serum (FBS)
(Gibco, Invitrogen, NY, USA) and 2% penicillin/streptomycin (P/S) (Gibco, Invitrogen, NY, USA).
In the case of BJ-5ta line, cells were maintained in DMEM:199, supplemented with 10% of FBS,
2% P/S and Hygromycin B (0.01mg/mL). All cell lines were grown at 37°C with 5% CO_2 for up to
one month. After this, the cells were discarded, and a new culture was started.

Crispr assay and edition validation:

HeLa cells were edited with a dual episomal system for homology directed repair (HDR) (Santa

Cruz Biotechnology, Inc., Dallas, USA). Cells were transfected with Lipofectamine 2000 (Thermo

Fisher, Waltham, EE. UU) with 3 gRNAs directed to *ALMS1* exons 1 and 3 (Supplementary Table

S1) following the manufacture´s protocol. After plasmid transfection, cells were selected with

puromycin (1 µg/mL) for 5 days.

Supplementary Table S1. gRNAs of the constructs used for gene editing. The forward and

reverse sequences are shown in the table. The red colour represents the base pairs added for

ligation in the plasmid.

ID	Cell line	Exon	Forward primer (gRNA sequence 5' - 3')	Reverse primer (5' - 3')	Strand	PAM	On-target Score	Off-Target Score
gRNA 1 HeLa	HeLa	1	GCCCCCGCAGCACCGCTACT	AGTAGCGGTGCTGCGGGGGC	+	CGG	51,4	89,3
gRNA 2 HeLa	HeLa	1	TCCGCTCGCCCTCCGAGTAG	CTACTCGGAGGGCGAGCGGA	-	CGG	56	91
gRNA 3 HeLa	HeLa	3	TTCTCACATTAAACCTAGTC	GACTAGGTTTAATGTGAGAA	-	TGG	29,1	76,7
gRNA 1 BJ	hTERT-BJ-5ta	3	CACCGCCTAGTCTGGGATGTATCCA	aaacTGGATACATCCCAGACTAGGCC	-	AGG	56,2	77,1
gRNA 2 BJ	hTERT-BJ-5ta	3	CACCGAACAGAAGAGATCTCTGTT	aaacAACAGAGATCTCTTCTGTTC	-	GGG	63,3	58,1

BJ-5ta cells were edited using a lentivirus system according to the protocol described by Sanjana

et al [23], with some modifications. Check supplemental material for more information. After

lentiviral transduction, cells were selected with puromycin (1µg/mL) for 5 days.

Surviving cells were expanded and isolated to obtain individual clones. Of the isolation was

performed using Fluorescence-Activated Cell Sorting (FACS) with Hoechst (BD Bioscience, San

José, USA) in the flow cytometer FACS ARIA III (BD Bioscience, San José, USA). Inhibition of gene

expression was validated by qPCR in a StepOnePlus instrument (Thermo Fisher, Waltham, USA).

A commercial probe TaqMan® Gene Expression Assays (Hs00367316_m1) and PrimeTime® qPCR

Primer Assays (Hs.PT.56a.488948) were used to detect *ALMS1* in the HeLa and BJ-5ta models,

respectively.

Mitochondrial activity assay:

Mitochondrial activity was evaluated using Presto Blue reagent (Thermo Fisher, Waltham, USA). Wild type (*ALMS1 +/+*) and Hela Knockout cells (*ALMS1 -/-*) were seeded in 96-well plates by triplicate in concentrations from 5×10^4 to 1.5×10^3 cells per well. The plates were incubated for 24 hours as previously described. After incubation, 10 µL of Presto Blue were added to each well and cells were incubated for 2 hours. Next, the plate was centrifuged at 1500 rpm for 5 min and the supernatant was transferred to a fluorescence plate (Corning, NY, EE. UU). Finally, the plates were measured in the Envision plate reader (Perkin-Elmer, Waltham, USA). This experiment was repeated 3 times on different days.

Apoptosis assay:

The apoptosis assay was performed in flat-bottom 96-well plates. HeLa *ALMS1 +/+* and *ALMS1 -/-* cells were seeded in triplicate at a concentration of 3×10^4 cells/per well and incubated 24 hours in DMEM 2%P/S and 2%FBS before treatment. Next, apoptotic stimuli were added at final concentration of 100 nM for thapsigargin (THAP) and 25µM for C2-ceramide (C2-C) in plates, followed by incubation at 37ºC for 24-48 hours respectively. After that, cells were washed twice with 100 µL PBS. 50 µL trypsin was added per well and the plate was incubated for 5 min at 37°C. After that, trypsin was inactivated with 50 µL DMEM 10% FBS 2% P/S, and cells were transferred to 96-well plates with U bottom. The plate was centrifugated at 1500 rpm for 5 min, the supernatant was discarded, and the cells were resuspended in 200 µL biding buffer with 2µL 7-AAD and 2µL annexin V-FITC (Biolegend Inc, San Diego, CA, USA). The plate was incubated for 15 min in darkness and measured in the flow cytometer FC 500 (Beckman-Coulter, Brea, CA, USA). This experiment was repeated 3 times on different days.

Cell cycle assay:

To determine cell cycle alterations, we studied sub-populations in the HeLa *ALMS1 -/-* model with Hoechst staining (BD Bioscience, San José, USA). *ALMS1 +/+* and *ALMS1 -/-* cells were seeded in a 6-well plate at a concentration of 3×10^5 cells/per well and incubated overnight. The next day, 1µL Hoechst staining was added to the medium and the plate was incubated for 1 hour. Then, the medium was removed, cells were washed twice with 1 mL PBS per well, and trypsinised following the protocol described above. Cells were resuspended in PBS and analysed in the flow cytometer FACS ARIA III.

Protein extraction for proteomic profiling:

HeLa cells were seeded in 6-well plates by triplicate in DMEM 2% P/S, 2%FBS and incubated overnight for serum starvation. One the next day, cells were stimulated with rhTGF-β1 (2ng/mL; R&D Systems; 240-B) for 24 hours. After that, DMEM was removed, and wells were washed twice with PBS. Cells were scrapped in PBS. Then, cells were pelleted at 11000 rpm for 5 min, and 100 μL of RIPA Buffer were added to each sample. Samples were incubated on ice for 20 min in constant agitation. Finally, cell debris was pelleted by centrifugation for 30 min at 12000 rpm and 4 °C. Samples were aliquoted and stored at -80 °C until the next protocol's step. Protein quantification was performed by Bradford microplate assay (Bio-Rad, Hercules, USA).

Protein concentration, reduction, and alkylation:

Initially, 20 μg of each sample was concentrated and stained with Coomassie in a 12% SDS-PAGE gel. Bands were cut and faded. Band distinction was performed by adding 100 μL of ammonium bicarbonate 25mM/50% acetonitrile and incubated 20 min. This step was repeated 4 times. After distinction, the supernatant was removed and 100 μL of ammonium bicarbonate 25mM-50% EtOH was added to samples, and incubated for 20 min. Then, the supernatant was removed and 100 μL EtOH was added and incubated for 15 min. In the next step, 200 μL 10mM DTT in ammonium bicarbonate were added and samples were incubated for 1 hour at 56°C. Then, DTT solution was removed and 200 μL iodoacetamide (IAA) 55mM in 50 mM ammonium bicarbonate was added. Samples were incubated again for 30 min at room temperature. Finally, the supernatant was removed, 200 μL ammonium bicarbonate 50mM were added and samples were incubated for 15 min.

Protein digestion:

After reduction and alkylation, samples were digested. 100 μL ammonium bicarbonate 25mM-50% acetonitrile (ACN) was added to each sample and incubated for 30 min. The supernatant was removed and 100 μL ACN 100% was added and incubated for 10 min. The supernatant was removed again and 15 μL Trypsin in ammonium bicarbonate 25mM were added to each sample. Samples were maintained at 4°C for 30 min. Finally, trypsin was removed, 5 μL ammonium bicarbonate 25 mM were added, and samples were incubated overnight.

Peptide extraction:

40 μL of trifluoroacetic acid (TFA) 0.5% were added to each sample and incubated for 30 min at room temperature. Then, we added 20 μL ACN 100% in each 1.5 mL tube and incubated for 15

188 min at 37°C. Finally, the samples were dried in a speed-vac at 45°C for 1 hour and stored at -

189 20°C until the analysis.

Mass Spectrometry and bioinformatic analysis:

Samples were resuspended in 24 µL formic acid 0.1%. Then, 10 µL were injected into the HPLC

(LC-ESI-MS/MS workflow using a C18 precolumn). Samples were analysed in a mass

spectrophotometry Orbitrap ELITE (Thermo-Fisher, Waltham, USA) in long chromatography

mode (4 hours). After that, protein identification was performed in the database

UniProtKB/TrEMBL for humans, using the Proteome Discoverer software. Downstream analysis

was performed with the R package Differential Enrichment analysis of Proteomics data (DEP

package) [24]. Finally, Go Terms analysis was performed with Clusterprofiler package [25], and

results were visualized with GOplot package [26]. All the code for the analysis was deposited in

the GitHub repository (https://github.com/BreisOne/CRISPR_Hela_KO_Alms1).

The mass spectrometry proteomics data have been deposited in the ProteomeXchange

Consortium via the PRIDE [27] partner repository with the dataset identifier PXD024964.

RNA extraction and RT-PCR:

HeLa cells were seeded in 6-well plates at a concentration of $2x10^5$ by triplicate in DMEM 2%P/S

and 2%FBS and were incubated overnight for serum starvation. The next day, cells were

stimulated with rhTGF-β1 (2 ng/mL) for 24, 48 and 72 hours. DMEM was removed and wells

were washed twice with PBS. Cells were scraped in PBS and collected in a 1.5 mL tubes. For the

RNA extraction, the NYZ total RNA isolation kit (NYZtech, Lisboa, Portugal) was used following

the manufacture´s protocol.

After RNA elution, the sample concentrations were measured with Nanodrop (Thermo Fisher,

Waltham, USA). Next, 300 ng of RNA were retrotranscribed to cDNA using the NZY First-Strand

cDNA Synthesis kit (NYZtech, Lisboa, Portugal) following the manufacture´s protocol.

For the experiment with the BJ-5ta line, the same protocol was followed but using DMEM:199

and stimulating for 24 and 48 hours only.

qPCR expression analysis:

Samples were prepared in 96-well plates (Thermo Fisher, Waltham, EE. UU) adding 1µL of each

primer, 12.5µL of PowerUp SYBR Green Master Mix (Thermo Fisher, Waltham, EE. UU), 1µL of

cDNA and 4.5µL of H_2O_d in a final volume of 20 µL per well. qPCRs were performed in

StepOnePlus instrument using the following, cycling conditions: 10 min at 95°C, 40 cycles of 15

s at 95°C and 1 min at 60°C.Primers were synthesized on demand (IDT) (Supplementary Table

S2). Samples were analysed by technical triplicates for each sample and gene. Cts for technical

triplicates were averaged using the geometric mean and ΔΔCT method was applied for the

normalization, using 2 housekeeping (HK) genes (*ALAS1* and *YWHAZ*).

Supplementary Table S2. Sequences of the primers used for the evaluation of qPCR expression

of epithelial-mesenchymal transition markers and genotyping of the Knockout lines.

Gene symbol	Forward primer	Reverse primer
POSTN	CACTCTTTGCTCCCACCAAT	GCCACTTTGTCTCCCATGAT
CDH2	CCTCCAGAGTTTACTGCCATGAC	GTAGGATCTCCGCCACTGATTC
ACTA2	CTATGCCTCTGGACGCACAACT	CAGATCCAGACGCATGATGGCA
SNAI1	GCTGCAGGACTCTAATCCAGAGTT	GACAGAGTCCCAGATGAGCATTG
TWIST1	GCCAGGTACATCGACTTCCTCT	TCCATCCTCCAGACCGAGAAGG
EDIL3	GCGAATGGAACTTCTTGGCTGTG	GAGCGTTCTGAAGATGCTGGAG
VIM	AGGCAAAGCAGGAGTCCACTGA	ATCTGGCGTTCCAGGGACTCAT
DSP	TGACAGACCGCTGGCAAAGGAT	GGCGTTTAGCATCATAGAGCCAC
ALMS1 (Exon1)	GCACTGCGCCTAAGCTG	gttgggggtggaggctg
ALMS1 (Exon3)	cagttaatgacttagcatgtttttcct	ctttcccctttttgagttaagga
ALMS1 (Exon 1 -3)	GAGCGAGACACCAACATGGA	ATTGCGTCAATATGCCCTCCT

Wound healing assay:

BJ-5ta and HeLa (WT and KO) were seeded in a 24-well plate (6 wells per cell line and genotype)

at a concentration of $1x10^5$ and cultured until a 90-95% of confluence. Cells were serum-

starvated (2%FBS for HeLa cells) for 24 hours. After that, rh-TGF-β1 (2 ng/mL) was added to half

of the wells and a wound performed with a 200 μL pipette tip. Cells were incubated at 37°C 5%

CO_2 and images were taken at 0 and 24 hours with the microscope inverted Nikon TiU. Images

were analysed with the imageJ program [28] and the plugin developed by Suarez-Armendo et al

[29]. Finally, the percentage of closure was measured with respect to the wound area at time 0.

SMADs activation assays:

For p-SMADs/SMADs activation assays, BJ-5ta and HeLa (WT and KO) cells were plated in 100mm

dishes (Corning, NY, USA) and incubated in DMEM:199 2P/S% without FBS and DMEM 2% P/S,

2%FBS for 24 hour, respectively. Then, TGF-β pathway were stimulated adding rhTGF-β1 ligand

at 2 ng/mL (final concentration in dish) for 0, 10, 30, and 90 min in BJ-5ta and 0 and 30 min in

HeLa. After stimulation, cells were washed 3 times with PBS, harvested using a scraper, and

centrifuged in a Sigma® 1–14 K at 4°C, 7,400×g for 10 min. Then, pellets were lysed on ice for 10

min in 200 μL of RIPA lysis buffer containing 1 mM sodium orthovanadate as phosphatase

inhibitor (Sigma-Aldrich, Misuri, USA) and 0.1% (v/v) protease inhibitor cocktail (Merck,
Darmstadt, Germany). Lysates were centrifuged at 4°C, 14,500×g for 30 min, and the
supernatant was collected in 1.5mL low binding tubes (Thermo Fisher, Waltham, EE. UU). The
remaining pellet was discarded. Quantification was performed using Bradford assay in
microplates with Bio-Rad protein assay and samples were stored at -80°C until analysis.

SDS-PAGE, Western Blotting, and Quantification:

A total amount of 20µg from each sample were mixed with 6,25µL of Laemmli Buffer 4X (Biorad,
Hércules, EE.UU), 1,25µL β-mercaptoethanol (BME) and a volume of H_2O_d up to a final volume
of 25µL. Then, the mixes were boiled at 97ºC for 5 min and loaded in a Mini-PROTEAN® TGX™
Precast Gel (Biorad, Hércules, EE.UU) at 12% acrylamide/bis-acrylamide. The gel was migrated
for 1 hour at 150V. Then, gels were transferred to a 0.2 µm PVDF membrane using the Trans-
Blot Turbo transfer system (Biorad, Hércules, EE.UU) following the mixed molecular weight
(MW) protocol. After transfer, membranes were blocked with in-house tris-buffered saline, 0.1%
Tween20 and 5% milk (TBS-T 5%) for 90 min. Incubation with the primary antibody was kept at
4°C overnight. The primary antibodies used in this assay were: anti-SMAD3 (1:1000, Abcam,
40854), anti-p-SMAD3 (1:1000, Abcam, 63403), anti-SMAD2 (1:1000, Abcam, 40855), anti p-
SMAD2 (1:1000, Cell Signaling Technology, 3101).

Next day, incubation with secondary antibody was performed by 45min in TBST 5%, after
washed the membrane 3 times in TBST 5%. The secondary antibody used in this assay was goat
anti-rabbit IgG H&L (HRP) (Abcam, 205718) at 1:5000 for p-SMAD2 and p-SMAD3 and 1:15000
for SMAD2 and SMAD3. Finally, for protein visualization Clarity western ECL substrate (Biorad,
Hércules, EE.UU) was used and membranes were exposed in the ChemiDoc system (Biorad,
Hércules, EE.UU). Quantification of protein bands was carried on Image Lab software (Biorad,
Hércules, EE.UU) using the total protein standardisation method.

Western Blotting membranes were stripped and re-probed using Restore ™ PLUS Western Blot
Stripping Buffer (Thermo Fisher, Waltham, EE. UU) according to the manufacturer's protocol.
Incubation for 15 min in the buffer was followed by 3 washes with 1X TBS. Then, the membranes
were re-blocked and re-labelled following the previously described protocol.

Statistical analysis:

On one hand, mitochondrial activity, apoptosis, cell cycle, qPCR, Western Blot and wound
healing data visualization and statistical analysis were performed with the software GraphPad
Prism 8.00.

Mitochondrial activity was analysed using a linear regression model. Comparison of fluorescence
increase with cell concentration did not show significant differences between both slopes (α =
0.05). The apoptosis experiment results were analysed with 2ways-Anova to compare each
different population between genotypes. A Sidak's multiple comparisons test was applied to
correct the analysis. Finally, cell cycle results were analysed with a two-tailed unpaired t-test for
each population (G2/M, S and G0/G1).

On the other hand, proteomic data analysis included different statistical analyses and data
transformations along the pipeline. Initially, the data were normalised by variance stabilizing
transformation (VSN). The data imputation for missing values was performed like missing not at
random (MNAR) using the Minprob method with FDR < 0.01. Finally, differential expression
analysis was performed using protein-wise linear models combined with empirical Bayes
statistics implemented in the limma package, FDR < 0.05.

A third analysis was performed for the qPCR data. After normalization, fold changes (FC) were
calculated concerning the geometric mean of the expression of each gene in the biological
triplicates for *ALMS1* +/+ at 24h. Significant differences were calculates using a multiple t-test
with a significant threshold of FDR < 0.05.

Finally western-blot data were normalized using total protein method following the
methodology used by Clement et al. [22], [30] and as modified by Gürtler et al. [31]. After
normalization FCs were calculated respect WT at 30 min of TGF-β1 stimulation. Significant
differences were calculates using 2-way-ANOVA with a significant threshold of adj. p-value (FDR)
< 0.05. For wound healing assay, a 2-way-ANOVA was applied and a significant threshold of adj.
p-value (FDR) <0.05 was set.

RESULTS:

ALMS1 depletion does not alter mitochondrial basal activity despite
generating G2/M cell cycle arrest in HeLa cells:

ALMS1 is a structural centromere protein [5] that controls the structure of the cytoskeleton [32].
Since *ALMS1* depletion causes cell cycle elongation [8], [33] it is expected that mitochondrial
activity will also be altered, as both processes are dependent on each other and regulated by
cytoskeletal dynamics. [34].

We evaluated the different phases of the cell cycle using the fox synchronous algorithm [35] to
determine the Gaussian distribution of the different populations. DAPI fluorescence (461 nm)

and events count (cells) were displayed in the plots (Figure 1.A). After 24h of incubation, HeLa

cells ALMS1 -/- shown a significantly reduction (p-value < 0.05) in S phase from 24.26 % ± 1.32

in *ALMS1 +/+* to 19.27 % ± 1.69 in *ALMS1 -/-*. The cell cycle arrest was caused by a significant

increase in the G2/M population (p-value < 0.01), from 1.6% ± 0.62 in *ALMS1 +/+* to 4.88 % ±

0.80 (Figure 1.B). The results in the G0/G1 population did not show significant differences

between genotypes. Despite this, the basal mitochondrial activity did not show a significant

difference between *ALMS1 +/+* and *ALMS1 -/-* (Figure 1.C).

Figure 1. Cell cycle assay and mitochondrial activity results. **(A)** Cytometry plots with DAPI fluorescence and events

count. Colors correspond with each population defined by fox synchronous algorithm. **(B)** Box-plot for each

population previously defined in the cytometry plot that represents the percentage of cell cycle populations from

parent population. **(C)** Linear regression between relative fluorescence units (RFU) and cell concentration in both

genotypes (*ALMS1 +/+* and *ALMS1 -/-*). This experiment was carried out in triplicate and repeated 3 times on different

days (N=3, n=3). *: p-value < 0.05, **: p-value < 0.01.

The lack of *ALMS1* in HeLa prevents apoptosis induced by C2-ceramides

and Thapsigargin:

Accumulation of cells in G2/M phase triggers apoptosis by different pathways such as caspases

[36], [37]. We evaluated whether these cells were sensitive to apoptosis in response to an ER

stress-inducing stimulus (Thapsigargin; THAP) [38], and a pro-apoptotic sphingolipid such as C2-

ceramide.

As in the cell cycle assay, initial isolation of the cell population (P1) of interest was performed.

Plots were represented with the FL4 (7-AAD) fluorescence in the y-axis and FL1 (Annexin V)

fluorescence in the x-axis, both on a logarithmic scale. Thresholds for different populations

(viable, early apoptotic, late apoptotic, and necrotic) were defined by a previous compensation

analysis using THAP-like positive control of cell death. Finally, experiments were performed by

triplicates in the plate (n=3) and in 3 alternative days (N=3).

Genotypes *ALMS1* +/+ and *ALMS1* -/- showed significant differences in viable (44.02% and

69.49%, respectively) and early apoptotic (43,47% and 23,44%) populations in cells stimulated

with C2-C (Figure 2.A-B). Moreover, cells stimulated with THAP also showed significant

differences between genotypes in viable (49,56% and 71,33%), early apoptotic (18,25% and

8,56%) and late apoptotic (25,09% and 11,93%) populations (Figure 2.C-D). These results

indicate a resistance to apoptosis for these stimuli due to the lack of the *ALMS1*.

Figure 2. Apoptosis assay by annexin-V IP. **(A)** Cytometry plots for C2-ceramide (C2-C) with Log FL4 channel (IP) vs Log

FL1 channel (Annexin-v). **(B)** Stacked-bar graph for cells treated with C2-C in a concentration of 25 µM. Percentages

of each population are represented from a parent population previously defined. **(C)** Cytometry plots for thapsigargin

(THAP) with Log FL4 channel (IP) vs Log FL1 channel (Annexin-v). **(D)** Stacked-bar graph for cells treated with THAP in

a concentration of 100nM. Percentages of each population are represented from a parent population previously

defined. This experiment was carried out in triplicate and repeated 3 times on different days (N=3, n=3). *: adj. p-

value < 0.05, **: adj. p-value < 0.01, ***: adj. p-value < 0.001, ****: adj. p-value < 0.0001.

The proteomic profile in HeLa cells reveals *ALMS1* like a regulator of cell adhesion through the TGF-β pathway:

TGF-β is a pathway that plays a key role in the control of cell proliferation and apoptotic processes [14]. Since signal transduction through this pathway occurs via the primary cilium [13], [39], the lack of *ALMS1* could explain the aberrant processes seen above.

We evaluated how *ALMS1* depletion affects the signal transduction through stimulated TGF-β pathway. In this bulk analysis, we obtained a total of 86 differentially expressed proteins (DEP) (Figure 3.A), 31 (36%) over-expressed and 55 (64%) under-expressed. The variation in protein expression in this dataset can be explained mainly by a single principal component (Figure 3B). This would make sense as the main difference between these two genotypes would be the presence or absence of the *ALMS1* gene. From this dataset, a total of 76 proteins were associated with genes (Figure 3.C).

Figure 3. Differential expression analysis for proteomic data in HeLa. **(A)** Heatmap for all proteins (associated and not associated with genes) differentially expressed. Plot shows the expression pattern for the 3 biological replicates of genotype *ALMS1 +/+* (CT) and genotype *ALMS1 -/-* (KO). **(B)** Principal components analysis (PCA) for the top 500 variable proteins with all biological replicates for each genotype. **(C)** Volcano-plot showing the top 40 most significant genes associated with proteins. Proteins with no clear gene association were removed from this plot. Significant thresholds: log2 FC > 1.5, adj p-value < 0.05. This experiment was carried out in triplicate (n=3).

Based on the DE of protein-coding genes (PCG) we performed an enrichment analysis obtaining

a total of 9 GO terms significantly overrepresented (adj. p-value < 0.05) (Figure 4. A-B). The

majority of these PCG were associated with focal adhesion, cell-cell or cell-substrate

interactions. This result suggests the implication of *ALMS1* in the control of adhesion processes.

In addition, we calculated the z-score of these GO terms (Figure 4.A), showed a significant

decrease in process related to the ability of these cells to properly migrate or adhere to each

other or their matrix.

Moreover, genes such as vimentin (*VIM*) and desmoplakin (*DSP*), common biomarkers of EMT,

were associated with these GO terms (Figure 4.B). Interestingly, these genes exhibited an

expression pattern opposite to that observed when EMT process is activated.

Finally, *ITGB5* and *ITGA11*, two integrins whose activation plays an important role in the normal

development of EMT, proliferation and invasion are inhibited [40], [41].

Figure 4. Enrichment analysis based on Go terms for proteomic data. **(A)** GoCircle-plot that includes significant over-

represented GoTerms in a table (ID and Name) and scatter plot with FC of any gene related with each GoTerm.

Moreover, this plot includes Z-score which is an estimation of up-regulation or down-regulation in the Goterm. **(B)**

Circus-Plot applied to Goterms to relate any Gene name with the corresponding GoTerm name. The plot also shows

the FC of each gene, and the list is sorted by FC from the most over-expressed (3) up to the most under-expressed (-

3).

ALMS1 depletion alters the expression pattern of TGF-β1-induced EMT markers in HeLa:

Proteomic analysis showed alterations in several adhesion processes coupled with aberrant

expression of EMT-related genes (*VIM*, *DSP*, *ITGB5* and *ITGA11*). We measure the main EMT

markers in our *ALMS1*-deficiency HeLa cells and compare it with another *ALMS1* model in the

hTERT-BJ-5ta line.

We evaluated the expression of the principal EMT biomarkers (*SNAI1*, *CDH2*, *TWIST1*, *ACTA2*,

VIM, and *DSP*) and other genes related to EMT (*POSTN*, *EDIL3*). We perform stimulations at

different time points (24, 48 and 72 h for HeLa and 24 and 48 h for BJ-5ta) with rh-TGF-β1

(2ng/mL) stimulation.

In HeLa (Figure 5.A) at 24 hours of stimulation *POSTN*, *SNAI1*, *VIM, ACTA2* and *EDIL3* were found

under-expressed, while *TWIST1* and *DSP* were over-expressed. This data confirms the expression

pattern previously detected in the proteome analysis related to the genes *VIM*, *DSP* and *EDIL3*

(Figure 5.B). At 48 hours of stimulation, the over-expressed genes *CDH2*, *TWIST1* and *DSP*, in

addition to *EDIL3* and *VIM* inhibition, showed significant differences (Figure 5.A). No other tested

gene were significant. Finally, at 72 hours of stimulation the expression pattern was like the

previous stimulation time points. However, the data dispersion was too high, and no significant

differences were detected.

Figure 5. Expression measurement by qPCR of different EMT biomarkers with TGF-β stimulation at **(A)** 0, 24, 48, and

72 hours in HeLa. **(B)** Correlation plot between the expression of three EMT markers in the HeLa proteomic analysis

and the HeLa qPCR assay. The expression is represented based on the log2 FC and the R coefficient using Pearson's

method. **(C)** Expression measurement of EMT markers by qPCR in BJ-5ta after 0, 24 and 48 hours of stimulation with

TGF-β1 ligand **(D)** Comparative heatmap with Z-score values per sample of EMT markers in HeLa and BJ-5ta after 24

hours of TGF-β1 stimulation. These experiments were performed by triplicate (n=3). *: adj. p-value < 0.05, **: adj. p-

value < 0.01.

In BJ-5ta (Figure 5.B) no inhibition of *VIM* or overexpression of *DSP* were detected. However,

inhibition of *POSTN* and the transcription factor (TF) *SNAI1* was constant in both cell lines as well

as the overexpression of *TWIST1*. *ACTA2* also showed decreased levels in HeLa but not as

significant as in BJ-5ta. After Z score normalization of log2FC both cell lines showed a similar

expression pattern of these markers (Figure 5C). EDIL3 is not shown in the heatmap as FC is too

high, so it could mask the differences of the rest of the markers.

Absence of *ALMS1* decreases migration capacity in HeLa and BJ-5ta and reduce SMAD3 phosphorylation in BJ-5ta:

We evaluated the canonical activation of the TGF-β pathway. SMAD2 and SMAD3

phosphorilation levels were analysed at 4 different times (0, 10, 30 and 90 min) in BJ-5ta and 2

different times (0 and 30 min) in HeLa after TGF-β1 stimuli. The data shown that the lack of

ALMS1 inhibits SMAD3 (Ser213) activation after 30min of TGF-β1 ligand exposure (p-value

<0.01) but not affects to SMAD2 phosphorylation in BJ-5ta (Figure 6 A-D). In HeLa cells no

alterations in canonical pathway signalling of TGF-β were detected (Figure 6 E, F). The p-

SMADs/SMADs ratios did not differ from the p-SMADs/total protein ratios, data not shown.

Finally, we measured cell migration capacity in both cell lines in the presence and absence of -

rh-TGF-β1 ligand. (Figure 6 G, H). *ALMS1* depletion reduced migration capacity compared to the

control line in both cell lines. This reduction is exacerbated after TGF-β stimulation in BJ-5ta

while in HeLa the tendency is the same in the presence or absence of the stimulus.

Figure 6. TGF-β canonical pathway activation assay and measurement of cell migration capacity in HeLa and BJ-5ta.

(A) Representative western-blot of p-SMAD3 and SMAD3 levels after 0, 10, 30 and 90 min of stimulation with TGF-β1

(2 ng/µL) ligand and their corresponding loading controls (coomassie). **(B)** Quantification of band intensity in WB for

p-SMAD3 and SMAD3 at time 0 and after 30 min stimulation with TGF-β1. (n=3). *: p-value < 0.05, **: p-value < 0.01

(C) Representative western-blot of p-SMAD2 and SMAD2 levels after 0, 10, 30 and 90 min of stimulation with TGF-β1

(2 ng/µL) ligand and their corresponding loading controls (coomassie). **(D)** Quantification of band intensity in WB for

p-SMAD2 and SMAD2 at time 0 and after 30 min stimulation with TGF-β1. (n=3). *: p-value < 0.05, **: p-value < 0.01.

(E) Representative western-blot of p-SMAD3, SMAD3, p-SMAD2 and SMAD2 levels after 0 and 30 min of stimulation

with TGF-β1 (2 ng/µL) ligand and their corresponding loading control (coomassie). **(F)** Quantification of band intensity

in WB for p-SMAD3, SMAD3, p-SMAD2 and SMAD2 at time 0 and after 30 min stimulation with TGF-β1. (n=2). *: p-

value < 0.05, **: p-value < 0.01 **(E)** Representative images of wound healing assay in HeLa (WT and KO) and BJ-5ta

(WT and KO) at 0 and 24 hours in absence or presence of 2 ng/µL TGF-β1. **(F)** Quantification of the cell area of the

image and how it decreased after 24 hours of incubation (n=3). *: p-value < 0.05, **: p-value < 0.01, ***: p-value <

0.001.

DISCUSSION:

In this study, we obtain similar results as previously described for the ALMS1 deficiency such as

cell cycle arrest [33] or resistance to apoptosis [8] in a new cell type. Moreover, we reported

new insights to understand the role of *ALMS1* in cell function and TGF-β signalling

Cell cycle arrest in the G2/M phase (Figure 1. A-B), also described in *ALMS1*-knockdown

cardiomyocytes [33], seems to be a cell process triggered by *ALMS1* depletion. Several studies

have already linked cell cycle arrest in the G2/M phase to the TGF-β pathway and the

development of fibrosis [42]–[44].

Normally, cell cycle arrest in both the G2/M and G0/G1 phases is often accompanied by the

activation of apoptotic processes [34], [45]–[47]. In contrast, depletion of *ALMS1* seems to

generate apoptosis resistance against THAP and C2-C in HeLa cells (Figure 2), which is in

accordance with that reported by Zulato et al., [8] in patient fibroblasts. THAP is a non-

competitive inhibitor for Ca^{2+}-ATPases located in the endoplasmic reticulum (SERCA). This,

inhibits Ca^{2+} reflux to the lumen of the endoplasmic reticulum, generating endoplasmic

reticulum stress (ER stress) [48]. At the same time, ER stress activates a signalling network called

Unfolded Protein Response (UPR) which alleviates this stress and promotes cell survival,

restoring ER homeostasis [49]. At this point, *ALMS1* depletion could be acting at two levels:

inhibiting THAP action (ER stress) or over-activating UPR and generating apoptosis resistance.

Because of the role of *ALMS1* in the centrosome and its relation with endocytic trafficking [32],

its depletion could probably alter the structure of the membrane, inhibiting the signalling

transduction of ligands like THAP. THAP also induces apoptosis impairing cytoskeleton dynamics

and disorganization of F-actin via mTORC1 inhibition in the mTORC1-RhoA-Limk-Cofilin-1 axis in

A549 cells [50]. *ALMS1* depletion already causes disorganization of the F-actin cytoskeleton [32]

but does not lead to cell death. Probably, the mTOR pathway through mTORC1 was inhibited in

ALMS1 -/- cells, but in this case, other mechanisms could prevent cells from reaching apoptotic

stages. This would be supported by the inhibition of LAMTOR3 (Figure 4.B), a protein involved

in the formation of the Ragulator complex that activates mTORC1 [51].

C2-C is a sphingolipid that induces caspases-dependent apoptosis, inhibiting PI3K/AKT pathway

and increasing the depolarisation of the mitochondrial membrane [52]–[54]. The over-activation

of the PI3K/AKT pathway could explain the apoptotic resistance in the case of the C2-C [55], [56].

This fact would agree with the previous proposed mTORC1 inhibition, since phosphorylation of

AKT/PKB at s473 leads to inhibition of mTORC1 mediated by FoxO transcription factors [57],

which could be related to F-actin filament disorganisation previously described in Alström

fibroblasts [32].

In addition to the role of *ALMS1* in G2/M phase cell cycle arrest and resistance to THAP- and C2-

C-mediated apoptosis, this study suggests that *ALMS1* depletion leads to inhibition of TGF-β

signalling pathway. This could be affecting downstream processes such as cell migration and

adhesion.

After TGF-β stimulation, the lack of *ALMS1* alters the expression of several genes related to focal

adhesion and cell-substrate adherens junction (Figure 4 A-B), two processes required for TGF-

β1-induced EMT [58], [59]. This fact could be conditioning the activation of EMT, preventing cells

from moving into a mesenchymal state characterised by an increased migratory capacity.

Previous studies have demonstrated the relationship between signalling through the primary

cilium and the EMT focusing mainly on signalling through the WNT and SHH pathway in various

organs such as epicardial tissue, kidney epithelial cells, and pancreatic β-cells [31]–[34].

However, signalling through the TGF-β pathway has not yet been extensively studied in this

context.

Moreover, we have found interesting candidates in the HeLa proteome profiling that could

explain or be related to the mentioned cellular disturbances (Figure 3C). *S100A6*, under-

expressed, is a positive regulator of proliferation, migration and angiogenesis [60]–[62], and also

a regulator of endothelial calcium signalling [63]. *PKM*, which was over-expressed in HeLa cells,

is associated with apoptosis resistance [64]. *ITGA11*, another gene under-expressed in this cell

line and related to fibrosis generation, is expressed due to activation of the TGF-β pathway [65].

Finally, two known EMT markers, *VIM* and *DSP*, showed a reverse pattern to that expected when

this process is activated. *VIM* is a key player in the regulation of EMT and its deficiency leads to

inhibition of this process [66]. On the other hand, *DSP* is a protein involved in cell-cell junctions

mediated by desmosomes. It is therefore considered a typical epithelial marker whose decrease

is related to the passage of the cell from an epithelial to a mesenchymal state [67]. This could

be first evidence suggesting inhibition of the TGF-β1-induced EMT when *ALMS1* is missing.

One of the most relevant detected genes was *SNAI1,* which is inhibited in both cell lines (HeLa

and BJ-5ta). *SNAI1* and *TWIST1* are transcription factors that repress E-cadherin (*CDH1*)

expression and activate the expression of N-cadherins (*CDH2*) [68], [69]. E/N-cadherins switch is

involved in the induction of EMT [70], [71] through the TGF-β pathway [72]. Inhibition of *SNAI1*

could be affecting the normal EMT in these cells. The decreased expression of *SNAI1,* together

with the inhibition of *ACTA2* and *POSTN,* also suggests that TGF-β1-induced EMT would be

down-regulated. *SNAI1* is up-regulated by SMAD3, which is phosphorylated when the TGF-β

pathway is activated and forms a complex with SMAD2 [17]. The phosphorylated SMAD2-

SMAD3 complex translocates to the nucleus, activating the expression of several genes related

to cell migration and EMT such as *SNAI1* [73]. Because we detected consistent inhibition in both

cell models of *SNAI1*, we sought to determine whether SMAD3 and SMAD2 activation was also

compromised in BJ-5ta and HeLa models. In BJ-5ta, we conclude that *ALMS1* depletion inhibits

SMAD3 phosphorylation but not affect SMAD2 phosphorylation (Figure 6 A-D). This fact could

lead to downstream inhibition of *SNAI1* and affect migration and adhesion capacity in this cell

line (Figure 6 G, H). In HeLa, no alterations in the activation of the canonical TGF-β pathway were

detected (Figure 6 E, F). In addition, a decrease in cell migration capacity was observed, but

independently of TGF-β, in contrast to BJ-5ta (Figure 6 G, H). This could explain that the

involvement of ALMS1 in cell migration is a conserved process, but its regulation is a cell type-

dependent process.

We have previously described that *ALMS1* depletion by siRNA inhibits SMAD2 phosphorylation

in the hTERT RPE-1 cell line [10] but this was not detected in any of our KO models. The only

alteration detected in the canonical pathway was the inhibition of SMAD3 activation in BJ-5ta.

This difference could be due to the cell type or to differences between the KO and siRNA models.

This facts leads us to hypothesise that inhibition of p-SMAD3 could prevent the formation of the

SMAD2/3 complex, affecting the expression of *SNAI1* in BJ-5ta. [20]. Whereas in HeLa, inhibition

of *SNAI1* and cell migration capacity appears to involve other regulatory mechanisms.

CONCLUSION:

ALMS1 appears to have a role in the regulation of cell cycle and apoptotic processes but does not affect mitochondrial reduction capacity in HeLa. *ALMS1* depletion affects signal transduction through the TGF-β pathway, altering the expression of several EMT-related genes in BJ-5ta and HeLa. However, the canonical (SMAD-dependent) pathway did not show major differences. Finally downstream processes such as cell migration and adhesion capacity appear to be compromised in the absence of *ALMS1*.

Author Contributions: BB-M and DV designed the study. BB-M performed the experiments and analysed the data. EN-G assisted in performing apoptosis, cell cycle and mitochondrial activity assays. AI-R assisted in the quantification of the expression of EMT markers in the BJ-5ta model and in the cell migration assay. BB-M and DV wrote the manuscript. All authors have read the draft and provided approval for publication.

Data Availability Statement: Proteomic data are available via ProteomeXchange with identifier PXD024964.

Acknowledgments: We sincerely thank the Proteomics and Genomics services from Centro de Apoyo Científico-Tecnológico a la Investigación (CACTI) of University of Vigo and its specialists Paula Álvarez Chaver, Ángel Sebastián Comesaña, Verónica Outeiriño and Manuel Marcos for their guidance and advise. We also thank Mercedes Peleteiro Olmedo from Centro de Investigacións Biomédicas (CINBIO) from University of Vigo for the flow cytometry service.

Funding: This work was funded by Instituto de Salud Carlos III (project PI15/00049 and PI19/00332), Xunta de Galicia (Centro de Investigación de Galicia CINBIO 2019-2022; Ref. ED431G-2019/06) and Consolidación e estructuración de unidades de investigación competitivas e outras accións de fomento (Xunta de Galicia, ED431C-2018/54). Brais Bea-Mascato (FPU17/01567) was supported by graduate studentship awards (FPU predoctoral fellowship) from the Spanish Ministry of Education, Culture and Sports.

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

BIBLIOGRAPHY:

[1] J. D. Marshall, P. Maffei, G. B. Collin, and J. K. Naggert, "Alström syndrome: genetics and clinical overview," *Curr. Genomics*, vol. 12, no. 3, pp. 225–235, May 2011, doi: 10.2174/138920211795677912.

[2] A. B. Jaykumar, P. Caceres, E. L. Henson, W. H. Beierwaltes, and P. A. Ortiz, "Deletion of ALMS1 (Alstrom Syndrome 1) enhances salt-sensitive hypertension, and induces insulin resistance and obesity in rats.," *FASEB J.*, vol. 31, no. 1_supplement, pp. 857.26-857.26, Apr. 2017, doi: 10.1096/fasebj.31.1_supplement.857.26.

[3] G. B. Collin *et al.*, "Mutations in ALMS1 cause obesity, type 2 diabetes and neurosensory degeneration in Alström syndrome," *Nat. Genet.*, vol. 31, p. 74, Apr. 2002, [Online]. Available: https://doi.org/10.1038/ng867.

[4] J. D. Marshall *et al.*, "Spectrum of ALMS1 variants and evaluation of genotype-phenotype correlations in Alström syndrome," *Hum. Mutat.*, vol. 28, no. 11, pp. 1114–1123, Nov. 2007, doi: 10.1002/humu.20577.

[5] T. Hearn *et al.*, "Subcellular Localization of ALMS1 Supports Involvement of Centrosome and Basal Body Dysfunction in the Pathogenesis of Obesity, Insulin Resistance, and Type 2 Diabetes," *Diabetes*, vol. 54, no. 5, pp. 1581 LP – 1587, May 2005, doi: 10.2337/diabetes.54.5.1581.

[6] D. Jagger *et al.*, "Alström Syndrome protein ALMS1 localizes to basal bodies of cochlear hair cells and regulates cilium-dependent planar cell polarity.," *Hum. Mol. Genet.*, vol. 20, no. 3, pp. 466–481, Feb. 2011, doi: 10.1093/hmg/ddq493.

[7] G. Li *et al.*, "A role for Alström syndrome protein, alms1, in kidney ciliogenesis and cellular quiescence," *PLoS Genet.*, vol. 3, no. 1, pp. e8–e8, Jan. 2007, doi: 10.1371/journal.pgen.0030008.

[8] E. Zulato *et al.*, "ALMS1-deficient fibroblasts over-express extra-cellular matrix components, display cell cycle delay and are resistant to apoptosis," *PLoS One*, vol. 6, no. 4, pp. e19081–e19081, Apr. 2011, doi: 10.1371/journal.pone.0019081.

[9] C. C. Leitch, S. Lodh, V. Prieto-Echagüe, J. L. Badano, and N. A. Zaghloul, "Basal body proteins regulate Notch signaling through endosomal trafficking.," *J. Cell Sci.*, vol. 127, no. Pt 11, pp. 2407–2419, Jun. 2014, doi: 10.1242/jcs.130344.

[10] M. Álvarez-Satta *et al.*, "ALMS1 Regulates TGF-β Signaling and Morphology of Primary
Cilia," *Frontiers in Cell and Developmental Biology*, vol. 9, p. 112, 2021.

[11] A. B. Roberts *et al.*, "Smad3 is key to TGF-β-mediated epithelial-to-mesenchymal
transition, fibrosis, tumor suppression and metastasis," *Cytokine Growth Factor Rev.*,
vol. 17, no. 1–2, pp. 19–27, Feb. 2006, doi: 10.1016/J.CYTOGFR.2005.09.008.

[12] W. KL, J. KE, and H. CA, "Targeting TGF-β Mediated SMAD Signaling for the Prevention
of Fibrosis," *Front. Pharmacol.*, vol. 8, no. JUL, Jul. 2017, doi:
10.3389/FPHAR.2017.00461.

[13] C. A. Clement *et al.*, "TGF-β Signaling Is Associated with Endocytosis at the Pocket
Region of the Primary Cilium," *Cell Rep.*, vol. 3, no. 6, pp. 1806–1814, 2013, doi:
https://doi.org/10.1016/j.celrep.2013.05.020.

[14] Y. Zhang, P. B. Alexander, and X. F. Wang, "TGF-β family signaling in the control of cell
proliferation and survival," *Cold Spring Harb. Perspect. Biol.*, vol. 9, no. 4, Apr. 2017, doi:
10.1101/cshperspect.a022145.

[15] L. K, "Signaling Cross Talk between TGF-β/Smad and Other Signaling Pathways," *Cold
Spring Harb. Perspect. Biol.*, vol. 9, no. 1, Jan. 2017, doi:
10.1101/CSHPERSPECT.A022137.

[16] J. Massagué, "TGFβ signalling in context," *Nature Reviews Molecular Cell Biology*, vol.
13, no. 10. NIH Public Access, pp. 616–630, Oct. 2012, doi: 10.1038/nrm3434.

[17] A. Nakao *et al.*, "TGF-β receptor-mediated signalling through Smad2, Smad3 and
Smad4," *EMBO J.*, vol. 16, no. 17, pp. 5353–5362, Sep. 1997, doi:
10.1093/EMBOJ/16.17.5353.

[18] S. Thuault, E.-J. Tan, H. Peinado, A. Cano, C.-H. Heldin, and A. Moustakas, "HMGA2 and
Smads Co-regulate SNAIL1 Expression during Induction of Epithelial-to-Mesenchymal
Transition," *J. Biol. Chem.*, vol. 283, no. 48, p. 33437, Nov. 2008, doi:
10.1074/JBC.M802016200.

[19] Y. Wang, J. Shi, K. Chai, X. Ying, and B. P. Zhou, "The Role of Snail in EMT and
Tumorigenesis."

[20] T. Vincent *et al.*, "A SNAIL1-SMAD3/4 transcriptional repressor complex promotes TGF-
β mediated epithelial-mesenchymal transition," *Nat. Cell Biol.*, vol. 11, no. 8, pp. 943–
950, 2009, doi: 10.1038/ncb1905.

[21] X. Yan *et al.*, "Smad7 Protein Interacts with Receptor-regulated Smads (R-Smads) to
Inhibit Transforming Growth Factor-β (TGF-β)/Smad Signaling *," *J. Biol. Chem.*, vol.
291, no. 1, pp. 382–392, Jan. 2016, doi: 10.1074/JBC.M115.694281.

[22] M. Mönnich *et al.*, "CEP128 Localizes to the Subdistal Appendages of the Mother
Centriole and Regulates TGF-β/BMP Signaling at the Primary Cilium," *Cell Rep.*, vol. 22,
no. 10, pp. 2584–2592, Mar. 2018, doi: 10.1016/J.CELREP.2018.02.043.

[23] N. E. Sanjana, O. Shalem, and F. Zhang, "Improved vectors and genome-wide libraries
for CRISPR screening," *Nat. Methods*, vol. 11, no. 8, pp. 783–784, 2014, doi:
10.1038/nmeth.3047.

[24] X. Zhang, A. H. Smits, G. B. van Tilburg, H. Ovaa, W. Huber, and M. Vermeulen,
"Proteome-wide identification of ubiquitin interactions using UbIA-MS.," *Nat. Protoc.*,
vol. 13, no. 3, pp. 530–550, Mar. 2018, doi: 10.1038/nprot.2017.147.

[25] G. Yu, L.-G. Wang, Y. Han, and Q.-Y. He, "clusterProfiler: an R package for comparing
biological themes among gene clusters," *OMICS*, vol. 16, no. 5, pp. 284–287, May 2012,
doi: 10.1089/omi.2011.0118.

[26] W. Walter, F. Sánchez-Cabo, and M. Ricote, "GOplot: an R package for visually
combining expression data with functional analysis.," *Bioinformatics*, vol. 31, no. 17,
pp. 2912–2914, Sep. 2015, doi: 10.1093/bioinformatics/btv300.

[27] Y. Perez-Riverol *et al.*, "The PRIDE database and related tools and resources in 2019:
improving support for quantification data.," *Nucleic Acids Res.*, vol. 47, no. D1, pp.
D442–D450, Jan. 2019, doi: 10.1093/nar/gky1106.

[28] C. A. Schneider, W. S. Rasband, and K. W. Eliceiri, "NIH Image to ImageJ: 25 years of
image analysis," *Nat. Methods 2012 97*, vol. 9, no. 7, pp. 671–675, Jun. 2012, doi:
10.1038/nmeth.2089.

[29] A. Suarez-Arnedo, F. T. Figueroa, C. Clavijo, P. Arbeláez, J. C. Cruz, and C. Muñoz-
Camargo, "An image J plugin for the high throughput image analysis of in vitro scratch
wound healing assays," *PLoS One*, vol. 15, no. 7, Jul. 2020, doi:
10.1371/JOURNAL.PONE.0232565.

[30] C. A. Clement *et al.*, "TGF-β Signaling Is Associated with Endocytosis at the Pocket
Region of the Primary Cilium," *Cell Rep.*, vol. 3, no. 6, pp. 1806–1814, Mar. 2013, doi:
10.1016/j.celrep.2013.05.020.

[31] A. Gürtler *et al.*, "Stain-Free technology as a normalization tool in Western blot
analysis," *Anal. Biochem.*, vol. 433, no. 2, pp. 105–111, Feb. 2013, doi:
10.1016/J.AB.2012.10.010.

[32] G. B. Collin *et al.*, "The Alström syndrome protein, ALMS1, interacts with α-actinin and
components of the endosome recycling pathway," *PLoS One*, vol. 7, no. 5, pp. e37925–
e37925, 2012, doi: 10.1371/journal.pone.0037925.

[33] L. T. Shenje *et al.*, "Mutations in Alström protein impair terminal differentiation of
cardiomyocytes," *Nat. Commun.*, vol. 5, p. 3416, Mar. 2014, doi: 10.1038/ncomms4416.

[34] R. Horbay and R. Bilyy, "Mitochondrial dynamics during cell cycling," *Apoptosis*, vol. 21,
no. 12. Springer New York LLC, pp. 1327–1335, Dec. 01, 2016, doi: 10.1007/s10495-016-
1295-5.

[35] M. H. Fox, "A model for the computer analysis of synchronous DNA distributions
obtained by flow cytometry.," *Cytometry*, vol. 1, no. 1, pp. 71–77, Jul. 1980, doi:
10.1002/cyto.990010114.

[36] L. M. Westrate, A. D. Sayfie, D. M. Burgenske, and J. P. MacKeigan, "Persistent
mitochondrial hyperfusion promotes G2/M accumulation and caspase-dependent cell
death," *PLoS One*, vol. 9, no. 3, p. 91911, Mar. 2014, doi:
10.1371/journal.pone.0091911.

[37] D. Arita, M. Kambe, C. Ishioka, and R. Kanamaru, "Induction of p53-independent
apoptosis associated with G2M arrest following DNA damage in human colon cancer
cell lines," *Japanese J. Cancer Res.*, vol. 88, no. 1, pp. 39–43, Jan. 1997, doi:
10.1111/j.1349-7006.1997.tb00299.x.

[38] I. Yoshida, A. Monji, K. Tashiro, K. Nakamura, R. Inoue, and S. Kanba, "Depletion of
intracellular Ca2+ store itself may be a major factor in thapsigargin-induced ER stress
and apoptosis in PC12 cells," *Neurochem. Int.*, vol. 48, no. 8, pp. 696–702, 2006, doi:
https://doi.org/10.1016/j.neuint.2005.12.012.

[39] S. T. Christensen, S. K. Morthorst, J. B. Mogensen, and L. B. Pedersen, "Primary cilia and
coordination of receptor tyrosine kinase (RTK) and transforming growth factor β (TGF-
β) signaling," *Cold Spring Harbor Perspectives in Biology*, vol. 9, no. 6. Cold Spring
Harbor Laboratory Press, p. a028167, Jun. 01, 2017, doi: 10.1101/cshperspect.a028167.

[40] A. Bianchi, M. E. Gervasi, and A. Bakin, "Role of β5-integrin in epithelial-mesenchymal

transition in response to TGF-β," *Cell Cycle*, vol. 9, no. 8, pp. 1647–1659, Apr. 2010, doi: 10.4161/cc.9.8.11517.

[41] P. Erusappan, J. Alam, N. Lu, C. Zeltz, and D. Gullberg, "Integrin α11 cytoplasmic tail is required for FAK activation to initiate 3D cell invasion and ERK-mediated cell proliferation.," *Sci. Rep.*, vol. 9, no. 1, p. 15283, Oct. 2019, doi: 10.1038/s41598-019-51689-6.

[42] L. Yang, T. Y. Besschetnova, C. R. Brooks, J. V Shah, and J. V Bonventre, "Epithelial cell cycle arrest in G2/M mediates kidney fibrosis after injury," *Nat. Med.*, vol. 16, no. 5, pp. 535–543, 2010, doi: 10.1038/nm.2144.

[43] T. L. White, N. Deshpande, V. Kumar, A. G. Gauthier, and U. V Jurkunas, "Cell cycle re-entry and arrest in G2/M phase induces senescence and fibrosis in Fuchs Endothelial Corneal Dystrophy.," *Free Radic. Biol. Med.*, vol. 164, pp. 34–43, Feb. 2021, doi: 10.1016/j.freeradbiomed.2020.12.445.

[44] H. Li *et al.*, "Atg5-mediated autophagy deficiency in proximal tubules promotes cell cycle G2/M arrest and renal fibrosis.," *Autophagy*, vol. 12, no. 9, pp. 1472–1486, Sep. 2016, doi: 10.1080/15548627.2016.1190071.

[45] J. JAKUBIKOVA, Y. BAO, and J. SEDLAK, "Isothiocyanates Induce Cell Cycle Arrest, Apoptosis and Mitochondrial Potential Depolarization in HL-60 and Multidrug-resistant Cell Lines," *Anticancer Res.*, vol. 25, no. 5, 2005.

[46] N. B. Madungwe *et al.*, "Mitochondrial inner membrane protein (Mitofilin) knockdown induces cell death by apoptosis via an AIF-PARP-dependent mechanism and cell cycle arrest," *Am. J. Physiol. - Cell Physiol.*, vol. 315, no. 1, pp. C28–C43, Jul. 2018, doi: 10.1152/ajpcell.00230.2017.

[47] Y. Liu *et al.*, "Calycosin induces mitochondrial-dependent apoptosis and cell cycle arrest, and inhibits cell migration through a ROS-mediated signaling pathway in HepG2 hepatocellular carcinoma cells," *Toxicol. Vitr.*, vol. 70, p. 105052, Feb. 2021, doi: 10.1016/j.tiv.2020.105052.

[48] P. Lindner, S. B. Christensen, P. Nissen, J. V. Møller, and N. Engedal, "Cell death induced by the ER stressor thapsigargin involves death receptor 5, a non-autophagic function of MAP1LC3B, and distinct contributions from unfolded protein response components," *Cell Commun. Signal.*, vol. 18, no. 1, p. 12, 2020, doi: 10.1186/s12964-019-0499-z.

[49]   C. M. Oslowski and F. Urano, "Measuring ER stress and the unfolded protein response
using mammalian tissue culture system," *Methods Enzymol.*, vol. 490, pp. 71–92, 2011,
doi: 10.1016/B978-0-12-385114-7.00004-0.

[50]   F. Wang *et al.*, "Thapsigargin Induces Apoptosis by Impairing Cytoskeleton Dynamics in
Human Lung Adenocarcinoma Cells," *Sci. World J.*, vol. 2014, p. 619050, 2014, doi:
10.1155/2014/619050.

[51]   Z. Mu, L. Wang, W. Deng, J. Wang, and G. Wu, "Structural insight into the Ragulator
complex which anchors mTORC1 to the lysosomal membrane," *Cell Discov.*, vol. 3, no.
1, p. 17049, 2017, doi: 10.1038/celldisc.2017.49.

[52]   M. J. Zinda, C. J. Vlahos, and M. T. Lai, "Ceramide Induces the Dephosphorylation and
Inhibition of Constitutively Activated Akt in PTEN Negative U87MG Cells," *Biochem.*
Biophys. Res. Commun., vol. 280, no. 4, pp. 1107–1115, 2001, doi:
https://doi.org/10.1006/bbrc.2000.4248.

[53]   R. M. Sánchez-Mora, H. Arboleda, and G. Arboleda, "PINK1 Overexpression Protects
Against C2-ceramide-Induced CAD Cell Death Through the PI3K/AKT Pathway," *J. Mol.*
Neurosci., vol. 47, no. 3, pp. 582–594, 2012, doi: 10.1007/s12031-011-9687-z.

[54]   A. S. Arora, B. J. Jones, T. C. Patel, S. F. Bronk, and G. J. Gores, "Ceramide induces
hepatocyte cell death through disruption of mitochondrial function in the rat,"
Hepatology, vol. 25, no. 4, pp. 958–963, Apr. 1997, doi:
https://doi.org/10.1002/hep.510250428.

[55]   S.-J. Jeong *et al.*, "PI3K/AKT inhibition induces caspase-dependent apoptosis in HTLV-1-
transformed cells," *Virology*, vol. 370, no. 2, pp. 264–272, 2008, doi:
https://doi.org/10.1016/j.virol.2007.09.003.

[56]   B. N. Fahy, M. Schlieman, S. Virudachalam, and R. J. Bold, "AKT inhibition is associated
with chemosensitisation in the pancreatic cancer cell line MIA-PaCa-2," *Br. J. Cancer*,
vol. 89, no. 2, pp. 391–397, 2003, doi: 10.1038/sj.bjc.6601037.

[57]   C.-C. Chen *et al.*, "FoxOs Inhibit mTORC1 and Activate Akt by Inducing the Expression of
Sestrin3 and Rictor," *Dev. Cell*, vol. 18, no. 4, pp. 592–604, 2010, doi:
https://doi.org/10.1016/j.devcel.2010.03.008.

[58]   C. Cicchini *et al.*, "TGFβ-induced EMT requires focal adhesion kinase (FAK) signaling,"
Exp. Cell Res., vol. 314, no. 1, pp. 143–152, 2008, doi:

https://doi.org/10.1016/j.yexcr.2007.09.005.

[59]   T. Baronsky *et al.*, "Cell-Substrate Dynamics of the Epithelial-to-Mesenchymal
Transition.," *Nano Lett.*, vol. 17, no. 5, pp. 3320–3326, May 2017, doi:
10.1021/acs.nanolett.7b01558.

[60]   A. Li *et al.*, "S100A6 promotes the proliferation and migration of cervical cancer cells via
the PI3K/Akt signaling pathway.," *Oncol. Lett.*, vol. 15, no. 4, pp. 5685–5693, Apr. 2018,
doi: 10.3892/ol.2018.8018.

[61]   A. Li *et al.*, "S100A6 promotes cell proliferation in human nasopharyngeal carcinoma via
the p38/MAPK signaling pathway.," *Mol. Carcinog.*, vol. 56, no. 3, pp. 972–984, Mar.
2017, doi: 10.1002/mc.22563.

[62]   P. Li, X. Lv, Z. Zhang, and S. Xie, "S100A6/miR193a regulates the proliferation, invasion,
migration and angiogenesis of lung cancer cells through the P53 acetylation.," *Am. J.
Transl. Res.*, vol. 11, no. 8, pp. 4634–4649, 2019.

[63]   B. Haldar *et al.*, "S100A6 is a positive regulator of PPP5C-FKBP51-dependent regulation
of endothelial calcium signaling.," *FASEB J. Off. Publ. Fed. Am. Soc. Exp. Biol.*, vol. 34,
no. 2, pp. 3179–3196, Feb. 2020, doi: 10.1096/fj.201901777R.

[64]   S. Calabretta *et al.*, "Modulation of PKM alternative splicing by PTBP1 promotes
gemcitabine resistance in pancreatic cancer cells.," *Oncogene*, vol. 35, no. 16, pp.
2031–2039, Apr. 2016, doi: 10.1038/onc.2015.270.

[65]   N. Lu, S. Carracedo, J. Ranta, R. Heuchel, R. Soininen, and D. Gullberg, "The human α11
integrin promoter drives fibroblast-restricted expression in vivo and is regulated by
TGF-β1 in a Smad- and Sp1-dependent manner," *Matrix Biol.*, vol. 29, no. 3, pp. 166–
176, Apr. 2010, doi: 10.1016/j.matbio.2009.11.003.

[66]   F. Cheng *et al.*, "Vimentin coordinates fibroblast proliferation and keratinocyte
differentiation in wound healing via TGF-β-Slug signaling," *Proc. Natl. Acad. Sci. U. S. A.*,
vol. 113, no. 30, pp. E4320–E4327, Jul. 2016, doi: 10.1073/pnas.1519197113.

[67]   S. J. Serrano-Gomez, M. Maziveyi, and S. K. Alahari, "Regulation of epithelial-
mesenchymal transition through epigenetic and post-translational modifications," *Mol.
Cancer*, vol. 15, p. 18, Feb. 2016, doi: 10.1186/s12943-016-0502-x.

[68]   W. Jiao, K. Miyazaki, and Y. Kitajima, "Inverse correlation between E-cadherin and Snail
expression in hepatocellular carcinoma cell lines in vitro and in vivo," *Br. J. Cancer*, vol.

86, no. 1, pp. 98–101, 2002, doi: 10.1038/sj.bjc.6600017.

[69] I. Poser, D. Domínguez, A. G. de Herreros, A. Varnai, R. Buettner, and A. K. Bosserhoff,

"Loss of E-cadherin Expression in Melanoma Cells Involves Up-regulation of the

Transcriptional Repressor Snail*," *J. Biol. Chem.*, vol. 276, no. 27, pp. 24661–24666,

2001, doi: https://doi.org/10.1074/jbc.M011224200.

[70] L. Hao, J. R. Ha, P. Kuzel, E. Garcia, and S. Persad, "Cadherin switch from E- to N-

cadherin in melanoma progression is regulated by the PI3K/PTEN pathway through

Twist and Snail," *Br. J. Dermatol.*, vol. 166, no. 6, pp. 1184–1197, Jun. 2012, doi:

https://doi.org/10.1111/j.1365-2133.2012.10824.x.

[71] K. Gravdal, O. J. Halvorsen, S. A. Haukaas, and L. A. Akslen, "A Switch from E-Cadherin

to N-Cadherin Expression Indicates Epithelial to Mesenchymal Transition and Is of

Strong and Independent Importance for the Progress of Prostate Cancer," *Clin. Cancer

Res.*, vol. 13, no. 23, pp. 7003 LP – 7011, Dec. 2007, doi: 10.1158/1078-0432.CCR-07-

1263.

[72] K. Araki *et al.*, "E/N-cadherin switch mediates cancer progression via TGF-β-induced

epithelial-to-mesenchymal transition in extrahepatic cholangiocarcinoma," *Br. J.

Cancer*, vol. 105, no. 12, pp. 1885–1893, 2011, doi: 10.1038/bjc.2011.452.

[73] A. B. Roberts *et al.*, "Smad3 is key to TGF-β-mediated epithelial-to-mesenchymal

transition, fibrosis, tumor suppression and metastasis," *Cytokine Growth Factor Rev.*,

vol. 17, no. 1–2, pp. 19–27, 2006, doi: 10.1016/j.cytogfr.2005.09.008.

Supplementary Material and Methods:

CRISPR plasmids construction:

Single guides RNAs (sgRNAs) for BJ-5ta model were designing using benchling to generate a

deletion of 113pb in the exon 3 of *ALMS1* (ENSG00000116127). The 2 sgRNA (5'-

CCTAGTCTGGGATGTATCCA – 3' and 5'- GAACAGAAGAGATCTCTGTT – 3') result from benchling

were cloned into the plasmid lentiCRISPR v2 [1] following a modified protocol from that

described by Shalem et al [2]. LentiCRISPR v2 was a gift from Feng Zhang (Addgene plasmid

#52961). Initially, 12.5µL (5µg) of LentiCRISPR V2 were digested with 3µL Esp3I (BsmBI), 3µL

FastAP and 6µL 10X FastDigest Buffer from Thermo Fisher. Final volume was set at 60µL. Plasmid

digestion was checked by migration in an agarose gel at 1%.

sgRNAs were phosphorylated and annealed with its correspondent oligo mixing as follow: 1µL
sgRNA (100µM), 1µL of its complement oligo, 2µL of 5X T4 Buffer (Canvax, Cordoba, Spain), 0.5
T4 PNK (NEB, Ipswich, EE.UU) and 5.5 µL distilled H_2O (H_2Od). Mix reaction was incubated at 37ºC
for 30min and 95ºC for 5min. The samples were then left to stand at room temperature for 2
hours and diluted at 1:50, 1:100 and 1:200 ratios.

The unpurified digested plasmid was ligated with the diluted sgRNAs as follows: 0.5µL digested
LentiCRISPR v2, 2µL diluted sgRNA, 1µL T4 DNA Ligase (Canvax, Cordoba, Spain), 4.5µL H_2Od.
The mix was incubated at 37ºC by 30min.

Finally, Stable Competent E.coli (NEB, Ipswich, EE.UU) were transformed by thermal shock
according to the manufactured protocol and incubated at 30ºC overnight. After that, construct
validation was carried out using forward primer for U6 promotor (5'- GACTATCATATGCTTACCGT
-3') and reverse sgRNA.

818 Plasmids Transfection and Lentivirus production:

293T were seeded in a concentration of $3x10^5$ into a 6-well plate and incubated overnight. Then,
293T cells were co-transfected with each CRISPR construct and the plasmids psPAX2 and pCMV-
VSV-G [3] using lipofectamine 3000 (Thermo Fisher, Waltham, EE. UU). pCMV-VSV-G was a gift
from Bob Weinberg (Addgene plasmid #8454) and psPAX2 was a gift from Didier Trono (Addgene
plasmid #12260)

Initially, 7.5µL of lipofectamine 3000 were diluted in 125µL of Opti-MEM medium (Thermo
Fisher, Waltham, EE. UU) by duplicate. Then a total amount of 5µg of ADN (1.6µL for each
plasmid) were diluted at 250µL Opti-MEM. After that, diluted DNA and diluted lipofectamine
were mixed in a 1:1 ratio, centrifugated at 1200rpm and incubated for 20min at room
temperature. Finally, the 250µL of mix were applied to the wells containing the cells.

The cells were incubated with the mixture for 24 hours. Then, the mixture was removed and the
cell was cultured for a further 72 hours for lentivirus production.

831 Lentiviral transduction:

After 72 hours of culturing 293T cells transfected for lentivirus production, the culture medium
was collected in 15 mL tubes and filtered through a 45µm filter (Sartorious,). In addition, a
solution of complete DMEM medium without P/S with polybrene was prepared at a dilution of
1:1000 (8µg/ml).

Then, 300µL lentiviral preparation were mixed with 200µL complete DMEM médium with
polybrene for each well. An additional 1 ml of DMEM with polybrene was added and the plates
were centrifuged at 800g for 30 minutes. This step was repeated one more time. After the
second centrifugation the mix with lentivirus was added again and the plates were incubated
for 24 hours. Finally, mix was removed and cells were allowed to settle for 48 hours in DMEM
médium. They were then selected with puromycin (1ng/mL) for 5 days.

842 Edition validation:

843 After puromycin selection, clonal isolation and expansion we checked the lack of *ALMS1* gene
844 by qPCR in StepOnePlus (Thermo Fisher, Waltham, EE. UU). Finally, we found that these cells
845 had the expected exon 3 deletion by PCR with resolution of samples on a 2% agarose gel and
846 followed by sanger sequencing.

Supplementary figures:

Supplementary Figure S1. Characterisation of *ALMS1* -/- cell line HeLa. **(A)** Design of the model.

(B) Relative gene expression of *ALMS1 +/+* and polyclonal cell line for *ALMS1 -/-* genotype. **(C)**

Relative gene expression of ALMS1 +/+ and a first batch of clones for *ALMS1 -/-* genotype. **(D)**

Relative gene expression of ALMS1 +/+ and second batch of clones for ALMS1 -/- genotype. **(E)**

Strategy for amplification of exons 1 and 3 from genomic DNA in two clones of the *ALMS1 -/-*

(KO) line and the *ALMS1 +/+* control (WT). **(F)** Amplification strategy of the resulting sequence

between exon 1 and 3 after removal of the sequencing between the 2 outermost gRNAs.

Supplementary Figure S2. Genetic characterisation of *ALMS1* -/- cell line BJ-5ta. **(A)** Clone

screening. **(B)** Relative gene expression of ALMS1 +/+ and a batch of clones for ALMS1 -/-

genotype. **(C)** Agarose gel (2%) showing the exon 3 amplicon of the control line (ALMS1 +/+) and

the F4 clone (ALMS1 -/-) which showed the lowest gene expression. **(D)** Characterisation of the

F4 clone mutation by sanger sequencing.

ID	Description
GO:0005743	Mitochondrial inner membrane
GO:0005925	Focal adhesion
GO:0044391	Ribosomal subunit
GO:0005924	Cell−substrate adherens junction
GO:0030055	Cell−substrate junction
GO:0005652	Nuclear lamina
GO:0034399	Nuclear periphery
GO:0097110	Scaffold protein binding
GO:0098631	Cell adhesion mediator activity

A

KO   KH   KO   NTC

KH   KO   NTC

KO: homozygous for the deletion
KH: heterozygous for the deletion

B

ALMS1 mRNA expression

Relative expression (%)

ALMS1 +/+   B9-50   E4-30   B11-50   F4-30   H1-30   B5-30   B10-30   D5-30

C

bp   WT   F4   NTC

Exon 3 *ALMS1*

D

ALMS1 5' 1 2 3 4 5 6 7 8 9 10 ··· 3'

PAM1   gRNA1   PAM2   gRNA2

ALMS1 +/+

AAACCTTGGATACATCCCAGACTAGGTTTAAT   CCCAACAGAGATCTCTTCTGTTCTCCACT

Clone F4

GGACTCTTCCCAAACCTTGG AGAGATCTCTTCTGTTCTCCACTGCT

PAM1   gRNA2

Deletion of 113 pb in exon 3 of ALMS1 gene, generating a stop codon 14 aa downstream

A ALMS1: 23 Exon(s) | 12844 bp | 4168aa | ENST00000613296.6   Exons are depicted on a log2 scale, introns on a log10 scale

CDS Start

CDS End

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23

ALMS1_CDS

300   400   500   600

Exon1   Exon2   Exon3

gRNA1 PAM
PAM gRNA2
PAM gRNA3

gRNA1 Exon 1
5' GTC GCCCCCGCAGCACCGCTACT CGG AGG 3' Strand +

gRNA3 Exon 3
5' CCG TTCTCACATTAAACCTAGTC TGG GAT 3' Strand -

gRNA2 Exon 1
5' AGG TCCGCTCGCCCTCCGAGTAG CGG TGC 3' Strand -

B *ALMS1* mRNA expression

C *ALMS1* mRNA expression

D *ALMS1* mRNA expression

E

F

Exon 1 - 3 *ALMS1*