Title: Assessing the impact of fire on spider abundance through a global comparative

analysis

Running title: The impact of fire on spiders

Claire A. McLean[a, 1], Jane Melville[a], Joseph Schubert[a], Rebecca Rose[a] & Iliana Medina[b]

[a] Sciences Department, Museums Victoria, 11 Nicholson Street, Carlton, VIC 3053, Australia

[b] School of BioSciences, The University of Melbourne, Royal Parade, Parkville, VIC 3010,

Australia

[1] Corresponding author: Claire McLean

Level 3 West, Sciences Department, Museums Victoria, 11 Nicholson Street, Carlton, VIC

3053, Australia, Ph. +61 3 8341 7444

CAM cmclean@museum.vic.gov.au

JM jmelv@museum.vic.gov.au

JS jschubert@museum.vic.gov.au

RR rrose@museum.vic.gov.au

IM iliana.medina@unimelb.edu.au

Statement of authorship: CAM, JM and IM conceived of the research; CAM and RR

performed the literature search and extracted data; CAM and JS compiled the trait data; CAM

and IM designed and performed the comparative analysis; CAM and IM interpreted results;

CAM wrote the first draft of the manuscript, and all authors contributed to revisions.

Keywords: Araneae, burn, phylogenetic signal, time since fire, trait-based approach

Abstract

Fire is an important ecological disturbance; however, in many regions fire regimes are changing due to anthropogenic factors. Understanding the responses of species and ecosystem to fire can help to develop predictive models and inform fire management decisions. Despite their massive diversity and abundance, relatively little is known about the impacts of fire on invertebrates. Spiders are a diverse and ubiquitous group, with variation in ecological, behavioural and life history traits, and can offer important insights into the impacts of fire on invertebrates and whether these depend on environmental factors, phylogenetic history, or species traits. We conducted phylogenetic comparative analyses of published data from studies investigating the impacts of fire on spiders. We investigated whether fire affects family level abundance and whether this effect depends on ecologically relevant traits or site-specific factors (vegetation, latitude, fire type, time since fire). We found that time since fire had a significant effect on the abundance of some spider families (Lycosidae, Linyphiidae, Agelenidae), and within forest vegetation types. There was no relationship between family level traits and response to fire, suggesting that a trait-based approach does not increase predictive power in this group at this taxonomic level. Overall, it is difficult to make broad generalisations about the impacts of fire on spider abundance due to variation in site- and fire-specific factors. We did, however, find evidence that short fire intervals may be a threat to some spider families, and in forest ecosystems, where abundance initially decreased after fire, which may have important implications for fire management strategies. Finally, we show that analyses of published data can be used to detect broad scale ecological patterns and provide an alternative to traditional meta-analytical approaches.

INTRODUCTION

Fire is an important ecological and evolutionary force that has shaped ecosystems worldwide (Bond & Keeley 2005; Pausas & Keeley 2009). While fire plays a role in maintaining species richness and ecological processes in fire-prone regions (reviewed in He et al. 2019), fire can also pose a significant threat to biodiversity and individual species in sensitive landscapes or under different regimes (Bond & Keeley 2005; Shlisky et al. 2009). One of the major consequences of global change has been the alteration of fire regimes; in many regions, wildfires are becoming more frequent and severe, as well as threatening sensitive ecosystems that have long been fire-free (Dowdy et al. 2019; Virgilio et al. 2019; Goss et al. 2020). Understanding the impacts of fire, and fire regimes, is therefore essential for effective conservation and natural resource management now and into the future (Driscoll et al. 2010).

Predictive models quantifying the impacts of fire on species and ecosystems can inform fire management strategies for biodiversity conservation. Such generalisation at broad scales is often difficult due to local variation in species assemblages and environmental factors. However, trait-based approaches, focusing on the properties of an organisms (its traits) rather than species and relating these traits to fire responses, may overcome this challenge (Keith 2012), and an understanding of which traits predict a species response to fire may be especially useful for the management of understudied taxa. Such predictive frameworks have been used to assess the impacts of fire regimes on plant communities (e.g. Menges 2007; Keith 2012; Enright et al. 2014; Clarke et al. 2015; Rodman et al. 2020), and to a lesser extent on animals (e.g. Moretti et al. 2009; Langlands et al. 2011; Driscoll et al. 2019). One constraint of this approach is that closely related species may share similar functional traits (though not always the case; Losos 2008), making it difficult to determine whether observed responses are due to traits or shared ancestry. However, analyses combining both trait and phylogenetic information can provide mechanistic insights into patterns of biodiversity across regions with different fire histories and help to identify species and ecosystems most at risk from changing fire regimes.

Despite their massive diversity and abundance, little is known about the impact of fire on invertebrates (New et al. 2010; Saunders et al. 2021). This is largely because, relative to vertebrate taxa, invertebrates have been poorly studied. It is estimated that around 80% of terrestrial arthropod species are yet to be described (Stork 2018), and fundamental information about their ecology, life history and geographic distribution remains largely

unknown (Cardoso et al. 2011a). Invertebrates play an important role in ecosystem function

and changes to invertebrate assemblages can also have knock-on effects across the food web

(Wurtsbaugh 1992; Benndorf et al. 2000). Therefore, understanding how invertebrates

respond to fire is important at both the species and ecosystem level. Currently, this major

knowledge gap makes it extremely difficult to make informed fire management decisions

concerning invertebrates (Cardoso et al. 2011a; Saunders et al. 2021).

Spiders (order Araneae) are one of the most diverse and successful groups of animals

(Agnarsson et al. 2013) and it is estimated that there are approximately 80,000 species

worldwide (Raven & Yeates 2007; Agnarsson et al. 2013; World Spider Catalog 2021). They

are found in almost all terrestrial ecosystems, on every continent except Antarctica, and are

important invertebrate predators (Nyffeler 2017). Spiders exhibit diverse ecological,

behavioural and life history traits (Pekár et al. 2021), including variation in microhabitat use,

hunting strategies and dispersal ability. The number of studies measuring the response of

spiders to fire is relatively high compared to other invertebrates, likely owing to their

ubiquity and high capture rates in pitfall traps (the most commonly used survey method for

invertebrates). Therefore, spiders are an ideal model system with the potential to be used as

indicators of ecological disturbance (Churchill 1997), and can offer important insights into

the impacts of fire on invertebrates and whether these depend on environmental factors,

phylogenetic history or species traits.

Fire may impact spiders directly by increasing mortality, or indirectly by altering habitat

structure and microclimate (Engstrom 2010; Underwood & Quinn 2010). A previous meta-

analysis on the effects of land management (including fire) on spiders found that fire

negatively affected spider abundance (Prieto-Benítez & Méndez 2011). However, the

conclusions of this study were limited as the analysis was performed for the order Araneae as

a whole and more informative impacts are expected at lower taxonomic levels (e.g. family,

species). Additionally, the number of studies investigating the impact of fire on spiders has

grown in the past ten years. Susceptibility to fire is likely to vary across taxonomic levels due

to differences in the behavioural, life-history and morphological traits of the family or

species, as well as their habitat (Engstrom 2010). For example, fire may have the greatest

impact on taxa associated with leaf-litter, as this is the substrate most often burned during

fires (York 1999; Hanula & Wade 2003; New et al. 2010), while burrowing species may be

sheltered from the heat and flames (Carrel 2008). Additionally, the simplified habitat

structure following removal of leaf-litter may favour the more active hunting species (Niwa

& Peck 2002), while burned habitats may be less favourable for web-building spiders if fire

changes the vegetation structure and reduces the availability of web anchor sites (Robinson

1981; Uetz 1992). Overall, the response of animals to fire, and adaptations and traits

underpinning this, is still poorly understood (Pausas & Parr 2018; Pausas 2019; Nimmo et al.

2021).

Here, we ask the question: can we utilise published data to detect global patterns of the

impacts of fire on spiders? We use a phylogenetically informed framework to determine

whether fire affects spider abundance at the family level. Since we can make some

generalisations about the presence of behavioural and life-history traits within families

(Cardoso et al. 2011b), we also investigate whether there is a relationship between response

to fire and family level traits, which could contribute to the development of predictive

frameworks for spiders and invertebrates more broadly. Additionally, ecosystems vary in

their fire history and susceptibility, so we consider study level variation in location,

vegetation type, fire type and time of sampling in our analyses. We show that there is huge

variation in spider family responses to fire between studies, but that there are some broad

scale patterns associated with time since fire which may have important implications for fire

frequency management under a changing climate.

METHODS

Literature search

We conducted a systematic literature search in Web of Science on 12th September 2020.

Titles, abstracts, and keywords were searched using the terms "(spider* OR araneae OR

arachnid* OR arthropod*) AND (*fire* OR burn*)" from any date and in any language. We

also performed further unstructured searches of published and grey literature (e.g. using

Google Scholar, SCOPUS and Fire Research Institute) as well as citations within returned

literature. Studies were included in our analysis if they: 1) used burned versus unburned,

before versus after, or before, after, control, impact (BACI) study designs, 2) identified

spiders to at least family level (but not studies that targeted a single species), 3) used pitfall

trapping, and 4) measured abundance. Of the 1,105 studies found, 49 met the criteria for

inclusion in the analysis and 33 of these provided usable abundance data with the manuscript

(Supplementary Information Table S1). There were two cases where two studies had used the

same underlying abundance data, so these were combined, resulting in 29 independent

datasets (see PRISMA 2020 diagram; Supplementary Information Figure S1). A list of the

data sources is found in Appendix 1.

Data collection

For each study we recorded the major vegetation type (forest, grassland, shrubland or

woodland), mean time since the last fire (in months, for the fire treatment), fire type

(controlled, wildfire) and latitude, and collated raw family level abundance for both the fire

(burned/after) and control (unburned/before) treatments. When a study contained multiple

fire treatments, we used the data for the single and/or most recent fire. Given our aim was to

analyse the effects of fire per family (and not per study), we did not extract the overall effect

sizes per study, as would be done in a traditional meta-analytical framework. Abundance data

were extracted from tables using Tabula (https://tabula.technology), or from figures using

metaDigitise (Pick et al. 2019) in R (version 4.1.0; R Core Team 2021). Taxonomy was

checked within each dataset and updated if necessary (i.e. we ensured the taxonomic

classification of species and genera within families was current) (World Spider Catalog

2021).

Abundance was standardised across and within studies by dividing absolute family

abundance by the number of plots, traps and sampling days (sampling effort), which allowed

us to compare abundance between treatments and studies. We recognise that this is not a

perfect calculation as it is based on our interpretation of the study design reported in the

literature, but it was necessary because there were multiple instances where experimental

designs were not uniform or directly comparable between treatments within studies. After

calculating a standardised abundance per family per study and per treatment, we derived a

measure of difference in abundance by subtracting the standardised abundance of the control

treatment from that of the fire treatment, where a negative value indicated lower abundance

in the fire treatment and a positive value indicated higher abundance in the fire treatment.

This resulted in 563 data points. For some statistical models (1 and 3, see below) we further

filtered the data to only include families with at least 5 data points (5 fire-control differences),

resulting in 485 data points.

To determine whether ecologically relevant traits predict a family's response to fire, we

compiled information on hunting strategies (whether species within the family use capture

webs, sensing webs, active hunting and/or ambushing), habitat use (whether species within

the family are active within vegetation and/or burrow) and dispersal (whether species within

the family use ballooning) based on the World Spider Trait database (Pekár et al. 2021) and

published literature (Figure 1; Supplementary Information Table S2). All traits were binary

(presence/absence). We investigated the relationships between these ecological traits using

the R package corrplot (Wei & Simko 2021), and detected significant ($P < 0.05$) correlations

between all hunting strategy traits and burrowing (Supplementary Information Figure S2).

Consequently, we performed a logistic principal component analysis using the R package

logisticPCA (Landgraf & Lee 2015) with the first two axes extracted explaining 88.1% of the

total variation in ecological traits. The first axis (hunt_PC1) was associated with ambushing

strategies (ambush, sensing web, burrow), while the second axis (hunt_PC2) separated

capture web use from active hunting (Supplementary Information Figure S3). Hunt_PC1,

hunt_PC2, vegetation and ballooning were used as predictors in Models 2 and 3 (see below).

Phylogeny and comparative analysis

Phylogenetically controlled mixed models were run using the R package MCMCglmm

(Hadfield 2010). As a random factor, we used the phylogeny of Fernández et al. (2018),

which used data generated by the Spider Tree of Life project (Wheeler et al. 2017; 3

mitochondrial and 3 nuclear markers) with transcriptomic data as a backbone constraint. This

was the most complete (at the family level) phylogeny available. We pruned all non-focal

families using the R package phytools (Revell 2012), and missing taxa were added to the tree

next to sister families using the minimum branch length (1.349), in particular the infraorder

mygalomorphae was updated based on a more recent phylogeny incorporating all recognised

mygalomorph families (Opatova et al. 2019; Figure 1). For each of the models described

below, we visually assessed convergence and ensured effective sample sizes were above 900.

With the exception of Model 1, all models were run for 150 000 iterations, thinning every

100 iterations, and the first 15 000 iterations were discarded as burnin. Weak priors (inverse

Wishart) were used for residuals (R) and random effects (G), with variance set to 1 and

different degrees of belief parameters (Supplementary Information Table S3), and for fixed

effects we used default Gaussian priors (B). Prior choice did not affect results. We extracted a

statistic that quantifies the percentage of variance explained by just the fixed factors, and by

both the fixed and random factors (equivalent to marginal and conditional R^2 respectively), in

each of our models (Nakagawa & Schielzeth 2013; Nakagawa et al. 2017).

We first tested whether family, time since fire (in months, for the fire treatment), or their

interaction predicted the difference in abundance between the fire and control treatment

(Model 1; Table 1). We included phylogeny and the study ID as random factors to control for

phylogenetic relatedness between families and multiple data points per study. Data points

were weighted by the sample effort for the study (sample effort = (plots*traps*days of fire

treatment + plots*traps*days of control treatment)/2) using the mev argument. We ran the

model for 500 000 iterations, thinning every 100 interactions, and discarded the first 50 000

iterations as burnin. To further investigate the relationship between family and difference in

abundance we ran an intercept only model (difference in abundance ~ 1) with 3 different

random factor structures, and compared them based on Akaike's information Criterion (AIC)

using the R package MuMIn (Bartoń 2020). The random factors used were A) phylogeny,

family and study ID, B) phylogeny and study ID, and C) study ID. This intercept-only model

allowed us to explicitly quantify the separate effect of phylogenetic relationships, family and

study ID. We also calculated the phylogenetic signal (lambda) and extracted the intercept for

each family from model 1 to identify differences across families in the effect of fire.

We next tested whether family or study level traits predicted the difference in abundance

between the fire and control treatment (Model 2; Table 1). For this model, we included the

interaction between all variables and time since fire (months), and phylogeny, family and

study ID as random factors, and again weighted data points by sample effort. Given the high

number of predictors in this model, to determine the best fitting model for the response

variable, we employed an all-subset model selection approach (dredge, MuMIn) based on

corrected AIC (AIC_C). Although models with $\Delta AIC_C < 2$ are generally considered to be as

good as the top model, we considered models with $\Delta AIC_C < 7$ to increase the probability of

retaining the best model (Symonds & Moussalli 2011). The cumulative AIC_C weight for

models with $\Delta AIC_C < 7$ was 0.77, equating to 77% confidence that the best model was

included in this subset (compared to 0.30 for models with $\Delta AIC_C < 2$). From this subset of

models we measured the relative importance of each variable (RIV) by summing the

normalised AIC_C weights across all models in which that variable appeared (Symonds &

Moussalli 2011).

Lastly, we extracted the slopes of the interaction term for each family from Model 1 (i.e.

change in difference in abundance over time since fire) and tested whether these relationships

were predicted by family level traits (Model 3, Table 1). Specifically, this model tested

whether the increase or decrease in abundance through time after fire could be explained by

family traits. We included phylogeny as a random factor.

RESULTS

Summary of literature used

We analysed abundance data from 33 studies (29 datasets; Supplementary Information Table

S1), incorporating 67 spider families. The data spanned 15 countries across 6 continents, with

North America having the most studies (13; Supplementary Information Figure S4). The

family Thomisidae had the most datapoints (51), followed by Lycosidae (47) and

Gnaphosidae (45), and the most common vegetation type was forest (16 studies). Most

studies used a burned versus unburned study design (28 studies) and/or measured the effect of

controlled burns (17 studies). Only a few studies provided details about fire intensity,

severity, frequency, or the fire history of study sites (e.g. the age of the long-unburned site)

and we were therefore unable to include this valuable information in our analysis.

Comparative analysis

Most families showed both positive and negative differences in abundance between

treatments (Figure 2). Time since fire (in months, for the fire treatment) had a significant

effect on the difference in abundance between fire and control treatment for the families

Lycosidae, Linyphiidae and Agelenidae (Model 1; marginal R^2 = 0.159, conditional R^2 =

0.505; Table 1). The relationship was positive for the families Lycosidae and Linyphiidae,

with abundance initially lower in the fire treatment for both families. However, from around

two years after fire onwards abundance was higher in the fire treatment for Lycosidae, and

around equal in both treatments for Linyphiidae (Figure 3). Conversely, the relationship was

negative for the family Agelenidae, with decreasing abundance in the fire treatment over

time, for at least 2 years (Figure 3). These relationships were significantly different to those

observed in multiple other families (Supplementary Information Table S4).

Phylogenetic relatedness between families was not a strong predictor of the difference in

abundance between fire and control treatments, as confirmed by examination of the random

effect of family (Supplementary Information Figure S5; Table S5), and the posterior

probability of the phylogenetic signal (l = 0.007, 95% CI = 1.08×10^{-10}–0.028) in the intercept

only model. Abundance tended to be lower in the fire treatment for the families Agelenidae,

Amaurobiidae and Anyphaenidae (in > 80% of studies including these families there was

lower abundance in the fire treatment and mean differences in abundance < 0; Figure 2)

although these families had a low number of datapoints, so we interpret this with caution (9, 6

and 10 datapoints respectively). Conversely, multiple families with a high number of

datapoints (≥30) showed mean differences of close to zero (Figure 2) and no change in

abundance over time (e.g. Gnaphosidae, Hahniidae, Salticidae, Theridiidae, Thomisidae;

Figure 3). Therefore, fire either does not affect the abundance of these families, or the effect

of fire differs between studies and can be either positive or negative.

Family level and study level traits (family: hunt_PC1, hunt_PC2, vegetation, ballooning;

study: latitude, time since fire, fire type, vegetation type) explained only a small proportion of

the variance in the difference in abundance between fire and control treatments (Model 2; full

model marginal R^2 = 0.00018, conditional R^2 = 0.00024; Table 1). Time since fire, vegetation

type, and their interaction were the most important predictors in the top-model set (models

with ΔAIC_C < 7; Supplementary Information Table S6). These three predictors appeared in

all 47 top models (RIV = 1.00), while other predictors in the top-model set had much lower

relative importance (RIV 0.02–0.42; Supplementary Information Table S6). Time since fire

was a significant predictor of difference in abundance within the forest vegetation type

(Supplementary Information Table S7). The relationship was such that abundance was

initially lower in the fire treatment, but higher in the fire treatment from around two years

after fire onwards (Figure 2). The change in difference in abundance over time within the

forest vegetation type was also significantly different from grassland and woodland

vegetation types (Figure 4; Supplementary Information Table S7). Results were qualitatively

the same when we used an alternative flat prior (Longdon et al. 2011) for this model

(Supplementary Information Table S3, Table S7).

No family level traits predicted the slope of the relationship between time since fire and

difference in abundance within each family (Model 3; Supplementary Information Table S8;

marginal R^2 = 0.121, conditional R^2 = 0.192).

DISCUSSION

We investigated the global impact of fire on spider abundance through a comparative analysis

of published data. Our results show that the effect of fire on spider families is independent of

phylogenetic relatedness, and closely related families do not tend to show similar responses.

In fact, we found that the effect of fire is highly variable among studies, and there is no

consistency in responses within families. It is therefore difficult to make broad

generalisations about the impacts of fire on spiders at the family level. We did, however, find

that time since fire had a significant effect on the abundance of some spider families, and

within forest vegetation types, which highlights the importance of sampling across temporal

scales. This is needed to fully understand the impact of fire and increasing fire frequencies

under climate change (Dowdy et al. 2019; Virgilio et al. 2019; Goss et al. 2020). In

particular, ecosystems may experience greater impacts when fire intervals are short, due to

reduced time for re-establishment and for ecological succession to progress (Kral et al. 2017;

Whitman et al. 2019). We found that any negative impacts of fire on spider abundance were

restricted to the first few years after fire, demonstrating at a broader scale what previous

studies have shown within invertebrate communities (e.g. Brennan et al. 2006; Underwood &

Quinn 2010; Pryke & Samways 2012; Radford & Andersen 2012; Podgaiski et al. 2013).

Most spider families in our study showed both positive and negative responses to fire. This is

likely due to variation in other site-specific factors that influence abundance, such as species

interactions, habitat structure, climate and seasonality, as well as fire history variables that we

were unable to include in our models (because they were not known/not reported), and other

ecological disturbances (e.g. logging). Time since fire did have a significant effect on

abundance, but only in three spider families: Lycosidae, Linyphiidae and Agelenidae.

Lycosids (wolf spiders) are active hunting spiders and are often reported to increase in

abundance following fire (Niwa & Peck 2002; Brennan et al. 2006; Haddad et al. 2015). Our

results showed a global pattern of higher abundance of lycosids in the fire treatment, but only

from around two years after fire onwards, suggesting that fire may have direct impacts

(mortality) but that burned habitats are beneficial to, or highly colonised by lycosids.

Similarly, the abundance of linyphiids (dwarf spiders) was initially lower in the fire treatment

but increased over time after fire. Linyphiids are very small, sheet web weaving spiders and

have been found to be effective colonisers of burned areas (Malumbres-Olarte et al. 2014).

Conversely, spiders of the family Agelenidae decreased in abundance over time since fire.

Agelenids (funnel weavers) build sheet webs in the open on grass and shrubs and the negative

relationship observed may reflect a lack of suitable vegetation for web sites in burned

habitats. Two other families showed a non-significant trend of a negative association with

fire: Amaurobiidae (hacklemesh weavers) and Anyphaenidae (ghost spiders). Amaurobiids

are sheet web weavers while anyphaenids are active hunters; however, along with agelenids,

all three families build silken retreats in vegetation. Therefore, response to fire may be

associated with finer details of vegetation structure that we were not able to test here.

Time since fire also had a significant effect on spider abundance in forests, with abundance

initially lower in the fire treatment but increasing over time, while there was no effect of fire

in grassland, shrubland or woodland habitats. A meta-analysis of the impact of fire on ants

found that ant diversity and abundance decreased in forest biomes, but not deserts, grasslands

or savannahs (Vasconcelos et al. 2017). This is hypothesised to be due to changes in

vegetation structure and habitat openness as the result of fire rather than fire *per se*

(Vasconcelos et al. 2017; Andersen 2019). Therefore, the impacts of fire are expected to be

greater for assemblages from more complex habitats (i.e. forests) where habitat modification

is more pronounced. Additionally, fire behaviour differs between vegetation types, with

grasslands and woodlands showing higher spread rates and intensities, more fine fuel biomass

consumption, and increased release of heat when compared to forest ecosystems (Gomes et

al. 2020). These differences in fire behaviour are likely to influence the extent of vegetation

modification and have direct and indirect impacts on invertebrate taxa.

We found no relationship between response to fire and family level traits, suggesting that a

trait-based approach does not increase predictive power at this taxonomic level. The traits

that we selected for this study can be somewhat generalised to the family level in spiders;

however, within families there will be variation between species in these and other

unmeasured traits (e.g. body size, longevity, diet). Previous studies, at the local scale, have

detected associations between response to fire and species-level traits. For example,

Malumbres-Olarte et al. (2013) found that ambushing, stalking and sheet web building

spiders were associated with unburned sites in New Zealand grasslands. Another study in an

Australian desert ecosystem found that recently burned sites had a higher abundance of

individuals with large body size, limited dispersal ability, burrowing habit and ambush

hunting strategies, but that patterns were not consistent across post-fire ages (Langlands et al.

2011). Additionally, species-level data would allow for detection of changes in species

composition and diversity, as well as abundance. For example, de Brito et al. (2011) found

that arachnid abundance was lower, but diversity was higher in burned areas of Brazilian

savannah. Similarly, Langlands et al. (2012) found no difference in spider abundance across

post-fire years, but significant changes in species composition, in Australian spinifex habitat.

Moretti et al. (2002) found that spider community composition (but not overall abundance)

varied between areas of Swiss chestnut forest experiencing different fire regimes. In all cases
there were species unique to certain post-fire ages. Here we analysed data at the family level
as this was the taxonomic resolution provided by most studies and gave us the largest dataset
to work with. One of the major challenges of working with invertebrate taxa is the high
proportion of unknown and undescribed species. It is estimated that fewer than half of all
spider species have been formally described (Agnarsson et al. 2013), meaning it is often not
possible to identify specimens to species level during surveys. Additionally, of the nearly
50,000 species currently recognised (World Spider Catalog 2021), trait data are available for
fewer than 8000 species (16%) (Pekár et al. 2021), and phylogenetic data is available for
fewer again. Therefore, although it would be more informative to investigate species-level
responses to fire (Brennan et al. 2006), this is often not possible given current knowledge
gaps.

It is important to note that our analysis reflects the available data, and studies testing the
response of spiders to fire are limited. Additionally, the data are biased towards ground-
dwelling taxa (which are captured most frequently in pitfall traps) but arboreal species
contribute to species richness in many ecosystems and may be more influenced by changes to
vegetation structure as the result of fire (Frizzo et al. 2012). Therefore, studies that employ
multiple sampling techniques designed to capture the full community assemblage, and that
identify specimens to species/morphospecies level, are likely to detect more informative
patterns in spider responses to fire, and life history traits underpinning responses. We also
recognise that replicated studies are challenging to achieve when working with large-scale
disturbances such as fire. Studies utilising future management burns (e.g. before-after-
control-impact studies) may provide the most robust design (Green 1974; Stewart-Oaten &
Bence 2001). These studies should sample at multiple time points post-fire and consider each
of those time points separately (i.e. not combine temporal sampling into one "after fire"
group). Additionally, we were not able to include fire regime characteristics in our analysis,
which are likely to be key factors affecting how animals respond to fire (Keeley et al. 2011;
Pausas 2019). Where possible, studies should report detailed information about fire severity,
intensity, frequency, season, size and time since last fire, for both the treatment and control
sites. Finally, we encourage authors to publish datasets along with their manuscripts so that
their studies can be included in analyses such as this one.

In conclusion, more research is needed to understand the impacts of fire on spiders and

invertebrates more broadly, and this will be most informative for management decisions

when conducted at the local scale due to site-specific factors. We show that caution is needed

when making broad-scale assumptions about spider family responses to fire, as fire can affect

closely related taxa differently. Currently, our ability to predict individual species and

community responses to fire is somewhat limited by a lack of knowledge, which highlights

the importance of further taxonomic research and fundamental studies of the biology and

ecology of invertebrate species. Although difficult to make global generalisations, we did

find evidence that short fire intervals may be a threat to some spider families, and in forest

ecosystems, where abundance initially decreased after fire. This is important for considering

the impact of changing fire regimes and designing management strategies. Finally, although

not without limitations (noted above), we show that analyses of published raw data can be

used to detect broad scale ecological patterns and provide a novel alternative to traditional

meta-analytical approaches, which is not limited by the questions asked in the original

studies.

ACKNOWLEDGEMENTS

This work was financially supported by the Ian Potter Foundation to JM and the Australian

Research Council, Discovery Early Career Researcher Award (200100500) to IM. We

acknowledge the authors of the original research analysed in this work.

COMPETING INTERESTS

The authors declare that they have no competing interests.

DATA ACCESSIBILITY

The datasets and code used during the current study are available from the Dryad Digital

Repository: doi XXXXXX. *Data will be archived upon acceptance of the manuscript.*

LITERATURE CITED

Agnarsson I, Coddington JA, Kuntner M. 2013. Systematics: progress in the study of spider

diversity and evolution. in Penney D, editor. Spider research in the 21st century:

trends & perspectives. Siri Scientific Press, Manchester, UK.

Andersen AN. 2019. Responses of ant communities to disturbance: Five principles for
understanding the disturbance dynamics of a globally dominant faunal group. Journal
of Animal Ecology **88**:350–362.

Bartoń K. 2020. Mu-MIn: Multi-model inference.

Benndorf J, Wissel B, Sell AF, Hornig U, Ritter P, Böing W. 2000. Food web manipulation
by extreme enhancement of piscivory: an invertebrate predator compensates for the
effects of planktivorous fish on a plankton community. Limnologica **30**:235–245.

Bond WJ, Keeley JE. 2005. Fire as a global 'herbivore': the ecology and evolution of
flammable ecosystems. Trends in Ecology and Evolution **20**:387–394.

Brennan KEC, Ashby L, Majer JD, Moir ML, Koch JM. 2006. Simplifying assessment of
forest management practices for invertebrates: How effective are higher taxon and
habitat surrogates for spiders following prescribed burning? Forest Ecology and
Management **231**:138–154.

Cardoso P, Erwin TL, Borges PAV, New TR. 2011a. The seven impediments in invertebrate
conservation and how to overcome them. Biological Conservation **144**:2647–2655.

Cardoso P, Pekár S, Jocqué R, Coddington JA. 2011b. Global patterns of guild composition
and functional diversity of spiders. PLoS ONE **6**:e21710.

Carrel JE. 2008. Differential survival of Geolycosa xera archboldi and G. hubbelli (Araneae,
Lycosidae) after fire in Florida scrub. Journal of Arachnology **36**:595–599.

Churchill TB. 1997. Spiders as ecological indicators: an overview for Australia. Memoirs of
Museum Victoria **56**:331–337.

Clarke PJ, et al. 2015. A synthesis of postfire recovery traits of woody plants inAustralian
ecosystems. Science of the Total Environment **534**:31–42.

De Brito Freire-Jr G, Motta PC. 2011. Effects of experimental fire regimes on the abundance
and diversity of cursorial arachnids of Brazilian savannah (cerrado biome). Journal of
Arachnology **39**:263–272.

Dowdy AJ, Ye H, Pepler A, Thatcher M, Osbrough SL, Evans JP, Di Virgilio G, McCarthy
N. 2019. Future changes in extreme weather and pyroconvection risk factors for
Australian wildfires. Scientific Reports **9**:1–12.

Driscoll DA, et al. 2010. Fire management for biodiversity conservation: Key research
questions and our capacity to answer them. Biological Conservation **143**:1928–1939.

Driscoll DA, Smith AL, Blight S, Sellar I. 2019. Interactions among body size, trophic level,
and dispersal traits predict beetle detectability and occurrence responses to fire.
Ecological Entomology **45**:300–310.

Engstrom RT. 2010. First-order fire effects on animals: review and recommendations. Fire
Ecology **6**:115–130.

Enright NJ, Fontaine JB, Lamont BB, Miller BP, Westcott VC. 2014. Resistance and
resilience to changing climate and fire regime depend on plant functional traits.
Journal of Ecology **102**:1572–1581.

Fernández R, Kallal RJ, Dimitrov D, Ballesteros JA, Arnedo MA, Giribet G, Hormiga G.
2018. Phylogenomics, Diversification Dynamics, and Comparative Transcriptomics
across the Spider Tree of Life. Current Biology **28**:1489–1497.

Frizzo TLM, Campos RL, Vasconcelos HL. 2012. Contrasting effects of fire on arboreal and
ground-dwelling ant communities of a neotropical savanna. Biotropica **44**:254–261.

Gomes L, Miranda HS, Silvério DV, Bustamante MMC. 2020. Effects and behaviour of
experimental fires in grasslands, savannas, and forests of the Brazilian Cerrado. Forest
Ecology and Management **458**:117804.

Goss M, Swain DL, Abatzoglou JT, Sarhadi A, Kolden C, Williams AP, Diffenbaugh NS.
2020. Climate change is increasing the risk of extreme autumn wildfire conditions
across California. Environmental Research Letters **15**:094016

Green RH 1974. Sampling design and statistical methods for environmental biologist. Wiley
Interscience, Chichester, UK.

Haddad CR, Foord SH, Fourie R, Dippenaar-Schoeman AS. 2015. Effects of a fast-burning
spring fire on the ground-dwelling spider assemblages (Arachnida: Araneae) in a
Central South African Grassland Habitat. African Zoology **50**:281–292.

Hadfield JD. 2010. MCMC methods for multi-response generalized linear mixed models: the
MCMCglmm R package. Journal of Statistical Software **33**:1–22.

Hanula JL, Wade DD. 2003. Influence of long-term dormant-season burning and fire
exclusion on ground-dwelling arthropod populations in longleaf pine flatwoods
ecosystems. Forest Ecology and Management **175**:163–184.

He T, Lamont BB, Pausas JG. 2019. Fire as a key driver of Earth's biodiversity. Biological
Reviews **94**:1983–2010.

Keeley JE, Pausas JG, Rundel PW, Bond WJ, Bradstock RA. 2011. Fire as an evolutionary
pressure shaping plant traits. Trends in Plant Science **16**:406–4011.

Keith DA. 2012. Functional traits: their roles in understanding and predicting biotic responses
to fire regimes from individuals to landscapes in Bradstock RA, M. GA, and Williams
RJ, editors. Flammable Australia: Fire Regimes, Biodiversity and Ecosystems in a
Changing World. CSIRO Publishing, Clayton.

Kral KC, Limb RF, Harmon JP, Hovick TJ. 2017. Arthropods and Fire: Previous Research
Shaping Future Conservation. Rangeland Ecology & Management **70**:589–598.

Landgraf AJ, Lee Y. 2015. Dimensionality Reduction for Binary Data through the Projection
of Natural Parameters.

Langlands PR, Brennan KEC, Framenau VW, Main BY. 2011. Predicting the post-fire
responses of animal assemblages: testing a trait-based approach using spiders. Journal
of Animal Ecology **80**:558–568.

Langlands PR, Brennan KEC, Ward B. 2012. Is the reassembly of an arid spider assemblage
following fire deterministic? Austral Ecology **37**:429–439.

Longdon B, Hadfield J, Webster CL, Obbard DJ, Jiggins FM. 2011. Host Phylogeny
Determines Viral Persistence and Replication in Novel Hosts. PLoS Pathogens
7:e1002260.

Losos JB. 2008. Phylogenetic niche conservatism, phylogenetic signal and the relationship
between phylogenetic relatedness and ecological similarity among species. Ecology
Letters **11**:995–1003.

Malumbres-Olarte J, Barratt BIP, Vink CJ, Paterson AM, Cruickshank RH, Ferguson CM,
Barton DM. 2013. Habitat specificity, Dispersal and burning season: Recovery
indicators in New Zealand native grassland communities. Biological Conservation
160:140–149.

Malumbres-Olarte J, Barratt BIP, Vink CJ, Paterson AM, Cruickshank RH, Ferguson CM,
Barton DM. 2014. Big and aerial invaders: dominance of exotic spiders in burned
New Zealand tussock grasslands. Biological Invasions **16**:2311–2322.

Menges ES. 2007. Integrating demography and fire management: an example from Florida
scrub. Australian Journal of Botany **55**:261–272.

Moretti M, Conedera M, Duelli P, Edwards PJ. 2002. The effects of wildfire on ground-
active spiders in deciduous forests on the Swiss southern slope of the Alps. Journal of
Applied Ecology **39**:321–336.

Moretti M, de Bello F, Roberts SPM, Potts SG. 2009. Taxonomical vs. functional responses
of bee communities to fire in two contrasting climatic regions. Journal of Animal
Ecology **78**:98–108.

Nakagawa S, Johnson PCD, Schielzeth H. 2017. The coefficient of determination R2 and
intra-class correlation coefficient from generalized linear mixed-effects models
revisited and expanded. Journal of the Royal Society Interface **14**:20170213.

Nakagawa S, Schielzeth H. 2013. A general and simple method for obtaining R2 from

generalized linear mixed effects models. Methods in Ecology and Evolution **4**:133–

142.

New TR, Yen AL, Sands DPA, Greenslade P, Neville PJ, York A, Collett NG. 2010. Planned

fires and invertebrate conservation in south east Australia. Journal of Insect

Conservation **14**:567–574.

Nimmo DG, Carthey AJR, Jolly CJ, Blumstein DT. 2021. Welcome to the Pyrocene : animal

survival in the age of megafire. Global Change Biology.

Niwa CG, Peck RW. 2002. Influence of prescribed fire on carabid beetle (Carabidae) and

spider (Araneae) assemblages in forest litter in Southwestern Oregon. Environmental

Entomology **31**:785–796.

Nyffeler M. 2017. An estimated 400 – 800 million tons of prey are annually killed by the

global spider community. The Science of Nature **104**:30.

Opatova V, Hedin M, Montes de Oca L, Král J, Bond JE. 2019. Phylogenetic systematics and

evolution of the spider infraorder Mygalomorphae using genomic scale data.

Systematic Biology **69**:671–707.

Pausas JG. 2019. Generalized fire response strategies in plants and animals. Oikos **128**:147–

153.

Pausas JG, Keeley JE. 2009. A burning story: The role of fire in the history of life.

BioScience **59**:593–601.

Pausas JG, Parr CL. 2018. Towards an understanding of the evolutionary role of fire in

animals. Evolutionary Ecology **32**:113–125.

Pekár S, Černecká Ľ, Wolff J, Mammola S, Cardoso P, Lowe E, Fukushima CS, Birkhofer K,

Herberstein ME. 2021. The world spider trait database v1.0, Masaryk University,

Brno. Available from https://spidertraits.sci.muni.cz.

Pick JL, Nakagawa S, A. NDW. 2019. Reproducible, flexible and high-throughput data

extraction from primary literature: The metaDigitise R package. Methods in Ecology

and Evolution **10**:426–431.

Podgaiski LR, Joner F, Lavorel S, Moretti M, Ibanez S, Mendonça MdS, Pillar VD. 2013.

Spider Trait Assembly Patterns and Resilience under Fire-Induced Vegetation Change

in South Brazilian Grasslands. PLoS ONE **8**:e60207.

Prieto-Benítez S, Méndez M. 2011. Effects of land management on the abundance and

richness of spiders (Araneae): A meta-analysis. Biological Conservation **144**:683–

691.

Pryke JS, Samways MJ. 2012. Importance of using many taxa and having adequate controls
for monitoring impacts of fire for arthropod conservation. Journal of Insect
Conservation **16**:177–185.

R Core Team. 2021. R: A language and environment for statistical computing. R Foundation
for Statistical Computing, Vienna, Austria.

Radford IJ, Andersen AN. 2012. Effects of fire on grass-layer savanna macroinvertebrates as
key food resources for insectivorous vertebrates in northern Australia. Austral
Ecology **37**:733–742.

Raven PH, Yeates DK. 2007. Australian biodiversity: threats for the present, opportunities for
the future. Australian Journal of Entomology **46**:177–187.

Revell LJ. 2012. phytools: an R package for phylogenetic comparative biology (and other
things). Methods in Ecology and Evolution **3**:217–223.

Robinson JV. 1981. The Effect of Architectural Variation in Habitat on a Spider Community:
An Experimental Field Study. Ecology **62**:73–80.

Rodman KC, Veblen TT, Andrus RA, Enright NJ, Fontaine JB, Gonzalez AD, Redmond MD,
Wion AP. 2020. A trait-based approach to assessing resistance and resilience to
wildfire in two iconic North American conifers. Journal of Ecology **109**:313–326.

Saunders ME, Barton PS, Bickerstaff JRM, Frost L, Latty T, Lessard BD, Lowe EC,
Rodriguez J, White TE, Umbers KDL. 2021. Limited understanding of bushfire
impacts on Australian invertebrates. Insect Conservation and Diversity **14**:285–293.

Shlisky A, Alencar A, Manta M, Curran LM. 2009. Overview: Global fire regime conditions,
threats, and opportunities for fire management in the tropics. Tropical Fire Ecology.
Springer Praxis Books, Springer, Berlin, Heidelberg.

Stewart-Oaten A, Bence JR. 2001. Temporal and spatial variation in environmental impact
assessment. Ecological Monographs **71**:305–339.

Stork NE. 2018. How Many Species of Insects and Other Terrestrial Arthropods Are There
on Earth? Annual Review of Entomology **63**:31–45.

Symonds MRE, Moussalli A. 2011. A brief guide to model selection, multimodel inference
and model averaging in behavioural ecology using Akaike's information criterion.
Behavioral Ecology and Sociobiology **65**:13–21.

Uetz GW. 1992. Foraging strategies of spiders. Trends in Ecology and Evolution **7**:155–159.

Underwood EC, Quinn JF. 2010. Response of ants and spiders to prescribed fire in oak
woodlands of California. Journal of Insect Conservation **14**:359–366.

Vasconcelos HL, Maravalhas JB, Cornelissen T. 2017. Effects of fire disturbance on ant
abundance and diversity: a global meta-analysis. Biodiversity and Conservation
26:177–188.

Virgilio GD, Evans JP, Blake SAP, Armstrong M. 2019. Climate Change Increases the
Potential for Extreme Wildfires. Geophysical Research Letters **46**.

Wei T, Simko V. 2021. R package "corrplot": visualization of a correlation matrix.

Wheeler WC, et al. 2017. The spider tree of life: phylogeny of Araneae based on target-gene
analyses from an extensive taxon sampling. Cladistics **33**:574–616.

Whitman E, Parisien M-A, Thompson DK, Flannigan MD. 2019. Short-interval wildfire and
drought overwhelm boreal forest resilience. Scientific Reports **9**:18796.

World Spider Catalog. 2021. World Spider Catalog ver. 22.0, Natural History Museum Bern.
Available from http://wsc.nmbe.ch.

Wurtsbaugh WA. 1992. Food-web modification by an invertebrate predator in the Great Salt
Lake (USA) Oecologia **89**:168–175.

York A. 1999. Long-term effects of frequent low-intensity burning on the abundance of litter-
dwelling invertebrates in coastal blackbutt forests of southeastern Australia. Journal
of Insect Conservation **3**:191–199.

TABLES AND FIGURES

Table 1. Summary of the three models tested in this study and their main findings.

Model	Response variable	Predictor variable/s	Random variable/s	R^2	Results	Summary
1	Difference in abundance (fire - control)	Family $\times$ Months	Phylogeny Study ID	$R^2m = 0.159$ $R^2c = 0.505$	Figure 3 Figure S6 Table S4	Change in difference in abundance over time for families Lycosidae, Linyphiidae, Agelenidae
2	Difference in abundance (fire - control)	Hunt_PC1 $\times$ Months Hunt_PC2 $\times$ Months Vegetation $\times$ Months Ballooning $\times$ Months Fire type $\times$ Months Vegetation type $\times$ Months Latitude $\times$ Months	Phylogeny Study ID Family	$R^2m = 0.00018$ $R^2c = 0.00024$	Figure 4 Table S6 Table S7	Vegetation type, months and their interaction are the most important predictors Change in difference in abundance over time for Forest vegetation type, which differs from Grassland and Woodland vegetation types
3	Slope (change in difference in abundance over time)	Hunt_PC1 Hunt_PC2 Vegetation Ballooning	Phylogeny	$R^2m = 0.121$ $R^2c = 0.192$	Table S8	No relationship

Figure 1. Reduced version of the spider phylogeny used in this study listing the use of capture web (cWeb), sensing web (sWeb), active (Act.), and ambush (Amb.) foraging strategies, burrowing (Burr.), vegetation (Veg.) and ballooning (Ball.). Images of unburned (top) and burned (bottom) woodland vegetation show how recent fire alters habitat structure. Species from some key families are shown: *Jotus* sp. (Salticidae; image credit: J. Schubert), *Tasmanicosa* sp. (Lycosidae; image credit: J. Schubert), *Eratigena duellica* (Agelenidae; image credit: T. Killick), *Neriene montana* on a sheet web (Linyphiidae; image credit: T. Killick) and *Idiommata* sp. in a burrow (Barychelidae; image credit: J. Schubert).

649 **Figure 2.** Difference in abundance between fire and control treatments for each family, for

650 each study. Blue = higher abundance in fire treatment, red = lower abundance in fire

651 treatment, small black dot = mean difference in abundance for that family. Only data for

652 families with 5 or more datapoints are presented. Four outlying datapoints were removed for

653 visualisation.

654
655

Figure 3. Relationship between time since fire (months) and difference in abundance

between fire and control treatment within each family (Model 1). There was a significant

effect of months for the families Agelenidae, Linyphiidae and Lycosidae (colours correspond

to images in Figure 1), and the relationship differed between multiple families (see

Supplementary Information Table S4). Three outlying datapoints removed for visualisation.

Grey shading = 95% confidence bounds

662
663

664 **Figure 4.** Relationship between time since fire (months) and difference in abundance

665 between the fire and control treatment in each of the four major vegetation types (Model 2).

666 The relationship was significant in the forest vegetation type, and this differed significantly

667 from grassland and woodland. Eight outlying datapoints were removed for visualisation.

668 Grey shading = 95% confidence bounds.

APPENDIX 1 – Data sources

Baretta, D., Brescovit, A. D., Knysak, I., & Cardosa, E. J. B. N. (2007). Trap and soil monolith sampled edaphic spiders (Arachnida: Araneae) in Araucaria angustifolia forest. *Scientia Agricola,* 64(4), 375–383. doi:10.1590/s0103-90162007000400008

Bess, E. C., Parmenter, R. R., McCoy, S., & Molles, M. C. (2002). Responses of a riparian forest-floor arthropod community to wildfire in the middle Rio Grande Valley, New Mexico. *Environmental Entomology,* 31(5), 774–784. doi:10.1603/0046-225X-31.5.774

Brantley, S. L. (2020). Responses of ground-dwelling spider (Arachnida: Araneae) communities to wildfire in three habitats in Northern New Mexico, USA, with notes on mites and harvestmen (arachnida: Acari, opiliones). *Diversity,* 12(10), 1–23. doi:10.3390/d12100396

De Brito Freire-Jr, G., & Motta, P. C. (2011). Effects of experimental fire regimes on the abundance and diversity of cursorial arachnids of Brazilian savannah (cerrado biome). *Journal of Arachnology,* 39(2), 263–272. doi:10.1636/CP10-85.

Gillette, N. E., Vetter, R. S., Mori, S. R., Rudolph, C. R., & Welty, D. R. (2008). Response of ground-dwelling spider assemblages to prescribed fire following stand structure manipulation in the southern Cascade Range. *Canadian Journal of Forest Research,* 38(5), 969–980. doi:10.1139/X07-245

Greenberg, C. H., Forrest, T. G., & Waldrop, T. (2010). Short-term response of ground-dwelling arthropods to prescribed fire and mechanical fuel reduction in a southern Appalachian Upland hardwood forest. *Forest Science,* 56(1), 112–121. doi:10.1093/forestscience/56.1.112

Haddad, C. R., Foord, S. H., Fourie, R., & Dippenaar-Schoeman, A. S. (2015). Effects of a fast-burning spring fire on the ground-dwelling spider assemblages (Arachnida: Araneae) in a Central South African Grassland Habitat. *African Zoology,* 50(4), 281–292. doi:10.1080/15627020.2015.1088400

Haskins, M. F., & Shaddy, J. H. (1986). The ecological effects of burning , mowing , and plowing on ground-inhabiting spiders (Araneae) in an old-field ecosystem. *The Journal of Arachnology,* 14(1), 1–13.

Higgins, J. W., Cobb, N. S., Sommer, S., Delph, R. J., & Brantley, S. L. (2014). Ground-dwelling arthropod responses to succession in a pinyon-juniper woodland. *Ecosphere,* 5(1). doi:10.1890/ES13-00270.1

Hore, U., & Uniyal, V. P. (2008). Effect of Prescribed Fire on Spider Assemblage in Terai Grasslands , India. *Turkish Journal of Arachnology,* 1(1), 15–36.

Jansen, R., Makaka, L., Little, I. T., & Dippenaar-Schoeman, A. (2013). Response of ground-dwelling spider assemblages (Arachnida, Araneae) to Montane Grassland management practices in South Africa. *Insect Conservation and Diversity,* 6(5), 572–589. doi:10.1111/icad.12013

Koponen, S. (1993). Ground-living spiders (Araneae) one year after fire in three subarctic forest types, Quebec (Canada). *Memoirs of the Queensland Museum,* 33(2), 575–578.

Koponen, S. (2003). Effects of intensive fire on the ground-living spider (Araneae) fauna of a pine forest. *European Arachnology,* 2003(1), 133–137.

Koponen, S. (2005). Early succession of a boreal spider community after forest fire. *The Journal of Arachnology,* 33(2), 230–235.

Krause, R. H., & Assmann, T. (2016). Impact of prescribed burning on a heathland inhabiting spider community. *Arachnology Letters,* 2016(51), 57–63. doi:10.5431/aramit5108

Kwok, A. B. C., & Eldridge, D. J. (2015). Does fire affect the ground-dwelling arthropod community through
changes to fine-scale resource patches? *International Journal of Wildland Fire,* 24(4), 550–559.
doi:10.1071/WF14088
Langlands, P. R., Brennan, K. E. C., Framenau, V. W., & Main, B. Y. (2011). Predicting the post-fire responses
of animal assemblages: testing a trait-based approach using spiders. *Journal of Animal Ecology,* 80(3),
558–568. doi:10.1111/j.1365-2656.2010.01795.x
Langlands, P. R., Brennan, K. E. C., & Ward, B. (2012). Is the reassembly of an arid spider assemblage
following fire deterministic? *Austral Ecology,* 37(4), 429–439. doi:10.1111/j.1442-9993.2011.02299.x
Larrivée, M., Fahrig, L., & Drapeau, P. (2005). Effects of a recent wildfire and clearcuts on ground-dwelling
boreal forest spider assemblages. *Canadian Journal of Forest Research,* 35(11), 2575–2588.
doi:10.1139/x05-169
Ma, Y., Li, Q., Feng, P., & Yang, Z. (2013). Diversity of ground-dwelling spider community in different
restoring times of post-fire forest, Cangshan Mountain, Yunnan Province. *Acta Ecologica Sinica,*
33(3).
Ma, Y., Yang, Z., Feng, P., & Li, Q. (2014). The influence of fire disturbance on the biotype structure and
seasonal dynamics of ground-dwelling spider on Cangshan Mountain, Yunnan Province. *Biodiversity*
Science, 22(2), 208–215. doi:10.3724/sp.j.1003.2014.13160
McFerran, D. M., McAdam, J. H., & Montgomery, W. I. (1995). The impact of burning and grazing on
heathland plants and invertebrates in County Antrim. *Biology and Environment-Proceedings of the*
Royal Irish Academy, 95B(1), 1–17.
Milne, M. A., Gonsiorowski, J., Tuft, N., Deno, B., Ploss, T., Acosta, J., . . . Venable, C. (2021). Effects of Fire
on Ground-Dwelling Spider (Araneae) Assemblages in Central Indiana Forests. *Environmental*
Entomology. doi:10.1093/ee/nvab051
Moretti, M. (2000). *Effects of winter fire on spiders.* Paper presented at the European Arachnology 2000, Århus.
Moretti, M., Conedera, M., Duelli, P., & Edwards, P. J. (2002). The effects of wildfire on ground-active spiders
in deciduous forests on the Swiss southern slope of the Alps. *Journal of Applied Ecology,* 39, 321–336.
doi:10.1016/S0378-1127(03)00314-1
Polchaninova, N., Savchenko, G., Ronkin, V., Drogvalenko, A., & Putchkov, A. (2019). Summer fire in steppe
habitats: Long-term effects on vegetation and autumnal assemblages of cursorial arthropods.
Hacquetia, 18(2), 213–231. doi:10.2478/hacq-2019-0006
Polchaninova, N., Tsurikov, M., & Atemasov, A. (2016). Effect of summer fire on cursorial spider (Aranei) and
beetle (Coleoptera) assemblages in meadow steppes of Central European Russia. *Hacquetia,* 15(2),
113–132. doi:10.1515/hacq-2016-0019
Pompozzi, G. A., Rodrigo Tizón, F., & Pelaéz, D. V. (2011). Effects of different frequencies of fire on an
epigeal spider community in southern Caldenal, Argentina. *50,* 6, 718–724.
Rose, S. J., & Goebel, P. C. (2015). Short-term impacts of prescribed burning on the spider community (order:
Araneae) in a small Ohio Grassland. *Ohio Journal of Science,* 115(2), 79–89.
doi:10.18061/ojs.v115i2.4901
Schaefer, V. M. (1980). Sukzession von Arthropoden in verbrannten Kiefernforsten. *Forstwissenschaftliches*
Centralblatt, 99, 341–356.

Spear, D. M., Adams, T. A., Boyd, E. S., Dipman, M. M., Staubus, W. J., & Meyer, W. M. (2017). The effects
of development, vegetation-type conversion, and fire on low-elevation Southern California spider
assemblages. *Invertebrate Biology,* 136(2), 134–145. doi:10.1111/ivb.12163
Underwood, E. C., & Quinn, J. F. (2010). Response of ants and spiders to prescribed fire in oak woodlands of
California. *Journal of Insect Conservation,* 14(4), 359–366. doi:10.1007/s10841-010-9265-7
Vickers, M. E., & Culin, J. D. (2014). Spider (O: Araneae) Responses to fire and fire surrogate fuel reduction in
a piedmont forest in upstate South Carolina. *Southeastern Naturalist,* 13(2), 396–406.
doi:10.1656/058.013.0219