TITLE: Assessing the impact of fire on spiders through a global comparative analysis

Claire A. McLean[a, 1], Jane Melville[a, b], Joseph Schubert[a], Rebecca Rose[a] & Iliana Medina[c]

[a] Sciences Department, Museums Victoria, 11 Nicholson Street, Carlton, VIC 3053, Australia

[b] School of Biological Sciences, Monash University, Clayton, VIC 3800, Australia

[c] School of BioSciences, The University of Melbourne, Royal Parade, Parkville, VIC 3010,

Australia

[1] Corresponding author:

CAM cmclean@museum.vic.gov.au

ORCID: 0000-0002-6855-5067

Level 3 West, Sciences Department, Museums Victoria, 11 Nicholson Street, Carlton, VIC

3053, Australia, Ph. +61 3 8341 7444

JM jmelv@museum.vic.gov.au; ORCID: 0000-0002-9994-6423

JS jschubert@museum.vic.gov.au; ORCID: 0000-0001-9257-5749

RR rrose@museum.vic.gov.au

IM iliana.medina@unimelb.edu.au; ORCID: 0000-0002-1021-5035

RUNNING TITLE: The impact of fire on spiders

ABSTRACT

In many regions fire regimes are changing due to anthropogenic factors. Understanding the responses of species to fire can help to develop predictive models and inform fire management decisions. Spiders are a diverse and ubiquitous group and can offer important insights into the impacts of fire on invertebrates and whether these depend on environmental factors, phylogenetic history, or functional traits. We conducted phylogenetic comparative analyses of data from studies investigating the impacts of fire on spiders. We investigated whether fire affects spider abundance or presence and whether ecologically relevant traits or site-specific factors influence species' responses to fire. Although difficult to make broad generalisations about the impacts of fire due to variation in site- and fire-specific factors, we find evidence that short fire intervals may be a threat to some spiders, and that fire affects abundance and species compositions in forests relative to other vegetation types. Orb and sheet web weavers were also more likely to be absent after fire than ambush hunters, ground hunters, and other hunters suggesting functional traits may affect responses. Finally, we show that analyses of published data can be used to detect broad scale patterns and provide an alternative to traditional meta-analytical approaches.

KEYWORDS: Araneae, burn, invertebrate, phylogenetic signal, time since fire, trait-based approach

BACKGROUND

Fire is an important ecological and evolutionary force that has shaped ecosystems worldwide [1, 2]. While fire plays a role in maintaining species richness and ecological processes in fire-prone regions [reviewed in 3], fire can also pose a significant threat to biodiversity and individual species in sensitive landscapes or under different fire regimes [2, 4]. One of the major consequences of global climate change has been the alteration of fire regimes; in many regions, wildfires are becoming more frequent and severe, as well as threatening sensitive ecosystems that have long been fire-free [5-7]. Understanding the impacts of fire, and different fire regimes, is therefore essential for effective conservation and natural resource management now and into the future [8].

Predictive models quantifying the impacts of fire on species and ecosystems can inform fire management strategies for biodiversity conservation, including prescribed fire regimes or fire suppression. Such generalisation at broad scales is often difficult due to local variation in species assemblages and environmental factors. However, trait-based approaches, focusing on the properties of an organisms (its traits) rather than its species and relating these traits to fire responses, may overcome this challenge [9]. Additionally, an understanding of which traits predict a species' response to fire may be useful for the management of understudied taxa. Such predictive frameworks have been used to assess the impacts of fire regimes on plant communities (e.g., [9-13]), and to a lesser extent on animals (e.g., [14-16]). One constraint of this approach is that closely related species may share similar functional traits (though not always the case; [17]), making it difficult to determine whether observed responses are due to traits or shared ancestry. However, analyses combining both trait and phylogenetic information can provide mechanistic insights into patterns of biodiversity across regions with different fire histories and help to identify species and ecosystems most at risk from changing fire regimes.

Despite their massive diversity and abundance, we have a limited understanding of the impact of fire on invertebrate communities [18, 19]. This is largely because, relative to vertebrate taxa, invertebrates have been poorly studied. It is estimated that around 80% of terrestrial arthropod species are yet to be described [20], and fundamental information about their ecology, life history, and geographic distributions remains largely unknown [21]. Invertebrates play important roles in ecosystem function and changes to invertebrate assemblages can also have knock-on effects across the food web [22, 23]. Therefore,

understanding how invertebrates respond to fire is important at both the species and

ecosystem level. Currently, this major knowledge gap makes it difficult to make informed

fire management decisions concerning invertebrates [18, 21].

Spiders (order Araneae) are one of the most diverse and successful groups of animals [24]

and it is estimated that there are approximately 80,000 species worldwide [24-26]. They are

found in almost all terrestrial ecosystems, on every continent except Antarctica, and are

important invertebrate predators [27]. Spiders exhibit diverse ecological, behavioural, and life

history traits [28], including variation in hunting strategies, dispersal ability, and body size.

The number of studies measuring the response of spiders to fire is relatively high compared

to other invertebrates, likely owing to their ubiquity and high capture rates in pitfall traps (the

most common survey method for invertebrates). Therefore, spiders are a potential model

system as indicators of ecological disturbance [29], and can offer important insights into the

impacts of fire on invertebrates and whether these depend on environmental factors,

phylogenetic history, or functional traits.

Fire may impact spiders directly by increasing mortality, or indirectly by altering habitat

structure and microclimate [30, 31]. A previous meta-analysis on the effects of land

management (including fire) on spiders found that fire negatively affected spider abundance

[32]. However, this analysis was performed for the order Araneae as a whole and more

informative impacts are expected at lower taxonomic levels (e.g., family, species).

Additionally, the number of studies investigating the impact of fire on spiders has grown in

the past ten years. Susceptibility to fire is likely to vary across taxonomic levels due to

differences in the behavioural, life-history, and morphological traits of the family or species,

as well as their habitat [31]. Spiders are often classified into guilds based on prey capture

methods, locomotion, and habitat use [33-35]. Such classification is useful if guild members

respond in similar ways to habitat disturbances such as fire. For example, fire may have the

greatest impact on taxa associated with leaf-litter, as this is the substrate most often burned

during fires [19, 36, 37], while burrowing species may be sheltered from the heat and flames

[38]. Additionally, the simplified habitat structure following fire may favour the more active

hunting species [39], while burned habitats may be less favourable for web-building spiders if

fire changes the vegetation structure and reduces the availability of web anchor sites [40, 41].

Overall, the response of animals to fire, and the adaptations and traits underpinning this, are

still poorly understood [42-44].

Here, we ask the question: does fire have a global impact on spiders, and can we utilise

published data to detect it? We use a phylogenetically informed framework to determine

whether fire affects ground-active spider abundance or presence. We also investigate whether

there is a relationship between response to fire and functional traits (hunting guild,

ballooning, body length), which could contribute to the development of predictive

frameworks for spiders and invertebrates more broadly. Additionally, ecosystems vary in

their fire history and susceptibility, so we consider study level variation in location,

vegetation type, fire type, and time of sampling in our analysis. We show that there is huge

variation in spider responses to fire among studies, but that despite this there are some broad-

scale patterns associated with hunting guild, vegetation type, and time since fire which may

have important implications. Such information may aid land managers and policymakers to

identify priority species for further study and make decisions for managing and conserving

fire-prone ecosystems under a changing climate.

METHODS

Literature search

We conducted a systematic literature search in Web of Science on 12[th] September 2020.

Titles, abstracts, and keywords were searched using the terms "(spider* OR araneae OR

arachnid* OR arthropod*) AND (*fire* OR burn*)" from any date and in any language. We

also performed further unstructured searches of published and grey literature (using Google

Scholar, SCOPUS, and Fire Research Institute) as well as citations within returned literature.

Studies were included in our analysis if they: 1) used burned versus unburned, before versus

after, or before, after, control, impact (BACI) study designs, 2) identified spiders to at least

family level, 3) recorded abundance or presence/absence, and 4) used pitfall trapping.

Requirement 4 was to allow us to standardise abundance measures and compare data between

studies (i.e., adjust abundance according to sampling effort; more details below). Of the 1,105

studies returned by the literature search, 139 were relevant, 47 met the above criteria for

inclusion in the analysis and 34 of these provided usable data with the manuscript (Table S1).

There were four cases where two studies had used the same underlying data, so these were

combined, resulting in 30 independent datasets (see PRISMA 2020 diagram; Figure S1). A

list of the data sources used in the study is provided in the supplementary material (Table S1).

Data collection

For each study we recorded the major vegetation type (forest, grassland, shrubland,

woodland), mean time since the last fire (in months, for the fire treatment), fire type

(controlled, wildfire), and latitude, and collated raw family and species level abundance and

species level presence/absence data for both the fire (burned/after) and control

(unburned/before) treatments. Studies with before versus after designs were only included if

they sampled in the same season/s before and after the fire, as timing affects spider

assemblages. For studies that sampled at multiple time points, we included data from all time

points up to ten years post fire. For the few studies that compared fire frequencies (single

versus multiple fires), we used data for the single fire only as many studies did not provide

fire frequency information and we were therefore unable to include this as a factor in the

models. Given our aim was to analyse the effects of fire per family/species (and not per

study), we did not extract the overall effect sizes per study, as would be done in a traditional

meta-analytical framework. Data were extracted from tables using Tabula

(https://tabula.technology), or from figures using metaDigitise [45] in R (version 4.1.0; [46]).

Taxonomy was checked for all datapoints and updated if necessary (i.e., we ensured the

taxonomic classification of species, genera, and families were current as per the World Spider

Catalog; [26]). We acknowledge that other landscape (e.g., species assemblages,

fragmentation, microhabitat structure) and fire regime (e.g., frequency, severity, size) factors

are likely to influence the response of spiders to fire. Testing relationships with these finer

details is challenging in a global study such as this due to a lack of available data and

insufficient statistical power to incorporate many factors. Here our aim was to detect broad-

scale patterns which may help to identify priority species and ecosystems and guide future

work at a local scale.

Sampling effort differed among studies, and there were multiple instances where

experimental designs were not uniform among treatments within studies (e.g., different

numbers of traps). Therefore, abundance (at the family and species level) was standardised by

dividing absolute abundance by the number of plots, traps, and sampling days (sampling

effort). We recognise that this is not a perfect calculation as it is based on our interpretation

of the study design reported in the literature, but it was necessary to make abundance

measures comparable. After calculating a standardised abundance (abundance/trap/day) per

family/species per study and per treatment, we derived a measure of difference in abundance

by subtracting the standardised abundance of the control treatment from that of the fire

treatment, where a negative value indicated lower abundance in the fire treatment and a

positive value indicated higher abundance in the fire treatment. We also classified species

into one of three categories: 1) present in control, absent in fire treatment ("disappeared"), 2)

present in both control and fire treatment ("no change"), 3) absent in control, present in fire

treatment ("colonised"). For the family level analysis, we removed families with fewer than

ten datapoints (ten fire-control differences). The presence/absence dataset was larger than the

abundance dataset because not all studies measured abundance or provided enough detail for

us to standardise abundance. This resulted in 564 datapoints for the family level analysis and

2,914 (presence/absence) and 2,595 (abundance) datapoints for the species level analyses.

We compiled information on hunting guild, ballooning ability, and female total body length

(mm) using the World Spider Trait Database [28] and taxonomic literature listed on the

World Spider Catalog [26] (Figure 1). These traits are expected to be ecologically relevant.

For example, changes in vegetation structure following fire may favour some behavioural

strategies over others [38-41], ballooning allows long distance dispersal and likely influences

the ability of species to colonise burned areas after fire [47], and there may be a relationship

between fire age and body length as size affects locomotion, dispersal ability, and desiccation

resistance [16, 48, 49]. Species were classified into one of eight hunting guilds as described

by Cardoso et al. [33] (ambush hunters, ground hunters, orb web weavers, other hunters,

sensing web weavers, sheet web weavers, space web weavers, specialists). Given the huge

diversity of spiders and the fact that many species are understudied, behavioural data were

not available for all species in our study (68.3 % of the dataset). In these cases, we assigned

the hunting guild of a congeneric species, or the family-level guilds of Cardoso et al. [33].

Similarly, the ballooning ability of individual species was missing for 79.9% of the dataset.

Again, in cases where species data were not available, ballooning (presence/absence) was

assigned based on whether it had been recorded in the genus or family [28, 47]. We must note

that a limitation of this approach is that congeneric and family-level guild data may not

reflect the actual hunting strategy or ballooning ability of species that lack behavioural data,

as there is variation within families and genera. Body length data were from species

descriptions or the World Spider Trait Database (averages were calculated if there were

multiple datapoints per species). Female body length was used as there were more data

available for female than male spiders, but again, we were unable to obtain body length

measurements for all the species included in our study (female body length was missing for

27.7% of the dataset). Rather than perform list-wise deletion (removing datapoints with

missing body length data), which can reduce statistical power and bias parameter estimates

[50], we employed a data imputation procedure. The Amelia II R package [51] was first used

to perform expectation-maximisation with bootstrapping for multiple imputation under an

assumption of missing at random (MAR). All variables from Models 2 and 3 were included

in the imputation step, and female body length was log transformed to meet the assumption of

normality. We produced 10 imputed datasets (m = 10) and confirmed convergence of the

chains visually. Each missing value was then replaced by the mean of its corresponding

imputed values as per the single centre imputation from multiple chained equation (SICE)

technique [52]. All analyses were performed with both the imputed and a reduced complete

case dataset, which did not affect inferences about female body length (the imputed variable).

Phylogeny and comparative analysis

Phylogenetically controlled mixed models were run using the R package MCMCglmm [53].

As a random factor, we used the phylogeny of Fernández et al. [54], which used data

generated by the Spider Tree of Life project ([55]; 3 mitochondrial and 3 nuclear markers)

with transcriptomic data as a backbone constraint. This was the most complete (at the family

level) phylogeny available. We pruned all non-focal families using the R package phytools

[56], and missing taxa were added to the tree next to sister families using the minimum

branch length (1.349), in particular the infraorder Mygalomorphae was updated based on a

more recent phylogeny incorporating all recognised mygalomorph families ([57]; Figure 1).

For each of the models described below, we visually assessed convergence and ensured

effective sample sizes were above 900. Weak priors (inverse Wishart) were used for residuals

(R) and random effects (G), with variance set to 1 and different degrees of belief parameters

(Table S2), and for fixed effects we used default Gaussian priors (B). Prior choice did not

affect results. We extracted a statistic that quantifies the percentage of variance explained by

just the fixed factors, and by both the fixed and random factors (equivalent to marginal and

conditional R^2 respectively), in each of our models [58, 59].

Using the family level dataset, we first tested whether family, vegetation type (vegetation),

fire type, absolute latitude (latitude), time since fire (months), or the interaction between

family and months predicted the difference in abundance between the fire and control

treatment (Model 1; Table 1). We included phylogeny and the study ID as random factors to

control for phylogenetic relatedness among families and multiple datapoints per study.

Datapoints were weighted by the sample effort for the study (sample effort =

(plots*traps*days of fire treatment + plots*traps*days of control treatment)/2) using the mev

argument. We ran the model (family = gaussian) for 400,000 iterations, thinning every 100

interactions, and discarded the first 40,000 iterations as burnin. To further investigate the

relationship between family and difference in abundance we ran an intercept only model

(difference in abundance ~ 1) with 3 different random factor structures, and compared them

based on Akaike's information Criterion (AIC) using the R package MuMIn [60]. The

random factors used were A) phylogeny, family, and study ID, B) phylogeny and study ID,

and C) study ID. This intercept-only model allowed us to explicitly quantify the separate

effects of phylogenetic relationships, family, and study ID. We also calculated the

phylogenetic signal (lambda) and extracted the intercept for each family from random factor

model A to identify differences across families in the effect of fire.

We next used the species level abundance dataset to test whether hunting guild (guild),

ballooning, female body length (body length), vegetation, fire type, latitude, or months

predicted the difference in abundance between the fire and control treatment (Model 2;

family = gaussian; Table 1). Phylogeny, species, and study ID were included as random

factors, and datapoints were again weighted by sample effort (mev). Finally, we tested

whether the same seven variables predicted the presence/absence of species in the control and

fire treatments (Model 3, family = threshold; Table 1). Given that the response variable was

an ordinal factor for this model, we fixed the residual variance to 1 (Table S2). Models 2 and

3 were both run for 150,000 iterations, thinning every 100 iterations, with the first 15,000

iterations discarded as burnin.

RESULTS

Summary of literature used

We analysed data from 34 studies (30 datasets; Table S1), incorporating 67 families and

1,329 species. The data spanned 16 countries across 6 continents, with North America having

the most studies (11; Figure S2). The most common vegetation type was forest (16 studies),

and most studies used a burned versus unburned study design (31 studies) and/or measured

the effect of controlled burns (16 studies). Only a few studies provided details about fire

intensity, severity, frequency, or the fire history of study sites (e.g., the age of the long-

unburned site) and we were therefore unable to include this valuable information in our

analysis.

Comparative analysis

Most families showed both positive and negative differences in abundance between

treatments (Figure S3). Time since fire (months) had a significant effect on the difference in

abundance between fire and control treatment for the family Lycosidae (Model 1; marginal

$R^2 = 0.483$, conditional $R^2 = 0.648$; Table 1). The relationship was positive with abundance

initially lower in the fire treatment, but higher in the fire treatment from around two years

after fire onwards (Figure 2). This relationship was significantly different to that observed in

multiple other families (Table S3). Additionally, forest vegetation differed from woodland

vegetation, and there was a marginally non-significant ($P = 0.067$) difference between forest

and grassland vegetation. The mean difference in abundance was positive (higher abundance

in the fire treatment) for forest, and negative (lower abundance in the fire treatment) for all

other vegetation types (Figure 3).

Phylogenetic relatedness among families was not a strong predictor of the difference in

abundance between fire and control treatments, as confirmed by examination of the random

effect of family (Figure S5; Table S4), and the posterior probability of the phylogenetic

signal (lambda = 0.016, 95% CI = $1.15 \times 10^{-7} - 0.057$) in the intercept only model.

Abundance tended to be lower in the fire treatment for the families Agelenidae and

Anyphaenidae (in > 80% of studies including these families there was lower abundance in the

fire treatment and mean differences in abundance < 0; Figure S3), although these families had

a relatively low number of datapoints, so we interpret this with caution (14 and 12 fire-

control differences respectively). Conversely, multiple families with a high number of

datapoints (≥ 30) showed mean differences of close to zero (Figure S3) and no change in

abundance over time (e.g., Gnaphosidae, Hahniidae, Salticidae, Theridiidae, Thomisidae;

Figure 2). Therefore, the effect of fire on each family likely differed between studies and

could be either positive or negative.

Guild, ballooning, body length, vegetation, fire type, latitude, and months explained only a

small proportion of the variance in the difference in abundance between fire and control

treatments (Model 2; full model marginal $R^2 = 0.0000046$, conditional $R^2 = 0.0000087$; Table

1). Sheet web weavers differed from other hunters, and there was a non-significant trend

between sheet web weavers and both ambush hunters and ground hunters ($P = 0.59$; Figure

4), and between forest and grassland vegetation types ($P = 0.086$; Figure S6; Table S5; Table

S6).

There was a significant relationship between presence/absence category and hunting guild

(Model 3; marginal $R^2 = 0.021$, conditional $R^2 = 0.214$; Table 1). Both orb web weavers and

sheet web weavers differed from ambush hunters, ground hunters, and other hunters in that a

greater proportion of the two web weaving guilds "disappeared" and fewer "colonised" after

fire (Figure 5; Table S7; Table S8). Time since fire and vegetation type also had a significant

effect on presence/absence (Table 1; Table S7). Species that "disappeared" were recorded

sooner after fire than those that "colonised" (Figure 6), while forests had more species that

"disappeared" and "colonised" than grasslands (Figure 7).

DISCUSSION

We investigated the global impact of fire on ground-active spiders through a comparative

analysis of published abundance and presence/absence data. Our results show that the effect

of fire on spider families is independent of phylogenetic relatedness, and closely related

families do not tend to show similar responses. In fact, we found that the effect of fire on

individual families is highly variable among studies. It is therefore difficult to make broad

generalisations about the impacts of fire on spiders at the family level; however, at the

species level, we detected relationships between presence/absence categories and hunting

guild, vegetation type, and time since fire. The relationship with time since fire highlights the

importance of sampling across temporal scales. An understanding of this is essential to

predict the impact of fire and increasing fire frequencies under climate change [5-7]. In

particular, ecosystems may experience greater impacts when fire intervals are short, due to

reduced time for re-establishment and for ecological succession to progress [61, 62].

It is important to note that our analyses reflect the available data, and studies testing the

response of spiders to fire are limited. Additionally, the data are biased towards ground-

dwelling taxa (which are captured most frequently in pitfall traps) but arboreal species

contribute to species richness in many ecosystems and may be more influenced by changes to

vegetation structure as a result of fire [63]. It was necessary to restrict our analysis to data

obtained from pitfall trapping as it allowed us to standardise abundance within and between

studies. However, studies that employ multiple sampling techniques designed to capture the

full spider assemblage are likely to detect more informative patterns in spider responses to

fire and life history traits underpinning those responses. We also acknowledge that pitfall trap

design varied among studies (size of trap, preservation fluid or lack thereof, traps with or

without a roof), which may influence the species recorded. Trap design was consistent among

treatments within studies; however, it is possible that the efficacy of pitfall traps differed

between unburned and burned habitat. For example, some species may be more attracted to

pitfall traps with roofs in recently burned sites where vegetation cover is minimal. These

potentially confounding factors should be considered when interpreting the results of this

study.

Most spider families in our analysis showed both positive and negative responses to fire. This

is likely due to differences in species assemblages and variation in other site-specific factors,

such as species interactions, habitat structure, climate, and seasonality, as well as fire history

variables that we were unable to include in our models (because they were not known/not

reported), and other ecological disturbances (e.g., logging). Time since fire did have a

significant effect on abundance for the family Lycosidae. Lycosids (wolf spiders) are ground-

dwelling active hunters and burrowing habits and high dispersal ability (ballooning) have

been recorded in some species [28]. Lycosids are often reported to increase in abundance

following fire [39, 64, 65]. Our results showed a global pattern of higher abundance of

lycosids in the fire treatment, but only from around two years after fire onwards. Prior to this,

abundance was lower in the fire treatment. This suggests that fire may have direct impacts

(mortality) but that burned habitats are beneficial to, or highly colonised by lycosids. Overall,

the family level analysis detected some relationships with time since fire, but this taxonomic

level is too coarse to investigate the predictors of response to fire.

Of the nearly 50,000 spider species currently recognised [26], trait data are available for

fewer than 8000 species (16%) [28]. Given this, we only included hunting guild, ballooning,

and female total body length in our analyses, three traits that are expected to influence

response to fire. Our results suggest that a trait-based approach could increase predictive

power when species level abundance or presence/absence data are analysed (as done in

Models 2 and 3). We found that orb web weavers and sheet web weavers differed from

ambush hunters, ground hunters, and other hunters in that a greater proportion of the two web

weaving guilds "disappeared" and fewer "colonised" in the fire treatment. Fire also had a

more negative effect on the abundance of sheet web weavers relative to other hunters. Web

weaving spiders rely on vegetation for web anchor points [66, 67], and vegetation structure is

affected by fire [68]. The higher proportion of species that "disappeared" may reflect a lack

of suitable vegetation for web sites, or decreased habitat complexity in burned habitats [40,

41]. These guilds have been found to be associated with unburned sites in previous studies,

for example, Malumbres-Olarte et al. [69] found that sheet web building spiders were
associated with unburned sites in New Zealand grasslands. Conversely, ambush hunters often
burrow or use retreat sites which may protect them during the fire event and/or be less altered
in the post fire environment (soil and rock do not burn) [38], while ground and other hunters
may benefit from increased hunting success in the more open habitat after fire [39]. Web
attachment points are also critical for space web weavers. There was a possible trend of more
species that "disappeared" in the space web weaver guild (Figure 5), but we did not detect
significant differences to the other guilds. It is also important to note that orb web weavers
are arboreal and therefore less likely to be captured in pitfall traps. However, this is unlikely
to affect differences between unburned and burned sites within studies, unless orb web
weavers move along the ground, and thus encounter pitfall traps, less in burned habitat, and it
is unknown whether habitat changes post-fire affect spider behaviour. Overall, we detected a
broad-scale relationship with hunting guild, but this is likely to interact with local site and fire
specific factors.

Family and species level responses to fire also differed between vegetation types. Mean
family level abundance was higher in the fire treatment for forests, but lower in the fire
treatment for grassland, shrubland, and woodland habitats. Additionally, a higher proportion
of species "disappeared" or "colonised" in the fire treatment for forest compared to grassland
vegetation types, suggesting there may be a higher turnover of species in forests following
fire. We did not analyse other diversity or community composition indices in this study, but
this would likely give a more complete picture of the effects of fire. For example, de Brito et
al. [70] found that arachnid abundance was lower, but diversity was higher in burned areas of
Brazilian savannah. Similarly, Langlands et al. [71] found no difference in spider abundance
across post-fire years, but significant changes in species composition in Australian spinifex
habitat. Moretti et al. [72] found that spider community composition (but not overall
abundance) varied between areas of Swiss chestnut forest experiencing different fire regimes.
In all cases there were species unique to certain post-fire ages. Changes in abundance and/or
species assemblages are predicted to be driven by changes to vegetation structure and habitat
openness as the result of fire rather than fire *per se* [73, 74]. Therefore, the impacts of fire
may be greater for assemblages from more complex habitats (i.e., forests) where habitat
modification by fire is more pronounced. Additionally, fire behaviour differs between
vegetation types, with grasslands and woodlands showing higher spread rates and intensities,
more fine fuel biomass consumption, and increased release of heat when compared to forest

ecosystems [75]. These differences in fire behaviour are likely to influence the extent of

vegetation modification and have direct and indirect impacts on invertebrate taxa. The

response of individual species to fire is also likely to be associated with finer details of

vegetation structure that we were not able to test here.

Due to variation in study designs and limited reporting, we were not able to include other fire

regime characteristics in our analysis, but these are likely to be key factors affecting how

spiders respond to fire [42, 76]. For example, Andersen and Müller [77] found that spider

abundance declined following late season, but not early season fires in northern Australia.

Mason et al. [78] found that no mygalomorph spiders survived a high intensity wildfire, while

nearly all survived a low intensity controlled burn in Western Australia. Additionally, Hanula

and Wade [37] found that the effect of fire frequency differed among spider genera in pine

stands in Florida. Many invertebrates also show seasonal variation in presence and abundance

[79, 80]; therefore, the timing of sampling will have a major influence on species

assemblages. Seasonality is difficult to incorporate into a global analysis due to geographic

variation in seasons and the timing of fires; however, to minimise its impacts, we only

included studies that surveyed control and fire treatments concurrently, or during the same

season before and after fire. Where possible, studies should report detailed information about

fire severity, intensity, frequency, season, size, and time since last fire, for both the treatment

and control sites, and these should be included in analyses. Additionally, we recognise that

replicated studies to test these factors are challenging to achieve when working with large-

scale and often unpredictable disturbances such as fire, and suggest studies utilising future

management burns (e.g., before-after-control-impact studies) may provide the most robust

design [81, 82]. Given that time since fire was found to be an important factor, these studies

should sample at multiple time points post-fire and consider each of those time points

separately (i.e., not combine temporal sampling into one "after fire" group).

CONCLUSIONS

More research is needed to understand the impacts of fire on spiders and invertebrates more

broadly, and this will be most informative for management decisions when conducted at a

local scale so that relationships with site-specific landscape and fire factors can be examined.

As such, caution is needed when making broad-scale assumptions, but despite this we did

find some global patterns in the response of spiders to fire and show that a trait-based

approach may increase predictive power. Currently, this predictive ability is somewhat

limited by a lack of knowledge, which highlights the importance of further taxonomic

research and fundamental studies of the biology and ecology of invertebrate species.

However, we did find evidence that hunting guilds differ in their response to fire, specifically

that fire has a negative impact on orb and sheet web weavers. Additionally, short fire

intervals may be a threat to some spiders, with abundance of lycosids initially lower after fire,

and "disappearing" species detected sooner after fire than "colonising" species. This is

important when considering the impact of changing fire regimes and designing fire

management strategies to conserve biodiversity (e.g., determining fire intervals), and may

help to identify priority species and ecosystems for more detailed study at a local scale.

Finally, although not without limitations (noted above), we show that analyses of published

raw data can be used to detect broad-scale ecological patterns and provide an alternative to

traditional meta-analytical approaches, which is not limited by the questions asked in the

original studies. We encourage authors to publish datasets along with their manuscripts so

that their studies can be easily included in analyses such as this one.

ACKNOWLEDGEMENTS

We acknowledge the authors of the original research analysed in this work.

FUNDING

This work was financially supported by the Ian Potter Foundation to JM and the Australian

Research Council, Discovery Early Career Researcher Award (200100500) to IM.

COMPETING INTERESTS

The authors declare that they have no competing interests.

DATA ACCESSIBILITY

The datasets and code used during the current study are available from the Dryad Digital

Repository: doi:xxxxxxx

AUTHORS' CONTRIBUTIONS

CAM, JM, and IM conceived of the research; CAM and RR performed the literature search;

CAM extracted the abundance/presence data; CAM and JS compiled the trait data; CAM and

IM designed and performed the comparative analysis; CAM and IM interpreted results; CAM

wrote the manuscript, and all authors contributed to revisions.

REFERENCES

1.   Pausas, J.G. & Keeley, J.E. 2009 A burning story: The role of fire in the history of life. *BioScience* **59**, 593–601. (doi:10.1525/bio.2009.59.7.10).

2.   Bond, W.J. & Keeley, J.E. 2005 Fire as a global 'herbivore': the ecology and evolution of flammable ecosystems. *Trends in Ecology and Evolution* **20**, 387–394. (doi:10.1016/j.tree.2005.04.025).

3.   He, T., Lamont, B.B. & Pausas, J.G. 2019 Fire as a key driver of Earth's biodiversity. *Biol. Rev.* **94**, 1983–2010. (doi:10.1111/brv.12544).

4.   Shlisky, A., Alencar, A., Manta, M. & Curran, L.M. 2009 Overview: Global fire regime conditions, threats, and opportunities for fire management in the tropics. In *Tropical Fire Ecology* (Springer, Berlin, Heidelberg, Springer Praxis Books.

5.   Virgilio, G.D., Evans, J.P., Blake, S.A.P. & Armstrong, M. 2019 Climate Change Increases the Potential for Extreme Wildfires. *Geophysical Research Letters* **46**. (doi:10.1029/2019GL083699).

6.   Goss, M., Swain, D.L., Abatzoglou, J.T., Sarhadi, A., Kolden, C., Williams, A.P. & Diffenbaugh, N.S. 2020 Climate change is increasing the risk of extreme autumn wildfire conditions across California. *Environmental Research Letters* **15**, 094016 (doi:10.1088/1748-9326/ab83a7).

7.   Dowdy, A.J., Ye, H., Pepler, A., Thatcher, M., Osbrough, S.L., Evans, J.P., Di Virgilio, G. & McCarthy, N. 2019 Future changes in extreme weather and pyroconvection risk factors for Australian wildfires. *Scientific Reports* **9**, 1–12. (doi:10.1038/s41598-019-46362-x).

8.   Driscoll, D.A., Lindenmayer, D.B., Bennett, A.F., Bode, M., Bradstock, R.A., Cary, G.J., Clarke, M.F., Dexter, N., Fensham, R., Friend, G., et al. 2010 Fire management for biodiversity conservation: Key research questions and our capacity to answer them. *Biol. Conserv.* **143**, 1928–1939. (doi:10.1016/j.biocon.2010.05.026).

9.   Keith, D.A. 2012 Functional traits: their roles in understanding and predicting biotic responses to fire regimes from individuals to landscapes. In *Flammable Australia: Fire*

Regimes, Biodiversity and Ecosystems in a Changing World (eds. R.A. Bradstock, G.A. M. &
R.J. Williams). Clayton, CSIRO Publishing.

10. Menges, E.S. 2007 Integrating demography and fire management: an example from
Florida scrub. *Aust. J. Bot.* **55**, 261–272. (doi:10.1071/BT06020).

11. Enright, N.J., Fontaine, J.B., Lamont, B.B., Miller, B.P. & Westcott, V.C. 2014
Resistance and resilience to changing climate and fire regime depend on plant functional
traits. *J. Ecol.* **102**, 1572–1581. (doi:10.1111/1365-2745.12306).

12. Clarke, P.J., Lawes, M.J., Murphy, B.P., Russell-Smith, J., Nano, C.E.M., Bradstock,
R., Enright, N.J., Fontaine, J.B., Gosper, C.R., Radford, I., et al. 2015 A synthesis of postfire
recovery traits of woody plants inAustralian ecosystems. *Sci. Total Environ.* **534**, 31–42.
(doi:10.1016/j.scitotenv.2015.04.002).

13. Rodman, K.C., Veblen, T.T., Andrus, R.A., Enright, N.J., Fontaine, J.B., Gonzalez,
A.D., Redmond, M.D. & Wion, A.P. 2020 A trait-based approach to assessing resistance and
resilience to wildfire in two iconic North American conifers. *J. Ecol.* **109**, 313–326.
(doi:10.1111/1365-2745.13480).

14. Langlands, P.R., Brennan, K.E.C., Framenau, V.W. & Main, B.Y. 2011 Predicting the
post-fire responses of animal assemblages: testing a trait-based approach using spiders.
Journal of Animal Ecology **80**, 558–568. (doi:10.1111/j.1365-2656.2010.01795.x).

15. Moretti, M., de Bello, F., Roberts, S.P.M. & Potts, S.G. 2009 Taxonomical vs.
functional responses of bee communities to fire in two contrasting climatic regions. *Journal
of Animal Ecology* **78**, 98–108. (doi:10.1111/j.1365-2656.2008.01462.x).

16. Driscoll, D.A., Smith, A.L., Blight, S. & Sellar, I. 2019 Interactions among body size,
trophic level, and dispersal traits predict beetle detectability and occurrence responses to fire.
Ecol. Entomol. **45**, 300–310. (doi:10.1111/een.12798).

17. Losos, J.B. 2008 Phylogenetic niche conservatism, phylogenetic signal and the
relationship between phylogenetic relatedness and ecological similarity among species.
Ecology Letters **11**, 995–1003. (doi:10.1111/j.1461-0248.2008.01229.x).

18.   Saunders, M.E., Barton, P.S., Bickerstaff, J.R.M., Frost, L., Latty, T., Lessard, B.D.,

Lowe, E.C., Rodriguez, J., White, T.E. & Umbers, K.D.L. 2021 Limited understanding of

bushfire impacts on Australian invertebrates. *Insect. Conserv. Divers.* **14**, 285–293.

(doi:10.1111/icad.12493).

19.   New, T.R., Yen, A.L., Sands, D.P.A., Greenslade, P., Neville, P.J., York, A. &

Collett, N.G. 2010 Planned fires and invertebrate conservation in south east Australia. *J.*

Insect Conserv. **14**, 567–574. (doi:10.1007/s10841-010-9284-4).

20.   Stork, N.E. 2018 How many species of insects and other terrestrial arthropods are

there on Earth? *Annual Review of Entomology* **63**, 31–45. (doi:10.1146/annurev-ento-020117-

043348).

21.   Cardoso, P., Erwin, T.L., Borges, P.A.V. & New, T.R. 2011 The seven impediments

in invertebrate conservation and how to overcome them. *Biol. Conserv.* **144**, 2647–2655.

(doi:10.1016/j.biocon.2011.07.024).

22.   Wurtsbaugh, W.A. 1992 Food-web modification by an invertebrate predator in the

Great Salt Lake (USA) *Oecologia* **89**, 168–175. (doi:10.1007/BF00317215.).

23.   Benndorf, J., Wissel, B., Sell, A.F., Hornig, U., Ritter, P. & Böing, W. 2000 Food

web manipulation by extreme enhancement of piscivory: an invertebrate predator

compensates for the effects of planktivorous fish on a plankton community. *Limnologica* **30**,

235–245. (doi:10.1016/S0075-9511(00)80053-5).

24.   Agnarsson, I., Coddington, J.A. & Kuntner, M. 2013 Systematics: progress in the

study of spider diversity and evolution. In *Spider research in the 21st century: trends &*

perspectives (ed. D. Penney). Manchester, UK, Siri Scientific Press.

25.   Raven, P.H. & Yeates, D.K. 2007 Australian biodiversity: threats for the present,

opportunities for the future. *Australian Journal of Entomology* **46**, 177–187.

(doi:10.1111/j.1440-6055.2007.00601.x).

26.   World Spider Catalog. 2021 World Spider Catalog ver. 22.0. Natural History Museum

Bern. (doi:http://wsc.nmbe.ch).

27.   Nyffeler, M. 2017 An estimated 400 – 800 million tons of prey are annually killed by

the global spider community. *The Science of Nature* **104**, 30. (doi:10.1007/s00114-017-1440-

1).

28.   Pekár, S., Wolff, J., Černecká, Ľ., Birkhofer, K., Mammola, S., Lowe, E.C.,

Fukushima, C.S., Herberstein, M.E., Kučera, A., Buzatto, B., et al. 2021 The World Spider

Trait database: a centralised global open repository for curated data on spider traits. Database

2021: baab064. Masaryk University, Brno. (doi:https://spidertraits.sci.muni.cz/).

29.   Churchill, T.B. 1997 Spiders as ecological indicators: an overview for Australia.

Mem. Mus. Vic. **56**, 331–337. (doi:10.24199/j.mmv.1997.56.21).

30.   Underwood, E.C. & Quinn, J.F. 2010 Response of ants and spiders to prescribed fire

in oak woodlands of California. *J. Insect Conserv.* **14**, 359–366. (doi:10.1007/s10841-010-

9265-7).

31.   Engstrom, R.T. 2010 First-order fire effects on animals: review and

recommendations. *Fire Ecology* **6**, 115–130. (doi:10.4996/fireecology.0601115).

32.   Prieto-Benítez, S. & Méndez, M. 2011 Effects of land management on the abundance

and richness of spiders (Araneae): A meta-analysis. *Biol. Conserv.* **144**, 683–691.

(doi:10.1016/j.biocon.2010.11.024).

33.   Cardoso, P., Pekár, S., Jocqué, R. & Coddington, J.A. 2011 Global patterns of guild

composition and functional diversity of spiders. *PLoS One* **6**, e21710.

(doi:10.1371/journal.pone.0021710).

34.   Uetz, G.W., Halaj, J. & Cady, A.B. 1999 Guild structure of spiders in major crops.

The Journal of Arachnology **27**, 270–280.

35.   Wolff, J.O., Wierucka, K., Paterno, G.B., Coddington, J.A., Hormiga, G., Kelly,

M.B.J., Herberstein, M.E. & Ramirez, M.J. 2022 Stabilized morphological evolution of

spiders despite mosaic changes in foraging ecology. *Systematic Biology* **71**, 1487–1503.

(doi:10.1093/sysbio/syac023).

36.   York, A. 1999 Long-term effects of frequent low-intensity burning on the abundance

of litter-dwelling invertebrates in coastal blackbutt forests of southeastern Australia. *J. Insect*

Conserv. **3**, 191–199. (doi:10.1023/A:1009643627781).

37.   Hanula, J.L. & Wade, D.D. 2003 Influence of long-term dormant-season burning and

fire exclusion on ground-dwelling arthropod populations in longleaf pine flatwoods

ecosystems. *For. Ecol. Manage.* **175**, 163–184. (doi:10.1016/S0378-1127(02)00130-5).

38.   Carrel, J.E. 2008 Differential survival of Geolycosa xera archboldi and G. hubbelli

(Araneae, Lycosidae) after fire in Florida scrub. *J. Arachnol.* **36**, 595–599. (doi:10.1636/T08-

14.1).

39.   Niwa, C.G. & Peck, R.W. 2002 Influence of prescribed fire on carabid beetle

(Carabidae) and spider (Araneae) assemblages in forest litter in Southwestern Oregon.

Environ. Entomol. **31**, 785–796. (doi:10.1603/0046-225X-31.5.785).

40.   Robinson, J.V. 1981 The effect of architectural variation in habitat on a spider

community: an experimental field study. *Ecology* **62**, 73–80. (doi:10.2307/1936670).

41.   Uetz, G.W. 1992 Foraging strategies of spiders. *Trends in Ecology and Evolution* **7**,

155–159. (doi:10.1016/0169-5347(92)90209-T).

42.   Pausas, J.G. 2019 Generalized fire response strategies in plants and animals. *Oikos*

128, 147–153. (doi:10.1111/oik.05907).

43.   Pausas, J.G. & Parr, C.L. 2018 Towards an understanding of the evolutionary role of

fire in animals. *Evolutionary Ecology* **32**, 113–125. (doi:10.1007/s10682-018-9927-6).

44.   Nimmo, D.G., Carthey, A.J.R., Jolly, C.J. & Blumstein, D.T. 2021 Welcome to the

Pyrocene : animal survival in the age of megafire. *Global Change Biology*.

(doi:10.1111/gcb.15834).

45.   Pick, J.L., Nakagawa, S. & Noble, D.W.A. 2019 Reproducible, flexible and high-

throughput data extraction from primary literature: The metaDigitise R package. *Methods in*

Ecology and Evolution **10**, 426–431. (doi:10.1111/2041-210X.13118).

46.   R Core Team. 2021 R: A language and environment for statistical computing.

(Vienna, Austria, R Foundation for Statistical Computing.

47.   Bell, J.R., Bohan, D.A., Shaw, E.M. & Weyman, G.S. 2005 Ballooning dispersal

using silk: world fauna, phylogenies, genetics and models. *Bulletin of Entomological*

Research **95**, 69–114. (doi:10.1079/BER2004350).

48.   Buckingham, S., Murphy, N. & Gibb, H. 2019 Effects of fire severity on the

composition and functional traits of litter-dwelling macroinvertebrates in a temperate forest.

For. Ecol. Manage. **434**, 279–288. (doi:10.1016/j.foreco.2018.12.030).

49.   Lazarina, M., Sgardelis, S.P., Tscheulin, T., Kallimanis, A.S., Devalez, J. &

Petanidou, T. 2016 Bee response to fire regimes in Mediterranean pine forests: The role of

nesting preference, trophic specialization, and body size. *Basic and Applied Ecology* **17**, 308–

320. (doi:10.1016/j.baae.2016.02.001).

50.   Nakagawa, S. 2015 Missing Data: Mechanisms, Methods and Messages. In

Ecological Statistics: Contemporary Theory and Application (eds. G.A. Fox, S. Negrete-

Yankelevich & S.V. J.), pp. 81–105. Oxford, Oxford University Press.

51.   Honaker, J., King, G. & Blackwell, M. 2011 Amelia II: A Program for Missing Data.

Journal of Statistical Software **45**, 1–47. (doi:10.18637/jss.v045.i07).

52.   Khan, S.I. & Hoque, A.S.M.L. 2020 SICE: an improved missing data imputation

technique. *Journal of Big Data* **7**, 37. (doi:10.1186/s40537-020-00313-w).

53.   Hadfield, J.D. 2010 MCMC methods for multi-response generalized linear mixed

models: the MCMCglmm R package. *Journal of Statistical Software* **33**, 1–22.

(doi:10.18637/jss.v033.i02).

54.   Fernández, R., Kallal, R.J., Dimitrov, D., Ballesteros, J.A., Arnedo, M.A., Giribet, G.

& Hormiga, G. 2018 Phylogenomics, Diversification Dynamics, and Comparative

Transcriptomics across the Spider Tree of Life. *Current Biology* **28**, 1489–1497.

(doi:10.1016/j.cub.2018.03.064).

55.   Wheeler, W.C., Coddington, J.A., Crowley, L.M., Dimitrov, D., Goloboff, P.A.,

Griswold, C.E., Hormiga, G., Prendini, L., Ramírez, M.J., Sierwald, P., et al. 2017 The spider

tree of life: phylogeny of Araneae based on target-gene analyses from an extensive taxon

sampling. *Cladistics* **33**, 574–616. (doi:10.1111/cla.12182).

56. Revell, L.J. 2012 phytools: an R package for phylogenetic comparative biology (and
other things). *Methods in Ecology and Evolution* **3**, 217–223. (doi:10.1111/j.2041-
210X.2011.00169.x).

57. Opatova, V., Hedin, M., Montes de Oca, L., Král, J. & Bond, J.E. 2019 Phylogenetic
systematics and evolution of the spider infraorder Mygalomorphae using genomic scale data.
Systematic Biology **69**, 671–707. (doi:10.1093/sysbio/syz064).

58. Nakagawa, S. & Schielzeth, H. 2013 A general and simple method for obtaining R2
from generalized linear mixed‐effects models. *Methods in Ecology and Evolution* **4**, 133–
142. (doi:10.1111/j.2041-210x.2012.00261.x).

59. Nakagawa, S., Johnson, P.C.D. & Schielzeth, H. 2017 The coefficient of
determination R2 and intra-class correlation coefficient from generalized linear mixed-effects
models revisited and expanded. *Journal of the Royal Society Interface* **14**, 20170213.
(doi:10.1098/rsif.2017.0213).

60. Bartoń, K. 2020 Mu-MIn: Multi-model inference. (

61. Whitman, E., Parisien, M.-A., Thompson, D.K. & Flannigan, M.D. 2019 Short-
interval wildfire and drought overwhelm boreal forest resilience. *Scientific Reports* **9**, 18796.
(doi:10.1038/s41598-019-55036-7).

62. Kral, K.C., Limb, R.F., Harmon, J.P. & Hovick, T.J. 2017 Arthropods and Fire:
Previous Research Shaping Future Conservation. *Rangel. Ecol. Manag.* **70**, 589–598.
(doi:10.1016/j.rama.2017.03.006).

63. Frizzo, T.L.M., Campos, R.L. & Vasconcelos, H.L. 2012 Contrasting effects of fire
on arboreal and ground-dwelling ant communities of a neotropical savanna. *Biotropica* **44**,
254–261. (doi:10.1111/j.1744-7429.2011.00797.x).

64. Haddad, C.R., Foord, S.H., Fourie, R. & Dippenaar-Schoeman, A.S. 2015 Effects of a
fast-burning spring fire on the ground-dwelling spider assemblages (Arachnida: Araneae) in a
Central South African Grassland Habitat. *Afr. Zool.* **50**, 281–292.
(doi:10.1080/15627020.2015.1088400).

65.   Brennan, K.E.C., Ashby, L., Majer, J.D., Moir, M.L. & Koch, J.M. 2006 Simplifying
assessment of forest management practices for invertebrates: How effective are higher taxon
and habitat surrogates for spiders following prescribed burning? *For. Ecol. Manage.* **231**,
138–154. (doi:10.1016/j.foreco.2006.05.035).

66.   Cunha, E.R., Thomaz, S.M., Mormul, R.P., Cafofo, E.G. & Bonaldo, A.B. 2012
Macrophyte structural complexity influences spider assemblages attributes in wetlands.
Wetlands **32**, 369–377. (doi:10.1007/s13157-012-0272-1).

67.   Mcnett, B.J. & Rypstra, A.L. 2008 Habitat selection in a large orb-weaving spider:
vegetational complexity determines site selection and distribution. *Ecol. Entomol.* **25**, 423–
432. (doi:10.1046/j.1365-2311.2000.00279.x).

68.   Foster, C.N., Barton, P.S., Robinson, N.M., MacGregor, C.I. & Lindenmayer, D.B.
2017 Effects of a large wildfire on vegetation structure in a variable fire mosaic. *Ecological*
Applications **27**, 2369–2381. (doi:10.1002/eap.1614).

69.   Malumbres-Olarte, J., Barratt, B.I.P., Vink, C.J., Paterson, A.M., Cruickshank, R.H.,
Ferguson, C.M. & Barton, D.M. 2013 Habitat specificity, Dispersal and burning season:
Recovery indicators in New Zealand native grassland communities. *Biol. Conserv.* **160**, 140–
149. (doi:10.1016/j.biocon.2013.01.004).

70.   De Brito Freire-Jr, G. & Motta, P.C. 2011 Effects of experimental fire regimes on the
abundance and diversity of cursorial arachnids of Brazilian savannah (cerrado biome). *J.*
Arachnol. **39**, 263–272. (doi:10.1636/CP10-85.).

71.   Langlands, P.R., Brennan, K.E.C. & Ward, B. 2012 Is the reassembly of an arid
spider assemblage following fire deterministic? *Austral Ecol.* **37**, 429–439.
(doi:10.1111/j.1442-9993.2011.02299.x).

72.   Moretti, M., Conedera, M., Duelli, P. & Edwards, P.J. 2002 The effects of wildfire on
ground-active spiders in deciduous forests on the Swiss southern slope of the Alps. *J. Appl.*
Ecol. **39**, 321–336. (doi:10.1016/S0378-1127(03)00314-1).

73.   Vasconcelos, H.L., Maravalhas, J.B. & Cornelissen, T. 2017 Effects of fire
disturbance on ant abundance and diversity: a global meta-analysis. *Biodivers. Conserv.* **26**,
177–188. (doi:10.1007/s10531-016-1234-3).

74. Andersen, A.N. 2019 Responses of ant communities to disturbance: Five principles

for understanding the disturbance dynamics of a globally dominant faunal group. *Journal of*

Animal Ecology **88**, 350–362. (doi:10.1111/1365-2656.12907).

75. Gomes, L., Miranda, H.S., Silvério, D.V. & Bustamante, M.M.C. 2020 Effects and

behaviour of experimental fires in grasslands, savannas, and forests of the Brazilian Cerrado.

For. Ecol. Manage. **458**, 117804. (doi:10.1016/j.foreco.2019.117804).

76. Keeley, J.E., Pausas, J.G., Rundel, P.W., Bond, W.J. & Bradstock, R.A. 2011 Fire as

an evolutionary pressure shaping plant traits. *Trends in Plant Science* **16**, 406–4011.

(doi:10.1016/j.tplan ts.2011.04.002).

77. Andersen, A.L. & Müller, W.J. 2000 Arthropod responses to experimental fire

regimes in an Australian tropical savannah: ordinal-level analysis. *Austral Ecol.* **25**, 199–209.

78. Mason, L., Bateman, P.W., Miller, B.P. & Wardell-Johnson, G.W. 2019 Ashes to

ashes: Intense fires extinguish populations of urban short-range endemics. *Austral Ecol.* **44**,

514–522. (doi:10.1111/aec.12685).

79. Frith, D. & Frith, C. 1990 Seasonality of litter invertebrate populations in an

Australian upland tropical rain forest. *Biotropica* **22**, 181–190.

80. Moeed, A. & Meads, M.J. 1985 Seasonality of pitfall trapped invertebrates in three

types of native forest, Orongorongo Valley, New Zealand. *New Zealand Journal of Zoology*

12, 17–53. (doi:10.1080/03014223.1985.10428264).

81. Green, R.H. 1974 *Sampling design and statistical methods for environmental*

biologist. Chichester, UK, Wiley Interscience.

82. Stewart-Oaten, A. & Bence, J.R. 2001 Temporal and spatial variation in

environmental impact assessment. *Ecological Monographs* **71**, 305–339.

(doi:10.2307/2657220).

Figure 1. Reduced version of the spider phylogeny used in this study listing the hunting guilds present within families: ambush hunters (Amb.), ground hunters (Ground), other hunters (Other), orb web weavers (Orb), sensing web weavers (Sens.), sheet web weavers (Sheet), space web weavers (Space), and specialists (Spec.). Species from highlighted families are shown: *Jotus* sp. (Salticidae; image credit: J. Schubert), *Tasmanicosa* sp. (Lycosidae; image credit: J. Schubert), *Eratigena duellica* (Agelenidae; image credit: T. Killick), *Neriene montana* on a sheet web (Linyphiidae; image credit: T. Killick), and *Idiommata* sp. in a burrow (Barychelidae; image credit: J. Schubert). Images of unburned (top) and burned (bottom) woodland vegetation show how recent fire can alter habitat structure.

Figure 2. Relationship between time since fire (months) and difference in abundance

between fire and control treatments within each family (Model 1). There was a significant

effect of months for the family Lycosidae, and this relationship differed to all other families

shown here (Table S3). Three outlying datapoints were removed for visualisation. Grey

shading = 95% confidence bounds (N = 564).

Figure 3. Difference in abundance between fire and control treatments for each vegetation

type, for each family in each study. Blue = higher abundance in fire treatment, red = lower

abundance in fire treatment, white = no difference, small black dot = mean difference in

abundance for that vegetation type. There was a significant difference between forest and

woodland vegetation types. Plot has been zoomed in to abundance differences between -0.1

and 0.1 for visualisation (N = 564).

Figure 4. Difference in abundance between fire and control treatments for each hunting

guild, for each species in each study. Blue = higher abundance in fire treatment, red = lower

abundance in fire treatment, white = no difference, small black dot = mean difference in

abundance for that guild. There was a significant difference between sheet web weavers and

other hunters. Plot has been zoomed in to abundance differences between -0.1 and 0.1 for

visualisation (N = 2595).

Figure 5. Proportion of datapoints (fire-control differences) for each hunting guild that fell

into the three presence/absence categories: present in control treatment only ("disappeared"),

present in both control and fire treatment ("no change"), present in fire treatment only

("colonised"). The total number of datapoints are shown within bars. There was a significant

difference between both orb and sheet web weavers and ambush hunter, ground hunters, and

other hunters (N = 2914).

Figure 6. Relationship between time since last fire and the three presence/absence categories: present in control treatment only ("disappeared"), present in both control and fire treatment ("no change"), present in fire treatment only ("colonised"; N = 2914).

Figure 7. Proportion of datapoints (fire-control differences) for each vegetation type that fell

into the three presence/absence categories: present in control treatment only ("disappeared"),

present in both control and fire treatment ("no change"), present in fire treatment only

("colonised"). The total number of datapoints are shown within bars. There was a significant

difference between forest and grassland vegetation types (N = 2914).

Table 1. Summary of the three models tested in this study and their main findings.

Model	Response variable	Predictor variable/s	Random variable/s	R^2	Results	Summary
1	Difference in abundance (fire - control)	Family Vegetation Fire type Latitude Months Family × months	Phylogeny Study ID	$R^2m = 0.483$ $R^2c = 0.647$	Figure 2 Figure 3 Figure S4 Table S3	Forest differs to woodland vegetation. Change in difference in abundance over time for Lycosidae.
2	Difference in abundance (fire - control)	Guild Ballooning Body length Vegetation Fire type Latitude Months	Phylogeny Study ID Species	$R^2m = 0.000005$ $R^2c = 0.000009$	Figure S6 Table S5 Table S6	Sheet web weavers differ to other hunters.
3	Difference in presence/absence (3 categories: disappeared, no change, colonised)	Guild Ballooning Body length Vegetation Fire type Latitude Months	Phylogeny Study ID Species	$R^2m = 0.026$ $R^2c = 0.211$	Figure 4 Figure 5 Figure 6 Figure 7 Table S7 Table S8	Orb web weavers and sheet web weavers differ to ambush hunters, ground hunters, and other hunters. Forest differs to grassland vegetation. Relationship with time since fire.