Integrated view and comparative analysis of baseline protein expression in mouse and rat tissues

Shengbo Wang[1†], David García-Seisdedos[1,2†], Ananth Prakash[1,2†*], Deepti Jaiswal Kundu[1],

Andrew Collins[3], Nancy George[1], Silvie Fexova[1], Pablo Moreno[1], Irene Papatheodorou[1,2],

Andrew R. Jones[3], Juan Antonio Vizcaíno[1,2 *]

[1] European Molecular Biology Laboratory - European Bioinformatics Institute (EMBL-EBI),

Wellcome Genome Campus, Hinxton, Cambridge, CB10 1SD. United Kingdom.

[2] Open Targets, Wellcome Genome Campus, Hinxton, Cambridge, CB10 1SD. United Kingdom.

[3] Institute of Systems, Molecular and Integrative Biology, University of Liverpool, Liverpool L69 7ZB, United Kingdom.

*Corresponding authors.

†All three authors have contributed equally and they wish to be considered as joint first authors.

Dr. Ananth Prakash. European Molecular Biology Laboratory, European Bioinformatics Institute (EMBL-EBI), Wellcome Trust Genome Campus, Hinxton, Cambridge, CB10 1SD, UK. Phone: + 44 (0) 1223 492610. Email: ananth@ebi.ac.uk

Dr. Juan Antonio Vizcaíno. European Molecular Biology Laboratory, European Bioinformatics Institute (EMBL-EBI), Wellcome Trust Genome Campus, Hinxton, Cambridge, CB10 1SD, UK. Phone: + 44 (0) 1223 492686. Email: juan@ebi.ac.uk.

Keywords

Mass spectrometry, quantitative proteomics, comparative protein expression, public data

reuse, rat proteome, mouse proteome, PRIDE database

Abstract

The increasingly large amount of proteomics data in the public domain enables, among other

applications, the combined analyses of datasets to create comparative protein expression

maps covering different organisms and different biological conditions. Here we have

reanalysed public proteomics datasets from mouse and rat tissues (14 and 9 datasets,

respectively), to assess baseline protein abundance. Overall, the aggregated dataset contained

23 individual datasets, including a total of 211 samples coming from 34 different tissues

across 14 organs, comprising 9 mouse and 3 rat strains, respectively.

In all cases, we studied the distribution of canonical proteins between the different organs.

The number of canonical proteins per dataset ranged from 273 (tendon) and 9,715 (liver) in

mouse, and from 101 (tendon) and 6,130 (kidney) in rat. Then, we studied how protein

abundances compared across different datasets and organs for both species. As a key point

we carried out a comparative analysis of protein expression between mouse, rat and human

tissues. We observed a high level of correlation of protein expression among orthologs

between all three species in brain, kidney, heart and liver samples, whereas the correlation of

protein expression was generally slightly lower between organs within the same species.

Protein expression results have been integrated into the resource Expression Atlas for

widespread dissemination.

Author summary

We have reanalysed 23 baseline mass spectrometry-based public proteomics datasets stored

in the PRIDE database. Overall, the aggregated dataset contained 211 samples, coming from

34 different tissues across 14 organs, comprising 9 mouse and 3 rat strains, respectively. We

analysed the distribution of protein expression across organs in both species. We also studied

how protein abundances compared across different datasets and organs for both species. Then

we performed gene ontology and pathway enrichment analyses to identify enriched biological

processes and pathways across organs. We also carried out a comparative analysis of baseline

protein expression across mouse, rat and human tissues, observing a high level of expression

correlation among orthologs in all three species, in brain, kidney, heart and liver samples. To

disseminate these findings, we have integrated the protein expression results into the resource

Expression Atlas.

1. Introduction

High-throughput mass spectrometry (MS)-based proteomics approaches have matured significantly in recent years, becoming an essential tool in biological research [1]. This has been the consequence of very significant technical improvements in MS instrumentation, chromatography, automation in sample preparation and computational analyses, among other areas. The most used MS-based experimental approach is Data Dependent Acquisition (DDA) bottom-up proteomics. Among the main quantitative proteomics DDA techniques, label-free intensity-based approaches remain very popular, although labelled-approaches, especially those techniques based on the isotopic labelling of peptides (MS^2 labelling), such as iTRAQ (Isobaric tag for relative and absolute quantitation) and TMT (Tandem Mass Tagging), are becoming increasingly used as well.

Following the steps initiated by genomics and transcriptomics, open data practices in the field have become embedded and commonplace in proteomics in recent years. In this context, datasets are now commonly available in the public domain to support the claims published in the corresponding manuscripts. The PRIDE database [2], located at the European Bioinformatics Institute (EBI), is currently the largest resource worldwide for public proteomics data deposition. PRIDE is also one of the founding members of the global ProteomeXchange consortium [3], involving five other resources, namely PeptideAtlas, MassIVE, iProX, jPOST and PanoramaPublic. ProteomeXchange has standardised data submission and dissemination of public proteomics data worldwide.

As a consequence, there is an unprecedented availability of data in the public domain, which is triggering multiple applications [4], including the joint reanalysis of datasets (so-called

meta-analysis studies) [5-7]. Indeed, public proteomics datasets can be systematically

reanalysed and integrated e.g., to confirm the results reported in the original publications,

potentially in a more robust manner since evidence can be strengthened if it is found

consistently across different datasets. Potentially, new insights different to the aims of the

original studies can also be obtained by reanalysing the datasets using different strategies, this

includes repurposing of public datasets [8], including for instance approaches such as

proteogenomics studies for genome annotation purposes [9-12].

In this context of reuse of public proteomics data, PRIDE has started to work on developing

data dissemination and integration pipelines into popular added-value resources at the EBI.

This is perceived as a more sustainable approach in the medium-long term than setting up

new independent bioinformatics resources. One of them is Expression Atlas [13], a resource

that has enabled over the years easy access to gene expression data across species, tissues,

cells, experimental conditions and diseases. Only recently, protein expression information

coming from reanalysed datasets has been integrated in the 'bulk' section of Expression

Atlas. As a result, proteomics expression data can be integrated with transcriptomics

information, mostly coming from RNA-Seq experiments. So far, we have performed two

meta-analysis studies involving the reanalysis and integration of: (i) 11 public quantitative

datasets coming from cell lines and human tumour samples [13]; and (ii) 24 human baseline

datasets coming from 31 different organs [14].

The next logical step is to perform an analogous study of baseline protein expression in two

of the main model organisms: *Mus musculus* and *Rattus norvegicus*. To date, there are only a

small number of bioinformatics resources providing access to reanalysed MS-based

quantitative proteomics datasets, and even fewer if one considers only mouse and rat data. In

this context, at the end of 2020, ProteomicsDB [15] released a first version of the mouse

proteome, based on the reanalysis of five label-free datasets. To the best of our knowledge,

there is no such public resource storing accurate MS-derived data for rat data yet. PaxDB is a

resource [16] that provides protein expression information coming from many species

(including mouse and rat) but the reported data relies on spectral counting, a technique that

generally does not provide the same level of accuracy than intensity-based label-free

approaches. Additionally, although antibody-based human protein expression information is

provided *via* the Human Protein Atlas [17] , their efforts are focused on human protein

expression.

Here, we report the reanalysis and integration of 23 public mouse (14 datasets) and rat (9

datasets) label-free datasets, and the incorporation of the results into the resource Expression

Atlas as baseline studies. Additionally, we report a comparative analysis of protein

expression across mouse, rat and human (in this case using the results reported at [14] using

the same methodology).

2. Results

2.1. Baseline proteomics datasets

Overall, we quantified protein expression from 34 healthy tissues in 14 organs coming from 23 datasets. The analyses covered a total of 1,173 MS runs from 211 samples that were annotated as healthy/control/non-treated samples, thus representing baseline protein expression. Non-control/disease samples associated with these datasets were also reanalysed but are not discussed here. Normalised protein abundances values (as ppb, parts per billion, see Methods for calculation) from both control/healthy/non-treated and disease/treated tissue samples are available to view as heatmaps in Expression Atlas. The protein abundances along with sample annotations, sample quality assessment summary and experimental parameter inputs for MaxQuant can be downloaded from Expression Atlas as text files. A summary of the data selection and reanalysis protocols is shown in Fig. 1. The total number of peptides and proteins identified in these datasets are shown in Table 1.

Figure 1. An overview of the study design and reanalysis pipeline. QA: Quality assessment.

Expression Atlas accession numbers	PRIDE dataset identifiers	Tissues	Organs	Species	Strains	Fractionation	Number of MS runs	Number of samples	Number of protein groups[†]	Number of peptides[†]	Number of unique peptides[†]	Number of unique genes mapped[†]
E-PROT-7[§]	PXD000867 [18]	Liver	Liver	*Mus musculus*	C57BL/6J	Yes	24	4	12,792	246,738	167,725	9,715
E-PROT-10[§]	PXD000288 [19]	Triceps muscles	Triceps Muscles	*Mus musculus*	C57BL/6	Yes	36	3	10,870	189,553	126,670	6,421
E-PROT-16	PXD003155 [20]	Cerebellum, Liver	Brain, Liver	*Mus musculus*	C57BL/6	No	24	12	4,508	59,696	45,728	3,797
E-PROT-74	PXD004612 [21]	Achilles and Plantaris tendon	Tendon	*Mus musculus*	C57BL/6	No	8	8	457	6,643	3,271	273
E-PROT-75	PXD005230 [22]	Hippocampus, Cerebellum, Cortex	Brain	*Mus musculus*	C57BL/10J	Yes	72	36	7,663	63,479	41,683	6,037
E-PROT-76	PXD009909 [23]	Retina	Eye	*Mus musculus*	ND4 Swiss Webster	Yes	12	1	5,002	29,454	24,961	3,686
E-PROT-77	PXD012307 [24]	Lung	Lung	*Mus musculus*	C57BL/6	No	32	2	6,809	106,391	73,950	5,795
E-PROT-78	PXD009639 [25]	Lens	Eye	*Mus musculus*	CD1	Yes	10	1	4,519	20,779	18,006	3,064
E-PROT-79	PXD019394 [26]	Heart, Kidney, Liver, Lung, Brain, Spleen, Testis, Pancreas	Heart, Kidney, Liver, Lung, Brain, Spleen, Testis, Pancreas	*Mus musculus*	Swiss-Webster	Yes	96	8	9,853	141,506	105,701	8,185
E-PROT-81	PXD012636 [27]	Left atrium, Left ventricle, Right atrium, Right ventricle	Heart	*Mus musculus*	C57BL/6	Yes	120	4	7,772	146,966	99,577	6,435
E-PROT-82	PXD019431 [28]	Articular cartilage	Articular cartilage	*Mus musculus*	BALB\ c	No	72	6	1,815	17,695	15,191	1,518
E-PROT-83	PXD022614 [29]	Brain	Brain	*Mus musculus*	C57BL/6J: Rj C57BL/6JR ccHsd	Yes	120	6	6,645	97,443	69,884	5,673

E-PROT-84	PXD004496 [30]	Hippocampus	Brain	*Mus musculus*	C57BL/6J	Yes	204	17	4,192	37,363	30,100	3,424
E-PROT-85	PXD008736 [31]	Right atrium, Sinus node	Heart	*Mus musculus*	C57BL/6J	Yes	143	6	7,906	144,926	94,379	6,554
E-PROT-86§	PXD012677 [32]	Amygdala	Brain	*Rattus norvegicus*	Sprague Dawley	No	3	3	1,872	15,326	12,367	1,382
E-PROT-87§	PXD006692 [33]	Lung	Lung	*Rattus norvegicus*	Sprague Dawley	No	10	10	2,079	14,440	11,696	1,398
E-PROT-88§	PXD016793 [34]	Liver	Liver	*Rattus norvegicus*	Sprague Dawley	No	8	8	4,787	57,998	46,411	3,743
E-PROT-89§	PXD004364 [35]	Testis	Testis	*Rattus norvegicus*	Sprague Dawley	No	3	3	2,351	15,880	13,674	1,601
E-PROT-91	PXD001839 [36]	Left ventricle	Heart	*Rattus norvegicus*	F344/BN	No	12	12	1,345	10,310	8,804	925
E-PROT-92§	PXD013543 [37]	Left ventricle	Heart	*Rattus norvegicus*	Wistar	No	8	8	1,858	17,303	13,622	1,340
E-PROT-93	PXD016958 [38]	First segment of proximal tubule, second segment of proximal tubule, third segment of proximal tubule, medullary thick ascending limb, cortical thick ascending limb, distal convoluted tubule, connecting tubule, cortical collecting duct, outer medullary collecting duct, inner medullary collecting duct	Kidney	*Rattus norvegicus*	Sprague Dawley	Yes	132	32	7,846	103,886	83,662	6,130

E-PROT-94	PXD003375 [39]	Caudal and rostral segments of spinal cord	Spinal cord	*Rattus norvegicus*	Wistar	Yes	21	18	2,477	29,213	22,025	1,926
E-PROT-95[§]	PXD015928 [40]	Tendon	Tendon	*Rattus norvegicus*	Wistar	No	3	3	199	1,253	1,063	101
TOTAL	23 datasets (Mouse: 14, Rat: 9)	34 tissues (Mouse: 21, Rat: 18)	14 organs (Mouse: 12, Rat: 8)				1,173 MS runs (Mouse: 973, Rat: 200)	211 samples (Mouse: 114, Rat: 97)				

153 **Table 1.** List of mouse and rat proteomics datasets that were reanalysed. [§]Only normal/untreated samples within this dataset are reported in this

154 study. However, results from both normal and disease samples are available in Expression Atlas. † Numbers after post-processing.

2.2. Protein coverage across organs and datasets

One of our main aims was to study protein expression across various organs. To enable a simpler comparison [14] we first grouped 34 different tissues into 14 distinct organs, as discussed in 'Methods'. We defined 'tissue' as a distinct functional or structural region within an 'organ'. We estimated the number of 'canonical proteins' identified across organs by first mapping all members of each protein group to their respective parent genes. We defined the parent gene as equivalent to the UniProt 'canonical protein' and we will denote the term 'protein abundance' to mean 'canonical protein abundance' from here on in the manuscript.

2.2.1. Mouse proteome

A total of 21,274 protein groups were identified from mouse datasets, among which 8,176 protein groups (38.4%) were uniquely present in only one organ and 70 protein groups (0.3%) were ubiquitously observed (see the full list in Supplementary File 2). This does not imply that these proteins are unique to these organs. Merely, this is the outcome considering the selected datasets. Mouse protein groups were mapped to 12,570 genes (canonical proteins) (Supplementary File 3). We detected the largest number of canonical proteins in samples coming from liver (9,920, 78.9% of the total) and the lowest numbers in samples from tendon (273, 2.2%) and articular cartilage (1,519, 12.1%) (Fig. 2A). In the case of tendon and articular cartilage, both experiments did not include sample fractionation in their sample preparation methodology, which can also explain the lower number of detected proteins. The comparatively even lower number of proteins identified in tendon could be attributed to the smallest sample size (only one sample out of 114, 0.9%). Also, tendon is a relatively hypocellular tissue, which has a low protein turnover rate. Dataset PXD000867, containing mouse liver samples, had the highest number of canonical proteins detected

(9,715, 77.3%), while the smallest number of proteins was detected in dataset PXD004612

(tendon, 273, 2.2%), as highlighted above (Fig. 2C).

We studied the normalised protein abundance distribution in organs (Fig. 2B) and found that

all organs, except tendon, had similar median abundances. However, one cannot attribute

further biological meaning to these observations, since by definition the method of

normalisation fixes each sample to have the same "total abundance", which then gets shared

out amongst all proteins. The normalised protein abundance distribution in datasets indicated

a higher than median abundances detected in datasets PXD004612 (tendon) and PXD003164

(testis) (Fig. 2D). A linear relationship was observed between the number of canonical

proteins detected in datasets and organs, when compared to the relative amount of their

spectral data (Fig. 2E). We found a significant number of proteins uniquely detected in one

organ (Fig. 2F). However, the list of concrete canonical proteins that were detected in just

one organ should be taken with caution since the list is subjected to inflated False Discovery

Rate (FDR), due to the accumulation of false positives when analysing the datasets

separately.

Some of the organs (liver, heart and brain) were represented across multiple mouse studies in

the aggregated dataset. A pairwise comparison of protein abundances in these organs

generally showed a good correlation in expression (heart: R^2 values ranged from 0.54 to 0.83;

brain: R^2 from 0.28 to 0.72; and liver: R^2 from 0.59 to 0.74) (Figure S1 in Supplementary File

4).

Figure 2. (A) Number of canonical proteins identified across different mouse organs. The

number within the parenthesis indicates the number of samples. (B) Range of normalised

iBAQ protein abundances across different organs. The number within the parenthesis

indicates the number of samples. (C) Canonical proteins identified across different datasets.

The number within the parenthesis indicate the number of unique tissues in the dataset. (D)

Range of normalised iBAQ protein abundances across different datasets. The number within

parenthesis indicate the number of unique tissues in the dataset. (E) Comparison of total

spectral data with the number of canonical proteins identified in each dataset and organ. (F)

Distribution of canonical proteins identified across organs.

2.2.2. Rat proteome

A total of 7,769 protein groups were identified across 8 different rat organs among which

3,649 (46.9%) protein groups were unique to one specific organ while 13 (0.16%) protein

groups were present among all organs (see full list in Supplementary File 2). The protein

groups were mapped to 7,116 genes (canonical proteins) (Supplementary File 3). The highest

number of canonical proteins (6,106, 85.1%) was found in rat kidney samples. The lowest

number of canonical proteins (101, 1.4%) was found in samples from tendon, as shown in

Fig. 3A. The largest number of canonical proteins identified in kidney is likely because of the

relatively large number of samples (32 samples), when compared to other organs. However, it

is interesting to note that large numbers of canonical proteins were detected in liver samples,

which relatively had fewer number of samples, when compared to the total number of

samples in heart and spinal cord.

Datasets PXD016958 and PXD016793 consisted entirely of kidney (where fractionation was

performed) and liver (no fractionation) samples, respectively, and as mentioned above had

the largest number of canonical proteins identified (Fig. 3C). The normalised protein

abundances were similar among the various organs and datasets (Fig. 3B, D). We also

observed a linear relation between the number of canonical proteins identified and the MS

spectra identified (Fig. 3E). As seen in the mouse datasets, we also observed a large number

of proteins uniquely detected in one organ (Fig. 3F). As highlighted above, the list of

concrete canonical proteins that were detected in just one organ should be taken with caution

since the list is subjected to inflated False Discovery Rate (FDR).

 In the case of rat datasets, left ventricle heart samples were the only ones represented in more

than one study (PXD001839 and PXD013543) in the aggregated dataset. A pairwise

comparison of protein abundances of heart between these two datasets was performed,

showing a strong correlation in protein expression ($R^2 = 0.9$) (Figure S1D in Supplementary

File 4).

Figure 3. (A) Number of canonical proteins identified across different rat organs. The

number within the parenthesis indicates the number of samples. (B) Range of normalised

iBAQ protein abundances across different organs. The number within the parenthesis

indicates the number of samples. (C) Canonical proteins identified across different datasets.

The number within the parenthesis indicate the number of unique tissues in the dataset. (D)

Range of normalised iBAQ protein abundances across different datasets. The number within

parenthesis indicate the number of unique tissues in the dataset. (E) Comparison of total

spectral data with the number of canonical proteins identified in each dataset and organ. (F)

Distribution of canonical proteins identified across organs.

2.3. Protein abundance comparison across organs

Next, we studied how protein abundances compared across different datasets and organs. The

presence of batch effects between datasets makes this type of comparisons challenging. To

aid comparison of protein abundances between datasets we transformed the normalised iBAQ

intensities into ranked bins as explained in 'Methods', i.e., proteins included in bin 5 are

highly abundant whereas proteins in bin 1 are expressed in the lowest abundances (among the

detected proteins).

2.3.1. Mouse proteome

We found that 1,086 (8.6%) proteins were found with their highest level of expression in at

least 3 organs, with a median bin value greater than 4 (Supplementary File 3). On the other

end of the scale, 138 (1.1%) canonical proteins were found with their lowest expression in at

least 3 organs, with a median bin value of less than 2. The bin transformed abundances in all

organs are provided in Supplementary File 3.

To compare protein expression across all organs, we calculated pairwise Pearson correlation

coefficients across 117 samples (Fig. 4A). We observed some correlation in protein

expression within brain (median $R^2 = 0.31$) and a higher one in heart (median $R^2 = 0.67$)

samples. We performed Principal Component Analysis (PCA) on all samples from mouse

datasets for testing the effectiveness of the bin transformation method in reducing batch

effects. Fig. 4B shows the clustering of samples from various organs of mouse. We observed

samples from the same organ generally clustered together. For example, we observed that

brain samples all clustered together in one group, even though they come from different

datasets, indicating decent removal of batch effects (Fig. 4C). However, we also observed

that samples from other organs such as liver did not cluster according to their organ types but

clustered together within the dataset they were part of, indicating some residual batch effects,

which are hard to remove completely.

Figure 4. (A) Heatmap of pairwise Pearson correlation coefficients across all mouse samples. The colour represents the correlation coefficient and was calculated using the bin transformed iBAQ values. The samples were hierarchically clustered on columns and rows using Euclidean distances. (B) PCA of all samples, using the binned protein abundances as input, coloured by the organ types. (C) PCA of all samples coloured by their respective dataset identifiers. The numbers in parenthesis indicate the number of datasets for each organ. Binned values of canonical proteins quantified in at least 50% of the samples were used to perform the PCA.

In addition, we compared the protein abundances generated in this study with the data available in the resource PaxDB generated using spectral counting across different mouse organs. We observed generally a strong correlation of protein abundances calculated using iBAQ from this study (fraction of total (FOT) normalised ppb) and spectral counting methods (Figure S2 in Supplementary File 4). However, the expression of low abundant proteins seemed to be underestimated in PaxDB when compared with our results, as shown by a S-shaped curve in the scatterplot in organs such as brain, heart, liver and lung. The 'dynamic exclusion' [41] setting used by modern mass spectrometers prevents the instrument from fragmenting abundant peptides multiple times when they are repeatedly observed in scans nearby in time. This has the effect that spectral counting approaches will limit the dynamic range observed, as high abundant proteins will be under sampled. This is a limitation when using spectral counting methods, and these days spectral counting is not commonly used as a truly quantitative data type in proteomics.

2.3.2. Rat proteome

Next, we studied the distribution of protein abundances across organs in rat. On one hand,

311 (4.3%) proteins were found with their highest expression in at least 3 organs with a

median bin value greater than 4. On the other hand, 27 (0.37%) canonical proteins were

found with their lowest expression in at least 3 organs, with a median bin value of less than 2.

The bin transformed abundances in all organs are provided in Supplementary File 3.

Overall, the samples from rat datasets showed a better correlation in protein expression (Fig.

5A) than in the case of mouse. We observed generally a strong correlation of protein

expression within samples from liver (median Pearson's correlation $R^2 = 0.85$), lung (median

$R^2 = 0.71$), spinal cord (median $R^2 = 0.65$), heart (median $R^2 = 0.71$) and brain (median $R^2 =$

0.86). We also observed the clustering in the PCA of samples coming from the same organ

(Fig. 5B). Kidney, lung, spinal cord and heart samples all clustered together according to

their organ type. Fig. 5C shows the samples based on the dataset they were part of. However,

most organ samples were part of individual datasets except in the case of samples from heart,

which came from two datasets (PXD001839 and PXD013543). Fig. 5C shows that the heart

samples clustered into two nearby groups (bottom left two clusters on Fig. 5B and 5C),

wherein each cluster included samples from a different dataset, indicating the presence of

small batch effects.

Figure 5. (A) Heatmap of pairwise Pearson correlation coefficients across all rat samples.

The colour represents the correlation coefficient and was calculated using the bin transformed

iBAQ values. The samples were hierarchically clustered on columns and rows using

Euclidean distances. (B) PCA of all samples coloured by the organ types. (C) PCA of all

samples coloured by their respective dataset identifiers. The numbers in parenthesis indicate

the number of datasets for each organ. Binned values of canonical proteins quantified in at

least 50% of the samples were used to perform the PCA.

2.4. The organ elevated proteome and the over-representative biological processes

Based on their expression, canonical proteins were classified into three different groups based

on their organ specificity: "mixed", "group-enriched" and "organ-enriched" (see

Supplementary File 5). We considered over-expressed canonical proteins in each organ as

those which were in "group-enriched" and "organ-enriched". The analysis showed that on

average, 20.8% and 26.0% of the total elevated canonical proteins were organ group-specific

in mouse and rat, respectively (Fig. 6). In addition, 4.3% and 14.2% were unique organ-

enriched in mouse and rat, respectively. The highest ratio of organ-enriched in mouse was

found in liver (13.6%), whereas in rat, it was found in kidney (39.8%).

We then performed a gene ontology (GO) enrichment analysis of those proteins that were

'organ-enriched' and group-enriched' using GO terms associated with biological processes.

We found 1,036 GO terms to be statistically significant in all organs, as seen in

Supplementary File 6. The most significant GO terms for each organ are shown in Table 2.

345

346 **Figure 6.** Organ specificity of canonical proteins in (A) mouse and (B) rat.

Organ	Species	GO ID	Description	adjusted p-value
Articular cartilage	*Mus musculus*	GO:0030198 GO:0043062 GO:0045229	Extracellular matrix organization Extracellular structure organization External encapsulating structure organization	$8.94*10^{-38}$ $8.94*10^{-38}$ $8.94*10^{-38}$
Brain	*Mus musculus*	GO:0050804 GO:0099177 GO:0050808	Modulation of chemical synaptic transmission Regulation of trans-synaptic signalling Synapse organization	$7.03*10^{-65}$ $7.03*10^{-65}$ $1.41*10^{-48}$

Heart	*Mus musculus*	GO:0060047 GO:0008016 GO:0060537	Heart contraction Regulation of heart contraction Muscle tissue development	$7.10*10^{-11}$ $4.43*10^{-10}$ $6.16*10^{-10}$
Kidney	*Mus musculus*	GO:0015711 GO:0044282 GO:0016054	Organic anion transport Small molecule catabolic process Organic acid catabolic process	$4.59*10^{-19}$ $4.91*10^{-15}$ $6.25*10^{-15}$
Eye	*Mus musculus*	GO:0007601 GO:0001654 GO:0099504	Visual perception Eye development Synaptic vesicle cycle	$7.54*10^{-50}$ $5.31*10^{-31}$ $8.36*10^{-18}$
Liver	*Mus musculus*	GO:0016569 GO:0016570 GO:0019369	Covalent chromatin modification Histone modification Arachidonic acid metabolic process	$6.26*10^{-10}$ $1.71*10^{-08}$ $1.71*10^{-08}$
Lung	*Mus musculus*	GO:0120031 GO:0030031 GO:0044782	Plasma membrane bounded cell projection assembly Cell projection assembly Cilium organization	$3.61*10^{-14}$ $3.61*10^{-14}$ $9.83*10^{-14}$
Pancreas	*Mus musculus*	GO:0007586 GO:0032328	Digestion Alanine transport	0.005 0.018
Spleen	*Mus musculus*	GO:0046649 GO:0050776 GO:0045087	Lymphocyte activation Regulation of immune response Innate immune response	$4.12*10^{-22}$ $2.00*10^{-20}$ $2.23*10^{-20}$
Tendon	*Mus musculus*	GO:0003012 GO:0050879 GO:0050881	Muscle system process Multicellular organismal movement Musculoskeletal movement	$1.46*10^{-25}$ $3.14*10^{-19}$ $1.46*10^{-25}$
Testis	*Mus musculus*	GO:0048232 GO:0003341 GO:0044782	Male gamete generation Cilium movement Cilium organization	$8.75*10^{-49}$ $3.04*10^{-38}$ $6.78*10^{-37}$
Triceps muscles	*Mus musculus*	GO:0061061 GO:0055002 GO:0003009	Muscle structure development Striated muscle cell development Skeletal muscle contraction	$1.56*10^{-14}$ $2.41*10^{-14}$ $3.53*10^{-14}$
Brain	*Rattus norvegicus*	GO:0099537 GO:0007268 GO:0098916	Trans-synaptic signalling Chemical synaptic transmission Anterograde trans-synaptic signalling	$1.79*10^{-60}$ $1.79*10^{-60}$ $1.79*10^{-60}$
Heart	*Rattus norvegicus*	GO:0061061 GO:0003012 GO:0055001	Muscle structure development Muscle system process Muscle cell development	$2.94*10^{-17}$ $6.30*10^{-16}$ $4.00*10^{-15}$
Kidney	*Rattus norvegicus*	GO:0006396 GO:0045944 GO:0006260	RNA processing positive regulation of transcription by RNA polymerase II DNA replication	$6.19*10^{-13}$ $7.29*10^{-06}$ $1.74*10^{-05}$
Liver	*Rattus norvegicus*	GO:0008202 GO:0016054 GO:0032787	Steroid metabolic process Organic acid catabolic process Monocarboxylic acid metabolic process	$2.74*10^{-10}$ $1.61*10^{-09}$ $1.64*10^{-09}$
Lung	*Rattus norvegicus*	GO:0031589 GO:0009617 GO:0030036	Cell-substrate adhesion Response to bacterium Actin cytoskeleton organization	$7.62*10^{-08}$ $7.62*10^{-08}$ $1.40*10^{-07}$
Spinal cord	*Rattus norvegicus*	GO:0061564 GO:0099537 GO:0007268	Axon development Trans-synaptic signalling Chemical synaptic transmission	$4.26*10^{-18}$ $5.93*10^{-16}$ $5.93*10^{-16}$
Tendon	*Rattus norvegicus*	GO:0030199 GO:0061448 GO:0001501	Collagen fibril organization Connective tissue development Skeletal system development	$1.23*10^{-13}$ $2.31*10^{-09}$ $3.39*10^{-09}$

| Testis | *Rattus norvegicus* | GO:0019953 GO:0051704 GO:0007018 | Sexual reproduction Multi-organism process Microtubule-based movement | $3.98*10^{-24}$ $1.61*10^{-18}$ $4.00*10^{-12}$ |

Table 2. Analysis of the top three GO terms for each organ in mouse and rat using the

elevated organ-specific and group-specific canonical proteins as described in the 'Methods'

section.

2.5. Protein abundances across orthologs in three species

In a previous study, we analysed 25 label-free proteomics datasets from healthy human

samples to assess baseline protein abundances in 14 organs following the same analytical

methodology [14]. We compared the expression of canonical proteins identified in all three

species (rat, mouse and human). Overall, 13,248 detected human genes (corresponding to the

canonical proteins) were compared with 12,570 genes detected in mouse and 7,116 genes

detected in rat. The number of orthologous mappings (i.e., "one-to-one" mappings, see

'Methods') between rat, mouse and human genes are listed in table 3. We only considered

one-to-one mapped orthologues for the comparison of protein abundances.

Species	Identified genes	Orthologs of human genes identified in [14]	Percentage of genes with different mapping against identified human genes				
			one-to-one	one-to-many	many-to-many	many-to-one	not mapped
Mus musculus	12,570	10,601	80.4%	1.9%	0.56%	1.46%	15.7%
Rattus norvegicus	7,116	6,058	82.0%	2.2%	0.70%	0.25%	14.9%

364 **Table 3.** Homologs identified in mouse and rat datasets when compared with the background

365 list of genes (corresponding to canonical proteins) identified in human datasets

366 (Supplementary File 2 in [14]).

Among human and mouse orthologues we observed relatively high levels of correlation of
protein abundances in brain ($R^2 = 0.61$), heart ($R^2 = 0.65$) and liver ($R^2 = 0.56$) (Fig. 7A).
Human and rat orthologs showed also relatively high levels of correlation in brain ($R^2 = 0.62$), kidney ($R^2 = 0.53$) and liver ($R^2 = 0.56$), but almost no correlation in lung ($R^2 = 0.12$)
and testis ($R^2 = 0.18$) (Fig. 7B). Between mouse and rat orthologs, the correlation of protein
abundances was higher in liver ($R^2 = 0.65$), kidney ($R^2 = 0.54$) and brain ($R^2 = 0.57$) samples,
when compared to the samples coming from the rest of the organs (Fig. 7C). Fig. 7D shows
an illustration of some example comparisons of individual orthologs using binned protein
abundances.

For the same corresponding subsets, we also investigated the correlation of protein
expression between various organs within each organism. We observed that in general the
correlation of protein expression was slightly lower between organs within the same species,
when compared to a higher correlation, which was observed among orthologs (Figure S3 in
Supplementary File 4). The found lower correlation of protein expression between different
organs was more apparent in mouse and rat.

Among the orthologs expressed in all organs in all three species, 747 (12.3%) orthologs were
detected with a median bin expression value of more than 4, i.e., proteins that appear to have
conserved high expression in all organs and all tissues. Additionally, 13 (0.2%) orthologs
were found with a median bin expression value less than 2 in all organs, although, it is harder
to detect consistently proteins with low abundances across all organs. A full list of the binned
protein abundances of orthologs is available in Supplementary File 7. The illustration of all
binned protein abundances across the three species is shown in Supplementary File 8.

Figure 7. Comparison of protein abundances (in ppb) between one-to-one mapped orthologs

of mouse, rat and human in various organs. (A) Pairwise correlation using normalised protein

abundances of human and mouse orthologues. (B) Human and rat orthologs. (C) Mouse and

rat orthologs. (D) As an example, the comparisons of binned protein expression of ten

randomly sampled orthologs are shown. Data corresponding to all cases (as reported in panel

D) are available in Supplementary File 7 and the corresponding illustration of binned values

is available in Supplementary File 8. Orthologs in (D) are shown using their human gene

symbol.

Since each sample contains potentially thousands of protein values this creates a high level of

dimensionality within the data. To reduce this, we used the non-linear dimension reduction

algorithm, Uniform Manifold Approximation and Projection (UMAP) (see Section 4.7 in the

'Methods' section). The UMAP algorithm enables the reduction of multidimensional data to

a two-dimensional space upon which the relationship between each sample can be visualised.

Specifically, it enables the visualisation of the relationships of proteins across individual

samples and organs. Should multiple samples be positioned near to each other, it allows for

us to predict that these samples shared similar properties (in this case, similar protein

abundance values). Consequently, by overlaying samples from various species UMAP

representations can be used to visualise the relationship of various orthologs across similar

organs.

Using the UMAP algorithm, we were able to visualise the relationships between individual

organs regardless of the involved species (human, mouse, rat) and to identify similar genes

(corresponding to canonical proteins) within those organs. The overall view of all samples

labelled by their respective organ is shown as Figure 8A. We chose to use the biological

system as the basis for the colouring scheme for each sample to reduce the overall complexity

of the visualisation, due the high number of organs included. By using this labelling scheme,

we could see that the clustering of each sample was deterministic. Each sample was

positioned within a clear region for the corresponding organs, despite the original layout

being unaware of this information. This indicates that not only do the samples within those

organs share common protein abundance values, but furthermore, that samples that come

from the same organs share similar protein expression (as three species are present).

Furthermore, in Figure 8B we show the representation of binned protein abundance values for

three example genes (SH3GL2, MYOZ2 and PYROXD2), providing information on the

abundance of them across different biological systems. These visualisations use the same

layout than within Figure 8A. In the example of SH3GL2, it can be seen that Figure 8B

shows multiple values that have been scored as bin 5. By referring to Figure 8A, we can see

that those points corresponding to highly abundant proteins, come from samples from the

nervous system (in all three species). Furthermore, using the same method, it can be seen that

MYOZ2 is highly abundant in the circulatory system, and that PYROXD2 is highly abundant

in the urinary system. The UMAP coordinates and our binned protein abundance data that is

used in these plots to allow for the generation of similar visualisations are provided in

Supplementary File 9.

Figure 8: Visualisations generated using the UMAP algorithm to show the relationships

between human, mouse, and rat samples. (**A**) Shows the relationship of all samples,

particularly showing strong relationship between biological systems. (**B**) Shows the protein

abundancy of 3 example gene orthologs (SH3GL2, MYOZ2 and PYROXD2), within each

sample. Human baseline protein expression data was generated in [14].

2.6. Pathway enrichment analysis

Based on the ortholog protein expression analysis described above, we mapped canonical proteins from mouse and rat to the corresponding ortholog human proteins, which were subsequently subjected to pathway-enrichment analysis using Reactome (Fig. 9). After filtering out the disease and statistically insignificant pathways, there were 2,990 pathways found in all the organs of mouse and 2,162 pathways in all the organs of rat. In mouse samples, the largest number of pathways (367) were found in articular cartilage, and the lowest number of pathways was found in liver (44). We also observed that Neuronal System-related pathways were predominantly present in the brain and eye, which is consistent with expectations. In rat samples, brain included the largest number of pathways (387), while the lowest number of pathways was found in tendon, with 117.

455   **Figure 9.** Pathway analysis performed using the canonical proteins, showing the statistically

456   significant representative pathways (p-value < 0.05) in (A) mouse and (B) rat organs.

3. Discussion

We have previously reported two meta-analysis studies involving the reanalysis and integration in Expression Atlas of public quantitative datasets coming from cell lines and human tumour samples [13], and from human baseline tissues [14], respectively. In this study, we reanalysed mouse and rat baseline proteomics datasets representing protein expression across 34 healthy tissues and 14 organs. We have used the same methodology as in the study involving baseline human tissues, which enabled a comparison of protein expression levels across the three species. Our main overall aim was to provide a system-wide baseline protein expression catalogue across various tissues and organs of mouse and rat and to offer a reference for future related studies.

We analysed each dataset separately using the same software (MaxQuant) and the same search protein sequence database. The disadvantage of this approach is that the FDR statistical thresholds are applied at a dataset level and not to all datasets together as a whole. However, as reported before [14], using a dataset per dataset analysis approach is in our view the only sustainable manner to reanalyse and integrate quantitative proteomics datasets, at least at present. The disadvantage of this approach is that the FDR statistical threshold are applied at a dataset level and not to all datasets together as a whole, with the potential accumulation of false positives across datasets. However, it is important to highlight that the number of commonly detected false positives is reduced in parallel with the increase in the number of common datasets where a given protein is detected. As also reported in previous studies, one of the major bottlenecks was the curation of dataset metadata, consisting of mapping files to samples and biological conditions. Very recently, the MAGE-TAB-Proteomics format has been developed and formalised to enable the reporting of the experimental design in proteomics experience, including the relationship between samples

and raw files, which is recorded in the SDRF-Proteomics section of the file [42]. Submission

of the SDRF-Proteomics files to PRIDE is now supported. The more well-annotated datasets

in the public domain, the easier these data reuse activities will become.

The generated baseline protein expression data can be used with different purposes such as

the generation of protein co-expression networks and/or the inference of protein complexes.

For the latter application, expression data can be alone or for potentially refining predictions

obtained using different methods such as the recently developed AlphaFold-based protein

complexes predictions [43]. Mouse and rat are widely used species in the context of drug

discovery, the latter especially, to undertake regulatory pre-clinical safety studies. Therefore,

it is important to know quantitative protein expression distribution in these species in

different tissues [44] to assist in the selection of species for such studies and also for the

interpretation of the final results.

In addition to the analyses reported, it would have also been possible to perform correlation

studies between gene and protein expression information. However, we did not find any

relevant public datasets in the context of this manuscript where the same samples were

analysed by both techniques, which is the optimal way to perform these studies. Future

directions in analogous studies will involve: (i) additional baseline protein expression studies

of other species, including other model organisms or other species of economic importance;

(ii) the inclusion of differential proteomics datasets (e.g. using TMT and/or iTRAQ); and (iii)

include relevant proteomics expression data coming from the reanalysis of Data Independent

Acquisition (DIA) datasets [45].

As mentioned above, we performed a comparative analysis of baseline protein expression across human, mouse and rat. It was possible to perform this analysis for six common organs (brain, heart, kidney, liver, lung and testis). Ortholog expression across species is useful to infer protein function across experimentally studied proteins. This is particularly useful as evolutionarily closely related species are likely to conserve protein function. We could not find in the literature an analogous comparative study performed at the protein level. However, expression from closely related orthologs across tissues or organs has been compared at the transcriptomics level, providing a complete picture of gene expression. In this context, many studies have compared gene-expression in mouse, rat and human orthologues and found that orthologues had generally a highly correlated expression tissue distribution profile in baseline conditions [46-50]. Gene expression levels among orthologs were found to be highly similar in muscle and heart tissues, liver and nervous system and less similar in epithelial cells, reproductive systems, bone and endocrine organs [48]. Studies have also shown that variability of gene expression between homologous tissues/organs in closely related species can be lower than the variability between unrelated tissues within the same organism [46, 47], in agreement with the results reported here at the protein level. Additionally, we showed an initial analysis of protein expression of orthologs across the three species using UMAP.

In conclusion we here present a meta-analysis study of public mouse and rat baseline proteomics datasets from PRIDE. We demonstrate its feasibility, perform a comparative analysis across the three species and show the main current challenges. Finally, the data is made available *via* Expression Atlas. Whereas there are several analogous studies performed at the gene expression level for mouse and rat tissues, to the best of our knowledge this is the first of this kind at protein expression level.

4. Materials and Methods

4.1. Datasets

As of May 2021, there were 2,060 mouse (*Mus musculus*) and 339 rat (*Rattus norvegicus*) MS proteomics datasets publicly available in the PRIDE database (https://www.ebi.ac.uk/pride/). Datasets were manually selected based on the selection criteria described previously [14]. Briefly, we selected datasets where baseline expression experiments were performed on (i) label-free samples from tissues not enriched for post-translational modifications; (ii) Thermo Fisher Scientific instruments such as LTQ Orbitrap, LTQ Orbitrap Elite, LTQ Orbitrap Velos, LTQ Orbitrap XL ETD, LTQ-Orbitrap XL ETD, Orbitrap Fusion and Q-Exactive, since they represent a large proportion of datasets in PRIDE and to avoid heterogeneity introduced by data from other vendor instruments; (iii) had suitable sample metadata available in the original publication or it was possible to obtain it by contacting the authors; and (iv) our previous experience in the team of some datasets deposited in PRIDE, which were discarded because they were not considered to be useful. Overall, 14 mouse and 9 rat datasets were selected from all mouse and rat datasets for further analysis. Table 1 lists the selected datasets. The 23 datasets contained a total of 211 samples from 34 different tissues across 14 organs (meaning groups of related tissues, more details below), comprising 9 different mouse and 3 rat strains, respectively.

The sample and experimental metadata were manually curated using the information provided in the respective publications or by contacting the original authors/submitters. Annotare [51] was used for annotating the metadata and stored using the Investigation Description Format (IDF) and Sample-Data Relationship Format (SDRF) file formats [42], which are required for integration in Expression Atlas. An overview of the experimental design including experimental factors, protocols, publication information and contact

information are present in the IDF file, and the SDRF includes sample metadata describing

the relationship between the various sample characteristics and the data files contained in the

dataset.

4.2. Proteomics raw data processing

All datasets were analysed with MaxQuant (version 1.6.3.4) [52, 53] on a Linux high-

performance computing cluster for peptide/protein identification and protein quantification.

Input parameters for each dataset, such as MS^1 and MS^2 tolerances, digesting enzymes, fixed

and variable modifications, were set as described in their respective publications, with two

missed cleavage sites. The FDR at the PSM (peptide spectrum match) and protein levels were

set to 1%. The MaxQuant parameters were otherwise set to default values: the maximum

number of modifications per peptide was 5, the minimum peptide length was 7, the maximum

peptide mass was set to 4,600 Da, and for the matches between runs the minimum match time

window was set to 0.7 seconds and the minimum retention time alignment window was set to

20 seconds. The MaxQuant parameter files are available for downloading from Expression

Atlas. The *Mus musculus* UniProt Reference proteome release-2021_04 (including isoforms,

63,656 sequences) and *Rattus norvegicus* UniProt Reference proteome release-2021_04

(including isoforms, 31,562 sequences) were used as the target sequence databases for mouse

and rat datasets, respectively. The built-in contaminant database within MaxQuant was used

and a decoy database was generated by MaxQuant by reversing the input database sequences

after the respective enzymatic digestion. The datasets were run separately in multi-threaded

mode.

4.3. Post-processing

The post-processing of results from MaxQuant is explained in detail in [14]. In brief, the

protein groups labelled as potential contaminants, decoys and those with fewer than 2 PSMs

were removed. Protein intensities in each sample were normalised by scaling the iBAQ

intensity values to the total amount of signal in each MS run and converted to parts per

billion (ppb).

$$ppb\_iBAQ_i = \left(iBAQ_i \middle/ \sum_{i=1}^{n} iBAQ_i \right) x\ 1{,}000{,}000{,}000$$

The 'majority protein identifiers' within each protein group were mapped to their Ensembl

gene identifiers/annotations using the Bioconductor package 'mygene'. For downstream

analysis only protein groups whose isoforms mapped to a single unique Ensembl gene ID

were considered. Protein groups that mapped to more than one Ensembl gene ID are provided

in Supplementary File 1. The protein intensity values from different protein groups with the

same Ensembl gene ID were aggregated as median values. The parent genes to which the

different protein groups were mapped to are equivalent to 'canonical proteins' in UniProt

(https://www.uniprot.org/help/canonical_and_isoforms) and therefore the term protein

abundance is used to describe the protein abundance of the canonical protein throughout the

manuscript.

4.4. Integration into Expression Atlas

The calculated canonical protein abundances (mapped to genes), together with the validated

SDRF files, summary files detailing the quality of post-processing and the input MaxQuant

parameter files (mqpar.xml) were integrated into Expression Atlas

(https://www.ebi.ac.uk/gxa/home) as proteomics baseline experiments (E-PROT identifiers

are available in Table 1).

4.5. Protein abundance comparison across datasets

To compare protein abundances, the normalised protein abundances (in ppb) from each group

of tissues in a dataset were converted into ranked bins. In this study, 'tissue' is defined as a

distinct functional or structural region within an 'organ'. For example, hippocampus,

cerebellum and cortex are defined as 'tissues' that are part of the brain (organ) and similarly

sinus node, left atria, left ventricle, right atria, right ventricle are defined as 'tissues' in heart

(organ). Protein abundances were transformed into bins by first grouping MS runs from each

tissue within a dataset as a batch. The normalised protein abundances (ppb) for each MS run

within a batch were sorted from lowest to highest abundance and ranked into 5 bins. Proteins

whose ppb abundances are ranked in the lowest bin (bin 1) represent lowest abundance and

correspondingly proteins within bin 5 are of highest abundance in their respective tissue.

When merging tissues into organs, median bin values were used.

Proteins that were detected in at least 50% of the samples were selected for PCA (Principal

Component Analysis) and was performed using R (The R Stats package) [54] using binned

abundance values. For generating heatmaps, a Pearson correlation coefficient for all samples

was calculated on pairwise complete observations of bin transformed values. Missing values

were marked as NA (not available). For each organ a median R^2 was calculated from all

pairwise R^2 values of their respective samples. Samples were hierarchically clustered on

columns and rows using Euclidean distances. To compare the correlation in protein

expression of shared organs between datasets, the FOT normalised protein abundances (ppb)

were aggregated by calculating the median over samples. The regression line was computed

using the 'linear model' (lm) method in R.

4.6 Comparison of protein abundances using iBAQ and spectral counting data available

in PaxDB

To compare protein abundances generated from iBAQ in this study and spectral counting

methods, protein abundance data from different mouse organs was obtained from PaxDB

(https://www.pax-db.org/) [16]. FOT normalised iBAQ abundances, as described above, were

compared with the spectral counting abundances for the matching mouse organs. Organs

from mouse labelled as 'integrated' in PaxDB were selected. It was not possible to perform

this comparison for rat organs since data in PaxDB for rat are available for either the 'whole

organism' or for "cell types" only. Abundances were compared across mouse adipose tissue,

brain, heart, kidney, liver, lung, pancreas and spleen. The Ensembl ENSG gene ids were

mapped to ENSP protein ids in PaxDB using the 'mygene' bioconductor package in R.

4.7. UMAP analysis

To generate the UMAP visualisations we used the binned protein abundance values generated

in this study from rat and mouse, as well as the binned human protein abundance values from

[14]. First, we reduced this data to only contain the orthologs found in all three species. For

the purpose of only the initial visualisation layout, we filtered the data to include those

proteins present in 90% of samples. Once the initial layout was generated, we then used the

full protein abundance values to generate protein-specific visualisations. We use R v4.1.0

with the package 'umap' (Uniform Manifold Approximation and Projection in R) [55]

v0.2.7.0 to generate the UMAP visualisations.

4.8. Organ-specific expression profile analysis

For comparison across organs, the tissues were aggregated into organs and their median bin

values were considered. As described previously [14] the classification scheme done by

Uhlén *et al.* [17] was modified to classify the proteins into one of the three categories: (1)

"Organ-enriched": present in one unique organ with bin values 2-fold higher than the mean

bin value across all organs; (2) "Group enriched": present in at least 7 organs in mouse or in

at least 4 organs in rat, with bin values 2-fold higher than the mean bin value across all

organs; and (3) "Mixed": the remaining canonical proteins that are not part of the above two

categories.

Enriched gene ontology (GO) terms analysis was carried out through over-representation test

described previously [14], it was combined with "Organ-enriched" and "Group enriched"

mapped gene lists for each organ. In addition, Reactome [56] pathway analysis was

performed using mapped gene lists and running pathway-topology and over-representation

analysis, as reported previously [14].

4.9. Comparison of protein expression across species

The g:Orth Orthology search function in the g:Profiler suite of programs [57] was used for

translating gene identifiers between organisms. Since a custom list of gene identifiers could

not be used as the background search set, the mouse and rat genes were first mapped against

the background Ensembl database. The resulting list of mouse and rat genes mapped to

human orthologs were then filtered so that they only included parent gene identifiers of the

protein groups from mouse and rat organs identified in this study and the parent genes of

human organs described in our previous study (Supplementary File 2 in [14]), respectively.

The orthologs were grouped into various categories denoting the resulting mapping between

identifiers: "one-to-one", "one-to-many", "many-to-one", "many-to-many", and "no

mappings" between gene identifiers. Only "one-to-one" mapped ortholog identifiers were

used to compare protein intensities between mouse, rat and human organs. The normalised

ppb protein abundances of the one-to-one mapped orthologues in 6 organs (brain, heart,

kidney, liver, lung and testis), that were studied across all three organisms were used to assess

the pairwise correlation of protein abundances. The linear regression was calculated using the

linear fit 'lm' method in R.

Data availability

Expression Atlas E-PROT identifiers and PRIDE original dataset identifiers are included in

Table 1.

Acknowledgements

First of all, we would like to thank all data submitters who made their datasets available via

PRIDE and ProteomeXchange. We would also like to thank Andrew Leach and the rest of the

team involved in the Open Targets "Target Safety" project, for helpful discussions.

Authors' contributions

SW, DGS, AP, DJK selected and curated the datasets. SW, AP, DGS performed analyses. AC

and ARJ helped in the interpretation of results and designed approach for data normalisation.

NG, SF, PM, and IP helped in the integration of the results into Expression Atlas. SW, AP,

DGS, JAV wrote the manuscript. JAV, ARJ and IP obtained the funding for performing the

study. All authors have read and approved the manuscript.

Funding

AP and DGS were funded by Open Targets (project OTAR-02-068),

https://www.opentargets.org/. AP, IP, NG, PM, SF and JAV were funded by BBSRC

BB/T019670/1, https://bbsrc.ukri.org/. AC and ARJ were funded by BBSRC BB/T019557/1,

https://bbsrc.ukri.org/. AP, DGS, SW and JAV were funded by Wellcome Trust [grant

698 number 208391/Z/17/Z], https://wellcome.org/. DJK, IP and JAV were funded by EMBL

699 core funding, https://www.embl.org/. The funders had no role in study design, data collection

700 and analysis, decision to publish, or preparation of the manuscript.

702 **Supporting information**

703 **Supplementary file 1:** Protein groups from all datasets that are mapped to more than one

704 Ensembl Gene ID.

705 **Supplementary file 2:** Median protein abundances (in ppb) for each protein group across

706 various tissue samples in each organ.

707 **Supplementary file 3:** Median binned protein abundances across various tissue samples in

708 each organ of mouse and rat.

709 **Supplementary file 4:** Supplementary figures (S1) illustrating correlation of protein

710 abundances in organs represented in different datasets. (S2) Correlation of protein

711 abundances generated using iBAQ and spectral counting methods in various mouse organs.

712 (S3) Correlation of protein expression between organs within human, mouse and rat.

713 **Supplementary file 5:** Organ distribution of canonical proteins in mouse and rat.

714 **Supplementary file 6:** Gene Ontology enrichment analysis of 'organ-enriched' and 'group-

715 enriched' proteins.

716 **Supplementary file 7:** Binned protein abundances of one-to-one mapped orthologs across all

717 organs studied.

718  **Supplementary file 8:** Figure illustrating binned protein abundances of all one-to-one

719 mapped orthologs across six common organs in mouse, rat and human.

720 **Supplementary file 9:** UMAP co-ordinates and source data of UMAP analysis.

Abbreviations

DDA: Data Dependent Acquisition

DIA: Data Independent Acquisition

FOT: Fraction of Total

GO: Gene Ontology

iBAQ: intensity-based absolute quantification

iTRAQ: Isobaric tag for relative and absolute quantitation

IDF: Investigation Description Format

MS: Mass Spectrometry

ppb: Parts per billion

PCA: Principal Component Analysis

SDRF: Sample and Data Relationship Format

TMT: Tandem Mass Tagging

UMAP: Uniform Manifold Approximation and Projection

References

References

1.   Aebersold R, Mann M. Mass-spectrometric exploration of proteome structure and
function. Nature. 2016;537(7620):347-55. Epub 2016/09/16. doi: 10.1038/nature19949.
PubMed PMID: 27629641.

2.   Perez-Riverol Y, Csordas A, Bai J, Bernal-Llinares M, Hewapathirana S, Kundu DJ,
et al. The PRIDE database and related tools and resources in 2019: improving support for
quantification data. Nucleic Acids Res. 2019;47(D1):D442-D50. Epub 2018/11/06. doi:
10.1093/nar/gky1106. PubMed PMID: 30395289; PubMed Central PMCID:
PMCPMC6323896.

3.   Vizcaino JA, Deutsch EW, Wang R, Csordas A, Reisinger F, Rios D, et al.
ProteomeXchange provides globally coordinated proteomics data submission and
dissemination. Nat Biotechnol. 2014;32(3):223-6. Epub 2014/04/15. doi: 10.1038/nbt.2839.
PubMed PMID: 24727771; PubMed Central PMCID: PMCPMC3986813.

4.   Martens L, Vizcaino JA. A Golden Age for Working with Public Proteomics Data.
Trends Biochem Sci. 2017;42(5):333-41. Epub 2017/01/26. doi: 10.1016/j.tibs.2017.01.001.
PubMed PMID: 28118949; PubMed Central PMCID: PMCPMC5414595.

5.   Romanov N, Kuhn M, Aebersold R, Ori A, Beck M, Bork P. Disentangling Genetic
and Environmental Effects on the Proteotypes of Individuals. Cell. 2019;177(5):1308-18 e10.
Epub 2019/04/30. doi: 10.1016/j.cell.2019.03.015. PubMed PMID: 31031010; PubMed
Central PMCID: PMCPMC6988111.

6.   Skinnider MA, Foster LJ. Meta-analysis defines principles for the design and analysis
of co-fractionation mass spectrometry experiments. Nat Methods. 2021;18(7):806-15. Epub
2021/07/03. doi: 10.1038/s41592-021-01194-4. PubMed PMID: 34211188.

7.   Ochoa D, Jarnuczak AF, Vieitez C, Gehre M, Soucheray M, Mateus A, et al. The
functional landscape of the human phosphoproteome. Nat Biotechnol. 2020;38(3):365-73.
Epub 2019/12/11. doi: 10.1038/s41587-019-0344-3. PubMed PMID: 31819260; PubMed
Central PMCID: PMCPMC7100915.

8.   Vaudel M, Verheggen K, Csordas A, Raeder H, Berven FS, Martens L, et al.
Exploring the potential of public proteomics data. Proteomics. 2016;16(2):214-25. Epub
2015/10/10. doi: 10.1002/pmic.201500295. PubMed PMID: 26449181; PubMed Central
PMCID: PMCPMC4738454.

9.   Kumar D, Yadav AK, Jia X, Mulvenna J, Dash D. Integrated Transcriptomic-
Proteomic Analysis Using a Proteogenomic Workflow Refines Rat Genome Annotation. Mol
Cell Proteomics. 2016;15(1):329-39. Epub 2015/11/13. doi: 10.1074/mcp.M114.047126.
PubMed PMID: 26560066; PubMed Central PMCID: PMCPMC4762527.

10.   Brunet MA, Brunelle M, Lucier JF, Delcourt V, Levesque M, Grenier F, et al.
OpenProt: a more comprehensive guide to explore eukaryotic coding potential and
proteomes. Nucleic Acids Res. 2019;47(D1):D403-D10. Epub 2018/10/10. doi:
10.1093/nar/gky936. PubMed PMID: 30299502; PubMed Central PMCID:
PMCPMC6323990.

11.   Levitsky LI, Kliuchnikova AA, Kuznetsova KG, Karpov DS, Ivanov MV, Pyatnitskiy
MA, et al. Adenosine-to-Inosine RNA Editing in Mouse and Human Brain Proteomes.
Proteomics. 2019;19(23):e1900195. Epub 2019/10/03. doi: 10.1002/pmic.201900195.
PubMed PMID: 31576663.

12.   Li H, Zhou R, Xu S, Chen X, Hong Y, Lu Q, et al. Improving Gene Annotation of the
Peanut Genome by Integrated Proteogenomics Workflow. J Proteome Res. 2020;19(6):2226-
35. Epub 2020/05/06. doi: 10.1021/acs.jproteome.9b00723. PubMed PMID: 32367721.

13.   Jarnuczak AF, Najgebauer H, Barzine M, Kundu DJ, Ghavidel F, Perez-Riverol Y, et
al. An integrated landscape of protein expression in human cancer. Sci Data. 2021;8(1):115.

Epub 2021/04/25. doi: 10.1038/s41597-021-00890-2. PubMed PMID: 33893311; PubMed
Central PMCID: PMCPMC8065022.

14. Prakash A, García-Seisdedos D, Wang S, Kundu DJ, Collins A, George N, et al. An
integrated view of baseline protein expression in human tissues. bioRxiv.
2021:2021.09.10.459811. doi: 10.1101/2021.09.10.459811.

15. Samaras P, Schmidt T, Frejno M, Gessulat S, Reinecke M, Jarzab A, et al.
ProteomicsDB: a multi-omics and multi-organism resource for life science research. Nucleic
Acids Res. 2020;48(D1):D1153-D63. Epub 2019/10/31. doi: 10.1093/nar/gkz974. PubMed
PMID: 31665479; PubMed Central PMCID: PMCPMC7145565.

16. Wang M, Herrmann CJ, Simonovic M, Szklarczyk D, von Mering C. Version 4.0 of
PaxDb: Protein abundance data, integrated across model organisms, tissues, and cell-lines.
Proteomics. 2015;15(18):3163-8. Epub 2015/02/07. doi: 10.1002/pmic.201400441. PubMed
PMID: 25656970; PubMed Central PMCID: PMCPMC6680238.

17. Uhlen M, Fagerberg L, Hallstrom BM, Lindskog C, Oksvold P, Mardinoglu A, et al.
Proteomics. Tissue-based map of the human proteome. Science. 2015;347(6220):1260419.
Epub 2015/01/24. doi: 10.1126/science.1260419. PubMed PMID: 25613900.

18. Azimifar SB, Nagaraj N, Cox J, Mann M. Cell-type-resolved quantitative proteomics
of murine liver. Cell Metab. 2014;20(6):1076-87. Epub 2014/12/04. doi:
10.1016/j.cmet.2014.11.002. PubMed PMID: 25470552.

19. Deshmukh AS, Murgia M, Nagaraj N, Treebak JT, Cox J, Mann M. Deep proteomics
of mouse skeletal muscle enables quantitation of protein isoforms, metabolic pathways, and
transcription factors. Mol Cell Proteomics. 2015;14(4):841-53. Epub 2015/01/27. doi:
10.1074/mcp.M114.044222. PubMed PMID: 25616865; PubMed Central PMCID:
PMCPMC4390264.

20. Meierhofer D, Halbach M, Sen NE, Gispert S, Auburger G. Ataxin-2 (Atxn2)-Knock-
Out Mice Show Branched Chain Amino Acids and Fatty Acids Pathway Alterations. Mol
Cell Proteomics. 2016;15(5):1728-39. Epub 2016/02/07. doi: 10.1074/mcp.M115.056770.
PubMed PMID: 26850065; PubMed Central PMCID: PMCPMC4858951.

21. Sarver DC, Kharaz YA, Sugg KB, Gumucio JP, Comerford E, Mendias CL. Sex
differences in tendon structure and function. J Orthop Res. 2017;35(10):2117-26. Epub
2017/01/11. doi: 10.1002/jor.23516. PubMed PMID: 28071813; PubMed Central PMCID:
PMCPMC5503813.

22. Duda P, Wojcicka O, Wisniewski JR, Rakus D. Global quantitative TPA-based
proteomics of mouse brain structures reveals significant alterations in expression of proteins
involved in neuronal plasticity during aging. Aging (Albany NY). 2018;10(7):1682-97. Epub
2018/07/22. doi: 10.18632/aging.101501. PubMed PMID: 30026405; PubMed Central
PMCID: PMCPMC6075443.

23. Harman JC, Guidry JJ, Gidday JM. Comprehensive characterization of the adult ND4
Swiss Webster mouse retina: Using discovery-based mass spectrometry to decipher the total
proteome and phosphoproteome. Mol Vis. 2018;24:875-89. Epub 2019/02/05. PubMed
PMID: 30713425; PubMed Central PMCID: PMCPMC6334985.

24. Angelidis I, Simon LM, Fernandez IE, Strunz M, Mayr CH, Greiffo FR, et al. An
atlas of the aging lung mapped by single cell transcriptomics and deep tissue proteomics. Nat
Commun. 2019;10(1):963. Epub 2019/03/01. doi: 10.1038/s41467-019-08831-9. PubMed
PMID: 30814501; PubMed Central PMCID: PMCPMC6393476.

25. Zhao Y, Wilmarth PA, Cheng C, Limi S, Fowler VM, Zheng D, et al. Proteome-
transcriptome analysis and proteome remodeling in mouse lens epithelium and fibers. Exp
Eye Res. 2019;179:32-46. Epub 2018/10/26. doi: 10.1016/j.exer.2018.10.011. PubMed
PMID: 30359574; PubMed Central PMCID: PMCPMC6360118.

26.   Huttlin EL, Jedrychowski MP, Elias JE, Goswami T, Rad R, Beausoleil SA, et al. A
tissue-specific atlas of mouse protein phosphorylation and expression. Cell.
2010;143(7):1174-89. Epub 2010/12/25. doi: 10.1016/j.cell.2010.12.001. PubMed PMID:
21183079.
27.   Linscheid N, Santos A, Poulsen PC, Mills RW, Calloe K, Leurs U, et al. Quantitative
proteome comparison of human hearts with those of model organisms. PLoS Biol.
2021;19(4):e3001144. Epub 2021/04/20. doi: 10.1371/journal.pbio.3001144. PubMed PMID:
33872299; PubMed Central PMCID: PMCPMC8084454.
28.   Dudek M, Angelucci C, Pathiranage D, Wang P, Mallikarjun V, Lawless C, et al.
Circadian time series proteomics reveals daily dynamics in cartilage physiology.
Osteoarthritis Cartilage. 2021;29(5):739-49. Epub 2021/02/22. doi:
10.1016/j.joca.2021.02.008. PubMed PMID: 33610821; PubMed Central PMCID:
PMCPMC8113022.
29.   Schroeder S, Hofer SJ, Zimmermann A, Pechlaner R, Dammbrueck C, Pendl T, et al.
Dietary spermidine improves cognitive function. Cell Rep. 2021;35(2):108985. Epub
2021/04/15. doi: 10.1016/j.celrep.2021.108985. PubMed PMID: 33852843.
30.   Bundy JL, Vied C, Nowakowski RS. Sex differences in the molecular signature of the
developing mouse hippocampus. BMC Genomics. 2017;18(1):237. Epub 2017/03/18. doi:
10.1186/s12864-017-3608-7. PubMed PMID: 28302071; PubMed Central PMCID:
PMCPMC5356301.
31.   Linscheid N, Logantha S, Poulsen PC, Zhang S, Schrolkamp M, Egerod KL, et al.
Quantitative proteomics and single-nucleus transcriptomics of the sinus node elucidates the
foundation of cardiac pacemaking. Nat Commun. 2019;10(1):2889. Epub 2019/06/30. doi:
10.1038/s41467-019-10709-9. PubMed PMID: 31253831; PubMed Central PMCID:
PMCPMC6599035.
32.   Alugubelly N, Mohammed AN, Edelmann MJ, Nanduri B, Sayed M, Park JW, et al.
Adolescent rat social play: Amygdalar proteomic and transcriptomic data. Data Brief.
2019;27:104589. Epub 2019/11/02. doi: 10.1016/j.dib.2019.104589. PubMed PMID:
31673590; PubMed Central PMCID: PMCPMC6817652.
33.   Roffia V, De Palma A, Lonati C, Di Silvestre D, Rossi R, Mantero M, et al. Proteome
Investigation of Rat Lungs subjected to Ex Vivo Perfusion (EVLP). Molecules. 2018;23(12).
Epub 2018/11/24. doi: 10.3390/molecules23123061. PubMed PMID: 30467300; PubMed
Central PMCID: PMCPMC6321151.
34.   Bernier M, Harney D, Koay YC, Diaz A, Singh A, Wahl D, et al. Elucidating the
mechanisms by which disulfiram protects against obesity and metabolic syndrome. NPJ
Aging Mech Dis. 2020;6:8. Epub 2020/07/28. doi: 10.1038/s41514-020-0046-6. PubMed
PMID: 32714562; PubMed Central PMCID: PMCPMC7374720.
35.   Huang Q, Luo L, Alamdar A, Zhang J, Liu L, Tian M, et al. Integrated proteomics
and metabolomics analysis of rat testis: Mechanism of arsenic-induced male reproductive
toxicity. Sci Rep. 2016;6:32518. Epub 2016/09/03. doi: 10.1038/srep32518. PubMed PMID:
27585557; PubMed Central PMCID: PMCPMC5009432.
36.   Kaushik G, Spenlehauer A, Sessions AO, Trujillo AS, Fuhrmann A, Fu Z, et al.
Vinculin network-mediated cytoskeletal remodeling regulates contractile function in the
aging heart. Sci Transl Med. 2015;7(292):292ra99. Epub 2015/06/19. doi:
10.1126/scitranslmed.aaa5843. PubMed PMID: 26084806; PubMed Central PMCID:
PMCPMC4507505.
37.   Vileigas DF, Harman VM, Freire PP, Marciano CLC, Sant'Ana PG, de Souza SLB, et
al. Landscape of heart proteome changes in a diet-induced obesity model. Sci Rep.
2019;9(1):18050. Epub 2019/12/04. doi: 10.1038/s41598-019-54522-2. PubMed PMID:
31792287; PubMed Central PMCID: PMCPMC6888820.

38.	Limbutara K, Chou CL, Knepper MA. Quantitative Proteomics of All 14 Renal Tubule Segments in Rat. J Am Soc Nephrol. 2020;31(6):1255-66. Epub 2020/05/03. doi: 10.1681/ASN.2020010071. PubMed PMID: 32358040; PubMed Central PMCID: PMCPMC7269347.

39.	Devaux S, Cizkova D, Quanico J, Franck J, Nataf S, Pays L, et al. Proteomic Analysis of the Spatio-temporal Based Molecular Kinetics of Acute Spinal Cord Injury Identifies a Time- and Segment-specific Window for Effective Tissue Repair. Mol Cell Proteomics. 2016;15(8):2641-70. Epub 2016/06/03. doi: 10.1074/mcp.M115.057794. PubMed PMID: 27250205; PubMed Central PMCID: PMCPMC4974342.

40.	Choi H, Simpson D, Wang D, Prescott M, Pitsillides AA, Dudhia J, et al. Heterogeneity of proteome dynamics between connective tissue phases of adult tendon. Elife. 2020;9. Epub 2020/05/13. doi: 10.7554/eLife.55262. PubMed PMID: 32393437; PubMed Central PMCID: PMCPMC7217697.

41.	Hodge K, Have ST, Hutton L, Lamond AI. Cleaning up the masses: exclusion lists to reduce contamination with HPLC-MS/MS. J Proteomics. 2013;88:92-103. Epub 2013/03/19. doi: 10.1016/j.jprot.2013.02.023. PubMed PMID: 23501838; PubMed Central PMCID: PMCPMC3714598.

42.	Dai C, Fullgrabe A, Pfeuffer J, Solovyeva EM, Deng J, Moreno P, et al. A proteomics sample metadata representation for multiomics integration and big data analysis. Nat Commun. 2021;12(1):5854. Epub 2021/10/08. doi: 10.1038/s41467-021-26111-3. PubMed PMID: 34615866; PubMed Central PMCID: PMCPMC8494749.

43.	Evans R, O'Neill M, Pritzel A, Antropova N, Senior A, Green T, et al. Protein complex prediction with AlphaFold-Multimer. bioRxiv. 2021:2021.10.04.463034. doi: 10.1101/2021.10.04.463034.

44.	Bregman CL, Adler RR, Morton DG, Regan KS, Yano BL, Society of Toxicologic P. Recommended tissue list for histopathologic examination in repeat-dose toxicity and carcinogenicity studies: a proposal of the Society of Toxicologic Pathology (STP). Toxicol Pathol. 2003;31(2):252-3. Epub 2003/04/17. doi: 10.1080/01926230390183751. PubMed PMID: 12696587.

45.	Walzer M, García-Seisdedos D, Prakash A, Brack P, Crowther P, Graham RL, et al. Implementing the re-use of public DIA proteomics datasets: from the PRIDE database to Expression Atlas. bioRxiv. 2021:2021.06.08.447493. doi: 10.1101/2021.06.08.447493.

46.	Sollner JF, Leparc G, Hildebrandt T, Klein H, Thomas L, Stupka E, et al. An RNA-Seq atlas of gene expression in mouse and rat normal tissues. Sci Data. 2017;4:170185. Epub 2017/12/13. doi: 10.1038/sdata.2017.185. PubMed PMID: 29231921; PubMed Central PMCID: PMCPMC5726313.

47.	Sudmant PH, Alexis MS, Burge CB. Meta-analysis of RNA-seq expression data across species, tissues and studies. Genome Biol. 2015;16:287. Epub 2015/12/24. doi: 10.1186/s13059-015-0853-4. PubMed PMID: 26694591; PubMed Central PMCID: PMCPMC4699362.

48.	Zheng-Bradley X, Rung J, Parkinson H, Brazma A. Large scale comparison of global gene expression patterns in human and mouse. Genome Biol. 2010;11(12):R124. Epub 2010/12/25. doi: 10.1186/gb-2010-11-12-r124. PubMed PMID: 21182765; PubMed Central PMCID: PMCPMC3046484.

49.	Liao BY, Zhang J. Evolutionary conservation of expression profiles between human and mouse orthologous genes. Mol Biol Evol. 2006;23(3):530-40. Epub 2005/11/11. doi: 10.1093/molbev/msj054. PubMed PMID: 16280543.

50.	Prasad A, Kumar SS, Dessimoz C, Bleuler S, Laule O, Hruz T, et al. Global regulatory architecture of human, mouse and rat tissue transcriptomes. BMC Genomics.

2013;14:716. Epub 2013/10/22. doi: 10.1186/1471-2164-14-716. PubMed PMID: 24138449;
PubMed Central PMCID: PMCPMC4008137.
51. Athar A, Fullgrabe A, George N, Iqbal H, Huerta L, Ali A, et al. ArrayExpress update
- from bulk to single-cell expression data. Nucleic Acids Res. 2019;47(D1):D711-D5. Epub
2018/10/26. doi: 10.1093/nar/gky964. PubMed PMID: 30357387; PubMed Central PMCID:
PMCPMC6323929.
52. Cox J, Mann M. MaxQuant enables high peptide identification rates, individualized
p.p.b.-range mass accuracies and proteome-wide protein quantification. Nat Biotechnol.
2008;26(12):1367-72. Epub 2008/11/26. doi: 10.1038/nbt.1511. PubMed PMID: 19029910.
53. Tyanova S, Temu T, Cox J. The MaxQuant computational platform for mass
spectrometry-based shotgun proteomics. Nat Protoc. 2016;11(12):2301-19. Epub 2016/11/04.
doi: 10.1038/nprot.2016.136. PubMed PMID: 27809316.
54. Team RC. R: A language and environment for statistical computing. R
Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0, URL
http://www.R-project.org/. 2013.
55. McInnes L HJ, Melville J. UMAP: Uniform Manifold Approximation and Projection
for dimension reduction. arXiv. 2018.
56. Wu G, Haw R. Functional Interaction Network Construction and Analysis for Disease
Discovery. Methods Mol Biol. 2017;1558:235-53. Epub 2017/02/06. doi: 10.1007/978-1-
4939-6783-4_11. PubMed PMID: 28150241.
57. Raudvere U, Kolberg L, Kuzmin I, Arak T, Adler P, Peterson H, et al. g:Profiler: a
web server for functional enrichment analysis and conversions of gene lists (2019 update).
Nucleic Acids Res. 2019;47(W1):W191-W8. Epub 2019/05/09. doi: 10.1093/nar/gkz369.
PubMed PMID: 31066453; PubMed Central PMCID: PMCPMC6602461.
958