Title: Mixture Density Regression reveals frequent recent adaptation in the human

genome

Authors: Diego F. Salazar-Tortosa[1,2], Yi-Fei Huang[3,4], David Enard[1]

Affiliations:

[1]Department of Ecology and Evolutionary Biology, University of Arizona, Tucson, AZ,

USA.

[2]PROFITH 'PROmoting FITness and Health through physical activity' research group,

Sport and Health University Research Institute (iMUDS), University of Granada,

Granada, Spain.

[3]Department of Biology, Pennsylvania State University, University Park, PA, USA.

[4]Huck Institutes of the Life Sciences, Pennsylvania State University, University Park,

PA, USA.

Corresponding Authors: David Enard: denard@email.arizona.edu; Yi-Fei Huang:

yuh371@psu.edu

Running head: MDR-based analysis of positive selection.

Keywords: human evolution, recent adaptation, mixture distributions, selection

determinants, recombination rate.

Abstract

How much genome differences between species reflect neutral or adaptive evolution is a central question in evolutionary genomics. In humans and other mammals, the prevalence of adaptive versus neutral genomic evolution has proven particularly difficult to quantify. The difficulty notably stems from the highly heterogenous organization of mammalian genomes at multiple levels (functional sequence density, recombination, etc.) that complicates the interpretation and distinction of adaptive vs. neutral evolution signals. Here, we introduce Mixture Density Regressions (MDRs) for the study of the determinants of recent adaptation in the human genome. MDRs provide a flexible regression model based on multiple Gaussian distributions. We use MDRs to model the association between recent selection signals and multiple genomic factors likely to affect positive selection, if the latter was common enough in the first place to generate these associations. We find that a MDR model with two Gaussian distributions provides an excellent fit to the genome-wide distribution of a common sweep summary statistic (iHS), with one of the two distributions likely capturing the positively selected component of the genome. We further find several factors associated with recent adaptation, including the recombination rate, the density of regulatory elements in immune cells and testis, GC-content, gene expression in immune cells, the density of mammal-wide conserved elements, and the distance to the nearest virus-interacting gene. These results support that strong positive selection was relatively common in recent human evolution and highlight MDRs as a powerful tool to make sense of signals of recent genomic adaptation.

Introduction

The characteristics of the human genome can influence both the occurrence and the detection of recent positive selection in a specific genomic region. Similar to other mammals, the human genome has a complex organization, with highly heterogeneous recombination rates and distribution of functional elements (regulatory or coding) along

chromosomes. This inherent heterogeneity in the factors that are likely to influence both

the occurrence of positive selection and our ability to detect it, has complicated the study

of the prevalence of recent adaptation in the human genome, and has made the topic

controversial (Enard et al. 2014; Hernandez et al. 2011). Even though there is growing

evidence that recent positive selection in the form of selective sweeps may have been

relatively common during recent human evolution (Enard and Petrov 2020; Souilmi et al.

2021), further resolution of the controversy will depend on our ability to understand what

factors govern the local genomic rate of recent adaptation along human chromosomes.

Indeed, if signatures of selective sweeps do not occur randomly, as expected if all sweep

signals are false positives only reflecting genetic drift, but instead associate non-

randomly with functional elements in the genome, then this provides evidence that

sweeps had to be common enough in the first place to generate the association.

What factors then are *a priori* expected to matter for recent adaptation? Among

all genomic factors, recombination is a key process determining the patterns of linkage

between alleles across the human genome (The International HapMap Consortium 2005),

that can strongly influence the probability to detect recent selection in the form of

selective sweeps. The lower the recombination rate, the larger the genomic region where

neutral variants will hitchhike to higher frequencies along with an advantageous variant,

before recombination breaks down their linkage (Sabeti et al. 2006; Pritchard et al. 2010).

The higher the recombination rate, the smaller the genomic region affected by the sweep.

Different sweep sizes in low and high recombination regions result in a higher statistical

power to detect sweeps in low than in high recombination regions (O'Reilly et al. 2008;

Booker et al. 2020; Johnson and Voight 2018). In parallel, larger neutral haplotypes make

low recombination regions more prone to detect false positive sweeps (O'Reilly et al.

2008; Booker et al. 2020). This heterogeneity, together with other factors such as

background selection, has greatly contributed to a persistent debate around the prevalence

of selective sweeps in the human genome.

If recent adaptation and selective sweeps in particular were relatively common

during recent human evolution, we expect that other factors on top of recombination

should be relevant for their occurrence and detection in the human genome. Generally,

we expect that selective sweeps should occur more frequently around functional

segments of the genome where adaptive mutations are expected to take place (Enard et al.

2014). Therefore, we should find enough signals of positive association between selective

sweeps and the overall density of functional elements, either coding or non-coding. We

also expect specific functions to be associated with an increased occurrence of selective

sweeps. For example, the presence of genes coding for proteins that interact with

pathogens, and viruses in particular (Virus-Interacting Proteins or VIPs), should influence

the frequency of sweeps in a given genomic region, independently of other genomic

features (Enard and Petrov 2020; Souilmi et al. 2021). More generally, genomic regions

with immune genes are expected to have experienced more sweeps (Barreiro et al. 2009;

Enard et al. 2016; Deschamps et al. 2016; Enard and Petrov 2018). Similarly,

reproduction-related functions show signals of positive selection (Voight et al. 2006;

Nielsen et al. 2005), thus tissues related with these functions should be associated with

the frequency of sweeps.

Several summary statistics are currently available to detect recent genomic

adaptation. Among all possible choices, the statistics that use the structure of haplotypes

along chromosomes are the most appropriate to investigate the prevalence and

determinants of recent positive selection. First, haplotype-based statistics have good

statistical power to detect strong and recent incomplete sweeps (Voight et al. 2006;

Ferrer-Admetlla et al. 2014; Garud et al. 2015; Enard and Petrov 2020). Second, they are

not confounded by other processes such as background selection (BGS) (Enard et al.

2014; Schrider 2020). This insensitivity to BGS is a particularly important attribute when

testing the specific association between recent positive selection and multiple genomic

factors that can also correlate with BGS. In this regard, the integrated Haplotype Score

(iHS) (Voight et al. 2006) is a haplotype-based statistic that has been extensively tested

(Voight et al. 2006; Barreiro et al. 2009; Enard et al. 2014; Johnson and Voight 2018),

and can detect selection signals both from *de novo* mutations and standing genetic

variation, provided that selection started from not too high initial allele frequency (Enard

and Petrov 2020). This summary statistic can be used to scan many genomes given the

availability of fast implementations (Maclean et al. 2015).

Here, we model the genomic determinants of recent positive selection in the

human genome. We measure recent selective sweeps with the iHS statistic from five

human populations represented in the 1,000 Genomes phase 3 dataset (The 1000

Genomes Project Consortium 2015). The iHS statistic has more power to detect recent

and incomplete sweeps, while it has reduced power to detect complete sweeps or sweeps

more than 30,000 years old (Sabeti et al. 2006; Enard and Petrov 2020). Therefore, we

restrict our analyses to selection that occurred after the main human migration out of

Africa (Henn et al. 2012). We use Mixture Density Regressions (MDRs) to test the

association between iHS and several genomic factors that are possible determinants of

selection. MDRs can fit a mixture of several distributions to the observed statistic. In our

case, we fit a mixture of two Gaussian distributions to the observed iHS, expecting that

one distribution matches the drift component of iHS, while the second one fits its selected

component. Then, we look at the genomic factors that influence iHS in the regression

model, but only considering the selected component (Methods). This differs from other

approaches trying to understand the determinants of selection in two important ways.

First, previous studies have focused on the detection of partial correlations between

selection and single genomic factors (e.g., Lohmueller et al. (2011)). This approach is

likely limited to model the impact of multiple genomic factors that could influence

selection simultaneously. Second, they analyze the overall correlation without

distinguishing between the selected and non-selected components of the genome.

Therefore, the true impact of positive selection could be diluted among values of iHS that

are not influenced by selective sweeps (i.e., drift component). Given that the impact of

hitchhiking on the human genome is likely localized and not expected to influence the

entire genome, an approach specifically distinguishing a selected component should have

more power to reveal the impact of positive selection.

As expected according to previous studies, we find a strong association between

recombination rate and recent positive selection signals in the human genome. In addition

to recombination, we find that other factors, such as the density of regulatory elements in

immune cells and testis, GC-content, gene expression in immune cells, the density of

mammal-wide conserved elements and the distance to VIPs, all show strong associations

with recent positive selection. These results imply that recent positive selection had to be

common enough to create these associations, and further highlight how Mixed Density

Regressions (MDRs) can be used to clarify the prevalence and determinants of recent

positive selection based on genome scans.

Results

We consider multiple genomic factors to model the association with recent selection

signals. These factors are possible determinants of selection according to previous

evidence (Siepel et al. 2005; Nielsen et al. 2005; O'Reilly et al. 2008; Enard et al. 2014;

Luisi et al. 2015; Booker et al. 2020; Enard and Petrov 2020; Voight et al. 2006). We use

a gene-centric perspective where each individual protein-coding gene in the human

genome is associated with a set of factors measured by centering a genomic window on

each gene (Methods). This approach is likely to detect the influence of factors that differ

between genes, rather than only differences between genic and intergenic regions. The set

of factors in the MDR model includes several genomic features, like the length of the

gene at the center of the window or recombination estimates from the latest deCODE

genetic map (Methods). We obtain other genomic factors, like the GC-content and the

densities of genes, coding sequences and conserved elements. Moreover, we include the

density of transcription factors binding sites according to the hypersensitivity to DNaseI

and chromatin immunoprecipitation (ChIP-seq) experiments (The ENCODE Project

Consortium 2012). In the latter case, we also obtain the density of regulatory elements for

specific subsets of cell lines: testis and immune cells. Table 1 provides a complete list of

the factors considered, and the Methods section details how these factors were obtained.

We also consider other functional factors like gene expression, including the

average gene expression across 53 GTEx v7 tissues (GTEx Consortium 2015). The

expression in testis and immune cells could impact the frequency of sweeps, thus we also

include them as independent expression variables (Nielsen et al. 2005; Voight et al.

2006). We also consider the number of protein-protein interactions (PPIs) of each gene,

given that this variable has been previously associated with the rate of sweeps (Luisi et al.

2015). Finally, we consider the influence of viruses on genomic adaptation, as previous

evidence suggests that they have acted as a strong selective pressure during recent human

evolution (Enard and Petrov 2020; Souilmi et al. 2021). We use the distance of each gene

to the closest gene coding for a protein that interacts with viruses. More details for all the

factors considered are given in the Methods.

Genomic factors that are calculated across the genomic windows can vary

depending on the window size used. Therefore, we infer the association between recent

selection and factors measured in genomic windows of varying sizes (Methods). For

example, we measure the association between the average iHS within 50kb windows and

the average recombination rate within the same 50kb windows, or between the average

iHS and recombination within the same 1000kb windows (Methods). Using different

window sizes also relaxes assumptions about the expected strength of selection, as larger

windows are likely more sensitive to strong selection compared to smaller windows, and

vice versa (Enard and Petrov 2020). In total, we use five different window sizes (50, 100,

200, 500 and 1,000 kb) centered at the genomic center of Ensembl v99 coding genes

(Yates et al. 2020). Using fixed size windows irrespective of gene length avoids a priori

biasing measures of iHS by gene length. We also account for gene length in our model

through the inclusion of multiple functional densities along with the genomic length of

the gene (between transcription start and stop) at the center of the window as potential

explanatory variables.

We estimate recent selection signals using the iHS statistic in five populations

represented in the 1,000 Genomes Project (The 1000 Genomes Project Consortium

2015). Each population represents a different continent: Yoruba for Africa, Han Chinese

for East Asia, Utah residents with Northern and Western European ancestry, Toscani for

Southern Europe, and Peruvians for Americas. We select Peruvians given that they show

the highest percentage of Native American ancestry among populations included in the

1,000 Genomes Project (The 1000 Genomes Project Consortium 2015; Harris et al.

2018). The iHS statistic measures how far the haplotypes carrying a specific derived

allele extend upstream and downstream of this focal allele, compared to how far

haplotypes carrying the ancestral allele extend (Voight et al. 2006). As such, the approach

provides one value of iHS for each biallelic Single Nucleotide Polymorphism (SNP) in

the genome with clearly defined derived and ancestral alleles (Methods). Although

isolated extreme values of iHS at a single SNP have often been used as a signature of

selection, we recently found that measuring the average absolute value of iHS across all

the SNPs in a genomic window provides high power to detect recent selection (Enard and

Petrov 2020). The average iHS in an entire window is also likely to better associate with

other factors similarly measured as averages in the same windows. To account for the

fact that the variance of the average iHS depends on the number of SNPs with individual

iHS values in a window, we include this number as a predictor in the tested MDR

models.

MDR fit to recent selection signals

For each separate human population and window size, we model the association between

genomic factors and iHS using a MDR with a mixture of two Gaussian distributions. We

design one distribution to fit the non-selective, drift component of iHS that is not

expected to correlate with genomic factors. In contrast, the second distribution is

designed to fit the selective component of iHS and it is expected to associate with the

tested genomic factors (Methods). We apply this approach to account for the fact that

only a part of the genome is expected to have undergone recent positive selection. In this

context, a very small second distribution and non-significant associations with the tested

predictive factors would suggest a very small contribution of recent positive selection.

The patterns of iHS would then be entirely dominated by genetic drift and past

demography. We use this modeling approach together with an optimization algorithm to

get the overall mixture distribution with the best fit for the observed iHS in each

population and window size (Methods).

Overall, the MDR models fit well the distribution of observed iHS (Figure 1). iHS

does not follow a Gaussian distribution, and our flexible approach combining a mixture

of two Gaussian distributions and linear regression clearly fits better the iHS distribution

than would a single distribution from a classic linear model. We find a clear separation

between the selected and non-selected components, with the selected component shifted

toward higher iHS values (Figure 1). Together, the close fit and separation of the two

gaussian distributions likely provide an improved ability of our model to estimate the

contributions of different factors to iHS.

Patterns of selection across populations and window sizes

We find different patterns of selection across five populations from four different

continents (Figure 1). The African Yoruba population shows the largest selected

component (Figure 1A). The selected component is clearly shifted towards high iHS

values, indicating stronger selection signals that stand out more from the non-selected

component. The selection components are less pronounced but still substantial in the four

other populations (Figure 1B-1E). The Peruvian population shows the smallest selected

component (Figure 1E). These results suggest that the visible (not necessarily the actual)

contribution of selection to the whole variation of iHS is smaller in populations that were

exposed to bottlenecks during migrations Out of Africa and consequently, more subject

to genetic drift (Henn et al. 2012; The 1000 Genomes Project Consortium 2015). In other

words, our results might be explained by the fact that selection is more visible in Yoruba

due to a reduced effect of genetic drift.

With respect to window size, we find the largest selective components when using

larger, 1,000 kb windows (Supplementary Data 1). This is an expected result, because it

is easier to distinguish strong selection from genetic drift and background noise in larger

windows. Smaller windows are less sensitive to strong selection, while larger windows

can include a great accumulation of large iHS values. In addition, small windows are

more influenced not only by the content of a genomic window, but also by genetically

linked surrounding regions. These surrounding areas are in linkage disequilibrium with

the genomic areas inside smaller windows as their center is closer to the edges.

Therefore, smaller windows do not differentiate well between inside and outside genomic

regions (Enard et al. 2014; Enard and Petrov 2020). Given these results, we mainly focus

the rest of our analyses on the Yoruba population and 1,000 kb windows.

Figure 1. Mixture of Gaussian distributions fitting observed iHS (1,000 kb windows)
for the five studied populations: A) Africa - Yoruba; B) Europe - Utah residents with
Northern and Western European ancestry; C) Europe - Toscani; D) East Asia - Han
Chinese; E) America - Peruvians. The figure shows the non-selected and selected
components (Component 1 and 2, respectively) for each population. In the latter
component, iHS linearly depends on the genomic factors considered. Legend: Light blue
= Observed iHS; Dark blue = Mixture model; Full red curve = Component 1 of the
mixture model without selection; Dashed red curve = Component 2 of the mixture model
with selection.

Determinants of strong positive selection in the Yoruba population

Several genomic factors show significant associations with iHS in the selected

component (Table 1). Unsurprisingly, we find the strongest association with

recombination rate, which negatively correlates with iHS (slope=-2.44; p-value<1E-16;

Table 1). This is congruent with the eroding effect of recombination on selection

signatures. Recombination can break haplotypes generated by selection, making it more

difficult to detect the selective events (O'Reilly et al. 2008; Booker et al. 2020). These

results confirm the role of recombination as a key determinant of the probability of

detecting selection with iHS (Johnson and Voight 2018). We also find an association

between GC-content and recent positive selection. When the effect of recombination is

not accounted for, that is, recombination is not included in the model, GC-content

negatively correlates with iHS (slope=-0.29; p-value=1.22E-03). This is expected given

that GC-content is positively associated with long-term recombination (Duret and Arndt

2008). However, adding the control for recombination reveals a positive association

between GC-content and iHS (slope=0.55; p-value=1.34E-06; Table 1). Given that

recombination already explains a great proportion of iHS variability including that shared

with GC-content, the positive association between GC-content and iHS could be

independent from recombination. Accounting for the eroding effect of recombination on

haplotypes could make more visible a positive (direct or indirect) influence of GC-

content on selection. Indeed, this association remains significant when considering only

low-recombination regions, further suggesting its independence from the eroding effect

of recombination (see below).

GC-content might be a better proxy of overall functional density than individual

functional factors included in the regression model, which could explain its positive

association with iHS. We test this hypothesis by removing different genomic factors from

the model (see Supplementary Data 2 for modeling results after removing different sets

of factors). The removal of GC-content increases the association of iHS with both coding

density (slope=0.22 vs 0.16; p-value=1.32E-02 vs. 8.41E-02) and DNAseI

hypersensitivity, which is a measure of the overall density of transcription-factor binding

sites (slope=0.44 vs. -0.047; p-value=1.77E-07 vs. 7.15E-01). Since GC-content has been

positively associated with the density of coding sequences and the hypersensitivity to

DNAseI (Lander et al. 2001; Di Filippo and Bernardi 2008), removing GC-content may

thus make more visible a positive association between iHS and functional densities. We

further test if this is also the case for more tissue-specific functional densities such as the

density of regulatory elements in testis or immune cells that are expected to exhibit more

positive selection (Methods), but we find no such evidence. The removal of GC-content

does not affect to the association between selection and regulatory density in testis

(slope=-0.765 vs. -0.795; p-value<1E-16 in both cases) and immune cells (slope=0.072

vs. 0.047; p-value=4.5E-01 vs. 6.23E-01). As shown later in this manuscript, fine-scale

patterns of recombination seem to hinder the detection of associations for these

regulatory variables. Finally, we find that the removal of coding density and DNAseI

hypersensitivity increases the significance for the association between GC-content and

iHS (p-value=2.40E-14 vs. 1.34E-06). From this model, we then further remove

additional functional factors whose association with selection is also affected by the

removal of GC-content, namely, gene number, gene length and ChIP-seq regulatory

density (Methods). Accordingly, the removal of these functional factors leads to a highly

significant association between GC-content and recent positive selection (slope=0.49; p-

value<1E-16). In summary, the removal of different functional genomic factors from the

model supports that the positive association between positive selection and GC-content is

mediated by the role of this factor as proxy of overall functional density. However, we

cannot exclude the implication of other processes related to recombination that would

require more detailed analyses (for example, see fine scale patterns related to regulatory

density below).

Table 1. Slopes and p-values of the association between iHS and genomic factors for

the Yoruba population in 1,000 kb windows.

Covariate	Slope	p-value
Intercept	-1.426	<1E-16
Recombination rate	-2.435	<1E-16

GC-content	0.55	1.34E-06
Density of conserved elements	0.159	6.40E-04
Number PPIs	-0.07	4.25E-02
Distance to VIPs	-0.17	4.16E-06
Gene expression	-0.139	3.77E-02
Gene expression in immune cells	0.265	1.51E-05
Gene expression in testis	0.001	9.90E-01
Regulatory density (ChIP-seq)	-0.147	2.37E-01
Regulatory density in immune cells (ChIP-seq)	0.047	6.24E-01
Regulatory density in testis (ChIP-seq)	-0.795	<1E-16
Regulatory density (DNaseI)	-0.047	7.15E-01
Gene length	-0.059	1.12E-01
Gene number	-0.086	3.46E-01
Coding density	0.159	8.41E-02
Number iHS data points	-0.007	8.19E-01

PPIs: Protein-Protein Interactions. VIPs: Virus Interacting Proteins. ChIP-seq: Density of
regulatory elements according to chromatin immunoprecipitation experiments. DNaseI:
Density of regulatory elements according to the hypersensitivity to DNaseI. iHS:
Integrated Haplotype Score.

Another genomic factor associated with selection is the density of conserved

elements (Methods), showing a positive association with iHS (slope=0.16; p-value=6.4E-

04; Table 1). This is an expected result since coding sequences and non-coding regulatory

elements tend to be conserved (Siepel et al. 2005). The density of conserved elements is

thus likely to be a good proxy of overall (coding and non-coding) functional density. The

inclusion of this factor in the model could then, however, explain the lack of association

for other factors related to functional density such as coding density. Indeed, the removal

of conserved elements density makes significant the positive association between coding

density and selection (slope=0.26 vs. 0.16; p-value=2.35E-03 vs. 8.41E-02;

Supplementary Data 2). In addition, the simultaneous removal of conserved elements

density and GC-content, which can also act as a proxy of functional density, increases

even more the positive association of coding density with iHS (slope=0.29; p-

value=7.76E-04). These, and previous results about GC-content suggest that the lack of

association for multiple functional factors could be explained, at least partially, by the

simultaneous consideration of better measures of overall functional density in our
models.

The number of protein-protein interactions is another factor that correlates with
iHS. This factor associates negatively with selection, suggesting the existence of a slight
depletion of selection for regions with higher number of PPIs (slope=-0.07; p-
value=4.25E-02; Table 1). This is opposite to that found by Luisi et al. (2015), as they
reported higher recent positive selection on central proteins within the human
interactome. Higher number of PPIs has been also associated with higher positive
selection in the chimpanzee lineage according to the McDonald-Kreitman (MK) test.
However, this factor does not seems to be a determinant of positive selection when other
genomic factors are simultaneously considered using a MK regression approach (Huang
2021). Given the inconsistency with previous evidence and the low strength of the
association, the weak negative relationship between selection and the number of PPIs
found in this study should be taken with caution.

The modeling approach presented in this study can also be used to test the
influence of specific selective pressures. We illustrate this feature by showing that
positive selection is associated with multiple variables related to viral interaction (Table
1). One of the factors more highly associated with iHS is the distance to VIPs (slope=-
0.17; p-value=4.16E-06). Selection decreases further away from VIPs, in other words, we
find an enrichment of selection around VIPs. Viruses have acted as a strong selective
pressure during human evolution, shaping genomic adaptation, and previous studies have
also found more positive selection at VIPs (Enard and Petrov 2020; Souilmi et al. 2021).
Gene expression is another functional factor significantly associated with selection (Table
1). The average expression across 53 tissues from GTEx (GTEx Consortium 2015) is
only slightly and negatively associated with iHS (slope=-0.14; p-value=3.77E-02), while
gene expression in immune cells shows a strong and positive association with selection
(slope=0.26, p-value=1.51E-05). Given their relevance in the response to pathogens,
genomic regions highly expressed in immune cells do represent expected targets of
positive selection.

The densities of regulatory elements in tissues expected to experience more

positive selection (testis and immune cells) show surprising patterns of association with

iHS (Table 1). In particular, a higher density of transcription factor binding sites in testis

correlates with weaker iHS selection signals (slope=-0.79; p-value<1E-16). This is a

counterintuitive result, given that more regulatory density would give more options to

modulate gene expression and hence more room for positive selection to act. An

explanation could be the following: hotspots of recombination are usually regions where

the chromatin is more open, and hence more accessible for transcription factors to bind

(Spruce et al. 2020). If the density of transcription factor binding sites in testis (where

meiosis and recombination occur) is higher in areas with high recombination, then

selection signals within these sites would be erased by recombination (O'Reilly et al.

2008; Booker et al. 2020). Fine scale patterns of recombination and selection support this

hypothesis, as recombination tends to increase close to regulatory elements, while iHS

tends to decrease (Supplementary Figures 1, 2, 3). Therefore, high levels of local

recombination could erode haplotypes close to the selected variants, although the average

recombination in the gene window covering that region is low. In other words, the 1,000

kb gene windows we use could have a low recombination rate on average, while

containing local peaks of recombination close to regulatory elements. Our original model

considers the average recombination in each 1,000 kb gene-centered window, so it is

likely unable to detect the effect of these fine-scale patterns of recombination (see below

for further analyses in this respect).

Robustness of recent selection patterns to varying genomic window sizes

Next, we ask if the trends observed with 1,000 kb windows in the Yoruba genomes also

hold or not when using smaller window sizes (Supplementary Data 1). Some patterns

visible in large 1000kb windows are also visible when using smaller genomic window

sizes, while some are not (100 kb vs 1,000 kb windows; Supplementary Data 1). The

strong negative association between recombination rate and selection is found regardless

of window size. On the contrary, the distance to VIPs for example shows a strong

association with iHS in large windows but not in smaller windows (100 kb windows:

slope=-0.02; p-value=5.11E-01). This is consistent with previous evidence showing that

VIPs are particularly enriched in strong selective sweeps, for which larger windows are

more sensitive (Enard and Petrov 2020). This is also the case for the density of conserved

elements (100 kb windows: slope=0.1; p-value=9.3E-02) and especially regulatory

elements in testis, with a slope going from -0.70 in 1,000 kb windows to -0.03 in 100 kb

windows. These results suggest that both factors may correlate with strong rather than

weak selection. In the case of testis regulatory density, this pattern might be explained by

the fact that smaller windows take better into account the fine scale patterns of

recombination, which limits the negative influence of using average recombination (see

below for fine-scale analyses of recombination). Note, however, that all these differences

might also be explained by the fact that smaller windows do not differentiate well

between inside and outside of the genomic regions they are supposed to delineate. That

is, they may be more influenced by the surrounding, genetically linked genomic regions

(Enard et al. 2014; Enard and Petrov 2020).

Robustness of the recent selection patterns across human populations

We next ask whether the selection patterns observed in the Yoruba genomes are

consistent across different human populations. Association patterns with iHS are largely

similar across the five populations studied. Recombination rate and GC-content show

strong associations with iHS in 1000kb window across all populations (Tables 1 and 2).

Similarly, functional factors consistently correlate with selection across populations. This

is the case of VIP distance, gene expression in immune cells and density of regulatory

elements in testis. Most patterns are significant across all groups, but to varying extents.

One notable difference is that the density of conserved elements associates with selection

in 1000kb windows for all populations except for the Peruvian population sample, which

might reflect the impact of past bottlenecks on the visibility of the selective patterns

(Tables 1 and 2). Regulatory density across multiple tissues according to ChIP-seq

experiments and hypersensitivity to DNaseI along with coding density correlate with

selection in all populations except in Yoruba (Tables 1 and 2). Despite we detect a

masking effect of GC-content and conserved elements on functional factors (see above),

it is still surprising that the association between selection and some functional factors is

not readily visible in the Yoruba population, while it is in others. The selected component

is larger (i.e., selection is more visible) in Yoruba than in other populations. Therefore,

the association between functional density and iHS should be more visible in this

compared to other populations that have been affected by large bottlenecks connected to

the Out of Africa migration. We will return to this apparent contradiction later in the

Discussion. See Supplementary Data 1 for the complete results across populations and

window sizes.

Table 2. Slopes and p-values of the association between iHS and genomic factors for

four of the five studied populations in 1,000 kb windows: Utah residents, Toscani,

Han Chinese and Peruvians.

Covariate	Utah residents		Toscani		Han Chinese		Peruvians	
	Slope	p-value	Slope	p-value	Slope	p-value	Slope	p-value
Intercept	-1.77	<1E-16	-1.66	<1E-16	-1.78	<1E-16	-1.55	<1E-16
Recombination rate	-2.80	<1E-16	-2.70	<1E-16	-2.51	<1E-16	-1.74	<1E-16
GC-content	0.75	1.35E-08	0.52	2.47E-05	1.17	<1E-16	0.35	2.63E-03
Density of conserved elements	0.38	7.74E-13	0.34	2.70E-11	0.31	4.20E-10	0.06	2.43E-01
Number PPIs	-0.05	1.95E-01	-0.04	2.67E-01	-0.09	1.92E-02	-0.04	3.00E-01
Distance to VIPs	-0.34	8.45E-14	-0.27	1.92E-10	-0.14	5.35E-04	-0.18	2.77E-05
Gene expression	-0.34	9.70E-06	-0.25	4.92E-04	-0.25	1.14E-03	-0.26	2.56E-04
Gene expression in immune cells	0.41	6.87E-09	0.37	5.34E-08	0.31	9.51E-06	0.25	1.56E-04
Gene expression in testis	-0.03	5.45E-01	-0.04	4.21E-01	0.07	2.06E-01	0.05	2.61E-01
Regulatory density (ChIP-seq)	0.59	3.45E-06	0.74	3.06E-10	0.49	3.92E-05	0.33	7.64E-03
Regulatory density in immune cells (ChIP-seq)	-0.05	6.01E-01	-0.08	3.94E-01	-0.16	1.26E-01	0.15	1.32E-01
Regulatory density in testis (ChIP-seq)	-1.03	<1E-16	-0.86	<1E-16	-0.39	1.65E-09	-0.11	4.76E-02
Regulatory density (DNaseI)	-0.61	5.45E-05	-0.64	8.24E-06	-0.84	1.66E-08	-0.55	8.27E-05
Gene length	-0.05	2.84E-01	-0.02	5.59E-01	0.00	9.13E-01	0.02	5.46E-01
Gene number	-0.52	1.28E-06	-0.40	5.91E-05	-0.46	6.23E-04	-0.02	7.96E-01
Coding density	0.54	8.72E-06	0.59	3.77E-08	0.40	1.93E-03	0.20	2.73E-02
Number iHS data points	0.32	<1E-16	0.33	<1E-16	0.16	7.29E-08	0.08	1.51E-03

Genomic determinants of selection in low-recombination regions

Given the strong influence of recombination on selection across populations and window

sizes, we further assess whether the rest of significant associations with other genomic

factors are truly independent of recombination in the MDR, and whether insufficient

accounting of recombination can explain the observed selected component in the MDR,

instead of actual recent positive selection. The risk indeed exists that because we use an

imperfect measure of recombination in our model (due to measurements errors in

recombination maps and variation across populations), the effect of recombination on

iHS may not be properly removed. The selected component of the distribution of iHS

might then reflect an effect of recombination, rather than actual signals of recent positive

selection.

Figure 2. Mixture of Gaussian distributions fitting observed iHS (1,000 kb windows) in low-recombination regions for the five studied populations: A) Africa - Yoruba; B) Europe - Utah residents with Northern and Western European ancestry; C) Europe - Toscani; D) East Asia - Han Chinese; E) America - Peruvians. The figure shows the non-selected and selected components (Component 1 and 2, respectively) for each population considering only genes in low-recombination regions. In the latter component, iHS linearly depends on the genomic factors considered. Legend: Light blue = Observed iHS; Dark blue = Mixture model; Full red curve = Component 1 of the mixture model without selection; Dashed red curve = Component 2 of the mixture model with selection.

Because of this concern, we test if our results are robust to recombination by

recalculating the association between all the genomic factors and iHS, but this time

considering only genes with more similar recombination rates overall, in particular lower

recombination rates. Indeed, by using genes with more homogeneous recombination rates

and a reduced range of recombination values, we should also reduce any indirect effect of

recombination. We focus on genes with low recombination rates because we can predict

that if the significant associations found are due to actual positive selection, then the

positive selection component predicted by the MDR should become more prominent.

This is expected because positive selection is more readily detectable in low

recombination regions, where the eroding effect of recombination is reduced (O'Reilly et

al. 2008; Booker et al. 2020). Conversely, if the selected component and the significant

associations are in fact due to residual correlations with recombination but not actual

selection, then the selective component should become less prominent when focusing on

regions with more homogeneous, lower recombination rates. Note, however, that for

specific associations between iHS and genomic factors, a decrease of significance could

be caused by a reduction in gene sample size. The consideration of only a subset of genes

could reduce the power to detect association for genomic factors that correlate with

recombination. Focusing only on lower recombination genes may restrict the range of

values taken by these genomic factors, which could in turn limit our ability to detect

positive associations with iHS.

We select those genes with a recombination rate equal to or lower than the lowest

tertile of recombination across genes (0.837 cM/Mb). Larger windows are more sensitive

to strong selection and the estimates of genomic factors are more precise in these

windows. In addition, the Yoruba population again shows the largest selected component

across all the studied populations (Figure 2). Therefore, in the main text we only describe

results with 1,000 kb windows in Yoruba. See Supplementary Data 3 for full results of

low-recombination regions across other populations and windows sizes.

As expected from reducing the range of recombination values, running the model

with similar levels of recombination substantially decreases the regression slope between

recombination rate and iHS compared to the whole set of genes (slope: -1.01 vs. -2.44;

Table 3). In contrast, the selected component (second distribution in the model) becomes

larger (Figure 2 compared to Figure 1). This is consistent with the expectation that recent

positive selection is more readily detectable in regions where recombination is less

frequent, and hence long haplotypes caused by selection can increase in frequency and

remain unbroken for a longer amount of evolutionary time. Congruent with this, the

distance to VIPs is more associated with selection in low-recombination regions (slope=-

0.34 vs. -0.17 in the entire gene set; p-value=2.18E-05 vs. 4.16E-06). An expected result

given that VIPs distance does not correlate strongly with recombination (Spearman's

ρ=0.028; p-value=4.40E-04; correlation calculated across 1000 kb windows), thus its

range of values is not restricted by focusing only on low recombination genes. Despite

GC-content is strongly correlated with recombination (Spearman's ρ=0.30; p-value<1E-

16; correlation calculated across 1000 kb windows), its association with iHS remains

very strong and positive at low recombination genes (slope=0.93 vs. 0.55; p-

value=1.17E-06 vs. 1.34E-06; Table 3). This latter association suggests that the positive

trend of GC-content may be independent of recombination, given that it increases when

focusing only on low-recombination regions. Regulatory density in testis maintains a

negative association with iHS (slope=-0.76 vs. -0.79; p-value=1.10E-11 vs. <1E-16;

Table 3), while regulatory density across multiple tissues becomes significant and

negatively associated with selection in low-recombination regions (ChIP-seq; slope=-

0.63 vs. -0.15, p-value=7.22E-04 vs. 2.37E-01). As previously explained, a higher

accessibility of transcription factors to DNA in recombination hotspots should mask the

influence of regulatory density on selection. Indeed, if the mismatch between

recombination at local and gene-window scale is the cause of this pattern, focusing on

low-recombination regions should increase this confounding effect, by increasing the

discrepancy between low window-wide average recombination rates and more elevated

fine-scale recombination rates around regulatory elements. We specifically focus on gene

windows having overall low recombination, but they could still have fine-scale local

peaks of recombination, resulting in an increased mismatch between scales. Results from

this and previous sections suggest that a fine control of recombination rate is needed in

order to limit its confounding effect on the detection of recent positive selection. In the

next section we further analyze the association of iHS with genomic factors, but this time

controlling for fine-scale patterns of recombination.

Table 3. Slopes and p-values of the association between iHS and genomic factors for

the Yoruba population in 1,000 kb windows. These associations were calculated

considering only low-recombination regions.

Covariate	Slope	p-value
Intercept	-0.153	3.56E-01
Recombination rate	-1.005	<1E-16
GC-content	0.929	1.17E-06
Density of conserved elements	0.085	3.14E-01
Number PPIs	-0.095	9.66E-02
Distance to VIPs	-0.344	2.18E-05
Gene expression	-0.113	3.31E-01
Gene expression in immune cells	0.275	1.14E-02
Gene expression in testis	-0.087	2.68E-01
Regulatory density (ChIP-seq)	-0.633	7.22E-04
Regulatory density in immune cells (ChIP-seq)	0.217	2.17E-01
Regulatory density in testis (ChIP-seq)	-0.757	1.09E-11
Regulatory density (DNaseI)	0.229	3.09E-01
Gene length	-0.094	1.46E-01
Gene number	-0.265	7.58E-02
Coding density	-0.01	9.49E-01
Number iHS data points	-0.01	8.37E-01

PPIs: Protein-Protein Interactions. VIPs: Virus Interacting Proteins. ChIP-seq: Density of

regulatory elements according to chromatin immunoprecipitation experiments. DNaseI:

Density of regulatory elements according to the hypersensitivity to DNaseI. iHS:

Integrated Haplotype Score.

Although some factors remain significant, in other cases there is a decrease of

significance (Table 3). For instance, gene expression in immune genes shows a similar

slope (slope=0.27 vs. 0.26), but decreased significance (p-value=1.14E-02 vs. 1.51E-05).

The association with selection of the density of conserved elements decreases even more,

completely losing significance (slope=0.085 vs. 0.16, p-value=3.14E-01 vs. 6.4E-04).

This could be caused by the smaller sample of genes included, as focusing only on low-

recombination genes could restricts the range of values taken by these factors. This is

supported by the fact that immune gene expression and especially the density of

conserved elements are correlated with recombination (Spearman's ρ=-0.044 and -0.17;

p-value = 3.94E-08 and <1E-16; correlation calculated across 1000 kb windows).

Fine scale patterns of recombination around regulatory elements

We further analyze the influence of local patterns of recombination around regulatory

elements in our results. Given that recombination increases closer to the regulatory

elements (Supplementary Figures 1, 2, 3), it is likely that whole window estimates of

recombination are not an accurate measure of its effect on the association between iHS

and regulatory element densities. To address this limitation, we use three variables related

to the recombination around regulatory elements in immune cells, testis and across

multiple tissues, respectively. We calculate the more local, average recombination around

regulatory elements within each gene window, up to a maximum distance of 5kb from

each side of regulatory elements. Each variable is included in the original model of

Yoruba 1000 kb separately, and replaces the previous, window-wide average

recombination. Given that some gene windows can overlap with more regulatory

elements than others, we also consider the number of recombination data points obtained

for regulatory elements inside each gene window. In this way, we account for the fact

that recombination may vary more broadly in windows that have less regulatory

elements. See Supplementary Data 4 for further details about these calculations.

Recombination sharply increases around regulatory elements in immune cells

while iHS sharply decreases, with a steeper slope compared to other tissues

(Supplementary Figures 1, 2, 3), which might have confounded the expected positive

association with selection in the original regression model with window-wide

recombination rates. The control for local patterns of recombination around immune

regulatory elements should increase our power to detect this association. In line with this

prediction, we find that the association between immune regulatory density and iHS

becomes positive and highly significant when considering the local patterns of

recombination around immune regulatory elements (slope=0.68 vs. 0.047, p-

value=4.60E-14 vs. 6.24E-01; Supplementary Data 4). In order to assess whether this

increase is caused just by the lack of consideration of recombination patterns at a

window-wide scale, we repeat the analyses but including the original window-wide

recombination variable. After including recombination rate at window-wide scale, the

positive association remains stronger compared to the original model (slope=0.28 vs.

0.047, p-value=5.72E-03 vs. 6.24E-01).

As explained above, we hypothesize that the confounding effect of recombination

is caused by a mismatch between different scales. Gene windows with low recombination

can have local peaks of recombination around transcription factor binding sites

(Supplementary Figures 1, 2, 3), as these peaks are more accessible to transcription

factors (Spruce et al. 2020). Therefore, recombination could erode signals of positive

selection around regulatory elements even if the region has an overall low recombination.

In contrast, regions with high recombination at the window-wide level should suffer less

from this confounding effect, as recombination rate is then high both at a local and

window-wide scale. Therefore, focusing on high-recombination regions should improve

our ability to detect the expected positive association between selection and regulatory

density. This approach, however, has the caveat of reducing the power to detect positive

selection due the higher probability that recombination breaks selected haplotypes. We

nevertheless repeat the analyses focusing on gene windows with a recombination rate

equal or higher than the second tertile (1.552 cM/Mb). Focusing on high-recombination

regions, regulatory density in immune cells shows a stronger association with selection in

a model including recombination rate around both gene windows and immune regulatory

elements compared to the same model run across all genes (slope=0.85 vs, 0.28, p-

value=1.27E-08 vs. 5.72E-03). Interestingly, regulatory density in testis also shows a

positive association with selection in a model including recombination around both testis

regulatory elements and gene windows and focused on high-recombination regions. Note

that this regulatory variable shows one of the strongest negative associations with

selection in the original model (slope=0.58 vs. -0.79, p-value=4.98E-05 vs. <1E-16), an

unexpected result given the role of regulatory sequences in positive selection (Enard et al.
2014). These results suggest that, indeed, a higher regulatory density can increase the
probability for positive selection to occur, especially in some tissues. However, this
signal is distorted by fine-scale patterns of recombination, which have to be carefully
taken into account.

Discussion

We find several functional factors independently associated with recent positive
selection. For instance, viral interactions along with gene expression and regulatory
density in immune cells are strongly and positively associated with selection. Viruses
have acted as key drivers of adaptation in humans and VIPs show higher expression in
lymphocytes compared to non-VIPs (Halehalli and Nagarajaram 2015; Enard et al. 2016;
Enard and Petrov 2018). These results confirm the existence of strong viral and immune
selective pressures during recent human evolution (Enard and Petrov 2020; Souilmi et al.
2021; Enard and Petrov 2018). We also detect a positive association between testis
regulatory density and selection, which is, congruent with the fact that reproduction-
related functions show signals of positive selection (Nielsen et al. 2005; Voight et al.
2006). Note, however, that this latter result has been more difficult to reveal in our
analyses, as it is only visible after controlling for fine-scale patterns of recombination
around testis regulatory elements and after focusing on high-recombination regions, thus
it is more questionable. Finally, the density of conserved elements and GC-content,
which are related to overall (coding and non-coding) functional density (Lander et al.
2001; Di Filippo and Bernardi 2008; Siepel et al. 2005), also show a positive association
with selection. In summary, recent and positive selection is associated with the
distribution of functional regions across the human genome.

If sweep signals are false positives only reflecting genetic drift, the signatures of
sweeps should occur randomly across the genome. In contrast, if selective sweeps were
relatively common during recent human evolution, we expect that other factors on top of
recombination should associate with sweep signals across the human genome. In that

case, we expect that selective sweeps should occur more frequently around functional
(coding or non-coding) elements where adaptive mutations are expected to take place
(Enard et al. 2014), also around genomic regions associated with specific functions. We
find strong evidence for the latter scenario, as multiple functional factors are associated
with recent positive selection. In other words, the distribution of sweep signals is not
random, and is associated with the functional characteristics of the human genome.
Therefore, the present study provides evidence to further clarify the controversy about the
frequency of positive selection during recent human evolution (Enard et al. 2014;
Hernandez et al. 2011). Our results support that selective sweeps were at least common
enough during recent human evolution to generate associations between selection and
functional factors.

Our MDR approach shows a good fit to the distribution of iHS across the human
genome, with its two components (selection and genetic drift) showing a clear
differentiation. This suggests that the consideration of a more biologically meaningful
model, which assumes two Gaussian distributions instead of one, works better to analyze
recent adaptation than simple correlations. The associations found using this approach are
more informative than classic correlations as this model represents more directly genome
evolution and considers multiple factors simultaneously. These characteristics make it
possible to reveal expected, but also unexpected patterns. For instance, GC-content is
strongly and positively associated with iHS. Regions with high GC-content also tend to
have high long-term recombination rates (Duret and Arndt 2008), which would predict a
negative association between iHS and GC content, not positive. A possible explanation
for this pattern is that the MDR approach can control for recombination as a confounder
by simultaneously analyzing recombination rate and other genomic factors. This can
unravel new patterns, like a positive association between GC-content and sweep signals,
which may be independent from the eroding effect of recombination. This is supported by
the fact that GC-content only shows a negative association with selection after removing
recombination from the model. GC-content could be a better proxy of overall functional
density than individual functional factors (Lander et al. 2001; Di Filippo and Bernardi
2008), explaining its positive association with selection. Results from MDRs support this
hypothesis as the removal of functional factors like coding density, gene number or

regulatory density strongly increases the significance of this association. This could also

explain another surprising result, the lack of association between iHS and coding regions,

which are expected targets of selection. The removal of GC-content and the density of

conserved elements, which is another factor associated with overall functional density

(Siepel et al. 2005), unmasks a significant positive association between coding density

and selection. These results suggest that considering measures of overall functional

density (i.e., GC-content and conserved elements density) could absorb the expected

positive association with selection for individual functional factors like coding density.

These results illustrate the utility of our approach. The simultaneous consideration of

multiple predictors lets us unravel novel associations between genomic factors and

adaptation. In addition, it enables easier testing of different hypotheses by modifying the

set of predictors considered. Therefore, the MDR approach is useful to model recent

genomic adaptation, and to analyze its prevalence and determinants.

681   We find other unexpected results using the MDR approach. For instance, we do

not find a positive association between selection and testis regulatory density in the

original model. This is an unexpected result given that a higher regulatory density would

lead to a larger mutational target for adaptation through changes in expression. Given that

genomic regions with local peaks of recombination are usually more accessible,

transcription factors are more likely to bind DNA in these regions (Spruce et al. 2020).

Therefore, any adaptative signal associated with regulatory density would be likely

eroded by recombination (Sabeti et al. 2006; O'Reilly et al. 2008; Booker et al. 2020). In

support of this hypothesis, we find higher recombination and lower iHS closer to

regulatory elements. Indeed, when fine-scale patterns of recombination are considered as

new predictors, regulatory density in immune cells and testis become strongly and

positively associated with selection. As previously mentioned, the positive association of

testis regulatory density only emerges after focusing on high-recombination regions, thus

it should be taken with more caution. These results support the role of regulatory

sequences as targets of positive selection independently of other genomic features (Enard

et al. 2014), along with the relevance of testis and specially immune cells in recent

human evolution (Nielsen et al. 2005; Voight et al. 2006). Our results also illustrate how

the detection of selection can be hindered by fine-scale characteristics of the genome, like

the sharp increase of recombination we observe around regulatory elements. This might

mask selection signals around regulatory elements, but not only because of the breaking

of haplotypes caused by recombination. An additional mechanism might be the existence

of biased gene conversion in these recombinant regions, which could specifically favor

GC mutations (gBGC) (Galtier et al. 2001; Duret and Arndt 2008; Dutta et al. 2018). This

could lead to an increase in the frequency of GC mutations that are associated with short

haplotypes (due to the eroding effect of recombination). BGC might therefore make it

even more difficult to detect recent positive selection around regulatory regions, where it

is expected to happen (Enard et al. 2014). Note, however, that the present study does not

provide direct evidence about the implication of gBGC in the observed patterns. It can be

difficult to disentangle the eroding effect of recombination from GC-biased gene

conversion on haplotypes because gBGC is caused and therefore very correlated with

recombination. Distinguishing the specific effect of biased-gene conversion may require

precise recombination maps for African populations, where we find the clearest selection

signals. Although we have not found evidence about the implication of gBGC yet, the

fact that this hypothesis has emerged from implementing our model illustrates its utility

to propose new hypotheses about the different processes that influence the detection and

occurrence of positive selection.

Although the MDR approach provides an excellent fit to the observed selection

statistic, we may still not detect associations with specific factors because they do not

have a simple linear relationship with recent selection. Although more biologically

meaningful than a model assuming just one distribution, our approach is still linear, thus

more complex relationships may not be detected. For example, the unexpected lack of

significance for coding density could be explained not only by the consideration of better

proxies of overall functional density, but also by the existence of non-linear relationships

with selection. Advantageous mutations are more likely to appear in regions with high

coding density, thus a high density of coding sequences should favour the appearance of

selection (Enard et al. 2014). However, high coding density can also mean a higher

frequency of deleterious mutations. These mutations can interfere with adaptation when

they are in linkage with advantageous mutations. This would be especially relevant for

genomic regions with low recombination, where the linkage between deleterious and

advantageous mutations is more likely (Di et al. 2021). Therefore, a higher coding

density could favor or hinder positive selection depending on the circumstances,

complicating the detection of an association between coding density and selection with

our approach. The existence of this genetic interference could also explain the different

results observed across populations in relation to coding density. Despite having the

largest selected component, the Yoruba population does not show a clear link between

selection and coding density. This association becomes visible only after removing

proxies of overall functional density (i.e., density of conserved elements and GC-

content). In contrast, this pattern is more readily visible in the other populations studied

despite showing smaller selected components. A potential, but still speculative

explanation at this point could be the fact that bottlenecks related to the migration out of

Africa may have removed segregating recessive deleterious variants, which may reduce

their interference over advantageous mutations, thus making it easier for positive

selection to act on coding regions (Di et al. 2021). In contrast, there were no similar

bottlenecks to decrease the genetic interference in the Yoruba, and regions of the genome

with high a coding density may still experience stronger interference from recessive

deleterious mutations. Interestingly, despite this potential limitation of positive selection,

we still see a larger selective component in the entire Yoruba genome, suggesting that

selection is either more abundant and/or more visible in this population. Note, however,

this is a potential explanation that is speculative at this point and will require further

investigation.

Conclusions

753   We found multiple functional factors highly associated with recent selection using MDRs

754   with two distributions meant to reflect the selected and drift components of the human

755   genome. Our results suggest that positive selection was relatively common in recent

756   human evolution and highlight MDRs as a novel approach to quantify the prevalence of

757   recent adaptation and its determinants in the human and other genomes.

Methods

Window sizes and genes coordinates

All analyses were performed using hg19 genomic coordinates and protein-coding gene annotations from Ensembl v99 (Yates et al. 2020). We considered windows of different sizes (50, 100, 200, 500 and 1,000 kb). Each window was centered at the genomic center of a gene, in the middle between the most upstream transcription start site and the most downstream transcription end site. We used a fixed window size to avoid biases related to gene length, as larger genes are more likely to overlap with high local iHS values just by chance compared to shorter genes. This would bias power in favour of larger genes, increasing the probability to detect sweeps in them just because of their length. In addition, the consideration of different window sizes provides the opportunity to detect different types of sweeps as larger windows are specifically sensitive to strong selective sweeps compare to smaller windows (Enard and Petrov 2020).

Genomic features

Multiple genomic factors were calculated inside the gene windows. We considered the following factors that are likely to influence the frequency of sweeps:

- Length of the gene at the center of each window.

- Number of genes overlapped with each window. We used this value as an estimate of gene density.

- Recombination rate. For each window, we calculated the genetic distance between the edges of the window and then divided it by the physical distance between them. Genetic position of each window edge was obtained from the deCODE 2019 genetic map (Halldorsson et al. 2019). In case no data was available for a window edge, we selected the closest genetic position data points within 50 kb at each side of the window edge to estimate its genetic position. We used linear interpolation to this end, considering the genetic and physical position of the two points around the

window edge. In other words, we assumed that genetic distance increased linearly

between the two selected data points, and hence we can estimate the genetic position

of the window edge based on its physical position respect to the other two positions.

If no genetic position data was available at 50 kb or closer to the window edge, no

genetic position was calculated and the whole window was discarded for

recombination calculations and subsequent analyses. Note that we searched for

genetic position data points only up to 50 kb from each window edge to avoid that

the total distance between points at each edge could be too large (specially for larger

windows). In that scenario, linear interpolation could be inappropriate because we

would use very distant data to calculate recombination rate. For example, if a 1,000

kb window had no genetic position for any of its edges and we looked for genetic

positions up to 1,000 kb at each side, it would possible that the selected data points

of genetic position downstream and upstream of the window were separated by

3,000 kb. In addition, the recombination rate would not correspond with the size of

the window.

• Density of coding sequences. We used Ensembl v99 coding sequences. The density

was calculated as the proportion of coding bases respect to the whole length of the

window without considering gaps (gap locations obtained from the UCSC Genome

Browser; https://genome.ucsc.edu/;

http://hgdownload.cse.ucsc.edu/goldenPath/hg19/database/). A similar approach was

used for the rest of density estimates.

• Density of mammalian phastCons conserved elements (Siepel et al. 2005), also

downloaded from the UCSC Genome Browser. Given that each conserved segment

had a score, we considered as conserved only those segments above a given

threshold. In order to minimize the inclusion of non-conserved elements, we used a

threshold that only considers 10% of the genome as conserved, as it is unlikely that

more than 10% of the whole genome is constrained (Siepel et al. 2005).

• Density of regulatory elements. We calculated several variables related to the

density of binding sites for transcription factors. In all cases, we calculated the

proportion of sequences that are considered binding sites within a window. The data

were also obtained from the UCSC Genome Browser (The ENCODE Project
Consortium 2012).

– Density of DNaseI hypersensitive sites. We considered as binding sites those
segments with a score higher than a given threshold. The selected threshold
was a score value according to which only 10% of the genome is considered
accessible to DNaseI, and hence a binding site for transcription factors. In
this way, we minimize the probability to consider non-binding sites (i.e.,
false positives).

– Density of binding sites according to the technique of chromatin
immunoprecipitation followed by sequencing (ChIP-seq). We calculated a
threshold to consider a given segment as a binding site following the same
approach used for DNaseI hypersensitive sites. This database includes 1264
experiments representing 338 transcription factors in 130 cell types. We
calculated the density of binding sites using the whole dataset, but also
considering two subsets, one for experiments performed on cells lines of
testis, and another for experiments performed on immune cells (lymphocytes
in most cases).

• GC-content, calculated as a percentage per window. It was also obtained from the
UCSC Genome Browser.

• Gene expression: We used the log (base 2) of TPM (Transcripts Per Million), which
was obtained from GTEx (GTEx Consortium 2015)
(https://www.gtexportal.org/home/). We considered the average gene expression
across 53 GTEx v7 tissues, but also expression in immune cells (lymphocytes) and
testis.

• Number of protein-protein interactions (PPIs) in the human protein interaction
network (Luisi et al. 2015). We used the log (base 2) of the number of PPIs of each
gene. We summed 1 to the original dataset to avoid problems applying the logarithm
to genes with no interactions (i.e., PPI=0).

• Proportion of genes that interact with viruses (VIPs). We used a previously
published dataset (Enard and Petrov 2020) that includes around 4500 VIPs with
evidence of physical interaction with viruses. 1920 of these VIPs were manually
curated from the virology literature, while the remaining 2600 VIPs were identified
using high-throughput methods and retrieved from the VirHostnet 2.0 database and
additional studies. See the original publication for further details (Enard and Petrov
2020). The variable included in the model was calculated as the distance of each
gene to the closest gene that interacts with viruses.

iHS calculation

In order to calculate iHS, we used polymorphism data from the 1,000 Genomes Project
phase 3 (The 1000 Genomes Project Consortium 2015). We calculated iHS for 5
populations: Yoruba, Utah residents with Northern and Western European ancestry,
Toscani, Han Chinese and Peruvians. We used the hapbin software (Maclean et al. 2015)
in order to perform fast scans of iHS across the genomes of these populations. Note that
the genetic maps used as input in hapbin were calculated using the deCODE 2019 genetic
map (Halldorsson et al. 2019). In case that no data was available for the focal variant, we
again used linear interpolation to estimate its genetic position. We followed the same
approach used to calculate the genetic position of window edges during recombination
calculations. In this case, we searched for genetic position data points at both sides of the
focal variant until 1,000 kb. We discarded variants with no data points around, as they
were present in genomic regions with low density of recombination data. The genetic
maps based on the deCODE 2019 map were used as an input in order to calculate the iHS
for the 5 populations.

The raw iHS values obtained with hapbin were standardized by frequency. For
each population, we divided variants in 50 bins of frequency (from 0 to 1) considering the
whole genome. We then calculated the mean and standard deviation of raw iHS within
each bin, using them to standardize raw iHS values: (raw iHS - mean iHS) / sd iHS. The
mean of standardized iHS was calculated per window, considering the absolute value of

iHS. In other words, we considered as high selection both very positive and very negative

iHS values, which indicates selection in favor of the haplotypes carrying the derived and

the ancestral allele, respectively. We also obtained the number of iHS values per window

to have an estimate of data density. Mean iHS in windows with low number of iHS

values can be more influenced by outliers (very low or very high iHS values). Therefore,

we included the number of iHS values per window as predictor in the models to control

for this.

Modeling

We modeled the association between genomic factors and iHS in each population and

window size by combining a Gaussian mixture distribution with linear regression.

Mixture distributions are useful to model data that cannot be fully described by a single

distribution and that are likely generated by multiple processes. In this case, a mixture of

two Gaussian distributions could fit better a scenario where selection influences some

genomic regions but not others, and hence a bimodal distribution of iHS would be

observed across the genome. Therefore, we posit that sweep signals in the human genome

may have a selected and a non-selected component. According to this, we assumed that

observed log iHS followed a mixture of two Gaussian distributions representing the two

selection components. Because selective sweeps often result in elevated iHS, we assumed

that the Gaussian distribution for the selected component had a higher mean than that for

the non-selected component. Furthermore, to model the effects of genomic factors on

iHS, we assumed that the probability of the selected component was a linear combination

of genomic factors followed by a sigmoid transformation. Specifically, for each factor,

we calculated the product of its observed value with a slope indicating its association

with iHS. We summed the product over all factors for a given gene to model the

accumulative effect of genomic factors on the distribution of iHS. Then, we added an

intercept indicating the baseline level of selection when all genomic factors are equal to 0

and transformed the result to a probability (0 to 1) using a sigmoid function. This

probability was obtained for each gene and can be regarded as the overall influence of all

genomic factors on the frequency of recent selective sweeps. These steps are described by

the equation $p = \sigma(\sum b_i * x_i + a) = \frac{1}{1+\exp(-\sum b_i * x_i - a)}$, where p is a probability

summarizing the influence of all studied genomic factors on iHS, σ is the sigmoid

function, b_i and x_i are the slope and the observed value for genomic factor i in a given

gene, respectively, and a is the intercept.

To obtain the likelihood of log iHS in each gene, we calculated a weighted sum of

Gaussian density functions, $P(Y) = p * f(Y|\mu_1, \sigma_1) + (1 - p) * f(Y|\mu_0, \sigma_0)$, where Y

and $P(Y)$ are the observed log iHS and its likelihood in a given gene, respectively, p is

the probability of the selected component calculated in the previous step, $f(Y|\mu_1, \sigma_1)$ is

the Gaussian density function with mean μ_1 and standard deviation σ_1 describing the

selected component, and $f(Y|\mu_0, \sigma_0)$ is the Gaussian density function with mean μ_0 and

standard deviation σ_0 describing the non-selected component. Finally, we summed the

logarithm of $P(Y)$ over all genes to obtain the log likelihood of the entire dataset.

We estimated model parameters by maximizing the log likelihood of the entire

dataset. These parameters included the mean and standard deviation of each Gaussian

distribution, along with the intercept and the slopes representing the association between

iHS and each genomic factor. For each genomic factor, we tested the significance of its

slope by comparing the log likelihood between the full model with all genomic factors

and a nested null model without the selected factor. We assumed that two times the

difference in log likelihood between the two models followed a chi-square distribution

with one degree of freedom.

Data access

925   Genetic polymorphism data was obtained from the 1,000 Genomes Project (The 1000

926   Genomes Project Consortium 2015). Data for the genomic factors were mostly obtained

927   from the UCSC genome browser (https://genome.ucsc.edu/), along with other public

928   sources. See Methods section for specific details about the source of each dataset.

Competing interest statement

The authors declare no competing interests.

Acknowledgments

We thank the members of the Enard, Masel and Gutenkunst labs for their helpful comments. DFST was supported by a Marie S. Curie Global Fellowship within the European Union research and innovation framework programme (2014-2020; ClimAHealth: 101030971). Y-FH was supported by the National Institute of General Medical Sciences of the National Institutes of Health under Award Number R35GM142560 and by startup funds from Pennsylvania State University. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health or the European Union.

Author contributions

Y-FH, DE and DFST designed the study. DE and DFST collected the data. Y-FH and DFST performed the analyses. DE and DFST wrote the paper and interpreted the results. Y-FH, DE and DFST edited and approved the paper.

References

Barreiro LB, Ben-Ali M, Quach H, Laval G, Patin E, Pickrell JK, Bouchier C, Tichit M, Neyrolles O, Gicquel B, et al. 2009. Evolutionary dynamics of human toll-like receptors and their different contributions to host defense. *PLoS Genet* **5**.

Booker TR, Yeaman S, Whitlock MC. 2020. Variation in recombination rate affects detection of outliers in genome scans under neutrality. *Mol Ecol* **29**: 4274–4279. https://doi.org/10.1111/mec.15501.

Deschamps M, Laval G, Fagny M, Itan Y, Abel L, Casanova JL, Patin E, Quintana-Murci

L. 2016. Genomic Signatures of Selective Pressures and Introgression from Archaic

Hominins at Human Innate Immunity Genes. *Am J Hum Genet* **98**: 5–21.

http://dx.doi.org/10.1016/j.ajhg.2015.11.014.

Di C, Murga Moreno J, Salazar-Tortosa DF, Lauterbur ME, Enard D. 2021. Decreased

adaptation at human disease genes as a possible consequence of interference

between advantageous and deleterious variants. *Elife* **10**: e69026.

https://doi.org/10.7554/eLife.69026.

Di Filippo M, Bernardi G. 2008. Mapping DNase-I hypersensitive sites on human

isochores. *Gene* **419**: 62–65.

Duret L, Arndt PF. 2008. The impact of recombination on nucleotide substitutions in the

human genome. *PLoS Genet* **4**: e1000071.

https://doi.org/10.1371/journal.pgen.1000071.

Dutta R, Saha-Mandal A, Cheng X, Qiu S, Serpen J, Fedorova L, Fedorov A. 2018. 1000

human genomes carry widespread signatures of GC biased gene conversion. *BMC*

Genomics **19**: 256.

Enard D, Cai L, Gwennap C, Petrov DA. 2016. Viruses are a dominant driver of protein

adaptation in mammals. *Elife* **5**: e12469.

Enard D, Messer PW, Petrov DA. 2014. Genome-wide signals of positive selection in

human evolution. *Genome Res* **24**: 885–895. https://doi.org/10.1101/gr.164822.113.

Enard D, Petrov DA. 2020. Ancient RNA virus epidemics through the lens of recent

adaptation in human genomes. *Philos Trans R Soc London Ser B* **375**: 20190575.

http://dx.doi.org/10.1098/rstb.2019.0575.

Enard D, Petrov DA. 2018. Evidence that RNA Viruses Drove Adaptive Introgression

between Neanderthals and Modern Humans. *Cell* **175**: 360–371.

https://doi.org/10.1016/j.cell.2018.08.034.

Ferrer-Admetlla A, Liang M, Korneliussen T, Nielsen R. 2014. On detecting incomplete

soft or hard selective sweeps using haplotype structure. *Mol Biol Evol* **31**: 1275–
1291. https://doi.org/10.1093/molbev/msu077.

Galtier N, Piganeau G, Mouchiroud D, Duret L. 2001. GC-Content Evolution in
Mammalian Genomes: The Biased Gene Conversion Hypothesis. *Genetics* **159**:
907–911. https://doi.org/10.1093/genetics/159.2.907.

Garud NR, Messer PW, Buzbas EO, Petrov DA. 2015. Recent Selective Sweeps in North
American Drosophila melanogaster Show Signatures of Soft Sweeps. *PLoS Genet*
11: e1005004. https://doi.org/10.1093/molbev/msu077.

GTEx Consortium. 2015. The Genotype-Tissue Expression (GTEx) pilot analysis:
Multitissue gene regulation in humans eds. K.G. Ardlie, D.S. Deluca, A. V Segrè,
T.J. Sullivan, T.R. Young, E.T. Gelfand, C.A. Trowbridge, J.B. Maller, T.
Tukiainen, M. Lek, et al. *Science (80-)* **348**: 648–660.
https://doi.org/10.1126/science.1262110.

Halehalli R, Nagarajaram HA. 2015. Molecular principles of human virus protein-protein
interactions. *Bioinformatics* **31**: 1025–1033.

Halldorsson B V., Palsson G, Stefansson OA, Jonsson H, Hardarson MT, Eggertsson HP,
Gunnarsson B, Oddsson A, Halldorsson GH, Zink F, et al. 2019. Characterizing
mutagenic effects of recombination through a sequence-level genetic map. *Science*
(80-) **363**. https://doi.org/10.1126/science.aau1043.

Harris DN, Song W, Shetty AC, Levano KS, Cáceres O, Padilla C, Borda V, Tarazona D,
Trujillo O, Sanchez C, et al. 2018. Evolutionary genomic dynamics of Peruvians
before, during, and after the Inca Empire. *Proc Natl Acad Sci U S A* **115**: E6526–
E6535.

Henn BM, Cavalli-Sforza LL, Feldman MW. 2012. The great human expansion. *Proc*
Natl Acad Sci U S A **109**: 17758–17764. https://doi.org/10.1073/pnas.1212380109.

Hernandez RD, Kelley JL, Elyashiv E, Melton SC, Auton A, McVean G, Sella G,
Przeworski M. 2011. Classic selective sweeps were rare in recent human evolution.

Science (80-) **331**: 920–924. https://doi.org/10.1126/science.1198878.

Huang Y-F. 2021. Dissecting genomic determinants of positive selection with an
evolution-guided regression model. *Mol Biol Evol* msab291.
ttps://doi.org/10.1093/molbev/msab291.

Johnson KE, Voight BF. 2018. Patterns of shared signatures of recent positive selection
across human populations. *Nat Ecol Evol* **2**: 713–720.
http://dx.doi.org/10.1038/s41559-018-0478-6.

Lander ES, Linton LM, Birren B, Nusbaum C, Zody MC, Baldwin J, Devon K, Dewar K,
Doyle M, Fitzhugh W, et al. 2001. Initial sequencing and analysis of the human
genome. *Nature* **409**: 860–921.

Lohmueller KE, Albrechtsen A, Li Y, Kim SY, Korneliussen T, Vinckenbosch N, Tian
G, Huerta-Sanchez E, Feder AF, Grarup N, et al. 2011. Natural selection affects
multiple aspects of genetic variation at putatively neutral sites across the human
genome. *PLoS Genet* **7**: e1002326. https:doi.org/10.1371/journal.pgen.1002326.

Luisi P, Alvarez-Ponce D, Pybus M, Fares MA, Bertranpetit J, Laayouni H. 2015. Recent
positive selection has acted on genes encoding proteins with more interactions
within the whole human interactome. *Genome Biol Evol* **7**: 1141–1154.
https://doi.org/10.1093/gbe/evv055.

Maclean CA, Chue Hong NP, Prendergast JGD. 2015. hapbin: An efficient program for
performing haplotype-based scans for positive selection in large genomic datasets.
Mol Biol Evol **32**: 3027–3029. https://doi.org/10.1093/molbev/msv172.

Nielsen R, Bustamante C, Clark AG, Glanowski S, Sackton TB, Hubisz MJ, Fledel-Alon
A, Tanenbaum DM, Civello D, White TJ, et al. 2005. A scan for positively selected
genes in the genomes of humans and chimpanzees. *PLoS Biol* **3**: e170.

O'Reilly PF, Birney E, Balding DJ. 2008. Confounding between recombination and
selection, and the Ped/Pop method for detecting selection. *Genome Res* **18**: 1304–
1313. https://doi.org/10.1101/gr.067181.107.

Pritchard JK, Pickrell JK, Coop G. 2010. The Genetics of Human Adaptation: Hard
        Sweeps, Soft Sweeps, and Polygenic Adaptation. *Curr Biol* **20**: R208–R215.
        http://doi.org/10.1016/j.cub.2009.11.055.

Sabeti PC, Schaffner SF, Fry B, Lohmueller J, Varilly P, Shamovsky O, Palma A,
        Mikkelsen TS, Altshuler D, Lander ES. 2006. Positive natural selection in the
        human lineage. *Science (80-)* **312**: 1614–1620.
        https://science.sciencemag.org/content/312/5780/1614.

Schrider DR. 2020. Background selection does not mimic the patterns of genetic diversity
        produced by selective sweeps. *Genetics* **216**: 499–519.

Siepel A, Bejerano G, Pedersen JS, Hinrichs AS, Hou M, Rosenbloom K, Clawson H,
        Spieth J, Hillier LDW, Richards S, et al. 2005. Evolutionarily conserved elements in
        vertebrate, insect, worm, and yeast genomes. *Genome Res* **15**: 1034–1050.
        https://doi.org/10.1101/gr.3715005.

Souilmi Y, Lauterbur ME, Tobler R, Huber CD, Johar AS, Varasteh Moradi S, Johnston
        WA, Krogan NJ, Alexandrov K, Enard D. 2021. An ancient viral epidemic
        involving host coronavirus interacting genes more than 20,000 years ago in East
        Asia. *Curr Biol* **31**: 1–11. https://doi.org/10.1016/j.cub.2021.05.067.

Spruce C, Dlamini S, Ananda G, Bronkema N, Tian H, Paigen K, Carter GW, Baker CL.
        2020. HELLS and PRDM9 form a pioneer complex to open chromatin at meiotic
        recombination hot spots. *Genes Dev* **34**: 398–412.
        https://doi.org/10.1101/gad.333542.119.

The 1000 Genomes Project Consortium. 2015. A global reference for human genetic
        variation. *Nature* **526**: 68–74. https://doi.org/10.1038/nature15393.

The ENCODE Project Consortium. 2012. An integrated encyclopedia of DNA elements
        in the human genome. *Nature* **489**: 57–74. https://doi.org/10.1038/nature11247.

The International HapMap Consortium. 2005. A haplotype map of the human genome.
        Nature **437**: 1299–1320. https://doi.org/10.1038/nature04226.

1063   Voight BF, Kudaravalli S, Wen X, Pritchard JK. 2006. A map of recent positive selection

1064   in the human genome. *PLoS Biol* **4**: 0446–0458.

1065   Yates AD, Achuthan P, Akanni W, Allen J, Allen J, Alvarez-Jarreta J, Amode MR,

1066   Armean IM, Azov AG, Bennett R, et al. 2020. Ensembl 2020. *Nucleic Acids Res* **48**:

1067   D682–D688. https://doi.org/10.1093/nar/gkz966.

1068