Title: Mixture Density Regression reveals frequent recent adaptation in the human

genome

Authors: Diego F. Salazar-Tortosa[1,2], Yi-Fei Huang[3,4], David Enard[1]

Affiliations:

[1]Department of Ecology and Evolutionary Biology, University of Arizona, Tucson, AZ,

USA.

[2]PROFITH 'PROmoting FITness and Health through physical activity' research group,

Sport and Health University Research Institute (iMUDS), University of Granada,

Granada, Spain.

[3]Department of Biology, Pennsylvania State University, University Park, PA, USA.

[4]Huck Institutes of the Life Sciences, Pennsylvania State University, University Park,

PA, USA.

Corresponding Authors: David Enard: denard@arizona.edu; Yi-Fei Huang:

yuh371@psu.edu

Running head: MDR-based analysis of positive selection.

Keywords: human evolution, recent adaptation, mixture distributions, selection

determinants, recombination rate.

Abstract

How much genome differences between species reflect neutral or adaptive evolution is a central question in evolutionary genomics. In humans and other mammals, the prevalence of adaptive versus neutral genomic evolution has proven particularly difficult to quantify. The difficulty notably stems from the highly heterogeneous organization of mammalian genomes at multiple levels (functional sequence density, recombination, etc.) that complicates the interpretation and distinction of adaptive vs. neutral evolution signals. Here, we introduce Mixture Density Regressions (MDRs) for the study of the determinants of recent adaptation in the human genome. MDRs provide a flexible regression model based on multiple Gaussian distributions. We use MDRs to model the association between recent selection signals and multiple genomic factors likely to affect positive selection, if the latter was common enough in the first place to generate these associations. We find that a MDR model with two Gaussian distributions provides an excellent fit to the genome-wide distribution of a common sweep summary statistic (iHS), with one of the two distributions likely enriched in positive selection. We further find several factors associated with recent adaptation, including the recombination rate, the density of regulatory elements in immune cells, GC-content, gene expression in immune cells, the density of mammal-wide conserved elements, and the distance to the nearest virus-interacting gene. These results support that strong positive selection was relatively common in recent human evolution and highlight MDRs as a powerful tool to make sense of signals of recent genomic adaptation.

Author Summary

Over the last 50,000 years, human populations have been exposed to selective pressures that can trigger adaptation in the genome. The search for signals of these selective events is however obscured by the substantial variation of factors that are relevant for the prevalence and detection of adaptation across the genome. Here, we analyze the impact of multiple factors on positive selection using a biologically meaningful approach that

considers the influence of adaptative and non-adaptive processes in the human genome.

Our results show that this novel approach is better suited to find genomic factors

associated with recent positive selection compared to the classic regression or correlation

approaches. We find multiple genomic functional elements associated with selection

across the genome, including novel associations that emerge only after controlling for

multiple confounding factors. This strongly suggests that adaptation was common enough

in recent evolutionary times to produce a widespread correlation between functional

elements and positive selection in the human genome.

Note on the language used in this manuscript

In this manuscript we use discrete population groups, such as the Yoruba. We want to

emphasize that these discrete groups are only used for convenience and clarity when

presenting our results, but in fact represent arbitrary human constructions, the same way

that the boundaries of countries are arbitrary. There is an unbroken continuum and

mixing of geographical ancestries across groups often identified as distinct populations

across the world. The discrete groups we use are, as such, by no mean discrete genetic

entities. Once grouped together, the grouped individuals only happen to be genetically

more similar, with their ancestries coming from specific geographic locations more

predominantly than individuals from the other groups. How much more predominantly is

completely arbitrary.

Introduction

The characteristics of the human genome can influence both the occurrence and the

detection of recent positive selection in a specific genomic region. Similar to other

mammals, the human genome has a complex organization, with highly heterogeneous

recombination rates and distribution of functional elements (regulatory or coding) along

chromosomes. This inherent heterogeneity in the factors that are likely to influence both

the occurrence of positive selection and our ability to detect it, has complicated the study

of the prevalence of recent adaptation in the human genome [1,2]. Another important

complicating factor is that statistics of recent adaptation have complex distributions

across the genome (see below), which limits the ability of classic correlation and

regression approaches to analyze the genomic factors that contributed to recent genomic

adaptation. There is nevertheless growing evidence from genomic scans suggesting that

recent positive selection in the form of selective sweeps may have been relatively

common during recent human evolution [3–9]. More recent and powerful selection scans

using machine (deep) learning approaches suggest that a non-negligible proportion of the

human genome may be affected by selective sweeps [10–13]. Advances on this topic will

depend on our ability to understand what factors govern the local genomic rate of recent

adaptation along human chromosomes. Indeed, if signatures of selective sweeps do not

occur randomly, as expected if all sweep signals are false positives only reflecting genetic

drift, but instead associate non-randomly with functional elements in the genome, then

this provides evidence that sweeps had to be common enough in the first place to

generate the association.

What factors then are *a priori* expected to matter for recent adaptation? Among

all genomic factors, recombination is a key process determining the patterns of linkage

disequilibrium between alleles across the human genome [14], that can strongly influence

the probability to detect recent selection in the form of selective sweeps. The lower the

recombination rate, the larger the genomic region where neutral variants will hitchhike to

higher frequencies along with an advantageous variant, before recombination breaks

down their linkage disequilibrium [15,16]. The higher the recombination rate, the smaller

the genomic region affected by the sweep. Different sweep sizes in low and high

recombination regions result in a higher statistical power to detect sweeps in low than in

high recombination regions [8,17,18]. In parallel, the increase of linkage disequilibrium

in low-recombination regions favors the appearance of larger neutral haplotypes making

these regions more prone to detect false positive sweeps [17,18]. In addition,

recombination rate influences the probability of deleterious variants interfering with the

adaptive ones given the increased probability of linkage disequilibrium between them

under low recombination [19–21]. This heterogeneity, together with other factors such as

background selection, has contributed to a persistent debate around the prevalence of

selective sweeps in the human genome [1,2].

If recent adaptation and selective sweeps in particular were relatively common

during recent human evolution, we expect that other factors on top of recombination

should be relevant for their occurrence and detection in the human genome. Generally,

we expect that selective sweeps should occur more frequently around functional

segments of the genome where adaptive mutations are expected to take place [1].

Therefore, we should find enough signals of positive association between selective

sweeps and the overall density of functional elements, either coding or non-coding. In a

pioneer analysis, Barreiro et al. (2008) showed for the first time a relationship between

genome-wide signatures of positive selection and the distribution of functional elements.

Among those highly differentiated Single Nucleotide Polymorphisms (SNPs), they found

an excess of non-synonymous SNPs compared to non-genic SNPs. Note, however, that

they did not explicitly control for the impact of background Selection (BGS), the

coincident removal of neutral variants together with genetically linked deleterious

mutations, which varies between the classes analyzed [2,23]. We also expect specific

functions to be associated with an increased occurrence of selective sweeps. For example,

the presence of genes coding for proteins that interact with pathogens, and viruses in

particular (Virus-Interacting Proteins or VIPs), should influence the frequency of sweeps

in a given genomic region, independently of other genomic features [9,24]. More

generally, genomic regions with immune genes are expected to have experienced more

sweeps [25–28]. Similarly, reproduction-related functions show signals of positive

selection [5,29], thus tissues related with these functions should be associated with the

frequency of sweeps.

Several summary statistics are currently available to detect recent genomic

adaptation. Among all possible choices, the statistics that use the structure of haplotypes

along chromosomes are the most appropriate to investigate the prevalence and

determinants of recent positive selection. First, haplotype-based statistics have good

statistical power to detect strong and recent incomplete sweeps [5,9,30,31]. Second, they

are not confounded by BGS [1,32]. This insensitivity is a particularly important attribute

when studying genomic regions subject to different levels of BGS. Given that multiple

genomic factors correlate with this process, their association with selection can be

confounded if the summary statistic used is also sensitive to BGS. In this regard, the

integrated Haplotype Score (iHS) [5] is a haplotype-based statistic that has been

extensively tested [1,5,8,25], and can detect selection signals both from *de novo*

mutations and to some extent from standing genetic variation, provided that selection

started from not too high initial allele frequency [9,30]. This summary statistic can be

used to scan many genomes given the availability of fast implementations [33].

Therefore, we use iHS as a measure of recent positive selection across the human

genome. As shown in Figure 1 (see also Methods), iHS has a complex, asymmetric

distribution across the genome that is not well captured by classic distributions

(Supplemental Results S1: Figure S1). This raises an important limitation of violated

assumptions when using classic regression or correlation analysis to find the genomic

determinants of iHS, with for example the assumption of statistics following a gaussian

distribution clearly not met. Furthermore, classic regression and correlation analyses may

not capture well the process of recent positive selection through selective sweeps

measured by iHS. Indeed, classic regression and correlation assume a linear association

between iHS and a factor contributing to recent positive selection in a distributed way

across the entire genome. This neglects the localized nature of recent selection events,

that have likely affected only a limited genomic portion, while leaving the rest of the

genome (and likely substantial majority) without any correlation between iHS and a

contributing factor, even though the latter did contribute to selection occurring locally in

the genome. This creates a situation where only the higher values of a statistic such as

iHS, those that are more likely to represent selection, might correlate at all with genomic

factors influencing selection. In this respect, the distribution of iHS is particularly

interesting, with an upper tail that is much heavier than the lower tail. We can

hypothesize that localized selection in a limited portion of the genome might have

generated this heavy tail, rather than a scenario where more generalized selection

covering the entire genome would have shifted the whole distribution. What tools then

can we use to properly account for this complex distribution of iHS?

Here, we model the genomic determinants of recent positive selection in the
human genome. We measure recent selective sweeps with the iHS statistic from five
human populations represented in the 1,000 Genomes phase 3 dataset [34]. The iHS
statistic has more power to detect recent and incomplete sweeps, while it has reduced
power to detect complete sweeps or sweeps more than 30,000 years old [9,15]. Therefore,
we restrict our analyses to selection that occurred after the main human migration out of
Africa [35]. To account for the observed complexity of the distribution of iHS, we use
Mixture Density Regressions (MDRs) to test the association between iHS and several
genomic factors that are possible determinants of selection (Methods). MDRs can fit a
mixture of several distributions to the observed statistic. In our case, we fit a mixture of
two Gaussian-distributed components to the observed iHS (Methods). The first
component fitting lower values of iHS is expected to match more the distribution of iHS
under drift, the portion of the genome that was not affected by localized selection. The
second component fitting higher iHS values is expected to be comparatively enriched in
positive selection. Then, we look at the genomic factors that influence iHS in the
regression model, interpreting the significant association of a genomic factor with the
selection-enriched component as evidence for positive selection (Methods). This differs
from other approaches trying to understand the determinants of selection. Previous
studies have focused on the detection of classic partial correlations between selection and
genomic factors (e.g., Lohmueller et al. [36]). Given that the impact of hitchhiking on the
human genome is likely localized and not expected to influence the entire genome, the
signal of recent sweeps could be diluted enough across the genome so that this classic
approach misses true selection signals. An approach that specifically uses a selection-
enriched component should have more power to reveal the impact of positive selection.

As expected according to previous studies, we find a strong association between
recombination rate and recent positive selection signals in the human genome. In addition
to recombination, we find that other factors, such as the density of regulatory elements in
immune cells, GC-content, gene expression in immune cells, the density of mammal-
wide conserved elements and the distance to VIPs, all show strong associations with
recent positive selection. These results imply that recent positive selection had to be
common enough to create these associations, and further highlight how MDRs can be

used to clarify the prevalence and determinants of recent positive selection based on

genome scans.

Results

We first validate that the MDR with two gaussian distributions fits the distribution of iHS

much better than a single gaussian (Supplemental Results S1: Figure S1). The tight,

visually obvious fit between the observed iHS distribution and the two gaussians in our

MDR model (Figure 1) also shows that more than two gaussians are very likely not

needed and would negligibly improve the fit. We consider multiple genomic factors to

model the association with recent selection signals. These factors are possible

determinants of selection according to previous evidence [1,5,9,17,18,29,37,38]. We use

a gene-centric perspective where each individual protein-coding gene in the human

genome is associated with a set of factors measured by centering a genomic window on

each gene (Methods). This approach is likely to detect the influence of factors that differ

between genes, rather than only differences between genic and intergenic regions. The set

of factors in the MDR model includes several genomic features, like the length of the

gene at the center of the window or recombination estimates from the latest deCODE

genetic map (Methods). We obtain other genomic factors, like the GC-content and the

densities of genes, coding sequences and conserved elements. Moreover, we include the

density of transcription factor binding sites according to the hypersensitivity to DNaseI

and chromatin immunoprecipitation (ChIP-seq) experiments [39]. In the latter case, we

also obtain the density of regulatory elements for specific subsets of cell lines: testis and

immune cells. Table 1 provides a complete list of the factors considered, and the Methods

section details how these factors were obtained.

Table 1. Slopes and p-values of the association between iHS and genomic factors for the five studied populations (Yoruba,
Utah residents, Toscani, Han Chinese and Peruvians) in 1,000 kb windows within the selection-enriched component

Covariate	Yoruba		Utah residents		Toscani		Han Chinese		Peruvians	
	Slope	p-value	Slope	p-value	Slope	p-value	Slope	p-value	Slope	p-value
Intercept	-1.43	<1E-16	-1.77	<1E-16	-1.66	<1E-16	-1.78	<1E-16	-1.55	<1E-16
Recombination rate	-2.44	<1E-16	-2.8	<1E-16	-2.7	<1E-16	-2.51	<1E-16	-1.74	<1E-16
GC-content	0.55	1.34E-06	0.745	1.35E-08	0.516	2.47E-05	1.171	<1E-16	0.351	2.63E-03
Density of conserved elements	0.159	6.40E-04	0.379	7.74E-13	0.337	2.70E-11	0.311	4.20E-10	0.057	2.43E-01
Number PPIs	-0.07	4.25E-02	-0.05	1.95E-01	-0.04	2.67E-01	-0.09	1.92E-02	-0.04	3.00E-01
Distance to VIPs	-0.17	4.16E-06	-0.34	8.45E-14	-0.27	1.92E-10	-0.14	5.35E-04	-0.18	2.77E-05
Gene expression	-0.14	3.77E-02	-0.34	9.70E-06	-0.25	4.92E-04	-0.25	1.14E-03	-0.26	2.56E-04
Gene expression in immune cells	0.265	1.51E-05	0.408	6.87E-09	0.366	5.34E-08	0.307	9.51E-06	0.251	1.56E-04
Gene expression in testis	0.001	9.90E-01	-0.03	5.45E-01	-0.04	4.21E-01	0.065	2.06E-01	0.053	2.61E-01
Regulatory density (ChIP-seq)	-0.15	2.37E-01	0.593	3.45E-06	0.742	3.06E-10	0.49	3.92E-05	0.327	7.64E-03
Regulatory density in immune cells (ChIP-seq)	0.047	6.24E-01	-0.05	6.01E-01	-0.08	3.94E-01	-0.16	1.26E-01	0.146	1.32E-01
Regulatory density in testis (ChIP-seq)	-0.8	<1E-16	-1.03	<1E-16	-0.86	<1E-16	-0.39	1.65E-09	-0.11	4.76E-02
Regulatory density (DNaseI)	-0.05	7.15E-01	-0.61	5.45E-05	-0.64	8.24E-06	-0.84	1.66E-08	-0.55	8.27E-05
Gene length	-0.06	1.12E-01	-0.05	2.84E-01	-0.02	5.59E-01	0.004	9.13E-01	0.023	5.46E-01
Gene number	-0.09	3.46E-01	-0.52	1.28E-06	-0.4	5.91E-05	-0.46	6.23E-04	-0.02	7.96E-01
Coding density	0.159	8.41E-02	0.535	8.72E-06	0.586	3.77E-08	0.396	1.93E-03	0.202	2.73E-02
Number iHS data points	-0.01	8.19E-01	0.315	<1E-16	0.33	<1E-16	0.157	7.29E-08	0.081	1.51E-03

We also consider other functional factors like gene expression, including the

average gene expression across 53 GTEx v7 tissues [40]. The expression in testis and

immune cells could impact the frequency of sweeps, thus we also include them as

independent expression variables [5,29]. We also consider the number of protein-protein

interactions (PPIs) of each gene, given that this variable has been previously associated

with the rate of sweeps [38]. Finally, we consider the influence of viruses on genomic

adaptation, as previous evidence suggests that they have acted as a strong selective

pressure during recent human evolution [9,24]. We use the distance of each gene to the

closest gene coding for a protein that interacts with viruses. More details for all the

factors considered are given in the Methods. Note that our analysis is not expected to

detect all selective pressures affecting the human genome. We test factors that should

associate with positive selection according to prior strong evidence, using them to test the

ability of our new method to detect the genome-wide action of recent adaptation (see

Discussion).

Genomic factors that are calculated across the genomic windows can vary

depending on the window size used. Therefore, we infer the association between recent

selection and factors measured in genomic windows of varying sizes (Methods). In total,

we use five different window sizes (50, 100, 200, 500 and 1,000 kb) centered at the

genomic center of Ensembl v99 coding genes [41]. For example, we measure the

association between the average iHS within 50kb windows and the average

recombination rate within the same 50kb windows, or between the average iHS and

recombination within the same 1000kb windows (Methods). Using different window

sizes also relaxes assumptions about the expected strength of selection, as larger windows

are likely more sensitive to strong selection compared to smaller windows, and vice versa

[9]. Using fixed size windows irrespective of gene length avoids a priori biasing

measures of iHS by gene length. We also account for gene length in our model through

the inclusion of multiple functional densities along with the genomic length of the gene

(between transcription start and stop) at the center of the window as potential explanatory

variables.

We estimate recent selection signals using the iHS statistic in five populations

represented in the 1,000 Genomes Project [34]. Each population represents a different

continent: Yoruba for Africa, Han Chinese for East Asia, Utah residents with Northern

and Western European ancestry, Toscani for Southern Europe, and Peruvians for

Americas. We select Peruvians given that they show the highest percentage of Native

American ancestry among populations included in the 1,000 Genomes Project [34,42].

The iHS statistic measures how far the haplotypes carrying a specific derived allele

extend upstream and downstream of this focal allele, compared to how far haplotypes

carrying the ancestral allele extend [5]. As such, the approach provides one value of iHS

for each biallelic SNP in the genome with clearly defined derived and ancestral alleles

(Methods). Although isolated extreme values of iHS at a single SNP have often been

used as a signature of selection, we recently found that measuring the average absolute

value of iHS across all the SNPs in a genomic window provides high power to detect

recent selection [9]. The average iHS in an entire window is also likely to better associate

with other factors similarly measured as averages in the same windows. To account for

the fact that the variance of the average iHS depends on the number of SNPs with

individual iHS values in a window, we include this number as a predictor in the tested

MDR models.

MDR fit to recent selection signals

For each separate human population and window size, we model the association between

genomic factors and iHS using a MDR with a mixture of two Gaussian distributions. We

design one distribution to match more the drift component of iHS that is not expected to

correlate with genomic factors. In contrast, the second distribution is designed to be

sensitive to positive selection and it is expected to associate with the tested genomic

factors (Methods). We apply this approach to account for the fact that only a part of the

genome is expected to have undergone recent positive selection. In this context, a small

second distribution and non-significant associations with the tested predictive factors

would suggest a very small contribution of recent positive selection. The patterns of iHS

would then be entirely dominated by genetic drift and past demography. We use this

modeling approach together with an optimization algorithm to get the overall mixture

distribution with the best fit for the observed iHS in each population and window size

(Methods).

290   Overall, the MDR models fit well the distribution of observed iHS (Figure 1).

iHS does not follow a Gaussian distribution, and our flexible approach combining a

mixture of two Gaussian distributions and linear regression clearly fits better the iHS

distribution than would a single distribution from a classic linear model. We find a clear

separation between the Gaussian components, with the second component shifted toward

higher iHS values, and thus likely enriched in positive selection (Figure 1). Together, the

close fit and separation of the two Gaussian distributions likely provide an improved

ability of our model to estimate the contributions of different factors to iHS.

299   Patterns of selection across populations and window sizes

We find different patterns of selection across five populations from four different

continents (Figure 1). The African Yoruba population shows the largest selection-

enriched component (Figure 1A). This component is clearly shifted towards high iHS

values, indicating stronger selection signals that stand out more from the first component.

The selection-enriched components are less pronounced but still substantial in the four

other populations (Figure 1B-1E). The Peruvian population shows the smallest selection-

enriched component (Figure 1E). These results suggest that the visible (not necessarily

the actual) contribution of selection to the whole variation of iHS is smaller in

populations that were exposed to bottlenecks during migrations Out of Africa and

consequently, more subject to genetic drift [34,35]. In other words, our results might be

explained by the fact that selection is more visible in Yoruba due to a reduced effect of

genetic drift.

Figure 1. Mixture of Gaussian distributions fitting observed iHS (1,000 kb windows) for the five studied populations: A) Africa - Yoruba; B) Europe - Utah residents with Northern and Western European ancestry; C) Europe - Toscani; D) East Asia - Han Chinese; E) America - Peruvians. For each population, the figure shows two Gaussian distributions, components 1 and 2 of iHS, being the latter enriched in positive selection.

In that component, iHS linearly depends on the genomic factors considered. Legend:
Light blue = Observed iHS; Dark blue = Mixture model; Full red curve = Component 1
of the mixture model; Dashed red curve = Component 2 of the mixture model enriched in
positive selection.

With respect to window size, we find the largest selective components when using

larger, 1,000 kb windows (Supplemental Results S2). This is an expected result, because

it is easier to distinguish strong selection from genetic drift and background noise in

larger windows. Smaller windows are less sensitive to strong selection, while larger

windows can include a great accumulation of large iHS values. In addition, small

windows are more influenced not only by the content of a genomic window, but also by

genetically linked surrounding regions. These surrounding areas are in linkage with the

genomic areas inside smaller windows as their center is closer to the edges. Therefore,

smaller windows do not differentiate well between inside and outside genomic regions

[1,9]. Given these results, we highlight primarily the results for the Yoruba population in

the main text (while providing the results for all tested populations in main Tables and

Figures), and we use 1,000 kb windows.

Determinants of strong positive selection in human populations

Several genomic factors show significant associations with iHS in the selection-enriched

component (Table 1). Unsurprisingly, we find the strongest association with

recombination rate, which negatively correlates with iHS (e.g., Yoruba slope=-2.44; p-

value<1E-16; Table 1). This is congruent with the eroding effect of recombination on

selection signatures, impacting the probability to detect selective sweeps. Recombination

can break haplotypes generated by selection, making it more difficult to detect the

selective events [17,18]. These results confirm the role of recombination as a key

determinant of the probability of detecting selection with iHS [8]. We also find an

association between GC-content and recent positive selection. When the effect of

recombination is not accounted for, that is, recombination is not included in the model,

GC-content negatively correlates with iHS (Yoruba slope=-0.29; p-value=1.22E-03;

Supplemental Results S3: Table S1). This is expected given that GC-content is positively

associated with long-term recombination [43]. However, adding the control for

recombination reveals a positive association between GC-content and iHS (Yoruba

slope=0.55; p-value=1.34E-06; Table 1). Given that recombination already explains a

great proportion of iHS variability including that shared with GC-content, the positive

association between GC-content and iHS could be independent from recombination.

Accounting for the eroding effect of recombination on haplotypes could make more

visible a positive (direct or indirect) influence of GC-content on selection.

GC-content might be a better proxy of overall functional density than individual

functional factors included in the regression model, which could explain its positive

association with iHS. We test this hypothesis by removing different genomic factors from

the model (see Supplemental Results S3 for modeling results after removing different sets

of factors in Yoruba). The removal of GC-content makes emerge a positive association of

iHS with DNAseI hypersensitivity, which is a measure of the overall density of

transcription-factor binding sites (Yoruba slope=0.44 vs. -0.047; p-value=1.77E-07 vs.

7.15E-01; Table 1; Supplemental Results S3: Table S2). Since GC-content has been

positively associated with the density of functional elements [44,45], removing GC-

content may thus make more visible a positive association between iHS and functional

density. We further test if this is also the case for more tissue-specific functional densities

such as the density of regulatory elements in testis or immune cells that are expected to

exhibit more positive selection (Methods), but we find no such evidence. The removal of

GC-content does not affect to the association between selection and regulatory density in

testis (slope=-0.765 vs. -0.795; p-value<1E-16 in both cases; Table 1; Supplemental

Results S3: Table S2) and immune cells (slope=0.072 vs. 0.047; p-value=4.5E-01 vs.

6.23E-01; Table 1; Supplemental Results S3: Table S2). As shown later in this

manuscript, fine-scale patterns of recombination seem to hinder the detection of

associations for these regulatory variables. Finally, we find that the removal of coding

density and DNAseI hypersensitivity increases the significance for the association

between GC-content and iHS (p-value=2.40E-14 vs. 1.34E-06; Table 1; Supplemental

Results S3: Table S3). From this model, we then further remove additional functional

factors whose association with selection is also affected by the removal of GC-content,

namely, gene number, gene length and ChIP-seq regulatory density (Methods).

Accordingly, the removal of these functional factors leads to a highly significant

association between GC-content and recent positive selection (slope=0.49; p-value<1E-

16; Supplemental Results S3: Table S4). In summary, the removal of different functional

genomic factors from the model supports that the positive association between positive

selection and GC-content is mediated by the role of this factor as proxy of overall

functional density. However, we cannot exclude the implication of other processes

related to recombination that would require more detailed analyses (for example, see fine

scale patterns related to regulatory density below).

Another genomic factor associated with selection is the density of conserved

elements (Methods), showing a positive association with iHS (Yoruba slope=0.16; p-

value=6.4E-04; Table 1). This is an expected result since coding sequences and non-

coding regulatory elements tend to be conserved [37]. This factor shows a significant

association in all populations except in the Peruvian population sample, which might

reflect the impact of past bottlenecks on the visibility of the selective patterns (Table 1).

The density of conserved elements is thus likely to be a good proxy of overall (coding

and non-coding) functional density. The inclusion of this factor in the model could then,

however, explain the lack of association for other factors related to functional density

such as coding density. Indeed, the removal of conserved elements density makes

significant the positive association between coding density and selection in Yoruba

(Yoruba slope=0.26 vs. 0.16; p-value=2.35E-03 vs. 8.41E-02; Table 1; Supplemental

Results S3: Table S5). All other populations have a clear significant positive correlation

of iHS with coding density even without removing the density of conserved elements

(Table 1). In addition, the simultaneous removal of conserved elements density and GC-

content, which can also act as a proxy of functional density, increases even more the

positive association of coding density with iHS in Yoruba (slope=0.29; p-value=7.76E-

04; Supplemental Results S3: Table S6). These, and previous results about GC-content

suggest that the lack of association for multiple functional factors could be explained, at

least partially, by the simultaneous consideration of better measures of overall functional

density in our models.

The number of protein-protein interactions is another factor that correlates with
iHS. This factor associates negatively with selection, suggesting the existence of a slight
depletion of selection for regions with higher number of PPIs (Yoruba slope=-0.07; p-
value=4.25E-02; Table 1). This is opposite to that found by Luisi et al. [38], as they
reported higher recent positive selection on central proteins within the human
interactome. Higher number of PPIs has been also associated with higher positive
selection in the chimpanzee lineage according to the McDonald-Kreitman (MK) test.
However, this factor does not seems to be a determinant of positive selection when other
genomic factors are simultaneously considered using a MK regression approach [46].
Given the inconsistency with previous evidence and the low strength of the association,
the weak negative relationship between selection and the number of PPIs found in this
study should be taken with caution.

The modeling approach presented in this study can also be used to test the
influence of specific selective pressures. We illustrate this feature by showing that
positive selection is associated with multiple variables related to viral interaction (Table
1). One of the factors more highly associated with iHS is the distance to VIPs in all tested
populations (Yoruba slope=-0.17; p-value=4.16E-06; Table 1). Selection decreases
further away from VIPs, in other words, we find an enrichment of selection around VIPs.
Viruses have acted as a strong selective pressure during human evolution, shaping
genomic adaptation, and previous studies have also found more positive selection at VIPs
[9,24]. Gene expression is another functional factor significantly associated with
selection (Table 1). The average expression across 53 tissues from GTEx [40] is
negatively associated with iHS (Yoruba slope=-0.14; p-value=3.77E-02), while gene
expression in immune cells shows a strong and positive association with selection in all
populations (Yoruba slope=0.26, p-value=1.51E-05; Table 1). Given their relevance in
the response to pathogens, genomic regions highly expressed in immune cells do
represent expected targets of positive selection.

Overall, we find positive associations between the density of regulatory elements
and iHS except in Yoruba where this association is not significant (Table 1; see below for
analyses under neutral conditions explaining these results). However, the densities of

regulatory elements in tissues expected to experience more positive selection (testis and

immune cells) show surprising patterns of association with iHS (Table 1). In particular, a

higher density of transcription factor binding sites in testis correlates with weaker iHS

selection signals (slope=-0.79; p-value<1E-16). This is a counterintuitive result, given

that more regulatory density would give more options to modulate gene expression and

hence more room for positive selection to act. An explanation could be the following:

hotspots of recombination are usually regions where the chromatin is more open, and

hence more accessible for transcription factors to bind [47]. If the density of transcription

factor binding sites in testis (where meiosis and recombination occur) is higher in areas

with high recombination, then selection signals within these sites would be erased by

recombination [17,18]. Fine scale patterns of recombination and selection support this

hypothesis, as recombination tends to increase close to regulatory elements, while iHS

tends to decrease (Supplemental Results S4: Figures S1-S3). Therefore, high levels of

local recombination could erode haplotypes close to the selected variants, although the

average recombination in the gene window covering that region is low. In other words,

the 1,000 kb gene windows we use could have a low recombination rate on average,

while containing local peaks of recombination close to regulatory elements. Our original

model considers the average recombination in each 1,000 kb gene-centered window, so it

is likely unable to detect the effect of these fine-scale patterns of recombination. We also

observe a surprising lack of association between immune regulatory density and iHS

(Table 1), even though we find a strong positive association between iHS and immune

gene expression. Here again we find that the fine-scale patterns of recombination close to

immune regulatory elements are likely to blame, with an increase in the fine-scale

recombination rate as one gets closer to immune regulatory elements (Supplemental

Results S4: Figure S3). It is surprising that we find a positive association between iHS

and regulatory density across multiple tissues, but not when focusing on specific tissues

where selection is expected to act (i.e., testis and immune cells). Although recombination

rate tends to concentrate close to all regulatory elements, as it does more specifically

around regulatory elements in testis and immune cells (Supplemental Results S4: Figure

S1-S3), recombination declines rapidly when considering regulatory elements of all

tissues compared to testis and immune cells (Supplemental Results S4: Figure S4). This

suggests that we could have more power to detect positive associations with iHS and
regulatory density in all tissues compared to testis or immune cells (see below for further
evidence).

Robustness of recent selection patterns to varying genomic window sizes in Yoruba

Next, we ask if the trends observed with 1,000 kb windows in the Yoruba genomes also
hold or not when using smaller window sizes (Supplemental Results S2: Tables S1-S5).
Some patterns visible in large 1000kb windows are also visible when using smaller
genomic window sizes, while some are not (100 kb vs 1,000 kb windows; Supplemental
Results S2: Tables S2, S5). The strong negative association between recombination rate
and selection is found regardless of window size. On the contrary, the distance to VIPs
for example shows a strong association with iHS in large windows but not in smaller
windows (100 kb windows: slope=-0.02; p-value=5.11E-01; Supplemental Results S2:
Tables S2, S5). This is consistent with previous evidence showing that VIPs are
particularly enriched in strong selective sweeps, for which larger windows are more
sensitive [9]. This is also the case for the density of conserved elements (100 kb
windows: slope=0.1; p-value=9.3E-02) and especially regulatory elements in testis, with
a slope going from -0.79 in 1,000 kb windows to -0.03 in 100 kb windows (Supplemental
Results S2: Tables S2, S5). These results suggest that both factors may correlate with
strong rather than weak selection. In the case of testis regulatory density, this pattern
might be explained by the fact that smaller windows take better into account the fine
scale patterns of recombination, which limits the negative influence of using average
recombination (see below for fine-scale analyses of recombination). Note, however, that
all these differences might also be explained by the fact that smaller windows do not
differentiate well between inside and outside of the genomic regions they are supposed to
delineate. That is, they may be more influenced by the surrounding, genetically linked
genomic regions [1,9]. See Supplemental Results S2 for the complete results across
populations and window sizes.

Population simulations of neutral expectations of associations with iHS

As described above, we find multiple strong functional associations with iHS in the directions expected under recent positive selection. That said, some observations remain unexplained at this point, with for example a lack of positive association between overall regulatory density and iHS in Yoruba. We also observe a lack of association between the overall sum of regulatory plus coding functional density and iHS in Yoruba (slope=-0.063; p-value=0.65; Supplemental Results S5: Figure S1). The association of this overall coding plus regulatory density with iHS is strongly positive in the other tested populations (Supplemental Results S5: Figure S1). This prompted us to better characterize the null expectations of the associations between this overall coding plus regulatory density (as a measure encompassing the different functional types of elements considered in our model) and iHS in the absence of positive selection. In particular, the heterogenous, fine scale patterns of recombination and their variation around functional elements (Supplemental Results S4: Figures S1-S5) might make null neutral expectations deviate from a simple lack of association with slopes centered around zero.

We therefore use forward SLiM [48] neutral simulations to estimate the neutral expected associations between the overall coding plus regulatory density and iHS in our analysis (Methods). Under neutral conditions (i.e., only neutral mutations), and recreating the actual distribution of recombination rate and functional (coding plus regulatory) elements (Methods), we find that the association between iHS and functional density tends to be negative, and in every case much less positive compared to the studied populations including Yoruba (Supplemental Results S5, Figure S1). In addition, the magnitude of the second Gaussian component (estimated with p, see Methods) is also lower under neutral expectations compared to Yoruba (Supplemental Results S5: Figure S2). This suggests that the results observed with the MDR approach cannot be explained by neutral evolution. Moreover, these results support that, although the second Gaussian distribution captures neutral loci as shown by its presence even under strictly neutral evolution (Supplemental Results S5: Figure S2), this component is also sensitive to adaptation, supporting its enrichment in recent positive selection. These results also show that the lack of association between iHS and functional coding plus regulatory density in

Yoruba does not represent a lack of support for positive selection. Indeed, the null

expected distribution for this association is negative and the observed Yoruba association

is clearly above the neutral distribution (Supplemental Results S5: Figure S1). In the

discussion we mention multiple reasons why recent positive selection may have been not

less, or even more common in the Yoruba, while still having a weaker impact on iHS.

Fine scale patterns of recombination around regulatory elements

The increase of recombination rate around regulatory elements may explain the fact that

the null expectations for the association between functional density and iHS is negative

(Supplemental Results S4: Figures S1-S5). Our neutral simulations recreate the

distribution of recombination rate and functional (coding plus regulatory) elements of the

human genome, thus regulatory and coding elements tend to be close to recombination

peaks also in the simulations (Supplemental Results S4: Figure S5), leading to lower iHS

and lowering the null expectation. This led us to further analyze the influence of local

patterns of recombination around regulatory elements. Given the local increase of

recombination around regulatory elements, it is likely that whole window estimates of

recombination are not a sufficient measure of its effect on the association between iHS

and regulatory densities. In particular, this might explain the surprising negative

association between iHS and testis regulatory density or the lack of association for

immune regulatory density (Table 1). Indeed, recombination rate tends to be higher

specifically around these regulatory elements compared to the whole set of regulatory

elements (Supplemental Results S4: Figure S4). We use three variables related to the

recombination around regulatory elements in immune cells, testis and across multiple

tissues, respectively. We calculate the more local, average recombination around

regulatory elements within each gene window, up to a maximum distance of 5kb from

each side of regulatory elements. Each variable is included in the original model of

Yoruba 1000 kb separately, and replaces the previous, window-wide average

recombination. Given that some gene windows can overlap with more regulatory

elements than others, we also consider the number of recombination data points obtained

for regulatory elements inside each gene window. In this way, we account for the fact

that recombination may vary more broadly in windows that have less regulatory

elements. See Supplemental Results S4 for further details about these calculations.

The iHS statistic decreases around regulatory elements in immune cells while

recombination increases, staying very high further from immune regulatory elements

compared to other tissues (Supplemental Results S4: Figures S1-S4). This might have

confounded the expected positive association with selection in the original regression

model with window-wide recombination rates. The control for local patterns of

recombination around immune regulatory elements should however increase our power to

detect this association. In line with this prediction, we find that the association between

immune regulatory density and iHS becomes positive and highly significant when

considering the local patterns of recombination around immune regulatory elements

(Yoruba slope=0.68 vs. 0.047, p-value=4.60E-14 vs. 6.24E-01; Table 1; Supplemental

Results S4: Table S1). In order to assess whether this increase is caused just by the lack

of consideration of recombination patterns at a window-wide scale, we repeat this

analysis but including the original window-wide average recombination variable. After

including the average recombination rate at window-wide scale, the positive association

remains stronger compared to the original model (slope=0.28 vs. 0.047, p-value=5.72E-

03 vs. 6.24E-01; Table 1; Supplemental Results S4: Table S2).

As explained above, we hypothesize that the confounding effect of recombination

is caused by a mismatch between different scales. Gene windows with low recombination

can have local peaks of recombination around transcription factor binding sites

(Supplemental Results S4: Figures S1-S4), as these peaks are more accessible to

transcription factors [47]. Therefore, recombination could erode signals of positive

selection around regulatory elements even if the region has an overall low recombination.

In contrast, regions with high recombination at the average window-wide level should

suffer less from this confounding effect, as recombination rate is then high both at a local

and window-wide scale. Therefore, focusing on high-recombination regions should

improve our ability to detect the expected positive association between selection and

regulatory density. This approach, however, has the caveat of reducing the power to

detect positive selection due the higher probability that recombination breaks selected

haplotypes. We nevertheless repeat the analysis focusing only on gene windows with a

recombination rate equal or higher than the second tertile (1.552 cM/Mb). Focusing on

high-recombination regions, regulatory density in immune cells shows a stronger

association with selection in a model including recombination rate around both gene

windows and immune regulatory elements compared to the same model run across all

genes (Yoruba slope=0.85 vs, 0.28, p-value=1.27E-08 vs. 5.72E-03; Supplemental

Results S4: Tables S2-S3). Interestingly, regulatory density in testis also shows a positive

association with selection in a model including recombination around both testis

regulatory elements and gene windows and focused on high-recombination regions. Note

that this regulatory variable shows one of the strongest negative associations with

selection in the original model (Yoruba slope=0.58 vs. -0.79, p-value=4.98E-05 vs. <1E-

16; Supplemental Results S4: Tables S4-S6), an unexpected result given the role of

regulatory sequences in positive selection [1]. These results suggest that, indeed, a higher

regulatory density can increase the probability for positive selection to occur, especially

in some tissues. However, this signal is distorted by fine-scale patterns of recombination,

which have to be carefully taken into account.

Discussion

We find several functional factors independently associated with recent positive

selection. For instance, viral interactions along with gene expression and regulatory

density in immune cells are strongly and positively associated with selection. Viruses

have acted as key drivers of adaptation in humans and VIPs show higher expression in

lymphocytes compared to non-VIPs [26,28,49]. These results confirm the existence of

strong viral and immune selective pressures during recent human evolution [9,24,28]. We

also detect a positive association between testis regulatory density and selection, which is

congruent with the fact that reproduction-related functions show signals of positive

selection [5,29]. Note, however, that this latter result has been more difficult to reveal in

our analyses, as it is only visible after controlling for fine-scale patterns of recombination

around testis regulatory elements and after focusing on high-recombination regions, thus

it is more questionable. Finally, the density of conserved elements and GC-content,

which are related to overall (coding and non-coding) functional density [37,44,45], also

show a positive association with selection. In summary, recent and positive selection is

associated with the distribution of functional regions across the human genome.

If sweep signals are false positives only reflecting genetic drift, the signatures

of sweeps should occur randomly across the genome. In contrast, if selective sweeps

were relatively common during recent human evolution, we expect that other factors on

top of recombination should associate with sweep signals across the human genome. In

that case, we expect that selective sweeps should occur more frequently around

functional (coding or non-coding) elements where adaptive mutations are expected to

take place [1], also around genomic regions associated with specific functions. We find

strong evidence for the latter scenario, as multiple functional factors are associated with

recent positive selection. In other words, the distribution of sweep signals is not random

relative to the distribution of functional elements and is associated with the functional

characteristics of the human genome. That said, stochasticity still plays a role in the

occurrence of advantageous mutations in the first place, and we still expect randomness

when considering which specific loci of the human genome with equivalent functional

characteristics have experienced positive selection. Our results thus support that selective

sweeps were at least common enough during recent human evolution to generate

associations between selection and functional factors.

Our MDR approach shows a good fit to the distribution of iHS across the human

genome, with its two components showing a clear differentiation and the second being

enriched in positive selection. This suggests that the consideration of a more biologically

meaningful model, which assumes two Gaussian distributions instead of one, works

better to analyze recent adaptation than simple correlations. Given that the MDR

approach represents more directly genome evolution, the associations detected by this

approach are more informative than classic linear models and partial correlations. The

associations detected by these classical approaches may be caused by a deficit of weak

iHS values, not by an enrichment of high iHS values as shown in the second component

of our MDR approach, a scenario more specifically expected under strong selection.

Indeed, we cannot replicate our results using classic linear models and partial correlations

in 1,000 kb windows. These classic approaches show much lower correlations between

iHS and functional factors expected to associate with positive selection (Supplemental

Results S1: Tables S1-S2). For example, they are unable to replicate the strong

enrichment in positive selection around VIPs reported by previous studies [9,24]. This is

also expected given the poor fit of a single Gaussian distribution to the whole iHS

distribution compared to our approach assuming two Gaussian distributions

(Supplemental Results S1: Figure S1). Therefore, our results support the consideration of

two distributions to model positive selection in the human genome. The characteristics of

our MDR approach make it possible to reveal expected, but also unexpected patterns. For

instance, GC-content is strongly and positively associated with iHS. Regions with high

GC-content also tend to have high long-term recombination rates [43], which would

predict a negative association between iHS and GC content, not positive. A possible

explanation for this pattern is that the MDR approach can control for recombination as a

confounder by simultaneously analyzing recombination rate and other genomic factors.

Therefore, it can control for the impact of recombination on the probability to detect

selection. This can unravel new patterns, like a positive association between GC-content

and sweep signals. This association only emerged after adding recombination in the

model, becoming even more significant after the removal of functional factors like

coding density, gene number or regulatory density. This supports the independence of

this association from recombination and the role of GC-content as a better proxy of

overall functional density than individual factors. These results illustrate the utility of our

approach. The simultaneous consideration of multiple predictors lets us unravel novel

associations between genomic factors and adaptation. In addition, it enables easier testing

of different hypotheses by modifying the set of predictors considered. Therefore, the

MDR approach is useful to model recent genomic adaptation, and to analyze its

prevalence and determinants.

We find other unexpected results using the MDR approach. For instance, we do

not find a positive association between selection and testis or immune regulatory density

in the original model. This is an unexpected result given that a higher regulatory density

would lead to a larger mutational target for adaptation through changes in expression and

these tissues are expected targets of positive selection [5,29]. As previously posited, the

adaptive signals associated with regulatory density could be eroded due to the higher

tendency of transcription factors to bind DNA at recombination hotspots [47]. This is

supported by the higher recombination and lower iHS found closer to regulatory

elements, along with the fact that the positive associations for immune and testis

regulatory density only emerge when considering the fine-scale patterns of

recombination. These results support the role of regulatory sequences as targets of

positive selection independently of other genomic features [1], along with the relevance

of testis and specially immune cells in recent human evolution [5,29]. Our results also

illustrate how the detection of selection can be hindered by fine-scale characteristics of

the genome, like the sharp increase of recombination we observe around regulatory

elements. This might mask selection signals around regulatory elements, but not only

because of the breaking of haplotypes caused by recombination. An additional

mechanism might be the existence of biased gene conversion in these recombinant

regions, which could specifically favor GC mutations (gBGC) [43,50,51]. This could lead

to an increase in the frequency of GC mutations that are associated with short haplotypes

(due to the eroding effect of recombination). BGC might therefore make it even more

difficult to detect recent positive selection around regulatory regions, where it is expected

to happen [1]. Note, however, that the present study does not provide evidence about the

implication of gBGC in the observed patterns. It can be difficult to disentangle the

eroding effect of recombination from GC-biased gene conversion on haplotypes because

gBGC is caused by and therefore very correlated with recombination. Distinguishing the

specific effect of biased-gene conversion may require precise recombination maps for

African populations, where we find the largest selection-enriched second Gaussian

distribution. Although we have not found evidence of the implication of gBGC yet, the

fact that this hypothesis has emerged from implementing our model illustrates its utility

to propose new hypotheses about the different processes that influence the detection and

occurrence of positive selection.

708   Although the MDR approach provides an excellent fit to the observed selection

statistic, we may still not detect associations with specific factors because they do not

have a simple linear relationship with recent selection. Although more biologically

meaningful than a model assuming just one distribution, our approach is still linear, thus

more complex relationships may not be detected. For example, the weaker association of

iHS with coding density in Yoruba could be explained not only by the consideration of

better proxies of overall functional density, but also by the existence of non-linear

relationships with selection. Advantageous mutations are more likely to appear in regions

with high coding density, thus a high density of coding sequences should favor the

appearance of selection [1]. However, high coding density can also mean a higher

frequency of deleterious mutations. These mutations can interfere with adaptation when

they are in linkage disequilibrium with advantageous mutations. This would be especially

relevant for genomic regions with low recombination, where the linkage disequilibrium

between deleterious and advantageous mutations is more likely [19,52]. Therefore, a

higher coding density could favor or hinder positive selection depending on the

circumstances, complicating the detection of an association between coding density and

selection with our approach. The existence of this genetic interference could also explain

the different results observed across populations in relation to coding density. Despite

having the largest selection-enriched component, the Yoruba population shows a weaker

association between selection and coding density compared to other populations. This

association becomes visible only after removing proxies of overall functional density

(i.e., density of conserved elements and GC-content). A potential, but still speculative

explanation at this point might be the fact that the bottleneck related to the migration out

of Africa may have removed segregating recessive deleterious variants, which may

reduce their interference over advantageous mutations, thus making it easier for positive

selection to act on regions with a high coding density [52]. In contrast, there was no

similar bottleneck to decrease the genetic interference in the Yoruba, and regions of the

genome with high a coding density may still experience stronger interference from

recessive deleterious mutations. Interestingly, despite this potential limitation of positive

selection, we still see a larger selection-enriched component in the entire Yoruba genome,

suggesting that selection might still be more abundant in this population. There are other

possible explanations for this apparent contradiction. For example, the ability of iHS to

detect selective variants could be more limited in African populations due to lower

overall linkage disequilibrium, together with the larger genetic diversity in these

populations, which could favor the existence of sweeps from standing genetic variants

that are harder to detect with iHS [30]. These potential explanations are speculative at

this point and will require further investigation. Our results nevertheless demonstrate that

the complex and heterogenous distributions of multiple genomic factors, together with

summary statistics of hitchhiking robust to background selection, need to be used in order

to quantify the determinants and importance of recent positive selection in the human

genome.

Methods

Window sizes and genes coordinates

All analyses were performed using hg19 genomic coordinates and protein-coding gene

annotations from Ensembl v99 [41]. We selected hg19 instead of GRCh38 mainly

because this was the assembly used by the 1,000 Genomes consortium to generate and

phase the genetic variants considered in our analysis. Note that the assembly selection

should not influence our results as we performed genome-wide scale analyses, thus

avoiding focusing on specific regions where the assembly could change after hg19. We

considered windows of different sizes (50, 100, 200, 500 and 1,000 kb). Each window

was centered at the genomic center of a gene, in the middle between the most upstream

transcription start site and the most downstream transcription end site. We used a fixed

window size to avoid biases related to gene length, as larger genes are more likely to

overlap with high local iHS values just by chance compared to shorter genes. This would

bias power in favour of larger genes, increasing the probability to detect sweeps in them

just because of their length. In addition, the consideration of different window sizes

provides the opportunity to detect different types of sweeps as larger windows are

specifically sensitive to strong selective sweeps compared to smaller windows [9].

Genomic features

Multiple genomic factors were calculated inside the gene windows. We considered the

following factors that are likely to influence the frequency of sweeps:

•   Length of the gene at the center of each window.

•   Number of genes overlapped with each window. We used this value as an estimate
of gene density.

•   Recombination rate. For each window, we calculated the genetic distance between
the edges of the window and then divided it by the physical distance between them.
Genetic position of each window edge was obtained from the deCODE 2019 genetic
map [53]. In case no data was available for a window edge, we selected the closest
genetic position data points within 50 kb at each side of the window edge to estimate
its genetic position. We used linear interpolation to this end, considering the genetic
and physical position of the two points around the window edge. In other words, we
assumed that genetic distance increased linearly between the two selected data
points, and hence we can estimate the genetic position of the window edge based on
its physical position respect to the other two positions. If no genetic position data
was available at 50 kb or closer to the window edge, no genetic position was
calculated and the whole window was discarded for recombination calculations and
subsequent analyses. Note that we searched for genetic position data points only up
to 50 kb from each window edge to avoid that the total distance between points at
each edge could be too large (especially for larger windows). In that scenario, linear
interpolation could be inappropriate because we would use very distant data to
calculate recombination rate. For example, if a 1,000 kb window had no genetic
position for any of its edges and we looked for genetic positions up to 1,000 kb at
each side, it would possible that the selected data points of genetic position
downstream and upstream of the window were separated by 3,000 kb. In addition,
the recombination rate would not correspond with the size of the window.

•   Density of coding sequences. We used Ensembl v99 coding sequences. The density
was calculated as the proportion of coding bases respect to the whole length of the
window without considering gaps (gap locations obtained from the UCSC Genome
Browser; https://genome.ucsc.edu/;

http://hgdownload.cse.ucsc.edu/goldenPath/hg19/database/). A similar approach was

used for the rest of density estimates.

• Density of mammalian phastCons conserved elements [37], also downloaded from

the UCSC Genome Browser. Given that each conserved segment had a score, we

considered as conserved only those segments above a given threshold. In order to

minimize the inclusion of non-conserved elements, we used a threshold that only

considers 4.17% of the genome as conserved, as it is unlikely that much more than

that is strongly constrained [37].

• Density of regulatory elements. We calculated several variables related to the

density of binding sites for transcription factors. In all cases, we calculated the

proportion of sequences that are considered binding sites within a window. The data

were also obtained from the UCSC Genome Browser [39].

– Density of DNaseI hypersensitive sites (wgEncodeRegDnaseClusteredV3

track). We considered as binding sites those segments with a score higher

than a given threshold. The selected threshold was a score value according to

which only 10% of the genome is considered accessible to DNaseI, and

hence a binding site for transcription factors. In this way, we minimize the

probability to consider non-binding sites (i.e., false positives).

– Density of binding sites according to the technique of chromatin

immunoprecipitation followed by sequencing (ChIP-seq;

encRegTfbsClustered track). We calculated a threshold to consider a given

segment as a binding site following the same approach used for DNaseI

hypersensitive sites. This database includes 1264 experiments representing

338 transcription factors in 130 cell types. We calculated the density of

binding sites using the whole dataset, but also considering two subsets, one

for experiments performed on cells lines of testis, and another for

experiments performed on immune cells (lymphocytes in most cases). Note

that the threshold was calculated considering the 1264 experiments, being

then applied to the whole dataset and the two subsets.

•   GC-content, calculated as a percentage per window. It was also obtained from the
UCSC Genome Browser.

•   Gene expression: We used the log (base 2) of TPM (Transcripts Per Million), which
was obtained from GTEx [40] (https://www.gtexportal.org/home/). We considered
the average gene expression across 53 GTEx v7 tissues, but also expression in
immune cells (lymphocytes) and testis.

•   Number of protein-protein interactions (PPIs) in the human protein interaction
network [38]. We used the log (base 2) of the number of PPIs of each gene. We
summed 1 to the original dataset to avoid problems applying the logarithm to genes
with no interactions (i.e., PPI=0).

•   Proportion of genes that interact with viruses (VIPs). We used a previously
published dataset [9] that includes around 4500 VIPs with evidence of physical
interaction with viruses. 1920 of these VIPs were manually curated from the
virology literature, while the remaining 2600 VIPs were identified using high-
throughput methods and retrieved from the VirHostnet 2.0 database and additional
studies. See the original publication for further details [9]. The variable included in
the model was calculated as the distance of each gene to the closest gene that
interacts with viruses.

iHS calculation

In order to calculate iHS, we used polymorphism data from the 1,000 Genomes Project
phase 3 [34]. We calculated iHS for 5 populations: Yoruba, Utah residents with Northern
and Western European ancestry, Toscani, Han Chinese and Peruvians. We used the
hapbin software [33] in order to perform fast scans of iHS across the genomes of these
populations. Note that the genetic maps used as input in hapbin were calculated using the
deCODE 2019 genetic map [53]. In case that no data was available for the focal variant,
we again used linear interpolation to estimate its genetic position. We followed the same
approach used to calculate the genetic position of window edges during recombination

calculations. In this case, we searched for genetic position data points at both sides of the
focal variant until 1,000 kb. We discarded variants with no data points around, as they
were present in genomic regions with low density of recombination data. The genetic
maps based on the deCODE 2019 map were used as an input in order to calculate the iHS
for the 5 populations.

The raw iHS values obtained with hapbin were standardized by frequency. For
each population, we divided variants in 50 bins of frequency (from 0 to 1) considering the
whole genome. We then calculated the mean and standard deviation of raw iHS within
each bin, using them to standardize raw iHS values: (raw iHS - mean iHS) / sd iHS. The
mean of standardized iHS was calculated per window, considering the absolute value of
iHS. In our case, large positive iHS values can be caused by unusually long haplotypes
carrying the derived or the ancestral allele, in the case the ancestral alleles hitchhiked
together with the actual, selected derived allele. Therefore, high iHS can be interpreted as
selection in favor of the derived or ancestral haplotype. We also obtained the number of
iHS values per window to have an estimate of data density. Mean iHS in windows with
low number of iHS values can be more influenced by outliers (very low or very high iHS
values). Therefore, we included the number of iHS values per window as predictor in the
models to control for this.

Modeling

We modeled the association between genomic factors and iHS in each population and
window size by combining a Gaussian mixture distribution with linear regression.
Mixture distributions are useful to model data that cannot be fully described by a single
distribution and that are likely generated by multiple processes. In this case, a mixture of
two Gaussian distributions could fit better a scenario where selection influences some
genomic regions but not others, and hence a bimodal distribution of iHS would be
observed across the genome. Therefore, we posit that sweep signals in the human genome
may have components differentially influenced by positive selection. According to this,
we assumed that observed log iHS followed a mixture of two Gaussian distributions

representing two components, one more influenced by drift and a second one enriched in

positive selection. Because selective sweeps often result in elevated iHS, we assumed that

the Gaussian distribution for the selection-enriched component had a higher mean than

that for the first component. Furthermore, to model the association of genomic factors

with the selection-enriched component of iHS, we assumed that the probability of the

selection-enriched component was a linear combination of genomic factors followed by a

sigmoid transformation. Specifically, for each factor, we calculated the product of its

observed value with a slope indicating its association with the selection-enriched

component of iHS. A positive slope indicates that the probability of the selection-

enriched component increases with the increasing value of a factor, whereas a negative

slope indicates the probability decreases with the factor. Then, we summed the product

over all factors for a given gene to model the accumulative effect of genomic factors on

the probability of the selection-enriched component. Then, we added an intercept

indicating the baseline level of selection when all genomic factors are equal to 0 and

transformed the result to a probability (0 to 1) using a sigmoid function. This probability

was obtained for each gene and can be regarded as the overall influence of all genomic

factors on the frequency of recent selective sweeps. These steps are described by the

equation $p = \sigma\left(\sum b_i * x_i + a\right) = \frac{1}{1+exp\left(-\sum b_i * x_i - a\right)}$, where p is a probability summarizing

the influence of all studied genomic factors on the selection-enriched component of iHS,

σ is the sigmoid function, b_i and x_i are the slope and the observed value for genomic

factor i in a given gene, respectively, and a is the intercept.

To obtain the likelihood of log iHS in each gene, we calculated a weighted sum of

Gaussian density functions, $P(Y) = p * f(Y|\mu_1, \sigma_1) + (1 - p) * f(Y|\mu_0, \sigma_0)$, where Y

and $P(Y)$ are the observed log iHS and its likelihood in a given gene, respectively, p is the

probability of the selection-enriched component calculated in the previous step,

$f(Y|\mu_1, \sigma_1)$ is the Gaussian density function with mean μ_1 and standard deviation σ_1

describing the selection-enriched component, and $f(Y|\mu_0, \sigma_0)$ is the Gaussian density

function with mean μ_0 and standard deviation σ_0 describing the first component. Finally,

we summed the logarithm of $P(Y)$ over all genes to obtain the log likelihood of the entire

dataset.

We estimated model parameters by maximizing the log likelihood of the entire

dataset (L-BFGS-B optimization). These parameters included the mean and standard

deviation of each Gaussian distribution, along with the intercept and the slopes

representing the association between the selection-enriched component of iHS and each

genomic factor. For each genomic factor, we tested the significance of its slope by

comparing the log likelihood between the full model with all genomic factors and a

nested null model without the selected factor. We assumed that two times the difference

in log likelihood between the two models followed a chi-square distribution with one

degree of freedom. The script and input data needed to run the MDR are publicly

available at https://github.com/dtortosa/Mixture_Density_Regression_pipeline.

Population simulations to recreate the neutral expectation

We used SLiM [48] to simulate the neutral expectations in our analyses and assess in that

way if our results could be fully explained by neutral patterns. First, we randomly

selected 100 genomic segments of 20 mb each one, totaling 2000 mb. We recreated the

functional (coding and regulatory) elements observed in the human genome using the

same coordinates considered in the calculation of functional densities for the main

analyses (see Genomic features section). We also considered the same recombination

map (deCODE 2019) to recreate the actual recombination heterogeneity in these

segments. Therefore, we simulated a total of 2000 mb using the observed distribution of

functional elements and recombination rate. We only simulated neutral mutations in all

the simulated genomic regions. The population size was set to 10,000 individuals. We

first run a burn-in period of 100,000 generations, running then 100 independent neutral

simulations with 2,000 additional generations. For each independent neutral simulation,

we calculated iHS using the same approach than in the actual genomes. As the simulated

segments had direct correspondence with the human genome, we extracted the genes

included in each segment, and calculate 1000 kb windows centered around them. Inside

each window, we calculated the average of iHS and the number of iHS data points, along

with recombination rate and functional density as done in the actual genomes. Finally, we

used the MDR to model iHS as a function of recombination rate, functional density and

the number of iHS data points measured all in 1000 kb windows. Therefore, we model

iHS under neutral conditions across 100 independent runs simulating each one 2000 mb.

In addition to the genomic factors already mentioned, we also included the GC-content

observed for each gene window in the actual genome. It can be useful to simulate GC to

control for varying mutations rates, however we used a uniform mutation rate across the

simulated regions, so we did not include it in the simulations. We are still adding it as a

predictor using the GC-content in the actual gene windows because this factor is

associated with both functional density and long-term recombination [43–45]. This can

help to make visible more local patterns of recombination. In addition, it makes fairer the

comparison between the neutral simulations and the observed genomes, as in the latter

iHS was also modeled using GC-content.

From the model of each neutral run, we obtain the slope for the association

between functional density and iHS, while controlling for the rest of genomic factors

calculated. We also took p, i.e., the accumulative effect of all genomic factors on the

probability of the second component of iHS (see Modeling section). We calculated its

average across all genes within each neutral run, using it as a measure of the magnitude

of the second component of iHS. We compared the functional associations and the

magnitude of the second Gaussian component between the neutral expectations and the

actual genomes. Note that in the latter comparison, we only considered Yoruba. As

previously noted, the distribution of iHS seems to be more influenced by drift in non-

African populations due to their greater exposition to bottlenecks. This explains the

reduction of the second Gaussian component in these populations. Indeed, the population

that suffered the most extreme bottlenecks is the one showing the smallest second

component, i.e., Peruvians. Our simulations reproduce a constant population size of

10,000 individuals and no bottlenecks, thus it is fairer to compare the magnitude of the

second Gaussian component in the simulations with the African Yoruba population,

which was not exposed to bottlenecks in the same degree than the rest of studied

populations.

Competing interest statement

The authors declare no competing interests.

Acknowledgments

We thank the members of the Enard, Masel and Gutenkunst labs for their helpful comments. DFST was supported by a Marie S. Curie Global Fellowship within the European Union research and innovation framework programme (2014-2020; ClimAHealth; https://doi.org/10.3030/101030971). DE was supported by the National Institute of General Medical Sciences of the National Institutes of Health under award number R35GM142677 (https://www.nigms.nih.gov/). Y-FH was supported by the National Institute of General Medical Sciences of the National Institutes of Health under award number R35GM142560 and by startup funds from Pennsylvania State University (https://www.nigms.nih.gov/; https://www.psu.edu/). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health or the European Union. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Author contributions

Y-FH, DE and DFST designed the study. DE and DFST collected the data. Y-FH and DFST performed the analyses. DE and DFST wrote the paper and interpreted the results. Y-FH, DE and DFST edited and approved the paper.

References

1.   Enard D, Messer PW, Petrov DA. Genome-wide signals of positive selection in human evolution. Genome Res. 2014;24: 885–895. doi:10.1101/gr.164822.113

2.   Hernandez RD, Kelley JL, Elyashiv E, Melton SC, Auton A, McVean G, et al.

Classic selective sweeps were rare in recent human evolution. Science (80-).

2011;331: 920–924. doi:10.1126/science.1198878

3.   Akey JM, Zhang G, Zhang K, Jin L, Shriver MD. Interrogating a high-density SNP

map for signitures of natural selection. Genome Res. 2002;12: 1805–1814.

doi:10.1101/gr.631202

4.   Tang K, Thornton KR, Stoneking M. A new approach for using genome scans to

detect recent positive selection in the human genome. PLoS Biol. 2007;5: 1587–

1602. doi:10.1371/journal.pbio.0050171

5.   Voight BF, Kudaravalli S, Wen X, Pritchard JK. A map of recent positive

selection in the human genome. PLoS Biol. 2006;4: 0446–0458.

doi:10.1371/journal.pbio.0040072

6.   Sabeti PC, Varilly P, Fry B, Lohmueller J, Hostetter E, Cotsapas C, et al. Genome-

wide detection and characterization of positive selection in human populations.

Nature. 2007;449: 913–918. doi:10.1038/nature06250

7.   Fumagalli M, Sironi M, Pozzoli U, Ferrer-Admettla A, Pattini L, Nielsen R.

Signatures of environmental genetic adaptation pinpoint pathogens as the main

selective pressure through human evolution. PLoS Genet. 2011;7.

doi:10.1371/journal.pgen.1002355

8.   Johnson KE, Voight BF. Patterns of shared signatures of recent positive selection

across human populations. Nat Ecol Evol. 2018;2: 713–720. doi:10.1038/s41559-

018-0478-6

9.   Enard D, Petrov DA. Ancient RNA virus epidemics through the lens of recent

adaptation in human genomes. Philos Trans R Soc London Ser B. 2020;375:

20190575. doi:10.1098/rstb.2019.0575

10.   Schrider DR, Kern AD. Soft sweeps are the dominant mode of adaptation in the

human genome. Mol Biol Evol. 2017;34: 1863–1877. doi:10.1093/molbev/msx154

11.   Hejase HA, Mo Z, Campagna L, Siepel A. A Deep-Learning Approach for

Inference of Selective Sweeps from the Ancestral Recombination Graph. Mol Biol

Evol. 2022;39: 1–18. doi:10.1093/molbev/msab332

12. Gower G, Picazo PI, Fumagalli M, Racimo F. Detecting adaptive introgression in

human evolution using convolutional neural networks. Elife. 2021;10: 1–45.

doi:10.7554/eLife.64669

13. Mughal MR, Koch H, Huang J, Chiaromonte F, DeGiorgio M. Learning the

properties of adaptive regions with functional data analysis. PLoS Genetics. 2020.

doi:10.1371/JOURNAL.PGEN.1008896

14. The International HapMap Consortium. A haplotype map of the human genome.

Nature. 2005;437: 1299–1320. doi:10.1038/nature04226

15. Sabeti PC, Schaffner SF, Fry B, Lohmueller J, Varilly P, Shamovsky O, et al.

Positive natural selection in the human lineage. Science (80-). 2006;312: 1614–

1620. doi:10.1126/science.1124309

16. Pritchard JK, Pickrell JK, Coop G. The Genetics of Human Adaptation: Hard

Sweeps, Soft Sweeps, and Polygenic Adaptation. Curr Biol. 2010;20: R208–R215.

doi:10.1016/j.cub.2009.11.055

17. O'Reilly PF, Birney E, Balding DJ. Confounding between recombination and

selection, and the Ped/Pop method for detecting selection. Genome Res. 2008;18:

1304–1313. doi:10.1101/gr.067181.107

18. Booker TR, Yeaman S, Whitlock MC. Variation in recombination rate affects

detection of outliers in genome scans under neutrality. Mol Ecol. 2020;29: 4274–

4279. doi:10.1111/mec.15501

19. Hill WG, Robertson A. The effect of linkage on limits to artificial selection. Genet

Res. 1966;8: 269–294. doi:10.1017/S0016672300010156

20. Castellano D, Coronado-Zamora M, Campos JL, Barbadilla A, Eyre-Walker A.

Adaptive evolution is substantially impeded by hill-Robertson interference in

Drosophila. Mol Biol Evol. 2016;33: 442–455. doi:10.1093/molbev/msv236

21. Presgraves DC. Recombination enhances protein adaptation in Drosophila
melanogaster. Curr Biol. 2005;15: 1651–1656. doi:10.1016/j.cub.2005.07.065

22. Barreiro LB, Laval G, Quach H, Patin E, Quintana-Murci L. Natural selection has
driven population differentiation in modern humans. Nat Genet. 2008;40: 340–
345. doi:10.1038/ng.78

23. McVicker G, Gordon D, Davis C, Green P. Widespread genomic signatures of
natural selection in hominid evolution. PLoS Genet. 2009;5: e1000471.
doi:10.1371/journal.pgen.1000471

24. Souilmi Y, Lauterbur ME, Tobler R, Huber CD, Johar AS, Varasteh Moradi S, et
al. An ancient viral epidemic involving host coronavirus interacting genes more
than 20,000 years ago in East Asia. Curr Biol. 2021;31: 1–11. Available:
https://doi.org/10.1016/j.cub.2021.05.067

25. Barreiro LB, Ben-Ali M, Quach H, Laval G, Patin E, Pickrell JK, et al.
Evolutionary dynamics of human toll-like receptors and their different
contributions to host defense. PLoS Genet. 2009;5.
doi:10.1371/journal.pgen.1000562

26. Enard D, Cai L, Gwennap C, Petrov DA. Viruses are a dominant driver of protein
adaptation in mammals. Elife. 2016;5: e12469. doi:10.7554/eLife.12469

27. Deschamps M, Laval G, Fagny M, Itan Y, Abel L, Casanova JL, et al. Genomic
Signatures of Selective Pressures and Introgression from Archaic Hominins at
Human Innate Immunity Genes. Am J Hum Genet. 2016;98: 5–21.
doi:10.1016/j.ajhg.2015.11.014

28. Enard D, Petrov DA. Evidence that RNA Viruses Drove Adaptive Introgression
between Neanderthals and Modern Humans. Cell. 2018;175: 360–371.
doi:10.1016/j.cell.2018.08.034

29. Nielsen R, Bustamante C, Clark AG, Glanowski S, Sackton TB, Hubisz MJ, et al.
A scan for positively selected genes in the genomes of humans and chimpanzees.

PLoS Biol. 2005;3: e170. doi:10.1371/journal.pbio.0030170

30. Ferrer-Admetlla A, Liang M, Korneliussen T, Nielsen R. On detecting incomplete
soft or hard selective sweeps using haplotype structure. Mol Biol Evol. 2014;31:
1275–1291. doi:10.1093/molbev/msu077

31. Garud NR, Messer PW, Buzbas EO, Petrov DA. Recent selective sweeps in North
American Drosophila melanogaster show signatures of soft sweeps. PLoS Genet.
2015;11: e1005004. doi:10.1371/journal.pgen.1005004

32. Schrider DR. Background selection does not mimic the patterns of genetic
diversity produced by selective sweeps. Genetics. 2020;216: 499–519.
doi:10.1534/genetics.120.303469

33. Maclean CA, Chue Hong NP, Prendergast JGD. hapbin: An efficient program for
performing haplotype-based scans for positive selection in large genomic datasets.
Mol Biol Evol. 2015;32: 3027–3029. doi:10.1093/molbev/msv172

34. The 1000 Genomes Project Consortium. A global reference for human genetic
variation. Nature. 2015;526: 68–74. doi:10.1038/nature15393

35. Henn BM, Cavalli-Sforza LL, Feldman MW. The great human expansion. Proc
Natl Acad Sci U S A. 2012;109: 17758–17764. doi:10.1073/pnas.1212380109

36. Lohmueller KE, Albrechtsen A, Li Y, Kim SY, Korneliussen T, Vinckenbosch N,
et al. Natural selection affects multiple aspects of genetic variation at putatively
neutral sites across the human genome. PLoS Genet. 2011;7: e1002326.
doi:10.1371/journal.pgen.1002326

37. Siepel A, Bejerano G, Pedersen JS, Hinrichs AS, Hou M, Rosenbloom K, et al.
Evolutionarily conserved elements in vertebrate, insect, worm, and yeast genomes.
Genome Res. 2005;15: 1034–1050. doi:10.1101/gr.3715005

38. Luisi P, Alvarez-Ponce D, Pybus M, Fares MA, Bertranpetit J, Laayouni H. Recent
positive selection has acted on genes encoding proteins with more interactions
within the whole human interactome. Genome Biol Evol. 2015;7: 1141–1154.

doi:10.1093/gbe/evv055

39.   The ENCODE Project Consortium. An integrated encyclopedia of DNA elements

in the human genome. Nature. 2012;489: 57–74. doi:10.1038/nature11247

40.   GTEx Consortium. The Genotype-Tissue Expression (GTEx) pilot analysis:

Multitissue gene regulation in humans. Ardlie KG, Deluca DS, Segrè A V,

Sullivan TJ, Young TR, Gelfand ET, et al., editors. Science (80-). 2015;348: 648–

660. doi:10.1126/science.1262110

41.   Yates AD, Achuthan P, Akanni W, Allen J, Allen J, Alvarez-Jarreta J, et al.

Ensembl 2020. Nucleic Acids Res. 2020;48: D682–D688. doi:10.1093/nar/gkz966

42.   Harris DN, Song W, Shetty AC, Levano KS, Cáceres O, Padilla C, et al.

Evolutionary genomic dynamics of Peruvians before, during, and after the Inca

Empire. Proc Natl Acad Sci U S A. 2018;115: E6526–E6535.

doi:10.1073/pnas.1720798115

43.   Duret L, Arndt PF. The impact of recombination on nucleotide substitutions in the

human genome. PLoS Genet. 2008;4: e1000071.

doi:10.1371/journal.pgen.1000071

44.   Lander ES, Linton LM, Birren B, Nusbaum C, Zody MC, Baldwin J, et al. Initial

sequencing and analysis of the human genome. Nature. 2001;409: 860–921.

doi:10.1038/35057062

45.   Di Filippo M, Bernardi G. Mapping DNase-I hypersensitive sites on human

isochores. Gene. 2008;419: 62–65. doi:10.1016/j.gene.2008.02.012

46.   Huang Y-F. Dissecting genomic determinants of positive selection with an

evolution-guided regression model. Mol Biol Evol. 2021; msab291.

doi:10.1093/molbev/msab291

47.   Spruce C, Dlamini S, Ananda G, Bronkema N, Tian H, Paigen K, et al. HELLS

and PRDM9 form a pioneer complex to open chromatin at meiotic recombination

hot spots. Genes Dev. 2020;34: 398–412. doi:10.1101/gad.333542.119

48.   Haller BC, Messer PW. SLiM 3: Forward Genetic Simulations Beyond the
Wright-Fisher Model. Mol Biol Evol. 2019;36: 632–637.
doi:10.1093/molbev/msy228

49.   Halehalli R, Nagarajaram HA. Molecular principles of human virus protein-protein
interactions. Bioinformatics. 2015;31: 1025–1033.
doi:10.1093/bioinformatics/btu763

50.   Galtier N, Piganeau G, Mouchiroud D, Duret L. GC-Content Evolution in
Mammalian Genomes: The Biased Gene Conversion Hypothesis. Genetics.
2001;159: 907–911. doi:10.1093/genetics/159.2.907

51.   Dutta R, Saha-Mandal A, Cheng X, Qiu S, Serpen J, Fedorova L, et al. 1000
human genomes carry widespread signatures of GC biased gene conversion. BMC
Genomics. 2018;19: 256. doi:10.1186/s12864-018-4593-1

52.   Di C, Murga Moreno J, Salazar-Tortosa DF, Lauterbur ME, Enard D. Decreased
adaptation at human disease genes as a possible consequence of interference
between advantageous and deleterious variants. Elife. 2021;10: e69026.
doi:10.7554/eLife.69026

53.   Halldorsson B V., Palsson G, Stefansson OA, Jonsson H, Hardarson MT,
Eggertsson HP, et al. Characterizing mutagenic effects of recombination through a
sequence-level genetic map. Science (80-). 2019;363.
doi:10.1126/science.aau1043

1155

Supporting information captions

Supplemental Results S1. Results obtained using classic linear models and partial
correlations.

Supplemental Results S2. Associations between genomic factors and iHS across
populations and window sizes.

Supplemental Results S3. Associations between genomic factors and iHS after removing

specific predictors.

Supplemental Results S4. Fine scale analyses of recombination rate, regulatory density

and positive selection.

Supplemental Results S5. Results of population simulations.