Inhibition of autophagy as a novel therapy for the treatment of neurofibromatosis type

1 tumors

Stevens, M.[1,2], Wang, Y.[1,3], Bouley, S.J.[4], Mandigo, T.R.[4], Sharma, A.[4], Sengupta, S.[1,2],

Housden, A.[1], Oltean S.[2], Perrimon, N.[5,6], Walker, J.A.[4,7]*, Housden, B.E.[1,2,*]

[1] Living Systems Institute, University of Exeter, Exeter, UK

[2] Department of Clinical and Biomedical Science, University of Exeter, Exeter, UK

[3] The First People's Hospital of Qinzhou, China

[4] Center for Genomic Medicine, Massachusetts General Hospital, Boston, MA 02114, USA

[5] Department of Genetics, Blavatnik Institute, Harvard Medical School, Boston, MA, 02115,

USA

[6] Howard Hughes Medical Institute

[7] Cancer Program, Broad Institute of MIT and Harvard, Cambridge, MA 02142, USA

* Correspondence to b.housden@exeter.ac.uk or jwalker1@mgh.harvard.edu

Key Words

Autophagy; Drosophila; Drug repurposing; Neurofibromatosis type 1; Synthetic lethality

ABSTRACT

Neurofibromatosis type 1 (NF1) is a genetic disorder associated with various symptoms including the formation of benign tumors along nerves. Drug treatments are currently limited. The MEK inhibitor selumetinib is used for a subset of cases but is not always effective and can cause side effects. Therefore, there is a clear need to discover new drugs to target *NF1*-deficient tumor cells. Using a *Drosophila* cell model of NF1, we performed synthetic lethal screens to identify novel drug targets. We identified 54 candidates, which were validated with variable dose analysis as a secondary screen. Five candidates could be targeted using existing drugs, with autophagy inhibitors (chloroquine (CQ) and bafilomycin A1) showing the greatest potential for selectively killing *NF1*-deficient *Drosophila* cells. When further investigating autophagy-related genes, we found that 14 out of 30 genes tested had a synthetic lethal interaction with *NF1*. These 14 genes are involved in the regulation of all aspects of the autophagy pathway and can be targeted with additional autophagy drugs, although none were as effective as CQ. The lethal effect of autophagy inhibitors was conserved in a panel of human *NF1*-deficient Schwann cell lines, highlighting their translational potential. The effect of CQ was also conserved in a *Drosophila NF1 in vivo* model and in a xenografted *NF1*-deficient tumor cell line in mice, with CQ treatment resulting in a more significant reduction in tumor growth than selumetinib treatment. Furthermore, combined treatment with CQ and selumetinib resulted in a further reduction in *NF1*-deficient cell viability. In conclusion, *NF1*-deficient cells are vulnerable to disruption of the autophagy pathway. This pathway represents a promising therapeutic target for *NF1*-associated tumors, and CQ was identified as a promising candidate drug for the treatment of *NF1* tumors.

INTRODUCTION

Neurofibromatosis type 1 (NF1) is a genetic disorder with autosomal-dominant inheritance affecting 1 in ~2,700 births (1, 2). Although the penetrance of NF1 is virtually complete after childhood, the disease is characterized by highly variable clinical expressivity. Symptoms include near universal benign, but often disfiguring, peripheral nerve associated tumors known as neurofibromas, as well as malignant tumors, including usually fatal malignant peripheral nerve sheath tumors (MPNSTs) (3). In part reflecting higher rates of vascular defects and cancer, the life expectancy of NF1 patients is reduced by approximately 15 years (4).

NF1 is caused by loss of neurofibromin, a 320 kDa protein whose only widely accepted function is to serve as a RAS GTPase Activating Protein (RASGAP) for H-, K-, N-RAS and R-RAS1, 2, and 3 (5-8). RASGAPs promote the conversion of active RAS-GTP into inactive RAS-GDP by stimulating the low intrinsic rate of RAS-GTP hydrolysis. Consequently, loss of neurofibromin can result in dysregulation of signaling downstream of RAS, the best documented being the RAF/MEK/ERK and PI3K/AKT/mTOR pathways (9). Although dysregulated RAS signaling is believed to be the proximal cause of NF1 symptoms, it is unclear which of the numerous effectors downstream of RAS are relevant for disease progression, as well as the identities of the disease-pertinent targets of the signaling pathways mediating their effects. The situation is further complicated since there is undoubtedly crosstalk between these different pathways. In patients with *NF1*-driven malignant tumors, targeting RAS pathway components such as MEK or ERK is a reasonable therapeutic option, although RAS is subject to highly robust regulation (9), which may explain why, despite considerable effort, effective therapies for RAS-driven cancers have been very challenging to develop. However, chronically blocking RAS may never be an appropriate strategy for treating the many serious but non-life-threatening symptoms of NF1, especially in children.

Currently, there are limited therapies for any *NF1*-associated tumors. The only available drug

is the MEK inhibitor selumetinib, which was approved for use in a subset of pediatric plexiform

neurofibromas in April 2020. However, not all tumors were responsive to treatment and serious

side effects can be associated with MEK inhibition (10-13). Therefore, there is a clear clinical

need to discover new drugs that specifically target *NF1*-deficient tumor cells either alone or in

combination with selumetinib*.*

One approach to identify candidate drug targets for tumorigenic diseases is the use of

synthetic lethal interaction screens. Synthetic lethal interactions are a type of genetic

interaction in which inhibition of either of two genes alone is viable, but the combined inhibition

of both genes is inviable. When one of these genes is mutated in tumor cells, such interactions

can be exploited to kill those cells exclusively by targeting the synthetic lethal partner gene

using a drug (14, 15). This approach is attractive because treatment is expected to be lethal

to tumor cells but have no effect on wild-type, healthy cells.

Despite long-term interest in the use of synthetic lethality as a therapeutic strategy to treat

tumors, few drugs have successfully progressed to clinical use. A major factor preventing

successful development of treatments against synthetic lethal interactions is a lack of

consistency between interactions identified in different genetic backgrounds (16). Therefore,

candidates identified from a single model system often fail to translate to other model systems

and subsequently to clinical use. To overcome this limitation, our approach makes use of

diverse model systems by first screening for synthetic lethal interactions with genes mutated

in tumors using *Drosophila* cells. The conservation of candidate interactions can then be

assessed in a range of other model systems, including human cells, providing a filter to remove

interactions that are specific to a single model system. This approach has previously proved

successful, leading to the discovery of mizoribine and palbociclib as promising candidates for

the treatment of tuberous sclerosis complex (TSC) and Von Hippel-Lindau (VHL)-linked

cancers, respectively (17-19). In both cases, hits from *Drosophila* synthetic lethal screens

were validated with a high success rate in both human cells and mouse models.

Given the previous success of using the *Drosophila* approach, we have applied this method

to identify candidate drug targets to treat *NF1*-deficient tumors. Here, we describe the

generation of a *dNF1* null mutant *Drosophila* cell line using CRISPR gene editing and its use

in 1) conducting synthetic lethal screens to identify perturbed pathways that confer

vulnerability of NF1-deficient cells, and 2) to identify candidate drug targets that might be used

for therapeutic benefit to specifically kill NF1-associated tumors. We find dNF1-deficient cells

are vulnerable to inhibition of autophagy. Importantly, we show that this selective effect can

be reproduced with multiple inhibitors and in several human tumor-derived cell lines, as well

as in a *Drosophila in vivo* NF1 model and xenografts of *NF1*-deficient tumor cell lines in mice,

indicating that these repurposed drugs may have promise for the treatment of *NF1* tumors.

Finally, we show that combined treatment with CQ or bafilomycin A1 and selumetinib results

in increased selective killing of *NF1*-deficient cells, indicating the potential for combinatorial

therapy.

RESULTS

Generation of *Drosophila* and human Schwann cell *NF1* models using CRISPR/Cas9

gene editing

Our previous studies have demonstrated the potential of using cross-species genetic screens

to identify candidate therapeutic targets for human disease (17-20). The *NF1* gene is well

conserved between *Drosophila* and humans with 68% identity at the amino acid level (**Figure**

S1). To use the same approach to find new targets for the treatment of *NF1*-associated

tumors, we first used CRISPR gene editing to generate indel mutations in *dNF1* in *Drosophila*

S2R+ cells. Sequencing was used to confirm that the induced frame-shift mutations resulted

in null *dNF1* alleles (**Figure 1A**). Of note, S2R+ cells are aneuploid (21), and our sequencing

results suggest that they have three copies of the *dNF1* gene.

The resulting dNF1-KO S2R+ cell line (hereafter called dNF1-KO) was characterized by

assessing the expression of neurofibromin using western blots. We found no detectable signal

in the dNF1-KO line compared to parental wild-type (WT) S2R+ cells (**Figure 1B, Figure S2A**).

However, we found no significant increase in pERK levels in dNF1-KO cells under normal

culturing conditions, suggesting that other growth conditions (i.e., reduced serum) may be

necessary to exacerbate differences in RAS signaling, as is necessary in other NF1 cell line

models (**Figure 1B-C, Figure S2B**). Given that neurofibromin is a negative regulator of RAS,

we assessed the growth and proliferative phenotypes of dNF1-KO cells compared to S2R+

cells. Consistent with deregulation of a mitogenic pathway, dNF1-KO cells showed an

increased rate of growth as measured using CellTiter-Glo assays to assess total ATP levels

in the population. This effect was observed in both the presence and absence of serum in the

culture media (**Figure 1D**), indicating that culture growth is both accelerated in the absence of

dNF1 and is decoupled from upstream growth factor signaling pathways. To determine

whether this increase in culture growth was due to increased proliferation, increased cell

growth, or both, we performed cell counts following culture in full serum and CellTiter-Glo

assays on normalized numbers of cells from each genotype (baseline readings). Cell counts

for dNF1-KO showed an increase in cell numbers following culture compared to S2R+ cells,

and the 'baseline' CellTiter-Glo showed no difference (**Figure 1D**). This suggests that the

difference in culture growth is primarily due to increased cell proliferation rather than an

increase in the cell size or ATP content of the cells. Together, these results indicate that the

dNF1-KO line represents a novel *dNF1* null mutant cell model, with properties consistent with

known effects of NF1 loss.

We also utilized CRISPR/Cas9 gene editing to generate indel mutations within exon 3 of *NF1*

in wild-type immortalized human Schwann cells (ipn02.3 2λ). Sequencing of single cell clones

was used to confirm out-of-frame deletions in one or both *NF1* alleles, resulting in *NF1-*

deficient cell lines (**Figure 1E**). The resulting heterozygous (C8) and homozygous (C23) cell

lines were characterized by assessing neurofibromin expression using western blots (**Figure**

1F, Figure S2C). Consistent with what we observed in dNF1-KO cells, there was no significant

increase in pERK expression under normal culturing conditions (**Figure 1F-G, Figure S2D)**.

These results suggest that these otherwise isogenic human cell lines are an appropriate model

to validate any results obtained in our dNF1-KO cell lines.

A

B

C

E

F

G

Figure 1. Generation and characterization of NF1-deficient Drosophila and human Schwann cells using CRISPR. (A) Drosophila S2R+ cells were transfected with Cas9 and sgRNA designed to target a double-stranded break in exon 2 of dNF1. Molecular analysis of the dNF1-KO line revealed deletions of 20, 11 and 13bp in three alleles (S2R+ cells are aneuploid). The position of the guide RNA is shown in blue and the PAM site in red. Predicted effect of deletions on the amino acids in each allele are shown, each resulting in premature termination. (B) Representative image of western blot showing loss of neurofibromin in dNF1-KO cells compared to WT S2R+ cells (two replicates performed) and pERK and total ERK levels in S2R+ and dNF1-KO cells under normal culturing conditions (10% serum). (C) Quantification of pERK/ERK ratio in S2R+ and dNF1-KO cells (in 10% serum) shows no significant change (from triplicate experiments). (D) Characterization of dNF1-KO cell population growth and proliferation rate as assessed using CellTiter-Glo assays (n=3-8, error bars indicate standard deviation, p values determined using unpaired (except for cell counts, which were paired), two-tailed t-tests) or cell counts. (E) CRISPR/Cas9 was used to target exon 3 of NF1 in a telomerase-immortalized human Schwann cell line (hTERT ipn02.3 2λ) (22) to generate isogenic NF1 knock out (NF1-/+ and NF1-/-) cell lines. Line C8 has a heterozygous 1 bp deletion (c.282delT) and line C23 is a transheterozygous combination of 1 bp (c.282delt) and 2 bp deletions (c.282-283 delTG). Position of the guide RNA (NF1-sg2) is shown in blue. (F) Western blot analysis of cell lines showing reduction (C8) and absence (C23) of neurofibromin compared to the wild-type ipn02.3 2λ progenitor Schwann cell line. pERK and total ERK levels were assessed in cells grown in 10% serum. (G) Quantification of pERK/ERK ratios shows no significant change between wild-type and C8 or C23 (from triplicate experiments).

Mapping synthetic lethal interactions in dNF1-KO cells using a genome-wide RNAi screen

We used a genome-wide dsRNA library to screen in both S2R+ and dNF1-KO cells for synthetic lethal interactions (**Figure S3**). Correlation coefficients ranged between 0.9 and 0.99 (average 0.93) for control wells and between 0.55 and 0.66 (average 0.61) for non-control wells, illustrating a high rate of reproducibility between replicates. Next, we identified synthetic lethal interactions by filtering the results for dsRNA reagents that reduced the viability of dNF1-KO cells (median $Z<-1.5$) to a greater extent than wild-type cells (median $Z\geqslant-1.5$). This analysis identified 134 candidate genes (**Table S1**).

Genetic screens are often associated with false-positive results due to off-target effects from dsRNA reagents or noise in the screen assay. To remove potential false positives, we overlaid the screen hits onto a protein-protein interaction network from the String database (23). Synthetic lethal interactions are generally similar between genes that have related functions; therefore, proteins that physically interact are expected to share synthetic lethal interactions. Using the combination of physical and genetic interaction data, we could remove false positives from the screen results by isolating only hits that have physical interactions with at least one other hit from the genetic interaction screen. In addition, we filtered the candidates to isolate only those with clear orthologs in humans. Following this process, 54 high-confidence candidate targets remained, corresponding to 74 human genes (**Figure 2A, Table S2**).

Validation of candidate synthetic lethal interactions using Variable Dose Analysis (VDA)

We used the VDA assay as an additional combinatorial screen to assess synthetic lethality between *dNF1* and all 54 candidate drug targets. Two shRNAs targeting each of the genes

were generated. These reagents were tested in S2R+ and dNF1-KO cells. Of the 54 genes,

46 showed a >10% reduction in viability in dNF1-KO cells compared to S2R+ controls (**Figure**

2B; ranked in order of effect on dNF1-KO viability). These results indicate that the network is

a reliable representation of the synthetic lethal interaction profile of the *dNF1* gene.

To identify potential drugs for repurposing to treat NF1 tumors, we filtered the candidate gene

list to those that could be targeted using existing drugs, resulting in eight candidate drug

targets (**Figure 2C**). We then removed candidates that had previously been studied in relation

to NF1, leaving five candidate drug targets. We also included MEK (selumetinib) as a control.

Note that three drugs that target autophagy were included, and some drugs inhibited multiple

targets. In total we tested seven candidate drugs inhibiting six different targets (**Table 1**).

Figure 2. A network of synthetic lethal interaction for Drosophila dNF1 and VDA

analysis of candidate drug targets. *Using wild-type S2R+ and dNF1-KO cells, we used a*

near genome-wide dsRNA library to screen approximately 10000 genes for difference in

viability when expression is knocked down as assessed by CellTiter-Glo assays, resulting in

134 genes identified as having a synthetic lethal interaction with NF1. ***(A)*** *A synthetic lethal*

interaction network for dNF1 in Drosophila S2R+ cells generated using Cytoscape (24). Solid

lines represent physical interactions within functional groups and dashed lines represent

physical interactions between functional groups. ***(B)*** *VDA assays were performed for all 54*

candidate genes, with two shRNAs per gene, in WT S2R+ (blue) and dNF1-KO (red) cells.

The best shRNA from the 46 genes that reduced dNF1-KO viability by >10% relative to S2R+

controls ranked in order of effect are shown. ***(C)*** *shRNA knockdowns of candidate genes that*

resulted in a >10% reduction in dNF1-KO viability compared to S2R+ controls and were

druggable targets (n=6-9, error bars indicate standard deviation). All eight shRNAs showed a

significant reduction in dNF1-KO viability relative to S2R+ controls assessed using two-tailed,

unpaired t-tests. Results in panel C are reproduced from panel B for clarity.

Table 1. Candidate genes that selectively decreased dNF1-KO viability that can be targeted with drugs, either directly or through pathway inhibition.

Drug	Target (direct or indirect): DM (Human)
Chloroquine*	Autophagy; Atg8b (GABARAP), RpS6 (RPS6)
EAD1	Autophagy; Atg8b (GABARAP), RpS6 (RPS6)
Selumetinib*	Dsor1 (MEK, MAPKK)
Metformin*	ND-MNLL (NDUFB1)
PYR41	Uba1 (UBA1)
LB100	Pp2A-29B (PPP2R1A and PPP2R1B)
Bafilomycin A1	Autophagy; Atg8b (GABARAP), RpS6 (RPS6)

*FDA-approved

Existing inhibitors selectively affect *dNF1*-KO *Drosophila* cells

Repurposing existing drugs represents the most efficient route to develop new therapeutics. Each of the seven drugs that target the *dNF1* synthetic lethal partner genes were first tested in S2R+ and dNF1-KO cells using CellTiter-Glo assays to measure viability after 48 h of treatment (**Figure S4**). Of the seven drugs tested in *Drosophila* cells, we selected the two autophagy inhibitors (CQ and bafilomycin A1) for further study, as both showed a significant and consistent effect on reducing dNF1-KO viability.

One of the strongest hits from the genetic screen was *Atg8b*, which encodes a key component

of the autophagy pathway. Autophagy is commonly inhibited experimentally using CQ, which

is clinically used as an anti-malarial and shows anti-viral properties, and bafilomycin A1. CQ

functions to inhibit autophagy by blocking the binding of autophagosomes to lysosomes by

diffusing into the lysosomes and altering the acidic environment, thereby inhibiting autophagic

lysosomal degradation (25). On the other hand, bafilomycin A1 disrupts autophagic flux by

independently inhibiting V-ATPase-dependent acidification and Ca-P60A/SERCA-dependent

autophagosome-lysosome fusion (26). We initially focused on CQ because it is generally well-

tolerated (27) and can inhibit autophagy *in vivo* at clinically achievable concentrations (28).

Bafilomycin A1 is a potent inhibitor of autophagy but is not clinically approved (29).

Nevertheless, we tested bafilomycin A1 to provide additional validation of the effects on

autophagy inhibition brought about through an independent mechanism.

First, we quantified autophagic flux in dNF1-KO an dS2R+ cells without drug treatment, i.e.,

how many autophagosomes form and then become degraded, by measuring the difference in

the number of autophagic vesicles in the presence versus the absence of a lysosomal inhibitor,

CQ. We observed significantly higher levels of autophagic flux in dNF1-KO cells under serum-

free conditions compared to S2R+ controls (**Figure 3A**)**.** We then tested whether CQ and

bafilomycin A1 would phenocopy the selective effect observed using genetic inhibition of

Atg8b in *Drosophila* cells. Both S2R+ and dNF1-KO cells were treated with varying doses of

CQ or bafilomycin A1 and cell viability was measured using CellTiter-Glo assays, PI staining,

and annexin V staining. A significantly greater effect on dNF1-KO cell viability was observed

at multiple concentrations of each drug in serum-free media after 48 h of treatment across all

three assays, further validating the interaction between autophagy and NF1 and

demonstrating that the effect is cytotoxic (**Figure 3B-F**).

A

B

C

D

E

F

Figure 3. CQ and bafilomycin A1 selectively affect dNF1-deficient Drosophila cells. (A)
dNF1-KO cells show a significant increase in autophagic flux compared to WT S2R+ control
cells in serum-free media as assessed by inhibiting autophagosome flux for 4 h using the
lysosomal inhibitor CQ (10 μM), and then measuring the initial rate of accumulation of the
fluorescent substrate labelling the autophagosomes using a fluorescent plate reader
(normalized to DAPI, and then to S2R+ controls) (n=6, p value obtained using the two-tailed,
unpaired, Student's t-test). (B) CQ reduced dNF1-KO cell viability compared to S2R+ cells
after 48 h in serum-free media as measured using CellTiter Glo assays (n=4; p values obtained
using a two-way ANOVA with Tukey for multiple comparisons). (C) Analysis of cell death using
PI staining with CQ in S2R+ and dNF1-KO cells grown in serum-free media (n=4; p values
obtained using a two-way ANOVA with Tukey for multiple comparisons). (D) Bafilomycin A1
reduced dNF1-KO cell viability compared to S2R+ cells after 48 h in serum-free media using
the CellTiter Glo assay (n=4; P values obtained using a two-way ANOVA with Tukey for
multiple comparisons). (E) Analysis of cell death using PI staining with bafilomycin A1 in S2R+
and dNF1-KO cells grown in serum free media (n=6; p values obtained using a two-way
ANOVA with Tukey for multiple comparisons). (F) CQ and bafilomycin A1 (B-A1) increased
annexin V staining in dNF1-KO cells relative to S2R+ controls after 48 h in serum-free media
(n=4-6; p values obtained using a two-way ANOVA with Tukey for multiple comparisons). In
all cases, bars represent the mean and error bars indicate standard deviation.

Both early- and late-stage autophagy genes have a synthetic lethal interaction with
***dNF1*, which can be targeted with other autophagy inhibitors**

Autophagy is a complex process involving multiple stages and many different proteins (**Figure
4A**). To determine whether more specific targeting of autophagy components would result in
a greater selective effect in *dNF1*-KO cells, we performed an additional VDA screen to assess
for synthetic lethal interactions between *dNF1* and 29 key autophagy genes (in addition to
Atg8b) using S2R+ and dNF1-KO cells (all 29 genes screened are shown in **Table S3**). In

total, 14 genes, in addition to the previously identified *Atg8b*, were found to significantly reduce

dNF1-KO viability by >10% relative to controls (**Figure 4B**). Interestingly, these 14 genes are

implicated across all stages of the autophagy pathway (**Figure 4A,** genes shown in red).

Therefore, we tested drugs targeting specific aspects of the autophagy pathway in dNF1-KO

cells, including MRT68921 (an early-stage autophagy inhibitor of ULK1 and ULK2), VPS34-

IN1 (a potent early-stage autophagy inhibitor of the PI3K-III complex), NSC 185058 (a mid-

stage inhibitor of *Atg4b*), CA5f (a late-stage inhibitor of autophagic flux), and obatoclax (a late-

stage autophagy inhibitor) (**Figure 4A**). In general, the autophagy inhibitors showed a

selective viability effect in dNF1-KO cells, except for MRT68921 and NSC 185058 (**Figure 4C-**

G); however, none of the drugs were deemed to be more effective in selectively killing dNF1-

KO cells than CQ and bafilomycin A1. CQ also has the advantage of being FDA-approved.

Therefore, although targeting autophagy at various stages of the pathway appears to have

selective viability effects in dNF1-KO cells, the late-stage autophagy inhibitor CQ could provide

the greatest potential for use in the clinic to treat NF1.

Figure 4. Both early- and late-stage autophagy genes have a synthetic lethal interaction with dNF1, which can be targeted with other autophagy-inhibitors. (A) Diagrammatic representation of the autophagy pathway in Drosophila. The genes found to have a synthetic lethal interaction with dNF1 are shown in red. Drugs used to target each stage of the pathway in subsequent viability assays are shown in blue. (B) VDA assays performed for 30 autophagy-related genes, with two shRNAs per gene, in S2R+ (blue) and dNF1-KO (red) cells. Shown is the most effective shRNA from each of the 14 genes that reduced dNF1-KO viability by >10% relative to S2R+ controls, ranked in order of effect. (C-G) Testing autophagy inhibitors for differences in cell viability between dNF1-KO and S2R+ cells after 48 h in serum-free media as measured using the CellTiter Glo assays (n=4; p values obtained using a two-way ANOVA with Tukey for multiple comparisons). VPS34-IN1 (D), CA5f (F), and obatoclax (G) selectively reduced dNF1-KO cell viability compared to S2R+ cells in a dose-dependent manner. By contrast, MRT68921 (C) and NSC 185058 (E) did not selectively affect dNF1-KO cell viability compared to S2R+ cells. In all cases, data represents the mean and error bars indicate standard deviation.

CQ and bafilomycin A1 selectively affect *NF1*-deficient human cells

To determine whether the selective effect of CQ and bafilomycin A1 was conserved in human cells, we tested the effects of drug treatment on a panel of human *NF1*-deficient cell lines. These included a pair of otherwise isogenic *NF1$^{+/-}$* (C8) and *NF1$^{-/-}$* (C23) immortalized Schwann cells generated using CRISPR/Cas9 gene editing (**Figure 1F-H**). In addition, we used two pairs of immortalized Schwann cell lines (pair 1: ipnNF95.11C (*NF1$^{+/-}$*) and ipNF95.11b 'C' (*NF1$^{-/-}$*) and pair 2: ipnNF09.4 (*NF1$^{+/-}$*) and ipNF05.5 (*NF1$^{-/-}$*)) derived from plexiform neurofibromas (22). Autophagic flux was significantly higher in *NF1$^{-/-}$* cells (ipNF95.11b 'C' and ipNF05.5) relative to *NF1$^{+/-}$* controls (ipnNF95.11C and ipnNF09.4), with a similar but not significant effect also observed in the CRISPR-generated *NF1$^{-/-}$* (C23) cells compared to *NF1$^{+/-}$* (C8) controls (**Figure 5A-C**). Similarly, lysosomal activation was increased

in ipNF95.11b 'C' cells compared to ipnNF95.11C controls, further indicating an increase in

baseline autophagy levels (**Figure S5**). Additionally, lysosomal activation was inhibited by CQ

and bafilomycin A1 in both $NF1^{+/-}$ and $NF1^{-/-}$ cells (**Figure S5**).

***Figure 5. CQ and bafilomycin A1 selectively affect NF1-deficient human cells.** (A-C)*

Autophagic flux was increased in C23, ipNF95.11b 'C', and ipNF05.5 $NF1^{-/-}$ cells compared to

heterozygous controls after 4 h of lysosomal inhibition with CQ (10 μM) in serum-free media

(n=5–9; p values obtained using the two-tailed, unpaired, Student's t-test). CQ significantly

*reduced $NF1^{-/-}$ cell viability relative to $NF1^{+/-}$ controls at varying doses in C8/C23 cells **(D)**,*

*ipnNF95.11C/ipNF95.11b 'C' cells **(E)**, and ipnNF09.4/ipNF05.5 cells **(F)** after 48 h in serum-*

free media as measured with CellTiter Glo assays (n=3-4; p values obtained using a two-way

*ANOVA with Tukey for multiple comparisons). **(G-I)** Bafilomycin A1 significantly reduced $NF1^{-/-}$*

*cell viability relative to $NF1^{+/-}$ controls at varying doses in C8/C23 cells **(G)**,*

*ipnNF95.11C/ipNF95.11b 'C' cells **(H)**, and ipnNF09.4/ipNF05.5 cells **(I)** after 48 h in serum-*

free media as measured with CellTiter Glo assays (n=3-4; p values obtained using a two-way

*ANOVA with Tukey for multiple comparisons). **(J-L)** We observed an increase in caspase*

activation in $NF1^{-/-}$ cells relative to $NF1^{+/-}$ control cells when treated with CQ or bafilomycin A1

*for 48 h in serum-free media: **(J)**, C8 and C23 cells (50 and 5 nM, respectively), **(K)***

*ipnNF95.11C/ipNF95.11b 'C' cells (1 and 5 nM, respectively), and **(L)** ipnNF09.4/ipNF05.5*

cells (50 and 5 nM, respectively) (n=4; p values obtained using a two-way ANOVA with Tukey

for multiple comparisons). In all cases, bars represent the mean and error bars indicate

standard deviation.

Both CQ and bafilomycin A1 resulted in a significantly greater reduction in the viability of

homozygous $NF1^{-/-}$ deficient cells compared to heterozygous $NF1^{+/-}$ controls after 48 h of

treatment under serum-free media conditions, as measured with the CellTiter-Glo and

caspase assays (**Figure 5D-L**), demonstrating that the selective effects are conserved

between *Drosophila* and human systems. Although the effective dose of bafilomycin A1 in

Drosophila cells appears to be very low, human cells were affected by doses in the range of

the previously demonstrated IC_{50} of 0.44 nM in human cells (30).

We also tested the additional five autophagy inhibitors in the two patient *NF1*-deficient cell

lines (**Figure S6**); however, no drug showed a consistent effect across the panel of human

cell lines that was comparable to that of CQ and bafilomycin A1, further highlighting the

reproducibility of our *Drosophila* model system.

Together, these results demonstrate that *NF1*-deficient cells have a vulnerability to disruption

of the autophagy pathway, which is conserved and reproducible with multiple inhibitors

between *Drosophila* and human Schwann cells derived from NF1-associated tumors. Not only

does autophagy represent a promising pathway for targeting NF1-associated tumors, but we

identified CQ as a candidate drug for the potential treatment of NF1 tumors.

CQ affects survival in a *Drosophila in vivo NF1* mutant model

As CQ is a well-tolerated FDA-approved therapeutic, and showed a significant effect on dNF1-

KO and human NF1$^{-/-}$ cell viability, we chose to take this drug forward to determine synthetic

lethality *in vivo*. *Drosophila dNf1* mutant flies show defective Ras signaling, which result in a

number of neurobehavioral phenotypes (31-35). For this study, we generated a novel *dNf1*

null mutant fly using CRISPR gene editing: *dNf1^{C1}* (delAT162-163) (**Figure 6A**). Western blots

using lysates prepared from adult heads from *dNf1^{C1}* homozygous mutants showed no

detectable expression of neurofibromin (**Figure 6B**). In addition, ELISAs showed a 4-fold

increase in pERK/ERK of *dNf1^{C1}* mutants compared to the WT parental line (**Figure 6C**).

To determine whether CQ affects survival in *NF1*-deficient *Drosophila*, we took two

approaches. Firstly, we compared the effect of CQ on *dNf1^{C1}* homozygous null mutant flies,

the WT parental line, and *dNf1^{C1}* with re-expression of dNf1 from a *UAS-dNF1* transgene

driven with a pan-neuronal (*nSyb-Gal4*) driver (*dNf1^{C1} + nSyb-Gal4>UAS-dNf1)*. CQ resulted

in increased lethality of *dNf1^{C1}* mutants compared to the WT control in a dose-dependent

manner (**Figure 6D, Figure S7**). Furthermore, we were able to rescue the CQ sensitivity of

dNf1^{C1} mutant flies by re-expression of dNf1 from a *UAS-dNf1* transgene (**Figure 6D**).

Secondly, we tested flies with pan-neuronal RNAi knock down of dNf1 (using *nSyb-Gal4*)

compared to a landing site control on food containing CQ (35 mM). Flies with *dNf1* RNAi

knockdown, showed significantly reduced survival time on CQ compared to CQ-treated

landing site control flies, and untreated flies, phenocopying the effects seen in *dNf1^{C1}* mutant

flies (**Figure 6E**). Together, these results demonstrate that *dNf1*-deficient flies have

vulnerability to disruption of the autophagy pathway, as shown to be conserved and

reproducible in *Drosophila* dNF1-KO cells and human Schwann cells derived from NF1-

associated tumors. This further highlights the autophagy pathway as a target for the potential

treatment of NF1-associated tumors, with CQ as a candidate drug.

CQ reduced *NF1*-deficient MPNST tumor xenograft growth *in vivo*

In order to test the effects of CQ *in vivo* in a mammalian model, we used the ST88-14 *NF1$^{-/-}$*

MPNST xenograft mouse model. Although some NF1 PN tumor cells have been shown to

form xenograft tumors, they are very slow growing and require interactions with the tumor

microenvironment (36). Therefore, we first assessed whether CQ selectively killed ST88-14

cells *in vitro* using the CellTiter-Glo assay and found CQ to affect cell viability at concentrations

slightly higher than those showing an effect in the previously tested *NF1*-deficient Schwann

cell lines (**Figure 6F** compared to **Figure 5D-F**). In addition, we found that the ST88-14 cells

were more sensitive to CQ than the C8 heterozygous NF1 Schwann cell line (**Figure 6F**).

In mice implanted with ST88-14 *NF1$^{-/-}$* xenografts, treatment with CQ (50 mg/kg in saline,

intraperitoneally, three times per week) or selumetinib (25 mg/kg in saline, oral gavage, three

times per week) once the tumors had started to grow, resulted in a significant reduction in

tumor xenograft growth over a period of 3 weeks in comparison to vehicle-treated controls

(**Figure 6G**). Furthermore, there was a significant reduction in tumor xenograft growth in CQ-

treated mice in comparison to selumetinib-treated mice, indicating CQ to have a superior effect

on *NF1*-deficient tumor xenograft growth. CQ and selumetinib treatment were found to have

no toxicity effects in these mice (**Figure S8**), which was also assessed in a prior study on

C57BL/6 mice (data not shown).

CQ and selumetinib were found to induce *NF1*-deficient cell apoptosis *in vitro*; therefore, we

used western blotting to assess the cleavage of a key apoptosis protein, PARP, in xenografts

from control, CQ, and selumetinib-treated mice. We found a significant increase in the 24 kDa

fragment of cleaved PARP/PARP in xenografts from CQ-treated mice (**Figure 6H-I**), indicating

that CQ was causing cell apoptosis in the NF1 xenografts, thus slowing tumor growth. This

increase was also observed in selumetinib-treated mice, although the difference was not

significant.

Figure 6. CQ affected lethality in NF1 mutant Drosophila and reduced NF1-deficient

MPNST tumor xenograft growth in vivo. (A) *We generated a novel dNf1 null mutants using*

*CRISPR gene editing: dNf1^{C1} (dNF1 delAT162-163). **(B)** Western blots of anti-dNf1*

immunoprecipitates from lysates prepared from adult heads from CRISPR mutants showed

*no expression of Nf1 in dNf1^{C1} homozygous animals **(C)** ELISA for pERK/ERK showed a 4-*

fold increase in pERK/ERK of dNf1^{C1} mutants compared to WT flies. Error bars indicate the

standard deviation between duplicate samples. A paired, two-tailed t-test was performed to

*determine significance; *P < 0.05. **(D)** Addition of CQ to 35 µM to food resulted in increased*

lethality of dNf1^{C1} mutants compared to WT flies (n=3, 20 flies per replicate). The sensitivity of

dNf1 mutant flies to CQ was rescued by re-expression of dNf1 from a UAS-dNF1 transgene

*using the nSyb-Gal4 driver (n=3, 20 flies per replicate) **(E)** dNf1 RNAi knockdown flies (nSyb-*

Gal4>v109637) show a similarly reduced survival time when cultured on food with 35 µM CQ

*compared to control flies (nSyb-Gal4>VIE-260B v60100). n=3, 20 flies per replicate. **(F)** CQ*

significantly reduced NF1$^{-/-}$ mutant cell viability relative to C8 NF1$^{+/-}$ controls at varying doses

in ST88-14 cells after 48 h in serum free media, as measured with the CellTiter Glo assay

*(n=9; P values obtained using a two-way ANOVA with Tukey for multiple comparisons). **(G)** In*

mice implanted with ST88-14 NF1$^{-/-}$ xenografts, intraperitoneal injections of CQ (50 mg/kg, 3x

weekly) or oral gavage with selumetinib (25 mg/kg, 3x weekly) significantly slowed tumor

growth compared to vehicle-treated controls. Furthermore, there was a significant reduction

in tumor growth in CQ-treated mice compared to selumetinib-treated mice (n=6; P value

*obtained using a two-way ANOVA). **(H-I)** Following extraction of the xenografts, western*

blotting revealed the increased protein expression of the 24 kDa fragment of cleaved PARP

relative to PARP in CQ-treated mice in comparison to controls (n=4; P values obtained using

the unpaired Student's t-test).

Selumetinib enhances to viability effect of CQ and bafilomycin A1

Selumetinib, a MEK1/2 inhibitor, is currently the only FDA-approved drug for the treatment of tumors associated with neurofibromatosis type 1 (37). We treated S2R+ and dNF1-KO cells with selumetinib (10 μM) with and without CQ (1 mM) or bafilomycin A1 (100 pM) for 48 h in serum-free media and performed CellTiter-Glo assays to measure cell viability (**Figure 7A**). When selumetinib was combined with CQ or bafilomycin A1, we saw a further reduction in dNF1-KO, but not S2R+ viability relative to the CQ/bafilomycin A1 only, indicating that selumetinib enhances the effects of CQ/bafilomycin A1 on dNF1-KO cell viability.

Similarly, in C8/C23, ipnNF9511.C/ipNF95.11b 'C', and ipnNF09.4/ipNF05.5 cell lines, selumetinib significantly enhanced the reduced viability effect of CQ in $NF1^{-/-}$, but not $NF1^{+/-}$ cells (**Figure 7B-D**). Therefore, combined treatment of CQ and selumetinib had a greater impact in $NF1$-deficient cell viability compared to either drug alone. In addition, selumetinib significantly enhanced the viability effect of bafilomycin A1 in patient $NF1^{-/-}$ (ipNF95.11b 'C' and ipNF05.5) but not $NF1^{+/-}$ cells (ipnNF9511.C and ipnNF09.4), although not in C8/C23 cells (**Figure 7B-D**).

To determine whether these combined effects were additive or synergistic, the coefficient of drug interaction (CDI) was calculated for each cell line treated with CQ + selumetinib or bafilomycin A1 + selumetinib. A CDI<1 indicates synergy, CDI=1 indicates additivity, and CDI>1 indicates antagonism. We observed a relatively strong synergistic effect with CQ + selumetinib in C23 cells, and with bafilomycin A1 + selumetinib in ipNF05.5 cells. Weaker evidence of synergy was also seen in other cell lines (**Figure 7E**).

E

Drug synergy	CDI: CQ + Selumetinib	CDI: Bafilomycin A1 + Selumetinib
dNF1-KO	0.99	0.99
C23	0.72	0.94
ipNF95.11b 'C'	1.0	0.96
ipNF05.5	0.93	0.75

Figure 7. Combined effect of selumetinib with CQ or bafilomycin A1 on NF1-deficient

cell viability. (A) *S2R+ and dNF1-KO cells were treated with 0 μM or 10 μM selumetinib*

combined with CQ (1 mM) or bafilomycin A1 (10 pM). Alone, selumetinib significantly reduced

the viability of S2R+ but not dNF1-KO cells after 48 h in serum-free media, as measured using

CellTiter Glo assays. In combination with CQ or bafilomycin A1, selumetinib resulted in a

significantly greater reduction in dNF1-KO cell viability compared to CQ alone, which was not

observed in S2R+ cells (n=3; p values obtained using a two-way ANOVA with Tukey for

*multiple comparisons). **(B)** In a similar experiment in human cell lines, selumetinib alone had*

no effect on the viability of C8, C23 (10 μM), ipnNF95.11C (5 μM), ipNF09.4, and ipNF05.5

*(10 μM) **(B-D)**; however, it did significantly decrease viability of ipNF95.11b 'C' NF1$^{-/-}$ (5 μM)*

(C). *Similar to observations in Drosophila cells, the effect of CQ was enhanced when*

combined with selumetinib in all three NF1$^{-/-}$ cell lines (50 nM, 100 pM, and 50 nM,

respectively), whereas the effect of bafilomycin A1 (100 pM, 1 nM, and 1 nM, respectively)

was enhanced by selumetinib only in the patient-derived NF1-deficient cell lines (ipNF95.11b

'C' and ipNF05.5). (n=3-4; p values obtained using a two-way ANOVA with Tukey for multiple

comparisons). In all cases, bars represent the mean and error bars indicate standard

*deviation. **(E)** Drug synergy between CQ + selumetinib or bafilomycin A1 + selumetinib was*

calculated using the coefficient of drug interaction (CDI). CDI<1 indicates synergy (with

CDI<0.7 indicating a strong synergistic effect), CDI=1 indicates additivity, and CDI>1 indicates

antagonism. In this case, drug pairs were classed as synergistic if CDI<0.95.

DISCUSSION

In this study, we used a *Drosophila* cell culture model of NF1 to identify new potential therapeutic targets and subsequently candidate drugs for the selective killing of *NF1*-deficient cells. Using genetic screening of *Drosophila* cells, we identified 54 candidate synthetic lethal genes that when knocked down resulted in the death of *NF1*-deficient cells without impacting the viability of wild-type, healthy cells. Of these genes, 85% (46/54) were validated in secondary assays, indicating the high quality of the screen results. A key outcome from this screen was the identification of *Atg8b* as a synthetic lethal partner of *NF1*. Atg8b (human ortholog: *GABARAP*) is an autophagy-related protein located in the autophagosome. Autophagy is the process where damaged organelles and unfolded proteins are sequestered into autophagosomes in the cytoplasm, which then undergo fusion with lysosomes (autolysosomes), resulting in degradation of the intracellular components (38). This is important in the regulation of cell growth, maturation, and death.

In cancer, autophagy is reported to take on two opposing roles: 1) degradation of damaged organelles and recycling of macromolecules to maintain a stable cellular environment, which prevents the formation of tumors (39); and 2) aiding in cancer cell survival in response to growth-limiting conditions, contributing to tumorigenesis (40). Furthermore, autophagy has been widely reported to promote cancer cell survival through a drug resistance mechanism (40, 41). Therefore, targeting the autophagy pathway is a potential therapeutic option in the treatment of cancer.

Autophagy is a complex process involving many different genes. In addition to *Atg8b*, *Uba1* was also identified in the screen, which is also linked to autophagy (42). While the selective effects of inhibiting *Atg8b* and *Uba1* were consistent and reproducible in secondary assays, it was surprising that more autophagy linked genes were not identified in our screens. Upon

further investigation, we found that approximately half of the 30 autophagy genes tested with

low throughput assays showed the selective effects observed with *Atg8b*. These genes have

roles from early-stage vesicle induction to late-stage fusion of the autophagosome and

lysosome. These results suggest that the entire autophagy pathway is dysregulated in *dNF1*-

KO cells, and it is likely that other autophagy genes were missed by the screen due to

ineffective RNAi reagents or noise.

Interestingly, *Atg5,* which is involved in the induction of autophagy via autophagic vesicle

formation, was found to have the most significant synthetic lethal interaction with *NF1.*

Previous studies have implicated ATG5 upregulation in RAS-induced autophagy and

malignant cell transformation (43, 44). However, at present, there are no commercially

available inhibitors of ATG5. Further studies are required to determine the exact autophagy

genes that are dysregulated in our panel of human *NF1*-deficient cell lines.

In total, we assessed seven autophagy inhibitors, and all reproduced the selective effects in

NF1-deficient cells. However, there was some variation in effects with some drugs not being

effective across all cell lines tested. CQ appears to be the most promising of the drugs that

we tested for clinical application. It is already approved for clinical use and produced the most

consistent and robust effects across the different NF1 cell models that we tested. CQ is a

widely used anti-malarial that functions to inhibit autophagy by blocking the binding of

autophagosomes to lysosomes by diffusing into the lysosomes and altering the acidic

environment, thereby inhibiting autophagic lysosomal degradation (25). Previous studies have

shown that CQ has anti-tumor properties in several types of cancer, including glioblastoma

(45, 46), hepatocellular carcinoma (47), prostate cancer (48), breast cancer (49), and

pancreatic cancer (50). In addition, a recent study reported that the expression of

metalloproteinase 1 (MMP1) was down-regulated in *NF1*-deficient fibroblasts, with a further

reduction associated with lysosomal degradation of MMP1. Interestingly, treatment of *NF1*-

deficient cells with CQ restored MMP1 expression via two mechanisms: activation of the

AHR/ERK pathway to enhance the mRNA and protein expression of MMP1, and inhibition of

the lysosomal degradation of MMP1 (51). Although this study did not assess whether

autophagy was dysregulated in the *NF1*-deficient fibroblasts and did not assess Schwann cells

or cells derived from NF1 tumors, it does highlight the therapeutic potential of autophagy

inhibition. Similarly, we found that CQ significantly reduced dNF1-KO cell viability relative to

wild-type S2R+ cells across a range of concentrations under conditions where autophagy is

induced (serum starvation of cells and *NF1* deficiency). This effect of CQ was conserved

across a panel of three human *NF1*-deficient cell lines, two of which were derived from

neurofibromas from NF1 patients. Furthermore, CQ altered the survival of *dNf1*-deficient

Drosophila in vivo and reduced the growth of *NF1*-mutant xenografts in mice to an even

greater extent than selumetinib. The doses of CQ and selumetinib administered to the mice

were deemed non-toxic, as has been widely reported previously (52, 53). Through the

inhibition of autophagy with CQ, we observed an increased level of apoptosis in the CQ-

treated xenografts, as assessed with the levels of cleaved PARP. Therefore, CQ shows great

potential as a therapeutic agent for *NF1*-associated tumors.

While there is a vast array of evidence for the efficacy and safety of CQ, the underlying

mechanisms of the tumor suppressive actions of CQ remain to be determined. One potential

mechanism is that under starvation conditions, such as those used in this study (i.e. serum

starvation), a reduction in glucose transport results in a release of mTOR inhibition of the ULK1

complex, inducing vesicle nucleation and facilitating the process of autophagy. Inhibition of

the lysosome using CQ has been shown to inhibit tumor growth and induce tumor cell death

in vitro (54, 55). We speculate that *NF1*-deficient cells are more susceptible to autophagy

inhibition with CQ because the baseline levels of autophagy are higher. RAS has previously

been shown to regulate autophagic flux (56). In addition, cancers associated with *RAS*

mutations have been reported to be dependent on autophagy, although this appears to be

tumor cell line dependent (57, 58). Therefore, cancers with aberrant RAS activity can be more

susceptible to autophagy inhibition with CQ. Neurofibromin functions as a RASGAP (GTPase-

activating protein), which facilitates RAS inactivation by enabling its GTPase activity (59). In

NF1-associated tumors, neurofibromin expression is downregulated or absent, resulting in

aberrant RAS activity and upregulation of the PI3K/Akt/mTORC1 pathway. Therefore,

aberrant RAS activity is expected to negatively regulate autophagy; however, RAS is

implicated in many signaling pathways and so it has a multifaceted role in autophagy

regulation. For example, RAS also activates the RAF1/MEK1/2/ERK and RAC1/MKK7/JNK

signaling pathways, both of which are known to activate autophagy (56). Furthermore,

upregulation of ATG5 and ATG7 has been implicated in RAS-induced autophagy and

malignant cell transformation (43, 44). Therefore, we speculate that aberrant RAS activity in

NF1-deficient cells results in the initiation of autophagy, and that inhibition of this pathway has

anti-tumor effects. When combining our data on the effects of CQ and bafilomycin A1, we

provide strong evidence for inhibition of the autophagy pathway as a target in the treatment of

NF1-associated tumors.

Finally, we found that inhibition of autophagy with CQ and bafilomycin A1 increased the

sensitivity of *NF1*-deficient cells (both dNF1-KO and human $NF1^{-/-}$ cells) to MEK1/2 inhibition

with selumetinib. Selumetinib is currently the only FDA-approved drug for the treatment of

tumors associated with neurofibromatosis type 1 (37). The phase 2 trial (SPRINT) for the use

of selumetinib in plexiform neurofibromas reported clinically meaningful improvements in 71%

of patients (13), prompting its FDA approval for patients ages 2 to 18 years with NF1 who have

symptomatic, inoperable plexiform neurofibromas. However, there are toxicity effects related

to long-term MEK inhibition. Interestingly, selumetinib had no selective effect in the dNF1-KO

cells compared to wild-type S2R+ controls when used alone. Furthermore, we observed no

difference in the phosphorylation of ERK between S2R+ and dNF1-KO cells under baseline

conditions, which suggests that RAS activity levels are similar in the two cell lines. Therefore,

while there is extensive evidence that RAS activation leads to dependence on autophagy, it is

possible that there is a second, RAS-independent, mechanism that also contributes to the

selective effect of autophagy inhibition. Such a mechanism may explain why we see a

combinatorial effect of autophagy inhibition and MEK1/2 inhibition. This finding is in line with

a previous study, which showed that combined inhibition of MEK1/2 plus autophagy had a

synergistic anti-proliferative effect in pancreatic ductal adenocarcinoma cell lines, which

display aberrant K-RAS activity, as well as patient-derived pancreatic ductal adenocarcinoma

xenograft tumors in mice (60).

In conclusion, we have shown using multiple techniques, reagents, and models that inhibition

of autophagy has potential as a novel therapeutic strategy for the treatment of NF1 tumors.

Given the existing clinical use of CQ and the robust and conserved effects that we observe

between cell culture models, this candidate drug has a high chance of successful translation

to the clinic, resulting in a positive impact on NF1 patients.

METHODS

Cell culture

Drosophila Schneider (S2R+) cells, both WT and dNF1-KO, were cultured at 25°C in

Schneider's media (Gibco) containing 1% antibiotic (Gibco) and 10% fetal bovine serum

(Gibco). Human cell lines used include: $NF1^{+/-}$ and $NF1^{-/-}$ immortalized human Schwann cell

(SC) lines, derived from the ipn02.3 2λ cell line using CRISPR/Cas9 gene editing as described

below; two pairs of immortalized human SC lines derived from plexiform neurofibromas from

NF1 patients (22), which included ipnNF95.11C ($NF^{+/-}$) and ipNF95.11b 'C' ($NF1^{-/-}$) cells

(germline *NF1* mutation: c.1756delACTA), and ipnNF09.4 ($NF1^{+/-}$) and ipNF05.5 ($NF1^{-/-}$) cells

(germline *NF1* mutation: c.3456_3457insA). All human cell lines were cultured at 37℃ in 5%

CO_2 in DMEM media (Merck) containing 1% antibiotic (Gibco) and 10% fetal bovine serum

(Gibco). For ipnNF09.4 and ipNF05.5 cells, the media was also supplemented with 50 ng/ml

neuregulin-1 (NRG-1) (Sigma).

Generation of dNF1-KO *Drosophila* cells

The dNF1-KO cell line was generated using methods described previously (19, 61). WT S2R+

cells were co-transfected with the pl018 plasmid to express Cas9 and the sgRNA

(CGCTTCTCCCTTGTCATATC) and pAct-GFP plasmid to mark transfected cells. Five days

after transfection, individual cells with the highest 15% GFP signal were isolated and seeded

into wells of 96-well plates using FACS. Single cells were cultured using conditioned media

for approximately 3 weeks. DNA was then extracted from each candidate cell population and

assessed using HRMA to identify those carrying mutations at the sgRNA target site. Positive

candidates were then sequenced to confirm frame-shift mutations were present in each *NF1*

allele (**Figure 1A**).

Generation of *NF1*-deficient human Schwann cell lines using CRISPR-Cas9

The wild-type human hTERT-immortalized Schwann cell line (hTERT ipn02.3 2λ) was a gift

from Dr. Peggy Wallace, University of Florida (22). CRISPR-Cas9 genome editing was

performed using NF1-sg2 (ACACTGGAAAAATGTCTTGC), a sgRNA targeting human *NF1*

exon 3 in the lentiCRISPR v2 plasmid, which was a gift from Dr. Feng Zhang (Broad Institute)

(62). hTERT ipn02.3 2λ cells were transfected by electroporation using the Amaxa Basic

Nucleofector Kit for Primary Mammalian Fibroblasts (Lonza) and the program U-023,

according to the manufacturer's instructions. After 48 h of selection on puromycin (1 μg/ml) in

DMEM/10% FBS, cells were re-plated in to allow single clone isolation. Genotyping of selected

single clones was performed using PCR using primers NF1-exon3-FOR

(CCCCAATTCAAGATTCTGGT) and NF1-exon3-REV (ATCGCACTCTCCCACAACTC). PCR

products were treated with ExoSAP (Affymetrix) and sequenced using NF1-exon3-SEQ

(TGCCATTTCTGTTTGCCTTA).

Synthetic lethal screen

Screens were performed using wild-type S2R+ or dNF1-KO cells using the genome-wide RNAi

library from the Sheffield RNAi Screening Facility. In total, 10,000 cells were seeded into each

library well in 10 μl serum-free Schneider's *Drosophila* media. Plates were then incubated at

room temperature for 45 minutes before addition of 35 μl media with 10% FBS. Libraries were

incubated at 25°C for 5 days before CellTiter-Glo assays (Promega) were performed. Screens

were performed in triplicate in each cell line (**Figure S3**).

Data were analyzed by first normalizing all values to the median of each row and column of

the library plate to allow direct comparison between plates. Z-scores were then calculated for

each RNAi reagent using the average and standard deviation of each replicate screen and

correlation between replicates was used to assess the quality of screen results. Reagents

were considered hits if the Z-score in at least 2/3 of replicates was below -1.5 in dNF1-KO

cells and above -1.5 in S2R+ cells. We note that some assay plates were affected by position

effects; these were identified manually and removed from the analysis prior to correlation

analysis.

Functional gene groups were determined using the in-build clustering tool in the STRING

database to group genes. Functions of each group were determined manually by searching

for associated GO terms or assessing annotated functions of human orthologs in cases where

the *Drosophila* gene was not sufficiently annotated.

Variable dose analysis (VDA)

VDA is an RNAi-based method in which each cell within a population receives a different dose

of shRNA (20, 63). The relative knockdown efficiency of each cell is then measured with a

fluorescent reporter. On the day of transfection S2R+ and dNF1-KO cells were plated at 1 x

10^4 cell/100 µl culture media, per well of a 96-well plate. Cells were incubated at 25℃ for 40

mins to allow adhesion. Cells were then transfected with 40 ng actin-GFP and 160 ng shRNA

expression plasmid using 0.6 µl FuGENE® HD transfection reagent in a total volume of 10 µl.

We used a positive (*thread*, an apoptosis inhibitor that induces cell death when inhibited) and

negative (*white*, known to have no viability effect in these cells) shRNA on each plate for

normalization of data. Plates were then sealed and incubated for 4 days at 25℃ in a

humidifying chamber.

Flow cytometry was used to identify GFP-positive cells (transfection efficiency). Area under

an inverted cumulative distribution curve was used as a readout of relative viability, normalized

to the positive and negative control. More detailed protocol and data analysis information can

be found in Sierzputowska et al. (63).

CellTiter-Glo assays

The CellTiter-Glo assay (Promega; G7570) is a luminescent viability assay based on the

quantification of ATP, which is an indicator of metabolically active cells. All cells were plated

in white 384-well plates at a seeding density of 5 x 10^3 cells/25 µl culture media in either

complete or serum-free media. S2R+ cells were left to adhere for 40 mins at 25℃ and human

cells were left to adhere for 4 h at 37℃ in 5% CO_2 before treatment. Cells were treated with

250 nl per well of each drug (CQ, bafilomycin A1, and selumetinib) in PBS at varying

concentrations in replicates of 5-8 using the Mosquito LV Genomics (SPT Labtech) and

incubated for 48 h. The CellTiter-Glo assay was then performed according to the

manufacturer's instructions, and luminescence was measured using a plate reader (TECAN

Infinite M200 Pro).

Autophagy assay and Caspase assay

An autophagy assay kit (Abcam; ab139484) was used to measure autophagic vacuoles in live

cells using a dye that selectively labels autophagic vacuoles. All cells were plated in 96-well

plates in either complete or serum-free media. S2R+ cells were left to adhere for 40 mins at

25℃ and human cells were left to adhere for 4 h at 37℃ in 5% CO_2 before treatment. Cells

were treated with 10 µM per well of CQ in PBS at varying concentrations and incubated for 4

h. The autophagy assay was then performed according to the manufacturer's instructions, and

fluorescence intensity was measured using a plate reader (TECAN Infinite M200 Pro).

Autophagic flux was quantified by inhibiting lysosomal fusion with CQ and then measuring the

initial rate of accumulation of autophagosomes using a fluorescent plate reader.

Caspase activity was assessed using the same plating and treatment method, except that

cells were treated for 48 h, via the generic caspase activity assay kit (Abcam; ab112130) per

the   manufacturer's   instructions.   In   both   the   autophagy   and   caspase   assay,

autophagy/caspase fluorescence was normalized to DAPI in each well.

Annexin V and PI staining

The Annexin V-FITC apoptosis kit (Abcam; ab14085) was used to detect apoptosis by staining

of phosphatidylserine molecules that have translocated to the outside of the cell membrane.

Cells were co-stained with propidium iodide (PI) to detect dead cells in the population. S2R+ cells were left to adhere for 40 mins at 25℃ before treatment. Cells were treated with 1 µl per well of each drug (CQ, bafilomycin A1) in PBS at varying concentrations, and incubated for 48 h. The annexin-V/PI assay was then performed according to the manufacturer's instructions, and fluorescence intensity was measured using flow cytometry.

Western blotting

Lysates from S2R+ cells, dNF-KO cells, human immortalized Schwann cells, and homogenized xenograft tissue were prepared in RIPA buffer supplemented with protease inhibitors (Complete Protease Inhibitor Cocktail, Sigma) and phosphatase inhibitors (NaF/Na$_3$VO$_4$/β-glycerophosphate). Lysates for testing expression of neurofibromin were resolved on NuPAGE 3–8% Tris-Acetate and for assessing levels of pERK/ERK on NuPAGE 4-12% Bis-Tris protein gels. After transfer to nitrocellulose membranes, blots were processed according to the Odyssey CLx protocol (LI-COR). Antibodies used for immunoblotting: anti-*Drosophila* neurofibromin (mouse, ascites purified mAb21 and mAb30, 1:500 each), anti-human neurofibromin (mouse, Infixion r07E, 1:1000), anti-phospho-ERK (mouse, Sigma M8159, 1:2500), total ERK (rabbit, CST 9102, 1:1000), anti-β-tubulin (mouse, Developmental Studies Hybridoma Bank E7, 1:10,000), GAPDH (rabbit, Proteintech 10494-1-AP, 1:15,000), and anti-PARP (rabbit, Cell Signaling 952, 1:1000). Secondary antibodies used were anti-mouse Alexa Fluor Plus 800 (goat, Invitrogen, A32730, 1:10,000) and anti-rabbit Alexa Fluor Plus 680 (goat, Invitrogen, A21109, 1:10,000).

***Drosophila* husbandry and stocks**

Flies were cultured on cornmeal/agar food medium according to standard protocol and housed at 25°C.

The *Nf1^C1* mutant fly line was generated by CRISPR/Cas9 gene editing (64). using the sgRNA

line: *y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8]=TKO.GS01796}attP40* (GS01796 sgRNA

sequence: CGCTTCTCCCTTGTCATATC) and a germline source of Cas9: *y[1]*

M{w[+mC]=nos-Cas9.P}ZH-2A w[*] (Bloomington Stock #54591). The *UAS-dNf1* transgene

encodes full-length *dNf1* cDNA corresponding to the RF isoform with addition of introns 9 and

10 and was previously published (65). The RNAi and landing site control lines were obtained

from the Vienna *Drosophila* RNAi Center: *dNf1 RNAi* line: P{KK101909}VIE-260B (VDRC

#109637) and VIE-260B (VDRC #60100). *UAS-Dicer2* was included to potentiate the RNAi

effect (66): *P{w[+mC]=UAS-Dcr-2.D}10* (Bloomington: 24651). Male flies were used for all

experiments.

Assessing drug sensitivity in flies

Flies were raised on standard fly food and kept at 25°C. For testing drugs, adult flies were

transferred to Formula 4-24 Instant *Drosophila* Medium (Carolina Biological) prepared using

water (control) or chloroquine (chloroquine diphosphate; Sigma C6628) in water.

Concentrations of CQ tested ranged from 0-40 µM (Figure S7) and 35 µM was used for the

experiments in Figure 6 D and E. For each genotype, three replicates of 20 flies were added

to each condition and kept at 25°C. Flies were monitored periodically to assess lethality.

Survival was reported as the percent of flies still alive at each time point.

NF1 tumor xenografts

All experiments were conducted in accordance with UK legislation and with local ethics

committee approval (University of Exeter AWERB). Two million ST88-14 cells mixed 1:1 with

Matrigel (ThermoFisher Scientific) were injected subcutaneously into the right flank of male

CD-1 nude mice (Charles River Laboratories). Tumors were measured three times weekly

with a caliper, and the tumor volume was calculated as follows: [(length + width) / 2] x length

x width. Once the tumor diameter began to increase over two separate measurements, mice

were intraperitoneally injected with either saline or CQ (50 mg/kg in saline) or received

selumetinib (25 mg/kg in saline) via oral gavage three times per week (n=6 mice per group).

Mice were culled by cervical dislocation (Schedule 1) when the control tumor sizes reached

the allowed endpoint (12 mm in diameter), and tumors were dissected. Tumors were flash

frozen for further analysis.

Competing Interest Statement: B.E.H. is a shareholder and founding director of Quest

Genetics Ltd. The remaining authors declare no competing interests.

Acknowledgements

This study was funded by a Medical Research Council project grant (MR/V009583/1) and a

Medical Research Council Confidence in Concept (CiC) award (MC_PC_19037) to B.H. and

NIH R21 NS096402, DOD-NFRP W81XWH-16-1-0220 and the Neurofibromatosis

Therapeutic Acceleration Program (NTAP) (J.A.W.). We would also like to thank Nerve

Tumours UK for their support as collaborators on the CiC award. S.J.B was supported by the

Dorothy and Spiro Latsis Fellowship for NF1 Research. N.P. is an investigator of HHMI. We

thank the Sheffield RNAi Screening Facility, Biomedical Sciences, University of Sheffield,

supported by the Wellcome Trust (grant reference number 084757) for providing RNAi

libraries.

REFERENCES

1.   Evans DG, Howard E, Giblin C, Clancy T, Spencer H, Huson SM, et al. Birth incidence

and prevalence of tumor-prone syndromes: estimates from a UK family genetic register

service. Am J Med Genet A. 2010;152A(2):327-32.

2.   Gutmann DH, Ferner RE, Listernick RH, Korf BR, Wolters PL, Johnson KJ.

Neurofibromatosis type 1. Nat Rev Dis Primers. 2017;3:17004.

3.   Lin AL, Gutmann DH. Advances in the treatment of neurofibromatosis-associated

tumours. Nat Rev Clin Oncol. 2013;10(11):616-24.

4.   Rasmussen SA, Yang Q, Friedman JM. Mortality in neurofibromatosis 1: an analysis

using U.S. death certificates. Am J Hum Genet. 2001;68(5):1110-8.

5.   Martin GA, Viskochil D, Bollag G, McCabe PC, Crosier WJ, Haubruck H, et al. The

GAP-related domain of the neurofibromatosis type 1 gene product interacts with ras p21. Cell.

1990;63(4):843-9.

6.   Xu GF, Lin B, Tanaka K, Dunn D, Wood D, Gesteland R, et al. The catalytic domain of

the neurofibromatosis type 1 gene product stimulates ras GTPase and complements ira

mutants of S. cerevisiae. Cell. 1990;63(4):835-41.

7.   Viskochil D, White R, Cawthon R. The neurofibromatosis type 1 gene. Annu Rev

Neurosci. 1993;16:183-205.

8.   Ratner N, Miller SJ. A RASopathy gene commonly mutated in cancer: the

neurofibromatosis type 1 tumour suppressor. Nat Rev Cancer. 2015;15(5):290-301.

9.   Simanshu DK, Nissley DV, McCormick F. RAS Proteins and Their Regulators in

Human Disease. Cell. 2017;170(1):17-33.

10.   Abdel-Rahman O, ElHalawani H, Ahmed H. Risk of Selected Cardiovascular Toxicities

in Patients With Cancer Treated With MEK Inhibitors: A Comparative Systematic Review and

Meta-Analysis. J Glob Oncol. 2015;1(2):73-82.

11.   Avery RA, Trimboli-Heidler C, Kilburn LB. Separation of outer retinal layers secondary

to selumetinib. J AAPOS. 2016;20(3):268-71.

12.   Baldo F, Magnolato A, Barbi E, Bruno I. Selumetinib side effects in children treated for

plexiform neurofibromas: first case reports of peripheral edema and hair color change. BMC

Pediatr. 2021;21(1):67.

13.   Gross AM, Wolters PL, Dombi E, Baldwin A, Whitcomb P, Fisher MJ, et al. Selumetinib

in Children with Inoperable Plexiform Neurofibromas. N Engl J Med. 2020;382(15):1430-42.

14.   Kaelin WG, Jr. The concept of synthetic lethality in the context of anticancer therapy.

Nat Rev Cancer. 2005;5(9):689-98.

15.   Lord CJ, Ashworth A. PARP inhibitors: Synthetic lethality in the clinic. Science.

2017;355(6330):1152-8.

16.   Ryan CJ, Bajrami I, Lord CJ. Synthetic Lethality and Cancer - Penetrance as the Major

Barrier. Trends Cancer. 2018;4(10):671-83.

17.   Valvezan AJ, Turner M, Belaid A, Lam HC, Miller SK, McNamara MC, et al. mTORC1

Couples Nucleotide Synthesis to Nucleotide Demand Resulting in a Targetable Metabolic

Vulnerability. Cancer Cell. 2017;32(5):624-38 e5.

18.   Nicholson HE, Tariq Z, Housden BE, Jennings RB, Stransky LA, Perrimon N, et al.

HIF-independent synthetic lethality between CDK4/6 inhibition and VHL loss across species.

Sci Signal. 2019;12(601).

19.   Housden BE, Valvezan AJ, Kelley C, Sopko R, Hu Y, Roesel C, et al. Identification of

potential drug targets for tuberous sclerosis complex by synthetic screens combining CRISPR-

based knockouts with RNAi. Sci Signal. 2015;8(393):rs9.

20.   Housden BE, Li Z, Kelley C, Wang Y, Hu Y, Valvezan AJ, et al. Improved detection of

synthetic lethal interactions in Drosophila cells using variable dose analysis (VDA). Proc Natl

Acad Sci U S A. 2017;114(50):E10755-E62.

21.   Lee H, McManus CJ, Cho DY, Eaton M, Renda F, Somma MP, et al. DNA copy number

evolution in Drosophila cell lines. Genome Biol. 2014;15(8):R70.

22.   Li H, Chang LJ, Neubauer DR, Muir DF, Wallace MR. Immortalization of human normal

and NF1 neurofibroma Schwann cells. Lab Invest. 2016;96(10):1105-15.

23.   Szklarczyk D, Gable AL, Lyon D, Junge A, Wyder S, Huerta-Cepas J, et al. STRING

v11: protein-protein association networks with increased coverage, supporting functional

discovery in genome-wide experimental datasets. Nucleic Acids Res. 2019;47(D1):D607-D13.

24.   Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, et al. Cytoscape: a

software environment for integrated models of biomolecular interaction networks. Genome

Res. 2003;13(11):2498-504.

25.   Homewood CA, Warhurst DC, Peters W, Baggaley VC. Lysosomes, pH and the anti-

malarial action of chloroquine. Nature. 1972;235(5332):50-2.

26.   Mauvezin C, Neufeld TP. Bafilomycin A1 disrupts autophagic flux by inhibiting both V-

ATPase-dependent acidification and Ca-P60A/SERCA-dependent autophagosome-lysosome

fusion. Autophagy. 2015;11(8):1437-8.

27.   Ursing J, Rombo L, Eksborg S, Larson L, Bruvoll A, Tarning J, et al. High-Dose

Chloroquine for Uncomplicated Plasmodium falciparum Malaria Is Well Tolerated and Causes

Similar QT Interval Prolongation as Standard-Dose Chloroquine in Children. Antimicrob

Agents Chemother. 2020;64(3).

28.   Levy JMM, Towers CG, Thorburn A. Targeting autophagy in cancer. Nat Rev Cancer.

2017;17(9):528-42.

29.   Kocaturk NM, Akkoc Y, Kig C, Bayraktar O, Gozuacik D, Kutlu O. Autophagy as a
molecular target for cancer treatment. Eur J Pharm Sci. 2019;134:116-37.

30.   Sorensen MG, Henriksen K, Neutzsky-Wulff AV, Dziegiel MH, Karsdal MA. Diphyllin,
a novel and naturally potent V-ATPase inhibitor, abrogates acidification of the osteoclastic
resorption lacunae and bone resorption. J Bone Miner Res. 2007;22(10):1640-8.

31.   King LB, Boto T, Botero V, Aviles AM, Jomsky BM, Joseph C, et al. Developmental
loss of neurofibromin across distributed neuronal circuits drives excessive grooming in
Drosophila. PLoS Genet. 2020;16(7):e1008920.

32.   Moscato EH, Dubowy C, Walker JA, Kayser MS. Social Behavioral Deficits with Loss
of Neurofibromin Emerge from Peripheral Chemosensory Neuron Dysfunction. Cell Rep.
2020;32(1):107856.

33.   Bai L, Lee Y, Hsu CT, Williams JA, Cavanaugh D, Zheng X, et al. A Conserved
Circadian Function for the Neurofibromatosis 1 Gene. Cell Rep. 2018;22(13):3416-26.

34.   Walker JA, Gouzi JY, Long JB, Huang S, Maher RC, Xia H, et al. Genetic and
functional studies implicate synaptic overgrowth and ring gland cAMP/PKA signaling defects
in the Drosophila melanogaster neurofibromatosis-1 growth deficiency. PLoS Genet.
2013;9(11):e1003958.

35.   Gouzi JY, Moressis A, Walker JA, Apostolopoulou AA, Palmer RH, Bernards A, et al.
The receptor tyrosine kinase Alk controls neurofibromin functions in Drosophila growth and
learning. PLoS Genet. 2011;7(9):e1002281.

36.   Liao CP, Pradhan S, Chen Z, Patel AJ, Booker RC, Le LQ. The role of nerve
microenvironment for neurofibroma development. Oncotarget. 2016;7(38):61500-8.

37.   Markham A, Keam SJ. Selumetinib: First Approval. Drugs. 2020;80(9):931-7.

38.   Zhang XJ, Chen S, Huang KX, Le WD. Why should autophagic flux be assessed? Acta
Pharmacol Sin. 2013;34(5):595-9.

39.   Mizushima N, Levine B, Cuervo AM, Klionsky DJ. Autophagy fights disease through

cellular self-digestion. Nature. 2008;451(7182):1069-75.

40.   O'Donovan TR, O'Sullivan GC, McKenna SL. Induction of autophagy by drug-resistant

esophageal cancer cells promotes their survival and recovery following treatment with

chemotherapeutics. Autophagy. 2011;7(5):509-24.

41.   Peng X, Gong F, Chen Y, Jiang Y, Liu J, Yu M, et al. Autophagy promotes paclitaxel

resistance of cervical cancer cells: involvement of Warburg effect activated hypoxia-induced

factor 1-alpha-mediated signaling. Cell Death Dis. 2014;5:e1367.

42.   Chang TK, Shravage BV, Hayes SD, Powers CM, Simin RT, Wade Harper J, et al.

Uba1 functions in Atg7- and Atg3-independent autophagy. Nat Cell Biol. 2013;15(9):1067-78.

43.   Byun JY, Yoon CH, An S, Park IC, Kang CM, Kim MJ, et al. The Rac1/MKK7/JNK

pathway signals upregulation of Atg5 and subsequent autophagic cell death in response to

oncogenic Ras. Carcinogenesis. 2009;30(11):1880-8.

44.   Kim MJ, Woo SJ, Yoon CH, Lee JS, An S, Choi YH, et al. Involvement of autophagy

in oncogenic K-Ras-induced malignant cell transformation. J Biol Chem. 2011;286(15):12924-

32.

45.   Golden EB, Cho HY, Jahanian A, Hofman FM, Louie SG, Schonthal AH, et al.

Chloroquine enhances temozolomide cytotoxicity in malignant gliomas by blocking autophagy.

Neurosurg Focus. 2014;37(6):E12.

46.   Zanotto-Filho A, Braganhol E, Klafke K, Figueiro F, Terra SR, Paludo FJ, et al.

Autophagy inhibition improves the efficacy of curcumin/temozolomide combination therapy in

glioblastomas. Cancer Lett. 2015;358(2):220-31.

47.   Mei L, Chen Y, Wang Z, Wang J, Wan J, Yu C, et al. Synergistic anti-tumour effects of

tetrandrine and chloroquine combination therapy in human cancer: a potential antagonistic

role for p21. Br J Pharmacol. 2015;172(9):2232-45.

48.   Farrow JM, Yang JC, Evans CP. Autophagy as a modulator and target in prostate

cancer. Nat Rev Urol. 2014;11(9):508-16.

49.   Lefort S, Joffre C, Kieffer Y, Givel AM, Bourachot B, Zago G, et al. Inhibition of

autophagy as a new means of improving chemotherapy efficiency in high-LC3B triple-negative

breast cancers. Autophagy. 2014;10(12):2122-42.

50.   Yang A, Kimmelman AC. Inhibition of autophagy attenuates pancreatic cancer growth

independent of TP53/TRP53 status. Autophagy. 2014;10(9):1683-4.

51.   Tsuji G, Takai-Yumine A, Kato T, Furue M. Metalloproteinase 1 downregulation in

neurofibromatosis 1: Therapeutic potential of antimalarial hydroxychloroquine and

chloroquine. Cell Death Dis. 2021;12(6):513.

52.   Tang MC, Wu MY, Hwang MH, Chang YT, Huang HJ, Lin AM, et al. Chloroquine

enhances gefitinib cytotoxicity in gefitinib-resistant nonsmall cell lung cancer cells. PLoS One.

2015;10(3):e0119135.

53.   Troiani T, Vecchione L, Martinelli E, Capasso A, Costantino S, Ciuffreda LP, et al.

Intrinsic resistance to selumetinib, a selective inhibitor of MEK1/2, by cAMP-dependent protein

kinase A activation in human lung and colorectal cancer cells. Br J Cancer.

2012;106(10):1648-59.

54.   Eng CH, Wang Z, Tkach D, Toral-Barza L, Ugwonali S, Liu S, et al. Macroautophagy

is dispensable for growth of KRAS mutant tumors and chloroquine efficacy. Proc Natl Acad

Sci U S A. 2016;113(1):182-7.

55.   Jia L, Wang J, Wu T, Wu J, Ling J, Cheng B. In vitro and in vivo antitumor effects of

chloroquine on oral squamous cell carcinoma. Mol Med Rep. 2017;16(5):5779-86.

56.   Schmukler E, Kloog Y, Pinkas-Kramarski R. Ras and autophagy in cancer

development and therapy. Oncotarget. 2014;5(3):577-86.

57. Guo JY, Chen HY, Mathew R, Fan J, Strohecker AM, Karsli-Uzunbas G, et al.

Activated Ras requires autophagy to maintain oxidative metabolism and tumorigenesis.

Genes Dev. 2011;25(5):460-70.

58. Morgan MJ, Gamez G, Menke C, Hernandez A, Thorburn J, Gidan F, et al. Regulation

of autophagy and chloroquine sensitivity by oncogenic RAS in vitro is context-dependent.

Autophagy. 2014;10(10):1814-26.

59. Downward J. Targeting RAS signalling pathways in cancer therapy. Nat Rev Cancer.

2003;3(1):11-22.

60. Kinsey CG, Camolotto SA, Boespflug AM, Guillen KP, Foth M, Truong A, et al.

Protective autophagy elicited by RAF-->MEK-->ERK inhibition suggests a treatment strategy

for RAS-driven cancers. Nat Med. 2019;25(4):620-7.

61. Housden BE, Nicholson HE, Perrimon N. Synthetic Lethality Screens Using RNAi in

Combination with CRISPR-based Knockout in Drosophila Cells. Bio Protoc. 2017;7(3).

62. Shalem O, Sanjana NE, Hartenian E, Shi X, Scott DA, Mikkelson T, et al. Genome-

scale CRISPR-Cas9 knockout screening in human cells. Science. 2014;343(6166):84-7.

63. Sierzputowska K, Baxter CR, Housden BE. Variable Dose Analysis: A Novel High-

throughput RNAi Screening Method for Drosophila Cells. Bio Protoc. 2018;8(24):e3112.

64. Ren X, Sun J, Housden BE, Hu Y, Roesel C, Lin S, et al. Optimized gene editing

technology for Drosophila melanogaster using germ line-specific Cas9. Proc Natl Acad Sci U

S A. 2013;110(47):19012-7.

65. Walker JA, Tchoudakova AV, McKenney PT, Brill S, Wu D, Cowley GS, et al. Reduced

growth of Drosophila neurofibromatosis 1 mutants reflects a non-cell-autonomous requirement

for GTPase-Activating Protein activity in larval neurons. Genes Dev. 2006;20(23):3311-23.

66.   Dietzl G, Chen D, Schnorrer F, Su KC, Barinova Y, Fellner M, et al. A genome-wide

transgenic RNAi library for conditional gene inactivation in Drosophila. Nature.

2007;448(7150):151-6.