CRISPR/Cas9 knockout of EPFL10 reduces stomatal density while maintaining photosynthesis and enhancing water conservation in rice

Nicholas G. Karavolias, Dhruv Patel, Kyungyong Seong, Michelle Tjahjadi, Gloria-Alexandra Gueorguieva, Jaclyn Tanaka, Douglas Dahlbeck, Myeong-Je Cho, Krishna K. Niyogi, Brian J. Staskawicz

Summary

Rice production is of paramount importance for global nutrition and potential yields will be detrimentally affected by climate change. Rice stomatal developmental genetics were explored as a mechanism to improve water use efficiency while maintaining yield under climate stress.

Gene-editing of STOMAGEN and its paralog, EPFL10, using CRISPR/Cas9 in rice cv. Nipponbare yielded lines with altered stomatal densities that were functionally characterized. CRISPR/Cas9 mediated knockouts of EPFL10 and STOMAGEN yielded lines with c. 80% and 25% of wild-type stomata, respectively.

***epfl10* lines with small reductions in stomatal densities are able to conserve water to similar extents as *stomagen* lines with large stomatal density reductions but do not suffer from any concomitant reductions in stomatal conductance, carbon assimilation, or thermoregulation.**

The duplicate of STOMAGEN, EPFL10, is a weak positive regulator of stomatal development in rice. *epfl10* lines maintained wild-type physiological characteristics while conserving more water. Modest reductions in stomatal densities may be a climate-adaptive approach in rice that can safeguard yield.

Introduction

The need to develop climate change adapted crops in the face of rapid global population increases and worsening climates is necessary and timely. Prolonged periods of drought and increased desertification are anticipated to become more prevalent in the next century (IPCC et al., 2021). Climate change modeling predicts increases in global temperatures by 2-4 °C by the end of the 21st century. Increased temperatures alone and in combination with limited water will negatively impact crop yields (Jagadish, Murty, & Quick, 2015; Rang, Jagadish, Zhou, Craufurd, & Heuer, 2011; Shah et al., 2011). Rice (*Oryza sativa)* is the most widely directly consumed crop globally and trails only wheat and maize in area harvested (Food and Agriculture Organisation (FAO), 2020). Originally domesticated in semi-aquatic habitats, rice is especially sensitive to drought relative to other C3 cereal crops (Bernier, Atlin, Serraj, Kumar, & Spaner, 2008; Jägermeyr et al., 2021; Lafitte, Ismail, & Bennett, 2004).

Future water limitations may necessitate transitions of fully flooded paddy conditions to water-
saving production schemes (Jagadish et al., 2015). Rain-fed production, which comprises about
45% of total rice grown, is particularly susceptible to drought as a result of unpredictable
precipitation (Khush, 1997; Pandey et al., 2007; Tuong & Bouman, 2009). Furthermore, most
regions where irrigated rice is produced are currently experiencing or are projected to experience
water scarcity (Tuong & Bouman, 2009). Thus, all rice, regardless of production method, would
benefit from improvements that maintain yields with lower water requirements. 90% of water
loss in rice occurs via transpiration from the stomata necessitating explorations of stomata-driven
improvements in water conservation (T. N. Buckley, 2005).

Stomata are at the nexus of plants and the atmosphere. They facilitate gaseous exchanges of
carbon dioxide, oxygen, and water vapor. The development of these essential structures has been
studied extensively in model systems with growing understandings of development in non-model
plant as well (Endo & Torii, 2019; Liu, Ohashi-Ito, & Bergmann, 2009; Tiago D.G. Nunens, Dan
Zhang, 2019) . Grass stomata are morphologically distinct from classically studied *Arabidopsis*
thaliana stomata, possessing physiologically linked subsidiary cells flanking dumbbell-shaped
guard cells among other unique features (Stebbins & Shah, 1960; Tiago D.G. Nunens, Dan
Zhang, 2019). Furthermore, grass stomata mature basipetally along the longitudinal axis of
leaves and are distributed in files adjacent to vasculature unlike the stomata of *Arabidopsis*
thaliana (Tiago D.G. Nunens, Dan Zhang, 2019). In parallel with their divergent structure, the
grass stomatal developmental framework contains variations from well-studied eudicot
regulatory pathways which account for some of the observed phenotypic variation. For example,
the *Brachypodium distachyon* ortholog of the conserved transcription factor MUTE has evolved
a novel role in specifying subsidiary cell identity through acquisition of cell-to-cell mobility
(Raissig et al., 2017).

Despite these differences, the core regulatory wiring of stomatal development is largely
conserved in angiosperms (Liu et al., 2009). Epidermal patterning factors (EPF) are essential to
the regulation of stomatal development. EPFs in the context of stomatal development, are mobile
peptides that regulate cell fate transitions and cell divisions to ensure proper spacing and number
of stomata (Hara, Kajita, Torii, Bergmann, & Kakimoto, 2007; Hara et al., 2009; Hunt & Gray,
2009; Shimada, Sugano, & Hara-Nishimura, 2011). These peptides are cysteine rich with
conserved cysteine residues near the C-terminal end. EPF1 and EPF2 function as negative
regulators. EPF2 in *Arabidopsis thaliana* is known to act before EPF1 to inhibit entry into the
stomatal lineage by limiting asymmetric divisions (Hara et al., 2009; Hunt & Gray, 2009). EPF1
regulates the differentiation of guard mother cells in *Arabidopsis thaliana.*(Hara et al., 2007).

Unlike EPF1 and EPF2 which serve as negative regulators, EPF-LIKE9, or STOMAGEN, is a
positive regulator of stomatal development in *Arabidopsis thaliana and O. sativa* (Hunt, Bailey,
& Gray, 2010; Sugano et al., 2010a; Yin et al., 2017) EPF1 and EPF2 are expressed in the
stomatal lineage cells, whereas EPFL9 is mesophyll derived (Hara et al., 2007, 2009; Hunt &
Gray, 2009; Shimada et al., 2011). EPFL9 is composed of three distinct regions: an N-terminal
signal peptide region, a pro-peptide region, and a C-terminal cysteine-rich active peptide region
(Ohki, Takeuchi, & Mori, 2011; Sugano et al., 2010a). The full-length peptide is processed in-
vivo to yield a 45 C-terminal amino acid active peptide (Lee et al., 2015; Sugano et al., 2010a) .
The active peptide encoded by EPFL9 possesses the conserved cysteine residues of other EPF1

and EPF2 and binds the same ERECTA (ER)-family receptors and co-receptor TOO MANY
MOUTHS (TMM) in *Arabaidopsis thaliana* (Hepworth, Caine, Harrison, Sloan, & Gray, 2018;
Lee et al., 2015; Shimada et al., 2011; Sugano et al., 2010b).

EPF2 and EPFL9 competitively bind to the ERECTA (ER)-family and co-receptor TOO MANY
MOUTHES (TMM) to mediate downstream effects (Lee et al., 2015). Binding of EPFL9 to the
TMM/ER complex prevents inhibition of stomatal development in *Arabidopsis thaliana.* (Lee et
al., 2015) STOMAGEN knockdown using RNAi in *A. thaliana* reduces stomatal densities to
40% of wild type (Sugano et al., 2010a). Likewise, a knockout of EPFL9 in rice using
CRISPR/Cas9 and CRISPR/Cpf1 yielded an eightfold reduction in abaxial stomatal density in
the IR64 background (Yin et al., 2017).

Interestingly, EPFL9 has undergone a duplication event in Poaceae leading to multiple copies in
all species surveyed (Hepworth et al., 2018). Overexpression of rice STOMAGEN and its
duplicate, previously named EPFL9-2, in *A. thaliana* revealed a common, though reduced,
function of EPFL9-2 as a positive regulator of stomatal development when ectopically expressed
(Lu et al., 2019). Ectopic expression of *Brachypodium distachyon* and *Triticum aestivum*
STOMAGEN and STOMAGEN paralogs in *Arabidopsis thaliana* resulted in similar stomatal
density increases (Jangra et al., 2021). In contrast, overexpression of negative regulators of
stomatal development reduced stomatal density and improved water use efficiency in
Arabidopsis thaliana, wheat, barley, and rice. (Caine et al., 2019; Dunn et al., 2019; Franks, W.
Doheny-Adams, Britton-Harper, & Gray, 2015; Hepworth, Doheny-Adams, Hunt, Cameron, &
Gray, 2015; Hughes et al., 2017; Mohammed et al., 2019). However, all stomatal density
reductions achieved by overexpressing negative regulators of stomatal development to any extent
also reduced stomatal conductance and carbon assimilation under physiologically relevant light
conditions (Caine et al., 2019; Dunn et al., 2019; Hughes et al., 2017). For example, rice lines
overexpressing EPF1 to reduce stomatal densities, exhibited lower stomatal conductance and
carbon assimilation at all light conditions that exceeded 1000 μmol photos $m^{-2}s^{-1}$ (Caine et al.,
2019).

Stomatal conductance is essential for crop productivity (Fischer et al., 1998; Kusumi,
Hashimura, Yamamoto, Negi, & Iba, 2017; Ohsumi, Kanemura, Homma, Horie, & Shiraiwa,
2007a; Richards, 2000; Roche, 2015; Taylaran, Adachi, Ookawa, Usuda, & Hirasawa, 2011;
Zhang et al., 2021). In rice specifically, higher stomatal conductance has been associated with
greater rates of leaf photosynthesis (Kusumi et al., 2017; Ohsumi, Kanemura, Homma, Horie, &
Shiraiwa, 2007b; Zhang et al., 2021). Efforts to enhance stomatal conductance have led to
increased leaf photosynthesis and biomass accumulation in a range of C3 plants (Kusumi et al.,
2017; Papanatsiou et al., 2019; Wang et al., 2014; Zhang et al., 2021). Thus, opportunity exists to
further fine-tune stomatal density reductions to maintain wild-type levels of carbon assimilation
and stomatal conductance while enhancing water conservation and drought resilience.

Here, we report further characterization of the rice STOMAGEN duplicate gene, subsequently
referred to as EPFL10, in its relationship to stomatal development. Furthermore, we explore the
effect of the reductions in stomatal density in *stomagen* and *epfl10* mutants on stomatal
conductance, carbon assimilation, water conservation, and yield in varying water regimes.

Methods

Plant Growth conditions:

Rice cultivar Nipponbare (Oryza sativa ssp. Japonica) seeds were germinated and grown for eight days in a petri dish with 20mL of water in a Conviron growth chamber at 28°C for day-length periods of 16hours in 100 μmol photons $m^{-2}s^{-1}$ of light and 80% relative humidity. Seedlings were transferred to a soil mixture comprised of equal parts turface (https://www.turface.com/products/infield-conditioners/mvp) and sunshine mix #4 (http://www.sungro.com/professional-products/fafard/).

Germinated seedlings used for stomatal phenotyping and growth chamber physiological assays were transferred to 10.16 cm, 0.75 L McConkey tech square pots and placed in growth chambers 28°C for day-length periods of 16 hours in 400 μmol photons $m^{-2}s^{-1}$ of light and 80% relative humidity.

Plants designated for yield trials, greenhouse physiological assays, and stomatal aperture measurements were moved to the greenhouse with temperature setpoints of 27°C/22°C at ambient light conditions in February 2020 with daylengths of 12h in Kord 15.24 cm, 1.835L pots.

All plants were fertilized with 125mL of 1% w/v iron solution one-week post-transplant. 1000mL of 5% w/v JR Peter's Blue 20-20-20 fertilizer (https://www.jrpeters.com/) was added to each flat at 3- and 11-weeks post-germination. Well-watered plants were provided a constant supply of water by maintaining a flooded condition in the tray.

Yield and water regimes:

Yield in three watering regimes were tested: well-watered, vegetative drought, and reproductive drought. Yield trials in varying water regimes were conducted using methods adapted and modified slightly from Caine et. al 2019. Well-watered flats were kept flooded for the entirety of the growth period. Vegetative drought was imposed by removing all water from flats containing pots for 7 days starting on day 28 after germination and for 9 days at day 56 after germination. In reproductive drought, water was removed from flats for 4 days at day 98 when panicles were undergoing grain filling. All grain and aboveground biomass from well-watered, vegetative and reproductive drought plants were harvested after 167 days, 177 days, and 181 days, respectively. Biomass measurements were completed on samples dried at 60°C for three days prior to weighing.

Generation of edited lines:

Guides for targeting EPFL10 and STOMAGEN were selected to minimize off-targets effects and maximize on-target efficiency in the first exon of the coding region. Guide sequences were selected using CRISPR-P 2.0 (http://crispr.hzau.edu.cn/CRISPR2/). Forward and reverse strand

guide sequence oligonucleotides with relevant sticky ends amenable for Golden Gate cloning
were ordered from IDT (IDTdna.com). Equal volumes of 10mM primers were annealed at room
temperature. Golden Gate cloning was used to insert guides into the PeGM entry clone
containing the tracrRNA and U3 promoter. LR clonase reactions were used to insert entry clone
into destination vectors for biolistic transformation and *Agrobacterium*-mediated transformation.
Plasmid maps are provided in the supporting information. *epfl10* lines were produced via
Agrobacterium-mediated transformation and *stomagen* lines via biolistic transformation.

Plant material and culture of explants

Mature seeds of rice (*Oryza sativa* L. japonica cv. Nipponbare) were de-hulled, and surface-
sterilized for 20 min in 20% (v/v) commercial bleach (5.25% sodium hypochlorite) plus a drop
of Tween 20. Three washes in sterile water were used to remove residual bleach from seeds. De-
hulled seeds were placed on callus induction medium (CIM) medium [N6 salts and vitamins
(Chu et al., 1975), 30 g/L maltose, 0.1 g/L myo-inositol, 0.3 g/L casein enzymatic hydrolysate,
0.5 g/L L-proline, 0.5 g/L L-glutamine, 2.5 mg/L 2,4-D, 0.2 mg/L BAP, 5 mM $CuSO_4$, 3.5 g/L
Phytagel, pH 5.8] and incubated in the dark at 28 °C to initiate callus induction. Six- to 8-week-
old embryogenic calli were used as targets for transformation.

Agrobacterium-mediated transformation

Embryogenic calli were dried for 30 min prior to incubation with an *Agrobacterium tumefaciens*
EHA105 suspension (OD_{600nm} = 0.1) carrying a binary vector of interest, OsEPFL10. After a 30
min incubation, the *Agrobacterium* suspension was removed. Calli were then placed on sterile
filter paper, transferred to co-cultivation medium [N6 salts and vitamins, 30 g/L maltose, 10 g/L
glucose, 0.1 g/L myo-inositol, 0.3 g/L casein enzymatic hydrolysate, 0.5 g/L L-proline, 0.5 g/L
L-glutamine, 2 mg/L 2,4-D, 0.5 mg/L thiamine, 100 mM acetosyringone, 3.5 g/L Phytagel, pH
5.2] and incubated in the dark at 21°C for 3 days. After co-cultivation, calli were transferred to
resting medium [N6 salts and vitamins, 30 g/L maltose, 0.1 g/L myo-inositol, 0.3 g/L casein
enzymatic hydrolysate, 0.5 g/L L-proline, 0.5 g/L L-glutamine, 2 mg/L 2,4-D, 0.5 mg/L
thiamine, 100 mg/L timentin, 3.5 g/L Phytagel, pH 5.8] and incubated in the dark at 28°C for 7
days. Calli were then transferred to selection medium (CIM plus 250 mg/L cefotaxime and 50
mg/L hygromycin B) and allowed to proliferate in the dark at 28°C for 14 days. Well-
proliferating tissues were transferred to CIM containing 75 mg/l hygromycin B. The remaining
tissues were subcultured at 3- to 4- week intervals on fresh selection medium. When a sufficient
amount (about 1.5 cm in diameter) of the putatively transformed tissues was obtained, they were
transferred to regeneration medium [MS salts and vitamins (Murashige & Skoog, 1962), 30 g/L
sucrose, 30 g/L sorbitol, 0.5 mg/L NAA, 1 mg/L BAP, 150 mg/L cefotaxime) containing 40
mg/L hygromycin B and incubated at 26 °C, 16-hr light, 90 μmol photons m^{-2} s^{-1}. When
regenerated plantlets reached at least 1 cm in height, they were transferred to rooting medium
(MS salts and vitamins, 20 g/L sucrose, 1 g/L myo-inositol, 150 mg/L cefotaxime) containing 20
mg/L hygromycin B and incubated at 26 °C under conditions of 16-hr light (150 μmol photons
m^{-2} s^{-1}) and 8-h dark until roots were established and leaves touched the Phytatray lid. Plantlets
were then transferred to soil.

Biolistic-mediated transformation

Embryogenic callus tissue pieces (3–4 mm) were transferred for osmotic pretreatment to CIM
medium containing mannitol and sorbitol (0.2 M each). Four hours after treatment with
osmoticum, tissues were bombarded as previously described (Cho et al., 2004) with
modifications. 7.5 ml of gold particles (0.6 μm), coated with 5 mg of
pU3Stomagen0.4:OsUbiCas9:HPT were divided equally among 10 macro-carriers and used for
bombardment with a Bio-Rad PDS-1000 He biolistic device (Bio-Rad, Hercules, Calif.) at 650
psi. Sixteen to 18 hr after bombardment, tissues were placed on osmotic-free CIM and incubated
at 28°C under dim light (10-30 μmol photons m^{-2} s^{-1}, 16-hr light). After 7 days, tissues were
transferred to selection medium (CIM containing 50 mg/l hygromycin B) and maintained/grown
using the same procedure as described above, but timentin or cefotaxime were not supplemented
in the media.

Validation of edits

T$_0$ plants genotypes at targeted loci were evaluated using PCR to amplify the region of interest
using primers listed in Supporting Table 3. PCR products were Sanger sequenced. Sequence data
was analyzed using the Synthego ICE tool (https://ice.synthego.com/#/) to detect alleles present.
Only lines with homozygous frame-shift mutations were retained for downstream experiments.
Plants from the second generation after transformation were used for experimental data
collection generation to account for somaclonal variation, which may have accumulated during
tissue culture (Bairu, Aremu, & Van Staden, 2010; Wei et al., 2016)

Phenotyping stomatal density, size, and aperture:

Stomatal densities were recorded from epidermal impressions of leaves using nail polish peels.
Stomatal densities of eight biological replicates of each leaf were measured. Impressions were
taken from the widest section of fully expanded leaves. Images were taken using a Leica
DM5000 B epifluorescent microscope at 10x magnification. Number of stomata in a single
stomatal band were counted and the area of each band was measured (Huang et al., 2019)
Stomatal densities were calculated by dividing stomatal counts by stomal band area (mm^2).

Epidermal peels of 21-day old plants were produced using a razorblade on the adaxial leaf to
remove tissues above the abaxial epidermal layer. Images of individual stomata at 100x
magnification were captured. Guard cell length was measured using ImageJ. 35 individual
stomata from five biological replicates of each genotype were measured.

Stomatal aperture measurements were generated using epidermal peels of flag leaves from 85-
day old plants. Leaves were harvested at 1:00pm and peels were generated immediately.
Epidermal peels were then fixed by submerging in 4% formaldehyde for 30s using a method
adapted by Eisele et al. (Eisele, Fäßler, Bürgel, & Chaban, 2016). Images of 20 individual
stomata from six biological replicates of each genotype were measured.

Quantifying EPFL10 and STOMAGEN transcript abundance:

Total RNA was extracted from seedlings eight days after germination, from leaf base and fully
expanded leaf of 21-day-old leaves using the Qiagen Total RNAeasy Plant Kit. RNA quality was

validated on an agarose gel prior to reverse transcription using the QuantiTect™ reverse
transcription kit to generate first-strand cDNA. Quantitative reverse transcription PCR was
performed using FAST SYBR on Applied Biosystem's QuantStudio 3 thermocycler. Relative
expression levels were calculated by normalizing to the rice UBQ5 housekeeping gene
(LOC_Os01g22490) (Jain, Vergish, & Khurana, 2018). Primers used for qPCR listed in
Supporting Table 3. Relative log fold expression was calculated using the $2^{-\Delta\Delta CT}$ method using
STOMAGEN in adult leaves as the control group.

Determining methylation profile of genes of interest:

252 Methylation profiles of rice genes of interest were viewed using the Plant Methylation Database,
253 (https://epigenome.genetics.uga.edu/PlantMethylome/) (Niederhuth et al., 2016). Snapshots of
254 CHH and CHG methylation 1.5 kb upstream of the start codon and 1.5 downstream of the stop
255 codon were taken.

Evolutionary analysis:

We collected complete single orthologs from the species used in this study (Table S1), using
BUSCO v4.0.6 (Seppey, Manni, & Zdobnov, 2019) and the viridiplantae_odb10 database. 82
orthologous groups present in at least 23 species were individually selected with MAFFT v7.487
(--maxiterate 1000 --globalpair) (Katoh & Standley, 2013). All multiple sequence alignments
were concatenated, trimmed with TrimAl v1.4.rev15 (-gt 0.2)(Capella-Gutiérrez, Silla-Martínez,
& Gabaldón, 2009) and then used to infer a species tree with FastTree v2.1.10 (Price, Dehal, &
Arkin, 2010). We determined the copy number variations of the STOMAGEN family by
searching for the stomagen domain (PF16851) from the protein annotation sets with hmmsearch
v3.3 (Eddy, 2011; Mistry et al., 2021) or from the genomes with exonerate v2.2.0 (Slater &
Birney, 2005) if genome annotations are absent. To understand the sequence variations of
STOMAGEN and EPFL10 orthologs at the species and family level, we collected non-redundant
Oryza or Poaceae species that have single copies of STOMAGEN or EPFL10 (Table S2). The
stomagen domain of STOMAGEN or EPFL10 orthologs was aligned with MAFFT, and we used
the filtered alignment to compute normalized Shannon's entropy, $-\sum_{i=1}^{20} p_i \log_2 p_i / \log_2 20$ where
p_i is the probability of observing ith amino acids among the twenty in the given position of the
alignment. Gaps were ignored.

Photosynthesis and stomatal conductance assays:

Physiological assays in Figures 3a, 3c were conducted on full expanded leaf 5 of 21-day old
plants. Stomatal conductance and CO_2 assimilation data for Fig 3c was captured using an infrared
gas analyzer (LI6400XT, LI-COR, Lincoln, NE, USA) with chamber conditions set to: light
intensity 1000 µmol photons m^{-2}s^{-1} (90% red light, 10% blue light); leaf temperature 27°C; flow
rate 500 µmol s^{-1}; relative humidity 40%; and CO_2 concentration of sample 400µmol mol^{-1}.

Light response curves in Figure 3b and 3d were generated using a LI6800 infrared gas analyzer
(LI-COR, Lincoln, NE, USA) with chamber conditions set to: leaf temperature 25°C; flow rate

500 µmol s^{-1}; water vapor pressure deficit 1.8 kPA; and CO_2 concentration of sample 400µmol
mol^{-1}. Light intensity was first increased to 2000 µmol photons m^{-2}s^{-1}, and with steady-state
waiting times of 5 to 10 minutes, subsequently decreased to 1500, 12000, 1000, 750, 500, 300,
200, 100, and 50 µmol photons m^{-2}s^{-1} light. Light was composed of at least 90% red light and at
maximum 40 µmol photons m^{-2}s^{-1} blue light to match equipment specifications. Measurements
were taken on fully expanded fifth leaves of 32-day-old plants grown in the greenhouse.
Measurements for Figure S1 were taken on full expanded leaf 5 of 28-day-old plants grown in
well-watered and vegetative drought conditions. All data collected were adjusted according to
leaf area within the gas exchange chamber. Intrinsic water use efficiency (iWUE) in Figure S3
was calculated by dividing photosynthesis by stomatal conductance for each biological replicate.
Specific stomatal conductance in Fig 3f was calculated by dividing stomatal conductance by the
average number of stomata within the probe area.

Thermal imaging:

Thermal images were captured using a FLIR E8-XT Infrared Camera (FLIR-DIRECT,
Wilmington, NC, USA). Images of well-watered and vegetative drought plants were taken 65
days after germination on the last day of the vegetative drought treatment. Images of
reproductive drought plants were captured 102 days post-germination on the last day of the
reproductive drought treatment. Images were captured between 1:00pm and 2:00pm to capture
the effects of transpiration-mediated cooling during the hottest part of the days. Images were
processed using FLIR Thermal Studio. Leaf temperatures of 4-6 leaves per biological replicate
were quantified.

Water loss

Two plants of identical genotype were placed in a 10"x10" flat covered with aluminum foil to
decrease evaporation from soil. Non-plant evaporation was estimated by measuring daily water
loss from covered flats containing pots without plants. Daily water loss of each flat was
calculated by taking the difference of flats with plants and without. Eight replicate flats for each
genotype were measured daily from 70-77 days after germination.

Graphs and statistics:

All graphs were produced using the ggplot2 package in R studio (Wickham, 2017). All statistics
were calculated in R-studio using post-hoc tests for significance between groups.

Results

Duplication of STOMAGEN in multiple plant families

The duplication of STOMAGEN(EPFL9;LOC_Os01g68598) in the Poaceae family was
previously reported (Hepworth et al., 2018) and is in agreement with our expanded gene tree in
eudicots and monocots (Fig. 1a) . Further resolution of the gene tree suggested that the orthologs

of STOMAGEN and its duplicate, EPFL10 (LOC_Os08g41360 hereafter referred to as EPFL10),
may have evolved differently, given the branch lengths of the two orthologous groups (Fig. 1b).
Furhter phylogenetic investigation of the duplication of STOMAGEN among angiosperms
revealed an additional putative family-level STOMAGEN duplication in the Asteraceae (Fig.
1c).

Peptide variation between STOMAGEN and EPFL10

Comparisons of the active 45 amino acid C-terminal sequences of STOMAGEN and EPFL10
indicated that some sequence divergence exists. (Fig. 1d). Sequence conservation of
STOMAGEN orthologs is much greater at the genus and family level relative to EPFL10
orthologs. Paralogous sequence variation of STOMAGEN and EPFL10 mapped onto the
TMM/ERL1, EPF1 complex highlighted the orientation of dissimilar and similar substitutions
within the peptide-receptor complex (Fig. 1e). The two dissimilar amino acids present in the
random coil in the beginning of the STOMAGEN domain are near the interface with TMM and
ERL1. The substitutions concentrated on the first beta sheet were all mapped to the residues near
ERL1, potentially altering binding capacities to the receptor.

Varied expression of EPFL10 and STOMAGEN

334   STOMAGEN mRNA abundance greatly exceeds EPFL10 expression in leaf base tissues where
335   stomatal development occurs and STOMAGEN and EPFL10 expression is greatest (Fig. 2a).

Stomatal density and morphology in knockout lines

CRISPR/Cas9-mediated knockout of STOMAGEN and EPFL10 was achieved by targeting
guides to the first exon of each gene to disrupt the open reading frame. Two unique homozygous
knockout alleles were generated in EPFL10 and in STOMAGEN in the T_0 generation using a
single guide sequence adjacent to a PAM motif (designated in green and purple, respectively in
Fig. 2b-c). *epfl10* exhibited reductions in stomatal densities which represented 80% of wild-type
densities, whereas *stomagen* possessed only 25% of wild-type densities in the fifth fully
expanded adult leaf and flag leaf, respectively (Fig. 2d-e). Stomatal size was measured to
determine if there was a relationship between stomatal density reductions and size increases in
the Nipponbare background. Consistent with previous reports, the guard cell length of stomata in
stomagen were longer relative to *epfl10* and wild type (Fig. 2f) (Caine et al., 2019; Mohammed
et al., 2019).

Stomatal conductance, carbon assimilation, and stomatal aperture

Infrared gas exchange analysis of reduced density lines was undertaken to determine if lines with
stomatal modifications altered carbon assimilation (A_n, µmol CO_2 $m^{-2}s^{-1}$) or stomatal
conductance (g_s, mol H_2O $m^{-2}s^{-1}$). At ambient CO_2 (400ppm) and saturating light (1000 µmol
photons $m^{-2}s^{-1}$), *stomagen* steady-state A_n and g_s were lower relative to the wild type grown in
growth chambers (Fig. 3a,c) . Interestingly, wild type and *epfl10* displayed similar steady-state
values (Fig. 3a-d). Likewise, *epfl10* maintained wild type levels of A_n and g_s whereas *stomagen*
did not in light response curves carried out on an independent cohort of plants in the greenhouse

(Fig. 3b,d). Similar reductions of A_n and g_s in *stomagen* lines were recapitulated in greenhouse
measurements at 1000 μmol photons $m^{-2}s^{-1}$ (Fig. S1). However, *stomagen* did not exhibit
lowered levels of carbon assimilation at 1000 μmol photons $m^{-2}s^{-1}$ after vegetative drought stress
despite still having reduced levels of stomatal conductance (Fig. S1c-d). Measurements of
stomatal apertures on the abaxial side of flag leaves indicated that *epfl10* lines maintained a
larger stomatal aperture than the wild type, and *stomagen* lines exhibited an even greater aperture
(Fig. 3e). Thus, *epfl10* and *stomagen* maintained even greater levels of stomatal conductance per
individual stoma mediated by a larger aperture (Fig. 3f).

Leaf temperature and water conservation

Thermal imaging was used to assess evaporative cooling in altered stomatal lines. In well-
watered conditions *stomagen* lines were warmer on average than wild type, whereas *epfl10* leaf
temperatures were intermediate (Fig. 4a). No difference in leaf temperature was detected during
vegetative drought (Fig. 4b). During reproductive drought, *eplf10* was marginally cooler than
stomagen (Fig. 4c). To test the impact of reduced stomatal densities on water conservation, daily
water loss was measured over the course of a week beginning with 70 days after germination.
Stomagen and *epfl10* both conserved greater volumes of water in a week by 48mL and 83mL,
respectively (Fig. 4d).

Yield Trials

To assess the impacts of stomatal modifications on crop performance, yield trials were conducted
using three watering regimes in the greenhouse. In well-watered, vegetative drought, and
reproductive drought conditions, there was no discernible difference in grain or biomass yield
among genotypes (Fig. S2).

Discussion:

In current climate conditions, drought is the most severe and widespread environmental stressor
in South and Southeast Asia ("Climate change - ready rice | International Rice Research
Institute," n.d.). Application of gene editing in crops for climate change could serve as a potent
mechanism for realizing actual technology transfer to growers (Jenkins, Dobert, Atanassova, &
Pavely, 2021; Karavolias, Horner, Abugu, & Evanega, 2021). Previous manipulation of
epidermal patterning factors generated rice with improved drought tolerance and maintenance of
yields in greenhouse conditions, albeit with reductions in A_n and g_s (Caine et al., 2019; Dunn et
al., 2019; Hughes et al., 2017). In this study, novel characterization of the rice epidermal
patterning factor EPFL10 in its native system enabled the development of climate change
adapted rice in this study.

In rice, we hypothesize that EPFL10 may bind to the same receptor partners as STOMAGEN
with a reduced affinity thereby limiting its capacity in positive regulation of stomatal
development. Substituted residues in EPFL10 relative to STOMAGEN may underlie the
phenotypic variation observed in previous overexpression data. As the random coil in the

beginning of the STOMAGEN domain appears to contact both ERL1 and TMM in the complex
structure of EPF1, the two substituted amino acids mapped to this region may be important for
alternate binding affinity of EPFL10 (Figure 1e). Additional substituted amino acids may also
play a role in lowered relative stability of EPFL10 in the TMM/ERL receptor complex. Lower
expression levels of EPFL10 relative to STOMAGEN in leaf base and seedling where new
stomatal complexes are forming in combination with a peptide-level difference likely account for
the modest reductions of stomatal density in EPFL10 knockouts relative to STOMAGEN (Figure
2a, Figure 1). Greater CHG and CHH repressive methylation marks vicinal to the EPFL10 gene
relative to STOMAGEN may account for differences in expression and function of these
duplicated genes (Figure S3) (Raju, Ledford, & Niederhuth, 2021). Overall, higher sequence
conservation across various species and greater expression levels in relevant tissues signify
fundamental roles of STOMAGEN in development. EPFL10 may be selectively utilized to fine-
tune stomatal densities, and relatively divergent sequence evolution may reflect species'
adaptations or relaxed purifying selection.

Stomatal density reductions mediated by single gene knockouts of EPFL10 and STOMAGEN
were equivalently capable of conserving water. Fewer stomata offer fewer sites for water loss to
occur. It is noteworthy that *stomagen* lines conserved water to a similar extent as *epfl10* despite
having much greater reductions in stomatal density. Our data indicates that fewer stomata with
larger apertures offer comparable water conservativity properties as lines possessing greater
number of stomata with smaller apertures. It is still unclear how more severe and/or field-
relevant drought stresses may affect water loss and assimilation when specific stomatal
conductance is limiting. Yet, stomatal reductions broadly offer water-savings relative to wild
type.

Despite a promising water conservation phenotype, *stomagen* lines were unable to maintain
wild-type levels of A_n and g_s whereas *epfl10* maintained levels equivalent to wild type across all
light and growth conditions. Stomatal conductance and photosynthesis of *stomagen* but not
epfl10 were lower at all light intensities greater than 1000 μmol photons $m^{-2}s^{-1}$ relative to wild-
type. Fluctuating light in field conditions tends to be present at photosynthetic flux densities
greater than 1000 μmol $m^{-2}s^{-1}$ in mid-canopy for the majority of the day (Slattery, Walker,
Weber, & Ort, 2018). No differences in grain yield were measured among genotypes despite
marked differences in steady state g_s and A_n of *stomagen* lines. A previous greenhouse study in
which g_s and A_n were lowered by EPF1 overexpression did not result in lowered yields (Caine et
al., 2019). However, most other literature suggests that high levels of leaf photosynthesis and
stomatal conductance are directly linked to higher yields (Kusumi et al., 2017; Ohsumi et al.,
2007a; Richards, 2000; Roche, 2015; Taylaran et al., 2011; Zhang et al., 2021). Additional
studies of yield in dynamic and substantially more stressful field conditions are thus necessary to
reconcile discrepancies with previous reports and more robustly assay the impacts of stomatal
density reductions on yield.

Larger stomatal apertures were measured in *stomagen* and *epfl10* lines relative to wild type
(Figure 3e). Pore area adjustments in *stomagen* lines were unable to physiologically compensate
for large reductions in stomatal densities, unlike *epfl10* lines, which maintained wild-type levels
of A_n and g_s (Figure 3a-d). The theoretical maximum stomatal conductance of grass stomata
greatly exceeds the measured stomatal conductance (Caine et al., 2019; Faralli, Matthews, &

Lawson, 2019). *epfl10* maintained high levels of specific stomatal conductance concurrent with overall A_n and g_s(Figure 3f, Figure 3a-3d, Figure 4d). Enhancing specific stomatal conductance in a reduced density background may thus provide a promising mechanism for maintaining photosynthetic capacities simultaneous with water-use efficiency (C. R. Buckley, Caine, & Gray, 2019; Faralli et al., 2019).

In accordance with anticipated global warming caused by climate change, maintained or improved thermoregulation will be vital for crop agronomic performance (Jagadish et al., 2015). *epfl10* maintained wild-type thermoregulation in all watering regimes whereas *stomagen* lines were warmer in well-watered and reproductive drought conditions (Figure 4a, 4c). A previous report indicated a similar trend wherein lines with reductions in stomatal densities were able to remain cooler during both vegetative and reproductive droughts (Caine et al., 2019). Greater water retention prior to drought may facilitate enhanced thermoregulatory capacities during reproductive drought despite having fewer stomata.

Characterization of developmental genes in crop plants can expand upon available developmental frameworks and provide tools for developing the next generation of crops (Hughes et al., 2017; Raissig et al., 2017). The genetics underlying the novel components of grass stomatal development can only be fully resolved by expanding developmental genetics explorations beyond the paradigm of *A. thaliana* investigations. In this case, explorations of the role of EPFL10 unraveled its native function as a weak positive regulator of stomatal development in rice plants. We report here the demonstration of stomatal density reductions with no concomitant reductions in stomatal conductance or carbon assimilation. *epfl10* lines maintained wild-type physiological capacities of stomatal conductance, carbon assimilation, thermoregulation, and yield while also conserving more water than wild-type. These attributes could contribute to improved climate resilience in current and future conditions where water is limiting, and temperatures are increased. Field-based investigations of *epfl10* and *stomagen* will further resolve the agronomic utility of these edited rice lines.

Author Contributions:

NGK developed project idea and coordinated research efforts. NGK constructed gene-editing vectors with assistance from DD. MT and JT with oversight from MJC produced *epfl10* and *stomagen* lines, respectively. KS generated dendrogram, gene tree and peptide models. NGK phenotyped stomatal densities, stomatal aperture, stomatal size, water loss, and yield. GAG repeated stomatal density phenotyping independently to confirm results. NGK and DP conducted gs and A assays. DP captured thermal images. NGK drafted the manuscript with edits from DP and KKN. KKN provided LICOR equipment, thermal imaging equipment, and technical expertise. BJS provided feedback on experimental design and the manuscript as well as the facilities to carry out experiments.

Funding statement: Funding provided by Open Philanthropy, Foundation for Food and Agriculture Research, Innovative Genomics Institute and the National Science Foundation

476   Graduate Research Fellowship Program. KKN is an investigator of the Howard Hughes Medical
477   Institute.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 1. The evolution of STOMAGEN and EPFL10

(a) A dendrogram of angiosperm species and copy number variations of STOMAGEN family

members. **(b-c)** Gene trees of the STOMAGEN family members in Poales and Asterales,

respectively. *Arabidopsis* was used as an outgroup. **(d)** Genus and family-level sequence

conservation of STOMAGEN and EPFL10 orthologs. The sequence conservation is calculated

by 1 - normalized Shannon's entropy, with 1 being perfect conservation and 0 being completely

variable. The secondary structure annotation originates from the solved structure of

STOMAGEN (PDB: 2LIY) (Ohki et al., 2011). The colored residues highlight variable positions

between rice STOMAGEN and EPFL10. Substitutions to similar and dissimilar amino acids

based on BLOSUM62 are indicated with blue and red. **(e)** The variable positions between

STOMAGEN and EPFL10 mapped to the solved complex structure of EPF1, ERL1 and TMM

(PDB: 5XJO)(Lin et al., 2017). The variable positions with similar or dissimilar substitutions

follow (d).

Figure 2. EPFL10 is a weak positive regulator of stomatal development in Nipponbare (*Oryza*

sativa spp. Japonica). (a) qRT-PCR determined expression levels of EPFL10 and STOMAGEN

in fully expanded leaf, leaf base, and seedling. Bars represent means and error bars represent

standard deviation from the mean. The asterisk indicates a significant difference between the

means ($P<0.05$). (b,c) Representative gene models of STOMAGEN (b) and EPFL10 (c) with

guide sequence in green and PAM sequence in blue. Homozygous edits generated in two mutant

lines by CRISPR/Cas9 are shown in red. (d,e) Stomatal density of *epfl10, stomagen,* and wild

type. d.) Stomatal density of the fifth fully expanded true leaf. Stomatal densities of 21-day old

plants were measured e.) Stomatal density of the flag leaf on the primary tiller during grain

filling. Flag leaves of 55-day old plants were measured. (f) Stomatal length of *epfl10, stomagen,*

and wild type. Graphs D-F are box-and-whisker plots where the center horizontal indicates the

median, upper and lower edges of the box are the upper and lower quartiles and whiskers extend

to the maximum and minimum values within 1.5 interquartile ranges. Outliers are represented by

black dots. Letters indicate a significant difference between means ($P<0.05$, one-way ANOVA

Duncan post-hoc test). (a) N=3,3, (d,e) N=9,9,9 (f)N=35,35,35.

Figure 3. Gas exchange measurements and stomatal pore area measurements in reduced stomatal

density backgrounds in Nipponbare (*Oryza sativa* spp. Japonica). (a,b) Gas exchange

measurements of carbon assimilation rate of *stomagen, epfl10,* and wild-type at 1000 µmol

photons m^{-2} s^{-1} in 21-day-old plants grown in growth chamber (a) and 32-day-old plants grown

in the greenhouse across a range of light intensities: 2000, 1500, 120, 1000, 750, 500, 300, 200,

100 µmol photons m^{-2} s^{-1} (b). (c,d) Gas exchange measurements of stomatal conductance in

stomagen, *epfl10*, and wild-type at 1000 $µmol^{-2}s^{-1}$ in 21-day-old plants grown in growth chamber

(c) and 32-day-old plants grown in the greenhouse across a range of light intensities 2000, 1500,

120, 1000, 750, 500, 300, 200, 100 µmol photons m^{-2} s^{-1} (d). (e) Representative images of

stomatal pore size variation from largest to smallest (left to right), top, and pore area

measurements of *stomagen, epfl10,* and wild-type, bottom. (f) Specific stomatal conductance of

epfl10 and *stomagen* lines. Specific conductance values calculated by dividing stomatal

conductance by the average stomatal density of the probe area of the respective genotype. In

graphs b, d, dots represent means and error bars are standard deviation from the mean. Graphs in

a, c, e, and f are box-and-whisker plots where the center horizontal indicates the median, upper

and lower edges of the box are the upper and lower quartiles and whiskers extend to the

maximum and minimum values within 1.5 interquartile ranges. Outliers are represented by black
dots. Letters indicate a significant difference between means ($P<0.05$, one-way ANOVA Duncan
post-hoc test). (a,c) N=8,8,8 (b,d) N=6,5,5 (e) N= 5,5,6.
697
Figure 4. Stomatal density reductions influence thermoregulation and water conservation in
Nipponbare (*Oryza sativa* spp. Japonica). (a-c) Leaf temperatures of *stomagen, epfl10*, and wild-
type, left, with representative thermal images of each genotype, right. (a) In well-watered, (b) In
vegetative drought, and (c) In reproductive drought. (d) Cumulative water loss of *stomagen,*
epfl10, and wild type from days 70-77 after germination. Graphs in A-C are box-and-whisker
plots where the center horizontal indicates the median, upper and lower edges of the box are the
upper and lower quartiles and whiskers extend to the maximum and minimum values within 1.5
interquartile ranges. Outliers are represented by black dots. In graph D dots represent means and
error bars are standard error from the mean. Letters indicate a significant difference between
means ($P<0.05$, one-way ANOVA Duncan post-hoc test). (a-d) N=8,8,8.

Works Cited:

Bairu, M. W., Aremu, A. O., & Van Staden, J. (2010). Somaclonal variation in plants: causes and detection methods. *Plant Growth Regulation 2010 63:2*, *63*(2), 147–173. https://doi.org/10.1007/S10725-010-9554-X

Bernier, J., Atlin, G. N., Serraj, R., Kumar, A., & Spaner, D. (2008). Breeding upland rice for drought resistance. *Journal of the Science of Food and Agriculture*, *88*(6), 927–939. https://doi.org/10.1002/JSFA.3153

Buckley, C. R., Caine, R. S., & Gray, J. E. (2019). Pores for thought: Can genetic manipulation of stomatal density protect future rice yields? *Frontiers in Plant Science*, *10*, 1783. https://doi.org/10.3389/FPLS.2019.01783

Buckley, T. N. (2005). The control of stomata by water balance. *New Phytologist*, *168*(2), 275–292. https://doi.org/10.1111/j.1469-8137.2005.01543.x

Caine, R. S., Yin, X., Sloan, J., Harrison, E. L., Mohammed, U., Fulton, T., … Gray, J. E. (2019). Rice with reduced stomatal density conserves water and has improved drought tolerance under future climate conditions. *New Phytologist*, *221*(1), 371–384. https://doi.org/10.1111/nph.15344

Capella-Gutiérrez, S., Silla-Martínez, J. M., & Gabaldón, T. (2009). trimAl: A tool for automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics*, *25*(15). https://doi.org/10.1093/bioinformatics/btp348

Cho, M. J., Yano, H., Okamoto, D., Kim, H. K., Jung, H. R., Newcomb, K., … Lemaux, P. G. (2004). Stable transformation of rice (Oryza sativa L.) via microprojectile bombardment of highly regenerative, green tissues derived from mature seed. *Plant Cell Reports*, *22*(7). https://doi.org/10.1007/s00299-003-0713-7

Chu, C. C., Want, C. C., Sun, C. S., Hsu, C., Yin, K. C., Chu, C. Y., & Bi, F. Y. (1975). Establishment of an efficient medium for anther culture of rice, through comparative experiments on the nitrogen sources. *Sci Sin*, *18*.

Climate change - ready rice | International Rice Research Institute. (n.d.). Retrieved December 9, 2021, from https://www.irri.org/climate-change-ready-rice

Dunn, J., Hunt, L., Afsharinafar, M., Meselmani, M. Al, Mitchell, A., Howells, R., … Gray, J. E. (2019). Reduced stomatal density in bread wheat leads to increased water-use efficiency. *Journal of Experimental Botany*, *70*(18), 4737–4748. https://doi.org/10.1093/JXB/ERZ248

Eddy, S. R. (2011). Accelerated profile HMM searches. *PLoS Computational Biology*, *7*(10). https://doi.org/10.1371/journal.pcbi.1002195

Eisele, J. F., Fäßler, F., Bürgel, P. F., & Chaban, C. (2016). A Rapid and Simple Method for Microscopy-Based Stomata Analyses. *PLOS ONE*, *11*(10), e0164576. https://doi.org/10.1371/JOURNAL.PONE.0164576

Endo, H., & Torii, K. U. (2019). Stomatal Development and Perspectives toward Agricultural Improvement. *Cold Spring Harbor Perspectives in Biology*, *11*(5), a034660. https://doi.org/10.1101/cshperspect.a034660

Faralli, M., Matthews, J., & Lawson, T. (2019). Exploiting natural variation and genetic manipulation of stomatal conductance for crop improvement. *Current Opinion in Plant*

Biology. https://doi.org/10.1016/j.pbi.2019.01.003

Fischer, R. A., Rees, D., Sayre, K. D., Lu, Z. M., Condon, A. G., & Larque Saavedra, A. (1998).
Wheat yield progress associated with higher stomatal conductance and photosynthetic rate,
and cooler canopies. *Crop Science*, *38*(6), 1467–1475.
https://doi.org/10.2135/cropsci1998.0011183X003800060011x

Food and Agriculture Organisation (FAO). (2020). FAOSTAT: Statistical database.

Franks, P. J., W. Doheny-Adams, T., Britton-Harper, Z. J., & Gray, J. E. (2015). Increasing
water-use efficiency directly through genetic manipulation of stomatal density. *New
Phytologist*, *207*(1), 188–195. https://doi.org/10.1111/nph.13347

Hara, K., Kajita, R., Torii, K. U., Bergmann, D. C., & Kakimoto, T. (2007). The secretory
peptide gene EPF1 enforces the stomatal one-cell-spacing rule.
https://doi.org/10.1101/gad.1550707

Hara, K., Yokoo, T., Kajita, R., Onishi, T., Yahata, S., Peterson, K. M., … Kakimoto, T. (2009).
Epidermal Cell Density is Autoregulated via a Secretory Peptide, EPIDERMAL
PATTERNING FACTOR 2 in Arabidopsis Leaves. *Plant and Cell Physiology*, *50*(6),
1019–1031. https://doi.org/10.1093/PCP/PCP068

Hepworth, C., Caine, R. S., Harrison, E. L., Sloan, J., & Gray, J. E. (2018). Stomatal
development: focusing on the grasses. *Current Opinion in Plant Biology*, *41*(Stage 5), 1–7.
https://doi.org/10.1016/j.pbi.2017.07.009

Hepworth, C., Doheny-Adams, T., Hunt, L., Cameron, D. D., & Gray, J. E. (2015). Manipulating
stomatal density enhances drought tolerance without deleterious effect on nutrient uptake.
New Phytologist, *208*(2), 336–341. https://doi.org/10.1111/NPH.13598

Huang, L., Chen, L., Wang, L., Yang, Y., Rao, Y., Ren, D., … Zeng, D. (2019). A Nck-
associated protein 1-like protein affects drought sensitivity by its involvement in leaf
epidermal development and stomatal closure in rice. *The Plant Journal*, tpj.14288.
https://doi.org/10.1111/tpj.14288

Hughes, J., Hepworth, C., Dutton, C., Dunn, J. A., Hunt, L., Stephens, J., … Gray, J. E. (2017).
Reducing Stomatal Density in Barley Improves Drought Tolerance without Impacting on
Yield. *Plant Physiology*, *174*(2), 776–787. https://doi.org/10.1104/pp.16.01844

Hunt, L., Bailey, K. J., & Gray, J. E. (2010). The signalling peptide EPFL9 is a positive regulator
of stomatal development. *New Phytologist*, *186*(3), 609–614.
https://doi.org/10.1111/J.1469-8137.2010.03200.X

Hunt, L., & Gray, J. E. (2009). The Signaling Peptide EPF2 Controls Asymmetric Cell Divisions
during Stomatal Development. *Current Biology*, *19*(10), 864–869.
https://doi.org/10.1016/J.CUB.2009.03.069

IPCC, Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S. L., Péan, C., … B., Z. (2021).
*Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the
Sixth Assessment Report of the Intergovernmental Panel on Climate Change*. *Cambridge
University Press*.

Jagadish, S. V. K., Murty, M. V. R., & Quick, W. P. (2015). Rice responses to rising
temperatures – challenges, perspectives and future directions. *Plant, Cell & Environment*,
38(9), 1686–1698. https://doi.org/10.1111/PCE.12430

Jägermeyr, J., Müller, C., Ruane, A. C., Elliott, J., Balkovic, J., Castillo, O., … Rosenzweig, C.
(2021). Climate impacts on global agriculture emerge earlier in new generation of climate
and crop models. *Nature Food 2021*, 1–13. https://doi.org/10.1038/s43016-021-00400-y

Jain, N., Vergish, S., & Khurana, J. P. (2018). Validation of house-keeping genes for
normalization of gene expression data during diurnal/circadian studies in rice by RT-qPCR.
Scientific Reports, *8*(1), 3203. https://doi.org/10.1038/S41598-018-21374-1

Jangra, R., Brunetti, S. C., Wang, X., Kaushik, P., Gulick, P. J., Foroud, N. A., … Lee, J. S.
(2021). Duplicated antagonistic EPF peptides optimize grass stomatal initiation.
https://doi.org/10.1242/dev.199780

Jenkins, D., Dobert, R., Atanassova, A., & Pavely, C. (2021). Impacts of the regulatory
environment for gene editing on delivering beneficial products. *In Vitro Cellular &*
Developmental Biology, 1. https://doi.org/10.1007/S11627-021-10201-4

Karavolias, N. G., Horner, W., Abugu, M. N., & Evanega, S. N. (2021). Application of Gene
Editing for Climate Change in Agriculture. *Frontiers in Sustainable Food Systems*, *0*, 296.
https://doi.org/10.3389/FSUFS.2021.685801

Katoh, K., & Standley, D. M. (2013). MAFFT multiple sequence alignment software version 7:
Improvements in performance and usability. *Molecular Biology and Evolution*, *30*(4).
https://doi.org/10.1093/molbev/mst010

Khush, G. S. (1997). Origin, dispersal, cultivation and variation of rice. *Plant Molecular*
Biology, *35*(1–2). https://doi.org/10.1007/978-94-011-5794-0_3

Kusumi, K., Hashimura, A., Yamamoto, Y., Negi, J., & Iba, K. (2017). Contribution of the S-
type anion channel SLAC1 to stomatal control and its dependence on developmental stage
in rice. *Plant and Cell Physiology*, *58*(12), 2085–2094. https://doi.org/10.1093/pcp/pcx142

Lafitte, H. R., Ismail, A., & Bennett, J. (2004). Abiotic stress tolerance in rice for Asia : progress
and the future. Crop Science.

Lee, J. S., Hnilova, M., Maes, M., Lin, Y. C. L., Putarjunan, A., Han, S. K., … Torii, K. U.
(2015). Competitive binding of antagonistic peptides fine-tunes stomatal patterning. *Nature*.
https://doi.org/10.1038/nature14561

Lin, G., Zhang, L., Han, Z., Yang, X., Liu, W., Li, E., … Chai, J. (2017). A receptor-like protein
acts as a specificity switch for the regulation of stomatal development. *Genes &*
Development, *31*(9), 927–938. https://doi.org/10.1101/GAD.297580.117

Liu, T., Ohashi-Ito, K., & Bergmann, D. C. (2009). Orthologs of Arabidopsis thaliana stomatal
bHLH genes and regulation of stomatal development in grasses. *Development*, *136*(13),
2265–2276. https://doi.org/10.1242/DEV.032938

Lu, J., He, J., Zhou, X., Zhong, J., Li, J., & Liang, Y. K. (2019). Homologous genes of epidermal
patterning factor regulate stomatal development in rice. *Journal of Plant Physiology*, *234–*
235(November 2018), 18–27. https://doi.org/10.1016/j.jplph.2019.01.010

Mistry, J., Chuguransky, S., Williams, L., Qureshi, M., Salazar, G. A., Sonnhammer, E. L. L., …
Bateman, A. (2021). Pfam: The protein families database in 2021. *Nucleic Acids Research*,
49(D1). https://doi.org/10.1093/nar/gkaa913

Mohammed, U., Caine, R. S., Atkinson, J. A., Harrison, E. L., Wells, D., Chater, C. C., …

Murchie, E. H. (2019). Rice plants overexpressing OsEPF1 show reduced stomatal density
and increased root cortical aerenchyma formation. *Scientific Reports*, *9*(1), 1–13.
https://doi.org/10.1038/s41598-019-41922-7

Murashige, T., & Skoog, F. (1962). A Revised Medium for Rapid Growth and Bio Assays with
Tobacco Tissue Cultures. *Physiologia Plantarum*, *15*(3). https://doi.org/10.1111/j.1399-
3054.1962.tb08052.x

Niederhuth, C. E., Bewick, A. J., Ji, L., Alabady, M. S., Kim, K. Do, Li, Q., … Schmitz, R. J.
(2016). Widespread natural variation of DNA methylation within angiosperms. *Genome*
Biology 2016 17:1, *17*(1), 1–19. https://doi.org/10.1186/S13059-016-1059-0

Ohki, S., Takeuchi, M., & Mori, M. (2011). The NMR structure of stomagen reveals the basis of
stomatal density regulation by plant peptide hormones. *Nature Communications*, *2*(1), 1–7.
https://doi.org/10.1038/ncomms1520

Ohsumi, A., Kanemura, T., Homma, K., Horie, T., & Shiraiwa, T. (2007a). Genotypic variation
of stomatal conductance in relation to stomatal density and length in rice (Oryza sativa L.).
Plant Production Science, *10*(3), 322–328. https://doi.org/10.1626/pps.10.322

Ohsumi, A., Kanemura, T., Homma, K., Horie, T., & Shiraiwa, T. (2007b). Genotypic variation
of stomatal conductance in relation to stomatal density and length in rice (Oryza sativa L.).
Plant Production Science, *10*(3), 322–328. https://doi.org/10.1626/pps.10.322

Pandey, S., Bhandari, H., Ding, S., Prapertchob, P., Sharan, R., Naik, D., … Sastri, A. (2007).
Coping with drought in rice farming in Asia: Insights from a cross-country comparative
study. In *Agricultural Economics* (Vol. 37). https://doi.org/10.1111/j.1574-
0862.2007.00246.x

Papanatsiou, M., Petersen, J., Henderson, L., Wang, Y., Christie, J. M., & Blatt, M. R. (2019).
Optogenetic manipulation of stomatal kinetics improves carbon assimilation, water use, and
growth. *Science (New York, N.Y.)*, *363*(6434), 1456–1459.
https://doi.org/10.1126/science.aaw0046

Price, M. N., Dehal, P. S., & Arkin, A. P. (2010). FastTree 2 – Approximately Maximum-
Likelihood Trees for Large Alignments. *PLOS ONE*, *5*(3), e9490.
https://doi.org/10.1371/JOURNAL.PONE.0009490

Raissig, M. T., Matos, J. L., Anleu Gil, M. X., Kornfeld, A., Bettadapur, A., Abrash, E., …
Bergmann, D. C. (2017). Mobile MUTE specifies subsidiary cells to build physiologically
improved grass stomata. *Science (New York, N.Y.)*, *355*(6330), 1215–1218.
https://doi.org/10.1126/science.aal3254

Raju, S. K. K., Ledford, S. M., & Niederhuth, C. E. (2021). DNA methylation signatures of
duplicate gene evolution in angiosperms. *BioRxiv*, 2020.08.31.275362.
https://doi.org/10.1101/2020.08.31.275362

Rang, Z. W., Jagadish, S. V. K., Zhou, Q. M., Craufurd, P. Q., & Heuer, S. (2011). Effect of high
temperature and water stress on pollen germination and spikelet fertility in rice.
Environmental and Experimental Botany, *70*(1), 58–65.
https://doi.org/10.1016/J.ENVEXPBOT.2010.08.009

Richards, R. A. (2000). Selectable traits to increase crop photosynthesis and yield of grain crops.
Journal of Experimental Botany, *51*(suppl_1), 447–458.

https://doi.org/10.1093/JEXBOT/51.SUPPL_1.447

Roche, D. (2015). Stomatal Conductance Is Essential for Higher Yield Potential of C $_3$ Crops.
Critical Reviews in Plant Sciences, *34*(4), 429–453.
https://doi.org/10.1080/07352689.2015.1023677

Seppey, M., Manni, M., & Zdobnov, E. M. (2019). BUSCO: Assessing genome assembly and
annotation completeness. In *Methods in Molecular Biology* (Vol. 1962).
https://doi.org/10.1007/978-1-4939-9173-0_14

Shah, F., Huang, J., Cui, K., Nie, L., Shah, T., Chen, C., & Wang, K. (2011). Impact of high-
temperature stress on rice plant and its traits related to tolerance. *The Journal of*
Agricultural Science, *149*(5), 545–556. https://doi.org/10.1017/S0021859611000360

Shimada, T., Sugano, S. S., & Hara-Nishimura, I. (2011). Positive and negative peptide signals
control stomatal density. *Cellular and Molecular Life Sciences*.
https://doi.org/10.1007/s00018-011-0685-7

Slater, G. S. C., & Birney, E. (2005). Automated generation of heuristics for biological sequence
comparison. *BMC Bioinformatics*, *6*. https://doi.org/10.1186/1471-2105-6-31

Slattery, R. A., Walker, B. J., Weber, A. P. M., & Ort, D. R. (2018). The Impacts of Fluctuating
Light on Crop Performance. *Plant Physiology*, *176*(2), 990–1003.
https://doi.org/10.1104/PP.17.01234

Stebbins, G. L., & Shah, S. S. (1960). Developmental studies of cell differentiation in the
epidermis of monocotyledons. II. Cytological features of stomatal development in the
Gramineae. *Developmental Biology*, *2*(6). https://doi.org/10.1016/0012-1606(60)90050-6

Sugano, S. S., Shimada, T., Imai, Y., Okawa, K., Tamai, A., Mori, M., & Hara-Nishimura, I.
(2010a). Stomagen positively regulates stomatal density in Arabidopsis. *Nature*, *463*(7278),
241–244. https://doi.org/10.1038/nature08682

Sugano, S. S., Shimada, T., Imai, Y., Okawa, K., Tamai, A., Mori, M., & Hara-Nishimura, I.
(2010b). Stomagen positively regulates stomatal density in Arabidopsis. *Nature*, *463*(7278),
241–244. https://doi.org/10.1038/nature08682

Taylaran, R. D., Adachi, S., Ookawa, T., Usuda, H., & Hirasawa, T. (2011). Hydraulic
conductance as well as nitrogen accumulation plays a role in the higher rate of leaf
photosynthesis of the most productive variety of rice in Japan. *Journal of Experimental*
Botany, *62*(11), 4067–4077. https://doi.org/10.1093/JXB/ERR126

Tiago D.G. Nunens, Dan Zhang, and M. T. R. (2019). *Form, development and function of grass*
stomata. https://doi.org/10.1111/tpj.14552

Tuong, T. P., & Bouman, B. A. M. (2009). Rice production in water-scarce environments. In
Water productivity in agriculture: limits and opportunities for improvement.
https://doi.org/10.1079/9780851996691.0053

Wang, Y., Noguchi, K., Ono, N., Inoue, S., Terashima, I., & Kinoshita, T. (2014).
Overexpression of plasma membrane H+-ATPase in guard cells promotes light-induced
stomatal opening and enhances plant growth. *Proceedings of the National Academy of*
Sciences, *111*(1), 533–538. https://doi.org/10.1073/PNAS.1305438111

Wei, F.-J., Kuang, L.-Y., Oung, H.-M., Cheng, S.-Y., Wu, H.-P., Huang, L.-T., … Hsing, Y.-I.

947  C. (2016). Somaclonal variation does not preclude the use of rice transformants for genetic
948   screening. *The Plant Journal*, *85*(5), 648–659. https://doi.org/10.1111/tpj.13132

949 Wickham, H. (2017). *ggplot2 - Elegant Graphics for Data Analysis | Hadley Wickham |*
950   *Springer*. *Springer Science & Business Media*.

951 Yin, X., Biswal, A. K., Dionora, J., Perdigon, K. M., Balahadia, C. P., Mazumdar, S., …
952   Bandyopadhyay, A. (2017). CRISPR-Cas9 and CRISPR-Cpf1 mediated targeting of a
953   stomatal developmental gene EPFL9 in rice, *36*, 745–757. https://doi.org/10.1007/s00299-
954   017-2118-z

955 Zhang, M., Wang, Y., Chen, X., Xu, F., Ding, M., Ye, W., … Zhu, Y. (2021). Plasma membrane
956   H+-ATPase overexpression increases rice yield via simultaneous enhancement of nutrient
957   uptake and photosynthesis. *Nature Communications*, *12*(1), 1–12.
958   https://doi.org/10.1038/s41467-021-20964-4

960
961
962