Expanded FLP toolbox for spatiotemporal protein degradation and transcriptomic
profiling in *C. elegans*

Adrián Fragoso-Luna[1,4], Raquel Romero-Bueno[1,4], Michael Eibl[1,4], Cristina Ayuso[1],
Celia Muñoz-Jiménez[1], Vladimir Benes[2], Ildefonso Cases[1], Peter Askjaer[3]

[1] Andalusian Centre for Developmental Biology, Consejo Superior de
Investigaciones Científicas (CSIC), Universidad Pablo de Olavide, Junta de
Andalucía, Carretera de Utrera, km 1, 41013 Sevilla, Spain
[2] EMBL GeneCore, Meyerhofstrasse 1, 69117 Heidelberg, Germany
[3] Corresponding author (pask@upo.es)
[4] These authors contributed equally

Keywords: DamID; FLP/FRT recombination; GFPdeg; RAPID; spatiotemporal gene
expression control; targeted protein degradation

20 Abstract

Control of gene expression in specific tissues and/or at certain stages of
development allows the study and manipulation of gene function with high precision.
Site-specific genome recombination by the Flippase (FLP) and Cre enzymes has
proven particularly relevant. Joint efforts of many research groups have led to the
creation of efficient FLP and Cre drivers to regulate gene expression in a variety of
tissues in *Caenorhabditis elegans*. Here, we extend this toolkit by the addition of FLP
lines that drive recombination specifically in distal tip cells, the somatic gonad,
coelomocytes and the epithelial P lineage. In some cases, recombination-mediated
gene knockouts do not completely deplete protein levels due to persistence of long-
lived proteins. To overcome this, we developed a spatiotemporally regulated
degradation system for GFP fusion proteins (GFPdeg) based on FLP-mediated
recombination. Using two stable nuclear pore proteins, MEL-28/ELYS and NPP-
2/NUP85 as examples, we report the benefit of combining tissue-specific gene
knockout and protein degradation to achieve complete protein depletion. We also
demonstrate that FLP-mediated recombination can be utilized to identify
transcriptomes in a *C. elegans* tissue of interest. We have adapted RNA polymerase
DamID (RAPID) for the FLP toolbox and by focusing on a well-characterized tissue,
the hypodermis, we show that the vast majority of genes identified by RAPID are
known to be expressed in this tissue. These tools allow combining FLP activity for
simultaneous gene inactivation and transcriptomic profiling, thus enabling the inquiry
of gene function in various complex biological processes.

Introduction

The nematode *C. elegans* is a popular model organism for genetic and biomedical research. A variety of methods have been developed to perturb gene expression and protein abundance in specific tissues and/or at particular developmental stages of *C. elegans*. These include tissue-specific RNA interference (RNAi) (WATTS *et al.* 2020), conditional protein degradation (ZHANG *et al.* 2015; WANG *et al.* 2017; HILLS-MUCKEY *et al.* 2021) and site-specific recombinases (NANCE AND FROKJAER-JENSEN 2019; DRIESSCHAERT *et al.* 2021). Especially Cyclization recombination (Cre) and Flippase (FLP) enzymes are popular tools for the control of gene expression in a spatiotemporal manner (HUBBARD 2014; DRIESSCHAERT *et al.* 2021). While Cre was derived from the P1 bacteriophage (SAUER 1987), FLP originates from *Saccharomyces cerevisiae* (SENECOFF *et al.* 1985). The Cre and FLP recombinases recognize 34 bp sequences, that are called loxP (locus of crossing-over in P1) and FRT (FLP recognition target), respectively. DNA located between two target sequences is either excised or inverted depending on their orientation. Both systems have been used efficiently in *C. elegans*, for example to induce gene knockout (KO) (RUIJTENBERG AND VAN DEN HEUVEL 2015; KONIETZKA *et al.* 2020), control ectopic expression (SCHMITT *et al.* 2012), ablation of neurotransmission (DAVIS *et al.* 2008), determination of gene expression (GOMEZ-SALDIVAR *et al.* 2020), mapping of chromatin organization (CABIANCA *et al.* 2019; HARR *et al.* 2020) and transgenesis (NONET 2020). Moreover, by utilizing both Cre and FLP, further versatility and cell specificity can be achieved (GE *et al.* 2020; NONET 2020). Thanks to community efforts, a growing number of characterized and stable Cre and FLP drivers are available, which facilitates the development and implementation of recombination-based tools (VOUTEV AND HUBBARD 2008; KAGE-NAKADAI *et al.* 2014; RUIJTENBERG AND VAN DEN HEUVEL 2015; MUNOZ-JIMENEZ *et al.* 2017; MACIAS-LEON AND ASKJAER 2018; AYUSO AND ASKJAER 2019). Most recently, a light-inducible Cre recombinase has been added to the toolbox (DAVIS *et al.* 2021).

In some cases, the study of gene function at specific developmental time points can be compromised by remaining protein levels after gene perturbation. At a time point of interest, remaining protein might occlude the true phenotype associated with the gene's loss of function by RNAi or conditional KO. This may be particularly relevant for long-lived proteins, such as histones and nuclear pore complex (NPC) proteins, that sometimes show lifelong persistence (SAVAS *et al.* 2012; TOYAMA *et al.* 2013). Besides, re-initiation of translation in the case of frameshift-mediated KO appears to be more prevalent than previously anticipated (COHEN *et al.* 2019). The combination of gene KO methods with conditional protein degradation may help to overcome this hurdle and completely deplete protein levels.

Tools for targeted protein degradation are constantly being developed and adapted for *C. elegans* and other model organisms (NANCE AND FROKJAER-JENSEN 2019). Auxin-inducible degradation (AID) of proteins tagged with a 44 amino acid AID degron offers both temporal and spatial control through exposure to auxin and expression of *Arabidopsis thaliana* TIR1 protein, respectively (ZHANG *et al.* 2015). Wang et al. (WANG *et al.* 2017) developed a system for the degradation of green fluorescent protein (GFP)-tagged proteins in *C. elegans* building on previous protein depletion systems (CAUSSINUS *et al.* 2011; ARMENTI *et al.* 2014). In this GFP degradation (GFPdeg) system, a camelid single-domain antibody fragment that specifically binds GFP ("GFP nanobody") is fused to the SOCS-box adaptor protein ZIF-1, that recruits CUL-2/Cullin-2, an E3 ligase. This leads to ubiquitination and

proteasomal degradation of the GFP-tagged protein. An advantage of GFPdeg is
that it is immediately applicable to any existing line expressing a GFP-tagged
protein, but it offers only spatial control in its current version. Degradation kinetics in
the AID or GFPdeg systems depend on the synthesis and turnover rates of the target
protein, which sometimes renders protein depletion incomplete. Therefore, using AID
or GFPdeg on its own may also not be sufficient for the inquiry of some biological
processes and could benefit from the combination with genetic tools, such as site-
specific recombination.

Here, we report an extension of the FLP-based toolbox for *C. elegans* by four
means. Firstly, we describe novel integrated single-copy FLP drivers that efficiently
induce recombination in the somatic gonad, distal tip cells, the P lineage or
coelomocytes. We also report an improved FLP driver for intestinal cells and
demonstrate that equally high recombination efficiency can be achieved for FLP
drivers on different autosomes. These drivers increase the number of cell types
amenable for easy FLP-mediated spatiotemporal control of gene expression.
Secondly, we adapt the GFPdeg system for FLP-based regulation, thereby enabling
spatiotemporal control of GFP fusion protein degradation. We show that inducible
GFPdeg is equally efficient as the original constitutive version. Thirdly, using two
endogenously GFP-tagged nuclear pore proteins, MEL-28/ELYS and NPP-2/NUP85,
as test cases, we provide evidence that combined FLP-mediated gene KO and
protein degradation is rapid and highly efficient to deplete stable proteins from a
specific tissue of interest. The possibility of using a single FLP driver to control both
conditional gene KO and protein degradation will facilitate the inquiry of tissue-
specific gene function in complex biological processes. Lastly, we present the
application of FLP-controlled RNA polymerase DamID (RAPID) (GOMEZ-SALDIVAR *et*
al. 2020) for identification of transcribed genes in the hypodermis. We propose that
combined with the growing list of tissue-specific FLP and Cre drivers, RAPID is an
efficient and attractive method to explore transcription profiles across tissues without
the need of physical isolation of cells.

Material and Methods

Plasmids

Promoter fragments for FLP expression constructs were PCR-amplified with *Pfu*
polymerase using wild type *C. elegans* genomic DNA as template and inserted as
*Not*I/*Kpn*I fragments into *Not*I/*Kpn*I of plasmid pBN338 (MUNOZ-JIMENEZ *et al.* 2017).
Details are in Supplementary Tables S2 and S3. Exceptions were (1) pBN527 *gpa-*
14p::FLP::SL2::mNG that was generated by first combining a 2.4 kb *Not*I/*Cla*I
restriction fragment from pNP259 (SCHMITT *et al.* 2012) with a 0.6 kb *Cla*I/*Kpn*I PCR
fragment followed by insertion into *Not*I/*Kpn*I of pBN338 and (2) pBN487 *unc-*
122p::FLP::SL2::mNG that was generated by inserting a *Not*I/*BsrG*I PCR fragment
into *Not*I/*Acc65*I of pBN338. For generation of pBN532 *hlh-*
12p::FLP::SL2::mTagBFP2, an intergenic *gpd-2/-3* sequence was amplified with
primers B1012 and B1364 (see Supplementary Table S1) using pBN338 as
template. A sequence encoding mTag2BFP2 was amplified with B1365 and B1366
using pMINIT mTagBFP2 (PORTA-DE-LA-RIVA *et al.* 2021) as template. The two PCR
fragments were PCR-stitched with primers B1012 and B1366 and inserted by
NEBuilder HiFi DNA Assembly (New England Biolabs) into *Mlu*I of pBN523. To
generate plasmids pBN548 *dat-1p::FLP::SL2::mNG* and pBN549 *hlh-*

12p::FLP::SL2::mTagBFP2 for integration into universal MosSCI landing sites
(FROKJAER-JENSEN *et al.* 2014), *Xma*I/*Not*I fragments from pBN338 and pBN532,
respectively, were inserted into *Ngo*MIV/*Not*I of pBN8 (RODENAS *et al.* 2012).

Targeting plasmids for SapTrap CRISPR/Cas9 gene modification were obtained as
described (SCHWARTZ AND JORGENSEN 2016). Details for construction of plasmids
pBN309 *mCh::baf-1*, pBN433 *g>f>p::npp-2* (">" denotes an FRT site) and pBN493
gf>p::baf-1 are provided in Supplementary Tables S1 and S3. The plasmids direct
the insertion of fluorescent tags immediately downstream of the endogenous start
codon. Plasmid pBN351 for insertion of *g>f>p* immediately upstream of the
endogenous *mel-28* start codon was generated from plasmids pMLS256, pMLS288
and pBN312 (SCHWARTZ AND JORGENSEN 2016; MUNOZ-JIMENEZ *et al.* 2017), sgRNA
oligonucleotides B1050 and B1051, 5' homology arm (499 bp; amplified with B1042
and B1043 using plasmid #1397 (GOMEZ-SALDIVAR *et al.* 2016) as template) and 3'
homology arm (258 bp; amplified with B1044 and B1045 using plasmid #1397 as
template) (see Supplementary Tables S1 and S3). Donor plasmid pBN477
containing FRT in the second *gfp* intron was constructed by inserting a 574 bp
*Xho*I/*Hin*dIII fragment from pBN312 into *Xho*I/*Hin*dIII of pMLS252 (SCHWARTZ AND
JORGENSEN 2016).

Plasmid pBN488 *hsp16.41p>mCh::his-58>vhhGFP4::zif-1* for targeted degradation
of GFP fusion proteins was constructed by amplifying *vhhGFP4::zif-1* with primers
B1333 and B1334 using plasmid pNH102 (a kind gift from Ngang Heok Tang and
Andrew Chisholm) as template. The PCR product was digested with *Xho*I and *Nhe*I
and inserted into plasmid pBN209 *hsp16.41p>mCh::his-58>dam::emr-1* (CABIANCA *et*
al. 2019) digested with the same enzymes to replace *dam::emr-1*.

Plasmid pBN537 *hsp16.41p>mCh::his-58>dam::rpb-6* for RAPID was constructed by
amplifying *rpb-6* with primers B1456 and B1457 using genomic DNA as template.
The PCR product was digested with *Ngo*MIV and *Nhe*I and inserted into plasmid
pBN209 digested with the same enzymes to replace *emr-1*.

Nematode culture, strains and genome engineering

Worm strains were maintained on Nematode Growth Medium (NGM) petri dishes
seeded with *Escherichia coli* OP50 (STIERNAGLE 2006). Unless otherwise noted,
strains were maintained at 16°C. The N2 strain was used as wild type reference.
Other strains are listed in Supplementary Table S4.

Single-copy insertion of FLP expression constructs into the *cxTi10882* locus on chrIV
or the *oxTi365* locus on chrV was done by microinjection into the gonads of EG6700
or EG8082 young adult hermaphrodites (FROKJAER-JENSEN *et al.* 2008; FROKJAER-
JENSEN *et al.* 2014; DOBRZYNSKA *et al.* 2016). Plasmids carrying FLP and *unc-119(+)*
as selection marker were injected at 50 ng/µl together with pCFJ601 *eft-3p::Mos1*
((FROKJAER-JENSEN *et al.* 2012), 50 ng/µl), pBN1 *lmn-1p::mCh::his-58* ((RODENAS *et*
al. 2012), 10 ng/µl), pCFJ90 *myo-2p::mCh* ((FROKJAER-JENSEN *et al.* 2008), 2.5 ng/µl)
and pCFJ104 *myo-3p::mCh* ((FROKJAER-JENSEN *et al.* 2008), 5 ng/µl). Integration of
the FLP expression constructs was confirmed by detection of fluorescent
mNeonGreen (mNG) or mTagBFP2 in the target tissue of wild-type moving animals
without ectopic mCherry expression from the co-injection markers. For FLP
constructs without fluorescent co-expression markers, correct expression pattern
was confirmed after crossing with the dual-color FRT reporter (see Confocal
Microscopy below).

Similarly, single-copy insertion of vhhGFP4::ZIF-1 and Dam::RPB-6 expression
constructs into the *ttTi5605* locus on chrII was done by microinjection into the
gonads of EG4322 young adult hermaphrodites. Injection markers and transposase
plasmid were injected at same concentrations as above. Integrity of the integrated
transgenes was confirmed by PCR, and, after crossing to FLP drivers, functional
assays (degradation of GFP fusion proteins for the GFPdeg system; see Confocal
Microscopy below).

CRISPR/Cas9-mediated genome modification to generate strains BN581 *baf-*
1(bq13[mCh::baf-1]) III, BN1164 *baf-1(bq47[gf>p::baf-1])* III, BN902 *mel-*
28(bq17[g>f>p::mel-28]) III and BN1044 *npp-2(bq38[g>f>p::npp-2])* I was performed
by microinjection of targeting plasmids pBN309, pBN493, pBN351 and pBN433,
respectively, into the gonads of HT1593 *unc-119(ed3)* III young adult
hermaphrodites (SCHWARTZ AND JORGENSEN 2016). Targeting plasmids were injected
at 65 ng/µl together with plasmids #1286 *eft-3p::Cas9* ((FRIEDLAND *et al.* 2013), 25
ng/µl), pBN1 (10 ng/µl), pCFJ90 (2.5 ng/µl) and pCFJ104 (5 ng/µl). Successful
modification of the target loci was confirmed by detection of fluorescent fusion
proteins in the nuclear envelope throughout the body of wild-type moving animals
without ectopic mCherry expression from the co-injection markers. The *unc-119(+)*
selection marker was next excised by microinjection with Cre expression plasmid
pMLS328 (SCHWARTZ AND JORGENSEN 2016). Finally, uncoordinated hermaphrodites
were outcrossed to N2 males to remove the *unc-119(ed3)* allele.

Insertion of an FRT in the *baf-1* 3'-UTR was performed by microinjection of BN1164
baf-1(bq47[gf>p::baf-1]) young adult hermaphrodites with a mixture containing 10 µM
baf-1 crRNA:trRNA duplex (B1312+B1067), 2.5 µM *dpy-10* crRNA:trRNA duplex
(B1068+B1067), 3 µM recombinant Cas9, 4 µM *baf-1* ssDNA repair template
(B1313) and 1 µM *dpy-10* ssDNA repair template (B725). Recombinant Cas9 was
produced by the CABD Proteomics and Biochemistry Facility. All DNA and RNA
oligonucleotides were purchased from IDT.

Confocal microscopy

To induce expression of either the dual-color FRT reporter or the GFPdeg system,
NGM plates with nematodes were incubated for 15 min floating in a 34°C water bath.
For evaluation of recombination efficiency and specificity, nematodes were left to
recover at 20°C for 2-3 hours prior to imaging. For GFP degradation experiments,
recovery was at 25°C for 1-24 hours as indicated for each experiment. For imaging,
nematodes were anesthetized in 5 µl 10 mM levamisole HCl on 2% agarose pads
and covered by a cover slip. Images were acquired with a Nikon Eclipse Ti
microscope equipped with Plan Fluor 40×/1.3 and Plan Apo VC 60×/1.4 objectives
and an A1R scanner using a pinhole of 1.2 airy units (MUNOZ-JIMENEZ *et al.* 2017).
Cells with GFP signal as an indication of a recombination event in the dual-color
reporter as well as mNG co-expression with FLP were evaluated from z-stacks
acquired with the Nikon NIS software. To analyze the reduction of protein levels of
GFP fusion proteins after FLP recombination and/or the induction of the GFP
degradation system, specific regions and focal planes were localized to acquire
reproducible sets of images. These are indicated in the text.

Fluorescence intensity was quantified in Fiji/ImageJ 2.1.0/1.53c (SCHINDELIN *et al.*
2012) by tracing and measuring the signal at the nuclear envelope with a freehand
line of width 3. After subtracting the background measured in animals without
expression of GFP, the data were analyzed in R Studio (1.3.1093 (TEAM 2020a);

running R 4.0.2 (TEAM 2020b)). Data were visualized either with base plotting or
using ggplot2 (WICKHAM 2016), mainly within the ggpubr framework (version 0.4.0
(KASSAMBARA 2020)). For statistical analysis, Shapiro-Wilk test of normality and
Levene's test for homogeneity of variance across groups were performed on the
data sets to determine if either ANOVA and pair-wise t tests or Kruskal-Wallis rank
sum test combined with Dunn's test for pairwise multiple comparisons of the ranked
data were most suitable. In all cases, p values were adjusted for multiple
comparisons (Benjamini & Hochberg method).

Brood size experiments

BN580 *g>f>p::baf-1*, BN1182 *mCh::baf-1/g>f>p::baf-1; nhx-2p::FLP* and BN1223
mCh::baf-1/gf>p::baf-1>; nhx-2p::FLP worms were cultured at 16°C until L4 larval
stage. For BN1182 and BN1223, homozygous *mCh::baf-1*, *g>f>p::baf-1* and
gf>p::baf-1> animals were selected under a fluorescence stereoscope. In each
replica, 8-10 L4 larvae per strain were transferred to individual NGM plates and
incubated at 20°C. During the reproductive period, adults were transferred to new
plates each day and the numbers of progenies were recorded. Adults were censored
if they desiccated on the walls of the plates or were damaged during transfer.
Statistical analysis was done in R using Wilcoxon rank sum tests and adjusted for
multiple comparisons (Benjamini & Hochberg method).

Lifespan experiments

Worms were cultured at 16°C until L4 larval stage. In each replica, 20 L4 larvae per
strain were transferred to newly seeded NGM plates and incubated for 15 min
floating in a 34°C water bath to induce the GFPdeg system. They were then
continuously incubated at 25°C. During the reproductive period, the live adults were
transferred to new plates each day to not confuse founders with their progeny.
Throughout the experiment, the number of live and dead animals were recorded.
Animals were censored if they disappeared, were damaged during transfer or died
due to internal hatching. Statistical analysis was done in R using the survival
package (version 3.2-7 (THERNEAU 2020)), which calculates Chi-square statistics
based on a log-rank test.

RAPID

Worms were grown at 20°C for at least two generations on NGM plates seeded with
E. coli GM119 *dam-* bacteria. Eggs were obtained by sodium hypochlorite treatment
(STIERNAGLE 2006) and incubated in M9 containing 0.01% Tween-20 (M9-T)
overnight at 20°C without food. For each strain and replica, four 100 mm NGM plates
with a lawn of GM119 were seeded each with approximately 1000 synchronized L1
larvae and incubated at 20°C for 48 h until reaching L4 stage. Larvae were washed
off the plates with M9-T, collected in 15 ml tubes and washed 8 times in M9-T. Next,
worms were transferred to 1.5 ml tubes washed twice in M9-T. The supernatant was
aspired to leave ~30 µl and tubes were snap frozen in liquid nitrogen and stored at -
80°C until used.

To purify genomic DNA, samples were lysed in 5 freeze/thaw cycles (1 min in liquid
nitrogen; 3 min in 37°C shaker) and processed with DNeasy Blood and Tissue Kit
(QIAGEN #69504). Two hundred ng of genomic DNA was used as input in the
DamID protocol that included an improved pool of AdR primers as described (DE LA
CRUZ RUIZ *et al.* 2022). Dam-methylated fragments were amplified with 14 PCR
cycles and quality checked on an Agilent Bioanalyzer 2100. Next, libraries were

prepared using 50 ng of PCR-amplified fragments and NEBNext® Ultra™ II DNA
Library Prep Kit for Illumina (E7645S). Library amplicons size range distribution was
checked as before, pooled and re-purified with 1x volume SpeedBeads (DE LA CRUZ
RUIZ *et al.* 2022) followed by sequencing at EMBL GeneCore using a NextSeq 500
(Illumina).

Sequencing fastq files were filtered to discard reads that do not start with the DamID
adapter followed by GATC ("CGCGGCCGAG*GATC*") using R Studio and the
damid.seq.R pipeline version 0.1.3 (available at github.com/damidseq/RDamIDSeq)
(SHARMA *et al.* 2016). Retained fragments were mapped to *C. elegans* UCSC
genome version ce11 with the same pipeline, which produced 9.5-41.9 million
uniquely mapped reads per sample. Normalization of Dam::RPB-6 reads to
GFP::Dam reads was performed using the damidseq_pipeline package
(https://owenjm.github.io/damidseq_pipeline/) (MARSHALL AND BRAND 2015) on the
bam files generated by the damid.seq.R script. Gene occupancy was computed as
log2(Dam::RPB-6/GFP::Dam) with polii.gene.call
(https://github.com/owenjm/polii.gene.call) using bedgraph output files from the
damidseq_pipeline and WBCel235 gene annotation. A false discovery rate (FDR) of
0.05 was used as threshold to call transcribed genes. Average profile plots and
heatmap were generated with SeqPlots (Version 3.0.12) using bedgraph output files
from the damidseq_pipeline and WBCel235 gene annotation; H3K36me3 signal
along either all protein coding genes or top20 transcribed genes on chromosome I
(included in the software) was used as comparison (STEMPOR AND AHRINGER 2016).
Genome browser views and Venn diagrams were generated with Integrative
Genomics Viewer (IGV Web App) (THORVALDSDOTTIR *et al.* 2013) and jvenn (Web
App) (BARDOU *et al.* 2014), respectively. Statistical significance of overlap between
protein coding gene lists was calculated with Fisher exact test for independence in
2x2 contingency tables using R Studio and a total count of protein coding genes of
19,975.

Data availability

See Table S1, Table S2, Table S3 and Table S4 for complete lists of primers,
promoters, plasmids and strains, respectively. Representative strains will be made
immediately available through the *Caenorhabditis* Genetics Center; plasmids and
other strains can be requested from the authors. Tables S5-S7 contain lists of
hypodermal genes identified by RAPID (Table S5), their overlap with other
hypodermal gene lists (Table S6) and their expression level in single-cell RNAseq
(Table S7). Raw and processed RAPID files are deposited at Gene Expression
Omnibus accession GSE214300 (https://www.ncbi.nlm.nih.gov/geo/).

Results

Characterization of new FLP lines

Previously, a series of efficient tissue-specific FLP-lines were generated by Mos1-
mediated Single Copy Insertion (MosSCI) (MUNOZ-JIMENEZ *et al.* 2017; MACIAS-LEON
AND ASKJAER 2018; AYUSO AND ASKJAER 2019; VAN DE WALLE *et al.* 2020). To expand
the existing toolkit, we created additional MosSCI lines that express FLP in specific
cell types and evaluated recombination efficiency with the heat-inducible dual color
reporter *bqSi294* that produces green nuclei (GFP::HIS-58) upon successful
recombination and red nuclei (mCherry::HIS-58) elsewhere (Figure 1A) (MUNOZ-

JIMENEZ *et al.* 2017). In most constructs (see below), an SL2 trans-splicing sequence followed by the gene encoding the fluorescent protein mNeonGreen (mNG) was inserted downstream of FLP (FLP::SL2::mNG), enabling visualization of FLP expression by co-expression of mNG (MUNOZ-JIMENEZ *et al.* 2017).

We first generated a line that expresses FLP under control of the *ckb-3* promoter, which is active in the Z1 and Z4 cells that compose the somatic gonadal primordium of the hermaphrodite in early L1 larvae (KROETZ AND ZARKOWER 2015). When crossed with the dual color reporter, we detected recombination in all 36 L1 larvae analyzed; in 35 larvae both Z1 and Z4 expressed GFP::HIS-58, and one larva expressed GFP::HIS-58 in a single cell (Figure 1B; Table 1). The majority of Z1/Z4 cells expressed GFP::HIS-58 only, indicating that recombination had taken place at both alleles (67/72 cells = 93%), whereas both green and red histones were observed in 4 cells. Of note, there were never more than the expected two green nuclei at this developmental stage (early L1), which highlights the tissue specificity of the *ckb-3p::FLP* line. The high efficiency of the line in the somatic gonadal primordium implies that it is very suitable for manipulation of all five tissues of the reproductive system: distal tip cells, gonadal sheath, spermatheca, spermatheca-uterine valve and the uterus (Figure 1B) (HUBBARD AND GREENSTEIN 2005).

The distal tip cells (DTCs) play important roles during development as the leading cells that guide migration of the gonad arms in hermaphrodites and also in control of germ cell proliferation (HUBBARD AND GREENSTEIN 2005). To express FLP specifically in DTCs, we used the *hlh-12* promoter which is expressed from L2 stage (TAMAI AND NISHIWAKI 2007). We observed recombination in both DTCs in all hermaphrodites analyzed (*n*=24, L2-adult stage; Figure 1B; Table 1). Except for one DTC in a young L2 larva, recombination had taken place at both alleles (47/48 cells = 98%) and was not observed in other cell types, demonstrating the high efficiency and specificity of the *hlh-12p::FLP* line.

We then generated a *lin-31p::FLP* line to enable recombination in the vulval precursor cells (VPCs). LIN-31 is expressed in P1.p-P11.p cells from the middle of L1 stage, when these cells are born (TAN *et al.* 1998). We focused our attention on the central VPCs in L2 larvae and on VPC descendants during vulval morphogenesis in L3 and L4 larvae. Importantly, recombination of both alleles was observed in all VPCs and VPC descendants in all animals (*n*=30, L2-adult stage; Figure 1B; Table 1). We also observed GFP::HIS-58 expression in a few head neurons, which is in agreement with a previous report on *lin-31* expression (Supplementary Figure S1A)(REECE-HOYES *et al.* 2007). Finally, we noted weak GFP::HIS-58 signal in hyp7 nuclei in late L3-L4 larvae, which we assume originates from fusion of descendants of P3.p, P4.p and P8.p with the hyp7 syncytium (Supplementary Figure S1A).

In the first version of our FLP toolkit, we used the *elt-2* promoter to drive recombination in the intestine with ~100% efficiency (MUNOZ-JIMENEZ *et al.* 2017). However, because recombination was also detected in the somatic gonad, we generated an alternative line based on the *nhx-2* promoter (NEHRKE AND MELVIN 2002). In young L1 larvae, the number of intestinal nuclei ranges from 19 to 22 and increases to 30-34 in L2 larvae (MCGHEE 2007). Recombination efficiency reached 98% based on green-only nuclei and 100% when also including nuclei with expression of both green and red histones (*n*=27, Figure 1B; Table 1). This demonstrates high recombination efficiency, comparable to the *elt-2p::FLP* line.

Importantly, we did not detect recombination outside the intestine with the *nhx-*
2p::FLP line, so we recommend to use this line.

We next generated an *unc-122p::FLP* line to enable recombination in coelomocytes
(CCs), which are specialized cells thought to be involved in the innate immune
response and scavenging (LORIA *et al.* 2004; ALTUN AND HALL 2009). Two anterior
CC pairs are born during embryogenesis whereas a posterior pair is born right
before the L1/L2 molt. Consistently, in 8 of 8 L1s we observed 4 GFP::HIS-58
expressing CCs, and in 23 of 23 L3-L4 larvae 6 GFP::HIS-58 expressing CCs
(Figure 1B; Table 1). Expression of mCh::HIS-58 was not detected in the CCs,
indicating that both alleles had recombined, whereas GFP::HIS-58 expression was
restricted to the CCs.

We also attempted to generate additional lines to target specific neuronal subtypes,
including cholinergic neurons (using the *unc-17* promoter) and *gpa-14*-expressing
neurons. However, although both lines expressed the marker mNG in the expected
tissues, they also showed either recombination in unexpected cell types (for *unc-*
17p::FLP) or inefficient recombination (for *gpa-14p::FLP*) and were not characterized
in detail (Supplementary Figure S1B-C; see Discussion).

In the strains described above, co-expression of mNG serves to visualize when and
where FLP is expressed. However, in some experiments, the signal from mNG might
interfere with other fluorescent proteins. To circumvent this, we replaced mNG with
mTagBFP2 that emits blue light. When placed under control of the *hlh-12* promoter,
we observed mTagBFP2 expression and FLP-mediated recombination in both DTCs
of all animals (*n*=12; Supplementary Figure S1D). The sequences encoding mNG
and mTagBFP2 in the MosSCI plasmids are flanked by *Mlu*I restriction sites, which
facilitates the exchange of the former for the latter in the creation of new strains.

All lines described so far involve FLP expression constructs on chromosome IV. To
increase the versatility of the FLP toolkit, we subcloned *dat-1p::FLP* and *hlh-*
12p::FLP constructs to *ttTi5605* targeting vectors which enable insertion into
universal MosSCI sites on any autosome (FROKJAER-JENSEN *et al.* 2014). After
insertion of the FLP expression constructs on chromosome V we tested their
efficiency on the dual-color reporter and found that they both induce specific
recombination in all expected cells (Table 1). This suggests that other existing FLP
constructs can easily be adapted for insertion into different loci without compromising
their efficiency.

Conditional gene knockout by GFP exon removal

We previously designed a conditional knockout (KO) cassette based on GFP
containing FRT sites in introns 1 and 2 (Figure 2A) (MUNOZ-JIMENEZ *et al.* 2017).
Excision of GFP exon 2 by FLP causes a frame shift and a premature termination
codon, which most likely targets the mRNA for nonsense-mediated decay (ARRIBERE
et al. 2020). The premature termination codon precedes three short open reading
frames that are out of frame with respect to the gene downstream of GFP. Together,
this should completely prevent expression of the tagged protein. To test this
experimentally, we compared the consequence of either complete removal of the
baf-1 gene (*gf>p::baf-1>*; ">" denotes the position of FRT; see Figure 2A) or removal
of GFP exon 2 in an endogenously tagged *gfp::baf-1* line (*g>f>p::baf-1*). BAF-1 is an
essential and ubiquitously expressed nuclear envelope protein (GORJANACZ *et al.*

2007). In both cases, we used the *nhx-2p::FLP* line described above to induce *baf-1*
KO specifically in intestinal cells, which negatively affects health (Munoz-Jimenez *et*
al. 2017). As controls we included (i) the *g>f>p::baf-1* line without FLP expression
and (ii) an endogenously tagged *mCh::baf-1* line without FRT sequences but with
FLP expression. While hermaphrodites of the two control lines produced an identical
number of progeny (*g>f>p::baf-1* 299±25 [mean±SD], n=10; *mCh::baf-1* 301±55,
n=15), fertility was reduced equally by 31% in the two conditional knockout lines
(*g>f>p::baf-1; nhx-2p::FLP* 207±71, n=15; *gf>p::baf-1>; nhx-2p::FLP* 206±68,
n=12)(Figure 2B). Based on this, we conclude that excision of the second exon of
the GFP KO cassette causes an equally strong loss of function phenotype as
complete gene removal. Moreover, it provides the advantage that insertion of the
GFP KO cassette in the 5'-end of a target gene in a single step is easier than
inserting both a GFP tag (with a single FRT) and a second FRT inside or
downstream of the gene.

While conditional gene KO prevents synthesis of new transcripts, mRNA and protein
molecules already existing in the cell might persist for days, weeks or even months
(Toyama *et al.* 2013). This may obscure both investigation of gene function and
potential therapeutic interventions. To address this, we evaluated the kinetics of
GFP::MEL-28 depletion from hypodermal cells of *g>f>p::mel-28; dpy-7p::FLP*
hermaphrodites. MEL-28 is a component of the long-lived NUP107 nuclear pore
subcomplex and is required for DNA segregation and post-mitotic nuclear assembly
(Fernandez and Piano 2006; Galy *et al.* 2006; Toyama *et al.* 2013). Although the
dpy-7 promoter is active already during embryogenesis to drive FLP expression,
~95% of GFP::MEL-28 was still present in L1 larvae as compared to the control
g>f>p::mel-28 line without FLP expression (Figure 3; compare Control and KO in
3B). In contrast, in L2 larvae and young adults a dramatic reduction of the median
GFP::MEL-28 signal to ~12% and ~3%, respectively, was observed specifically in
the hypodermis (Figure 3). This demonstrates the potential of conditional KO
methods to evaluate the function of long-lived proteins in late larvae and adults (see
also below), whereas they might be less suitable in early larvae.

FLP-mediated protein degradation

To enable analysis of long-lived proteins during development, we developed a
system for FLP-mediated, inducible and rapid protein degradation from specific cells.
A potent system for constitutive tissue-specific degradation of GFP-tagged proteins
based on expression of vhhGFP4::ZIF-1, a GFP-binding nanobody fused to ZIF-1,
was recently developed in *C. elegans* (Wang *et al.* 2017). To achieve both temporal
and spatial control of degradation of GFP-tagged proteins (hereafter termed
GFPdeg), we replaced the *gfp::his-58* sequence of the dual color reporter with a
sequence encoding vhhGFP4::ZIF-1. In this design, tissue-specificity is controlled by
the choice of a particular FLP-expressing line, whereas temporal control is provided
by the heat-inducible *hsp-16.41* promoter (Figure 4A).

We first determined that basal expression from the *hsp-16.41* promoter in the
absence of heat induction does not cause depletion of GFP::MEL-28 in the
hypodermis (Figure 4B). This implies that the system can be used to evaluate the
function of essential proteins late in development by raising the animals at 16°C until
induction. We next investigated whether the heat-induced GFPdeg system is equally
efficient as the original constitutive variant (Wang *et al.* 2017). We focused on

degradation of GFP::MEL-28 in intestinal cells. GFP intensity was quantified in one
or two intestinal cells closest to the pharynx and to the microscope lens to ensure
consistent measurements. Other intestinal cells often lie below the gonad, impeding
reproducible quantification. Three hours after a short heat pulse (15 min; 34°C) the
median GFP::MEL-28 intensity was reduced to ~27% in the inducible GFPdeg strain,
whereas slightly less protein remained in the constitutive GFPdeg strain (~18%;
Figure 4C, left graph). By exposing the animals to two 15 min heat pulses at 34°C,
one 24h and another 3h prior to imaging, the efficiencies of the inducible and
constitutive GFPdeg systems were equal (p=0.91; Figure 4C, right graph; Figure
4D). This demonstrates, that additional temporal control of the tissue-specific
GFPdeg module can be achieved without compromising degradation efficiency.
Depending on the protein of interest, induction conditions may have to be optimized.
For instance, the efficiency of endogenous GFP::MEL-28 degradation in the intestine
by either the constitutive or the inducible system (Figure 4C-D) is lower than the
depletion reported for endogenous GFP::MDF-1 based on the identical constitutive
GFPdeg strain (WANG *et al.* 2017). To evaluate the degradation dynamics in more
details, we measured GFP::MEL-28 levels in hypodermal cells at different time
frames after a single induction of GFPdeg (15 min; 34°C). Hypodermis-specific
expression of vhhGFP4::zif-1 was ensured by excision of *mCh::his-58* with FLP
under the control of the *dpy-7* promoter. All images were taken from a region of the
hypodermis closest to the lens and just posterior to the pharynx. One hour after
induction, the median level of GFP::MEL-28 was reduced to ~55% and further down
to ~19-20% after 2-4 h (Figure 5). A possible explanation for the remaining
GFP::MEL-28 even at the optimal time points (2-4 h) could be novel protein
synthesis since the degradation system will only be active for a finite time after
induction. Indeed, substantial protein re-synthesis was observed 24 h after heat
shock (Figure 5). The time frame of optimal degradation efficiency allows some
flexibility in protocol design. However, these protocols, especially number and timing
of heat inductions, presumably need to be tailored to the protein of interest and its
turnover.

Combination of FLP-mediated gene knockout and protein degradation

Having available two FLP-controlled systems for inhibition of protein expression, we
decided to test the efficiency of combining them. In this setup, the same FLP driver
controls both conditional gene KO and protein degradation (Figure 6A). As a proof of
principle, we evaluated the combined action of the GFP KO cassette and GFPdeg on
the expression of endogenously GFP-tagged MEL-28 in the hypodermis. GFPdeg
was induced by exposing young L1s to a short heat pulse (15 min; 34°C) followed by
incubation at 16°C. The effect was first evaluated in L1 larvae after 3 h and again in
L2 larvae as well as young adults. In contrast to the GFP KO system which alone
had insignificant effects in L1 larvae, the combination of the two systems led to
~80% reduction of median GFP::MEL-28 signal (Figure 3). In L2 larvae, GFP::MEL-
28 levels in the hypodermis were further decreased to ~3% and undetectable in most
young adults. Thus, in all three life stages tested, the combination of conditional
gene knockout and protein degradation provided the most efficient inhibition of
protein expression. We also compared the efficiency of either conditional gene KO
alone or in combination with targeted protein degradation by inserting the GFP KO
cassette in frame after the start codon of another stable nuclear pore protein, NPP-

2/NUP85. In young adults, the conditional KO caused a reduction of ~97% of
GFP::NPP-2 in the hypodermis, whereas only ~1.5% remained when conditional KO
was combined with the GFPdeg system (Figure 6B-C).

To test if these dramatic reductions of endogenous GFP::NPP-2 in the hypodermis
had consequences on the health of the animals, we determined their lifespan at
25°C. We followed >175 animals for each genotype and observed a 22-33%
reduction in median lifespan from 9 days post L4 in GFP::NPP-2 control animals
(95% confidence interval [CI] = 9-10 days) to 7 days in conditional KO animals (95%
CI = 6-7 days) and to 6 days in animals with combined conditional KO and GFPdeg
systems (95% CI = 6-7 days) (Figure 6D. Expression of FLP and GFPdeg in the
absence of the GFP KO cassette had no effect on the lifespan (p=0.3). The
statistically insignificant shift to the left of the lifespan curve for the animals with the
combined conditional KO and GFPdeg (red line) as compared to the animals with the
conditional KO only (dark red line; p=0.5) is in line with the unsubstantial difference
in GFP::NPP-2 signal intensity between the two strains. We conclude that conditional
NPP-2 KO in the hypodermis causes a dramatic shortening of lifespan, illustrating
the suitability of the combination of the GFP KO cassette and the GFPdeg system for
evaluation of physiological consequences of protein depletion in specific tissues.

Tissue-specific transcriptomic profiling by RAPID

Various complementary methods are available for identification of gene expression
profiles in specific cell types in *C. elegans*, including physical isolation of cells or
nuclei of interest (HAENNI *et al.* 2012; STEINER *et al.* 2012; KALETSKY *et al.* 2018),
affinity purification of RNAs from specific cells (BLAZIE *et al.* 2017), trans-splicing-
based RNA tagging (MA *et al.* 2016), and mapping of RNA polymerase occupancy
(GOMEZ-SALDIVAR *et al.* 2020; KATSANOS *et al.* 2021). RNA polymerase DamID
(RAPID) (GOMEZ-SALDIVAR *et al.* 2020) and Targeted DamID (TaDa) (KATSANOS *et al.*
2021) are based on tissue-specific basal expression of *E. coli* Dam
methyltransferase fused to the RNA polymerase subunit RPB-6. The Dam::RPB-6
fusion protein provides an *in vivo* footprint of RNA polymerases on transcribed genes
by specific methylation of GATC sequences that are identified by enzymatic and
deep sequencing techniques (MARSHALL *et al.* 2016; ASKJAER AND HARR 2021).

RAPID relies on recombinases to achieve tissue specificity and has been used in
body wall muscles, intestine and specific neurons (GOMEZ-SALDIVAR *et al.* 2020). To
increase the number of tissues immediately amenable to RAPID, we adapted it for
FLP-mediated spatiotemporal control. We cloned and inserted a *dam::rpb-6* fusion
gene downstream of the FRT-flanked *mCh::his-58* cassette used above and under
control of the *hsp-16.41* promoter (Figure 7A). We introduced a single copy of this
construct into the *C. elegans* genome and crossed the resulting line with a *dpy-*
7p::FLP driver. Nematodes were cultured at 20°C to ensure low levels of Dam::RPB-
6 expression in the hypodermis and total genomic DNA was purified from L4 larvae.
DNA from animals expressing GFP::Dam was used to control for unspecific
methylation. The genome-wide association profile of Dam::RPB-6 was determined
by deep sequencing (see Materials and Methods), which revealed a list of 2331
protein coding genes with FDR < 0.05 (Table S5). When plotting the RAPID signal
along all protein coding genes, an increase in signal is observed from the
transcriptional start site (TSS) to the transcriptional end site (TES), similar to
previous RAPID and TaDa observations (GOMEZ-SALDIVAR *et al.* 2020; KATSANOS *et*

al. 2021)(Figure 7B, left). However, clustering the signal from individual genes using
k-means algorithm (with default number of clusters = 5 (STEMPOR AND AHRINGER
2016)) suggests that the average signal masks multiple different patterns of RAPID
signal along genes (Figure 7B, middle). A more uniform RAPID signal from TSS to
TES is also observed when averaging the 2331 genes determined as expressed in
the hypodermis with FDR < 0.05 (Figure 7B, right). Among the transcribed genes are
several known and expected hypodermal genes, such as *col-39*, *col-79*, *col-80* and
sqt-1 that express collagens as well as ubiquitously expressed genes *eef-1A.1* and
fib-1 (Figure 7C).

To evaluate the specificity of RAPID, we compared the list of 2331 transcribed genes
to three datasets of genes reported to be expressed in the hypodermis (BLAZIE *et al.*
2017; KALETSKY *et al.* 2018; KATSANOS *et al.* 2021) (Figure 7C-D; Table S6). Our list
overlapped significantly with each of the individual datasets (54-66% overlap;
p<2.2e-16 Fisher's exact test), and 1860 genes (80%) overlapped with at least one
of the three datasets. These overlaps are comparable to those observed when
comparing the three other datasets among each other (Figure 7D). Moreover, when
exploring a small arbitrarily chosen subset of the 471 genes not overlapping with any
of the three other datasets, at least 22/50 (44%) are reliably detected in hypodermal
cells in single cell RNAseq performed on embryos (PACKER *et al.* 2019) (Table S7).
This argues that many of the genes identified uniquely by RAPID are expressed in
the hypodermal lineage during development. We conclude that RAPID is an
attractive and versatile method to identify transcribed genes in specific tissues with
high confidence.

Discussion

An increasing number of specific and efficient FLP and Cre drivers generated by the
C. elegans community are facilitating a variety of precise functional assays to
interrogate neural circuits (DAVIS *et al.* 2008; SCHMITT *et al.* 2012; KONIETZKA *et al.*
2020; DAVIS *et al.* 2021), cell proliferation and differentiation (RUIJTENBERG AND VAN
DEN HEUVEL 2015), gene expression programs (GOMEZ-SALDIVAR *et al.* 2020; VAN DE
WALLE *et al.* 2020) and chromatin organization (CABIANCA *et al.* 2019; HARR *et al.*
2020). In this study, we report four new FLP lines that reliably induce recombination
specifically in either distal tip cells, the somatic gonad, coelomocytes or in epithelial
Pn.p cells. The efficiency of recombination was ~100% in all animals evaluated for
the four lines. We also demonstrate that similarly high recombination efficiency is
achieved when FLP is expressed from single-copy transgenes on different
autosomes. In the case of the *ckb-3p::FLP* line, recombination was observed already
in the earliest progenitors Z1 and Z4, which implies that the line is suitable to target
all cell types of the somatic gonad, including distal tip cells, gonadal sheath,
spermatheca, spermatheca-uterine valve and the uterus (HUBBARD AND GREENSTEIN
2005). In contrast, the activity of the *hlh-12p::FLP* line is restricted to the distal tip cell
and may thus be useful to investigate its stem cell niche function (KIMBLE AND SEIDEL
2008). Similarly, biological processes in coelomocytes are amenable to manipulation
by the *unc-122p::FLP* line, whereas the *lin-31p::FLP* line induces recombination in
the cells producing the vulva. We previously created a FLP line to induce
recombination in intestinal cells based on the *elt-2* promoter. However, because this
promoter can also drive expression in other tissues (RUIJTENBERG AND VAN DEN
HEUVEL 2015; MUNOZ-JIMENEZ *et al.* 2017), we developed here a novel line based on

the *nhx-2* promoter. The new line is equally efficient to produce recombination in the intestine (Table 1) and with the advantage of not provoking recombination elsewhere. We also attempted to construct lines for specific neurons using the promoters of *gpa-14* and *unc-17*. In the case of *gpa-14*, we observed variable recombination rates in FLP-expressing cells (0, 1 or 2 alleles of the single-copy reporter). It seems that the behavior of the specific cells is reproducible between individuals, but we do not know if this is due to differences in FLP expression levels and/or the FRT sites being less accessible in certain cells. For the *unc-17p::FLP* line, we observed widespread recombination in cholinergic neurons as expected but also in the hypodermis. We speculate that this could be due to an inability of our construct to precisely reflect *unc-17* expression (PEREIRA *et al.* 2015), although we also note that two cholinergic neurons, PVNL and PVNR, are sister cells to two hypodermal cells (TLappa and TRappa). Thus, if *unc-17p::FLP* is activated already in the progenitor cells TLapp and TRapp, GFP::HIS-58 expression would be expected in both cell types.

We previously proposed that inserting a GFP KO cassette (with FRT sites in *gfp* introns 1 and 2) immediately after the start codon of a gene provides an informative GFP fusion protein and a conditional KO allele in a single step (MUNOZ-JIMENEZ *et al.* 2017). In this study we have further validated this approach by comparing strains that either carry the GFP KO cassette in the endogenous *baf-1* locus or a GFP cassette with a single FRT and a second FRT downstream of *baf-1*. When combined with an *nhx-2p::FLP* driver to KO *baf-1* specifically in the intestine, identical reductions of fertility were detected in the two strains. Furthermore, using the GFP KO cassette, we determined that conditional KO of *npp-2*/NUP85 in the hypodermis caused a dramatic lifespan shortening. We therefore conclude that insertion of the GFP KO cassette into an endogenous locus by CRISPR provides an attractive dual tool to study protein dynamics and KO phenotypes in a single workflow. A limitation of the approach is that the GFP KO cassette should be inserted in the 5'-end of the open reading frame. As with other gene tagging experiments, tests for hypomorph phenotypes should be performed.

Although gene KO is often considered an ultimate test for gene function, slow mRNA and protein turnover can delay the appearance of loss-of-function phenotypes. This is particularly relevant for organisms with rapid development and relatively short lifespan, such as *C. elegans*. Moreover, the irreversible nature of gene KO may be disadvantageous in some experimental designs. Targeted protein degradation is a relevant alternative and auxin-inducible degradation (AID) stands out as an attractive and popular option due to its fast kinetics and both inducible and reversible nature (ZHANG *et al.* 2015). Concerns have been raised regarding auxin-independent depletion of AID-tagged proteins (see references in (NANCE AND FROKJAER-JENSEN 2019)) and potential effects of auxin on endoplasmic reticulum stress (BHOI *et al.* 2021), but recent optimization of AID tools may have overcome this (HILLS-MUCKEY *et al.* 2021). We show here that GFPdeg, an alternative protein degradation system based on a GFP-binding nanobody (WANG *et al.* 2017) and adapted for FLP-mediated spatiotemporal control, can efficiently deplete 80% of a long-lived nuclear pore protein, MEL-28/ELYS, fused to GFP. By comparing the degradation of GFP::MEL-28 in intestinal cells using either the original, constitutive version of GFPdeg (WANG *et al.* 2017) or the heat inducible version of GFPdeg developed in this study, we conclude that they are equally efficient. Importantly, the GFPdeg system is immediately applicable to study the function of any GFP-tagged fusion

protein. We chose to focus on stable nuclear pore proteins (SAVAS *et al.* 2012; TOYAMA *et al.* 2013) in our study as they are probably more challenging to deplete but we assume that higher depletion efficiencies will be observed for proteins with shorter half-lives. The original GFPdeg version (WANG *et al.* 2017) has the advantage that it involves only two components (GFP-tagged fusion protein and constitutive, tissue-specific GFPdeg) and does not require induction. In contrast, our FLP-based GFPdeg system includes three components (GFP-tagged fusion protein, tissue-specific FLP driver and heat-inducible, FRT-flanked GFPdeg) and requires one or two short 15 min exposures to 34°C. On the other hand, our inducible system allows to evaluate processes at any stage in development or during aging by raising the nematodes at standard temperatures until induction at the desired time point. This cannot be achieved with the constitutive system where the animal's development potentially arrests once the concentration of the target protein is below a required threshold. We envision that the FLP-controlled GFPdeg system can be further modified to incorporate other inducers (MONSALVE *et al.* 2019; WURMTHALER *et al.* 2019; DRIESSCHAERT *et al.* 2021) in situations where short heat pulses are incompatible with the experimental design.

Depending on the experimental setup, either site-specific gene KO or targeted protein degradation may be sufficient to elucidate gene function. However, bimodal control of protein levels may be particularly valuable for the study of long-lived proteins during early developmental stages as highlighted in this study. By combining the GFP KO cassette and the GFPdeg system, we have generated a protocol that provides two-fold control of tissue-specific protein depletion. On one hand, a specific FLP driver induces gene KO in a tissue of interest and thus prevents synthesis of mRNA from the target gene. On the other hand, the same FLP driver excises tissue-specifically the FRT-flanked *mCh::his-58* cassette from the GFPdeg construct, thus enabling heat-induced and rapid degradation of existing GFP-tagged target protein. We show that the combined KO and GFPdeg systems can deplete GFP::MEL-28 and GFP::NPP-2 to non-detectable levels, thus offering a highly efficient approach for spatiotemporal control of protein levels in a multicellular organism.

One of the strengths of recombinase-based genome engineering is its versatility: the possibility of using collections of stable FLP and Cre drivers to control a variety of different effectors through straight-forward genetic crosses (HUBBARD 2014; NANCE AND FROKJAER-JENSEN 2019; DRIESSCHAERT *et al.* 2021). The Cre/loxP system was recently utilized to generate tissue-specific transcription footprints by RNA polymerase DamID (RAPID) in *C. elegans* (GOMEZ-SALDIVAR *et al.* 2020). In RAPID and other DamID methods for transcriptional profiling *E. coli* Dam is fused to an RNA polymerase subunit and expressed at low levels to mark *in vivo* transcribed genes (SOUTHALL *et al.* 2013; ASKJAER AND HARR 2021). In this study we have widened the range of tissues immediately accessible to RAPID by adapting RAPID for FLP-mediated spatial control. We have validated the potential of RAPID in *C. elegans* by expressing Dam::RPB-6/POLR2F specifically in the hypodermal lineage through the use of a *dpy-7p::FLP* driver and identified 2331 transcribed genes. We compared our list with hypodermal data from another recent DamID-based (TaDa) study (KATSANOS *et al.* 2021) and two RNAseq-based studies (BLAZIE *et al.* 2017; KALETSKY *et al.* 2018) and found a significant overlap in all cases (p<2.2e-16). We propose that the set of 847 protein coding genes identified in all four studies (Table S6) can be considered a high-confidence transcriptional signature of the hypodermis. A pilot

comparison of genes found uniquely in RAPID (i.e. not confirmed by any of the three other methods) revealed a relevant overlap with genes detected in hypodermal cells by single-cell RNA-seq (PACKER *et al.* 2019), suggesting that many of the unique RAPID genes are true positive hits.

Although the overlap between the gene lists identified in our RAPID experiment and the TaDa study is much higher than would be expected by chance, ~50% of the genes identified in one study were not found in the other. In both cases, Dam::RPB-6 was used to mark transcribed genes and samples were prepared at the L4 stage so how can the many unique genes be explained? An important difference is that Katsanos and co-workers used a mutated version of the *dpy-7* promoter ("*dpy7syn1*") for TaDa in the hypodermis to avoid signal from seam cells (KATSANOS *et al.* 2021), whereas we used a wild type *dpy-7p* sequence to drive FLP expression (MUNOZ-JIMENEZ *et al.* 2017). Katsanos and co-workers also determined the transcriptomic profile for seam cells and found that ~1/3 of the genes in their hypodermis and seam cell datasets are unique (KATSANOS *et al.* 2021). Our RAPID gene list overlaps significantly with both the entire seam cell gene list (1013 overlapping genes; p<2.2e-16) and the unique seam cell gene list (106 overlapping genes; p=0.016), suggesting that part of the RAPID signal in our experiments is derived from seam cells (Figure S2). It should be noted that the lineages of the two cell types are interconnected because many of the daughter cells from dividing seam cell precursors fuse with the hypodermis during larval development. Differences in the protocols for preparation of deep sequencing libraries (sequences of adapters, PCR conditions, +/- removal of adapters and fragmentation, etc.) are also likely to introduce variation in the lists of identified genes. Finally, the strategies underlying tissue specificity is also different between RAPID and TaDa (ASKJAER AND HARR 2021). Spatial control of basal Dam::RPB-6 expression is achieved in RAPID by combining FLP or Cre-mediated genome recombination with a non-induced *hsp* promoter (Figure 7A), whereas TaDa is based on a tissue-specific promoter and translation re-initiation of a bicistronic mRNA containing an upstream open reading frame and *Dam::rpb-6* separated by one or several stop codons. TaDa is feasible with a single transgene whereas RAPID relies on a recombinase transgene and a *Dam::rpb-6* transgene typically integrated on separate chromosomes. Although this appears to favor TaDa versus RAPID, the growing list of stable and efficient FLP and Cre drivers implies that different tissues can be explored with RAPID through a simple cross to combine the recombinase and *Dam::rpb-6* transgenes. In contrast, TaDa requires the design, cloning and transgenesis for each new cell type to be analyzed.

Because the methylation on GATC by Dam is only erased when cells enter S phase and replicate their DNA, RAPID and TaDa provide the history of transcription during the life of the cell. PATseq and FACS produce instead a compilation of all mRNAs present in the cell at the time of harvest. Depending on the biological question to be addressed, a particular method might be most suitable. The PATseq and FACS studies report the largest hypodermal gene sets (BLAZIE *et al.* 2017; KALETSKY *et al.* 2018), but they also suffer higher intrinsic risks of including false positive hits. For instance, FACS isolation of a particular cell type might include contaminants in the form of cells from other tissues. Similarly, in the PATseq protocol, mRNA from unlabeled cells might potentially co-purify with the mRNA from the target cells. Such false positives are difficult to control for in the FACS and PATseq protocols. In contrast, RAPID and TaDa are based on isolation of genomic DNA from the entire

organism followed by amplification of methylated GATC fragments from a tissue of interest, library preparation and a statistical analysis to identify genes with a higher Dam::RPB-6/GFP::Dam methylation ratio. Importantly, although the signal from Dam::RPB-6 reflects RNA polymerase occupancy on transcribed genes rather than transcription itself, the signal intensity correlates with transcript abundance (SOUTHALL *et al.* 2013; GOMEZ-SALDIVAR *et al.* 2020; KATSANOS *et al.* 2021). Furthermore, comparing Dam::RPB-6 methylation values across experimental conditions enables identification of differentially expressed genes (WIDMER *et al.* 2018).

We conclude that FLP-controlled RAPID has the capacity to identify transcribed genes in a tissue of interest with high confidence. We have tested the method in the hypodermis, but it is easily applied to other cell types for which an appropriate FLP driver is available. In particular, we envision that RAPID will be advantageous for cell types that are hard to isolate in an intact state, such as neurons (GOMEZ-SALDIVAR *et al.* 2020).

Table 1 Efficiency of FLP-mediated recombination

Promoter[a]	Cell type	Stage[b]	Observed[c]	Expected	Efficiency
ckb-3	Somatic gonad	Early L1	2.0±0.2 (36)	2	99%
dat-1	Dopaminergic neurons	L3-L4	8.0±0.0 (11)	8	100%
hlh-12	Distal tip cell	L2-adult	2.0±0.0 (24; chrIV[a]) 2.0±0.0 (10; chrV[a])	2	100%
lin-31	P1.p-P11.p (P5.p-P7p)	L2-L4	3.0±0.0 (14)	3[d]	100%
		L3	12.0±0.0 (13)	12[d]	100%
		L4	22.0±0.0 (3)	22[d]	100%
nhx-2	Intestine	L1	20.0±0.0 (10)	20 (19-22)	100%
		L2	31.9±0.6 (17)	32 (19-22)	100%
unc-122	Coelomocytes	L1	4.0±0.0 (8)	4	100%
		L3-L4	6.0±0.0 (23)	6	100%

[a] Single-copy FLP constructs inserted on chrIV except *dat-1p::FLP* and one of the *hlp-12p::FLP* constructs that were inserted on chrV.

[b] Developmental stage that was used for determination of recombination efficiency.

[c] Average number of GFP positive nuclei ± SD; number in brackets: *n*.

[d] Efficiencies were calculated based on recombination in central Pn.p cells and their descendants.

Acknowledgments

We are grateful to Andrew Chisholm, Carmen Ortiz González, Michael Krieg,
Montserrat Porta, Ngang Heok Tang, Patricia Regina Rider García and Sara
Castaño Díaz for reagents and advice. We also wish to acknowledge Ana Miñan
Fernández from the CABD Functional Genomics Facility for microinjections, Laura
Tomás from the CABD Proteomics and Biochemistry Facility for recombinant
proteins, Katherina García from the CABD Imaging Facility for technical assistance;
the Caenorhabditis Genetics Center (CGC), which is funded by the National
Institutes of Health (NIH) Office of Research Infrastructure Programs (P40
OD010440) and WormBase. This project was funded by the Spanish State Research
Agency and the European Regional Development Fund (BFU2016-79313-P,
CEX2020-001088-M and PID2019-105069GB-I00).

799 **Figure legends**

Figure 1. Characterization of novel FLP lines. (A) Schematic representation of the
dual color reporter used to evaluate recombination activity. Upon temperature
induction of the *hsp-16.41* promoter, the reporter expresses red (mCh) or green
(GFP) histone HIS-58 in cell lineages without or with FLP-mediated recombination,
respectively. (B) Evaluation of FLP lines crossed with the dual color reporter.
Although the FLP lines co-express diffuse mNG in the cytoplasm, GFP::HIS-58 is
clearly visible in the expected nuclei. (Ba) *ckb-3p::FLP::SL2::mNG*: maximum
projections of 2-3 confocal sections of BN854 L1 larva (Ba') and larva at L3/L4 molt
(Ba''). The Z1 and Z4 somatic gonad precursor cells are indicated. (Bb) *hlh-
12p::FLP::SL2::mNG*: single confocal sections of BN1204 L2 larva (Bb') and young
adult (Bb''). The distal tip cells (DTCs) are indicated. (Bc) *lin-31p::FLP::SL2::mNG*:
maximum projections of three confocal sections of BN1023 L2 larva (Bc') and L4
larva (Bc''). The posterior daughters of P1-P7 and vulA-vulF forming the vulva are
indicated. (Bd) *nhx-2p::FLP::SL2::mNG*: maximum projection of five confocal
sections of BN999 L1 larva (Bd'; indicated area shown enlarged in Bd''). (Be) *unc-
122p::FLP::SL2::mNG*: maximum projections of four confocal sections of BN1029 L1
larva (Be') and L4 larva (Be''). The three pairs of coelomocytes are indicated. Scale
bars 10 µm.

Figure 2. Comparison of GFP KO cassette and complete gene KO. (A) Schematic
representation of the two KO strategies. In the first strategy (top), a GFP KO
cassette with FRTs in *gfp* introns 1 and 2 is inserted immediately downstream of,
and in frame with, the start codon of a gene of interest (here *baf-1*). FLP-mediated
recombination leads to a frame shift and a premature termination codon ("Stop"). In
case the ribosome remains associated with the mRNA, the three downstream open
reading frames are all out-of-frame with *baf-1* (purple lines). In the second strategy
(bottom), the GFP cassette contains a single FRT and a second FRT is inserted in
the 3'-UTR of the gene of interest. FLP-mediated recombination leads to complete
gene removal. (B) Brood sizes were determined for control animals (*g>f>p::baf-1* and
mCh::baf-1; nhx-2p::FLP) and for animals with conditional KO of *baf-1* in the
intestine (*g>f>p::baf-1; nhx-2p::FLP* [first strategy] and *gf>p::baf-1>; nhx-2p::FLP*
[second strategy]; strains BN580, BN1182 and BN1223). Each point represents the
number of progenies produced by an individual worm; *n* for each genotype is
indicated in parentheses. P values from pair-wise t tests and adjusted for multiple
comparisons (Benjamini & Hochberg method) are indicated. In these and
subsequent boxplots, the midline represents the median and the upper and lower

limits of the box correspond to the third and first quartile, respectively. The whiskers
extend to the lowest and higher observations but maximum up to 1.5 times the
interquartile range.

Figure 3. KO of GFP::MEL-28 in hypodermal nuclei. (A) Confocal micrographs of
g>f>p::mel-28 L1 larvae, L2 larvae and young adults either not expressing FLP (top
panels; strain BN793) or expressing FLP in the hypodermis either without (middle
panels; *dpy-7p::FLP* driver; strain BN1121) or with GFPdeg in the hypodermis
(bottom panels; *dpy-7p::FLP* driver; strain BN1122). A single focal plane of the
hypodermis is shown. Scale bar 10 µm. (B) Quantification of GFP::MEL-28 at the
nuclear envelope of hypodermal nuclei. Each point represents a single nucleus. Data
were acquired from at least 10 animals in two independent experiments; *n* for each
genotype is indicated in parentheses. P values from Kruskal-Wallis rank sum tests
and adjusted with Dunn's test for multiple comparisons (Benjamini & Hochberg
method) are indicated.

Figure 4. Degradation of GFP::MEL-28 in intestinal cells. (A) Schematic
representation of the GFPdeg strategy. The vhhGFP4::ZIF-1 fusion protein targets
GFP-tagged proteins for rapid degradation. Temporal and spatial control is provided
by the *hsp-16.41* promoter and tissue-specific FLP expression, respectively. (B)
Quantification of GFP::MEL-28 at the nuclear envelope of hypodermal nuclei of
young adults either without the GFPdeg system (Control; strain BN452) or with the
GFPdeg system but without induction (GFPdeg; strain BN1084). P value from t test
is indicated. (C) Quantification of GFP::MEL-28 at the nuclear envelope of anterior
intestinal nuclei of young adults after 1 or 2 rounds of GFPdeg induction in the
intestine (*nhx-2p::FLP* driver; strain BN1118). Control (strain BN452) and constitutive
degradation (*elt-2p::vhhGFP4::zif-1*; strain BN746) are included for comparison.
Graph details are explained in Figure 3. P values from pair-wise t tests and adjusted
for multiple comparisons (Benjamini & Hochberg method) are indicated. (D) Confocal
micrographs of *gfp::mel-28* young adults without GFPdeg (top panels; strain BN452),
inducible intestinal GFPdeg (middle panels; strain BN1118) or constitutive intestinal
GFPdeg (bottom panels; strain BN746). A single focal plane of the anterior pair of
intestinal nuclei is shown. Scale bar 10 µm.

Figure 5. Kinetics of GFP::MEL-28 degradation in hypodermal cells. (A) Confocal
micrographs of *gfp::mel-28* young adults either without induction or 1-24 h after
induction of GFPdeg in the hypodermis (*dpy-7p::FLP* driver; strain BN1084). A single
focal plane of the hypodermis is shown. Scale bar 10 µm. (B) Quantification of
GFP::MEL-28 at the nuclear envelope of hypodermal nuclei. Graph details are
explained in Figure 3.

Figure 6. Efficient protein depletion by combined gene KO and GFPdeg. (A)
Schematic representation of the combined KO and GFPdeg strategy. A single FLP
driver controls in a tissue-specific manner both KO (left) and protein degradation
(right) of a gene tagged with the GFP KO cassette. (B) Confocal micrographs of
g>f>p::npp-2 L4 larvae either not expressing FLP (left panel; strain BN1082),
expressing FLP in the hypodermis (middle panel; *dpy-7p::FLP* driver; strain BN1111)
or expressing FLP and vhhGFP4::ZIF-1 in the hypodermis (right panel; *dpy-7p::FLP*
driver; strain BN1086). A single focal plane of the hypodermis is shown. Scale bar 10
µm. (C) Quantification of GFP::NPP-2 at the nuclear envelope of hypodermal nuclei.
Graph details are explained in Figure 3. (D) Survival curves for animals expressing
GFP::NPP-2 alone (blue; strain BN1082) or combined with either FLP in the

hypodermis (dark red; strain BN1111) or FLP and GFPdeg in the hypodermis (red;
strain BN1086). A strain with hypodermal FLP and GFPdeg but without target was
included as control (green; BN1097). The lifespan of the two strains with depletion of
GFP::NPP-2 in the hypodermis was significantly shorter compared to the two control
strains (*** indicates p<1e-08). More than 175 animals were analysed for each
genotype.

Figure 7. Tissue-specific identification of transcribed genes by RAPID. (A) Schematic
representation of the RAPID method. Spatial control is provided by tissue-specific
FLP expression that leads to excision of the *mCherry::his-58* cassette and basal
transcription of *Dam::rpb-6* from the *hsp-16.41* promoter. Dam::RPB-6 incorporates
into RNA polymerases and marks GATC sites in transcribed genes. A *GFP::Dam*
transgene is used in parallel to control for differences in chromatin accessibility and
unspecific methylation (not shown). (B-E) Data based on two replicas of L4s
expressing either Dam::RPB-6 (strain BN1414) or GFP::Dam (strain BN561) in the
hypodermis. The analysis was restricted to 19,975 protein coding genes annotated in
WBCel235. (B) RAPID signal (blue) averaged in 10 bp bins across all genes (left
graph) or across 2331 expressed genes (right graph) from 2 kb upstream of the
transcriptional start site (TSS) to 2 kb downstream of the transcriptional end site
(TES). The signal from TSS to TES was extrapolated to an arbitrary length of 3 kb
for each gene. The distribution of the active chromatin mark H3K36me3 (BANNISTER
et al. 2005) is included for comparison. The heatmap (middle graph) represents the
individual genes from the left graph. (C) Representation of Dam::RPB-6/GFP::Dam
ratios for individual replicas (Rep 1, Rep 2) at selected loci. Genes with significant
RAPID signal are indicated (FDR<0.05). (D-E) Comparison of the 2331 transcribed
genes identified by RAPID with three alternative methods for transcriptomic profiling
(see main text for details). Numbers in parentheses indicate the number of protein
coding genes reported in each study.

Supplementary Figure S1. Characterization of novel FLP drivers. Upon temperature
induction of the *hsp-16.41* promoter, the reporter expresses red (mCh) or green
(GFP) histone HIS-58 in cell lineages without or with FLP-mediated recombination,
respectively. (A) Recombination activity by *lin-31p::FLP::SL2::mNG* is observed in
the head (left panel) and central region (right panel) of BN1023 animals. Bright
GFP::HIS-58 expression is observed in the nuclei of cells forming the vulva (filled
arrows), whereas faint expression is detected in the hypodermis (open arrows; right
panel). Presumably, the faint expression is derived from Pn.p cells that fuse with
hyp7. (B) Recombination activity by *gpa-14p::FLP::SL2::mNG* is observed in the
head (left panel) and tail (right panel) of BN1208 animals. Filled arrows point to
nuclei expressing GFP::HIS-58 but not mCh::HIS-58 (i.e. recombination of both
alleles); a double arrow points to a nucleus expressing both fluorescent proteins (i.e.
recombination of one allele) and open arrows indicate cells that express diffuse mNG
(and hence also FLP) but not GFP::HIS-58. (C) Expression of mNG from the *unc-*
17p::FLP::SL2::mNG transgene overlaps with mKate2 signal from endogenously
tagged *unc-17* (lower left panel) and recombination (GFP::HIS-58) is detected as
expected in multiple neurons in the head (upper left panel) and ventral nerve cord
(upper right panel) of BN1133 animals. Unexpectedly, GFP::HIS-58 is also observed
in hypodermal nuclei (lower right panel). (D) Recombination activity by *hlh-*
12p::FLP::SL2::mTagBFP2 is observed in the distal tip cell of BN1319 animals. The
distal tip cell is marked by co-expressed mTagBFP2. Scale bars 10 µm.

Supplementary Figure S2. Overlap between protein coding gene sets identified by
TaDa in seam cells and hypodermis (KATSANOS *et al.* 2021) and by RAPID in
hypodermis (this study).

References

Altun, Z. F., and D. H. Hall, 2009 Coelomocyte system, pp. in *WormAtlas.*
Armenti, S. T., L. L. Lohmer, D. R. Sherwood and J. Nance, 2014 Repurposing an
endogenous degradation system for rapid and targeted depletion of C.
elegans proteins. Development 141**:** 4640-4647.
Arribere, J. A., H. Kuroyanagi and H. A. Hundley, 2020 mRNA Editing, Processing
and Quality Control in Caenorhabditis elegans. Genetics 215**:** 531-568.
Askjaer, P., and J. C. Harr, 2021 Genetic approaches to revealing the principles of
nuclear architecture. Curr Opin Genet Dev 67**:** 52-60.
Ayuso, C., and P. Askjaer, 2019 Spatiotemporal control of genome recombination
through combined FLP-Frt and GAL4-UAS technologies. MicroPubl Biol 2019.
Bannister, A. J., R. Schneider, F. A. Myers, A. W. Thorne, C. Crane-Robinson *et al.*,
2005 Spatial distribution of di- and tri-methyl lysine 36 of histone H3 at active
genes. J Biol Chem 280**:** 17732-17736.
Bardou, P., J. Mariette, F. Escudie, C. Djemiel and C. Klopp, 2014 jvenn: an
interactive Venn diagram viewer. BMC Bioinformatics 15**:** 293.
Bhoi, A., F. Palladino and P. Fabrizio, 2021 Auxin confers protection against ER
stress in Caenorhabditis elegans. Biol Open 10.
Blazie, S. M., H. C. Geissel, H. Wilky, R. Joshi, J. Newbern *et al.*, 2017 Alternative
Polyadenylation Directs Tissue-Specific miRNA Targeting in Caenorhabditis
elegans Somatic Tissues. Genetics 206**:** 757-774.
Cabianca, D. S., C. Munoz-Jimenez, V. Kalck, D. Gaidatzis, J. Padeken *et al.*, 2019
Active chromatin marks drive spatial sequestration of heterochromatin in C.
elegans nuclei. Nature 569**:** 734-739.
Caussinus, E., O. Kanca and M. Affolter, 2011 Fluorescent fusion protein knockout
mediated by anti-GFP nanobody. Nat Struct Mol Biol 19**:** 117-121.
Cohen, S., L. Kramarski, S. Levi, N. Deshe, O. Ben David *et al.*, 2019 Nonsense
mutation-dependent reinitiation of translation in mammalian cells. Nucleic
Acids Res 47**:** 6330-6338.
Davis, L., I. Radman, A. Goutou, A. Tynan, K. Baxter *et al.*, 2021 Precise optical
control of gene expression in C elegans using improved genetic code
expansion and Cre recombinase. Elife 10.
Davis, M. W., J. J. Morton, D. Carroll and E. M. Jorgensen, 2008 Gene activation
using FLP recombinase in C. elegans. PLoS Genet 4**:** e1000028.
de la Cruz Ruiz, P., R. Romero-Bueno and P. Askjaer, 2022 Analysis of Nuclear
Pore Complexes in Caenorhabditis elegans by Live Imaging and Functional
Genomics. Methods Mol Biol 2502**:** 161-182.
Dobrzynska, A., P. Askjaer and Y. Gruenbaum, 2016 Lamin-Binding Proteins in
Caenorhabditis elegans. Methods Enzymol 569**:** 455-483.
Driesschaert, B., L. Mergan and L. Temmerman, 2021 Conditional gene expression
in invertebrate animal models. J Genet Genomics 48**:** 14-31.
Fernandez, A. G., and F. Piano, 2006 MEL-28 is downstream of the Ran cycle and is
required for nuclear-envelope function and chromatin maintenance. Curr Biol
16**:** 1757-1763.

Friedland, A. E., Y. B. Tzur, K. M. Esvelt, M. P. Colaiacovo, G. M. Church *et al.*, 2013 Heritable genome editing in C. elegans via a CRISPR-Cas9 system. Nat Methods 10**:** 741-743.

Frokjaer-Jensen, C., M. W. Davis, M. Ailion and E. M. Jorgensen, 2012 Improved Mos1-mediated transgenesis in C. elegans. Nat Methods 9**:** 117-118.

Frokjaer-Jensen, C., M. W. Davis, C. E. Hopkins, B. J. Newman, J. M. Thummel *et al.*, 2008 Single-copy insertion of transgenes in Caenorhabditis elegans. Nat Genet 40**:** 1375-1383.

Frokjaer-Jensen, C., M. W. Davis, M. Sarov, J. Taylor, S. Flibotte *et al.*, 2014 Random and targeted transgene insertion in Caenorhabditis elegans using a modified Mos1 transposon. Nat Methods 11**:** 529-534.

Galy, V., P. Askjaer, C. Franz, C. Lopez-Iglesias and I. W. Mattaj, 2006 MEL-28, a novel nuclear-envelope and kinetochore protein essential for zygotic nuclear-envelope assembly in C. elegans. Curr Biol 16**:** 1748-1756.

Ge, M. H., W. Wang, T. H. Wu, X. Wen, U. Al-Sheikh *et al.*, 2020 Dual Recombining-out System for Spatiotemporal Gene Expression in C. elegans. iScience 23**:** 101567.

Gomez-Saldivar, G., A. Fernandez, Y. Hirano, M. Mauro, A. Lai *et al.*, 2016 Identification of Conserved MEL-28/ELYS Domains with Essential Roles in Nuclear Assembly and Chromosome Segregation. PLoS Genet 12**:** e1006131.

Gomez-Saldivar, G., J. Osuna-Luque, J. I. Semple, D. A. Glauser, S. Jarriault *et al.*, 2020 Tissue-Specific Transcription Footprinting Using RNA Pol DamID (RAPID) in Caenorhabditis elegans. Genetics.

Gorjanacz, M., E. P. Klerkx, V. Galy, R. Santarella, C. Lopez-Iglesias *et al.*, 2007 Caenorhabditis elegans BAF-1 and its kinase VRK-1 participate directly in post-mitotic nuclear envelope assembly. EMBO J 26**:** 132-143.

Haenni, S., Z. Ji, M. Hoque, N. Rust, H. Sharpe *et al.*, 2012 Analysis of C. elegans intestinal gene expression and polyadenylation by fluorescence-activated nuclei sorting and 3'-end-seq. Nucleic Acids Res 40**:** 6304-6318.

Harr, J. C., C. D. Schmid, C. Munoz-Jimenez, R. Romero-Bueno, V. Kalck *et al.*, 2020 Loss of an H3K9me anchor rescues laminopathy-linked changes in nuclear organization and muscle function in an Emery-Dreifuss muscular dystrophy model. Genes Dev.

Hills-Muckey, K., M. A. Q. Martinez, N. Stec, S. Hebbar, J. Saldanha *et al.*, 2021 An engineered, orthogonal auxin analog/AtTIR1(F79G) pairing improves both specificity and efficacy of the auxin degradation system in Caenorhabditis elegans. Genetics.

Hubbard, E. J., 2014 FLP/FRT and Cre/lox recombination technology in C. elegans. Methods 68**:** 417-424.

Hubbard, E. J., and D. Greenstein, 2005 Introduction to the germ line. WormBook**:** 1-4.

Kage-Nakadai, E., R. Imae, Y. Suehiro, S. Yoshina, S. Hori *et al.*, 2014 A conditional knockout toolkit for Caenorhabditis elegans based on the Cre/loxP recombination. PLoS One 9**:** e114680.

Kaletsky, R., V. Yao, A. Williams, A. M. Runnels, A. Tadych *et al.*, 2018 Transcriptome analysis of adult Caenorhabditis elegans cells reveals tissue-specific gene and isoform expression. PLoS Genet 14**:** e1007559.

Kassambara, A., 2020 ggpubr: 'ggplot2' Based Publication Ready Plots, pp.

Katsanos, D., M. Ferrando-Marco, I. Razzaq, G. Aughey, T. D. Southall *et al.*, 2021
Gene expression profiling of epidermal cell types in C. elegans using
Targeted DamID. Development 148.
Kimble, J., and H. Seidel, 2008 C. elegans germline stem cells and their niche in
StemBook, Cambridge (MA).
Konietzka, J., M. Fritz, S. Spiri, R. McWhirter, A. Leha *et al.*, 2020 Epidermal Growth
Factor Signaling Promotes Sleep through a Combined Series and Parallel
Neural Circuit. Curr Biol 30**:** 1-16 e13.
Kroetz, M. B., and D. Zarkower, 2015 Cell-Specific mRNA Profiling of the
Caenorhabditis elegans Somatic Gonadal Precursor Cells Identifies Suites of
Sex-Biased and Gonad-Enriched Transcripts. G3 (Bethesda) 5**:** 2831-2841.
Loria, P. M., J. Hodgkin and O. Hobert, 2004 A conserved postsynaptic
transmembrane protein affecting neuromuscular signaling in Caenorhabditis
elegans. J Neurosci 24**:** 2191-2201.
Ma, X., G. Zhan, M. C. Sleumer, S. Chen, W. Liu *et al.*, 2016 Analysis of C. elegans
muscle transcriptome using trans-splicing-based RNA tagging (SRT). Nucleic
Acids Res 44**:** e156.
Macias-Leon, J., and P. Askjaer, 2018 Efficient FLP-mediated germ-line
recombination in C. elegans. MicroPubl Biol 2018.
Marshall, O. J., and A. H. Brand, 2015 damidseq_pipeline: an automated pipeline for
processing DamID sequencing datasets. Bioinformatics 31**:** 3371-3373.
Marshall, O. J., T. D. Southall, S. W. Cheetham and A. H. Brand, 2016 Cell-type-
specific profiling of protein-DNA interactions without cell isolation using
targeted DamID with next-generation sequencing. Nat Protoc 11**:** 1586-1598.
McGhee, J. D., 2007 The C. elegans intestine. WormBook**:** 1-36.
Monsalve, G. C., K. R. Yamamoto and J. D. Ward, 2019 A New Tool for Inducible
Gene Expression in Caenorhabditis elegans. Genetics 211**:** 419-430.
Munoz-Jimenez, C., C. Ayuso, A. Dobrzynska, A. Torres-Mendez, P. C. Ruiz *et al.*,
2017 An Efficient FLP-Based Toolkit for Spatiotemporal Control of Gene
Expression in Caenorhabditis elegans. Genetics 206**:** 1763-1778.
Nance, J., and C. Frokjaer-Jensen, 2019 The Caenorhabditis elegans Transgenic
Toolbox. Genetics 212**:** 959-990.
Nehrke, K., and J. E. Melvin, 2002 The NHX family of Na+-H+ exchangers in
Caenorhabditis elegans. J Biol Chem 277**:** 29036-29044.
Nonet, M. L., 2020 Efficient Transgenesis in Caenorhabditis elegans Using Flp
Recombinase-Mediated Cassette Exchange. Genetics 215**:** 903-921.
Packer, J. S., Q. Zhu, C. Huynh, P. Sivaramakrishnan, E. Preston *et al.*, 2019 A
lineage-resolved molecular atlas of C. elegans embryogenesis at single-cell
resolution. Science.
Pereira, L., P. Kratsios, E. Serrano-Saiz, H. Sheftel, A. E. Mayo *et al.*, 2015 A
cellular and regulatory map of the cholinergic nervous system of C. elegans.
Elife 4.
Porta-de-la-Riva, M., A. C. Gonzalez, N. Sanfeliu-Cerdán, S. Karimi, S. González-
Bolívar *et al.*, 2021 Deploying photons for communication within neuronal
networks. bioRxiv**:** 2021.2008.2002.454613.
Reece-Hoyes, J. S., J. Shingles, D. Dupuy, C. A. Grove, A. J. Walhout *et al.*, 2007
Insight into transcription factor gene duplication from Caenorhabditis elegans
Promoterome-driven expression patterns. BMC Genomics 8**:** 27.
Rodenas, E., C. Gonzalez-Aguilera, C. Ayuso and P. Askjaer, 2012 Dissection of the
NUP107 nuclear pore subcomplex reveals a novel interaction with spindle

assembly checkpoint protein MAD1 in Caenorhabditis elegans. Mol Biol Cell 23**:** 930-944.

Ruijtenberg, S., and S. van den Heuvel, 2015 G1/S Inhibitors and the SWI/SNF Complex Control Cell-Cycle Exit during Muscle Differentiation. Cell 162**:** 300-313.

Sauer, B., 1987 Functional expression of the cre-lox site-specific recombination system in the yeast Saccharomyces cerevisiae. Mol Cell Biol 7**:** 2087-2096.

Savas, J. N., B. H. Toyama, T. Xu, J. R. Yates, 3rd and M. W. Hetzer, 2012 Extremely long-lived nuclear pore proteins in the rat brain. Science 335**:** 942.

Schindelin, J., I. Arganda-Carreras, E. Frise, V. Kaynig, M. Longair *et al.*, 2012 Fiji: an open-source platform for biological-image analysis. Nat Methods 9**:** 676-682.

Schmitt, C., C. Schultheis, N. Pokala, S. J. Husson, J. F. Liewald *et al.*, 2012 Specific expression of channelrhodopsin-2 in single neurons of Caenorhabditis elegans. PLoS One 7**:** e43164.

Schwartz, M. L., and E. M. Jorgensen, 2016 SapTrap, a Toolkit for High-Throughput CRISPR/Cas9 Gene Modification in Caenorhabditis elegans. Genetics 202**:** 1277-1288.

Senecoff, J. F., R. C. Bruckner and M. M. Cox, 1985 The FLP recombinase of the yeast 2-micron plasmid: characterization of its recombination site. Proc Natl Acad Sci U S A 82**:** 7270-7274.

Sharma, R., D. Ritler and P. Meister, 2016 Tools for DNA adenine methyltransferase identification analysis of nuclear organization during C. elegans development. Genesis 54**:** 151-159.

Southall, T. D., K. S. Gold, B. Egger, C. M. Davidson, E. E. Caygill *et al.*, 2013 Cell-type-specific profiling of gene expression and chromatin binding without cell isolation: assaying RNA Pol II occupancy in neural stem cells. Dev Cell 26**:** 101-112.

Steiner, F. A., P. B. Talbert, S. Kasinathan, R. B. Deal and S. Henikoff, 2012 Cell-type-specific nuclei purification from whole animals for genome-wide expression and chromatin profiling. Genome Res 22**:** 766-777.

Stempor, P., and J. Ahringer, 2016 SeqPlots - Interactive software for exploratory data analyses, pattern discovery and visualization in genomics. Wellcome Open Res 1**:** 14.

Stiernagle, T., 2006 Maintenance of C. elegans. WormBook**:** 1-11.

Tamai, K. K., and K. Nishiwaki, 2007 bHLH transcription factors regulate organ morphogenesis via activation of an ADAMTS protease in C. elegans. Dev Biol 308**:** 562-571.

Tan, P. B., M. R. Lackner and S. K. Kim, 1998 MAP kinase signaling specificity mediated by the LIN-1 Ets/LIN-31 WH transcription factor complex during C. elegans vulval induction. Cell 93**:** 569-580.

Team, R., 2020a RStudio: Integrated Development Environment for R, pp. RStudio, PBC.

Team, R. C., 2020b R: A Language and Environment for Statistical Computing, pp. R Foundation for Statistical Computing.

Therneau, T., 2020 A Package for Survival Analysis in R, pp.

Thorvaldsdottir, H., J. T. Robinson and J. P. Mesirov, 2013 Integrative Genomics Viewer (IGV): high-performance genomics data visualization and exploration. Brief Bioinform 14**:** 178-192.

Toyama, B. H., J. N. Savas, S. K. Park, M. S. Harris, N. T. Ingolia *et al.*, 2013
Identification of long-lived proteins reveals exceptional stability of essential
cellular structures. Cell 154**:** 971-982.
Van de Walle, P., C. Munoz-Jimenez, P. Askjaer, L. Schoofs and L. Temmerman,
2020 DamID identifies targets of CEH-60/PBX that are associated with neuron
development and muscle structure in Caenorhabditis elegans. PLoS One 15**:**
e0242939.
Voutev, R., and E. J. Hubbard, 2008 A "FLP-Out" system for controlled gene
expression in Caenorhabditis elegans. Genetics 180: 103-119.
Wang, S., N. H. Tang, P. Lara-Gonzalez, Z. Zhao, D. K. Cheerambathur *et al.*, 2017
A toolkit for GFP-mediated tissue-specific protein degradation in C. elegans.
Development 144**:** 2694-2701.
Watts, J. S., H. F. Harrison, S. Omi, Q. Guenthers, J. Dalelio *et al.*, 2020 New
Strains for Tissue-Specific RNAi Studies in Caenorhabditis elegans. G3
(Bethesda) 10**:** 4167-4176.
Wickham, H., 2016 *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag
New York.
Widmer, Y. F., A. Bilican, R. Bruggmann and S. G. Sprecher, 2018 Regulators of
Long-Term Memory Revealed by Mushroom Body-Specific Gene Expression
Profiling in Drosophila melanogaster. Genetics 209**:** 1167-1181.
Wurmthaler, L. A., M. Sack, K. Gense, J. S. Hartig and M. Gamerdinger, 2019 A
tetracycline-dependent ribozyme switch allows conditional induction of gene
expression in Caenorhabditis elegans. Nat Commun 10**:** 491.
Zhang, L., J. D. Ward, Z. Cheng and A. F. Dernburg, 2015 The auxin-inducible
degradation (AID) system enables versatile conditional protein depletion in C.
elegans. Development 142**:** 4374-4384.

1154

A

Frame shift and nonsense-mediated mRNA decay

Complete gene knockout

B

Figure 3

A

B

A

hsp-16.41p mCherry his-58 3'UTR Dam rpb-6 3'UTR

Frt Frt

+ FLP

basal expression

Dam::RPB-6

Dam::RPB-6

RNA polymerase

Tissue-specific labeling of transcribed genes (RNA polymerase DamID = RAPID)

B

All genes

RAPID (all genes)

Expressed genes

RAPID
H3K36me3

RAPID
H3K36me3

C

col-39; col-79; col-80

eef-1A.1; F31E.12

sqt-1

fib-1; T01C3.11

D

TaDa (2657)   PATseq (5984)

RAPID (2331)   FACS (7145)

E

In common with Fraction of	RAPID	TaDa	FACS	PATseq
RAPID genes		54%***	66%***	59%***
TaDa genes	48%***		78%***	72%***
FACS genes	21%***	29%***		64%***
PATseq genes	23%***	32%***	77%***	