Biofilm matrix proteome of clinical strain of *P. aeruginosa*

Daria A. Egorova [1],*, Andrey I. Solovyev [1], Nikita B. Polyakov [1], Ksenya V. Danilova [1], Anastasya A. Scherbakova [1], Ivan N. Kravtsov [1], Maria A. Dmitrieva [1], Valentina S. Rykova [1], Irina L. Tutykhina [1], Yulia M. Romanova [1, 2], Alexander L. Gintsburg [1, 2]

[1] National Research Center of Epidemiology and Microbiology n. a. N.F. Gamaleya, Russian Ministry of Health, Moscow, 123098 Russia; info@gamaleya.org
[2] I.M. Sechenov First Moscow State Medical University, Moscow, 119992 Russia; international@staff.sechenov.ru
* Correspondence: egorova.daria@gamaleya.org

Abstract: Extracellular matrix plays a pivotal role in biofilm biology. Despite the importance of matrix proteins as potential targets for the development of antibacterial therapeutics little is known about matrix proteomes. As matrix responsible for some extracellular functions, it is expected that protein composition should be different compared to embedded in biofilm cells. This difference reflects possible active processes in the matrix. While *P. aeruginosa* is one of the most important pathogens with emerging antibiotic resistance only a few studies are devoted to matrix proteomes and there are no studies describing matrix proteome for any clinical isolates. Here we report the first matrix proteome in comparison with embedded cells for the clinical isolate of *P. aeruginosa*. We have identified the largest number of proteins in the matrix among all published studies devoted to *P. aeruginosa* biofilms. Functional classification of enriched in matrix proteins resulted in several bioprocess groups. Bioprocess groups with the largest number of overrepresented in matrix proteins were oxidation-reduction processes, proteolysis, and transmembrane transport. The top three represented in matrix bioprocesses concerning the size of the GO annotated database were: cell redox homeostasis, nucleoside metabolism, and fatty acid synthesis. Finally, we discuss obtained data in a prism of possible directions for antibiofilm therapeutics development.

Keywords: biofilm; matrix; proteome; matrixome; *Pseudomonas aeruginosa*; nucleoside metabolism; cell redox homeostasis; fatty acid synthesis, proteolysis; transmembrane transport

1. Introduction

Biofilms are the most common lifestyle of microorganisms including both pathogenic and environmental bacterial species. From a clinical perspective, biofilms are often represented by microbial aggregates with tolerance to host defense mechanisms and antimicrobials, so they cause difficult-to-treat recurrent diseases. Currently, there is an urgent need to discover new targets and strategies to overcome the tolerance for the effective treatment of biofilm-associated infections.

The key feature of biofilms is the presence of an extracellular matrix that covers all members of biofilm and creates a microenvironment for communication, protects against different threats, provides an opportunity for spatial organization and functional diversification within the community. The matrix consists of a broad range of biopolymers, metabolites, and signal molecules. Also, it usually includes organized compartments like outer membrane vehicles (OMVs). To stress the idea of rich and complex matrix organization the term «matrixome» was proposed [1]. Proteins in the biofilm matrix might be actively secreted, passively leaked from cells, and entrapped into the matrix from the environment [2, 3]. Proteins in the matrix have a structural role in maintaining biofilm organization, play a role as a protective barrier, creates microenvironment with limited diffusion. The barrier role is somewhere similar to structural function but also includes hydrophobic features, charge, and so on rather than just being mechanically stable structure. Moreover, matrix proteins may bind antimicrobial molecules and reduce their diffusion and effect on bacterial cells i. e. extracellular ribosomal proteins bind antibiotics and prevent their penetration in bacterial cells. Other functions of matrix proteins rely on their enzymatic activity and include degradation of biopolymers, participation in biochemical processes, and signaling function.

Biofilms are dynamic communities with lifecycle consisting of attachment, maturation, and dispersion stages. During the maturation stage matrix may serve as depo for virulence factors which then come out together with dispersed bacteria. In the case of chronic infections dispersion stage is associated with the recurrence of symptomatic infections and colonization of new sites in the body.

Importantly bacteria inside biofilm have diverse phenotypes and some extracellular molecules (public goods) are consumed by non-producers. Matrix in general and some components in matrix are examples of public goods. So, a part of bacterial population, especially inside multispecies biofilms, may not have resistance mechanisms, but due to matrix still be irresponsible for the treatment.

Targeting of matrix proteins as a strategy to combat infections may act in the following ways: (1) destabilization/disruption of biofilm matrix to improve the action of host defense and/or antibiotic therapy; (2) targeting of extracellular biochemical processes to reduce overall biofilm succeed in survival and virulence; (3) targeting of extracellular virulence factors associated with biofilm matrix to decrease virulence in a case of mass dispersion.

Despite the accepted idea of a pivotal role of matrix in bacterial biofilms little is known about the protein composition of the matrix at a level of proteomes. Moreover, for the well-studied and clinically important biofilm-forming bacteria *P. aeruginosa* there are only a few studies devoted to matrix proteome of reference strains (mainly PAO1, Table 1), while proteomics of the whole biofilm is better described [4, 5].

Table 1. Studies devoted to biofilm matrix proteome. Biofilm`s compartments and MS methods were named by original publications.

Bacteria	Biofilm compartments	Method	Number of proteins	Reference
P. aeruginosa **PAO1**	**Matrix** OMVs	Gel-based vs Gel-free 2D LC-MS/MS	**Matrix 327** OMVs 207	[9]
P. aeruginosa **PAO1**	**Matrix** Total biofilm	LC-MS/MS	Total biofilm 857 **Matrix 60**	[8]
P. aeruginosa **PAO1**	**Matrix** OMVs Cells	1D-SDS-PAGE combined with nano-LC-ESI-MS/MS	**Matrix 178** OMVs 57 Cells 764	[7]
P. aeruginosa **PAO1 at different time points**	Cells OMVs	LC-MS/MS	Cells 2443 in total OMVs 1142 in total	[10]
P. aeruginosa **ATCC27853 at different time points**	**Matrix**	iTRAQ-labeled peptides and LC-MS/MS analysis	**Matrix** 389 in total	[11]
Dynamics in dual-species biofilm – *P. aeruginosa* **PAO1 and** *S. aureus* **ATCC 25923**	Surfaceome Exoproteome	Orbitrap QExactive Plus LC-MS/MS	Surfaceome by PAO1 495 Exoproteome by PAO1 762	[12]
P. aeruginosa **CF clinical isolate KB6**	Cells **Matrix**	Orbitrap LC-MS/MS	Cells 1652 **Matrix 957**	This study
S. aureus **UAMS-1** *in vivo* **chronic implant infection**	Secretome Surfactome	GeLC-MS/MS	Secretome 33 Surfactome 72	[13]
S. aureus **USA300 CA-MRSA strain LAC**	DNA-binding proteins in the biofilm matrix	quadrupole time of flight (Q/TOF) mass spectrometer (Agilent 6520) with a nanospray ionization source	49	[14]
S. aureus **HG001**	intracellular extracellular (ECM and flow-through)	LC-MS/MS on LTQ-Orbitrap-Velos mass spectrometer coupled to an EASY-nLC 1000	ECM 1407 flow-through 1400 intracellular 1621	[63]
B. multivorans **C1575**	Matrix	Gel-based LC-MS/MS	Matrix 161	[15]

	OMVs		OMVs	64	
S. acidocaldarius	EPS	nanoRSLC-Orbitrap LC-MS/MS	85		[16]
***Shewanella sp.* HRCR-1**	EPS	LC-MS/MS	58		[17]
Cutibacterium acnes	Matrix	Orbitrap MS	447		[18]
Vibrio cholerae	Matrix	LC electrospray ionization and then entered into an LTQ linear ion-trap mass spectrometer (ThermoFisher)	74		[19]
Haemophilus influenzae (NTHi)	ECM	Gel-based LC/MS/MS	60		[20]

In a comparative study of liquid culture of PAO1 strain and clinical isolates, it was shown that cystic fibrosis (CF) isolates expressed a narrower range of transporters and a broader set of enzymes of metabolic pathways for the biosynthesis of amino acids, carbohydrates, nucleotides, and polyamines, but this study did not cover biofilm mode of life as well as extracellular matrix composition [6]. Only one study described the proteome of the matrix in comparison with embedded cells [7] and one else study compared the matrix with the total biofilm proteome [8], both studies were devoted to reference strain PAO1. The gap in understanding the difference in protein composition between matrix and embedded cells frustrates the development of antibiofilm therapeutics and the overall understanding of biofilm biology. Here we performed a proteomic study of matrix composition in comparison with embedded cells for the clinical strain of *P. aeruginosa* to identify bioprocesses taking place in the matrix as probable targets or as factors or as barriers during pharmacological development of antibiofilm therapeutics.

2. Results

3.1. General overview of proteomes

P. aeruginosa KB6 (exoT+; exoY+; exoU-; exoS+) is a clinical isolate from bronchoalveolar lavage of cystic fibrosis patient with strong biofilm-forming phenotype [21]. To investigate protein composition of extracellular biofilm matrix we used previously published method of separation of biofilm matrix from embedded cells with high ionic solution of NaCl [29, 30]. To confirm the absence of cell lysis for KB6 strain during matrix separation we evaluated CFU number after NaCl treatment for both biofilm and suspension bacterial culture. There was no difference in the recovery of CFU with and without NaCl treatment for suspension bacteria, while CFU number for biofilm samples was comparable after NaCl treatment and after common biofilm-disruption approach – mild sonication for 5 minutes (data not shown). So, matrix isolation with NaCl was used for protein sample preparation. For the proteomic study, we have grown static biofilm for 18 h in LB medium. This time point corresponds to biofilm in the early stationary phase when the number of embedded bacterial cells reaches the plateau, and the matrix is already formed. Further increase in biofilm biomass occurs mainly due to mass of extracellular components rather than increasing number of bacterial cells. During the further biofilm growth mass of extracellular substance significantly impacted with lysed cells. To decrease the number of proteins that may represent passive cell leakage or death during the biofilm stationary phase and to avoid interference of these «archeological» proteins with secreted extracellular proteins we choose this early time point (18 h).

Two independent and separated in time biofilms were grown in a liquid medium. Extracellular matrix and embedded cells were separated with NaCl and processed for protein isolation. Also, one more biological replicate (the third) representing embedded cells only was added to the analysis from another set of experiments (data not shown). Proteomic analysis was performed for each sample in three technical replicates. For protein identification, *P. aeruginosa* KB6 strain-specific protein dataset was created based on genome sequence (available in GenBank under accession number NZ_CP034429). Protein identification was made with MSFragger. The full list of all identified proteins is available in Table S1.

In total, more than 1600 proteins were identified in all samples. After initial manual inspection of LFQ intensities we observed matrix-specific proteins, cell-specific proteins, and proteins with presence in both compartments. While an inspection of intensity signals is not applicable for comparison of the representation of different proteins it is still gave us an additional consciousness about proper separation

matrix from cells. The absence of several intracellular proteins in matrix samples confirms that our approach to separate matrix from cells does not cause significant cell leakage during sample preparation. For example, S18, L29, L33, S15, S20 proteins involved in ribosome assembly and function were absent in the matrix (LFQ intensities and spectral counts were zero in all matrix samples), but these proteins were observed in all cell's samples with LFQ intensities more than $1*10^8$. So, in general, the absence of this ribosomal and some other proteins in matrix samples is associated neither with cell disruption during matrix separation nor with low detection efficacy as a specific feature for some proteins.

Spearman correlation coefficient for proteins LFQ intensities between cells and matrix was 0.4959 (95% CI 0,4555 to 0,5343) (correlation matrix is presented in Figure 1).

Figure 1. Correlation of individual protein intensities (LFQ) between embedded cells and matrix.

While the correlation was expectedly positive there were several examples of outfitters like lytic murein transglycosylase with well-known localization on the outer surface of the bacterial cell wall [31].

Cells and matrix differ in overall biomolecule content, physicochemical and other properties. That was our premise for rigorous statistical analysis of comparative protein representation. For getting a quantitative comparison of the representation of proteins we proceeded to logarithm transformation and quantile normalization of LFQ intensities. Statistical analysis was made with the Limma R package [32]. All identified proteins are listed in table S1, and proteins differing in their representation in cells and matrix are listed in Table S2. For fold difference presentation volcano plot was created (Figure 2).

Figure 2. Volcano plot of representation of proteins between cells and matrix. A – overall protein representation;
B – an enlarged area with proteins with fold change between Log2 from 1 to 10. Red color indicates proteins with fold
change more than 2 and significance value p<0.01; green color indicates proteins with fold change more than 2 and
significance value p>0.01; blue color indicates proteins with fold change less than 2 but significance value p<0.01; grey
color indicates proteins with fold change less than 2 and significance value p>0.01.

In total there were more than 1600 proteins, and 957 of them were found in the biofilm matrix, 766
matrix proteins were present in both biological replicates.

3.2. Unique proteins in the extracellular matrix

Matrix unique proteins were defined if they were observed (LFQ intensity was nonzero) in both
experiments in the matrix, while LFQ intensities in cells samples were zero. Totally ten proteins were
present in the extracellular matrix only. We have proceeded with literature search and manual biofilm-
related functional annotation for these matrix unique proteins (Table 2). Nonetheless, for most extracellular
matrix-only proteins we did not find clear evidence of their importance for biofilm structure or any
extracellular function.

Table 2. Matrix unique proteins. Matrix unique proteins had nonzero intensities in all matrix samples and zero
intensities in all cell samples.

Protein	Protein ID	Role in biofilm	Reference
two-partner secretion system transporter CdrB	WP_010895680.1	Secretion partner of CdrA adhesin	[33]
DUF3298 and DUF4163 domain-containing protein	WP_003102069.1	Not described*	-
hypothetical protein	WP_003113151.1	Not described	-
hypothetical protein	WP_003102379.1	Not described	-
ABC transporter substrate-binding protein (Probable amino acid-binding protein)	WP_003113778.1	Not described, proposed transport function	-
NAD(P)/FAD-dependent oxidoreductase	WP_003114698.1	Not described	-
nucleoside hydrolase Nuh (catabolizes adenosine)	WP_003147076.1	QS-controlled private good	[34]
YgdI/YgdR family lipoprotein	WP_003089729.1	Not described	-
dGTPase	WP_003112492.1	Not described	-
ATP-dependent zinc protease	WP_003113269.1	Not described, but some envelope proteases play role in many biological processes	[35]

*Not described means absence of evidence of significant involvement specifically in biofilm formation and/or
pathogenesis. That does not reflect a described role in any other non-related to biofilm functions.

Additionally to unique matrix proteins, there were proteins with significant and extremely high log2
fold change (Log2FC) – more than 20. We consider these proteins as semi-unique matrix proteins (Table 3).

Table 3. Semi-unique matrix proteins. Semi-unique proteins had nonzero intensitiesLFQ in both cells and matrix
samples and log2FC more than 20.

Protein	Protein ID	Role in biofilm	Reference
malonate decarboxylase subunit alpha	WP_003112666.1	Malonate improve biofilm formation	[36]
OprD family porin	WP_003114177.1	Not described*, proposed transport function	-
ABC transporter substrate-binding protein	WP_003098136.1	Not described, proposed transport function	-
carbon-nitrogen hydrolase family protein	WP_003112739.1	Not described	-
bifunctional riboflavin kinase/FAD synthetase	WP_003102619.1	Not described; flavin nucleotide biosynthesis	-
M48 family metallopeptidase	WP_003112546.1	Not described	-

transporter substrate-binding domain-containing protein	WP_003115039.1	Not described, proposed transport function	-
division/cell wall cluster transcriptional repressor MraZ	WP_003103101.1	Not described	-

*Not described means absence of evidence of significant involvement specifically in biofilm formation and/or
pathogenesis. That does not reflect a described role in any other non-related to biofilm functions.

3.3. Functional classification of proteins in the extracellular matrix

All overrepresented in matrix proteins are displayed in Figure 2A. We observed distinguishable
clusters of overrepresented proteins depending on their fold change (Log2FC). While a separate cluster of
highly overrepresented proteins with log2FC more than 10 is visible on the main volcano plot (Figure 2A),
for better resolution of the area within log2FC 1-10 frame we also provide an enlarged area of the volcano
plot (Figure 2B). Analysis of bioprocess classification of all found in matrix proteins and overrepresented
in matrix proteins is displayed in Figure 3. Many bioprocess groups include at least one overrepresented
in biofilm matrix protein.

Figure 3. Bioprocess group classification of proteins found in the biofilm matrix. A – number of represented
proteins in different bioprocess groups concerning log2 fold change in comparison with embedded cells: grey color –
proteins found in the matrix but not overrepresented in comparison with embedded cells; green color –
overrepresented proteins with log2 fold change from 1 to 10; orange color – overrepresented proteins with log2 fold
change more than 10, only groups with at least 2 found in matrix proteins and at least one overrepresented protein are
displayed; B – distribution of overrepresented proteins in biofilm matrix, each color represents individual bioprocess
group, the legend indicates top ten of bioprocess groups.

We focus on groups with the largest number of found proteins and the proportion of overrepresented
proteins from all found in matrix proteins. For this reason, we applied the following criteria: (1) at least 10
proteins per group found in the matrix; (2) more than 3 found in matrix proteins are overrepresented (4 to
28 overrepresented proteins from 8 bioprocess groups). Such bioprocesses belong to oxidation-reduction
and cell redox homeostasis, proteolysis, transmembrane transport, amino acid, and nucleoside metabolic
processes, fatty acids biosynthesis. An important consideration is that GO annotated bioprocesses consist

of groups with highly different numbers of proteins. Number of proteins involved in bioprocess reflects complex nature and flexibility of protein interactions. As an absolute number of identified proteins may not reflect bioprocess representation, we count for each selected bioprocess group proportion of found proteins from all GO annotated group members as a measure of bioprocess representation (Table 4).

Table 4. Bioprocess representation in biofilm matrix based on the proportion of found proteins from GO-annotated for the bioprocess proteins.

	number of found in matrix proteins	number of overrepresented in matrix proteins	GO-annotated number of proteins in the database	%% of found in matrix proteins from GO-annotated in database
oxidation-reduction process	128	27	448	29
proteolysis	23	8	87	26
biosynthetic process	20	5	985	2
transmembrane transport	18	8	905	2
cellular amino acid metabolic process	16	7	141	11
cell redox homeostasis	14	5	26	54
fatty acid biosynthetic process	12	4	26	46
nucleoside metabolic process	10	5	11	91

In such analysis, the top three represented bioprocesses were cell redox homeostasis, nucleoside metabolism, and fatty acid synthesis. Considering annotated number of proteins for each group nucleoside metabolic process was represented with ten proteins in the matrix out of eleven GO annotated proteins. In contrast to the nucleoside metabolic process, DNA-templated regulation of transcription (this group includes only 2 overrepresented in matrix proteins) and biosynthetic process groups were represented in matrix with a greater number of proteins (22 and 20, respectively), but it was less than 5 % from GO annotated in database proteins for these groups (22 proteins from 472 and 20 from 985, respectively).

3. Discussion

Biofilm matrix in most cases contain many proteins. Extracellular proteins in the biofilm matrix provide their functions for the whole bacterial community, so they may be considered as public goods. Extracellular functions of matrix proteins include (but are not limited to) the external digestive system, signaling, protection, and maintaining the stability of the matrix [1, 37–39]. Moreover, many bacterial proteins may have moonlight functions.

Understanding protein composition is critical for resolving both fundamental questions in bacterial lifestyle and the development of tools for manipulating biofilms. Most proteomic studies focus on the whole biofilm and usually compare the whole biofilm proteome with planktonic cells proteome. At the same time, the identification of matrix protein composition may open new opportunities for the development of antibiofilm drugs. It is well known that biofilms of pathogenic microorganisms are tolerant to antibiotics, many other therapeutics, and host immune factors due to the matrix [40]. Matrix-disrupting or interrupting agents may reverse tolerant phenotype and increase the efficacy of antibiotic therapy [41]. For example, antibody-mediated destabilization of the matrix through the disruption of IHF-DNA complexes was effective *in vitro* and *in vivo* as a single therapy and combined with antibiotics [42–44]. Moreover, extracellular targets less probable cause selection of resistant mutants, so targeting of matrix proteins is a perspective way to combat chronic infections especially infections caused by ESKAPE pathogens. Despite all of these it is still little known about matrix proteomes. Only numerous studies described the matrix proteome of reference strain PAO1 of *P. aeruginosa* and some other bacteria (Table 1). Here we for the first time describe the matrix proteome of CF clinical isolate of *P. aeruginosa* in comparison with embedded cells and discuss obtained data in a prism of possible targets for therapeutic development. In comparison with other studies, we identified the largest number of proteins in the matrix. While some proteins might be invisible due to their low concentrations and individual limits of detection, we believe that further improvement of MS equipment and technics will get a more comprehensive picture of bacterial proteomics.

In our study, only a small number of proteins were unique for matrix only. One of them – CdrB protein
is involved in the transport of CdrA adhesin. This adhesin is important for biofilm formation and its
binding to Psl results in increased biofilm structural stability. Also, it is known that *cdrAB* is regulated
together with Psl. CdrA binding to Psl protects it from endogenous and exogenous protease digestion [45].
Surprisingly we did not find any nonzero LFQ for CdrA protein neither in cells nor in the matrix. In a less
sensitive gel-based proteomic study authors have observed CdrA protein in the matrix [7, 20], so the
absence of the protein in our data is unlikely due to limits of detection but seems to be under the limit of
quantitation (LFQ calculation). One of the possible explanations is that CdrA/CdrB components are not
important for KB6 strain due to the limited role of Psl polysaccharides in the biofilm matrix, so in the
absence of Psl CdrA might be degraded by proteases to quantities under the detection limit.

Another matrix unique protein is nucleoside hydrolase (Nuh) – an enzyme that hydrolyzes adenosine
and inosine, allowing the cell to grow on these nucleosides as the sole carbon or nitrogen source. This
protein was studied as an intracellular (periplasmic) private good in a prism of quorum sensing in bacterial
communities [34]. In our study, we found Nuh in the matrix, but not inside bacterial cells. That means Nuh
might be an extracellular public good at least for some strains like ours. Also, transporters responsible for
adenosine transport to periplasmic space in *P. aeruginosa* remain still undiscovered, so if Nuh works outside
the cell such a factor is not needed. In an environment with adenosine as the sole carbon source Nuh mutant
has impaired growth [46], but therapeutic potential of direct or indirect inhibition of Nuh activity remains
elusive due to the nutrient-rich nature of infected tissues, and further research is needed.

For the rest 8 matrix-unique proteins there is no clear evidence of their possible role in biofilm.
Meanwhile, as matrix is considered a nutrient-rich environment presence of substrate-binding protein from
ABC transporter and its role in nutrient (probably amino acids) acquisition is obvious. Antibody-mediated
blocking of some ABC transporters was shown to be effective in vitro and in vivo against *M. hominis* and
S. aureus [47] [48]. Monoclonal antibody Aurograb® entered phase III clinical trial as an addition to
vancomycin therapy for deep-seated staphylococcal infections (NCT00217841), but the trial was stopped
due to lack of reaching the primary endpoint. Anyway, somewhere positive results in targeting eukaryotic
ABC transporters for cancer treatment support the idea of a broader evaluation of the similar capability for
prokaryotes. Also, transmembrane transport is one of the prevalent bioprocess groups in terms of all found
in matrix proteins (n=18) as well as in terms of several overrepresented in matrix protein (n=8).

ATP-dependent Zn proteases are common enzymes in the cell envelope of *P. aeruginosa*, they
participate in several processes including metabolism, protein transport and removal of misfolded
proteins, and adaptation to environmental conditions [35]. Also, some proteases may act as a virulence
factor - Zn^{2+}-dependent protease *Bacillus anthracis* called Lethal Factor is required for infection [49]. The
proteolysis bioprocess group was the second represented in the matrix with 8 overrepresented proteins.
Proteolysis is a part of the external digestive system and provides peptides and amino acids for bacterial
nutrition, so the presence of some proteins involved in amino acids metabolism was expected (this
bioprocess group includes 16 found in matrix proteins with 7 overrepresented proteins). Protease`s
functions could be crucial for both bacterial survival and infection process and targeting bacterial proteases
could be a perspective way to combat bacterial infections [50]. Moreover, for Zn proteases, host nutritional
immunity (including Zn-dependent processes) was shown to be effective against infections caused by *P.*
aeruginosa [51]. Also, deprivation of Zn ions was proposed to combat infections caused by another common
pathogen - *S. aureus* [52].

Despite a small number of matrix unique proteins, we found a lot of overrepresented proteins in the
matrix. Eight proteins with extremely high log2 fold change (more than 20) were considered as semi-unique
for matrix, but we did not find any related to biofilm role for these proteins in literature. So as for unique
matrix proteins, there is an unexplored area in biofilm biology.

Bioprocesses classification of represented in matrix proteins reveals several groups. Considering log2
fold change we found that groups of oxidation-reduction processes, biosynthetic processes, and nucleoside
metabolism had the largest number of highly overrepresented proteins. Obviously, each bioprocess might
vary in the number of involved proteins, so we also introduced bioprocess representation as a part of all
GO-bioprocess annotated proteins.

In the matrix, the most reach group of proteins (128 proteins found in matrix and 27 of them are
overrepresented in comparison with embedded cells) belongs to oxidation-reduction processes. Also, a
group of proteins involved in cell redox homeostasis was one of the most represented GO-annotated
bioprocesses in the matrix (14 proteins found in matrix, 8 were overrepresented from 26 annotated in GO
bioprocess database). Biofilms of *P. aeruginosa* are often enriched in molecules involved in virulence and

competition with other microorganisms, including redox-active molecules. Self-produced factors involved
in the generation of reactive oxygen species might be harmful to the internal bacterial community. *P.*
aeruginosa is balancing to maintain oxidation-reduction processes at the appropriate level, so maintaining
the balance of oxidation-reduction reactions and redox homeostasis likely play important role in the biofilm
matrix as an environment with limited diffusion. Several effective therapeutic approaches utilize oxidative
stress to combat bacterial biofilms including photodynamic therapy (PDT) and sanitizers like hydrogen
peroxide [53, 54]. Also, recently was shown that extracellular electron transfer (EET) exists inside biofilm
matrix [55]. In matrix EET mainly involves pyocyanin and eDNA, but the role of the protein component is
not yet studied.

The second represented bioprocess included proteins involved in the fatty acid biosynthetic process
(4 overrepresented proteins from 12 found in the matrix). Fatty acids are one of the major components of
the cell envelope. Also, it is a well-known signal function of cis-2-docenoic acid messenger (DSM) as well
as the importance of the fatty acid component of AHL [56]. In a recently published study, Altay et al. made
a comprehensive analysis of essential reactions and affected pathways in *B. cenocepacia* (both planktonic
and biofilm) using a systems biology approach. From all identified essential reactions lipid metabolism
was accounted for more than half of the single lethal reactions; among this fatty acid biosynthesis was most
frequently found [57]. That data support the further development of fatty acids metabolism inhibitors as
promising therapeutics against bacterial infections, including bacterial biofilms.

The most represented bioprocess was nucleoside metabolism – 10 out of 11 GO-annotated proteins
were found in the matrix. Biofilms are often enriched with extracellular nucleic acids which act not only as
a structural component or component of EET, but also were considered as a nutrient depo [58]. Nucleosides
act as substrates and cofactors in many biosynthetic processes, as signal molecules, and are involved in
regulating bacterial community inside biofilm [59], so the presence of nucleoside metabolic proteins is
required. Therapeutic targeting of proteins involved in these processes is theoretically possible, but at
present is not clear. At the same time, nucleoside analogs are common drugs in other nonbacterial diseases,
and evaluation of their possible role as antibacterial drugs may open new opportunities [60]. So antimycotic
drug 5-fluorocytosine was able to suppress virulence of *P. aeruginosa* in a murine model of lung infection
[61].

Obviously, sources of extracellular matrix proteins belong to two main categories: (1) active secretion
of biomolecules and (2) passive way to increase extracellular content as a consequence of cell leakage or
lysis. While some extracellular proteins are «passive» products of bacterial cell lysis, they still might be
active outside the cell and provide their function to the bacterial community extracellularly. As protein
degradation rates are varied in broad ranges, some proteins may present in the matrix for a long time after
leakage or secretion from the cell, so their occurrence in the matrix does not (1) match the actual situation
inside cells or (i. e. active transcription and translation), or (2) reflect any real extracellular needs for biofilm
(i. e. be structurally or/and physiologically involved in extracellular processes). The limitation of our study
is an inability to conclude if the protein is «archeological», bystander, or functionally active in the matrix
and how these proteins are distributed in the matrix (are they cell-attached, part of OMVs, or associated
with other matrix components). Moreover, some proteins may act as distracting extracellular targets in a
way of absorption of inhibitor and distract from intracellular targets, so further research with a focus on
the function and dynamics of every single protein is needed.

4. Materials and Methods

2.1. Biofilm growth and separation matrix from cells

P. aeruginosa KB6 (CF clinical isolate) was a gift from Zigangirova N. A. (Gamaleya NRCEM, Moscow,
Russia) [21]. For biofilm preparation single colony from TSA plate was picked in liquid LB medium and
grown 24 h at 37 C, 210 rpm. The liquid culture was diluted 50 times with LB medium in a volume of 20
ml and placed in Petri dishes for 18 h at 37 C under static conditions. Then the medium was removed, and
biofilm was exposed to 10 ml of 1.8 M NaCl. After 5 min of incubation bacterial suspension and dissolved
matrix were separated with centrifugation at 5000 g. Liquid phase (dissolved matrix) was filtered through
a 0.22 mkm syringe filter and protein was precipitated with cold acetone (up to 80 %) 18 h at -20 C. Cells
pellet was resuspended in lysis buffer (2% SDS; 50 mM Tris-HCl; 180 mM NaCl; 0,1 mM EDTA; 1 mM
MgCl2) and boiled for 15 min in a water bath. Cell debris was removed by centrifugation at 10000 g, 15
min, and proteins from the liquid phase were precipitated with 80% cold acetone as for matrix samples.
Precipitated proteins were pelleted with centrifugation at 10000 g, 20 min, 4 C. Pellet was washed two times
with 80 % cold acetone and proceeded for proteomic sample preparation.

2.2. Proteomic sample preparation and peptide identification

Proteomic sample preparation and peptide identification were made in Advanced Mass Spectrometry Core Facility (Skolkovo Innovation Center, Moscow, Russia). The protein pellet was subjected to tryptic in-solution digestion. LC-MS/MS was carried out on a Q Exactive HF (Thermo Scientific) with a nanoESI interface in conjunction with an Ultimate 3000 RSLC nano HPLC (Dionex Ultimate 3000). Peptides were loaded onto a trap column and separated on an analytical column (C18) using an H2O/acetonitrile gradient with 0.1% formic acid for 150 min. The Q Exactive HF spectrometer was operated in the data-dependent mode with a nanoESI spray voltage of 1.8 kV, capillary temperature of 210 °C, and S-lens RF value of 55%. All spectra were acquired in positive mode with full scan MS spectra scanning from m/z 310–1500 in the FT mode at 120,000 resolution. A lock mass of 445.120025 was used. The top 25 most intense precursors were subjected to rapid collision induced dissociation (rCID). Dynamic exclusion with of 70 seconds was applied for repeated precursors.

2.3. Data analysis

To identify and quantify tryptic peptides and the proteins from which the peptides are derived, spectra from the MS/MS experiments were analyzed by GUI FragPipe v. 17.1 (https://github.com/Nesvilab/FragPipe). Peptide identification was performed by MSFragger search engine [22, 23] using protein sequence database extracted from NCBI (*Pseudomonas aeruginosa* strain GIMC5015:PAKB6 chromosome, complete genome NZ_CP034429) with decoys and contaminants. Oxidation of methionine and acetylation of protein N-termini were set as variable modifications, carbamidomethylation of cysteine was set as a fixed modification. The maximum allowed variable modifications per peptide was set to 3, mass tolerance was set as 20 ppm for precursor and 0.02 Da for fragment ions. Philosopher kit tools [24, 25] were used to estimate identification FDR. The PSMs were filtered at 1% PSM and 1% protein identification FDR. Quantification by label-free protein quantitation method and MBR was performed with IonQuant [26]. Obtained quantified data (intensities) were processed for differential expression analyses with limma package R [27], with followed visualization result by EnhancedVolcano package R ("EnhancedVolcano: Publication-ready volcano plots with enhanced coloring and labeling." https://github.com/kevinblighe/EnhancedVolcano).

Bioprocess classification and functional annotation were made with the Pseudomonas Genome Database (https://pseudomonas.com/primarySequenceFeature/list?strain_ids=10430&term=Pseudomonas+aerugino sa+GIMC5015%3APAKB6&c1=name&v1=&e1=1&assembly=complete) [28].

For correlation analysis, we used GraphPad Prism 9 desktop software (version 9.2.0). For the Figures arrangement, we used bio-render online software (Biorender.com).

5. Conclusions

Biofilm matrix of clinical strain *P. aeruginosa* is enriched with proteins. There are several unique for matrix and many overrepresented in matrix proteins, which reflects several bioprocesses. Development of antibiofilm therapeutics may benefit in the case of targeting proteins and processes taking place in biofilm matrix as the sole mechanism of action or in combination with antibiotics. Altogether matrix protein composition is important for choosing successful strategy in antibacterial drug development and reaching unmet needs of curing biofilm infections.

Supplementary Materials: The following supporting information can be downloaded:

Table S1: All identified proteins.

Table S2: Differentially represented proteins.

Author Contributions: Conceptualization, D.E., A.S., N.P.; methodology, A.S., K.D.; software, A.S., N.P.; validation, A.S.; investigation, D.E., I.K., V.R.; resources, Y.R.; data curation, M.D.; writing—original draft preparation, D.E., M.D., I.T.; writing—review and editing, Y.R.; project administration, D.E.; funding acquisition, A.G. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by Government Grant from the Ministry of Health of the Russian Federation № 056-00119-21-00-1F.

Data Availability Statement: The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE [62] partner repository with the dataset identifier PXD031480.

Acknowledgments: We thank Nailia A. Zigangirova (Gamaleya NRCEM, Moscow, Russia) for the *P. aeruginosa* KB6
strain. We thank Viktor G. Zgoda from Advanced Mass Spectrometry Core Facility (Skolkovo Innovation Center,
Moscow, Russia) for technical consulting and MS data obtaining.

Conflicts of Interest: The authors declare no conflict of interest.

References

1.   Karygianni L, Ren Z, Koo H, Thurnheer T. Biofilm Matrixome: Extracellular Components in Structured Microbial
Communities. Trends Microbiol 2020.
2.   Mirzaei R, Ranjbar R. Hijacking host components for bacterial biofilm formation: An advanced mechanism. Int
Immunopharmacol 2022; 103: 108471.
3.   Fong JNC, Yildiz FH. Biofilm Matrix Proteins. Microbiol Spectr 2015; 3.
4.   Erdmann J, Thöming JG, Pohl S, Pich A, Lenz C, Häussler S. The Core Proteome of Biofilm-Grown Clinical
Pseudomonas aeruginosa Isolates. Cells 2019; 8: 1129.
5.   Reales-Calderón JA, Sun Z, Mascaraque V, Pérez-Navarro E, Vialás V, Deutsch EW, et al. A wide-ranging
Pseudomonas aeruginosa PeptideAtlas build: A useful proteomic resource for a versatile pathogen. J Proteomics
2021; 239: 104192.
6.   Penesyan A, Kumar SS, Kamath K, Shathili AM, Venkatakrishnan V, Krisp C, et al. Genetically and
Phenotypically Distinct Pseudomonas aeruginosa Cystic Fibrosis Isolates Share a Core Proteomic Signature. Plos
One 2015; 10: e0138527.
7.   Toyofuku M, Roschitzki B, Riedel K, Eberl L. Identification of Proteins Associated with the Pseudomonas
aeruginosa Biofilm Extracellular Matrix. J Proteome Res 2012; 11: 4906–4915.
8.   Tseng B, Reichhardt C, Merrihew GE, Araujo-Hernandez SA, Harrison JJ, MacCoss MJ, et al. A Biofilm Matrix-
Associated Protease Inhibitor Protects Pseudomonas aeruginosa from Proteolytic Attack. Mbio 2018; 9: e00543-
18.
9.   Couto N, Schooling SR, Dutcher JR, Barber J. Proteome Profiles of Outer Membrane Vesicles and Extracellular
Matrix of Pseudomonas aeruginosa Biofilms. J Proteome Res 2015; 14: 4207–4222.
10. Park AJ, Murphy K, Surette MD, Bandoro C, Krieger JR, Taylor P, et al. Tracking the Dynamic Relationship
between Cellular Systems and Extracellular Subproteomes in Pseudomonas aeruginosa Biofilms. J Proteome Res
2015; 14: 4524–4537.
11. Zhang W, Sun J, Ding W, Lin J, Tian R, Lu L, et al. Extracellular matrix-associated proteins form an integral and
dynamic system during Pseudomonas aeruginosa biofilm development. Front Cell Infect Mi 2015; 5: 40.
12. Reigada I, San-Martin-Galindo P, Gilbert-Girard S, Chiaro J, Cerullo V, Savijoki K, et al. Surfaceome and
Exoproteome Dynamics in Dual-Species Pseudomonas aeruginosa and Staphylococcus aureus Biofilms. Front
Microbiol 2021; 12: 672975.
13. Lei MG, Gupta R, Lee CY. Proteomics of Staphylococcus aureus biofilm matrix in a rat model of orthopedic
implant-associated infection. PLOS ONE 2017; 12: e0187981.
14. Kavanaugh JS, Flack CE, Lister J, Ricker EB, Ibberson CB, Jenul C, et al. Identification of Extracellular DNA-
Binding Proteins in the Biofilm Matrix. mBio 2019; 10.
15. Terán LC, Distefano M, Bellich B, Petrosino S, Bertoncin P, Cescutti P, et al. Proteomic Studies of the Biofilm
Matrix including Outer Membrane Vesicles of Burkholderia multivorans C1576, a Strain of Clinical Importance
for Cystic Fibrosis. Microorg 2020; 8: 1826.
16. Jachlewski S, Jachlewski WD, Linne U, Bräsen C, Wingender J, Siebers B. Isolation of Extracellular Polymeric
Substances from Biofilms of the Thermoacidophilic Archaeon Sulfolobus acidocaldarius. Frontiers in
Bioengineering and Biotechnology 2015; 3: 123.
17. Cao B, Shi L, Brown RN, Xiong Y, Fredrickson JK, Romine MF, et al. Extracellular polymeric substances from
Shewanella sp. HRCR-1 biofilms: characterization by infrared spectroscopy and proteomics. Environmental
Microbiology 2011; 13: 1018–1031.
18. Gannesen AV, Zdorovenko EL, Botchkova EA, Hardouin J, Massier S, Kopitsyn DS, et al. Composition of the
Biofilm Matrix of Cutibacterium acnes Acneic Strain RT5. Front Microbiol 2019; 10: 1284.
19. Absalon C, Dellen K, Watnick PI. A Communal Bacterial Adhesin Anchors Biofilm and Bystander Cells to
Surfaces. Plos Pathog 2011; 7: e1002210.
20. Wu S, Baum MM, Kerwin J, Guerrero D, Webster S, Schaudinn C, et al. Biofilm-specific extracellular matrix
proteins of nontypeable Haemophilus influenzae. Pathog Dis 2014; 72: 143–160.
21. Sheremet AB, Zigangirova NA, Zayakin ES, Luyksaar SI, Kapotina LN, Nesterenko LN, et al. Small Molecule
Inhibitor of Type Three Secretion System Belonging to a Class 2,4-disubstituted-4H-[1,3,4]-thiadiazine-5-ones
Improves Survival and Decreases Bacterial Loads in an Airway Pseudomonas aeruginosa Infection in Mice.
Biomed Res Int 2018; 2018: 1–13.
22. Kong AT, Leprevost FV, Avtonomov DM, Mellacheruvu D, Nesvizhskii AI. MSFragger: ultrafast and
comprehensive peptide identification in mass spectrometry–based proteomics. Nat Methods 2017; 14: 513–520.
23. Teo GC, Polasky DA, Yu F, Nesvizhskii AI. Fast Deisotoping Algorithm and Its Implementation in the MSFragger
Search Engine. J Proteome Res 2020; 20: 498–505.
24. Leprevost F da V, Haynes SE, Avtonomov DM, Chang H-Y, Shanmugam AK, Mellacheruvu D, et al. Philosopher:
a versatile toolkit for shotgun proteomics data analysis. Nat Methods 2020; 17: 869–870.
25. Keller A, Nesvizhskii AI, Kolker E, Aebersold R. Empirical Statistical Model To Estimate the Accuracy of Peptide
Identifications Made by MS/MS and Database Search. Anal Chem 2002; 74: 5383–5392.
26. Yu F, Haynes SE, Nesvizhskii AI. IonQuant Enables Accurate and Sensitive Label-Free Quantification With FDR-
Controlled Match-Between-Runs. Mol Cell Proteomics 2021; 20: 100077.

27.   Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, et al. limma powers differential expression analyses for
445   RNA-sequencing and microarray studies. Nucleic Acids Res 2015; 43: e47–e47.
446   28.   Winsor GL, Griffiths EJ, Lo R, Dhillon BK, Shay JA, Brinkman FSL. Enhanced annotations and features for
447   comparing thousands of Pseudomonas genomes in the Pseudomonas genome database. Nucleic Acids Res 2016;
448   44: D646–D653.
449   29.   Scherbakova A, Rykova V, Danilova K, Solovyev A, Egorova D. Extracellular RNA Isolation from Biofilm Matrix
450   of Pseudomonas aeruginosa. Bio-protocol 2020; 10.
451   30.   Chiba A, Sugimoto S, Sato F, Hori S, Mizunoe Y. A refined technique for extraction of extracellular matrices from
452   bacterial biofilms and its applicability. Microbial Biotechnology 2015; 8: 392–403.
453   31.   Lamers RP, Nguyen UT, Nguyen Y, Buensuceso RNC, Burrows LL. Loss of membrane-bound lytic
454   transglycosylases increases outer membrane permeability and β-lactam sensitivity in Pseudomonas aeruginosa.
455   Microbiologyopen 2015; 4: 879–895.
456   32.   Kammers K, Cole RN, Tiengwe C, Ruczinski I. Detecting significant changes in protein abundance. Eupa Open
457   Proteom 2015; 7: 11–19.
458   33.   Borlee BR, Goldman AD, Murakami K, Samudrala R, Wozniak DJ, Parsek MR. Pseudomonas aeruginosa uses a
459   cyclic-di-GMP-regulated adhesin to reinforce the biofilm extracellular matrix. Mol Microbiol 2010; 75: 827–842.
460   34.   Schuster M, Sexton DJ, Hense BA. Why Quorum Sensing Controls Private Goods. Front Microbiol 2017; 8: 885.
461   35.   Heywood A, Lamont IL. Cell envelope proteases and peptidases of Pseudomonas aeruginosa: multiple roles,
462   multiple mechanisms. Fems Microbiol Rev 2020; 44: 857–873.
463   36.   Elmassry MM, Bisht K, Colmer-Hamood JA, Wakeman CA, Francisco MJS, Hamood AN. Malonate utilization by
464   Pseudomonas aeruginosa affects quorum-sensing and virulence and leads to formation of mineralized biofilm-
465   like structures. Mol Microbiol 2021; 116: 516–537.
466   37.   Egorova DA, Voronina OL, Solovyev AI, Kunda MS, Aksenova EI, Ryzhova NN, et al. Integrated into
467   Environmental Biofilm Chromobacterium vaccinii Survives Winter with Support of Bacterial Community.
468   Microorganisms 2020; 8.
469   38.   Flemming H-C, Wingender J. The biofilm matrix. Nat Rev Microbiol 2010; 8: 623–633.
470   39.   Romaní AM, Fund K, Artigas J, Schwartz T, Sabater S, Obst U. Relevance of Polymeric Matrix Enzymes During
471   Biofilm Formation. Microbial Ecol 2008; 56: 427–436.
472   40.   Uruén C, Chopo-Escuin G, Tommassen J, Mainar-Jaime RC, Arenas J. Biofilms as Promoters of Bacterial Antibiotic
473   Resistance and Tolerance. Antibiotics 2020; 10: 3.
474   41.   Gunn JS, Bakaletz LO, Wozniak DJ. What's on the Outside Matters: The Role of the Extracellular Polymeric
475   Substance of Gram-negative Biofilms in Evading Host Immunity and as a Target for Therapeutic Intervention*. J
476   Biol Chem 2016; 291: 12538–12546.
477   42.   Estellés A, Woischnig A-K, Liu K, Stephenson R, Lomongsod E, Nguyen D, et al. A High-Affinity Native Human
478   Antibody Disrupts Biofilm from Staphylococcus aureus Bacteria and Potentiates Antibiotic Efficacy in a Mouse
479   Implant Infection Model. Antimicrobial Agents and Chemotherapy 2016; 60: 2292–2301.
480   43.   Brockson EM, Novotny LA, Mokrzan EM, Malhotra S, Jurcisek JA, Akbar R, et al. Evaluation of the kinetics and
481   mechanism of action of anti-integration host factor-mediated disruption of bacterial biofilms. Mol Microbiol 2014;
482   93: 1246–1258.
483   44.   Novotny L, Goodman S, Bakaletz L. Redirecting the immune response towards immunoprotective domains of a
484   DNABII protein resolves experimental otitis media. npj Vaccines 2019; 4.
485   45.   Reichhardt C, Wong C, Silva DP da, Wozniak DJ, Parsek MR. CdrA Interactions within the Pseudomonas
486   aeruginosa Biofilm Matrix Safeguard It from Proteolysis and Promote Cellular Packing. Mbio 2018; 9: e01376-18.
487   46.   Heurlier K, Dénervaud V, Haenni M, Guy L, Krishnapillai V, Haas D. Quorum-Sensing-Negative (lasR) Mutants
488   of Pseudomonas aeruginosa Avoid Cell Lysis and Death. J Bacteriol 2005; 187: 4875–4883.
489   47.   Burnie JP, Matthews RC, Carter T, Beaulieu E, Donohoe M, Chapman C, et al. Identification of an
490   Immunodominant ABC Transporter in Methicillin-Resistant Staphylococcus aureusInfections. Infect Immun
491   2000; 68: 3200–3209.
492   48.   Burmistrova DA, Tillib SV, Shcheblyakov DV, Dolzhikova IV, Shcherbinin DN, Zubkova OV, et al. Genetic
493   Passive Immunization with Adenoviral Vector Expressing Chimeric Nanobody-Fc Molecules as Therapy for
494   Genital Infection Caused by Mycoplasma hominis. PLOS ONE 2016; 11.
495   49.   Moayeri M, Leppla SH, Vrentas C, Pomerantsev A, Liu S. Anthrax Pathogenesis. Annu Rev Microbiol 2014; 69:
496   1–24.
497   50.   Culp E, Wright GD. Bacterial proteases, untapped antimicrobial drug targets. J Antibiotics 2017; 70: 366–377.
498   51.   Wang S, Cheng J, Niu Y, Li P, Zhang X, Lin J. Strategies for Zinc Uptake in Pseudomonas aeruginosa at the Host–
499   Pathogen Interface. Front Microbiol 2021; 12: 741873.
500   52.   Elhakim YA, Ali AE, Hosny AE-DMS, Abdeltawab NF. Zinc Deprivation as a Promising Approach for Combating
501   Methicillin-Resistant Staphylococcus aureus: A Pilot Study. Pathogens 2021; 10: 1228.
502   53.   Tiganova IG, Zhizhimova YS, Philipova NI, Tolordava ER, Alekseeva NV, Makarova EA, et al. Antibacterial
503   Properties of Synthetic Cationic Bacteriochlorin Derivatives as Photosensitizers. Mol Genetics Microbiol Virology
504   2020; 35: 248–256.

54. Lineback CB, Nkemngong CA, Wu ST, Li X, Teska PJ, Oliver HF. Hydrogen peroxide and sodium hypochlorite
disinfectants are more effective against Staphylococcus aureus and Pseudomonas aeruginosa biofilms than
quaternary ammonium compounds. Antimicrob Resist Infect Control 2018; 7: 154.

55. Saunders SH, Tse ECM, Yates MD, Otero FJ, Trammell SA, Stemp EDA, et al. Extracellular DNA Promotes
Efficient Extracellular Electron Transfer by Pyocyanin in Pseudomonas aeruginosa Biofilms. Cell 2020; 182: 919-
932.e19.

56. Williams P. Quorum sensing, communication and cross-kingdom signalling in the bacterial world. Microbiology+
2007; 153: 3923–3938.

57. Altay O, Zhang C, Turkez H, Nielsen J, Uhlén M, Mardinoglu A. Revealing the Metabolic Alterations during
Biofilm Development of Burkholderia cenocepacia Based on Genome-Scale Metabolic Modeling. Metabolites
2021; 11: 221.

58. Finkel SE, Kolter R. DNA as a Nutrient: Novel Role for Bacterial Competence Gene Homologs. J Bacteriol 2001;
183: 6288–6293.

59. Fontaine BM, Duggal Y, Weinert EE. Exploring the Links between Nucleotide Signaling and Quorum Sensing
Pathways in Regulating Bacterial Virulence. Acs Infect Dis 2018; 4: 1645–1655.

60. Goncheva MI, Chin D, Heinrichs DE. Nucleotide biosynthesis: the base of bacterial pathogenesis. Trends
Microbiol 2022.

61. Imperi F, Massai F, Facchini M, Frangipani E, Visaggio D, Leoni L, et al. Repurposing the antimycotic drug
flucytosine for suppression of Pseudomonas aeruginosa pathogenicity. Proc National Acad Sci 2013; 110: 7458–
7463.

62. Perez-Riverol Y, Bai J, Bandla C, Hewapathirana S, García-Seisdedos D, Kamatchinathan S, Kundu D, Prakash A,
Frericks-Zipper A, Eisenacher M, Walzer M, Wang S, Brazma A, Vizcaíno JA (2022). The PRIDE database
resources in 2022: A Hub for mass spectrometry-based proteomics evidence. Nucleic Acids Res 50(D1):D543-
D552.

63. Graf AC, Leonard A, Schäuble M, Rieckmann LM, Hoyer J, Maass S, Lalk M, Becher D, Pané-Farré J, Riedel K.
Virulence Factors Produced by Staphylococcus aureus Biofilms Have a Moonlighting Function Contributing to
Biofilm Integrity. Mol Cell Proteomics. 2019 Jun;18(6):1036-1053.

1652 proteins

957 proteins
in matrix