Molecular features underlying *Pseudomonas aeruginosa* persistence in human plasma

Manon Janet-Maitre[1], Stéphane Pont[1*], Frerich Masson[2], Julian Trouillon[1**], Mylène Robert-Genthon[1], Chantal Dumestre-Perrard[3,4], Sylvie Elsen[1], Christine Moriscot[5], Bart Bardoel[2], Suzan Rooijakkers[2], François Cretin[1] and Ina Attrée[1]

[1] University Grenoble Alpes, Institute of Structural Biology, Team Bacterial Pathogenesis and Cellular Responses, 38054 Grenoble, France

[2] Medical Microbiology, University Medical Center Utrecht, Utrecht University, 3584 CX Utrecht, The Netherlands

[3] Laboratory of immunology, Grenoble Alpes University Hospital, cedex 9, 38043 Grenoble, France

[4] University Grenoble Alpes, CNRS, CEA, Institute of Structural Biology (IBS), cedex 9, 38044 Grenoble, France

[5] University Grenoble Alpes, CNRS, CEA, ISBG, F-38000 Grenoble, France

* present address: Laboratory of Pathogen-Host Interactions (LPHI), Université Montpellier, CNRS, UMR5235, 34095 Montpellier cedex 5, France.

** present address: Institute of Molecular Systems Biology, ETH Zurich, Zurich, Switzerland

Abstract

Persistence of bacterial pathogens is a main cause of treatment failure and establishment of chronic bacterial infection. *Pseudomonas aeruginosa*, an opportunistic Gram-negative pathogen is a leading cause of bacteremia with high associated mortality. In a previous study, we found that plasma-sensitive *P. aeruginosa* strains were able to form a persister-like sub-population named evaders. However, the molecular mechanisms underlying emergence and biology of evaders remained unknown. Here, using a gain-of-function genetic screen we examined the molecular determinants of persistence in plasma. In addition to known virulence factors (long O-specific antigens and exopolysaccharides), we found that ATP and biotin availability greatly impact bacterial survival in plasma. Mutants in genes of purine and biotin pathways display higher tolerance and persistence, respectively. Moreover, we identified a novel small protein named SrgA whose expression leads to 100-fold increase in survival in plasma. The analysis of different steps of complement activity and the use of outer-membrane impermeable drug nisin suggest that the mutants impede MAC activity *per se*. Through this work we highlight the multifactorial origin of bacterial resilience to plasma and pave the way to a more comprehensive picture of the complex interplay between *P. aeruginosa* and the complement system in the blood.

Introduction

Human bacterial pathogens employ sophisticated strategies to escape from the immune system and to resist or tolerate antibiotic treatments. Bacterial persistence towards antibiotics is a major obstacle in treatment of life-threatening infections and is frequently associated with the establishment of chronic infections, as it allows the survival of a sub-population that tolerates otherwise lethal antibiotic stress (Fauvart, De Groote, and Michiels 2011; Bartell et al. 2020). Unlike antibiotic resistance, persistence is a non-heritable, fully reversible trait, characterized by a biphasic killing curve with the rapid killing of the bulk population and the survival of a sub-population over a long period of time. Among different factors proposed to lead to antibiotic persistence (Balaban *et al.*, 2019), low intracellular levels of ATP (Shan *et al.*, 2017; Manuse *et al.*, 2021) and production of the "alarmone" signaling molecule guanosine (penta) tetraphosphate, (p)ppGpp (Korch, Henderson and Hill, 2003; Pacios *et al.*, 2020) are the most cited. Upon external antibiotic stress, the stochastic or responsive activation of the stringent response leads to accumulation of (p)ppGpp, which in turn activates toxin-antitoxin modules and degradation of the antitoxins through a regulation cascade, leading to inhibition of cell growth (Gaca, Colomer-Winter and Lemos, 2015; Harms, Maisonneuve and Gerdes, 2016). The host immune system also triggers persistence in different bacterial species. Vacuolar acidification after *Salmonella* internalization by macrophages induced persister cell formation and could contribute to the establishment of a bacterial reservoir for infection relapse (Helaine *et al.*, 2014). In a similar manner, human serum triggers the formation of antibiotic persisters as well as so called "viable but non-culturable" forms in *Vibrio vulnificus* (Ayrapetyan *et al.*, 2015) and (Putrinš *et al.*, 2015) also highlighted *Escherichia coli* phenotypic heterogeneity leading to serum and antibiotic killing evasion. We previously

demonstrated that a persister-like sub-population of evaders forms when globally

sensitive bacteria are incubated in human blood or human plasma. The evaders were

documented in several human pathogens including *Pseudomonas aeruginosa* (Pont

et al., 2020).

P. aeruginosa is a leading nosocomial pathogen extremely difficult to eradicate due to

its high adaptability to different environments, its intrinsic and acquired antibiotic

resistance (Moradali, Ghods and Rehm, 2017). The bloodstream infections (BSI) due

to *P. aeruginosa* strains hold the highest mortality rate with about 40% (Kang *et al.*,

2003; Vitkauskienė *et al.*, 2010). Interestingly, not all BSI isolates are fully resistant to

plasma. Although the complement system (CS) is the major immune component

responsible for *P. aeruginosa* killing in human blood and is globally efficient towards

sensitive strains, a tiny fraction of the initial bacterial population persists over prolonged

incubation in blood and plasma (Pont *et al.*, 2020). Kinetics of *P. aeruginosa* killing by

human plasma displays a classical "persister-type" biphasic profile (Balaban *et al.*,

2019), with a rapid elimination of more than 99.9% of the population reaching a plateau

at on average 0.1% after a 2h-incubation. We found that out of eleven *P. aeruginosa*

isolates tested, three were plasma-resistant, three displayed a tolerant phenotype and

five were able to form evaders, suggesting that the evaders may represent a common

strategy for *P. aeruginosa* to escape membrane attack complex (MAC)-mediated

bactericidal activity in plasma. The plasma evaders differed from antibiotic persisters

since a stationary phase population did not give rise to a higher population of evaders

as it was shown for antibiotic persisters (Keren *et al.*, 2004) and the mechanisms

permitting the evaders to resist the MAC-induced lysis remained unknown. In this

work, we refer to resistance, tolerance and persistence as defined in a context of

antibiotics in (Balaban *et al.*, 2019). Briefly, resistance is due to the heritable carriage

of a resistance factor allowing survival to a higher stress conditions. As opposed to resistance, tolerance is a phenomenon that allows survival of the population to an otherwise lethal stress for a longer period of time without any resistance mechanism. Persistence refers to a heterogeneous bacterial population composed of a rapidly eliminated sensitive population and a tolerant sub-population and is characterized by a biphasic killing curve. We use the term of resilience when including several of these phenomena.

In this work we used a genome-wide approach to get first insights into molecular mechanism(s) of *P. aeruginosa* resilience in human plasma. A screen based on transposon-insertion sequencing (Tn-seq) was applied to a recent clinical isolate that showed sensitivity to plasma and displayed 0.01-0.1% of evaders following incubation in plasma for 6h. We hypothesized that the transposon mutants displaying increased percentage of evaders or those with global increased plasma tolerance/resistance could reveal novel determinants of bacterial resilience to plasma and pathogen immune escape. In addition to known *P. aeruginosa* factors involved in plasma resistance, such as lipopolysaccharides (LPS) and exopolysaccharides, we identified an uncharacterized small periplasmic protein whose expression increased complement tolerance up to 100 fold. Moreover, our screen and phenotypic characterization of mutants revealed that the levels of biotin and ATP modulate pathogen sensitivity to MAC-induced lysis.

Results

Gain-of-function screen in plasma

In our previous study we found that the urinary isolate IHMA879472/AZPAE15042 (IHMA87) persisted in human blood and plasma. While the global population was

sensible to complement killing with 99.9% of the population eliminated within 60 minutes, approximately 0.1% evaders survived prolonged incubation in plasma (Pont *et al.*, 2020 and this work). In order to examine molecular determinants leading to persistence in plasma, we searched for hyper-evaders using a genome-wide approach. A transposon library of about 300,000 mutants was constructed using the Himar-1 transposon and insertion mutants with increased levels of survival in plasma were selected using the phenotypic screen schematically presented in Fig. 1A. The transposon library was challenged in native human plasma (output) or heat inactivated plasma (HIP, input) for three hours and recovered. The bacterial survival was determined by colony forming unit (CFU) counts for the whole library before and after the challenge, showing that the global survival increased by about 2-log in the mutant library compared to the parental strain. The screen set-up selects for mutants with global increased survival in plasma. The discrimination between resistant, hyper-evaders and tolerant phenotypes could be appreciated by performing killing curves over a long plasma incubation period.

Seven mutants were isolated directly during the screen and their survival was assayed in plasma killing assays. They survived 10 to 200 fold better than the parental strain (Fig. 1B), validating the screen. Among the isolated mutants, two mutants with hyper mucoid phenotypes reflecting alginate overproduction, carried insertions within *mucD* and in the intergenic region upstream of *mucA*, two negative regulators of alginate synthesis (reviewed in Damron and Goldberg, 2012). We also isolated a mutant with the transposon inserted in the intergenic region upstream of *mucE*, a positive regulator of alginate production (Qiu *et al.*, 2007; Yin *et al.*, 2013). The orientation of the transposon suggested that this mutant could overexpress *mucE,* which also led to a mucoid phenotype. The alginate indeed may interfere with the CS by decreasing C3b-

dependent opsonization and by interfering with CS activation when acetylated (Pedersen *et al.*, 1990; Pier *et al.*, 2001). A transposon insertion in *ladS*, encoding the histidine kinase positively affecting GacS/GacA-RsmA pathway (Ventre *et al.*, 2006; Broder, Jaeger and Jenal, 2017), resulted in increased mutant survival, by yet unknown effector expression. To note, the bioinformatics analysis of the sequencing data highlighted significant enrichment of transposon insertions in *retS*, a gene encoding for an antagonist to LadS, whose inactivation would lead to increased synthesis of polysaccharides Psl and Pel (Table 1). Indeed, Psl was shown to improve bacterial survival in serum in a mucoid strain (Jones and Wozniak, 2017). The role of Psl in complement resistance might be isolate-dependent, as it was initially shown using the reference strain PAO1, that Psl reduces bacterial opsonization without affecting C9 insertion and survival in serum (Mishra *et al.*, 2012). Moreover, insertion in the promoter of *pprB,* encoding the response regulator acting on type IVb pili/Tad, BapA adhesin and CupE fimbriae expression (Giraud *et al.*, 2011; de Bentzmann *et al.*, 2012) also led to increased survival in plasma. As shown, type IV pili contribute to serum resistance and recruitment of host negative complement regulator C4BP in *Vibrio cholerae* and *Neisseria gonorrheae*, respectively (Chiang *et al.*, 1995; Blom *et al.*, 2001). Our data suggests that either type IVb pili or CupE-fimbriae could mediate plasma resistance in *P. aeruginosa*. Random picking of colonies did not allow to obtain LPS-related mutants. However, the importance of LPS and more specifically O-specific antigens (OSA) biosyntheses in the interaction with plasma was highlighted by the screen and is in accordance with the literature (Goldberg and Pler, 1996; Priebe *et al.*, 2004; Kintz *et al.*, 2008; Osawa *et al.*, 2013). Two isolated clones with increased survival carried the transposon in *bioA* and *bioB*, encoding enzymes of biotin biosynthesis, to our knowledge, not previously linked to resilience to plasma or

complement.

Analysis of sequencing data was performed for both the insertions in coding regions

and insertions in intergenic regions, as the transposon used possesses an outward

tac promoter, possibly modulating the neighboring gene expression. In both analysis,

transposon insertions strongly increasing bacterial survival were associated with four

main pathways; *i.e.* biosynthetic pathways of purine and biotin, biosynthesis,

production and regulation of the exopolysaccharide alginate (Fig. 1C and Table 1,

complete results Table 1S).

Figure 1. Gain-of-function Tn-seq reveals common as well as novel pathways involved in *P. aeruginosa* survival in plasma. A. Schematic representation of the screen. A transposon mutant library was generated in the plasma-sensitive clinical isolate *P. aeruginosa* IHMA87 and incubated either in human plasma or in HIP (input) for 3h. Aliquots were plated on LB and individual transposon mutants were stored for further analysis. Transposon-insertion sequencing was performed by Illumina-based next generation sequencing. **B.** Survival of isolated transposon mutants in plasma following 3h-incubation. The survival was calculated using CFUs measurements. A

log-transformation was performed on the dataset and all mutants were shown significantly different from the wild-type strain (p-value < 0.005). Log_2(Fold-Change(FC)) obtained from the Tn-seq analysis for each gene or intergenic region is indicated above. **C.** Bioinformatics analysis presented as Volcano plots. Insertions in coding regions (left) and intergenic regions (right) were analyzed separately. Significant hits in genes of the same pathway are in the same color.

IHMA87 ID	PAO1 ID	Name/prediction	Log_2(FC)	P_{adj}
Purine biosynthetic pathway				
IHMA87_01837	*PA3108*	*purF*	17.26	3.64E-68
IHMA87_05343	*PA4855*	*purD*	14.06	3.85E-27
IHMA87_04292	*PA0944*	*purN*	12.99	3.14E-42
IHMA87_04291	*PA0945*	*purM*	14.04	2.51E-47
IHMA87_01191	*PA3763*	*purL*	12.75	3.02E-38
IHMA87_04214	*PA1013*	*purC*	7.67	1.23E-5
IHMA87_02487	*PA2629*	*purB*	5.35	7.23E-4
Biotin biosynthetic pathway				
IHMA87_00400	*PA0420*	*bioA*	14.15	2.29E-49
IHMA87_00477	*PA0501*	*bioF*	14.94	2.29E-49
IHMA87_00476	*PA0500*	*bioB*	14.10	1.39E-52
IHMA87_00479	*PA0503*	*bioC*	12.92	1.90E-39
IHMA87_00480	*PA0504*	*bioD*	10.80	4.23E-23
Alginate biosynthetic/regulation genes				
IHMA87_04494	*PA0762*	*algU*	11.25	3.01E-21
IHMA87_04493	*PA0763*	*mucA*	14.60	6.76E-46
IHMA87_04492	*PA0764*	*mucB*	10.69	5.56E-33
IHMA87_04490	*PA0766*	*mucD*	13.00	1.10E-45
IHMA87_00895	*PA4033*	P_{mucE}	13.07	5.88 E-22
IHMA87_05847	*PA5322*	*algC*	2.66	4.76 E-2
Other top hits				
IHMA87_02894		-	14.60	6.18E-20
IHMA87_02666	*PA2484*	-/transcriptional regulator	13.20	1.33E-36
IHMA87_06317		-	12.34	1.02E-45
IHMA87_01786	*PA3161*	*himD*	12.13	2.38E-16
IHMA87_02554	*PA2573*	-/chemotaxis transducer	11.13	1.02E-30
IHMA87_06033	*PA5490*	*cc4*	10.63	4.10E-26

IHMA87_04274		*slyX*	10.60	4.67E-13
IHMA87_04133	*PA1078*	*flgC*	10.40	4.59E-28
IHMA87_04779	*PA4457*	*kdsD*	10.24	3.15E-14
IHMA87_04779	*PA4457*	-/arabinose-5-phosphate isomerase	10.24	3.15E-14
IHMA87_00896	*PA4032*	*walR*	9.89	3.67E-24
IHMA87_00319	*PA0341*	*lgt*	9.95	5.21E-16
IHMA87_03447	*PA1758*	*pabB*	9.79	2.55E-22
IHMA87_03022	*PA2122*	-/lipid biosynthetic process	9.65	1.27E-13
IHMA87_01755		-	9.51	7.37E-23
IHMA87_04617	*PA4315*	*mvaT*	9.41	1.21E-23
IHMA87_02603		-	9.34	4.46E-17
IHMA87_02570		-	9.28	3.29E-13
IHMA87_05344	*PA4856*	*retS*	9.11	2.94E-18

Table 1. Insertion mutants enriched in plasma compared to HIP in the four main pathways. The genes and pathways showing the most significant (P_{adj}) fold change (Log$_2$(FC)) in the Tn-seq screen. IHMA87 gene identity (ID) (NCBI accession numbers CP041354 and CP04135), for the chromosome and plasmid sequences respectively (Trouillon *et al.*, 2020) are indicated with PAO1 equivalents where applicable. For each gene, the Log$_2$(FC) in plasma compared to HIP (input) condition is indicated together with corresponding adjusted *p*-value (P_{adj}), obtained by the DeSeq2 program. The genes are sorted according to their predicted pathway taken from the KEGG database and the *Pseudomonas* genome database (Kanehisa *et al.*, 2008; Winsor *et al.*, 2011).

For several mutants showing >9 Log$_2$(FC), the transposon was inserted in uncharacterized genes, representing potentially novel key factors of CS evasion (Table 1). Together, those results confirmed the role of LPS, alginates and Psl in resistance to the CS, which validated the Tn-seq approach and pointed toward several novel determinants to explore.

Deficiency in purine and biotin pathways leads to increased persistence

We further focused on hits in genes and intergenic regions of biotin and purine biosynthesis. Mutants with insertions in loci of the entire biotin biosynthesis, the *bioBFHCD* operon and the *bioA* gene, were significantly overrepresented in plasma (Table 1, Fig. 2A), suggesting that the decreased level of biotin leads to better survival in plasma. This finding was surprising, as the biotin pathway was found essential for survival of several pathogens in human serum including *Klebsiella pneumoniae*, *Mycobacterium tuberculosis* (Woong Park *et al.*, 2011; Carfrae *et al.*, 2020; Weber *et al.*, 2020). In addition to the mutants isolated from the screen (Tn::*bioA* and Tn::*bioB*), and in order to confirm this result we engineered a *bioB* deletion mutant (Δ*bioB*) and examined its behavior in plasma over a 6h-challenge (Fig. 2B). Compared to the wild-type strain, the fraction of Δ*bioB* that survived in plasma was over 10% compared to about 0.01% of parental strain. Moreover, the survival of IHMA87Δ*bioB* displayed a biphasic killing curve, suggesting that the low level of biotin triggered the persistence in human plasma, in accordance with higher survival of Tn::*bioA* and Tn::*bioB* in plasma (Fig. 1C).

A. Biotin biosynthesis genes

Purine biosynthesis genes

B.

Figure 2. Inactivation of biotin and purine biosynthetic pathways leads to higher survival rates of IHMA87 in plasma. A. Zoom on Tn-seq profiles of *bio* and *pur* genes and operons, showing number of normalized reads in input (HIP) and output (plasma) samples. **B.** Survival kinetics of IHMA87 wild-type strain, Δ*bioB* and Δ*purD* mutants in plasma over 6h incubation measured by CFU counting. Note the biphasic killing curves for the parental strain and Δ*bioB* mutant, indicating increased persistence of Δ*bioB*. The Δ*purD* mutant displayed increased tolerance compared to the parental strain (n=5).

De novo purine biosynthesis is essential for bacterial nucleotide metabolism together

with the pyrimidine pathway and is important for bacterial growth and survival in plasma

and serum (Samant *et al.*, 2008; Andersen-Civil *et al.*, 2018; Poulsen *et al.*, 2019; Zhu

et al., 2021). Similarly to biotin synthesis, in our model, mutants in the purine

biosynthetic pathway clearly displayed increased levels of *P. aeruginosa* survival in

plasma (Fig. 2A). Transposon-insertion mutants in three main operons encoding

enzymes of purine biogenesis were found overrepresented in the screen. Notably,

mutants in *purF*, encoding an amido-phospho-ribosyltransferase were significantly

enriched with a Log_2(Fold Change (FC)) of 17.26. To further study the link between

purine biosynthesis and increased plasma resistance, we designed a *purD* deletion

mutant (Δ*purD*) and investigated its killing kinetics over 6-hours in plasma. In

accordance with (Hill *et al.*, 2021) in an antibiotic context, a Δ*purD* mutant displayed

a tolerant phenotype in plasma (Fig. 2B). To note, no genes were found to be enriched

to the same level in the pyrimidine pathway, suggesting that the specific product of the

purine pathway and not global nucleotide metabolism, contributed to increased

survival in human plasma.

ATP levels influence *P. aeruginosa* tolerance to plasma

Recent studies showed that low ATP levels resulted in increased antibiotic persistence

(Shan *et al.*, 2017; Manuse *et al.*, 2021). In the same line, bacteria from the stationary

phase of growth have lower levels of intracellular ATP and form a higher proportion of

persisters when treated with antibiotics (Conlon *et al.*, 2016). In our hands, growth

phase did not contribute significantly to evaders' population suggesting that the initial

metabolic status of bacteria had no important role in persistence (Pont *et al.*, 2020).

However, because ATP is a final product of the purine pathway (Fig. 3A), we

hypothesized that it could also play a role in evaders' formation.

To tackle this question, we first measured intracellular ATP levels in ΔpurD both in rich

medium and after incubation in HIP. As expected, ΔpurD exhibited 5 times lower ATP

level than the parental strain (Fig. 3B), and no difference could be observed between

HIP and LB. To confirm that the low ATP concentration was indeed responsible for

higher resilience to human plasma, we modulated ATP levels by supplementing the

plasma with exogenous ATP (Fig. 3C) during plasma killing assay. Indeed, the addition

of ATP during the incubation in plasma restored a wild-type-like sensitivity to plasma,

further demonstrating the ATP-dependent phenotype of *pur* mutants. To further

characterize the *purD* mutant, we investigated potential envelope changes due to the

ATP depletion that could account for resistance to MAC killing at the cellular level. We

imaged and compared the different strains grown in LB medium or challenged in

plasma by transmission electron microscopy (TEM) (Fig. 3E). When grown in LB, the

wild-type strain and the Δ*bioB* and Δ*purD* mutants displayed similar size, shape and

apparent envelope thickness. However, the Δ*purD* mutant cells occasionally harbor

small electron dense granules (Fig. 3E, blue arrow). As expected from its higher

sensitivity to plasma, more cell debris was visible on the samples of the wild-type strain

incubated in plasma. Interestingly, when Δ*purD* cells were challenged by human

plasma, the size and number of the electron dense granules increased (Fig. 3E, red

arrows). In wild-type and Δ*bioB* backgrounds, plasma-challenged bacteria displayed

small granules, similar to the ones present in the Δ*purD* mutant grown in LB (Fig. 3E,

blue arrows). Bibliography mining indicated that these objects may be polyphosphate

(Poly-P) storage granules as suggested in (Racki *et al.*, 2017), whose formation is

accelerated in stress and starvation conditions. The biotin biosynthesis pathway (Fig.

3D) had no apparent link to ATP synthesis. However, as biotin impacts many biological

processes, we tested whether a Δ*bioB* mutant could lead to evader formation through

a different mechanism. As shown in Figure 3C, biotin but not ATP was able to restore

a wild-type-like sensitivity of Δ*bioB* suggesting a distinct molecular mechanism for

evader formation in this mutant.

Figure 3. ATP and biotin influence bacterial resistance to plasma. A. Schematic view of purine pathway (adapted from KEGG database (Kanehisa *et al.*, 2008)). Significantly enriched insertions in corresponding genes of the Tn-seq screen are indicated. **B.** Measurement of intra-bacterial ATP levels in LB and after 2h-incubation in HIP normalized to the CFU counts. **C.** Trans-complementation of Δ*bioB* and Δ*purD* phenotype by exogenous biotin or ATP, respectively. Biotin (1 µM) and ATP (1.5 mM) were added at the beginning of the incubation of bacteria in plasma. The survival was estimated by CFU counting. **D.** The biotin biosynthetic pathway. Significantly enriched hits in Tn-seq are highlighted in green. **E.** Transmission electron microscopy images of indicated strains after growth in LB (left) or 1h-incubation in human plasma (right). Red and blue arrow show big- and small-sized granules respectively. Scale bar = 500 nm.

Expression of the three-gene operon *srgABC* leads to better survival in plasma

One mutant isolated during the screen was significantly enriched in plasma with a Log$_2$(FC) of 13.2 (Fig. 4A). The Sanger sequencing determined the exact transposon insertion within the intergenic region between an uncharacterized predicted three-gene operon *IHMA87_01573–IHMA87_01571* (predicted in PA14 by (Wurtzel *et al.*, 2012), corresponding to *PA3369-PA3371* in PAO1) and *panM* (*IHMA87_01574/PA3368*), encoding a shorter version of a probable acetyltransferase (Fig. 4B). The isolated mutant displayed tolerant phenotype over a 6-hour period (Fig. S1) and a 100 fold increase in survival compared to the parental strain after 3h of incubation (Fig. 4C). The position of the transposon and the preliminary RT-qPCR analysis did not allow concluding on the effect of the insertion. In addition, strains carrying deletions either in *panM* or in *IHMA87_01573– IHMA87_01571* in the IHMA87 background showed no difference in survival in plasma compared to the wild-type strain (not shown). Therefore, to globally apprehend the transposon-induced changes in the mutant, we determined the transcriptomic profile of the mutant and compared it to the parental strain in LB. The RNA-Seq data showed the overexpression of the predicted *IHMA87_01573-IHMA87_01571* operon, named *srgABC* for <u>s</u>erum <u>r</u>esistance <u>g</u>enes, with about a 20-fold change compared to the parental strain. Thus, the transposon insertion mutant will be referred to as Tn::P$_{srg}$. All together 90 genes were differentially expressed (|FC|>2, p$_{adj}$<1x10^{-10}) between Tn::P$_{srg}$ and the wild-type strain (Table 2S), including *algU* encoding the alternative sigma factor AlgU and five other alginate genes (*algR* and *mucABCD* operon, Fig. 4D). Consistent with the observed upregulation of *algU*, colonies displayed characteristic mucoïd phenotype on solid medium. The *srg* operon encodes three small putative proteins of 9.9 (SrgA),

5.3 (SrgB) and 6.5 (SrgC) kDa. Based on structural predictions, SrgA is a periplasmic

or secreted protein presenting the signal peptide cleavage site at position Gly27. Both

SrgB and SrgC are predicted membrane proteins with one and two transmembrane

alpha-helices for SrgB and SrgC, respectively. The *srg* genes are highly conserved

over the 232 complete *P. aeruginosa* genomes available on *Pseudomonas* Genome

database (Winsor *et al.*, 2016) with 98-100% amino acid sequence identity.

The individual contributions of the *srg* genes in *P. aeruginosa* resilience to plasma

were evaluated by testing single, double and triple deletion mutants directly in the

Tn::P_{srg} background strain. Plasma killing assay of the mutants showed that SrgA was

required and sufficient for increased survival in plasma (Fig. 4C).

Figure 4. Overexpression of *srgA* leads to a 100-fold increase in survival. A. Zoom into the Tn-seq profiles in the region around the *srg* operon showing number of normalized reads in input (HIP) and output (plasma). **B.** Schematic representation of the predicted *srgABC* operon and position of the transposon in the isolated Tn::P_{srg} mutant as revealed by Sanger sequencing. Corresponding genes from PAO1 are indicated. Note that the transposon is inserted 13 bp upstream of the TSS **C.** Role of individual *srg* genes in the plasma resilience phenotype of the Tn::P_{srg} strain. Note that the deletion of *srgA* restores the sensitivity to plasma. **D.** Differential gene expression between Tn::P_{srg} and the parental IHMA87 strain represented in a volcano plot

showing the overexpression of *srgABC* and alginate related genes. RNA was obtained from LB-grown cultures and the whole transcriptomes were determined by RNA-seq pipeline **E-F.** Survival in plasma and alginate production of indicated strains. Alginate synthesis was visualized from colony morphology and quantified by carbazole assay, as described in Materials and methods. Note that alginate overproduction appears as darker bacteria spot (**F**). The data are from one representative experiment, performed in biological triplicates.

Contribution of *srgABC* and alginates in tolerance to complement killing

To establish the relationship between the *srg* operon, alginate production and bacterial resilience to complement, we inactivated *algD*, encoding GDP-mannose 6-dehydrogenase, a checkpoint in alginate biosynthesis (Deretic, Gill and Chakrabarty, 1987; Berry, DeVault and Chakrabarty, 1989; Wood and Ohman, 2012) both in the parental strain and in Tn::P_{srg}, and investigated relevant phenotypes. In the assay, we included alginate-overproducing mutant Tn::*mucB* isolated during the screen. We also designed a deletion mutant in the gene *IHMA87_01134* (*PA3819*), an uncharacterized gene that was overexpressed (Log$_2$(FC)=2.35, p_{adj}=2x10^{-153}) in the Tn::P_{srg} strain. For each strain, we established the survival rate by CFU counting, the quantities of alginates using carbazole assay and assessed the colony morphology on plates (Fig. 4E-F). The mutant Tn::P_{srg} displayed a 100-fold higher survival in human plasma compared to the parental strain, along with an increased alginate production. Although alginate overproduction can lead to higher resistance to human plasma, as seen for the mutant Tn::*mucB*, the alginates in Tn::P_{srg} did not account significantly for its resistance in plasma. Finally, the deletion of *IHMA87_01134* had no impact neither on alginate production nor on plasma resistance. Overall, these results demonstrate that increased plasma resistance in a *srg* overexpressing mutant is alginate-independent, and that SrgA alone modulates the plasma resistance.

The specificity of mutants toward MAC-induced killing

In order to examine the possible mechanisms leading to increased resistance to bactericidal activity of the MAC in plasma, we set up experiments to survey different steps of the complement cascade. The complement activation of both classical and alternative pathways by bacteria was examined using the measure of residual complement activity toward sensitized sheep or rabbit erythrocytes respectively, as described (Dumestre-Pérard *et al.*, 2008). Plasma incubated with the different mutants exhibited the same level of residual complement capacity to lyse red blood cells as that incubated with the parental strain (Fig. 5A-B), suggesting that complement activation was not altered by the mutants. To investigate whether the mutants are impaired in the formation of C3b and MAC (C5b-9) complexes, we first assessed the deposition of CS components at the bacterial surface by flow cytometry (Fig. 5C-H). As shown on Figure 5C-D, an increase in C3b deposition was observed at 15 min for the the Δ*bioB* mutant. When we exposed bacteria to pooled human serum supplemented with fluorescently labelled C9 (Figure 5E-F), we observed similar deposition of C9, with an apparent increase for the Tn::P_{srg} mutant. Finally, we investigated MAC polymerisation using an anti-C5b-9 antibody and flow cytometry (Fig. G-H). No significant variation in the quantity of the C5b-9 complex was detected between the different mutants, suggesting that globally, the mutants had similar amount of MAC at their surface. To assess the functionality of the inserted MAC pore, we compared serum survival of the different bacteria in the presence of nisin. Normally nisin cannot act on Gram-negative bacteria when they have an intact outer membrane (Heesterbeek *et al.*, 2019). However, MAC-induced damage to the outer-membrane allows the influx of nisin resulting in bacterial killing. As previously shown for plasma, the wild-type displays high sensitivity to killing in serum, with a very efficient bacterial

elimination (Fig. 5I). In contrast, survival of all mutants was increased. Interestingly,

nisin further killed most of the evaders' population. In a similar manner, the Tn::P_{srg}

mutant showed a reduction in numbers in presence of nisin, indicating MAC-induced

outer membrane damage. This suggests that outer membrane damage by the MAC

itself was not sufficient to kill this mutant or the evaders' population. In contrast, the

Δ$bioB$ and Δ$purD$ strain showed no decreased survival upon nisin addition to the

serum. All bacteria could be rescued by addition of C5 inhibitor OmCI, confirming that

the killing is complement-driven. Taken together, these data suggest that while the

Δ$bioB$, Δ$purD$, and Tn::P_{srg} mutants display increased survival to serum compared to

the wild-type, the Tn::P_{srg} mutant interferes at a different stage of MAC-dependent

killing than the Δ$bioB$ and Δ$purD$ mutants.

Figure 5. MAC-dependent killing in plasma resilient mutants. Residual complement activity of the classical (**A.**) and alternative (**B.**) complement pathways was measured after bacterial challenge in plasma. n=3. **C.** C3b deposition on bacterial surface after 5 or 15 min of incubation in plasma determined by FACS and **D.** plotted mean fluorescence intensities n=3 p-value < 0.001 when indicated by **. **E.** Bacterial C5b-9 deposition on bacterial surface after 5 or 15min of incubation in plasma determined by FACS and plotted mean fluorescence intensities (**F.**). n=3. **G.** C9-AF647 insertion into bacterial membrane after 30 min in 0% or 3% human pooled serum and **H.** plotted mean fluorescence intensities p-value < 0.05 when indicated by *. **I.** Bacterial survival in human pooled serum, with nisin or in presence of nisin and OmCI complement inhibitor.

Discussion

The variability of *P. aeruginosa* strains concerning their survival in human plasma is high, ranging from fully resistant to sensitive strains. The majority of strains displaying sensitivity at the whole population level form a persistent sub-population with still uncharacterized features (Pont *et al.*, 2020). In this work, we identified ATP, biotin and a small periplasmic protein, named SrgA, as novel molecular determinants of *P. aeruginosa* resilience to the CS in human plasma. In accordance with genome-wide screens performed in complement-resistant strains of different pathogens in human serum (Phan *et al.*, 2013; Sanchez-Larrayoz *et al.*, 2017; McCarthy, Stabler and Taylor, 2018; Poulsen *et al.*, 2019; Short *et al.*, 2020), our screen confirmed several previously identified factors, such as LPS, long OSA and exopolysaccharides, as being important for survival of *P. aeruginosa* in human plasma. Also, in agreement with genetic screens that identified bacterial persistence determinants facing antibiotic treatment (Molina-Quiroz *et al.*, 2016; Cameron *et al.*, 2018), we found that persistence in plasma is a multifactorial and probably a stochastic phenomenon. The mutant depleted in intracellular ATP (by a mutation in the purine pathway) showed a higher tolerance profile in plasma and became more MAC-sensitive upon external addition of ATP. In the context of higher tolerance/persistence to antibiotics, it has been proposed that the ATP-depleted conditions would decrease the global protein synthesis and availability of antibiotic targets (Shan *et al.*, 2017). Similarly, the slow growth *per se*, due to any restricted conditions triggers a persistent state (Pontes and Groisman, 2019). Although incapable of growing, Δ*bioB* and Δ*purD* mutants were able to survive and persist in HIP for more than 24 hours. Although *in vivo* mutations in the purine and biotin pathways are unlikely to occur, bacteria could use regulatory mechanisms to modulate purine and biotin synthesis and enter in an antibiotic or complement tolerant

state. Several reports highlighted the potential of targeting biotin biosynthesis as an

alternative antibacterial strategy (Carfrae *et al.*, 2020). Our results showing that the

mutants in the biotin biosynthetic pathway are more resistant to microbicidal activity of

CS call for more careful assessment of the role of biotin in BSI.

The number of plasma evaders following the 2-3h incubation varies between

experiments, strains and origin of plasma (Pont *et al.*, 2020); and this work). Therefore,

it is possible that the initial incubation in plasma, which is considered as a poor nutritive

and stressful milieu for bacteria may stochastically trigger formation of evaders.

However, our preliminary results show that pre-incubation of bacteria in HIP does not

modify the level of evaders, suggesting that the hostile nutritional environment of HIP

is not sufficient to increase the persistence. The adaptation of bacteria to human serum

includes upregulation of hundreds of genes, depending on the species. Notably,

human blood and serum upregulate quorum sensing genes in plasma-resistant *P.*

aeruginosa strain PAO1, partly through the global virulence regulator Vfr (Kruczek *et*

al., 2014, 2016; Beasley *et al.*, 2020). The serum factors involved in this regulation and

downstream effectors are still unknown.

In our search for mechanisms underlying escape of here-identified mutants from the

complement attack, we found that Tn::P_{srg}, Δ*bioB* and Δ*purD* mutants do not impede

complement activation, but probably modulate the function of MAC itself. MAC is a

multiprotein pore whose oligomerization occurs by sequential addition of C5b, C6, C7,

C8 and finally C9 which forms an oligomeric ring of 18 units inserted into bacterial

membranes (Bayly-Jones, Bubeck and Dunstone, 2017). Although the composition of

MAC has been known for years, its contribution to bacterial cell death and the way the

bacteria die upon MAC insertion is still a debate (Doorduijn, Rooijakkers and

Heesterbeek, 2019), and recent work suggested that complement-resistant bacteria

can specifically block polymerization of C9 (Doorduijn *et al.*, 2021). SrgA is a protein

of 7.1 kDa after predicted signal peptide cleavage, probably localized in the bacterial

periplasm and/or exported to the bacterial surface. We can speculate that SrgA directly

interferes with MAC assembly. Alternatively, SrgA may act on peptidoglycan or other

components of the Gram-negative envelope, which impede MAC function. Our efforts

to detect SrgA using protein tags in *P. aeruginosa* were until now unsuccessful, but

preliminary visualization of the protein in *E. coli* confirms that the *srgA* gene product is

of peptidic nature. However, the nature of *srgB* and *srgC* gene products is completely

speculative. The role of ATP in MAC activity is intriguing. No data are available on the

requirement of energy in MAC assembly, but we can speculate that it may be required

for one of the steps during pore formation. Interestingly, the mutants displayed

resistance to an outer-membrane-impermeable polycyclic antibacterial peptide nisin

(Heesterbeek *et al.*, 2019), while the majority of the evaders were sensitive. This

indicates that, in these mutants, the MAC pore did not allow the nisin incorporation

through the permeabilized outer membrane, either due to defective pore insertion or

to more resistant inner membranes. On the contrary, the evaders' outer membrane

seems to have been damaged by the MAC, suggesting that evaders possess a more

resistant envelope or have a quicker ability to repair envelope damage. Further

characterization of evaders' biology compared to the bulk population is needed to

identify molecular mechanisms triggering the evader state.

Taking into account the genetic and phenotypic diversity of *P. aeruginosa* strains

(Hilker *et al.*, 2015; Freschi *et al.*, 2018) and their variable levels of survival in human

plasma (Pont *et al.*, 2020), we can expect both common and strain-specific

determinants that counteract the MAC-dependent killing. Only parallel screens on

several representative bacterial strains may deliver the global picture on *P. aeruginosa* plasma and blood resistome/persistome.

In an era where bacterial resistance to antibiotics has become the global health concern, any alternative for pathogen elimination could be interesting to explore. The use of a complement system or MAC may be one of those, in combination with classical antibiotics, phages or antibodies that enhance MAC formation (Abd El-Aziz, Elgaml and Ali, 2019; Cruz *et al.*, 2019; Gulati *et al.*, 2019). However, we still have a very limited knowledge on different pathogens strategies to escape the complement. Better understanding of molecular mechanisms involved in pathogen-complement interplay should help designing strategies to fight multi-drug resistant bacteria.

Materials and methods

Bacterial strains and genetic manipulations

Bacterial strains, plasmids and primers are listed in Tables 3S and 4S. *E. coli* and *P. aeruginosa* were grown in Lysogeny Broth (LB) at 37°C with shaking (300 rpm). *P. aeruginosa* was selected on LB plates containing 25 µg/mL irgasan. Antibiotics concentrations were: 75 µg/mL gentamicin, 75 µg/mL tetracycline and 300 µg/mL carbenicillin for *P. aeruginosa* and 50 µg/mL gentamicin 10 µg/mL tetracycline, 100 µg/mL ampicillin for *E. coli*. To create deletion mutants, approximately 500 bp upstream and downstream flanking regions were amplified from genomic DNA by PCR using appropriate primer pairs (sF1/sR1, sF2/sR2). Overlapping fragments were cloned into *Sma*I-digested pEXG2, pEX100T or pEX18Tc by Sequence- and Ligation-Independent Cloning (SLIC, Li and Elledge, 2007). Plasmids were transformed into *E. coli* Top10 competent cells and their sequence was verified (Eurofins). Allelic

exchange vectors derived from pEXG2, pEX100T or pEX18Tc were introduced into *P.*
aeruginosa by triparental mating, using pRK600 as a helper plasmid. Merodiploids,
resulting from homologous recombinations, were selected on LB agar plate containing
irgasan and the appropriate antibiotic. Single merodiploid colonies were streaked on
NaCl-free LB plates with 10 % sucrose (w/v), selecting for plasmid loss. Resulting
sucrose resistant clones were screened for antibiotic sensitivity and gene deletion by
PCR with appropriate primer pairs (F0/R0).

Generation of the IHMA87 transposon library

The IHMA87 transposon mutant library was constructed essentially as previously
described (Skurnik *et al.*, 2013; Trouillon *et al.*, 2020). Briefly, IHMA87, grown
overnight at 42°C with shaking, was mixed with two *E. coli* strains carrying pRK2013
(Figurski and Helinski, 1979) and pBTK24. The conjugation mixtures containing 100
µl of each bacterial culture adjusted to optical densities measured at 600 nm (OD_{600})
of 1, were centrifuged, washed with LB, deposited onto pre-warmed LB plates and let
at 37°C for 5 h. Approximately 100 conjugation mixtures yielded a library of > 300,000
mutants. Bacteria were scraped off into liquid LB and aliquots were plated on LB plates
containing irgasan (25 µg/ml) and gentamicin (75 µg/ml). Following growth at 37°C,
the colonies were collected directly into LB containing 20% glycerol, aliquoted and
stored at -80 °C.

Plasma pools preparation

Pools of plasma were prepared from heparinized human blood from healthy donors
provided by the French National Blood Institute (EFS, Grenoble, France). Fresh blood
was centrifuged for 10 min at 1,000 x g at room temperature. The supernatants of ten
different donors were pooled, filtered with a 0.45 µm membrane filter and aliquoted

prior to storage at -80 °C until needed. A pool was prepared using the same procedure

from citrated blood from 30 healthy donors. Prior to use, plasma pool aliquots were

thawed on ice centrifuged 10 min at 10,000 rpm prior to filtration through 0.22 μm

membrane. For the heat inactivated plasma (HIP) condition, plasma was thawed and

heat inactivated for 30 min at 56°C and centrifuged 10 min at 10,000 rpm prior to

filtration through 0.22 μm membrane.

Screen in plasma

A vial containing the IHMA87 mutant library was thawed on ice, transferred into a flask

containing 30 mL of LB and grown 16h under agitation at 27°C to limit bacterial growth.

The next day, bacterial suspension was diluted to OD_{600nm} ~0.1 and grown at 37°C

under agitation. Meanwhile, plasma and HIP were prepared. At an OD_{600nm}~1, bacteria

were harvested and resuspended in PBS supplemented with calcium and magnesium

ions (Thermofisher Scientific) before infecting plasma, HIP or LB samples to a

concentration of 2.25×10^7 CFU/mL (90% final plasma concentration). The precise

initial CFU count of each experiment was determined by serial dilution. Samples were

incubated at 37°C for 3h on a rotating wheel. At the end of the challenge, part of the

sample was transferred to LB and grown overnight, part was spread on LB agar plates

to isolate individual mutants. Finally, bacterial survival was calculated by CFU counts

of each sample. In parallel, the same protocol was performed with the wild-type strain

for survival comparison. The next day, vials were taken from the culture and stored for

further mutant isolation. About 1×10^9 bacteria were harvested for each sample and

subjected to DNA extraction.

Illumina library construction, sequencing and data analysis

Genomic DNA. Bacterial pellets from the screen in plasma with either the HIP samples (Input) or the plasma samples (Output) were used for gDNA preparations. Classical protocol using SDS-NaCl lysis (2% SDS, 0.15M NaCl, 0.1M EDTA pH8, 0.6M Sodium perchlorate) step followed by two successive phenol-chloroform extractions was carried out. The gDNA was precipitated with 100% cold ethanol and diluted in a 200µl TE buffer (10mM Tris-HCl pH8, 1mM EDTA) to final concentration of 50-100 ng/µl. Mechanical DNA shearing was done at 4°C with a QSonica sonicator (Q700) during 20 min with 15 second pulses. Fragments were concentrated on columns (Monarch PCR & DNA Cleanup Kit, NEB). The size of generated DNA fragments (150-400 bp) was controlled on 2% agarose gel.

Library construction. Fragmented gDNA (1 µg) were end-repaired using End-repair module (NEB#E6050) followed by a dA-tailing with Klenow (NEB#6053). Short and Long adaptors (Table 1S) were annealed in 10 mM $MgCl_2$ in thermocycler with a program allowing decrease of 1°C/cycle, from 95°C to 20°C. Ligation of gDNA fragments with annealed adaptors was achieved overnight at 16°C with T4 DNA ligase (NEB#M0202). A size selection was performed on a 2% agarose gel by extracting the bands from 200 to 400 bp (Monarch DNA Gel Extraction Kit, NEB#T1020L). Amplification of purified adaptor-DNA fragments was carried out with Phusion polymerase (NEB#M0530) and specific primers (PCR1 Tn-spe direct and PCR1 Adaptor comp.). A second amplification step was performed with a primer carrying a P5 illumina sequence and Indexed-P7 illumina primers (NEB#E7335S). Primer sequences are available on request. After each step, quantification was performed using the Qubit dsDNA HS Assay kit (Q33230). The quality of the libraries was assessed on the Agilent Bioanalyzer 2100 using high-sensitivity DNA chips (Ref

#5067-4626). Sequencing was performed on an Illumina NextSeq High (I2BC, Saclay

Paris).

Data analysis. Sequencing reads were trimmed and aligned to the genome of IHMA87

using Bowtie2 (Langmead and Salzberg, 2012). Htseq-count (Anders, Pyl and Huber,

2015) was then used to obtain read counts per feature. Differential representation of

insertion mutants between the plasma and HIP sample was performed using (Love,

Huber and Anders, 2014). For the analysis on insertions in intergenic regions, an

annotation file was created with intergenic regions attributed to the downstream gene

independently of its orientation. The same analysis as above was performed using

Htseq-count and DESeq2. Tn-seq was performed in biological duplicates.

Plasma killing assay

Killing assay was performed as described previously (Pont *et al.*, 2020). Unless stated

otherwise, bacterial pre-cultures were diluted in LB to OD_{600nm} = 0.1 and grown at 37°C

under agitation until OD_{600nm} ~ 1. Bacteria were resuspended into PBS +/+

(Thermofisher Scientific) and incubated in plasma to a concentration of 2.25×10^7

CFU/mL (90 % final plasma concentration). The precise initial CFU count of each

experiment was determined by serial dilution. Bacterial survival was calculated by CFU

counts at the indicated time(s) and compared to the initial count. ATP, biotin and purine

were added to the plasma prior to bacterial addition at the indicated concentrations.

Transcriptomics

RNA extraction. Bacterial cultures were diluted to an OD_{600nm} of 0.1 and grown at

37°C under agitation until they reached an OD_{600nm} of 1. RNA isolation was performed

using the hot phenol-chloroform method. Briefly, bacteria were lysed in a hot phenol

lysis solution (2 mM EDTA, 1 % SDS, 40mM sodium acetate in acid phenol (Invitrogen

#15594-047)). RNA was then isolated by successive steps of phenol-chloroform

extraction, followed by one cold chloroform extraction and ethanol precipitations. An

additional Turbo DNAse (Invitrogen #AM1907) treatment was performed to remove

genomic DNA contamination. Total RNA quantity and quality were assessed using a

Agilent Bioanalyzer.

RNA-seq libraries construction. Following RNA extraction, ribosomal RNAs (rRNAs)

depletion was performed using ribominus kit (Invitrogen) according to manufacturer's

instructions. Depletion quality was verified on an Agilent Bioanalyzer using RNA pico

chips and RNA was concentrated using ethanol precipitation.

NEBNext Ultra II directional RNA library prep kit for illumina (NEB) was used to build

cDNA libraries for sequencing starting from 50ng of rRNA depleted RNA according to

manufacturer's instructions. cDNA libraries quality and concentration were assessed

using an Agilent Bioanalyzer High sensitivity DNA chips.

Sequencing and data analysis. Sequencing was performed by the Institute for

Integrative Biology of the Cell (I2BC http://www.i2bc.paris-saclay.fr) high-throughput

sequencing facility using an illumina NextSeq500 instrument. More than 12 millions

75-bp single-end reads were obtained per sample. Processed reads were mapped

onto the IHMA87 genome (available on NCBI CP041354 and CP041355 accession

numbers, for the chromosome and plasmid sequences respectively (Trouillon *et al.*,

2020)) using Bowtie2 (Langmead and Salzberg, 2012) and the readcount per feature

was calculated using Ht-seq count (Anders, Pyl and Huber, 2015). Finally, differential

gene expression between the Tn::P_{srg} and the wild-type strains was performed using

DESeq2 (Love, Huber and Anders, 2014). RNA-seq experiment was performed on

biological duplicates.

Alginate dosage

Equivalent amount of OD_{600nm} adjusted bacterial cultures were spread onto

Pseudomonas isolation agar (PIA) plates and incubated for 24h at 37°C. Bacterial

lawns were flushed with saline solution (0.9% NaCl) and scraped off the plates.

Bacteria were pelleted and supernatant was stored in a separated tube. Bacterial pellet

was resuspended in LB, incubated on a rotating wheel for 1-2 hours to avoid bacterial

aggregates and bacterial titer was calculated by serial dilutions and subsequent CFU

counting. The alginate dosage was done as described in Knutson and Jeanes, 1968

and May and Chakrabarty, 1994. Briefly, 600μL of ice cold borate working solution

(borate stock solution (4 M BO_3^{3-}) diluted to a final concentration of 100 mM in

H_2SO_4) were added to 70 μL of the bacteria supernatant, saline solution for blank or

commercial sodium alginate solution as standards. Keeping on ice, 20 μL of carbazole

solution (0.1% carbazole (Sigma #442506) in 100% ethanol) were added, and samples

were incubated at 55°C for 30 min after mixing. The absorbance was measured at 530

nm.

Quantitative measurement of intracellular bacterial ATP

For ATP measurement in LB, about 1×10^8 bacteria were pelleted and resuspended

in PBS. HIP was inoculated with a final concentration of 2.025×10^7 bacterial/mL in

PBS (90% final HIP concentration). Samples were incubated for 2h at 37°C on a

rotating wheel. At the end of the incubation, samples were centrifuged and

resuspended in PBS prior to a gentle sonication step using Qsonica sonicator (Q700)

for 45 sec with 10% intensity and 5 sec ON-5 sec OFF intervals in order to dissociate

the bacterial aggregates. For the two conditions, 10-fold serial dilutions were prepared in 96-well plates and precise bacterial counts were determined by counting CFUs. The concentration of ATP was determined using the BacTiter-Glo kit (Promega, #G8230). Briefly, equal volumes of bacterial suspension and BacTiter-Glo reagent were mixed in white 96-well Lumitrac plates (BioRad #655075) and incubated for 5 min with shaking. An ATP standard curve (0.1 to 100 nM) was established for each experiment. Bioluminescence was measured on a plate reader (Spark, TECAN). ATP concentration was then reported to the CFU counts. Experiments were performed in more than 3 biological replicates.

Residual complement activity

Sample preparation. Bacteria grown to $0D_{600nm} \sim 1$ were resuspended into PBS +/+ (Thermofisher Scientific) and incubated in plasma to a concentration of 2.25×10^9 CFU/mL (90% final plasma concentration) or PBS only for the reference. Samples were incubated for 3 hours at 37°C on a rotating wheel to allow complement activation by bacteria. The initial CFU count was determined by serial dilution. At the end of the incubation, samples were centrifuged and supernatant was filtered through 0.22 µm membrane to remove any remaining bacteria. Samples were aliquoted and frozen at -80°C until use. The remaining complement activity in the bacterial/plasma supernatant was measured using a total hemolytic assay for classical and alternative pathways, as described by (Dumestre-Pérard *et al.*, 2008).

Classical complement hemolytic activity. Briefly, sheep erythrocytes (Orgentec) were sensitized using anti-erythrocytes antibodies (1/40000 - Hemolysin, Orgentec) for 15min at 37°C. Bacterial/plasma supernatant (25 µL) was mixed with 3 mL of sensitized erythrocytes (4×10^6 cells/mL) diluted in DGVB^{2+} (2.5% glucose, 0.05%

gelatin, 2.5 mM Veronal, 72.5 mM NaCl, 0.15 mM Ca^{2+}, 0.5 mM Mg^{2+}) and lysis kinetic

was monitored by OD_{660nm} measurement using a spectrophotometer (safas UVmc2).

Area under the curve was measured for each series and reported to the 90% plasma

pool control.

Alternative complement hemolytic activity. Briefly, rabbit erythrocytes (Ecole

Nationale Vétérinaire de Toulouse) were washed 3 times using DGVB-Mg-EGTA

(2.5% glucose, 0.05% gelatin, 2.5 mM Veronal, 72.5 mM NaCl, 5 mM Mg^{2+}, 3 mM

EGTA). Bacterial/plasma supernatant (150 µL) was added to 3 mL of rabbit

erythrocytes suspension (diluted in DGVB-Mg-EGTA at 1/250) at 37°C and lysis kinetic

was measured similarly to classical complement activity.

C3b and C5b-9 binding assays

Forty millions of bacteria were incubated during 15 min in 90% heparinized plasma

pool at 37°C. Incubation was stopped by dilution of bacteria suspension in ten volumes

of PBS containing 20 mM of EDTA pH8. C3b binding was detected with PE-conjugated

mouse monoclonal anti-C3b/iC3b (5 µg/mL, BioLegend #846103) and C5b-9

membrane deposition/insertion was detected with mouse monoclonal anti-C5b-9

antibody (8 µg/mL, Abcam #ab66768) next revealed with PE-conjugated goat anti-

mouse antibodies (2.5 µg/mL, Abcam #ab97041). Bacteria in 0% plasma were used

as negative control. Samples were analyzed in a FACSCalibur instrument (Becton

Dickinson) and flow cytometry analysis were performed using the FCSExpress6

Software.

C9 binding

P. aeruginosa strains expressing green fluorescent protein (GFP) under the control of the constitutive P_{X2} promoter were grown overnight in LB supplemented with 10 µg/ml tetracycline at 37°C shaking. Bacteria were subcultured 1:30 in LB containing the antibiotic and let grow to a mid-log OD_{600nm} of 0.5 at 37°C with shaking. Bacteria were pelleted by centrifugation, washed, and resuspended to a final OD_{600nm} of 0.05 in RPMI + 0.05% human serum albumin (HSA) for the experiments. Bacteria were then exposed to pooled human serum supplemented with 100 nM C9-AF647 and incubated for 30 minutes at 37°C shaking. For CFU counting, serial bacterial dilutions were made in MilliQ water, spotted on LB plates, and incubated overnight at 37°C. For flow cytometry, bacteria were diluted 10-fold in RPMI-HSA and analyzed in a MACSquant VYB (Miltenyi Biotech) flow cytometer. Bacteria were gated on GFP-expression and subsequently analyzed for AF647 fluorescence. Flow cytometry data was analyzed in FlowJo V.10.

Nisin assay

Bacteria were grown overnight in LB at 37°C shaking, subcultured 1:30 in LB to a mid-log OD_{600nm} of 0.5 at 37°C shaking, pelleted by centrifugation, washed, and resuspended to a final OD_{600nm} of 0.05 in RPMI + 0.05% HSA for the experiments. Bacteria were then exposed to human pooled supplemented with buffer, 3 µg/ml nisin, or 3 µg/ml nisin and 10 µg/ml OmCI, and incubated for 3 hours at 37°C. Serial dilutions were made in MilliQ water, and spotted on LB plates for incubation overnight at 37°C for determination of CFU the next day.

Transmission electron microscopy

Bacterial pre-cultures were diluted to an OD_{600nm} of 0.1 in flasks containing 30 mL of LB and incubated at 37°C under agitation. Plasma was then inoculated with bacterial suspension to reach a bacterial concentration of 1.125×10^9 bacteria/mL and incubated on a rotating wheel for 1 hour at 37°C. At the end of the challenge, bacteria were pelleted by centrifugation and resuspended in LB prior to fixation. At the same time, bacteria grown in LB to an OD_{600nm} of 1 were pelleted by centrifugation and resuspended in LB prior to fixation.

Sample preparation was treated as described in (Mohamed *et al.*, 2021). Briefly, the cell pellet was spread on the 200-µm side of a 3-mm type A gold plate (Leica Microsystems) covered with the flat side of a 3-mm type B aluminium plate (Leica Microsystems) and vitrified by high-pressure freezing using an HPM100 system (Leica Microsystems). After freezing at -90°C in acetone with 1% OsO4, the samples were slowly warmed to -60°C, stored for 12 hours, then the temperature was increased to -30°C and the samples stored for a further 12 hours before a step at 0°C for 1 hour before being cooled again to -30°C (AFS2; Leica Microsystems). The samples were rinsed 4 times in pure acetone before being infiltrated with progressively increasing concentrations of resin (Epoxy embedding medium, Sigma) in acetone while increasing the temperature to 20°C. Pure resin was added at room temperature. After polymerisation, 70 nm thin sections were obtained using a UC7 ultramicrotome (Leica Microsystems) and collected on 100 mesh copper grids coated with Formvar carbon. The thin sections were post-stained for 5 minutes with 2% uranyl acetate, rinsed with water and incubated for 2 minutes with lead citrate. Samples were observed using a Tecnai G2 Spirit BioTwin microscope (FEI) operating at 120 kV with an Orius SC1000B CCD camera (Gatan).

Bioinformatic analysis and protein conservation

Protein sequence files from all *P. aeruginosa* with a closed assembled genome were retrieved from Pseudomonas genome database and *srg* conservation was investigated using Blast Reciprocal Best Hits program (Cock *et al.*, 2015) computed with a 90% coverage and 50% identity.

Signal peptide prediction was perfomed using SignalP website (Almagro Armenteros *et al.*, 2019).

Statistical analysis

Sigmaplot software wasused to perform statistical analysis. Multiple groups comparison was performed using one-way ANOVA or Kruskal-Wallis tests depending on data normality followed by a Student-Newman-Keuls pairwise comparison. Non-normally distributed data were converted to normally-distributed dataset by $\log_{10}$-transformation where indicated. Figures were designed using Graphpad Prism9.

Data availability

The Tn-seq and RNA-seq data generated in this work is available at GEO accession numbers GSExxxxxx and GSExxxxxx respectively.

Acknowledgments

The work was supported by grants from Agence Nationale de la Recherche (ANR-15-CE11-0018-01), the Laboratory of Excellence GRAL, financed within the University Grenoble Alpes graduate school (Écoles Universitaires de Recherche) CBH-EUR-GS (ANR-17-EURE-0003), the Fondation pour la Recherche Médicale (Team FRM 2017, DEQ20170336705) to I.A., the European Union's Horizon 2020 research programs H2020-EU-ITN-EJD (CORVOS #860044 to F.M. and SHMR). The sequencing part

has benefited from the facilities and expertise of the high throughput sequencing core

facility of I2BC (Centre de Recherche de Gif – http://www.i2bc.paris-saclay.fr/). *P.*

aeruginosa IHMA87 was obtained from the International Health Management

Association, USA. This work used the platforms of the Grenoble Instruct-ERIC center

(ISBG; UAR 3518 CNRS-CEA-UGA-EMBL) within the Grenoble Partnership for

Structural Biology (PSB), supported by FRISBI (ANR-10-INBS-0005-02) and GRAL,

financed within the University Grenoble Alpes graduate school (Ecoles Universitaires

de Recherche) CBH-EUR-GS (ANR-17-EURE-0003). M. JM received a Ph.D.

fellowship from the French Ministry of Education and Research. The IBS Electron

Microscope facility is supported by the Auvergne Rhône-Alpes Region, the Fonds

Feder, the Fondation pour la Recherche Médicale and GIS-IBiSA. The authors thank

Benoît Gallet and Emily Rey for their help for TEM and mutant analyses, respectively.

M.JM thanks Tweeter followers for indicating literature on PolyP granules.

Bibliography

Abd El-Aziz, A.M., Elgaml, A. and Ali, Y.M. (2019) 'Bacteriophage Therapy Increases
Complement-Mediated Lysis of Bacteria and Enhances Bacterial Clearance After
Acute Lung Infection With Multidrug-Resistant Pseudomonas aeruginosa', *The Journal*
of Infectious Diseases, 219(9), pp. 1439–1447. doi:10.1093/infdis/jiy678.

Almagro Armenteros, J.J. *et al.* (2019) 'SignalP 5.0 improves signal peptide predictions
using deep neural networks', *Nature Biotechnology*, 37(4), pp. 420–423.
doi:10.1038/s41587-019-0036-z.

Anders, S., Pyl, P.T. and Huber, W. (2015) 'HTSeq—a Python framework to work with
high-throughput sequencing data', *Bioinformatics*, 31(2), pp. 166–169.
doi:10.1093/bioinformatics/btu638.

Andersen-Civil, A.I.S. *et al.* (2018) 'The impact of inactivation of the purine biosynthesis
genes, purN and purT, on growth and virulence in uropathogenic E. coli', *Molecular*
Biology Reports, 45(6), pp. 2707–2716. doi:10.1007/s11033-018-4441-z.

Ayrapetyan, M. *et al.* (2015) 'Viable but Nonculturable and Persister Cells Coexist Stochastically and Are Induced by Human Serum', *Infection and Immunity*, 83(11), pp. 4194–4203. doi:10.1128/IAI.00404-15.

Balaban, N.Q. *et al.* (2019) 'Definitions and guidelines for research on antibiotic persistence (vol 17, pg 441, 2019)', *Nature Reviews Microbiology*, 17(7), pp. 460–460. doi:10.1038/s41579-019-0207-4.

Bartell, J.A. *et al.* (2020) 'Bacterial persisters in long-term infection: Emergence and fitness in a complex host environment', *PLoS pathogens*, 16(12), p. e1009112. doi:10.1371/journal.ppat.1009112.

Bayly-Jones, C., Bubeck, D. and Dunstone, M.A. (2017) 'The mystery behind membrane insertion: a review of the complement membrane attack complex', *Philosophical Transactions of the Royal Society B: Biological Sciences*, 372(1726), p. 20160221. doi:10.1098/rstb.2016.0221.

Beasley, K.L. *et al.* (2020) 'During bacteremia, Pseudomonas aeruginosa PAO1 adapts by altering the expression of numerous virulence genes including those involved in quorum sensing', *PLOS ONE*. Edited by R.M. Roop, 15(10), p. e0240351. doi:10.1371/journal.pone.0240351.

de Bentzmann, S. *et al.* (2012) 'Unique Biofilm Signature, Drug Susceptibility and Decreased Virulence in Drosophila through the Pseudomonas aeruginosa Two-Component System PprAB', *PLoS Pathogens*. Edited by D.S. Schneider, 8(11), p. e1003052. doi:10.1371/journal.ppat.1003052.

Berry, A., DeVault, J.D. and Chakrabarty, A.M. (1989) 'High osmolarity is a signal for enhanced algD transcription in mucoid and nonmucoid Pseudomonas aeruginosa strains', *Journal of Bacteriology*, 171(5), pp. 2312–2317. doi:10.1128/jb.171.5.2312-2317.1989.

Blom, A.M. *et al.* (2001) 'A Novel Interaction Between Type IV Pili of Neisseria gonorrhoeae and the Human Complement Regulator C4b-Binding Protein', *The Journal of Immunology*, 166(11), pp. 6764–6770. doi:10.4049/jimmunol.166.11.6764.

Broder, U.N., Jaeger, T. and Jenal, U. (2017) 'LadS is a calcium-responsive kinase that induces acute-to-chronic virulence switch in Pseudomonas aeruginosa', *Nature Microbiology*, 2(1), p. 16184. doi:10.1038/nmicrobiol.2016.184.

Cameron, D.R. *et al.* (2018) 'A Genetic Determinant of Persister Cell Formation in Bacterial Pathogens', *Journal of Bacteriology*. Edited by G. O'Toole, 200(17). doi:10.1128/JB.00303-18.

Carfrae, L.A. *et al.* (2020) 'Mimicking the human environment in mice reveals that inhibiting biotin biosynthesis is effective against antibiotic-resistant pathogens', *Nature Microbiology*, 5(1), pp. 93–101. doi:10.1038/s41564-019-0595-2.

Chiang, S.L. *et al.* (1995) 'Single amino acid substitutions in the N-terminus of Vibrio cholerae TcpA affect colonization, autoagglutination, and serum resistance', *Molecular Microbiology*, 17(6), pp. 1133–1142. doi:10.1111/j.1365-2958.1995.mmi_17061133.x.

Cock, P.J.A. *et al.* (2015) 'NCBI BLAST+ integrated into Galaxy', *GigaScience*, 4(1), pp. s13742-015-0080–7. doi:10.1186/s13742-015-0080-7.

Conlon, B.P. *et al.* (2016) 'Persister formation in Staphylococcus aureus is associated with ATP depletion', *Nature Microbiology*, 1(5), pp. 1–7. doi:10.1038/nmicrobiol.2016.51.

Cruz, J.W. *et al.* (2019) 'A novel bispecific antibody platform to direct complement activity for efficient lysis of target cells', *Scientific Reports*, 9(1), p. 12031. doi:10.1038/s41598-019-48461-1.

Damron, F.H. and Goldberg, J.B. (2012) 'Proteolytic regulation of alginate overproduction in Pseudomonas aeruginosa: Proteolytic regulation of alginate', *Molecular Microbiology*, 84(4), pp. 595–607. doi:10.1111/j.1365-2958.2012.08049.x.

Deretic, V., Gill, J.F. and Chakrabarty, A.M. (1987) '*Pseudomonas aeruginosa* infection in cystic fibrosis: nucleotide sequence and transcriptional regulation of the *algD* gene', *Nucleic Acids Research*, 15(11), pp. 4567–4581. doi:10.1093/nar/15.11.4567.

Doorduijn, D.J. *et al.* (2021) 'Polymerization of C9 enhances bacterial cell envelope damage and killing by membrane attack complex pores', *PLOS Pathogens*, 17(11), p. e1010051. doi:10.1371/journal.ppat.1010051.

Doorduijn, D.J., Rooijakkers, S.H.M. and Heesterbeek, D.A.C. (2019) 'How the Membrane Attack Complex Damages the Bacterial Cell Envelope and Kills Gram-Negative Bacteria', *BioEssays*, 41(10), p. 1900074. doi:10.1002/bies.201900074.

Dumestre-Pérard, C. *et al.* (2008) '*Aspergillus* Conidia Activate the Complement by the Mannan-Binding Lectin C2 Bypass Mechanism', *The Journal of Immunology*, 181(10), pp. 7100–7105. doi:10.4049/jimmunol.181.10.7100.

Fauvart, M., De Groote, V.N. and Michiels, J. (2011) 'Role of persister cells in chronic infections: clinical relevance and perspectives on anti-persister therapies', *Journal of Medical Microbiology*, 60(Pt 6), pp. 699–709. doi:10.1099/jmm.0.030932-0.

Figurski, D.H. and Helinski, D.R. (1979) 'Replication of an origin-containing derivative of plasmid RK2 dependent on a plasmid function provided in trans', *Proceedings of the National Academy of Sciences of the United States of America*, 76(4), pp. 1648–1652.

Freschi, L. *et al.* (2018) 'Genomic characterisation of an international Pseudomonas aeruginosa reference panel indicates that the two major groups draw upon distinct mobile gene pools', *FEMS Microbiology Letters*, 365(14). doi:10.1093/femsle/fny120.

Gaca, A.O., Colomer-Winter, C. and Lemos, J.A. (2015) 'Many Means to a Common End: the Intricacies of (p)ppGpp Metabolism and Its Control of Bacterial Homeostasis', *Journal of Bacteriology*, 197(7), pp. 1146–1156. doi:10.1128/JB.02577-14.

Giraud, C. *et al.* (2011) 'The PprA-PprB two-component system activates CupE, the first non-archetypal Pseudomonas aeruginosa chaperone-usher pathway system

assembling fimbriae: P. aeruginosa CupE fimbriae', *Environmental Microbiology*, 13(3), pp. 666–683. doi:10.1111/j.1462-2920.2010.02372.x.

Goldberg, J.B. and Pler, G.B. (1996) 'Pseudomonas aeruginosa lipopolysaccharides and pathogenesis', *Trends in Microbiology*, 4(12), pp. 490–494. doi:10.1016/s0966-842x(97)82911-3.

Gulati, S. *et al.* (2019) 'Complement alone drives efficacy of a chimeric antigonococcal monoclonal antibody', *PLOS Biology*. Edited by M.K. Waldor, 17(6), p. e3000323. doi:10.1371/journal.pbio.3000323.

Harms, A., Maisonneuve, E. and Gerdes, K. (2016) 'Mechanisms of bacterial persistence during stress and antibiotic exposure', *Science*, 354(6318). doi:10.1126/science.aaf4268.

Heesterbeek, D. a. C. *et al.* (2019) 'Complement-dependent outer membrane perturbation sensitizes Gram-negative bacteria to Gram-positive specific antibiotics', *Scientific Reports*, 9(1), p. 3074. doi:10.1038/s41598-019-38577-9.

Helaine, S. *et al.* (2014) 'Internalization of Salmonella by macrophages induces formation of nonreplicating persisters', *Science (New York, N.Y.)*, 343(6167), pp. 204–208. doi:10.1126/science.1244705.

Hilker, R. *et al.* (2015) 'Interclonal gradient of virulence in the Pseudomonas aeruginosa pangenome from disease and environment', *Environmental Microbiology*, 17(1), pp. 29–46. doi:10.1111/1462-2920.12606.

Hill, P.W.S. *et al.* (2021) 'The vulnerable versatility of Salmonella antibiotic persisters during infection', *Cell Host & Microbe* [Preprint]. doi:10.1016/j.chom.2021.10.002.

Jones, C.J. and Wozniak, D.J. (2017) 'Psl Produced by Mucoid Pseudomonas aeruginosa Contributes to the Establishment of Biofilms and Immune Evasion', *mBio*. Edited by J.B. Goldberg, 8(3), pp. e00864-17, /mbio/8/3/e00864-17.atom. doi:10.1128/mBio.00864-17.

Kanehisa, M. *et al.* (2008) 'KEGG for linking genomes to life and the environment', *Nucleic Acids Research*, 36(suppl_1), pp. D480–D484. doi:10.1093/nar/gkm882.

Kang, C.-I. *et al.* (2003) 'Pseudomonas aeruginosa Bacteremia: Risk Factors for Mortality and Influence of Delayed Receipt of Effective Antimicrobial Therapy on Clinical Outcome', *Clinical Infectious Diseases*, 37(6), pp. 745–751. doi:10.1086/377200.

Keren, I. *et al.* (2004) 'Persister cells and tolerance to antimicrobials', *FEMS Microbiology Letters*, 230(1), pp. 13–18. doi:10.1016/S0378-1097(03)00856-5.

Kintz, E. *et al.* (2008) 'Lipopolysaccharide O-Antigen Chain Length Regulation in Pseudomonas aeruginosa Serogroup O11 Strain PA103', *Journal of Bacteriology*, 190(8), pp. 2709–2716. doi:10.1128/JB.01646-07.

Knutson, C.A. and Jeanes, A. (1968) 'A new modification of the carbazole analysis: application to heteropolysaccharides', *Anal Biochem*, 24(3), pp. 470–81. doi:10.1016/0003-2697(68)90154-1.

Korch, S.B., Henderson, T.A. and Hill, T.M. (2003) 'Characterization of the hipA7 allele of Escherichia coli and evidence that high persistence is governed by (p)ppGpp synthesis', *Molecular Microbiology*, 50(4), pp. 1199–1213. doi:10.1046/j.1365-2958.2003.03779.x.

Kruczek, C. *et al.* (2014) 'Serum influences the expression of Pseudomonas aeruginosa quorum-sensing genes and QS-controlled virulence genes during early and late stages of growth', *MicrobiologyOpen*, 3(1), pp. 64–79. doi:10.1002/mbo3.147.

Kruczek, C. *et al.* (2016) 'Major Transcriptome Changes Accompany the Growth of Pseudomonas aeruginosa in Blood from Patients with Severe Thermal Injuries', *PLOS ONE*. Edited by R.M. Roop II, 11(3), p. e0149229. doi:10.1371/journal.pone.0149229.

Langmead, B. and Salzberg, S.L. (2012) 'Fast gapped-read alignment with Bowtie 2', *Nature Methods*, 9(4), pp. 357–359. doi:10.1038/nmeth.1923.

Li, M.Z. and Elledge, S.J. (2007) 'Harnessing homologous recombination in vitro to generate recombinant DNA via SLIC', *Nature Methods*, 4(3), pp. 251–256. doi:10.1038/nmeth1010.

Love, M.I., Huber, W. and Anders, S. (2014) 'Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2', *Genome Biology*, 15(12), p. 550. doi:10.1186/s13059-014-0550-8.

Manuse, S. *et al.* (2021) 'Bacterial persisters are a stochastically formed subpopulation of low-energy cells', *PLOS Biology*. Edited by T. Bollenbach, 19(4), p. e3001194. doi:10.1371/journal.pbio.3001194.

May, T.B. and Chakrabarty, A.M. (1994) '[22] Isolation and assay of Pseudomonas aeruginosa alginate', in *Methods in Enzymology*. Academic Press (Bacterial Pathogenesis Part A: Identification and Regulation of Virulence Factors), pp. 295–304. doi:10.1016/0076-6879(94)35148-1.

McCarthy, A.J., Stabler, R.A. and Taylor, P.W. (2018) 'Genome-Wide Identification by Transposon Insertion Sequencing of Escherichia coli K1 Genes Essential for In Vitro Growth, Gastrointestinal Colonizing Capacity, and Survival in Serum', *Journal of Bacteriology*, 200(7). doi:10.1128/JB.00698-17.

Mishra, M. *et al.* (2012) 'Pseudomonas aeruginosa Psl polysaccharide reduces neutrophil phagocytosis and the oxidative response by limiting complement-mediated opsonization', *Cellular Microbiology*, 14(1), pp. 95–106. doi:10.1111/j.1462-5822.2011.01704.x.

Mohamed, A.M.T. *et al.* (2021) 'Chromosome Segregation and Peptidoglycan Remodeling Are Coordinated at a Highly Stabilized Septal Pore to Maintain Bacterial Spore Development', *Developmental Cell*, 56(1), pp. 36-51.e5. doi:10.1016/j.devcel.2020.12.006.

Molina-Quiroz, R.C. *et al.* (2016) 'Transposon-Sequencing Analysis Unveils Novel Genes Involved in the Generation of Persister Cells in Uropathogenic Escherichia coli', *Antimicrobial Agents and Chemotherapy*, 60(11), pp. 6907–6910. doi:10.1128/AAC.01617-16.

Moradali, M.F., Ghods, S. and Rehm, B.H.A. (2017) 'Pseudomonas aeruginosa Lifestyle: A Paradigm for Adaptation, Survival, and Persistence', *Frontiers in Cellular and Infection Microbiology*, 7, p. 39. doi:10.3389/fcimb.2017.00039.

Osawa, K. *et al.* (2013) 'Modulation of O-antigen chain length by the wzz gene in Escherichia coli O157 influences its sensitivities to serum complement', *Microbiol Immunol*, 57(9), pp. 616–23. doi:10.1111/1348-0421.12084.

Pacios, O. *et al.* (2020) '(p)ppGpp and Its Role in Bacterial Persistence: New Challenges', *Antimicrobial Agents and Chemotherapy*, 64(10), pp. e01283-20. doi:10.1128/AAC.01283-20.

Pedersen, S.S. *et al.* (1990) 'Pseudomonas aeruginosa alginate in cystic fibrosis sputum and the inflammatory response', *Infection and Immunity*, 58(10), pp. 3363–3368. doi:10.1128/iai.58.10.3363-3368.1990.

Phan, M.-D. *et al.* (2013) 'The serum resistome of a globally disseminated multidrug resistant uropathogenic Escherichia coli clone', *PLoS genetics*, 9(10), p. e1003834. doi:10.1371/journal.pgen.1003834.

Pier, G.B. *et al.* (2001) 'Role of alginate O acetylation in resistance of mucoid Pseudomonas aeruginosa to opsonic phagocytosis', *Infection and Immunity*, 69(3), pp. 1895–1901. doi:10.1128/IAI.69.3.1895-1901.2001.

Pont, S. *et al.* (2020) 'Bacterial behavior in human blood reveals complement evaders with some persister-like features', *PLoS Pathog*, 16(12), p. e1008893. doi:10.1371/journal.ppat.1008893.

Pontes, M.H. and Groisman, E.A. (2019) 'Slow growth determines nonheritable antibiotic resistance in Salmonella enterica', *Sci Signal*, 12(592). doi:10.1126/scisignal.aax3938.

Poulsen, B.E. *et al.* (2019) 'Defining the core essential genome of Pseudomonas aeruginosa', *Proceedings of the National Academy of Sciences*, 116(20), pp. 10072–10080. doi:10.1073/pnas.1900570116.

Priebe, G.P. *et al.* (2004) 'The galU Gene of Pseudomonas aeruginosa is required for corneal infection and efficient systemic spread following pneumonia but not for infection confined to the lung', *Infection and Immunity*, 72(7), pp. 4224–4232. doi:10.1128/IAI.72.7.4224-4232.2004.

Putrinš, M. *et al.* (2015) 'Phenotypic Heterogeneity Enables Uropathogenic Escherichia coli To Evade Killing by Antibiotics and Serum Complement', *Infection and Immunity* [Preprint]. Available at: https://journals.asm.org/doi/abs/10.1128/IAI.02725-14 (Accessed: 20 December 2021).

Qiu, D. *et al.* (2007) 'Regulated proteolysis controls mucoid conversion in Pseudomonas aeruginosa', *Proceedings of the National Academy of Sciences*, 104(19), pp. 8107–8112. doi:10.1073/pnas.0702660104.

Racki, L.R. *et al.* (2017) 'Polyphosphate granule biogenesis is temporally and functionally tied to cell cycle exit during starvation in Pseudomonas aeruginosa', *Proceedings of the National Academy of Sciences*, 114(12), pp. E2440–E2449. doi:10.1073/pnas.1615575114.

Samant, S. *et al.* (2008) 'Nucleotide Biosynthesis Is Critical for Growth of Bacteria in Human Blood', *PLoS Pathogens*. Edited by J.E. Galán, 4(2), p. e37. doi:10.1371/journal.ppat.0040037.

Sanchez-Larrayoz, A.F. *et al.* (2017) 'Complexity of Complement Resistance Factors Expressed by Acinetobacter baumannii Needed for Survival in Human Serum', *Journal of Immunology (Baltimore, Md.: 1950)*, 199(8), pp. 2803–2814. doi:10.4049/jimmunol.1700877.

Shan, Y. *et al.* (2017) 'ATP-Dependent Persister Formation in *Escherichia coli*', *mBio*. Edited by K. Bush, 8(1). doi:10.1128/mBio.02267-16.

Short, F.L. *et al.* (2020) 'Genomic Profiling Reveals Distinct Routes To Complement Resistance in Klebsiella pneumoniae', *Infection and Immunity*, 88(8). doi:10.1128/IAI.00043-20.

Skurnik, D. *et al.* (2013) 'A Comprehensive Analysis of In Vitro and In Vivo Genetic Fitness of Pseudomonas aeruginosa Using High-Throughput Sequencing of Transposon Libraries', *PLoS Pathogens*. Edited by B.I. Kazmierczak, 9(9), p. e1003582. doi:10.1371/journal.ppat.1003582.

Trouillon, J. *et al.* (2020) 'Species-specific recruitment of transcription factors dictates toxin expression', *Nucleic Acids Res*, 48(5), pp. 2388–2400. doi:10.1093/nar/gkz1232.

Ventre, I. *et al.* (2006) 'Multiple sensors control reciprocal expression of Pseudomonas aeruginosa regulatory RNA and virulence genes', *Proceedings of the National Academy of Sciences*, 103(1), pp. 171–176. doi:10.1073/pnas.0507407103.

Vitkauskienė, A. *et al.* (2010) 'Pseudomonas aeruginosa bacteremia: resistance to antibiotics, risk factors, and patient mortality', *Medicina*, 46(7), p. 490. doi:10.3390/medicina46070071.

Weber, B.S. *et al.* (2020) 'Genetic and Chemical Screening in Human Blood Serum Reveals Unique Antibacterial Targets and Compounds against Klebsiella pneumoniae', *Cell Reports*, 32(3), p. 107927. doi:10.1016/j.celrep.2020.107927.

Winsor, G.L. *et al.* (2011) 'Pseudomonas Genome Database: improved comparative analysis and population genomics capability for Pseudomonas genomes', *Nucleic Acids Research*, 39(suppl_1), pp. D596–D600. doi:10.1093/nar/gkq869.

Winsor, G.L. *et al.* (2016) 'Enhanced annotations and features for comparing thousands of Pseudomonas genomes in the Pseudomonas genome database', *Nucleic Acids Research*, 44(D1), pp. D646–D653. doi:10.1093/nar/gkv1227.

Wood, L.F. and Ohman, D.E. (2012) 'Identification of genes in the sigma(2)(2) regulon
of Pseudomonas aeruginosa required for cell envelope homeostasis in either the
planktonic or the sessile mode of growth', *mBio*, 3(3). doi:10.1128/mBio.00094-12.

Woong Park, S. *et al.* (2011) 'Evaluating the Sensitivity of Mycobacterium tuberculosis
to Biotin Deprivation Using Regulated Gene Expression', *PLoS Pathogens*. Edited by
M.A. Behr, 7(9), p. e1002264. doi:10.1371/journal.ppat.1002264.

Wurtzel, O. *et al.* (2012) 'The Single-Nucleotide Resolution Transcriptome of
Pseudomonas aeruginosa Grown in Body Temperature', *PLOS Pathogens*, 8(9), p.
e1002945. doi:10.1371/journal.ppat.1002945.

Yin, Y. *et al.* (2013) 'Expression of mucoid induction factor MucE is dependent upon
the alternate sigma factor AlgU in Pseudomonas aeruginosa', *BMC Microbiology*,
13(1), p. 232. doi:10.1186/1471-2180-13-232.

Zhu, B. *et al.* (2021) 'Genome-wide identification of Streptococcus sanguinis fitness
genes in human serum and discovery of potential selective drug targets', *Molecular*
Microbiology, 115. doi:10.1111/mmi.14629.

1066

Supplementary material

Figure S1. Overexpression of srg operon leads to a population tolerant to plasma killing. Survival kinetics of IHMA87 wild-type strain (dame data as presented in Fig. 2B) and of Tn::P_{srg} in plasma over 6h incubation measured by CFU counting (n=5).

Table 1S. Tn-seq data

Table 2S. RNA-seq differential gene expression data

Table S3. Bacterial strains and plasmids

Bacteria	Features	Reference/origin
Pseudomonas aeruginosa		
IHMA879472/AZPAE15042	Urinary isolate, plasma sensitive	IHMA[1] collection (Kos *et al.*, 2015; Pont *et al.*, 2020; Trouillon *et al.*, 2020) (Kos, Trouillon, Pont)
IHMA87Δ*bioB*	IHMA87 with *bioB* deletion	This work
IHMA87Δ*purD*	IHMA87 with *purD* deletion	This work
IHMA87Δ*algD*	IHMA87 with *algD* deletion	This work
IHMA87Δ*01134*	IHMA87 with *01134* deletion	This work
IHMA87 Tn::P_{srg}	IHMA87 with mariner transposon inserted in the promoter of *srg* operon (Gm^R)	This work
IHMA87 Tn::P_{srg} Δ*srgABC*	IHMA87 Tn::P_{srg} with *srgABC* deletion (Gm^R)	This work

IHMA87 Tn::P_{srg} Δ$srgA$	IHMA87 Tn::P_{srg} with $srgA$ deletion (Gm^R)	This work
IHMA87 Tn::P_{srg} Δ$srgB$	IHMA87 Tn::P_{srg} with $srgB$ deletion (Gm^R)	This work
IHMA87 Tn::P_{srg} Δ$srgC$	IHMA87 Tn::P_{srg} with $srgC$ deletion (Gm^R)	This work
IHMA87 Tn::P_{srg} Δ$srgBC$	IHMA87 Tn::P_{srg} with $srgBC$ deletion (Gm^R)	This work
IHMA87 Tn::P_{srg} Δ$algD$	IHMA87 Tn::P_{srg} with $algD$ deletion (Gm^R)	This work
IHMA87 Tn::P_{srg} Δ01134	IHMA87 Tn::P_{srg} with 01134 deletion (Gm^R)	This work
IHMA87 $ladS$::Tn	IHMA87 with mariner transposon inserted in $ladS$ gene (Gm^R)	This work
IHMA87 $bioA$::Tn	IHMA87 with mariner transposon inserted in $bioA$ gene (Gm^R)	This work
IHMA87 $bioB$::Tn	IHMA87 with mariner transposon inserted in $bioB$ gene (Gm^R)	This work
IHMA87 $mucE$::Tn	IHMA87 with mariner transposon inserted in $mucE$ gene (Gm^R)	This work
IHMA87 $mucA$::Tn	IHMA87 with mariner transposon inserted in $mucA$ gene (Gm^R)	This work
IHMA87 $mucD$::Tn	IHMA87 with mariner transposon inserted in $mucD$ gene (Gm^R)	This work
IHMA87 $pprB$::Tn	IHMA87 with mariner transposon inserted in $pprB$ gene (Gm^R)	This work
IHMA87 $mucB$::Tn	IHMA87 with mariner transposon inserted in $mucB$ gene (Gm^R)	This work
Escherichia coli		
DH5α	Laboratory strain	Lab collection
TOP10	Cloning strain	Invitrogen
Plasmids		
pBTK24	Plasmid with Himar-1 mariner transposon and C9 transposase (Amp^R, Gm^R)	(Kulasekara *et al.*, 2005)
pRK600	Helper plasmid with conjugative properties (Cm^R)	(Kessler, de Lorenzo and Timmis, 1992)
pEXG2	Allelic exchange vector (Gm^R), $sacB$	(Rietsch *et al.*, 2005)
pEX18Tc	Allelic exchange vector (Tc^R), $sacB$	(Li *et al.*, 2011)
pEX100T	Allelic exchange vector (Cb^R), $sacB$	(Schweizer and Hoang, 1995)
pminiCTX-$pX2$-GFP	pminiCTX harboring the gfp gene under the control of a constitutive promoter ($attP$ site, Tc^R)	(Thibault *et al.*, 2009)
pminiCTX-$Psrg$-$lacZ$	pminiCTX-$lacZ$ harboring the srg promoter fused to $lacZ$ reporter gene ($attP$ site, Tc^R)	This work
pEXG2-mut-$bioB$	pEXG2 carrying DNA fragment for $bioB$ deletion obtained by SLIC (Gm^R)	This work
pEXG2-mut-$purD$	pEXG2 carrying DNA fragment for $purD$ deletion obtained by SLIC (Gm^R)	This work
pUC57-mut-$srgABC$	pUC57 carrying the synthetic DNA fragment for $srgABC$ deletion in Tn::P_{srg} (Genewiz)	This work
pEX18Tc-mut-$srgABC$	pEX18Tc carrying DNA fragment for srgABC deletion in Tn::P_{srg} mutant obtained by SLIC (Tc^R)	This work
pEX18Tc-mut-01134	pEX18Tc carrying DNA fragment for IHMA87_01134 deletion obtained by SLIC (Tc^R)	This work
pEX100T-mut-$srgA$	pEX100T carrying DNA fragment for $srgA$ deletion obtained by SLIC (Cb^R)	This work
pEX100T-mut-$srgB$	pEX100T carrying DNA fragment for $srgB$ deletion obtained by SLIC (Cb^R)	This work

pEX100T-mut-*srgC*	pEX100T carrying DNA fragment for *srgC* deletion obtained by SLIC (Cb^R)	This work
pEX100T-mut-*srgAB*	pEX100T carrying DNA fragment for *srgAB* deletion obtained by SLIC (Cb^R)	This work
pEX100T-mut-*srgBC*	pEX100T carrying DNA fragment for *srgBC* deletion obtained by SLIC (Cb^R)	This work
pEX100T-mut-*algD*	pEX100T carrying DNA fragment for *algD* deletion obtained by SLIC (Cb^R)	This work

[1] International Health Management Association, USA

Table S4. Oligonucleotides used for PCRs

Primers	Sequence (5'-3')	
pEXG2-mut-*bioB*-sF1	**GGTCGACTCTAGAGGATCCCC**GTTCGGACCCCTCGAACATCG	*bioB* deletion
pEXG2-mut-*bioB*-sR1	GGTGGCGACGGCTGCGGAT	*bioB* deletion
pEXG2-mut-*bioB*-sF2	**GCGCATCCGCAGCCGTCGCCACC**CAGCTGTTCTATAACGCCGCCT	*bioB* deletion
pEXG2-mut-*bioB*-sR2	**ACCGAATTCGAGCTCGAGCCC**CTTGCCGACCAGCGCGGTGA	*bioB* deletion
F0-mut-*bioB*	GCGAGCAGCCGTTCGGCG	PCR verification for *bioB* deletion
R0-mut-*bioB*	TTGAGGCGGTCCTCCAGCA	PCR verification for *bioB* deletion
pEXG2-mut-*purD*-sF1	**GGTCGACTCTAGAGGATCCCC**CAGGAGATCCACGACCTGATCT	*purD* deletion
pEXG2-mut-*purD*-sR1	ACCGCCGCTGCCGATGATGA	*purD* deletion
pEXG2-mut-*purD*-sF2	**TACTCATCATCGGCAGCGGCGGT**GAGCGCGGCGAGTCCTGAC	*purD* deletion
pEXG2-mut-*purD*-sR2	**ACCGAATTCGAGCTCGAGCCC**TCGCGGACCAGCTGGCCGT	*purD* deletion
F0-mut-*purD*	CTGAAGATCGTCACTCGCCG	PCR verification for *purD* deletion
R0-mut-*purD*	GACTCGCCGGTGTGCTGGT	PCR verification for *purD* deletion
pEX18Tc-mut-*srgABC*-sF1	**GTCGACTCTAGAGGATCCC**CGGATCCCCGACATTCGGCTA	Transfer DNA fragment from pUC57 to pEX18Tc for *srgABC* deletion in Tn::*P*~*srg*~
pEX18Tc-mut-*srgABC*-sR1	**CGAATTCGAGCTCGGTACCC**AAGCTTTGATCTACGTGCAAGC	
F0–mut-*srgABC*	GCCTCGCCGACCTCTACA	PCR verification for *srgA*, *srgB*, *srgC* and *srgABC* deletion
RO-Tn::*P*~*srg*~-mut-*srgABC*	GCTTGCTGCCTTCGACCAAG	PCR verification for *srgA*, *srgB*, *srgC* and *srgABC* deletion
pEX18Tc-mut-01134-sF1	**GTCGACTCTAGAGGATCCCC**CACCTCGGTGTCCACGCTGC	IHMA87_01134 deletion
pEX18Tc-mut-01134-sR1	CGACGAGCATCGACTTGTTCAC	IHMA87_01134 deletion
pEX18Tc-mut-01134-sF2	**TGAACAAGTCGATGCTCGTC**GACAAGAACGGTCGGCTGGT	IHMA87_01134 deletion
pEX18Tc-mut-01134-sR2	**CGAATTCGAGCTCGGTACCC**ATGTTCCGCAGCCTGGTCGG	IHMA87_01134 deletion
F0-mut-01134	CTGCTGTACGTCGGCTTCCG	PCR verification for IHMA87_01134 deletion
R0-mut-01134	CTGGATGGCACCACCAACTTC	PCR verification for IHMA87_01134 deletion
pEX100T-mut-*srgA*-sF1	**ACCCTGTTATCCCTACCC**GTGATCTACG	*srgA* and *srgAB* deletion in

	TGCAAGCAGA	Tn::P_{srg}
pEX100T-mut-*srgA*-sR1	CAGGTTGCGTGAGCTGCTCA	*srgA* and *srgAB* deletion in Tn::P_{srg}
pEX100T-mut-*srgA*-sF2	**TGAGCAGCTCACGCAACCTGTGA**CCTGGACATTCTGACGAGGTA	*srgA* deletion in Tn::P_{srg}
pEX100T-mut-*srgA*-sR2	**GGATAACAGGGTAATCCC**CACCAGCGCAAGCAGCAGC	*srgA* deletion in Tn::P_{srg}
pEX100T-mut-*srgB*-sF1	**ACCCTGTTATCCCTACCC**ATGGAGGTGAACATGAGCAGC	*srgB* and *srgBC* deletion in Tn::P_{srg}
pEX100T-mut-*srgB*-sR1	AGCCAGAATCCACATGTTTC	*srgB* and *srgBC* deletion in Tn::P_{srg}
pEX100T-mut-*srgB*-sF2	**GAAACATGTGGATTCTGGCTTGA**ACGGAAGCTCGGCGCCACT	*srgB* deletion in Tn::P_{srg}
pEX100T-mut-*srgB*-sR2	**GGATAACAGGGTAATCCC**CGTGTTGACTCACGTCGGAC	*srgB* deletion in Tn::P_{srg}
pEX100T-mut-*srgC*-sF1	**ACCCTGTTATCCCTACCC**CGACGAACAGGCCTGGACAT	*srgC* deletion in Tn::P_{srg}
pEX100T-mut-*srgC*-sR1	AAGAATGACGGCGGATCGTGA	*srgC* deletion in Tn::P_{srg}
pEX100T-mut-*srgC*-sF2	**TCACGATCCGCCGTCATTCTTTGA**GATCCCAGCCTGGACTGATC	*srgC* deletion in Tn::P_{srg}
pEX100T-mut-*srgC*-sR2	**GGATAACAGGGTAATCCCC**CACTCGAAGCCGACATTCG	*srgC* deletion in Tn::P_{srg}
pEX100T-mut-*srgAB*-sF2	**TGAGCAGCTCACGCAACCTGTGA**ACGGAAGCTCGGCGCCACT	*srgAB* deletion in Tn::P_{srg}
pEX100T-mut-*srgAB*-sR2	**GGATAACAGGGTAATCCC**CGTGTTGACTCACGTCGGAC	*srgAB* deletion in Tn::P_{srg}
pEX100T-mut-*srgBC*-sF2	**GAAACATGTGGATTCTGGCTTGA**GATCCCAGCCTGGACTGATC	*srgBC* deletion in Tn::P_{srg}
pEX100T-mut-*srgBC*-sR2	**GGATAACAGGGTAATCCCC**CACTCGAAGCCGACATTCG	*srgBC* deletion in Tn::P_{srg}
pEX100T-mut-*algD*-sF1	**ACCCTGTTATCCCTACCC**CGGAAACGCCATCAAGTTGGTA	*algD* deletion
pEX100T-mut-*algD*-sR1	GCCAGCACATACCGCACCGA	*algD* deletion
pEX100T-mut-*algD*-sF2	**TCGGTGCGGTATGTGCTGGC**CAGGCCGAGGGCATCTGCT	*algD* deletion
pEX100T-mut-*algD*-sR2	**GGATAACAGGGTAATCCC**GCTGTCGATGGCTTCGCGGA	*algD* deletion
F0-mut-*algD*	CTCGTGGCGAATAGGCCTAC	PCR verification for *algD* deletion
R0-mut-*algD*	GCGAAGCGCAGCTTGTGCC	PCR verification for *algD* deletion
Primers for Tn-seq		
Short adpator	TACCACGACCA-NH2	
Long adaptor	GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTGGTCGTGGTAT	
PCR1 Tn-specific	CACAGGAAACAGGACTCTAGAGG	
PCR2 adaptor complementary	GTGACTGGAGTTCAGACGTGTG	
P5+ Illumina	AATGATACGGCGACCACCGAGATCTACACTCTTTCCCTACACGACGCTCTTCCGATCTCTAGAGACCGGGGACTTATCAGC	
P7-index	CAAGCAGAAGACGGCATACGAGAT**CGTGAT**	

Supplementary references

Kessler, B., de Lorenzo, V. and Timmis, K.N. (1992) 'A general system to integrate *lacZ* fusions into the chromosomes of gram-negative eubacteria: regulation of thePm promoter of theTOL plasmid studied with all controlling elements in monocopy', *Molecular and General Genetics MGG*, 233(1–2), pp. 293–301. doi:10.1007/BF00587591.

Kos, V.N. *et al.* (2015) 'The resistome of Pseudomonas aeruginosa in relationship to phenotypic susceptibility', *Antimicrob Agents Chemother*, 59(1), pp. 427–36. doi:10.1128/AAC.03954-14.

Kulasekara, H.D. *et al.* (2005) 'A novel two-component system controls the expression of Pseudomonas aeruginosa fimbrial cup genes', *Molecular Microbiology*, 55(2), pp. 368–380. doi:10.1111/j.1365-2958.2004.04402.x.

Li, S. *et al.* (2011) 'Physiological role of the novel salicylaldehyde dehydrogenase NahV in mineralization of naphthalene by Pseudomonas putida ND6', *Microbiological Research*, 166(8), pp. 643–653. doi:10.1016/j.micres.2011.01.003.

Pont, S. *et al.* (2020) 'Bacterial behavior in human blood reveals complement evaders with some persister-like features', *PLoS Pathog*, 16(12), p. e1008893. doi:10.1371/journal.ppat.1008893.

Rietsch, A. *et al.* (2005) 'ExsE, a secreted regulator of type III secretion genes in Pseudomonas aeruginosa', *Proceedings of the National Academy of Sciences*, 102(22), pp. 8006–8011. doi:10.1073/pnas.0503005102.

Schweizer, H.P. and Hoang, T.T. (1995) 'An improved system for gene replacement and xylE fusion analysis in Pseudomonas aeruginosa', *Gene*, 158(1), pp. 15–22. doi:10.1016/0378-1119(95)00055-B.

Thibault, J. *et al.* (2009) 'Anti-activator ExsD forms a 1:1 complex with ExsA to inhibit transcription of type III secretion operons', *The Journal of Biological Chemistry*, 284(23), pp. 15762–15770. doi:10.1074/jbc.M109.003533.

Trouillon, J. *et al.* (2020) 'Species-specific recruitment of transcription factors dictates toxin expression', *Nucleic Acids Res*, 48(5), pp. 2388–2400. doi:10.1093/nar/gkz1232.