SARS-CoV-2 Omicron Spike recognition by plasma from individuals receiving BNT162b2 mRNA vaccination with a 16-weeks interval between doses

Debashree Chatterjee[1], Alexandra Tauzin[1,2], Lorie Marchitto[1,2], Shang Yu Gong[1,3], Marianne Boutin[1,2], Catherine Bourassa[1], Guillaume Beaudoin-Bussières[1,2], Yuxia Bo[4], Shilei Ding[1,2], Annemarie Laumaea[1,2], Dani Vézina[1,2], Josée Perreault[5], Laurie Gokool[1], Chantal Morrisseau[1], Pascale Arlotto[1], Éric Fournier[6], Aurélie Guilbault[6], Benjamin Delisle[6], Inès Levade[6], Guillaume Goyette[1], Gabrielle Gendron-Lepage[1], Halima Medjahed[1], Gaston De Serres[7], Cécile Tremblay[1,2], Valérie Martel-Laferrière[1,2], Daniel E. Kaufmann[1,8], Renée Bazin[5], Jérémie Prévost[1,2], Sandrine Moreira[6], Jonathan Richard[1,2,*], Marceline Côté[4,*], and Andrés Finzi[1,2,3,9,*]

[1]Centre de Recherche du CHUM, Montreal, QC, H2X 0A9 Canada
[2]Département de Microbiologie, Infectiologie et Immunologie, Université de Montréal, Montreal, QC, H2X 0A9, Canada
[3]Department of Microbiology and Immunology, McGill University, Montreal, QC, H3A 2B4, Canada
[4]Department of Biochemistry, Microbiology and Immunology, and Center for Infection, Immunity, and Inflammation, University of Ottawa, Ottawa, ON K1H 8M5, Canada
[5]Héma-Québec, Affaires Médicales et Innovation, Quebec, QC G1V 5C3, Canada
[6]Laboratoire de Santé Publique du Québec, Institut Nationale de Santé Publique du Québec, Sainte-Anne-de-Bellevue, QC H9X 3R5, Canada.
[7]Institut National de Santé Publique du Québec, Quebec, QC, H2P 1E2, Canada
[8]Département de Médecine, Université de Montréal, Montreal, QC, H3T 1J4, Canada
[9]Lead contact

*Correspondence:

jonathan.richard.1@umontreal.ca (J.R.)
marceline.cote@uottawa.ca (M.C.)
andres.finzi@umontreal.ca (A.F.)

Summary word count: **148**

Character count: **18816**

SUMMARY

Continuous emergence of SARS-CoV-2 variants of concern (VOC) is fueling the COVID-19 pandemic. Omicron (B.1.1.529), is rapidly spreading worldwide. The large number of mutations in its Spike raised concerns about a major antigenic drift that could significantly decrease vaccine efficacy and infection-induced immunity. A long interval between BNT162b2 mRNA doses was shown to elicit antibodies that efficiently recognize Spikes from different VOCs. Here we evaluated the recognition of Omicron Spike by plasma from a cohort of SARS-CoV-2 naïve and previously-infected individuals that received their BNT162b2 mRNA vaccine 16-weeks apart. Omicron Spike was recognized less efficiently than D614G, Alpha, Beta, Gamma and Delta Spikes. We compared to plasma activity from participants receiving a short (4-weeks) interval regimen. Plasma from individuals of the long interval cohort neutralized better the Omicron Spike compared to those that received a short interval. Whether this difference confers any clinical benefit against Omicron remains unknown.

KEYWORDS: Coronavirus, COVID-19, SARS-CoV-2, Spike glycoproteins, Delayed mRNA vaccine regimen, Variants of concern, Humoral responses, Neutralization, Omicron

INTRODUCTION

SARS-CoV-2 variants are constantly evolving under immune selective pressure. Ongoing mutational events in the viral genome leads to the emergence of variants with unique properties including increased transmission capabilities and resistance to antibodies elicited by both natural infection and vaccination. Based on transmission capabilities, virulence, and vaccine effectiveness, SARS-CoV-2 variants are classified as variants of concern (VOCs), variant of interest (VOIs) or variants under monitoring (VUMs)(WHO, 2021). In late 2020, the Alpha (B.1.1.7) variant emerged. The N501Y Spike mutation increased its affinity for the ACE2 receptor, leading to increased transmissibility (Davies et al., 2021; Prevost et al., 2021; Rambaut et al., 2020). The accumulation of E484K and K417N/T mutations along with N501Y in the receptor binding domain (RBD) led to the emergence of Beta (B.1.351) and Gamma (P.1) lineages, which rapidly spread worldwide (Amanat et al., 2021; ECDC, 2020; Tang et al., 2021). In April 2021, the Delta (B.1.617.2) variant emerged and quickly spread to most countries (Allen et al., 2022; Planas et al., 2021b), but is rapidly being replaced by Omicron (B.1.1.529). The World Health Organization designated Omicron as a VOC on November 26, 2021(WHO, 2021). Omicron accumulated more than 30 mutations in its Spike, raising concerns about a major antigenic drift that could significantly decrease vaccine efficacy.

Here we evaluated the recognition of the Omicron Spike by plasma from a cohort of SARS-CoV-2 naïve and previously-infected individuals that received the two BNT162b2 mRNA vaccine doses 16-weeks apart. We compared these responses to those elicited in individuals receiving a short dose interval regimen (4-weeks). Plasma from vaccinated previously-infected individuals recognized more efficiently all tested Spikes (D614G, Alpha, Beta, Gamma, Delta and Omicron) than those from naïve vaccinated individuals. Omicron Spike was recognized less efficiently than D614G, Alpha, Beta, Gamma and Delta Spikes. However, plasma from individuals receiving a long interval neutralized better the Omicron Spike compared to those that received a short interval.

RESULTS

Recognition of Spike variants by plasma from vaccinated individuals

The antigenic profile of D614G, Alpha, Beta, Gamma, Delta and Omicron Spikes was assessed

with plasma collected three weeks (V3) and 4 months (V4) after the second dose of the BNT162b2

mRNA vaccine administered with a 16-weeks interval between doses (Figure 1A) (Tauzin et al.,

2021a). Briefly, 293T cells were transfected with plasmids coding for full-length Spike variants.

Two days post-transfection, cells were incubated with the indicated plasmas followed by flow

cytometry analysis, as described (Anand et al., 2021; Beaudoin-Bussieres et al., 2020; Gasser et

al., 2021; Prevost et al., 2020; Tauzin et al., 2021a; Tauzin et al., 2021b). VOCs Spike expression

levels were normalized to the signal obtained with the conformationally independent anti-S2

neutralizing CV3-25 antibody (Li et al., 2021; Prevost et al., 2021; Ullah et al., 2021) that efficiently

recognized and neutralized Omicron (Supplemental Figure 1). Using plasma from previously-

infected individuals or from naïve double vaccinated individuals we observed a significant

increase of recognition of all tested Spikes upon vaccination (Figure 1B-G), in agreement with

previous observations (Stamatatos et al., 2021; Tauzin et al., 2021a; Tauzin et al., 2021b). In all

cases, the Omicron Spike was significantly less recognized than all other Spikes, with the

exception of the Beta variant (Figure 1D, H-J). At V3 and V4, levels of plasma binding against

Omicron Spike in previously-infected individuals were similar to those against Delta Spike in naïve

individuals (Figure 1 I and J). In agreement with previous observations (Tauzin et al., 2021a),

Spike recognition declined more rapidly in the naïve group compared to previously-infected

individuals.

Impact of the interval between mRNA vaccine doses on Omicron Spike recognition and neutralization

Recent reports suggested that vaccine regimens with a delayed boost elicits stronger humoral responses than the approved, short interval, regimen (Grunau et al., 2021; Tauzin et al., 2021a). The long regimen interval has been associated with good vaccine efficacy against different VOCs (Skowronski et al., 2021). We therefore compared the capacity of plasma from naïve vaccinated individuals that received the second dose with an interval of sixteen weeks (median [range]: 111 days [109–114 days]) to those obtained from 12 SARS-CoV-2 naïve donors who received their two doses four weeks apart (median [range]: 30 days [22–34 days]) (Table 1 and Figure 2A). As shown in Figure 2B, plasma from naïve individuals that received a 16-weeks interval between the two doses, recognized slightly better, albeit this difference was not significant, the Omicron Spike than plasma from individuals who received a short interval between doses (4 weeks). The 16-weeks interval regimen elicited significantly better neutralization activity against pseudoviral particles bearing the Omicron Spike (Figure 2C). This suggests that the delayed boosting in naïve individuals facilitates antibody maturation resulting in enhanced breadth able to provide detectable levels of neutralization against Omicron.

DISCUSSION

In the province of Québec, Canada, like in other jurisdiction worldwide, the prevalence of Omicron increased dramatically from the first case detected on November 23 to being the dominant variant less than one month later. Compared to the reference Wuhan-Hu-1 strain, the Omicron variant carries over 50 non-synonymous mutations within its genome, more than 30 of which are located in the gene coding for the Spike glycoprotein. Several of these mutations affecting the RBD, the N-terminal domain (NTD) and the furin cleavage domain were observed in other VOCs (Viana R., 2021), which is consistent with positive selection of favorable mutations. Previous *in vitro* studies already showed the association of some of these mutations with increased infectivity, ACE2 interaction (N501Y, P681H) (Gong et al., 2021; Saito et al., 2021) or immune evasion (K417N, N440K, G446S, S477N, E484A/K, Q493R) (Baum et al., 2020; Clark et al., 2021; Greaney et al., 2021a; Greaney et al., 2021b; Greaney et al., 2021c; Liu et al., 2020; Rappazzo et al., 2021; Starr et al., 2021; Weisblum et al., 2020) . This unprecedented accumulation of Spike mutations raised concern about a major antigenic drift that could significantly decrease the efficacy of current vaccines (Andrews et al., 2021; Khoury et al., 2021; Schmidt et al., 2021b).

To get a better understanding of the antigenic profile, we compared the antigenicity of the Omicron Spike to those from D614G, Alpha, Beta, Gamma and Delta VOCs. We used plasma from naïve and previously-infected individuals who received their two doses of the BNT162b2 mRNA vaccine 16 weeks apart. In agreement with previous observations, we found that previously-infected vaccinated individuals recognized more efficiently all Spikes than naïve individuals at the two timepoints analyzed (3 weeks and 4 months post second dose) (Stamatatos et al., 2021; Tauzin et al., 2021a; Tauzin et al., 2021b). Interestingly, we observed that recognition of all Spikes, including Omicron, decreased more rapidly in naïve than previously-infected individuals, as reported (Tauzin et al., 2021a). The three antigenic exposures (infection + 2 doses) of previously-

infected individuals compared to the two exposures in double-vaccinated naïve individuals

possibly explain their more sustained humoral response, suggesting that an additional exposition

to the Spike antigen in the form of a third vaccine dose could elicit similar responses. Alternatively,

if infection elicits a qualitatively broader humoral response linked to epitopes located outside the

Spike glycoprotein, the effect of a third dose in naïve individuals may remain qualitatively different.

Independently of their infection history, all plasma recognized significantly less efficiently the

Omicron Spike compared to Spikes from other VOCs (Figure 1).

Since recent studies have shown that recognition of full-length Spikes at the surface of transfected

293T cells strongly correlates with recognition of primary airway epithelial cells infected with

authentic viruses as well as with antibody-dependent cellular cytotoxicity (Ding et al., 2022), these

results suggest that Fc-mediated effector functions against Omicron could also be affected.   Of

note, low Spike recognition translated into increased Omicron neutralization resistance (Figure

2C). In agreement with previous observations (Stamatatos et al., 2021; Tauzin et al., 2021a;

Tauzin et al., 2021b), plasma from vaccinated previously-infected individuals recognized more

efficiently Omicron and all other VOCs than vaccinated naïve individuals (Figure 1). As naïve

double-vaccinated individuals have been well protected against the delta variant, the observation

of similar levels of plasma binding against Delta Spike in naïve individuals and those against

Omicron Spike in previously-infected individuals may be important. This suggests that the benefits

of hybrid immunity also apply to Omicron but this hypothesis will need confirmation through

vaccine effectiveness studies.

Several reports (Cele et al., 2021; Garcia-Beltran et al., 2021; Planas et al., 2021a; Schmidt et al.,

2021a; Zhang et al., 2021) have shown neutralization resistance using plasma from naïve donors

who received the approved regimen of the BNT162b2 mRNA vaccine (3-4 weeks vaccine interval).

Strikingly, we observed that plasma from naïve vaccinated donors who received their two doses

according to the approved short three-four-week interval, recognized and neutralized less

efficiently Omicron compared to the long (16-weeks interval). For all individuals, the level of

Omicron Spike recognition remains dramatically lower than for the ancestral Spike, the antigen

used in the current vaccines and these levels decrease over time.   Therefore, it will be important

to determine in epidemiological studies if the vaccine interval advantage, as measured by these

in vitro parameters, confers any clinical benefit against Omicron.

ACKNOWLEDGMENTS

The authors are grateful to the donors who participated in this study. The authors thank the CRCHUM BSL3 and Flow Cytometry Platforms for technical assistance. We thank Dr. Stefan Pöhlmann (Georg-August University, Germany) for the plasmid coding for SARS-CoV-2 Spike Wuhan-Hu-1 strain. We also thank Amélie Boivin and Yves Grégoire at Héma-Québec for helping to access the samples from the PLASCOV Biobank and all the plasma donors who participate in this biobank. This work was supported by le Ministère de l'Économie et de l'Innovation du Québec, Programme de soutien aux organismes de recherche et d'innovation to A.F. and by the Fondation du CHUM. This work was also supported by a Canadian Institutes of Health Research (CIHR) foundation grant #352417, by a CIHR operating Pandemic and Health Emergencies Research grant #177958, a CIHR stream 1 and 2 for SARS-CoV-2 Variant Research to A.F., and by an Exceptional Fund COVID-19 from the Canada Foundation for Innovation (CFI) #41027 to A.F. and D.E.K. Work on variants presented was also supported by the Sentinelle COVID Quebec network led by the LSPQ in collaboration with Fonds de Recherche du Québec Santé (FRQS) to A.F. This work was also partially supported by a CIHR COVID-19 rapid response grant (OV3 170632) and CIHR stream 1 SARS-CoV-2 Variant Research to MC. A.F. is the recipient of Canada Research Chair on Retroviral Entry no. RCHS0235 950-232424. MC is a Tier II Canada Research Chair in Molecular Virology and Antiviral Therapeutics. V.M.L. is supported by a FRQS Junior 1 salary award. C,T. is the recipient of the Pfizer/Université de Montréal Chair on HIV translational research. D.E.K. is a FRQS Merit Research Scholar. G.B.B. is the recipient of a FRQS PhD fellowship and J.P. is the recipient of a CIHR PhD fellowship. A.L. was supported by MITACS Accélération postdoctoral fellowships. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. We declare no competing interests.

AUTHOR CONTRIBUTIONS

D.C., J.R., S.M., M.C., and A.F. conceived the study. D.C., A.T., J.R., M.C. and A.F. designed experimental approaches. D.C., A.T., L.M., S.Y.G., M.B., C.B., G.B.-B., Y.B., S.D., A.L., D.V., E.F., A.G., B.D., I.L., J.Prévost, J.R. and A.F. performed, analyzed, and interpreted the experiments. J.Perreault, L.G., C.M., P.A., C.T., V.M.-L., D.E.K. and R.B. collected and provided clinical samples. G.D.S., C.T., V.M.-L., and D.E.K. provided scientific input related to VOCs and vaccine efficacy. D.C., J.R., and A.F. wrote the manuscript with inputs from others. Every author has read, edited, and approved the final manuscript.

DECLARATION OF INTERESTS

A.F. has filed a provisional patent application on the anti-Spike monoclonal antibody CV3-25.

FIGURE LEGENDS

Figure 1. Binding of vaccine-elicited antibodies to SARS-CoV-2 Spike variants

(**A**) SARS-CoV-2 vaccine cohort design. (**B-G**) 293T cells were transfected with the indicated full-

length Spike from different SARS-CoV-2 variants (D614G, Alpha, Beta, Gamma, Delta and

Omicron) and stained with the CV3-25 Ab or with plasma collected 3 weeks (V3) or 4 months (V4)

after a second dose administered with a 16 week interval. Samples were analyzed by flow

cytometry. The values represent the median fluorescence intensities (MFI) normalized by CV3-

25 Ab binding and presented as percentages of CV3-25 binding (**B-G, Left panels**). Each curve

represents the normalized MFIs obtained with the plasma of one donor at every time point. The

mean of each group is represented by a bold line. (**Right panels**) Plasma samples were grouped

in different time points (V0, V3 and V4). (**H-J**) Comparison of Spike recognition by plasma from

naïve and previously-infected donors, represented by red and black points respectively. Error

bars indicate means ± SEM. For naïve donors, n=10 at V3 and V4. For previously infected donors

vaccinated with two doses, n=10 at V0, V3 and V4. Statistical significance was tested using (**B-G**

Left panels) a Wilcoxon test, (**B-G Right panels**) Mann-Whitney test or (**H-J**) a Friedman test. (*

$P < 0.05$; ** $P < 0.01$; *** $P < 0.001$; **** $P < 0.0001$; ns, non-significant)

Figure 2. Omicron Spike recognition and neutralization with plasma from naïve individuals

that received a short versus a long mRNA vaccine dose interval.

(**A**) SARS-CoV-2 vaccine cohort design. (**B**) 293T cells were transfected with the full-length

Omicron Spike and stained with the CV3-25 Ab or with plasma from naïve donors who received

a short (4 weeks, yellow) or long (16 weeks, red) interval between doses collected three weeks

after the second dose (V3) and analyzed by flow cytometry. The values represent the MFI

normalized by CV3-25 Ab binding and presented as percentages of CV3-25 binding. (**C**)

Neutralizing activity was measured by incubating pseudoviruses bearing SARS-CoV-2

glycoproteins of Omicron, with serial dilutions of plasma for 1 h at 37°C before infecting 293T-

ACE2 cells. Neutralization half maximal inhibitory serum dilution (ID_{50}) values were determined

using a normalized non-linear regression using GraphPad Prism software. Undetectable

measures are represented as white symbols, and limits of detection are plotted. Error bars

indicate means ± SEM. (**** P < 0.0001). For naïve donors vaccinated with the short interval,

n=12. For naïve donors vaccinated with the long interval, n=10. Statistical significance was tested

using a Mann-Whitney test (**** P < 0.0001).

Table 1. Characteristics of the vaccinated SARS-CoV-2 cohorts

		SARS-CoV-2 Naïve		SARS-CoV-2 Previously infected
		Two doses Short interval (n=12)	Two doses Long interval (n=10)	Two doses Long interval (n=10)
Age		45 (24-60)	53 (40-59)	46 (39-65)
Sex	Male (n)	8	3	6
	Female (n)	4	7	4
Days between symptom onset and V0 [a]		N/A	N/A	169 (126-234)
Days between symptom onset and the 1st dose [a]		N/A	N/A	266 (185-321)
Days between the 1st and 2nd dose [a]		30 (22-34)	111 (109-114)	112 (90-134)
Days between the 2nd dose and V3 [a]		24 (12-37)	21 (17-27)	20 (13-29)
Days between the 2nd dose and V4 [a]		N/A	113 (105-118)	112 (90-127)

[a] Values displayed are medians, with ranges in parentheses.

STAR METHODS

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact, Andrés Finzi (andres.finzi@umontreal.ca).

Materials availability

All unique reagents generated during this study are available from the Lead contact without restriction.

Data and code availability

- All data reported in this paper will be shared by the lead contact (andres.finzi@umontreal.ca) upon request.

- This paper does not report original code.

- Any additional information required to reanalyze the data reported in this paper is available from the lead contact (andres.finzi@umontreal.ca) upon request.

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Ethics Statement

All work was conducted in accordance with the Declaration of Helsinki in terms of informed consent and approval by an appropriate institutional board. Blood samples were obtained from donors who consented to participate in this research project at CHUM (19.381) and from plasma donors who consented to participate in the Plasma Donor Biobank at Hema-Quebec (PLASCOV; REB-B-6-002-2021-003). Plasma was isolated by centrifugation, and samples stored at -80°C and in liquid nitrogen, respectively, until use.

Human subjects

The study was conducted in 10 SARS-CoV-2 naïve individuals (3 males and 7 females ; age range : 40-59 years) vaccinated with a long interval, 12 SARS-CoV-2 naïve individuals (8 males and 4 females ; age range : 24-60 years) vaccinated with a short interval and 10 SARS-CoV-2 previously-infected individuals (6 males and 4 females ; age range : 39-65 years) vaccinated with a long interval. All this information is summarized in Table 1. No specific criteria such as number of patients (sample size), gender, clinical or demographic were used for inclusion, beyond PCR confirmed SARS-CoV-2 infection in adults.

Plasma and antibodies

Plasma from SARS-CoV-2 naïve and previously-infected donors were collected, heat-inactivated for 1 hour at 56°C and stored at -80°C until ready to use in subsequent experiments. The conformationally independent S2-specific monoclonal antibody CV3-25 (Gong et al., 2021; Jennewein et al., 2021; Li et al., 2021; Prevost et al., 2021; Ullah et al., 2021) was used as a positive control and to normalize Spike expression in our flow cytometry assays, as described (Gong et al., 2021; Tauzin et al., 2021a; Tauzin et al., 2021b). Alexa Fluor-647-conjugated goat

anti-human Abs (Invitrogen) were used as secondary antibodies to detect plasma binding in flow

cytometry experiments.

Cell lines

293T human embryonic kidney cells (obtained from ATCC) were maintained at 37°C under 5%

CO_2 in Dulbecco's modified Eagle's medium (DMEM) (Wisent) containing 5% fetal bovine serum

(FBS) (VWR) and 100 µg/ml of penicillin-streptomycin (Wisent). The 293T-ACE2 cell line was

previously reported (Prévost et al., 2020).

METHOD DETAILS

Plasmids

The plasmids encoding the SARS-CoV-2 Spike variants; D614G, B.1.1.7, B.1.351, P.1 and

B.1.617.2 were previously described (Beaudoin-Bussieres et al., 2020; Gong et al., 2021; Li et al.,

2021; Tauzin et al., 2021a; Tauzin et al., 2021b). The plasmids encoding the B.1.1.529 Spike

was generated by overlapping PCR using a codon-optimized wild-type SARS-CoV-2 Spike gene

(GeneArt, ThermoFisher) that was synthesized (Biobasic) and cloned in pCAGGS as a template.

The B.1.1.529 Spike coding sequence was derived from the sequence ID EPI_ISL_6640919 that

initially contained the Q493K substitution as previously reported (Cameroni et al., 2021; Schmidt

et al., 2021a; Shah and Woo, 2021).

Cell surface staining and flow cytometry analysis

293T were transfected with full-length SARS-CoV-2 Spikes and a green fluorescent protein (GFP)

expressor (pIRES2-eGFP; Clontech) using the calcium-phosphate method. Two days post-

transfection, Spike-expressing 293T cells were stained with the CV3-25 Ab (5 µg/mL) as control

or plasma from vaccinated individuals (1:250 dilution) for 45 min at 37ºC. AlexaFluor-647-

conjugated goat anti-human IgG (1/1000 dilution) were used as secondary Abs. The percentage

of Spike-expressing cells (GFP+ cells) was determined by gating the living cell population based

on viability dye staining (Aqua Vivid, Invitrogen). Samples were acquired on a LSR II cytometer

(BD Biosciences), and data analysis was performed using FlowJo v10.7.1 (Tree Star). The

conformationally-independent anti-S2 antibody CV3-25 was used to normalize Spike expression,

as reported (Gong et al., 2021; Li et al., 2021; Prevost et al., 2021; Ullah et al., 2021). CV3-25

was shown to be effective against all Spike variants (Gong et al., 2021; Li et al., 2021; Prevost et

al., 2021; Ullah et al., 2021) and (Supplemental figure 1). The Median Fluorescence intensities

(MFI) obtained with plasma were normalized to the MFI obtained with CV3-25 (Gong et al., 2021;

Li et al., 2021; Prevost et al., 2021; Ullah et al., 2021) and presented as percentage of CV3-25

binding.

Virus neutralization assay

To produce SARS-CoV-2 pseudoviruses, 293T cells were transfected with the lentiviral vector

pNL4.3 R-E- Luc (NIH AIDS Reagent Program) and a plasmid encoding for the indicated S

glycoprotein (D614G, Alpha, Beta, Gamma, Delta or Omicron) at a ratio of 10:1. Two days post-

transfection, cell supernatants were harvested and stored at -80°C until use. For the neutralization

assay, 293T-ACE2 target cells were seeded at a density of 1×10^4 cells/well in 96-well

luminometer-compatible tissue culture plates (Perkin Elmer) 24h before infection. Pseudoviral

particles were incubated with several plasma dilutions (1/50; 1/250; 1/1250; 1/6250; 1/31250) for

1h at 37°C and were then added to the target cells followed by incubation for 48h at 37°C. For

CV3-25 neutralization, pseudoviral particles were incubated with increasing concentrations of

CV3-25 (0.01, 0.0316, 0.1, 0.316, 1 and 3.16 µg/mL) for 1h at 37°C and were then added to the

target cells followed by incubation for 48h at 37°C. Cells were lysed by the addition of 30 µL of

passive lysis buffer (Promega) followed by one freeze-thaw cycle. . An LB942 TriStar luminometer

(Berthold Technologies) was used to measure the luciferase activity of each well after the addition

of 100 µL of luciferin buffer (15mM $MgSO_4$, 15mM KH_2PO_4 [pH 7.8], 1mM ATP, and 1mM

dithiothreitol) and 50 µL of 1mM d-luciferin potassium salt (Prolume). The neutralization half-

maximal inhibitory dilution (ID_{50}) represents the plasma dilution to inhibit 50% of the infection of

293T-ACE2 cells by pseudoviruses.

QUANTIFICATION AND STATISTICAL ANALYSIS

Statistical analysis

Symbols represent biologically independent samples from SARS-CoV-2 naïve or PI individuals.

Lines connect data from the same donor. Statistics were analyzed using GraphPad Prism version

8.0.1 (GraphPad, San Diego, CA). Every dataset was tested for statistical normality and this

information was used to apply the appropriate (parametric or nonparametric) statistical test. P

values <0.05 were considered significant; significance values are indicated as * P<0.05, **

P<0.01, *** P<0.001, **** P<0.0001.

References

Allen, H., Vusirikala, A., Flannagan, J., Twohig, K.A., Zaidi, A., Chudasama, D., Lamagni, T.,
Groves, N., Turner, C., Rawlinson, C.*, et al.* (2022). Household transmission of COVID-19 cases
associated with SARS-CoV-2 delta variant (B.1.617.2): national case-control study. Lancet Reg
Health Eur *12*, 100252.

Amanat, F., Thapa, M., Lei, T., Ahmed, S.M.S., Adelsberg, D.C., Carreño, J.M., Strohmeier, S.,
Schmitz, A.J., Zafar, S., Zhou, J.Q.*, et al.* (2021). SARS-CoV-2 mRNA vaccination induces
functionally diverse antibodies to NTD, RBD, and S2. Cell.

Anand, S.P., Prevost, J., Nayrac, M., Beaudoin-Bussieres, G., Benlarbi, M., Gasser, R., Brassard,
N., Laumaea, A., Gong, S.Y., Bourassa, C.*, et al.* (2021). Longitudinal analysis of humoral
immunity against SARS-CoV-2 Spike in convalescent individuals up to 8 months post-symptom
onset. Cell Rep Med *2*, 100290.

Andrews, N., Stowe, J., Kirsebom, F., Toffa, S., Rickeard, T., Gallagher, E., Gower, C., Kall, M.,
Groves, N., O'Connell, A.-M.*, et al.* (2021). Effectiveness of COVID-19 vaccines against the
Omicron (B.1.1.529) variant of concern. 2021.2012.2014.21267615.

Baum, A., Fulton, B.O., Wloga, E., Copin, R., Pascal, K.E., Russo, V., Giordano, S., Lanza, K.,
Negron, N., Ni, M.*, et al.* (2020). Antibody cocktail to SARS-CoV-2 spike protein prevents rapid
mutational escape seen with individual antibodies. Science *369*, 1014-1018.

Beaudoin-Bussieres, G., Laumaea, A., Anand, S.P., Prevost, J., Gasser, R., Goyette, G.,
Medjahed, H., Perreault, J., Tremblay, T., Lewin, A.*, et al.* (2020). Decline of Humoral Responses
against SARS-CoV-2 Spike in Convalescent Individuals. mBio *11*.

Cameroni, E., Saliba, C., Bowen, J.E., Rosen, L.E., Culap, K., Pinto, D., De Marco, A., Zepeda,
S.K., di Iulio, J., Zatta, F.*, et al.* (2021). Broadly neutralizing antibodies overcome SARS-CoV-2
Omicron antigenic shift. 2021.2012.2012.472269.

Cele, S., Jackson, L., Khan, K., Khoury, D.S., Moyo-Gwete, T., Tegally, H., Scheepers, C.,
Amoako, D., Karim, F., Bernstein, M.*, et al.* (2021). SARS-CoV-2 Omicron has extensive but
incomplete escape of Pfizer BNT162b2 elicited neutralization and requires ACE2 for infection.
medRxiv.

Clark, S.A., Clark, L.E., Pan, J., Coscia, A., McKay, L.G.A., Shankar, S., Johnson, R.I., Brusic, V.,
Choudhary, M.C., Regan, J.*, et al.* (2021). SARS-CoV-2 evolution in an immunocompromised
host reveals shared neutralization escape mechanisms. Cell *184*, 2605-2617 e2618.

Davies, N.G., Abbott, S., Barnard, R.C., Jarvis, C.I., Kucharski, A.J., Munday, J.D., Pearson,
C.A.B., Russell, T.W., Tully, D.C., Washburne, A.D.*, et al.* (2021). Estimated transmissibility and
impact of SARS-CoV-2 lineage B.1.1.7 in England. Science *372*.

Ding, S., Adam, D., Beaudoin-Bussières, G., Tauzin, A., Gong, S.Y., Gasser, R., Laumaea, A.,
Anand, S.P., Privé, A., Bourassa, C.*, et al.* (2022). SARS-CoV-2 Spike Expression at the Surface
of Infected Primary Human Airway Epithelial Cells. Viruses *14*, 5.

ECDC (2020). Rapid increase of a SARS-CoV-2 variant with multiple spike protein mutations
observed in the United Kingdom. In Threat Assessment Brief.

Garcia-Beltran, W.F., St. Denis, K.J., Hoelzemer, A., Lam, E.C., Nitido, A.D., Sheehan, M.L.,
Berrios, C., Ofoman, O., Chang, C.C., Hauser, B.M.*, et al.* (2021). mRNA-based COVID-19
vaccine boosters induce neutralizing immunity against SARS-CoV-2 Omicron variant.
2021.2012.2014.21267755.

Gasser, R., Cloutier, M., Prevost, J., Fink, C., Ducas, E., Ding, S., Dussault, N., Landry, P.,
Tremblay, T., Laforce-Lavoie, A.*, et al.* (2021). Major role of IgM in the neutralizing activity of
convalescent plasma against SARS-CoV-2. Cell reports, 108790.

Gong, S.Y., Chatterjee, D., Richard, J., Prévost, J., Tauzin, A., Gasser, R., Bo, Y., Vézina, D.,
Goyette, G., Gendron-Lepage, G.*, et al.* (2021). Contribution of single mutations to selected
SARS-CoV-2 emerging variants Spike antigenicity. 2021.2008.2004.455140.

Greaney, A.J., Loes, A.N., Crawford, K.H.D., Starr, T.N., Malone, K.D., Chu, H.Y., and Bloom,
J.D. (2021a). Comprehensive mapping of mutations in the SARS-CoV-2 receptor-binding domain
that affect recognition by polyclonal human plasma antibodies. Cell Host Microbe *29*, 463-476
e466.

Greaney, A.J., Starr, T.N., Barnes, C.O., Weisblum, Y., Schmidt, F., Caskey, M., Gaebler, C.,
Cho, A., Agudelo, M., Finkin, S.*, et al.* (2021b). Mapping mutations to the SARS-CoV-2 RBD that
escape binding by different classes of antibodies. Nat Commun *12*, 4196.

Greaney, A.J., Starr, T.N., Gilchuk, P., Zost, S.J., Binshtein, E., Loes, A.N., Hilton, S.K.,
Huddleston, J., Eguia, R., Crawford, K.H.D.*, et al.* (2021c). Complete Mapping of Mutations to the
SARS-CoV-2 Spike Receptor-Binding Domain that Escape Antibody Recognition. Cell Host
Microbe *29*, 44-57 e49.

Grunau, B., Goldfarb, D.M., Asamoah-Boaheng, M., Golding, L., Kirkham, T.L., Demers, P.A.,
and Lavoie, P.M. (2021). Immunogenicity of Extended mRNA SARS-CoV-2 Vaccine Dosing
Intervals. JAMA : the journal of the American Medical Association.

Jennewein, M.F., MacCamy, A.J., Akins, N.R., Feng, J., Homad, L.J., Hurlburt, N.K., Seydoux,
E., Wan, Y.H., Stuart, A.B., Edara, V.V.*, et al.* (2021). Isolation and characterization of cross-
neutralizing coronavirus antibodies from COVID-19+ subjects. Cell reports *36*, 109353.

Khoury, D.S., Steain, M., Triccas, J.A., Sigal, A., Davenport, M.P., and Cromer, D. (2021). A meta-
analysis of Early Results to predict Vaccine efficacy against Omicron. 2021.2012.2013.21267748.

Li, W., Chen, Y., Prevost, J., Ullah, I., Lu, M., Gong, S.Y., Tauzin, A., Gasser, R., Vezina, D.,
Anand, S.P.*, et al.* (2021). Structural Basis and Mode of Action for Two Broadly Neutralizing
Antibodies Against SARS-CoV-2 Emerging Variants of Concern. Cell reports *S2211-*
1247(21)01714-9.

Liu, L., Wang, P., Nair, M.S., Yu, J., Rapp, M., Wang, Q., Luo, Y., Chan, J.F., Sahi, V., Figueroa,
A.*, et al.* (2020). Potent neutralizing antibodies against multiple epitopes on SARS-CoV-2 spike.
Nature *584*, 450-456.

Planas, D., Saunders, N., Maes, P., Guivel-Benhassine, F., Planchais, C., Buchrieser, J., Bolland,
W.-H., Porrot, F., Staropoli, I., Lemoine, F.*, et al.* (2021a). Considerable escape of SARS-CoV-2
variant Omicron to antibody neutralization. 2021.2012.2014.472630.

Planas, D., Veyer, D., Baidaliuk, A., Staropoli, I., Guivel-Benhassine, F., Rajah, M.M., Planchais,
C., Porrot, F., Robillard, N., Puech, J.*, et al.* (2021b). Reduced sensitivity of SARS-CoV-2 variant
Delta to antibody neutralization. Nature.

Prevost, J., Gasser, R., Beaudoin-Bussieres, G., Richard, J., Duerr, R., Laumaea, A., Anand, S.P.,
Goyette, G., Benlarbi, M., Ding, S.*, et al.* (2020). Cross-Sectional Evaluation of Humoral
Responses against SARS-CoV-2 Spike. Cell Rep Med *1*, 100126.

Prevost, J., Richard, J., Gasser, R., Ding, S., Fage, C., Anand, S.P., Adam, D., Gupta Vergara,
N., Tauzin, A., Benlarbi, M.*, et al.* (2021). Impact of temperature on the affinity of SARS-CoV-2
Spike glycoprotein for host ACE2. J Biol Chem *297*, 101151.

Rambaut, A., Holmes, E.C., O'Toole, A., Hill, V., McCrone, J.T., Ruis, C., du Plessis, L., and
Pybus, O.G. (2020). A dynamic nomenclature proposal for SARS-CoV-2 lineages to assist
genomic epidemiology. Nat Microbiol *5*, 1403-1407.

Rappazzo, C.G., Tse, L.V., Kaku, C.I., Wrapp, D., Sakharkar, M., Huang, D., Deveau, L.M.,
Yockachonis, T.J., Herbert, A.S., Battles, M.B.*, et al.* (2021). Broad and potent activity against
SARS-like viruses by an engineered human monoclonal antibody. Science *371*, 823-829.

Saito, A., Irie, T., Suzuki, R., Maemura, T., Nasser, H., Uriu, K., Kosugi, Y., Shirakawa, K.,
Sadamasu, K., Kimura, I.*, et al.* (2021). SARS-CoV-2 spike P681R mutation, a hallmark of the
Delta variant, enhances viral fusogenicity and pathogenicity. 2021.2006.2017.448820.

Schmidt, F., Muecksch, F., Weisblum, Y., Silva, J.D., Bednarski, E., Cho, A., Wang, Z., Gaebler,
C., Caskey, M., Nussenzweig, M.C.*, et al.* (2021a). Plasma neutralization properties of the SARS-
CoV-2 Omicron variant. 2021.2012.2012.21267646.

Schmidt, F., Weisblum, Y., Rutkowska, M., Poston, D., DaSilva, J., Zhang, F., Bednarski, E., Cho,
A., Schaefer-Babajew, D.J., Gaebler, C.*, et al.* (2021b). High genetic barrier to SARS-CoV-2
polyclonal neutralizing antibody escape. Nature *600*, 512-516.

Shah, M., and Woo, H.G. (2021). Omicron: A heavily mutated SARS-CoV-2 variant exhibits
stronger binding to ACE2 and potently escape approved COVID-19 therapeutic antibodies.
2021.2012.2004.471200.

Skowronski, D.M., Setayeshgar, S., Febriani, Y., Ouakki, M., Zou, M., Talbot, D., Prystajecky, N.,
Tyson, J.R., Gilca, R., Brousseau, N.*, et al.* (2021). Two-dose SARS-CoV-2 vaccine effectiveness
with mixed schedules and extended dosing intervals: test-negative design studies from British
Columbia and Quebec, Canada. 2021.2010.2026.21265397.

Stamatatos, L., Czartoski, J., Wan, Y.H., Homad, L.J., Rubin, V., Glantz, H., Neradilek, M.,
Seydoux, E., Jennewein, M.F., MacCamy, A.J.*, et al.* (2021). mRNA vaccination boosts cross-
variant neutralizing antibodies elicited by SARS-CoV-2 infection. Science.

Starr, T.N., Greaney, A.J., Addetia, A., Hannon, W.W., Choudhary, M.C., Dingens, A.S., Li, J.Z.,
and Bloom, J.D. (2021). Prospective mapping of viral mutations that escape antibodies used to
treat COVID-19. Science *371*, 850-854.

Tang, J.W., Toovey, O.T.R., Harvey, K.N., and Hui, D.D.S. (2021). Introduction of the South
African SARS-CoV-2 variant 501Y.V2 into the UK. J Infect *82*, e8-e10.

Tauzin, A., Gong, S.Y., Beaudoin-Bussières, G., Vézina, D., Gasser, R., Nault, L., Marchitto, L.,
Benlarbi, M., Chatterjee, D., Nayrac, M.*, et al.* (2021a). Strong humoral immune responses
against SARS-CoV-2 Spike after BNT162b2 mRNA vaccination with a 16-week interval between
doses. Cell Host Microbe *S1931-3128(21)00569-2*, 2021.2009.2017.21263532.

Tauzin, A., Nayrac, M., Benlarbi, M., Gong, S.Y., Gasser, R., Beaudoin-Bussieres, G., Brassard,
N., Laumaea, A., Vezina, D., Prevost, J.*, et al.* (2021b). A single dose of the SARS-CoV-2 vaccine
BNT162b2 elicits Fc-mediated antibody effector functions and T cell responses. Cell Host Microbe
29, 1137-1150 e1136.

Ullah, I., Prevost, J., Ladinsky, M.S., Stone, H., Lu, M., Anand, S.P., Beaudoin-Bussieres, G.,
Symmes, K., Benlarbi, M., Ding, S.*, et al.* (2021). Live imaging of SARS-CoV-2 infection in mice
reveals that neutralizing antibodies require Fc function for optimal efficacy. Immunity *54*, 2143-
2158 e2115.

Viana R., M.S., Amoako DG., Tegally H., Scheepers C., et al Gaseitsiwe S., von Gottberg AV.,
de Oliveira T. (2021). Rapid epidemic expansion of the SARS-CoV-2 Omicron variant in southern
Africa. medRxiv *https://krisp.org.za/manuscripts/ZHTOWa-MEDRXIV-2021-268028v1-
deOliveira.pdf*.

Weisblum, Y., Schmidt, F., Zhang, F., DaSilva, J., Poston, D., Lorenzi, J.C., Muecksch, F.,
Rutkowska, M., Hoffmann, H.H., Michailidis, E.*, et al.* (2020). Escape from neutralizing antibodies
by SARS-CoV-2 spike protein variants. eLife *9*.

WHO (2021). "Tracking SARS-CoV-2 variants".

Zhang, L., Li, Q., Liang, Z., Li, T., Liu, S., Cui, Q., Nie, J., Wu, Q., Qu, X., Huang, W.*, et al.* (2021).
The significant immune escape of pseudotyped SARS-CoV-2 Variant Omicron. Emerg Microbes
Infect, 1-11.

Supplemental Figure 1. Recognition and neutralization of different VOC Spikes by the anti-S2 neutralizing CV3-25 antibody.
(**A**) 293T cells were transfected with the full-length Spikes from different VOCs (D614G, Alpha, Beta, Gamma, Delta and Omicron), stained with the CV3-25 Ab and analyzed by flow cytometry. Shown are histograms showing a representative staining on Spike-expressing GFP+ cells. (**B**) CV3-25 neutralizing activity against pseudoviral particles bearing the different VOC Spikes.