Title

Intrinsic apoptosis is evolutionarily divergent among metazoans

Authors

Gabriel Krasovec [1,2*], Éric Quéinnec [1**], Jean-Philippe Chambon [3**]

* Authors for correspondence:

GK, gabriel.krasovec@universityofgalway.ie; gabrielkrasovec@hotmail.fr

** These authors share senior authorship.

Affiliations

[1]ISYEB, Institut de Systématique, Evolution et Biodiversité, Sorbonne Université, CNRS, MNHN,

F-75252 Paris Cedex 05, France.

[2]Centre for Chromosome Biology, School of Natural Sciences, National University of Ireland

Galway, Galway, Ireland.

[3]CRBM, Univ. Montpellier, CNRS, 34000 Montpellier, France

Abstract

Apoptosis is regulated cell death that depends on caspases. Upstream of each apoptotic signalling pathway is involved a specific initiator caspase. Characterised in nematode, fly and mammals, intrinsic apoptosis is considered to be ancestral and conserved among animals, and depends on shared initiators; caspase-9, Apaf-1 and Bcl-2. However, the biochemical role of mitochondria, the pivotal function of cytochrome c and the modality of caspase activation remain highly heterogeneous and hide profound molecular divergences among apoptotic pathways in animals. Uncovering the phylogenetic history of apoptotic actors, especially caspases, is crucial to shed light on intrinsic apoptosis evolutionary history. Here, we demonstrate by phylogenetic analyses, that caspase-9, the fundamental key of intrinsic apoptosis, is deuterostome-specific, while caspase-2 is ancestral to bilaterians. Our analysis of Bcl-2 and Apaf-1 confirm heterogeneity in functional organisation of apoptotic pathways in animals. Our results support emergence of distinct intrinsic apoptotic pathways during metazoan evolution.

INTRODUCTION

Apoptosis, regulated cell death defined by a set of morphological features and dependent on caspases, occurs during metazoan development, tissue homeostasis, and regeneration[1–3]. Pioneering works in *Caenorhabditis elegans* established a molecular network of apoptosis decision, execution, engulfment-degradation, and more fundamentally the first description of intrinsic apoptosis, previously known as the mitochondrial apoptotic pathway[4–6]. Next, investigation of apoptotic cascade key components from *Drosophila melanogaster* and mammals imposed the paradigm that the intrinsic apoptotic "molecular program" is conserved throughout animal evolution[4,5,7–9]. Intrinsic apoptosis depends on the key initiator caspase-9 (named Ced-3 and Dronc in *Caenorhabditis* and *Drosophila*, respectively), the activator Apaf-1 (named Ced-4 and Dark in *Caenorhabditis* and *Drosophila*, respectively) and the Bcl-2 multigenic family[4,8]. However, recent research has revealed evolutionary divergences between these models, with major functional diversifications[10,11]. Importantly and independently of functional approaches, the phylogenetic history of the signal transduction actors, mostly the caspase family, is the Rosetta Stone to shed light on intrinsic apoptosis evolution. Here, we investigate the evolutionary genetic history of the major actors of intrinsic apoptosis. We conducted phylogenetic analyses of initiator CARD-caspases, Bcl-2 multigenic family, and Apaf-1 proteins from major animal phyla and describe their evolutionary patterns. Despite common functional similarities, these actors, especially initiator caspases, are engaged in relatively independent evolutionary histories that involve different signalling pathways. Consequently, we consider the structural similarities of intrinsic apoptotic components to be the result of convergent evolution among animals. Convergent evolution is defined here as a set of patterns (i.e. named intrinsic apoptosis) produced by similar but not identical processes (i.e. pathways driven by at least two groups of paralogous

68   genes). Thus, the classical animal models (nematode, fly, mouse) which were used to establish

69   a unified concept of intrinsic apoptosis are markedly distinguished by distinct molecular origin

70   of their apoptotic actors but also by major functional divergences.

72   **RESULTS AND DISCUSSION**

73   **Initiator caspases of intrinsic apoptotic pathways are not homologous genes**

74   Initiation and execution of apoptotic signalling pathways are fundamentally linked to

75   the complex diversification of caspases which are widely distributed among all metazoan

76   phyla[12–15].

77   Caspases are a class of proteases composed of three protein domains; the pro-domain,

the small P10, and the large P20[16,17]. Initiator caspases, specific to each apoptotic pathway,

have a Caspase Recruiting Domain (CARD) pro-domain or two Death Effector Domains (DED)

pro-domains. Intrinsic and extrinsic apoptosis are distinct and involve specific initiators which

are caspase-9 (CARD pro-domain) or caspase-8 or -10 (DED pro-domains), respectively. Both

pathways trigger activation of common executioner caspases, leading to apoptosis

execution[4,18].

84   Due to the pivotal role of caspase-9 in initiating intrinsic apoptosis, our main analyses

focused on caspases having a CARD pro-domain (CARD-caspases). We confirmed their

distribution in most animal phyla and reconstructed their phylogenetic relationships (Figure 1,

Table S1). Regardless of the phylogenetic methodology used for the analysis, two strongly

supported (PP> 0.99) monophyletic groups were identified: caspase-9 and caspase-2 (Figure

1, Figure S1). A third and more heterogeneous group named here [Inflammatory Caspases +

Caspase-Y] is monophyletic in Bayesian analysis, the method which produced the most likely

topology (Table S2), and splits into two groups in maximum likelihood estimation (Figure 1,

Figure S1). Although relationships among bilaterian clades differ depending on the

methodology employed, monophyly of caspase-9 and caspase-2 families remains robust. They

group together with Inflammatory Caspases and Caspase-Y in a clade strictly diversified within

bilaterian animals (PP = 0.87; BS>90). The sequences of non-bilaterians [Cnidaria + Ctenophora

+ Placozoa] form a divergent paraphyletic group named here Caspase-X (Figure 1). Switching

between metazoan putative "basal" out groups [i.e. *Amphimedon queenslandica* (Porifera) or

Pleurobrachia bachei (Ctenophora)] does not affect caspase topology (Figure S2). All

bilaterians sequences, caspase-9, caspase-2 and [Inflammatory Caspases + Caspase-Y] remain

monophyletic (Figure S2).

Caspase-2 appears to be widely distributed among bilaterians and sequences evolution

is congruent with the bilaterian groups [Deuterostomia + Ecdysozoa +

Lophotrochozoa/Spiralia] reflecting its ancestral origin. Surprisingly caspase-9 is strictly

restricted to deuterostome animals. This distribution can be interpreted either as the early

emergence of caspase-9 in all bilaterians followed by a secondary loss in protostomes, or as a

specific derived character of deuterostomes. The diversity of the genomes explored, and their

depth of sequencing unambiguously confirms this result. More specifically, the robustness of

the deuterostome caspase-9 clade was explored (Figure S3), and its specific diversification

among [Vertebrata + Cephalochordata + Echinodermata + Hemichordata] was confirmed.

However, the relative position of the cephalochordate *Branchiostoma belcheri* 9A paralogous

gene remains unstable. Consistent with previous studies on the ascidian *Ciona intestinalis*[19],

the five representatives of ascidian genomes studied are devoid of any caspase-9, likely due

to a unique secondary loss in urochordates, the sister-group of vertebrates[20].

Unexpectedly, *Drosophila* Dronc and *Caenorhabditis* Ced-3 are distinctly identified as

orthologous (i.e. coming from speciation) to caspase-2 of vertebrates, and not to caspase-9,

as was previously reported and largely accepted[21–28]. All Dronc/Ced3 proteins of insects,

horseshoe crab (Xiphosura), and nematodes form a strongly supported monophyletic clade

(PP = 1) revealing single gene conservation without duplication (except horseshoe crab). Thus,

the caspase-2 clade of ecdysozoans is the sister group of caspase-2 clade of [Lophotrochozoa

+ Deuterostomia]. The absence of identifiable caspase-2 in echinoderms and hemichordates

(i.e. Ambulacraria) can probably be interpreted as a clade-specific derived loss.

Conversely, the clade [Caspase-Y + Inflammatory Caspase] appears to be restricted to

[Vertebrata and Lophotrochozoa].

To test evolutionary history and shuffling of different domains, we inferred

recombination rate from CARD-Caspase coding sequences (reflecting intra-codon

recombination). We investigated multiple alignments of domains sequences using the

approximate Bayesian computation (ABC) approach based on coalescent evolutionary history

of protein sequences, i.e. the framework *ProteinEvolverABC*[29]. The remarkable low level of

recombination rate (Rho mean = 2.0328712) we estimated during CARD-Caspase family

evolution rule out potential phylogenetic reconstruction artefacts (Table S3). As illustrative

examples, this low recombination rate is comparable to those observed for some

Coronaviruses protein families such as NS12.7 or astroviruses viroporins probably related to

the maintenance of viral pathogenicity[29–31]. The multiple and essential roles played by the

CARD-Caspase family within the cell is likely one factor explaining the extremely low

recombination rate that can be observed.

Taken globally, our analysis shows that the traditionally considered conserved initiator

caspases of intrinsic apoptosis are not orthologous, but are divergent genes putatively issued

from ancestral duplication (i.e. paralogous genes) among bilaterians.

The caspase activation regulator apoptosome is structurally divergent in metazoans

In intrinsic apoptosis, initiator caspase activation depends on their recruitment by a pivotal, and considered shared component, the apoptosome platform[4,10]. Apoptosome formation results on critical protein binding between *Caenorhabditis* Ced-4, *Drosophila* Dark, and human Apaf-1 with their respective initiator caspases Ced-3, Dronc, and procaspase-9, respectively[10]. CARD and other domains (NOD, arm) are highly conserved in Apaf-1, Dark and Ced-4 proteins[32–34]. However, excepted the majority of nematode species[11], Apaf-1 possesses WD40 repeats at its C-terminus which bind to cytochrome c. This binding is required in mammals for Apaf-1 oligomerisation and apoptosome formation[34,35], while Ced-4 and Dark do not require cytochrome c for their general assembly into an apoptosome[10,36]. Taken with major structural and regulating assembly differences between the octameric Dark, tetrameric Ced-4 and heptameric Apaf-1 apoptosomes, it reveals evolutionary divergence between animal apoptosomes formation and procaspases activation mechanisms, probably differentially modulating the cell death execution pathway[37].

We conducted exhaustive studies by reciprocal BLAST and phylogenetic analyses of Apaf-1 homologs and confirmed their conservation in the majority of metazoan phyla (Figure S4, Table S4)[32,33]. We conducted our analysis with all sequences issued from a previous study[33] supplemented with ones we have specifically identified. However, we have differentiated in our study Apaf-1 genes sharing a CARD domain, and likely involved in apoptosis regulation, from Apaf-1-like devoid of CARD domains but which possess a panel of unusual domains. NB-ARC being the most conserved and shared domain by almost all Apaf-1 and Apaf-1-like genes among metazoans, we conducted the analysis on the NB-ARC alignment as it was previously done[33].

Our topology presents two main clades that are common between Bayesian inference

method and maximum likelihood estimation, both of which are inconsistent with species

phylogeny. One of them grouped ecdysozoan sequences including Ced-4 and Dark in addition

to several Apaf-1-like from the cnidarian *Nematostella vectensis* and the cephalochordate

Branchiostoma floridae, while the other one contains deuterostomian and non-bilaterian

sequences. Unexpectedly, deuterostomians are not monophyletic but split into two groups,

suggesting classic Long Branch attraction artifact and a very fast sequence evolution.

Remarkably, Apaf-1 was absent from all available urochordate and mollusc genomes, in

accordance with previous work[19,33,38–41], and from *Capitella teleta* (Annelida). Even if

differences in the quality of the genomes might account for some of the putative losses, these

results suggest independent losses in urochordates and lophotrochozoans or, a less

parsimonious hypothesis, complex and independent modular protein evolution from the

common ancestor of metazoans (Figure 2). In non-bilaterians, the Apaf-1 gene family has not

been found in the *Pleurobrachia pileus* genome (Ctenophora), but seems present in other

early diverging metazoan phyla [Porifera, Cnidaria and Placozoa] ([33,42,43], our data).

Importantly, a previous analysis from several genomes (cnidarian, nematode, fly,

amphioxus, sea urchin, human) clearly identified three well-defined independent clades of

genes among metazoans, highlighting that Ced-4, Dark and Apaf-1 are not homologous genes

between ecdysozoans and vertebrates[33].

Despite its fundamental role in the apoptotic cell death pathways and also in non-

apoptotic functions[44–46], diverse metazoan phyla have independently experienced adapter

protein Apaf-1 losses. Apaf-1 with CARD-domain, likely involve in apoptosis, are under strong

purifying selection as illustrated by the metric non-synonymous/synonymous substitution rate

ratio (dN/dS) close to 0 (model M0; $\omega\theta = 0.07$), perhaps promoted by their crucial and specific

role as caspase activator. Consequently, we hypothesise a relaxation of functional constrains

on unusual Apaf-1 molecules (i.e. deprived of Card-domain, and probably not involved in

oligomerisation processes) that specifically drive the diversification of Apaf-1/Apaf-1-like

paralogous genes in the cnidarian *Nematostella vectensis* and the cephalochordate

Branchiostoma floridae[33].

These elements argue in favour of distinct evolutionary trajectory between species and

convergence of the modality of apoptosome formation, which could explain functional

analyses showing that its formation and the subsequent caspase activation differs among

taxa[11].

The regulation of apoptosis by the Bcl-2 family is divergent among metazoans

Intrinsic apoptosis is ultimately regulated by the Bcl-2 proteins, composed of several

Bcl-2 homologous (BH) domains[47,48]. In mammals, the balance between pro-survival (four

BH1-BH4 domains) and pro-apoptotic proteins (Bax/Bak/Bok and BH3-only) of Bcl-2 controls

initiation of intrinsic apoptosis. Conformational changes of the three-dimensional structures

and interactions between Bcl-2 actors enables the assembly of pore-like structures controlling

MOMP[49].

Multiple sequence alignments of metazoan Bcl-2 family proteins (Figure S5, Table S5)

(but with over-represented chordates reflective of greater availability of vertebrate genomes)

confirm the wide distribution and early origin of Bcl-2 in Metazoa[11,32,48,50]. Proteins clustered

into five monophyletic groups, consistent with classical functional Bcl-2 classification (i.e.

three 'pro-apoptotic' clades: Bok, Bak, Bax and two less supported pro-survival - 'anti-

apoptotic' - groups: Bcl-2/W/XL and a more complex Bcl-B/Mcl-1/Bfl-1 clade) (Figure S5).

However, relationships among the five well-supported groups were not clearly resolved

despite the use of various methodologies and outgroup selections (Figure S5A-B). Consequently, each of the five Bcl-2 clades seems conserved across evolution, and homology (orthology) relationships inside each clade can be established. However, due to the impossibility of addressing the relationship between these five monophyletic groups (Bok, Bak, Bax, Bcl-2/W/XL, Bcl-B/Mcl-1/Bfl-1), it seems difficult to hypothesises their evolutionary pattern, history, and determine a potential ancestral group. In addition, very divergent sequences from molluscs such as *Biomphalaria glabrata* (Bcl-like2, Bcl-like3), from the cnidarian *Hydra vulgaris* (Bcl-like1) or from the urochordate *Ciona intestinalis* (Bcl-like1) were not assigned to a particular class.

Caenorhabditis is deprived of pro-apoptotic Bcl-2 and possesses only the pro-survival Ced-9 (homolog to vertebrates Bcl-2/w/xl), and two BH-3 only proteins (Egl-1 and Ced-13). Conversely, only two pro-apoptotic Bok-like paralogs (Debcl and Buffy) were present in *Drosophila* (Figure 2, Figure S5), but their functions remain unclear[51]. Presence of both anti- and pro-apoptotic Bcl-2 in molluscs underlines the divergent particularities observed within protostomes. Contrarily, similarities in Bcl-2 family composition are observed within some deuterostomes (mammals and echinoderms).

Finally, if multiple Bcl-2 genes were acquired early in metazoan evolution, and despite conservation of almost homologous genes, key differences accumulate, making initiation mechanisms in intrinsic apoptotic signalling pathways divergent among animals.

Apoptotic mitochondrial pathways are divergent among metazoans

Functional evidence shows that caspase-2 members play a critical role in various cell death processes, but are independently involved in a range of non-apoptotic functions, including cell cycle regulation, DNA repair and tumour suppression[52–55]. This implication of

caspase-2 in a myriad of signalling pathways, and its ability to interact with a panel of molecules, demonstrates its functional versatility[52,53,56–59]. As previously reported, our phylogenetic analyses corroborate the wide distribution of caspase-2 in bilaterians and suggest a probable ancestral multifunctionality.

The major structural and functional similarities that led to an erroneous interpretation of the phylogenetic position of Ced-3 and Dronc underlies a probable common evolutionary origin of caspase-2 and -9 genes. We hypothesised here that caspase-9 originates from bilaterians-specific duplication of a caspase-like ancestor gene, followed by a loss of caspase-9 in protostomians (Figure 2). Among deuterostomians the two families of paralogs have been preserved in vertebrates and probably in cephalochordates (Figure 1, 2). Caspase-2 retains multifunctional activity, and in mammals can interact with the PIDDosome platform containing P53, adapter molecule RAIDD, and signalling complex DISC, activating both extrinsic and intrinsic apoptosis, and DNA damage pathways[58,60–62]. Conversely, the caspase-9 gene underwent a functional divergence in connection with its specialisation in allosteric interactions with the apoptosome[18].

Due to its pivotal role as a mediator of genomic stability through involvement in cell proliferation, oxidative stress, aging and cell death, the molecular divergence of the caspase-2 gene is highly constrained during evolution, probably because destabilisation of any signalling cascade is sufficient to initiate tumorigenesis. Indeed, purifying selection is likely important during caspase-2 evolution as illustrated by a dN/dS close to 0 (model M0 $\omega\theta$ = 0.07)[63,64]. However, during the radiation of deuterostomes, both presence of caspase-2 and -9 certainly facilitated the loss of one of them (cf. DCC Force's model -Duplication-Degeneration-Complementation model)[65–67]. Interestingly, a caspase-2 duplication co-occurred with the loss of caspase-9 in urochordates that may indicate a clade-specific

relaxation of purifying selection on caspase-2 gene (Figure 1, Figure 2). Similarly, caspase-2

has been lost, or strongly diverged, while caspase-9 presents a *de novo* relative expansion in

Strongylocentrotus purpuratus (Echinodermata) and *Saccoglossus kowalevskii*

(Hemichordata). These expansions certainly result from a less powerful purifying selection of

caspase-9, showing a higher dN/dS (model M0 $\omega\theta$ = 0.22) than caspase-2.

Due to generalised gene losses in ecdysozoans and in comparison with other

bilaterians, *Caenorhabditis* and *Drosophila* apoptotic pathways are generally considered

simpler than those of vertebrates[17,68]. However, comparing to mammals, ecdysozoans have

pathways organised around different paralogous genes in addition to a smaller number of

genes present. The absence of orthologous genes relationships results in a very different

structural organisation of apoptosome platforms but also generates important functional

divergences (i.e. mechanisms of regulating assembly, CARD-CARD interactions with

procaspases)[10].

Consistent with mammalian caspase-2 functions, Ced-3 has both initiator as well as

executioner activities[58,69–71]. Involvement of Dronc in various processes such as compensatory

cell proliferation, inhibition of cell migration or spermatid differentiation, brings this protein

closer to the functionalities of mammalian caspase-2[72,73]. Due to its interaction with Dark

(Apaf-1 paralog), the only CARD-caspase in *Drosophila* (Dronc) has been erroneously classified

as caspase-9[8,74]. Unlike the organisation of the mammal apoptosome, both *Caenorhabditis*

and *Drosophila* present neither MOMP nor the release and necessity of cytochrome c to

activate Ced-3 and Dronc via Ced-4 and Dark, respectively[5,68].

Like other protostomes, molluscs are devoid of caspase-9 but caspase-2 has been

identified in bivalves and was suspected to function in "a caspase-9-like manner"[75]. Hence,

caspase-2 seems involved in apoptotic processes in molluscs, but more surprisingly, despite

the absence of caspase-9 and Apaf-1 (and thus, of a mammalian-like apoptosome), this

peculiar pathway is surprisingly associated with cytochrome c release[38,41,76–79]. The complexity

of intrinsic apoptosis in molluscs seems to be important, but divergent from what was

observed in ecdysozoans or vertebrates, with a putative expansion of caspases that

participate both in immunity, stress responses, and apoptosis (Figure 2)[41,75,80–83].

Unexpectedly, mammals present an almost unique case (possibly with the

cephalochordates) in which both caspase-2 and caspase-9 are conserved and involved in

apoptosis. This putative functional redundancy (i.e. recruitment, autoactivation or

transactivation, homodimerisation and subsequent interchain proteolytic cleavage)

undoubtedly led to the functional specialisation observed for caspase-9. This may results from

the specificity of the mammalian mitochondrial pathway and the non-apoptotic function of

caspase-9[84–87]. Finally, echinoderms seem to uniquely have an intrinsic apoptosis similar to

mammals, with a caspase-9, Bcl-2, Apaf-1, and a MOMP with cytochrome c release (Figure

2)[7,88].

Although we can envisage a weak parsimonious scenario that showcases a common

ancestral apoptotic pathway in deuterostomes (but implying independent secondary losses in

hemichordates, cephalocordates, and urochordates), the similarities observed between

echinoderms and mammals more probably reflect "functional convergences" based on

independent recruitment of apoptotic actors.

CONCLUSION

The apoptotic networks of *Caenorhabditis* and *Drosophila* do not exemplify ancestral

conditions from which mammalian-grade apoptotic complexity emerged but are on the

contrary, and as suggested recently, the result of a derived condition specific to ecdysozoans

among animals [32,80,89,90].

The core components of intrinsic apoptotic pathways, especially initiator caspases and

the apoptosome platform, are not ancestral in metazoans. Our phylogenetic analyses highlight

an unexpected evolutionary history: while the bilaterian caspase-2-mediated apoptotic toolkit

emerged ancestrally and remains multifunctional, the caspase-9 mediator of the mammalian

apoptosome is specific to deuterostomes.

The major functional divergences in mitochondrial apoptotic pathways observed in

animals[70,91–93] mainly originated in the recruitment of paralogous actors from the same

multigenic families. The evolution of the different pathways underlined by non-homologous

molecular actors could be the reflection of putative adaptive processes or constrains specific

to each taxon, which ultimately led to divergent evolutionary histories. Interestingly, the

richness of the apoptotic genetic repertoire was suggested to be linked to the persistence of

stem cells in adults from different phyla[94,95].

Finally, mitochondria-mediated apoptosis, like other programmed cell deaths, has

likely evolved before and throughout metazoan diversification to shape developmental

processes, immune response, or to adapt cellular environment to environmental constraint.

MATERIAL AND METHODS

Sequence dataset construction

Putative metazoan caspases with a CARD pro-domain (CARD-caspases) were identified

using tBLASTn and BLASTp searches with human caspases, Ced-3, and Dronc as queries on

NCBI, ANISEED (ascidians), EchinoBase (*Strongylocentrotus purpuratus*), and

neurobase.rc.ufl.edu (*Pleurobrachia bachei*) databases, and followed by reciprocal BLAST.

After identification of CARD-caspases in target species, sequences were added as queries to

conduct BLAST searches in close relatives (i.e. identified CARD-caspases of *Crassostrea gigas*

were used as queries to search in other molluscs). Genomes and transcriptomes of species'

were downloaded when possible, to search a second time. Sequences with an e-value inferior

to 1e-10 were retained. All identified sequences were analysed with ScanProsite (ExPaSy)[96]

and InterProScan (EMBL-EBI)[97] to verify the presence of specific caspase domains. Another

sequence was added to verify its identification proposed as a caspase-2 in the literature:

Crassostrea angulata caspase-2[79]. Caspase family proteins are short (containing the large

common P20 and the small P10 domains) with a high number of genes per species which

rapidly limits the relevance of the phylogenetic analyses. To reduce artefact branching and

unreadable topologies, and to maximize phylogenetic diversity across Metazoa, the dataset

was built using CARD-caspase gene repertoires of selected species. A full list of all caspase

sequences is provided in Table S1. Taxon and genes were selected to have a wide diversity of

major metazoan phyla and to have an equilibrium inside and between each group. We choose

representative species for each phylum and took care to don't unreasonably increase the

number of sequences by redundant choice. We privileged species where we detected

caspases with CARD domains, DED domains, and executioner caspases, suggesting a high

quality of sequences allowing identification of all members of the caspase family. For each

taxon, all caspases having a CARD pro-domains have been include in the analysis.

Identification of metazoan Apaf-1 was made using tBLASTn and BLASTp using human

APAF-1, nematode Ced-4, and fly Dark as queries on NCBI, ANISEED (ascidians), and

neurobase.rc.ufl.edu (*Pleurobrachia bachei*) databases, and followed by reciprocal BLAST.

Potential resulting sequences were analysed with ScanProsite and also InterProScan. A full list

of all Apaf-1 sequences is provided in Table S4. Apaf-1 data set comprise sequences from a

previous study[33] in addition to our new sequences.

Metazoan Bcl-2 were identified by using tBLASTn and BLASTp searches with human

Bcl-2 as the query sequence on NCBI, and then again on downloaded genomes and

transcriptomes, followed by reciprocal BLAST. All identified sequences were analysed with

ScanProsite (ExPaSy) and InterProScan (EMBL-EBI) to verify the presence of BH domains. Due

to short sequence length, BH3-only were not taken into account. A full list of all Bcl-2

sequences is provided in Table S5.

Multiple alignments of protein sequences were generated using the MAFFT software

version 7[98] with default parameters and also Clustal Omega[99] to verify the congruence of the

different alignments. All sequences were then manually checked in BioEdit 7.2 software[100] to

verify the presence of specific domains previously identified. Gblocks version 0.91b[101] was

used to remove vacancies and blur sites. Final alignments comprise of 230, 235, 147, and 306

amino acids for metazoan CARD-caspases alignment, deuterostomian CARD-caspases

alignment, metazoan Bcl-2 alignment, and Apaf-1 alignment, respectively. Apaf-1 alignment

was done on the NB-ARC domain only.

Phylogenetic analysis

Phylogenetic analyses were carried out from the amino-acid alignment with

Maximum-Likelihood (ML) method in PhyML 3.1[102], a combined ML tree search with 1000

bootstrap replicates was produced and then visualised using Seaview[103]. The best amino-acid

evolution models to conduct analyses were determined using MEGA11[104] and determined to

be WAG for sequences of CARD-caspases alignments (CARD, P20, P10) and LG for both Bcl-2

and Apaf-1 alignments.

Bayesian analyses were performed using MrBayes (v3.2.6)[105] under mixed model. For

each analysis, one fourth of the topologies were discarded as burn-in values, while the

remaining ones were used to calculate posterior probability. The run for metazoan CARD-

caspases alignment were carried out for 2,000,000 generations with 15 randomly started

simultaneous Markov chains (1 cold chain, 14 heated chains) and sampled every 100

generations. The run for deuterostomian CARD-caspases alignment was carried out for

500,000 generations with 5 randomly started simultaneous Markov chains (1 cold chain, 4

heated chains) and sampled every 100 generations. The run for metazoan Apaf-1 alignment

was carried out for 500,000 generations with 5 randomly started simultaneous Markov chains

(1 cold chain, 4 heated chains) and sampled every 100 generations. The run for metazoan Bcl-

2 alignment was carried out for 5,000,000 generations with 20 randomly started simultaneous

Markov chains (1 cold chain, 19 heated chains) and sampled every 100 generations. ML

bootstrap values higher than 50% and Bayesian posterior probabilities are indicated on the

Bayesian tree.

The outgroup for the metazoan CARD-caspase phylogeny is the only caspase with a

CARD pro-domain of the Porifera *Amphimedon queenslandica* (XP_003383519) (Figure 1, S4,

S5) or the only one of the ctenophore *Pleurobrachia bachei* (Sb 2658116) (Figure S2). Analyses

of CARD-caspases were made independently at the deuterostome scale with four different

outgroups to test their effect on the stability of the topology: *i*) CARD-caspase-Y of the annelid

Capitella teleta (ELT97848.1), *ii*) CARD-caspase-2 of the mollusk *Aplysia californica*

(XP_005113266), *iii*) CARD-caspase-X2 of cnidarian *Hydra vulgaris* (NP_001274285.1) *iv*)

CARD-caspase Ced-3 of the ecdysozoan *Caenorhabditis elegans* (AAG42045.1) (Figure S3). For

the metazoan Apaf-1 phylogenies, the selected outgroup is Apaf-1 (XP_019855714.1) of

Porifera *Amphimedon queenslandica* (Figure S4). For the metazoan Bcl-2 phylogenies,

outgroups used to test their effect on stability of the topology are: *i*) Bcl-2-like1

(XP_003383425.1) and Bcl-2-like2 (XP_003387574.1) of Porifera *Amphimedon queenslandica*

(Figure S5).

Topology tests between metazoan CARD-caspase phylogenies from Bayesian inference

method and maximum likelihood estimation were evaluated using IQ-TREE[106]. We conducted

Kishino-Hasegawa test[107], Shimodaira-Hasegawa test, and approximately unbiased test[108].

Estimation of dN/dS and recombination rate

Codon aligned nucleotide sequences were built using PAL2NAL[109] from clustalo aligned

amino acid sequences and gene ORFs. The dN/dS ratio was calculated using EasyCodeML

under settings "site model", nuc-2, and standard icode[64], from the codon aligned nucleotide

sequences and newick format trees.

Co-estimation of recombination rate was conducted using ProteinEvolverABC[29]. We

run the analysis on the same multiple protein alignment we used for the metazoan CARD-

caspases phylogenetic analysis, including CARD, P20, and P10 domains. Analysis was

conducted with 10 000 simulations and an acceptance rate of 0.2% to optimise the accuracy

of the estimations (Beaumont 2010), and under WAG amino acid substitution model. Prior

parameters were estimated by ProteinEvolverABC prior to the analysis such as Rho = Uniform

(0, $3.0e^{-6}$) and Theta = Uniform ($9.1e^{-5}$, $4.0e^{-4}$). Proportion of invariable sites was 0.026 and

amino acid frequencies were fix.

Acknowledgements

Authors particularly acknowledge Helen R Horkan (University of Galway, Ireland) for

help in bioinformatics analysis and helpful comments, and Miguel Arenas (University of Vigo,

Spain) for discussion and assistance with ProteinEvolverABC. Authors thank Sébastien Darras

(Sorbonne Université, Banyuls-sur-Mer), Christine Vesque (Sorbonne Université, Paris),

Jérôme Gros (Institut Pasteur, Paris), Sabine Hennequin (Sorbonne Université, Paris) and Uri

Frank (University of Galway, Ireland) for helpful comments. GK was supported by a Ph.D.

fellowship from the French Ministry of Education, Research and Innovation.

Author Contributions

JP and EQ managed the project. GK made BLAST, phylogenetic and sequences analysis, and

figures. GK and EQ wrote the manuscript.

Competing Interest Statement

The authors declare no competing interests.

Data Availability

All data needed to evaluate the conclusions in this study are present in the paper and the

Supplementary Materials. Any requests can be addressed to the corresponding author.

BIBLIOGRAPHY

1. Jacobson, M. D., Weil, M. & Raff, M. C. Programmed cell death in animal development. *Cell* **88**,

347–354 (1997).

2. Hipfner, D. R. & Cohen, S. M. Connecting proliferation and apoptosis in development and

disease. *Nat Rev Mol Cell Biol* **5**, 805–815 (2004).

3. Vriz, S., Reiter, S. & Galliot, B. Cell death: a program to regenerate. *Curr. Top. Dev. Biol.* **108**, 121–

151 (2014).

4. Hengartner, M. O. The biochemistry of apoptosis. *Nature* **407**, 770–776 (2000).

5. Lettre, G. & Hengartner, M. O. Developmental apoptosis in C. elegans: a complex CEDnario. *Nat*

Rev Mol Cell Biol **7**, 97–108 (2006).

6. Ellis, H. M. & Horvitz, H. R. Genetic control of programmed cell death in the nematode C.

elegans. *Cell* **44**, 817–829 (1986).

7. Bender, C. E. *et al.* Mitochondrial pathway of apoptosis is ancestral in metazoans. *Proc. Natl.*

Acad. Sci. U.S.A. **109**, 4904–4909 (2012).

8. Steller, H. Regulation of apoptosis in Drosophila. *Cell Death Differ.* **15**, 1132–1138 (2008).

9. Driscoll, M. Cell death in C. elegans: molecular insights into mechanisms conserved between

nematodes and mammals. *Brain Pathol* **6**, 411–425 (1996).

10. Dorstyn, L., Akey, C. W. & Kumar, S. New insights into apoptosome structure and function. *Cell*

Death Differ **25**, 1194–1208 (2018).

11. Young, N. D. *et al.* Diversity in the intrinsic apoptosis pathway of nematodes. *Commun Biol* **3**, 478

(2020).

12. Bell, R. A. V. & Megeney, L. A. Evolution of caspase-mediated cell death and differentiation:

twins separated at birth. *Cell Death Differ* **24**, 1359–1368 (2017).

13. Shrestha, S. *et al.* Caspases from scleractinian coral show unique regulatory features. *J Biol Chem*

295, 14578–14591 (2020).

14. Uren, A. G. *et al.* Identification of Paracaspases and Metacaspases: Two Ancient Families of

Caspase-like Proteins, One of which Plays a Key Role in MALT Lymphoma. *Molecular Cell* **6**, 961–

967 (2000).

15. Crawford, E. D. *et al.* Conservation of caspase substrates across metazoans suggests hierarchical

importance of signaling pathways over specific targets and cleavage site motifs in apoptosis. *Cell*

Death Differ **19**, 2040–2048 (2012).

16. Thornberry, N. A. & Lazebnik, Y. Caspases: enemies within. *Science* **281**, 1312–1316 (1998).

17. S, E. Apoptosis: a review of programmed cell death. *Toxicologic pathology* vol. 35

https://pubmed.ncbi.nlm.nih.gov/17562483/ (2007).

18. Galluzzi, L. *et al.* Molecular mechanisms of cell death: recommendations of the Nomenclature

Committee on Cell Death 2018. *Cell Death Differ.* **25**, 486–541 (2018).

19. Terajima, D. *et al.* Identification of candidate genes encoding the core components of the cell

death machinery in the Ciona intestinalis genome. *Cell Death Differ.* **10**, 749–753 (2003).

20. Delsuc, F., Brinkmann, H., Chourrout, D. & Philippe, H. Tunicates and not cephalochordates are

the closest living relatives of vertebrates. *Nature* **439**, 965–968 (2006).

21. Lee, T. V., Kamber Kaya, H. E., Simin, R., Baehrecke, E. H. & Bergmann, A. The initiator caspase

Dronc is subject of enhanced autophagy upon proteasome impairment in Drosophila. *Cell Death*

Differ **23**, 1555–1564 (2016).

22. Fogarty, C. E. *et al.* Extracellular Reactive Oxygen Species Drive Apoptosis-Induced Proliferation

via Drosophila Macrophages. *Curr Biol* **26**, 575–584 (2016).

23. Fan, Y. *et al.* Genetic models of apoptosis-induced proliferation decipher activation of JNK and

identify a requirement of EGFR signaling for tissue regenerative responses in Drosophila. *PLoS*

Genet **10**, e1004131 (2014).

24. Fan, Y. & Bergmann, A. The cleaved-Caspase-3 antibody is a marker of Caspase-9-like DRONC

activity in Drosophila. *Cell Death Differ* **17**, 534–539 (2010).

25. Napoletano, F. *et al.* p53-dependent programmed necrosis controls germ cell homeostasis

during spermatogenesis. *PLoS Genet* **13**, e1007024 (2017).

26. Lim, Y., Dorstyn, L. & Kumar, S. The p53-caspase-2 axis in the cell cycle and DNA damage

response. *Exp Mol Med* **53**, 517–527 (2021).

27. Bao, Q. & Shi, Y. Apoptosome: a platform for the activation of initiator caspases. *Cell Death Differ*

14, 56–65 (2007).

28. Jaroszewski, L., Rychlewski, L., Reed, J. C. & Godzik, A. ATP-activated oligomerization as a

mechanism for apoptosis regulation: fold and mechanism prediction for CED-4. *Proteins* **39**, 197–

203 (2000).

29. Arenas, M. ProteinEvolverABC: Coestimation of Recombination and Substitution Rates in Protein

Sequences by approximate Bayesian computation. *Bioinformatics* btab617 (2021)

doi:10.1093/bioinformatics/btab617.

30. Lulla, V. & Firth, A. E. A hidden gene in astroviruses encodes a viroporin. *Nat Commun* **11**, 4070

(2020).

31. Nieva, J. L., Madan, V. & Carrasco, L. Viroporins: structure and biological functions. *Nat Rev*

Microbiol **10**, 563–574 (2012).

32. Zmasek, C. M. & Godzik, A. Evolution of the animal apoptosis network. *Cold Spring Harb Perspect*

Biol **5**, a008649 (2013).

33. Zmasek, C. M., Zhang, Q., Ye, Y. & Godzik, A. Surprising complexity of the ancestral apoptosis

network. *Genome Biol.* **8**, R226 (2007).

34. Zou, H., Henzel, W. J., Liu, X., Lutschg, A. & Wang, X. Apaf-1, a human protein homologous to C.

elegans CED-4, participates in cytochrome c-dependent activation of caspase-3. *Cell* **90**, 405–413

(1997).

35. Zou, H., Li, Y., Liu, X. & Wang, X. An APAF-1.cytochrome c multimeric complex is a functional

apoptosome that activates procaspase-9. *J. Biol. Chem.* **274**, 11549–11556 (1999).

36. Dorstyn, L. *et al.* The role of cytochrome c in caspase activation in Drosophila melanogaster cells.

J Cell Biol **156**, 1089–1098 (2002).

37. Shi, Y. Mechanical aspects of apoptosome assembly. *Curr Opin Cell Biol* **18**, 677–684 (2006).

38. Li, Y. *et al.* Conservation and divergence of mitochondrial apoptosis pathway in the Pacific oyster,

Crassostrea gigas. *Cell Death Dis* **8**, e2915 (2017).

39. Kiss, T. Apoptosis and its functional significance in molluscs. *Apoptosis* **15**, 313–321 (2010).

40. Romero, A., Novoa, B. & Figueras, A. The complexity of apoptotic cell death in mollusks: An

update. *Fish Shellfish Immunol.* **46**, 79–87 (2015).

41. Estévez-Calvar, N., Romero, A., Figueras, A. & Novoa, B. Genes of the mitochondrial apoptotic

pathway in Mytilus galloprovincialis. *PLoS ONE* **8**, e61502 (2013).

42. Srivastava, M. *et al.* The Amphimedon queenslandica genome and the evolution of animal

complexity. *Nature* **466**, 720–726 (2010).

43. Srivastava, M. *et al.* The Trichoplax genome and the nature of placozoans. *Nature* **454**, 955–960

(2008).

44. Zermati, Y. *et al.* Nonapoptotic role for Apaf-1 in the DNA damage checkpoint. *Mol Cell* **28**, 624–

637 (2007).

45. Ferraro, E. *et al.* Apaf1 plays a pro-survival role by regulating centrosome morphology and

function. *J Cell Sci* **124**, 3450–3463 (2011).

46. Shakeri, R., Kheirollahi, A. & Davoodi, J. Apaf-1: Regulation and function in cell death. *Biochimie*

135, 111–125 (2017).

47. Lee, E. F. *et al.* Discovery and molecular characterization of a Bcl-2-regulated cell death pathway

in schistosomes. *Proc. Natl. Acad. Sci. U.S.A.* **108**, 6999–7003 (2011).

48. Banjara, S., Suraweera, C. D., Hinds, M. G. & Kvansakul, M. The Bcl-2 Family: Ancient Origins,

Conserved Structures, and Divergent Mechanisms. *Biomolecules* **10**, (2020).

49. Kalkavan, H. & Green, D. R. MOMP, cell suicide as a BCL-2 family business. *Cell Death Differ* **25**,

46–55 (2018).

50. Dunn, S. R., Phillips, W. S., Spatafora, J. W., Green, D. R. & Weis, V. M. Highly conserved caspase

and Bcl-2 homologues from the sea anemone Aiptasia pallida: lower metazoans as models for

the study of apoptosis evolution. *J. Mol. Evol.* **63**, 95–107 (2006).

51. Clavier, A., Rincheval-Arnold, A., Colin, J., Mignotte, B. & Guénal, I. Apoptosis in Drosophila:

which role for mitochondria? *Apoptosis* **21**, 239–251 (2016).

52. Krumschnabel, G., Sohm, B., Bock, F., Manzl, C. & Villunger, A. The enigma of caspase-2: the

laymen's view. *Cell Death & Differentiation* **16**, 195–207 (2009).

53. Olsson, M., Forsberg, J. & Zhivotovsky, B. Caspase-2: the reinvented enzyme. *Oncogene* **34**,

1877–1882 (2015).

54. Lassus, P., Opitz-Araya, X. & Lazebnik, Y. Requirement for caspase-2 in stress-induced apoptosis

before mitochondrial permeabilization. *Science* **297**, 1352–1354 (2002).

55. Zhivotovsky, B. & Orrenius, S. Caspase-2 function in response to DNA damage. *Biochem. Biophys.*

Res. Commun. **331**, 859–867 (2005).

56. Krumschnabel, G., Manzl, C. & Villunger, A. Caspase-2: killer, savior and safeguard--emerging

versatile roles for an ill-defined caspase. *Oncogene* **28**, 3093–3096 (2009).

57. Olsson, M. *et al.* DISC-mediated activation of caspase-2 in DNA damage-induced apoptosis.

Oncogene **28**, 1949–1959 (2009).

58. Lavrik, I. N., Golks, A., Baumann, S. & Krammer, P. H. Caspase-2 is activated at the CD95 death-

inducing signaling complex in the course of CD95-induced apoptosis. *Blood* **108**, 559–565 (2006).

59. Braga, M. *et al.* Involvement of oxidative stress and caspase 2-mediated intrinsic pathway

signaling in age-related increase in muscle cell apoptosis in mice. *Apoptosis* **13**, 822–832 (2008).

60. Duan, H. & Dixit, V. M. RAIDD is a new 'death' adaptor molecule. *Nature* **385**, 86–89 (1997).

61. Tinel, A. & Tschopp, J. The PIDDosome, a protein complex implicated in activation of caspase-2 in

response to genotoxic stress. *Science* **304**, 843–846 (2004).

62. Haupt, S., Berger, M., Goldberg, Z. & Haupt, Y. Apoptosis - the p53 network. *J Cell Sci* **116**, 4077–

4085 (2003).

63. Eyre-Walker, A. The genomic rate of adaptive evolution. *Trends Ecol Evol* **21**, 569–575 (2006).

64. Gao, F. *et al.* EasyCodeML: A visual tool for analysis of selection using CodeML. *Ecol Evol* **9**, 3891–

3898 (2019).

65. Force, A. *et al.* Preservation of duplicate genes by complementary, degenerative mutations.

Genetics **151**, 1531–1545 (1999).

66. Conant, G. C. & Wolfe, K. H. Turning a hobby into a job: how duplicated genes find new

functions. *Nat Rev Genet* **9**, 938–950 (2008).

67. Fares, M. A., Keane, O. M., Toft, C., Carretero-Paulet, L. & Jones, G. W. The roles of whole-

genome and small-scale duplications in the functional specialization of Saccharomyces cerevisiae

genes. *PLoS Genet* **9**, e1003176 (2013).

68. Meier, P., Finch, A. & Evan, G. Apoptosis in development. *Nature* **407**, 796–801 (2000).

69. Degterev, A., Boyce, M. & Yuan, J. A decade of caspases. *Oncogene* **22**, 8543–8567 (2003).

70. Fuchs, Y. & Steller, H. Live to die another way: modes of programmed cell death and the signals

emanating from dying cells. *Nat. Rev. Mol. Cell Biol.* **16**, 329–344 (2015).

71. Mancini, M. *et al.* Caspase-2 Is Localized at the Golgi Complex and Cleaves Golgin-160 during

Apoptosis. *J Cell Biol* **149**, 603–612 (2000).

72. Ouyang, Y. *et al.* Dronc caspase exerts a non-apoptotic function to restrain phospho-Numb-

induced ectopic neuroblast formation in Drosophila. *Development* **138**, 2185–2196 (2011).

73. Huh, J. R., Guo, M. & Hay, B. A. Compensatory proliferation induced by cell death in the

Drosophila wing disc requires activity of the apical cell death caspase Dronc in a nonapoptotic

role. *Curr. Biol.* **14**, 1262–1266 (2004).

74. Kumar, S. & Doumanis, J. The fly caspases. *Cell Death Differ* **7**, 1039–1044 (2000).

75. Vogeler, S., Carboni, S., Li, X. & Joyce, A. Phylogenetic analysis of the caspase family in bivalves:

implications for programmed cell death, immune response and development. *BMC Genomics* **22**,

80 (2021).

76. Pirger, Z., Rácz, B. & Kiss, T. Dopamine-induced programmed cell death is associated with

cytochrome c release and caspase-3 activation in snail salivary gland cells. *Biol. Cell* **101**, 105–116

(2009).

77. Romero, A., Estévez-Calvar, N., Dios, S., Figueras, A. & Novoa, B. New insights into the apoptotic

process in mollusks: characterization of caspase genes in Mytilus galloprovincialis. *PLoS ONE* **6**,

e17003 (2011).

78. Zhang, L., Li, L. & Zhang, G. Gene discovery, comparative analysis and expression profile reveal

the complexity of the Crassostrea gigas apoptosis system. *Dev. Comp. Immunol.* **35**, 603–610

(2011).

79. Yang, B. *et al.* Molecular cloning of two molluscan caspases and gene functional analysis during

Crassostrea angulata (Fujian oyster) larval metamorphosis. *Mol. Biol. Rep.* **42**, 963–975 (2015).

80. Plachetzki, D. C., Pankey, M. S., MacManes, M. D., Lesser, M. P. & Walker, C. W. The Genome of

the Softshell Clam Mya arenaria and the Evolution of Apoptosis. *Genome Biol Evol* **12**, 1681–

1693 (2020).

81. Piquet, B., Shillito, B., Lallier, F. H., Duperron, S. & Andersen, A. C. High rates of apoptosis

visualized in the symbiont-bearing gills of deep-sea Bathymodiolus mussels. *PLoS ONE* **14**,

e0211499 (2019).

82. Sokolova, I. M., Evans, S. & Hughes, F. M. Cadmium-induced apoptosis in oyster hemocytes

involves disturbance of cellular energy balance but no mitochondrial permeability transition. *J.*

Exp. Biol. **207**, 3369–3380 (2004).

83. Sokolova, I. M. Apoptosis in molluscan immune defense. *Invertebrate Survival Journal* **6**, 49–58

(2009).

84. An, H.-K. *et al.* CASP9 (caspase 9) is essential for autophagosome maturation through regulation

of mitochondrial homeostasis. *Autophagy* **16**, 1598–1617 (2020).

85. Madadi, Z., Akbari-Birgani, S., Monfared, P. D. & Mohammadi, S. The non-apoptotic role of

caspase-9 promotes differentiation in leukemic cells. *Biochim Biophys Acta Mol Cell Res* **1866**,

118524 (2019).

86. Hollville, E. & Deshmukh, M. Physiological functions of non-apoptotic caspase activity in the

nervous system. *Semin Cell Dev Biol* **82**, 127–136 (2018).

87. Tran, H. T. *et al.* Caspase-9 has a nonapoptotic function in Xenopus embryonic primitive blood

formation. *J Cell Sci* **130**, 2371–2381 (2017).

88. Tamura, R. *et al.* Starfish Apaf-1 activates effector caspase-3/9 upon apoptosis of aged eggs. *Sci*

Rep **8**, 1611 (2018).

89. Belyi, V. A. *et al.* The origins and evolution of the p53 family of genes. *Cold Spring Harb Perspect*

Biol **2**, a001198 (2010).

90. Lu, W.-J., Amatruda, J. F. & Abrams, J. M. p53 ancestry: gazing through an evolutionary lens. *Nat*

Rev Cancer **9**, 758–762 (2009).

91. Berthelet, J. & Dubrez, L. Regulation of Apoptosis by Inhibitors of Apoptosis (IAPs). *Cells* **2**, 163–

187 (2013).

92. Ribeiro Lopes, M., Parisot, N., Callaerts, P. & Calevro, F. Genetic Diversity of the Apoptotic

Pathway in Insects. in *Evolution, Origin of Life, Concepts and Methods* (ed. Pontarotti, P.) 253–

285 (Springer International Publishing, 2019). doi:10.1007/978-3-030-30363-1_13.

93. Ribeiro Lopes, M. *et al.* Evolutionary novelty in the apoptotic pathway of aphids. *Proc Natl Acad*

Sci U S A **117**, 32545–32556 (2020).

94. Aubrey, B. J. *et al.* Mutant TRP53 exerts a target gene-selective dominant-negative effect to drive

tumor development. *Genes Dev* **32**, 1420–1429 (2018).

95. Aubrey, B. J., Kelly, G. L., Janic, A., Herold, M. J. & Strasser, A. How does p53 induce apoptosis

and how does this relate to p53-mediated tumour suppression? *Cell Death Differ* **25**, 104–113

(2018).

96. Gattiker, A., Gasteiger, E. & Bairoch, A. ScanProsite: a reference implementation of a PROSITE

scanning tool. *Appl Bioinformatics* **1**, 107–108 (2002).

97. Quevillon, E. *et al.* InterProScan: protein domains identifier. *Nucleic Acids Res* **33**, W116-120

(2005).

98. Katoh, K. & Standley, D. M. MAFFT multiple sequence alignment software version 7:

improvements in performance and usability. *Mol. Biol. Evol.* **30**, 772–780 (2013).

99. Sievers, F. *et al.* Fast, scalable generation of high-quality protein multiple sequence alignments

using Clustal Omega. *Mol. Syst. Biol.* **7**, 539 (2011).

100.   HALL, T. A. BioEdit : a user-friendly biological sequence alignment editor and analysis

program for Windows 95/98/NT. *Nucleic Acids Symp. Ser.* **41**, 95–98 (1999).

101.   Castresana, J. Selection of conserved blocks from multiple alignments for their use in

phylogenetic analysis. *Mol. Biol. Evol.* **17**, 540–552 (2000).

102.   Guindon, S. *et al.* New algorithms and methods to estimate maximum-likelihood phylogenies:

assessing the performance of PhyML 3.0. *Syst. Biol.* **59**, 307–321 (2010).

103.   Gouy, M., Guindon, S. & Gascuel, O. SeaView version 4: A multiplatform graphical user

interface for sequence alignment and phylogenetic tree building. *Mol Biol Evol* **27**, 221–224

(2010).

104.   Tamura, K., Stecher, G. & Kumar, S. MEGA11: Molecular Evolutionary Genetics Analysis

Version 11. *Mol Biol Evol* **38**, 3022–3027 (2021).

105.   Ronquist, F. & Huelsenbeck, J. P. MrBayes 3: Bayesian phylogenetic inference under mixed

models. *Bioinformatics* **19**, 1572–1574 (2003).

106.   Nguyen, L.-T., Schmidt, H. A., von Haeseler, A. & Minh, B. Q. IQ-TREE: A Fast and Effective

Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. *Mol Biol Evol* **32**, 268–274

(2015).

107.   Kishino, H. & Hasegawa, M. Evaluation of the maximum likelihood estimate of the

673   evolutionary tree topologies from DNA sequence data, and the branching order in hominoidea. *J*

674   *Mol Evol* **29**, 170–179 (1989).

675   108.   Shimodaira, H. An approximately unbiased test of phylogenetic tree selection. *Syst Biol* **51**,

676   492–508 (2002).

677   109.   Suyama, M., Torrents, D. & Bork, P. PAL2NAL: robust conversion of protein sequence

678   alignments into the corresponding codon alignments. *Nucleic Acids Res* **34**, W609-612 (2006).

Bayesian posterior probabilities 0 to 1.00
- ● ML Bootstrap   >90
- ● ML Bootstrap 70-90
- ○ ML Bootstrap 50-70

690
691
Figure 1:

Topology of metazoan CARD-caspase phylogeny obtained by Bayesian inference using full

sequences alignment. Three strongly supported monophyletic groups are identified: caspase-

9, caspase-2 and a more heterogeneous group named here [Inflammatory Caspases +

Caspase-Y]. Together they form a clade strictly corresponding to bilaterian animals. Caspase-

2 is widely distributed among bilaterians [Deutostomia + Ecdysozoa +

Lophotrochozoa/Spiralia] reflecting an ancestral origin. Conversely, caspase-9 is strictly

restricted to deuterostomian animals. Sequences of non-bilaterians (Cnidaria + Ctenophora +

Placozoa) form divergent paraphyletic groups. The selected outgroup is the unique CARD-

caspase of Porifera *Amphimedon queenslandica*.

Figure 2

Reconstruction of convergent hypothetical intrinsic apoptotic pathways among metazoans

according to molecular actors detected and identified in their genomes. Variability of intrinsic

apoptotic pathways among animals emerged from convergences and recruitment of distinct

actors with independent evolutionary history. Caspase-2 is bilaterian-specific and the initiator

of ecdysozoans intrinsic apoptosis. Caspase-9 is restricted to deuterostomes and the specific

initiator of mammalian intrinsic apoptosis. Deuterostomes exhibit several losses (i.e. caspase

2 in cephalochordates, caspase 9 in urochordates) or duplication (i.e. caspase 9 in

echinoderms), highlighting a putative evolutionary flexibility in apoptotic pathway

establishment. Mitochondrial functions diverge among phyla and cytochrome c release thanks

to mitochondrial outer membrane permeabilisation (MOMP) is specific only to mammals and

possibly echinoderms. The convergent evolutionary histories reflect a probable phylum

specific adaptive process leading to parallel evolution of mitochondrial apoptotic pathways

observed among animals.

Supplementary Figure 1:

Monophyly of each CARD-caspase group is conserved using Bayesian inference and maximum

likelihood analyses. Caspase-9, caspase-2 and [Inflammatory Caspases + Caspase-Y] groups

remain conserved, confirming the bilaterian-specificity of caspase-2 and deuterostomian-

specificity of caspase-9. Relationships among these three clades differ depending on the

methodology employed. Divergent sequences of non-bilaterians animals (Cnidaria +

Ctenophora + Placozoa) always form a paraphyletic group of caspases.

Bayesian posterior probabilities 0 to 1.00

● ML Bootstrap >90
● ML Bootstrap 70-90
○ ML Bootstrap 50-70

Supplementary Figure 2:

Topology of metazoan CARD-caspase phylogeny obtained by Bayesian inference. Selected

outgroup is the unique CARD-caspase of Ctenophora *Pleurobrachia bachei.* The three strongly

supported monophyletic groups identified, caspase-9, caspase-2 and [Inflammatory Caspases

+ Caspase-Y], are the same as when we used the unique CARD-caspase of the Porifera

Amphimedon queenslandica as outgroup. Also, sequences from bilaterian animals remain

monophyletic.

Nodes robustness
Bayesian posterior probabilities 0 to 1.00
- ● ML Bootstrap   >90
- ◉ ML Bootstrap 70-90
- ○ ML Bootstrap 50-70

B.belcheri Casp.9A ←

Supplementary Figure 3:

Phylogeny of CARD-caspases at the deuterostomian scale. Maximum likelihood and Bayesian

inference methods produce similar topologies. Despite the unstable position of

cephalochordate *Branchiostoma belcheri* caspase-9A (green arrow), it unequivocally appears

to belong to the caspase-9 group. Numbers given correspond to posterior probabilities. We

used respectively as outgroups *Capitella teleta* caspase-Y (ELT97848.1) (**A1**), *Aplysia*

760   *californica* caspase-2 (XP_005113266) (**A2**), *Hydra vulgaris* caspase-X2 (NP_001274285.1) (**A3**),

761   and *Caenorhabditis elegans* Ced3 (AAG42045.1) (**A4**).

Bayesian posterior probabilities 0 to 1.00

- ● ML Bootstrap >90
- ● ML Bootstrap 70-90
- ○ ML Bootstrap 50-70

Supplementary Figure 4:

Phylogeny of Apaf-1 at the metazoan scale made with the NB-ARC domain alignment. Two

monophyletic groups are identified: Apaf-1 of ecdysozoans, and apaf-1 of [Deutostomia +

Cnidaria + Placozoa]. Globally, the topology is not consistent with the species phylogeny, and

nodes are poorly supported. Among metazoans, Apaf-1 is characterised by high sequence

divergence and fast independent evolution, making homology hypotheses difficult to establish.

The selected outgroup is Apaf-1 of Porifera *Amphimedon queenslandica*.

Bayesian posterior probabilities 0 to 1.00

● ML Bootstrap   >90
● ML Bootstrap 70-90
○ ML Bootstrap 50-70

Bayesian posterior probabilities 0 to 1.00
- ● ML Bootstrap   >90
- ◉ ML Bootstrap 70-90
- ○ ML Bootstrap 50-70

Supplementary Figure 5:

Topology of Bcl-2 family phylogeny from Bayesian inference and maximum likelihood at the

metazoan scale using outgroups *Amphimedon queenslandica* Bcl-like 1 (XP_003383425.1) (**A**)

and Bcl-like 2 (XP_003387574.1) (**B**). Bcl-2 proteins strictly clustered into five monophyletic

groups: three 'pro-apoptotic' clades (Bok, Bak, Bax) and two less supported 'anti-apoptotic'

groups (Bcl-2/W/XL) and a more complex (Bcl-B/Mcl-1/Bfl-1) clade. Each group includes

bilaterian as well as non-bilaterian sequences which suggests a deep origin of this complex

multigenic family. Clustering into these five groups is consistent between analyses, while

relationships among them is not well resolved.

823   Supplemental table 1: List of CARD-caspases used for phylogenetic analysis.
824

Species		Accession number	Name
Porifera	*Amphimedon queenslandica*	XP_003383519.1	Casp. Card
Cnidaria	*Acropora digitifera*	XP_015766400.1	Casp. X2
		XP_015768208.1	Casp. X1
	Exaiptasia pallida	KXJ12672.1	Casp. X1
		KXJ20965.1	Casp. X2
		KXJ12683.1	Casp. X3
	Hydra vulgaris	ACY95435.1	Casp. X1
		ACY95436.1	Casp. X2
Ctenophora	*Pleurobrachia bachei*	Sb 2658116	Casp. X
Placozoa	*Trochoplax adherens*	RDD44781.1	Casp. X
Lophotrocozoa	*Aplysia californica*	XP_005113266.2	Casp. 2
		XP_012945422.1	Casp. Y
	Biomphalaria glabrata	XP_013082356.1	Casp. 2
	Capitella teleta	ELU00616.1	Casp. 2
		ELT97848.1	Casp. Y
	Crassostrea angulata	AGN75137.1	Casp. 2
	Crassostrea gigas	XP_011419292.2	Casp. 2
		XP_011414267.3	Casp. Y1
		XP_011449817.2	Casp. Y3
		XP_011432762.2	Casp. Y2
	Octopus bimaculoides	XP_014783442.1	Casp. 2
		XP_014790087.1	Casp. Y1
		XP_014784582.1	Casp. Y2
Chelicerata	*Limulus polyphemus*	XP_013788138.1	Casp. 2A

		XP_013776997.1	Casp. 2B
Insecta	*Agrilus planipennis*	XP_018324425.1	Casp. 2
	Apis melifera	XP_016771440.1	Casp. 2
	Drosophila melanogaster	CAB53565.1	Casp. 2
	Habropoda laboriosa	XP_017788772.1	Casp. 2
	Helicoverpa armigera	AEK20835.1	Casp. 2
	Papilio machaon	XP_014362236.1	Casp. 2
	Tribolium castaneum	XP_001813274.1	Casp. 2
Nematoda	*Ascaris suum*	ERG86894.1	Casp. 2
	Caenorhabditis elegans	AAG42045.1	Casp. 2
Vertebrata	*Salmo salar*	XP_014009539.1	Casp. 2
		XP_013991583.1	Casp. 9
	Mus musculus	NP_033937.2	Casp. 1
		EDL13489.1	Casp. 2
		NP_031635.2	Casp. 4
		AAH56447.1	Casp. 9
		AAT91067.1	Casp. 12
	Xenopus laevis	NP_001081223.1	Casp. 1
		NP_001081404.1	Casp. 2
		NP_001079035.1	Casp. 9
Echinodermata	*Strongylocentrotus purpuratus*	XP_789183.3	Casp. 9A
		XP_011661242.1 1	Casp. 9B
		XP_011661359.1	Casp. 9C
Hemichordata	*Saccoglossus kowalevskii*	XP_006811879.1	Casp. 9
Cephalochardata	*Branchiostoma belcheri*	XP_019646757.1	Casp. 2A
		XP_019642903.1	Casp. 2B
		XP_019623612.1	Casp. 9A
		XP_019644208.1	Casp. 9B
	Branchiostoma floridae	EEN68002.1	Casp. 2
Urochordata	*Botryllus schlosseri*	Boschl.CG.Botznik2013.chrUn.g02816.01.p	Casp. 2A1
		Boschl.CG.Botznik2013.chrUn.g08767.01.p	Casp. 2A2
		Boschl.CG.Botznik2013.chrUn.g09831.01.p	Casp. 2B
	Ciona intestinalis	KH2012:KH.C4.463.v1.A.ND1-1	Casp. 2A
	Ciona savignyi	CISAVI-CG-ENS81-R15-461426-463207-09349-P	Casp. 2A
		CISAVI-CG-ENS81-R54-461275-467018-10343-P	Casp. 2B
	Molgula oculata	Moocul-CG-ELv1_2-S113854-g13164-01-p	Casp. 2A
		Moocul-CG-ELv1_2-S103067-g09526-01-p	Casp. 2B
	Phallusia mammillata	PHMAMM-CG-MTP2014-S310-G07060-01-P	Casp. 2A
		PHMAMM-CG-MTP2014-S92-G02989-01-P	Casp. 2B
	Halocynthia roretzi	HARORE-CG-MTP2014-S35-G03390-01-P	Casp. 2A
		HARORE-CG-MTP2014-S130-G07320-01-P	Casp. 2B

827   Supplemental Table 2: Topology tests between Bayesian inference method (BI topology) and
828   maximum likelihood estimation (ML topology). p-KH : p-value of one sided Kishino-Hasegawa

829 test. p-SH : p-value of Shimodaira-Hasegawa test. p-AU : p-value of approximately unbiased
830 (AU) test. In all cases, BI topology is more likelihood than the ML topology.
831

	p-KH	p-SH	p-AU
BI topology	0.714	1	0.702
ML topology	0.286	0.286	0.298

832
833
834
835
836 Supplemental Table 3: Estimation of the recombination (Rho) and substitution (Theta) rates
837 using ProteinEvolverABC on metazoans CARD-caspases alignment. The low rho values
838 indicate an absence of recombination and consequently no domain shuffling.
839

	Rho	Theta
Min.:	0.2762848	225.9003910
2.5% Perc.:	0.6311461	257.7492870
Median:	2.1534998	338.5860425
Mean:	2.0328712	332.1133675
Mode:	2.4251897	353.2953718
97.5% Perc.:	2.6993927	366.5018469
Max.:	2.7559795	367.8813704

848 Supplemental Table 4: List of Apaf-1 among metazoans. N/A: No APAF-1 has been identified.
849

Species		Accession number	Name
Porifera	*Amphimedon queenslandica*	XP_019855714.1	Apaf-1
Ctenophora	*Pleurobrachia bachei*	N/A	
Placozoa	*Trichoplax sp.*	RDD40813.1	Apaf-1
Cnidaria	*Acropora digitifera*	XP_015776356.1	Apaf-1
	Exaiptasia pallida	KXJ17575.1	Apaf-1
	Hydra vulgaris	CDG72123.1	Apaf-1
	Nematostella vectensis	XP_032220711.2	Apaf-1
		XP_032223290.2	Apaf-1 like1
		XP_001626725.1	Apaf-1 like2
		XP_032225984.2	Apaf-1 like3
Nematoda	*Caenorhabditis elegans*	CAA48781.1	Ced-4
	Caenorhabditis briggsae	XP_002642027.2	Ced-4
Insecta	*Aedes aegypti*	EAT48066.2	Dark
	Agrilus planipennis	XP_025834330.1	Dark
	Drosophila melanogaster	NP_725637.1	Dark

	Drosophila pseudoobscura	XP_001360832.3	Dark
	Habropoda laboriosa	XP_017794472.1	Dark
	Helicoverpa armigera	XP_021181657.1	Dark
	Papilio machaon	XP_014359375.1	Dark
	Tribolium castaneum	XP_015840766.1	Dark
Lophotrocozoa	Aplysia californica	N/A	
	Biomphalaria glabrata	N/A	
	Crassostrea angulata	N/A	
	Crassostrea gigas	N/A	
	Octopus bimaculoides	N/A	
	Capitella teleta	N/A	
Vertebrata	Homo sapiens	XP_047284715.1	Apaf-1
	Mus musculus	NP_033814.2	Apaf-1
	Canis lupus familiaris	XP_038544278.1	Apaf-1
	Gallus gallus	XP_040518239.1	Apaf-1
	Xenopus laevis	NP_001085834.1	Apaf-1
	Takifugu rubripes	XP_011611631.2	Apaf-1
	Tetraodon nigroviridis	CAG01878.1	Apaf-1
Echinodermata	Strongylocentrotus purpuratus	XP_030844937.1	Apaf-1
		XP_030854919.1	Apaf-1 like1
		XP_030828596.1	Apaf-1 like2
Hemichordata	Saccoglossus kowalevskii	XP_006818297.1	Apaf-1
Cephalochordata	Branchiostoma belcheri	XP_019621685.1	Apaf-1
	Branchiostoma floridae	XP_035681983.1	Apaf-1-1
		XP_035678916.1	Apaf-1-2
		XP_035672288.1	Apaf-1 like1
		XP_035690210.1	Apaf-1 like2
		XP_035667061.1	Apaf-1 like3
		XP_035696241.1	Apaf-1 like4
		XP_035693308.1	Apaf-1 like5
		XP_035673101.1	Apaf-1 like6
		XP_035692718.1	Apaf-1 like7
Urochordata	Botryllus schlosseri	N/A	
	Ciona intestinalis	N/A	
	Ciona savignyi	N/A	
	Molgula oculata	N/A	
	Phallusia mammillata	N/A	
	Halocynthia roretzi	N/A	

850
851
852

855   Supplemental table 5: List of Bcl-2 used for phylogenetic analysis.

	Species	Accession number	Names
Porifera	*Amphimedon queenslandica*	XP_003383425.1	Bcl-2 like 1
		XP_003387574.1	Bcl-2 like 2
Cnidaria	*Hydra vulgaris*	EF104645.1	Bak 1
		EU035760.1	Bak 2
		XP_012562061.1	Bax
		EU035764.1	Bcl-2 like 1
		EF104646.1	Bcl-2 like 2
		EU035765.1	Bcl-2 like 3
		EU035763.1	Bcl-2 like 4
		EU035761.1	Bok
		EF104647.1	Mcl-1 1
		EU035762	Mcl-1 2
Nematoda	*Caenorhabditis elegans*	AAA20080.1	Ced-9
Insecta	*Drosophila melanogaster*	AAF44120.1	Buffy (Bok-1)
		AAF26289.1	Buffy (Bok-2)
Lophotrocozoa	*Biomphalaria glabrata*	XP_013085524.1	Bax 1
		XP_013096338_1	Bax 2
		XP_013086802.1	Bax 3
		XP_013070177.1	Bak
		XP_013081872.1	Bcl-2 like 1
		XP_013093137.1	Bcl-2 like 2
		XP_013068612.1	Bcl-2 like 3
		XP_013083436.1	Bcl-2 like 4
		XP_013069706.1	Bok
		XP_013065969.1	Mcl-1
	Crassostrea gigas	XP_011424481.1	Bax
		XP_011439700.1	Bak
		XP_011449013.1	Bcl-2 like 1
		ACH42081.1	Bcl-2 like 2
		EKC18663.1	Bcl-2 like 3
		XP_011436990.1	Mcl-1 1
		EKC40007.1	Mcl-1 2
Vertebrata	*Homo sapiens*	AAA74466.1	Bak
		NP_001278357.1	Bax
		API71171.1	Bcl-2
		AAK48715.1	Bcl-B
		AAB09055.1	Bcl-w
		CAA80661.1	Bcl-xL
		NP_004040.1	Bfl-1
		NP_115904.1	Bok
		AAF64255.1	Mcl-1
	Mus musculus	NP_031549.2	Bak
		NP_031553.1	Bax
		AAH95964.1	Bcl-2
		Q9Z0F3.1	Bcl-B

		AAA51039.1	Bcl-xL
		AAB09056.1	Bcl-w
		Q07440.1	Bfl-1
		NP_058058.1	Bok
		NP_032588.1	Mcl-1
	Xenopus laevis	NP_001089587.1	Bak
		AAR84081.1	Bax
		BAH28834.1	Bcl-2
		AAI10791.1	Bcl-xl
		XP_018089640.1	Bcl-w
		NP_001139563.1	Bok
		ACI47310.1	Mcl-1
Urochordata	*Ciona intestinalis*	KH2012:KH.C4.794.v1.A.SL1-1	Bax
		KH2012:KH.S653.2.v2.A.SL1-1	Bcl-xl
		KH2012:KH.L87.39.v1.A.ND1-1	Bok
	Molgula oculata	Moocul.CG.ELv1_2.S96550.g08147.01.p	Bax
		Moocul.CG.ELv1_2.S112899.g12639.01.p	Bok

857
858
859
860