Using antibody synergy to engineer a
high potency biologic cocktail against *C. difficile*

Hui Zhao[1], Michael Tasch[1], Michael Dodds[1], Mesfin Gewe[1], Anissa Martinez[1], Melanie Hutton[2], Kristie Keeney[1], Alex Pollock[1], Ben Jester[1], Nhi Khuong[1], Mia Zhang[1], Colin Brady[1], Mark Heinnickel[1], Hannah Tabakh[1], Nathan Sanjaya[1], Kendra Cruickshank[1], Troy Paddock[1], Stacey Ertel[1], Sarah Struyvenberg[1], Jason Dang[1], Tracy Saveria[1], Chelsea Shanitta[1], David Fletcher[1], Lauren Goetsch[1], Caitlin Gamble[1], Steven J. Mileto[2], Ryan Heselpoth[3], Dena Lyras[2], Craig A. Behnke[1], Vincent Fischetti[3], Brian Finrow[1], and James M. Roberts[1,4]

[1]Lumen Bioscience, Seattle, WA
[2]Biomedicine Discovery Institute and Department of Microbiology, Monash University, Victoria, Australia
[3]Laboratory of Bacterial Pathogenesis and Immunity, Rockefeller University, New York, NY
[4]Corresponding author: James M. Roberts, Lumen Bioscience, 1441 N. 34th Street, Seattle, WA, 98103, jroberts@lumen.bio

Keywords: *Arthrospira platensis*; spirulina; *C. difficile*; VHH; lysin; oral therapeutic; synergy

ABSTRACT

We applied a mathematical framework originally used to model the effects of multiple

inhibitors on enzyme activity to guide the development a therapeutic antibody cocktail, LMN-

201, to prevent and treat *C. difficile* infection (CDI). CDI causes hundreds of thousands of cases

of severe, often recurrent diarrhea and colitis in the United States annually and is associated

with significant morbidity and mortality worldwide. Current therapies for preventing recurrent

CDI are only partially successful, and there are no options available to prevent initial bouts of

CDI in at-risk populations. Almost all antibody therapies have been developed and administered

as monotherapies. Antibody cocktails are relatively rare even though they have the potential to

greatly increase efficacy. One reason for this is our limited understanding of how antibody

interactions can enhance potency, which makes it difficult to identify and develop antibodies

that can be assembled into optimally effective cocktails. In contrast to the view that antibody

synergies depend on unusual instances of cooperativity or allostery, we show that synergistic

efficacy requires nothing more than that the antibodies bind independently to distinct epitopes

on a common target. Therefore, synergy may be achieved much more readily than is generally

appreciated. Due to synergy the LMN-201 antibody cocktail, which targets the *C. difficile*

exotoxin B (TcdB), is 300- to 3000-fold more potent at neutralizing the most clinically prevalent

TcdB toxin types than bezlotoxumab, the only monoclonal antibody currently approved for

treatment or prevention of CDI. The efficacy of LMN-201 is further enhanced by inclusion of a

phage-derived endolysin that destroys the C. difficile bacterium, and which therefore has a

complementary mechanism of action to the antibody cocktail. These observations may serve as

a paradigm for the development of high potency biologic cocktails against targets that have

proven challenging for single-agent therapies.

INTRODUCTION

Clostridioides difficile is a Gram-positive spore-forming anaerobic bacterium[1]. *C. difficile*

infection (CDI) causes severe, recurrent diarrhea and colitis and is associated with significant

morbidity and mortality world-wide[2]. According to the U.S. Centers for Disease Control and

Prevention, *C. difficile* is the most common cause of health care-associated infections in U.S

hospitals, where nearly half a million CDI cases occur each year[3]. Though traditionally

considered a nosocomial infection, community-associated CDI is on the rise and now comprises

most cases in some regions[4]. Treatment costs are estimated to exceed $5 billion a year, but the

true economic burden of the disease is far higher when lost quality-adjusted life years are

considered[4].

C. difficile virulence is mediated by expression of two homologous, multidomain exotoxins,

TcdA and TcdB[5–8]. The toxins initially engage cell surface receptors on intestine epithelial cells,

which results in internalization by endocytosis[9,10]. Acidification of internalized endosomes

triggers a conformational change in the toxin that promotes its translocation into the

cytoplasm[11]. Cytoplasmic toxin undergoes autoproteolysis[12], releasing its N-terminal

glucosyltransferase domain that enzymatically inhibits Rho family-GTPases by mono-

glucosylation[13,14]. Inactivation of Rho disaggregates the cellular actin cytoskeleton thereby

disrupting the colonic epithelium[15–17], leading to severe inflammation and watery diarrhea

characteristic of CDI. The relative importance of TcdA and TcdB for causing *C. difficile* pathology

varies by host species, and current balance of opinion is that human pathophysiology is

primarily caused by TcdB[7,18–23].

Antibiotic therapy is often successful for initial cases of CDI. However, approximately 20%-40%

of CDI patients will have a recurrent episode[24], and the chance of additional episodes in those

patients exceeds 40%[25,26]. Microbial dysbiosis is a consequence of the antibiotics most often

used to treat CDI, which creates a niche favorable to *C. difficile* proliferation[27] and promotes

recurrent disease. Due to side effects and selection for antimicrobial resistance, antibiotics are

no longer used prophylactically.

One alternative to antibiotics for treatment of microbial pathogens is passive immunization by

direct infusion of neutralizing antibodies. Intravenous or intraperitoneal infusion of monoclonal

and polyclonal antibodies targeting TcdA and TcdB are protective against primary infection and

relapse in *C. difficile* challenge models in mice[8,21,28,29], hamsters,[30,31] piglets[22,32]. It is important

that immunotherapies used clinically address the natural antigenic variation displayed by

TcdB[33]. Bezlotoxumab, a monoclonal antibody against TcdB, has been adopted for clinical use in

conjunction with antibiotics for high-risk patients[23,34]. However, it poorly neutralizes the

prevalent TcdB2 variant, and clinically it is only partially effective in reducing recurrent CDI in

humans. Oral administration of polyclonal antibodies from cows hyperimmunized with TcdA

and TcdB has likewise been evaluated in human trials with reported success but has not

progressed to commercial development[35–37]. Most recently, fecal microbiota transplantation

(FMT) has also proven effective in reducing recurrence of CDI, presumably by helping to correct

the intestinal dysbiosis that predisposes to relapsing disease[38]. Despite these advances there

currently are no clinically approved therapeutics to prevent primary CDI. There are no FDA

approved CDI vaccines[39], and the most recent Phase 3 clinical trial of an anti-toxin vaccine was

terminated due to futility[40].

We report here the preclinical development of a four-component therapeutic cocktail to both

prevent and treat CDI. The cocktail is comprised of homodimeric versions of three toxin-binding

proteins (5D, E3, and 7F) derived from camelid single-domain antibody fragments known as

VHHs or nanobodies. Each protein was previously shown to inactivate TcdB[8,28,29,41]. Binding of

5D to TcdB prevents its pH-driven conformational change that enables pore formation and

transit into the cytoplasm from internalised endosomes. E3 binds to the glucosyltransferase

domain (GTD) of TcdB, interfering with the autoproteolysis step necessary for enzyme

activation. 7F also binds to the GTD, inhibiting catalytic glucosylation/inactivation of Rho-family

GTPases by the mature N-terminal glucosyltransferase. The fourth cocktail component is a

C. difficile-specific endolysin[42]. These proteins are highly specific, phage-encoded antimicrobial

lytic enzymes that, due their specificity and rapid effect, hold considerable promise for treating

bacterial infectious diseases, including CDI[42–44]. We showed that the selected combination of

three dimeric VHHs had a synergistic neutralization potency that vastly exceeded the potency

of any single dimeric VHH and had far greater potency and broader strain coverage than

bezlotoxumab[23,34]. We further demonstrated that the therapeutic was enhanced by combining

the dimeric VHHs that neutralize TcdB with an endolysin that destroys the *C difficile* bacterium.

LMN-201 was manufactured using the edible microorganism spirulina (*Arthrospira platensis*) as

the expression platform. It was delivered orally as a powder of whole, unrefined spirulina

biomass[45], and is easily scalable, shelf-stable, simple to administer, and broadly neutralizing

against all common TcdB types.

RESULTS

Spirulina strain engineering

The *C. difficile* therapeutic LMN-201 contained four therapeutic proteins: three homodimeric

fusion proteins designated pp1005, pp1006, and pp1007, which incorporated homodimeric

derivatives of VHHs 5D, E3 or 7F, respectively; and fourthly, the catalytic domain of the

bacteriophage-derived PlyCD endolysin, designated pp1092.

The primary structures of each monomer subunit in the VHH homodimers were similar. From

amino to carboxy terminus this was: the maltose-binding protein (MBP) chaperone (366 amino

acids)[46], a Gly-Gly-Gly-Gly tetrapeptide linker, a 5HVZ dimerization domain (49 amino acids), a

Gly-Ser-Gly-Ser-Gly-Gly peptide linker, a TcdB-binding VHH (111-127 amino acids), a Gly-Ser-Gly

tripeptide linker and a hexahistidine tag (543-562 amino acids total) (Figure 1A). The 5HVZ

dimerization domain was derived from the mammalian cAMP-dependent protein kinase A

(PKA) holoenzyme[47,48].

The endolysin PlyCD has been previously demonstrated to be *C. difficile* specific[42-44]. Full-length

PlyCD (pp1093) its catalytic domain alone, $PlyCD_{1-174}$ (pp1092), were expressed in spirulina

strains sp1286 and SP1287, respectively, with an N-terminal 6x his-tag (Figure 1A).

Genes encoding pp1005, pp1006, pp1007, pp1093 and pp1092 were engineered into the

spirulina chromosome by homologous recombination (Figure 1B)[45]. The final strains—SP1308,

SP1312, SP1313, SP1286 and SP1287, respectively—contained the gene of interest integrated

into the spirulina chromosome at a single, pre-selected location and no other exogenous

genetic information. Clonal spirulina filaments were then isolated by micromanipulation,

verified as being homozygous with respect to the transgene, and used to create a frozen strain

bank (Figure 1B, C)[45].

Characterization of spirulina-expressed anti-TcdB VHHs: expression and binding kinetics

Expression levels of all 5 candidate therapeutic proteins in spirulina were quantified using a

capillary electrophoresis-based immunoassay (Figure 2A). After growth in a bioreactor, dried

whole SP1308 biomass was comprised of 0.5% by weight pp1005; SP1312 was comprised of

2.3% pp1006; SP1313 was comprised of 3.4 % pp1007; and SP1287 was comprised of 0.5%

pp1092. At least 90% of each of these proteins was soluble following a one-hour extraction in

an aqueous buffer at neutral pH. The dimeric state of the VHHs on the 5HVZ scaffold was

confirmed by comparing their relative mobilities during size-exclusion chromatography, and by

capillary electrophoresis in reducing versus non-reducing conditions (Figure 2B). Intra-scaffold

disulfide bonds were present in the extracted proteins (Figure 2B) but were dispensable for

stable assembly of the 5HVZ-VHH dimers (Figure 2C).

Binding kinetics were measured by biolayer interferometry using recombinant subdomains of

TcdB that corresponded to the specific antigen-binding sites of each VHH homodimer

(Figure 3A, B). The apparent K_D's were in the mid- to low-picomolar range. The dimeric versions

of all three proteins displayed substantially greater apparent avidity for TcdB than the

corresponding monomeric versions (Figure 3B).

Characterization of TcdB's effect on cell adhesion

Disruption of the actin cytoskeleton by TcdB causes changes in cell shape and cell attachment.

In vitro, these cytoskeletal changes can be measured in real-time as a change in impedance

using cells cultured on a surface embedded with microelectrodes[49,50]. TcdB1 is the TcdB toxin

type most frequently present in worldwide *C. difficile* isolates from infected patients[51]. The

source of TcdB1 for these analyses was ribotype (RT) 087 (type strain 10463). The concentration

range of TcdB1 tested spanned the amounts reported in human patients with *C. difficile*

infection[52]. Data describing TcdB1 toxicity were collected and fit to a four-parameter sigmoid

using nonlinear regression to minimize the sums of residuals squared:

159
$$Adhesion = Top - (Top - Bottom) \cdot \frac{T^H}{T^H + IC_{50}^{\ H}}$$

where Adhesion indicates the results of the above cell attachment assay; Top and Bottom are

the highest (no TcdB1) and lowest (maximal TcdB1) assay values; T is the concentration of

TcdB1; IC_{50} is the concentration of TcdB1 eliciting half maximal inhibition in Adhesion; H is the

steepness of the concentration-response relationship.

The concentration of half-maximal inhibition of cell adhesion by TcdB1 was 1.268 fM with a Hill

slope coefficient of 1.495 (Table 1; Figure 4).

Synergistic Inhibition of TcdB by anti-TcdB VHH homodimers

Data were collected to evaluate inhibition of TcdB toxicity by each VHH homodimer, and all

two- and three-way combinations (Figure 4A; Supplemental Figure 1). Individually, each of the

dimeric VHHs had sub-nanomolar EC_{50} values for TcdB neutralization (Table 1). However,

compared to the single VHHs, the cocktail of the three dimeric VHHs showed remarkable

synergy, providing robust neutralization across the very wide range of TcdB concentrations

found in patients with CDI[52] (Figure 4; Supplemental Table 2). To illustrate, a concentration of

1.5 nM of any one of the three dimeric VHHs shifted the IC_{50} of TcdB1 10- to 60-fold, whereas a

1.5 nM cocktail of all three combined (0.5 nM each) shifted the IC_{50} of TcdB1 almost 3,000-fold.

For TcdB2 the cocktail synergy was similarly striking. At a concentration of 1.5 nM the individual

dimeric VHHs shifted the IC_{50} of TcdB2 10- to 150-fold, while a 1.5 nM cocktail of the three

dimeric VHHs (0.5 nM each) shifted the IC_{50} of TcdB2 more than 16,000-fold.

Surprisingly, the dramatic synergy of antibody combinations requires nothing more than that

they bind independently to distinct epitopes on the target. Quantitative modeling using

measured K_D's and the effects of dimeric VHH mixtures across a large range of TcdB

concentrations indicated that this was likely to be the entire explanation for the synergy we

observed (Supplemental Text and Supplemental Figures 1,2,3). This suggested that synergy

might be much more easily achieved than is generally appreciated, which has important

implications for the development of therapeutic antibody cocktails (see Discussion).

Broad coverage of TdcB toxin-types

Mansfield et al.[33] used *C. difficile* genomic sequences to define 12 distinct TcdB groups that

differed from each other by 3% or greater. 96% of the TcdB genes present in that dataset were

defined as belonging to groups 1, 2, 3, and 5. Group 1 is the most clinically prevalent group

worldwide, and group 2 is expressed by *C. difficile* ribotype 027, the single most prevalent strain

type in the United States, and second most prevalent in Europe[3,51,53,54].

The high degree of conservation of the pp1005, pp1006, and pp1007 binding sites[41] suggested

that each of the dimeric VHHs would neutralize most, if not all, of the four prevalent TcdB toxin

types, except for neutralization of TcdB group 5 by pp1005. We measured the IC_{50} values of

TcdB types 1.1, 2.1, 3.1, and 5.1 alone and when incubated in 1.5 nM pp1005, pp1006, pp1007,

or a cocktail comprised of all three VHH homodimers (0.5 nM of each) and compared these

with the neutralizing potency of 3 nM bezlotoxumab (Figure 4). Each VHH homodimer exhibited

strong neutralizing activity against all tested toxin types except pp1005, which was only weakly

active against toxin type 5. The triple VHH cocktail synergistically neutralized all tested toxin

types, with apparent relative potencies of against each toxin type of 2>1>3>5. Compared to the

triple VHH cocktail, bezlotoxumab was significantly less potent at neutralizing all tested toxin

types, and its neutralizing activity against toxin type 2 (ribotype 027) was notably weak.

Characterization of spirulina-expressed anti-*C. difficile* lysin

PlyCD, a prophage lysin, was found to have a high degree of specificity for degrading *C. difficile*

peptidoglycan[42]. The catalytic domain of this protein, $PlyCD_{1-174}$, efficiently lysed various strains

of *C. difficile* while leaving commensal bacteria unaffected, with rare exceptions that included

B. subtilis and *C. sordellii*.

Genes encoding pp1093 (full-length lysin PlyCD) and pp1092 ($PlyCD_{1-174}$) were engineered into

spirulina to produce strains SP1286 and SP1287, respectively. Expression of both proteins in

spirulina was quantified using a capillary electrophoresis-based immunoassay. After growth in a

bioreactor pp1093 comprised 0.1% of dry biomass and was approximately 50% soluble

following aqueous extraction. pp1092 comprised 0.5% of total SP1287 biomass, and after

aqueous extraction greater than 90% of pp1092 was soluble (Figure 5A). Following extraction

from SP1286 and SP1287 biomass the pp1093 and pp1092 proteins were found to be

monomeric by size-exclusion chromatography and non-reducing PAGE (Figure 5B). However,

pp1093 exhibited notable size heterogeneity, with lower molecular size species.

The lytic activities of pp1093 and pp1092 were compared in a turbidity reduction assay using

extracts from SP1286 and SP1287, and *B. subtilis* as the target bacterium. In these spirulina cell

extracts, the specific activity of pp1092 was 8-fold greater than pp1093 (Figure 5C). The

reduced lysin activity of the pp1093 compared to pp1092 was also observed using *C. difficile*

strain ATCC 43255 as the target bacterium (Supplemental Figure 4). The lytic activities of

pp1093 and pp1092 isolated from spirulina were not significantly different from the same

proteins isolated from *E. coli.* (Supplemental Figure 4). SP1287 (expressing pp1092) was

selected for inclusion in LMN-201 based upon its superior expression level, homogeneity and

bioactivity compared to SP1286. As a control for interference, it was shown that pp1092 did not

affect toxin neutralization by the anti-TcdB VHH cocktail (Figure 4).

Stabilities of anti-TcdB VHHs and anti-*C. difficile* lysin

The four therapeutic proteins within LMN-201 all demonstrated resistance to three common

intestinal proteases: trypsin, elastase, and chymotrypsin (Figures 6A, 7). When exposed to

conditions simulating constitutive levels of these proteases for one hour, pp1005, pp1006, and

pp1007 retained nearly 100% ELISA binding activity compared to undigested protein. At a

higher concentration of protease that simulated the induced protease amounts following a

meal, pp1005 and pp1006 exhibited some sensitivity to digestion with 0.1 mg/mL trypsin but

still retained 63% and 75% of their antigen-binding activities, respectively. SDS-PAGE and

western blotting demonstrated that chymotrypsin cleaved the MBP domain of these dimeric

toxin-binding proteins within 30 minutes, leaving the VHH-5HVZ dimer intact. This was

consistent with the observation that chymotrypsin did not reduce binding affinity as measured

by ELISA, and removal of the MBP moiety may explain the paradoxically increased ability to

detect bound VHH in an ELISA after chymotrypsin digestion. To confirm that digestion did not

affect bioactivity, chymotrypsin-digested pp1005, pp1006, and p1007 were also evaluated in a

cell-adhesion assay. No significant changes in toxin neutralizing activity were observed after

exposure to chymotrypsin for at least one hour (Figure 6B).

Protease treated pp1092 was analyzed for structural integrity by SDS-PAGE (Figure 7A), and for

bioactivity in a turbidity reduction assay using intact *B. subtilis* as the substrate (Figure 7B).

pp1092 showed no detectable change in molecular size (25kDa) following digestion with any of

the three proteases. In contrast, full length PlyCD (37kDa) was almost 100% cleaved to a smaller

25 kDa product by all three proteases, consistent with release of the smaller, protease-stable

catalytic subdomain. Further, pp1092 retained 100% bioactivity following digestion with any of

the three proteases at the lower level, and greater than 70% bioactivity following digestion with

higher (induced) protease amounts.

***In vivo* efficacy**

Efficacies of cocktail therapeutics containing anti-TcdB VHH homodimers and anti-*C. difficile*

lysins were evaluated in mouse (Figure 8) and hamster (Figure 9) pathogen challenge models.

Cohorts of male C57BL/6J mice were infected with *C. difficile* spores 24 hours after

administration of one of the following treatments by oral gavage: vancomycin (positive

control); wild-type spirulina (negative control); Cocktail 1; or Cocktail 2. Cocktail 1 was

comprised of equal proportions of spirulina strains SP1308, SP1312, and SP1313 (expressing

pp1005, pp1006, and pp1007, respectively) and SP1286 (expressing pp1093). Cocktail 2 was the

same but omitted SP1286. Each treatment was then administered once daily following

C. difficile challenge.

The largest number of *C. difficile* spores was shed by mice treated with wild-type spirulina

(Figure 8A). Treatment with Cocktail 1 significantly reduced fecal spore shedding at 24 hours

post infection and at the time of euthanasia ($10^{2.206}$ and $10^{1.867}$ cfu/g lower, respectively), than

wild-type spirulina controls and significant by t-test with p=0.0175 and p<0.001 (Supplemental

Table 3). Fewer spores were observed shed by animals treated with only the toxin-neutralizing

antibodies (Cocktail 2), but the difference did not reach statistical significance at either

timepoint.

The largest maximum weight loss was observed in the negative control group, mice treated

with wild-type spirulina, 3.922 g (Figure 8B; Supplemental Table 4). Treatment with either

spirulina cocktail or vancomycin all reduced the maximum weight loss significantly. On average,

the maximum weight lost by an individual treated with Cocktail 1 was 2.057 g less than the

group treated with wild-type spirulina and significant by t-test with p<0.01. The maximum

weight lost by subjects treated with Cocktail 2 was 2.102 g, a result significant by t-test with

p<0.001.

Treatment with the spirulina therapeutic Cocktails 1 and 2 also increased survival rates, with

borderline significance possibly due to small sample size (Figure 8C). None of the mice treated

with Cocktail 1 perished due to *C. difficile*-related causes (4/4, 100% survival) which was

equivalent to the vancomycin-treated group (10/10, 100% survival). The increased survival rate

was borderline significant relative to the wild-type spirulina-treated group by log-rank test with

p=0.079. The group treated with Cocktail 2 also exhibited higher survival rates than the

negative control (6/7, 87% compared to 2/5, 40%) with borderline significance by log-rank test

with p=0.14. The mouse efficacy study suggests that a spirulina anti-TcdB cocktail

supplemented with the lysin is effective at reducing bacterial load and improving clinical

outcomes *in vivo*.

The Golden Syrian Hamster model of CDI has been used previously for evaluating orally

delivered antibodies against *C. difficile* toxins[30]. The experimental groups we used were

Cohort 1: no treatment, Cohort 2: wild-type spirulina, Cohort 3: anti-TcdB VHHs (3) + Anti-TcdA

VHHs (2) + lysin PlyCD, all delivered within whole spray-dried spirulina biomass. Each treatment

was administered twice daily by oral gavage, from days -1 through day 12. It was necessary to

include anti-TcdA VHHs in this protocol because TcdA contributes significantly to disease in

hamsters (unlike in humans)[8]. We observed that the therapeutic protein cocktail significantly

extended survival time compared to the wild-type spirulina control (p=0.015) (Figure 9A), and

significantly reduced *C. difficile* colonization of the cecum by more than an order of magnitude

when compared to untreated or wild-type treated groups (p=0.046 and p=0.0041, respectively)

(Figure 9B).

DISCUSSION

We report here the first use of quantitative analyses of drug synergy for preclinical and clinical

development of a therapeutic antibody cocktail. These analyses build on a general framework

for evaluating the effects of multiple drugs acting on a single target that was initially developed

to model enzyme inhibition[55]. In that context it was shown that mutually exclusive inhibitors

(i.e., competitive) have an additive inhibitory effect, whereas mutually nonexclusive inhibitors

(i.e., independent) have a multiplicative (i.e., synergistic) effect. This framework can be applied

more generally, for example to antibody inhibitors of a protein[56], and has the potential to

inform decisions around drug cocktail formulation, including selection of lead candidates and

optimization of their relative concentrations.

Synergy is usually not considered in the development of cocktail therapeutics. To some degree

this is due to the costs and complexities associated with developing drug combinations, but it is

also because such synergies are often thought to be rare, arising from cooperative or unusual

allosteric effects[57]. A small number of anti-viral therapeutics have been intentionally developed

as antibody cocktails[58,59], but this was done to reduce the likelihood of pathogen escape.

Moreover, if drug concentrations are only evaluated within a 2- or 3-fold range of their

respective EC_{50} values, synergy can be difficult to distinguish from simple additivity, and is

therefore easily overlooked. Accordingly, almost all clinically approved antibody cocktails were

first developed as solo therapeutics, and subsequently only a small subset have been

empirically evaluated in combination, often without regard to understanding whether

underlying synergies might have contributed to improved efficacy.

In contrast to the prevailing view that synergy is rare and elusive we have demonstrated here,

using the specific example of anti-*C. difficile* toxin antibodies, that synergy can simply be the

result of the laws of mass action governing the independent interactions of drugs with a

common target, and therefore may be more readily achieved than is generally thought.

Moreover, the synergistic efficacy of independently acting drugs increases nonlinearly with

concentration and can be enormous when concentrations substantially exceed the EC_{50}. We

emphasize that this is not a complex function of allostery or cooperativity, but a simple

consequence of independent Michaelis-Menten binding. This has general and important

implications for therapeutic development. Synergistically acting antibodies will be particularly

useful for treating a disease like CDI, where single antibody therapeutics struggle to achieve the

very high level of target inhibition required. That the cocktail also offers broader strain

coverage and protection against escape variants adds to its appeal.

Beyond infectious diseases, antibody synergy may also be important for treating diseases with

complex etiologies where single-agent interventions have only been moderately successful, for

example inflammatory diseases of the gastrointestinal tract and cardiometabolic disease. The

results here offer antibody drug developers a paradigm for developing high potency cocktail-

based interventions against diseases refractory to single-antibody approaches.

CDI is an appealing initial target for clinically evaluating these principles, because current

therapies for preventing recurrent CDI are only partially successful, and there are no options

available to prevent initial bouts of CDI in at-risk populations. LMN-201 is a spirulina-expressed

and orally delivered cocktail of four therapeutic proteins that target both the *C. difficile*

bacterium and the key driver of its virulence, exotoxin TcdB. Orally delivered protein

therapeutics may offer significant advantages over their injected counterparts for treatment

and prevention of recurrent CDI. Oral administration can deliver high amounts of therapeutic

proteins directly to the intestine epithelium, whereas systemically delivered antibodies have

poor partitioning coefficients from the vasculature to the gut lumen[60]. Moreover, the risk of off-

target toxicity is reduced: the therapeutic proteins within LMN-201 are too large for intact

systemic absorption[45]. At least three prior clinical trials have reported safe and efficacious oral

administration of bovine antibodies against *C. difficile* toxins derived from milk or colostrum[35–

37], supporting the feasibility of this approach.

The harsh environment of the gastrointestinal tract is considered challenging for orally

delivered protein therapeutics. LMN-201 addresses this challenge in several ways. First, the

therapeutic proteins in LMN-201 showed significant resistance to the major GI proteases.

Second, for human administration the product is delivered orally in polymer-coated capsules

designed to release the drug product in the ileum. Third, spirulina-based cGMP manufacturing

is highly scalable, making feasible cost-effective administration of the drug product throughout

the period of high risk for CDI recurrence, even if the harsh GI environment necessitates daily or

twice daily administration to maintain therapeutic levels at the site of action[42].

Perhaps of greatest significance is the extraordinary potency of LMN-201: each dimeric VHH

within LMN-201 had a sub-nanomolar EC_{50} for TcdB inhibition, and this was substantially

enhanced by cocktail formulation. An oral dose of 1,500 mg of spirulina biomass (500 mg of

each VHH-spirulina strain) will deliver to the intestine a combined dimeric VHH concentration of

approximately 2.6 µM, which is more than 1,000-fold higher than minimally required[61,62].

Pharmacokinetic data from a Phase 1 trial of LMN-201 confirmed that the amount of bioactive

VHH recovered in the distal ileum following oral administration was more than sufficient to

neutralize TcdB-levels previously reported in patients at risk for CDI recurrence (manuscript in

preparation).

The fourth LMN-201 component—a bacteriophage-derived endolysin—not only broadens

C. difficile strain coverage but also enhances efficacy. While TcdB is the key virulence factor in

CDI, approximately 5% of patients are infected by *C. difficile* strains that secrete TcdA but not

TcdB[52]. Other virulence factors are also associated with CDI, including the *C. difficile* binary

toxin (CDT), which is associated with increased 30-day mortality[63]. Inclusion of the lysin may

offer protection against these non-canonical *C. difficile* strains, as well as novel strains with new

toxin variants that might emerge in the future.

The lysin may also increase therapeutic potency in at least two ways. First, clinical performance

is augmented by the lysin's previously demonstrated synergy with standard-of-care

vancomycin, which act in concert to disrupt *C. difficile* cell wall structure[42]. Second, these

agents independently intervene at two steps in the pathway that controls virulence—*C. difficile*

proliferation and toxin action on the gastrointestinal epithelium. The lysin thus complements

the VHH homodimers with a potential for orthogonal synergy to enhance LMN-201's clinical

real-world effectiveness.

MATERIALS AND METHODS

Engineering of spirulina

Markerless spirulina strains were generated as described[45]. In brief, the natural spirulina gene

WP_006618409.1, which confers resistance to kanamycin, was replaced with an exogenous

gene encoding resistance to streptomycin to generate spirulina strain SP205. SP205 was then

transformed with constructs containing the gene of interest, WP_006618409.1, as well as the

homology arms. The transformants were selected with kanamycin. Single filaments were

isolated and propagated to generate cell banks. Cells were then thawed from the cell banks and

DNA was extracted to be used to confirm complete segregation and verify sequence of the

WP_006618409.1 via PCR and Sanger Sequencing.

Illumina sequencing of spirulina

Spirulina genomic DNA was extracted using a Blood and Tissue Kit (Qiagen, 69504) with minor

modifications. Spirulina biomass was resuspended in buffer ATL. It was digested with

proteinase K (2 mg/mL) overnight at 56 °C and then treated with RNase A (10 mg/mL) for

20 minutes at 56 °C. 400 µL of buffer AL was added and the biomass was pelleted for 10

minutes at 13,000g. The supernatant was then transferred to a fresh tube with 250 µL of 100%

ethanol, mixed and applied to DNeasy columns. The samples were washed once with 500 µL

buffer AW1 and once with 500 µL AW2 and eluted using molecular grade water. The amount of

genomic DNA was quantified by a Qubit dsDNA HS Kit (Thermo, Q33230) and prepared for

Illumina analysis using the Illumina DNA prep kit (Illumina, 20018704). Illumina sequencing of

the spirulina genome was performed on a MiSeq instrument using a MiSeq Reagent Kit v2

(Illumina, MS-102-2003).

Illumina dataset analysis for copy number determination

Copy numbers of the genes of interest in SP1287, SP1308, SP1312, and SP1313 were

determined by comparing expression of the transgenes to 10 randomly selected genes

(referred to as "normalization genes": *ppaX, rpsA, chlM, degU2, radA, rbcL, sdhA, psbC, cysA1,*

smc6) in the genome. A copy number of about 1, within plus/minus 0.15, was expected and set

as the benchmark for a full genomic integration of 1 gene per genome copy. As a validation

control, we also chose a second set of 10 random spirulina genes (*thrB1, hisC1, isfD2, pilA, purB,*

apcE, nblA, gpmB, bicA2, cpcBA) which we compared pairwise to the "normalization genes" set

in a manner identical to the transgene. Gene copy number per genome was determined by first

using samtools (http://www.htslib.org/doc/) to generate an mpileup file with Illumina reads per

base pair. Then, the median read per base pair was determined across the 10 "normalization

genes" and the transgene. The median read per base pair of the transgene was then divided by

the 10 "normalization genes" median read-per-base-pair value.

Preparation of spirulina extracts for analysis of soluble protein

Soluble extracts from spray-dried or paste samples were prepared by bead beating in a

Percellys evolution bead beater (Bertin Technologies). Beads for this procedure were purchased

in pre-packed tubes from MP-Bio (cat# 6911500). Dried spirulina biomass was resuspended in

PBS containing Pierce Protease Inhibitor minitablets and 1 mM PMSF at a biomass

concentration of 2-10 mg/mL. Biomass samples were then centrifuged at 15,000g at 4 °C for 30

minutes, and the soluble fraction was transferred to a separate tube for downstream

applications.

Expression analysis of soluble recombinant proteins in spirulina

Recombinant protein expression in spirulina was measured by capillary electrophoresis

immunoassay (CEIA) using a Jess system (ProteinSimple). The Jess system was run as

recommended by the manufacturer. Briefly, dried biomass samples were diluted to a

concentration of 0.1-1 mg/mL using water and a 5X master mix prepared from an EZ Standard

Pack 1 (Bio-Techne). Purified protein controls used to generate standard curves were typically

loaded at a range of concentrations from 0.25-8 µg/mL. A 12-230 kDa Jess/Wes Separation

Module (ProteinSimple) was used for separation. A mouse anti-His-Tag antibody (GenScript)

was diluted 1:100 and used as the primary antibody. An anti-mouse NIR fluorescence-

conjugated secondary antibody (ProteinSimple) was primarily used for detection. Non-reducing

denaturing gels were performed in the same manner; however, reductant was withheld from

the 5X master mix. Data analysis was performed using the Protein Simple Compass software

and Microsoft Excel.

Expression analysis of total recombinant proteins (soluble and insoluble) in spirulina

Spray dried or paste samples were prepared by bead beating as above. The samples were

solubilized in PBS buffer mentioned above with a Triton X-100 concentration at 0.02%.

Following lysis, samples were diluted to 0.1-1 mg/mL in extraction buffer (60 mM dithiothreitol,

60 mM Na_2CO_3, 6 mM ethylenediamine tetraacetic acid, 2% (w/v) sodium dodecyl sulfate,

12% (w/v) sucrose and 0.02% Triton X-100). After the addition of the lysis buffer, samples were

immediately heated at 100 °C for 120 seconds, then centrifuged at 16,000g for 5 minutes.

Extracted samples were displayed on the Jess system as described above.

Cell lines, bacterial strains, toxins, and antibodies

Vero cells (ATCC) were cultured in Dulbecco's Modified Eagle Medium (DMEM) supplemented

with 10% heat-inactivated fetal bovine serum and 1% penicillin-streptomycin (Gibco). The cells

were grown in a high-humidity incubator at 37 °C and 5% CO_2. *C. difficile* toxin TcdB-10463 was

expressed in *B. megaterium* (a gift from C. Shoemaker, Tufts University) and purified using Ni-

NTA affinity and ion exchange chromatography. Purified *C. difficile* toxins TcdB-027, TcdB-1470,

and TcdB-078 were purchased from tgcBIOMICS. Bezlotoxumab was sourced from Henry Schein

Medical.

Expression of TcdB fragments in bacterial systems

The VHH-binding domains on TcdB[41] were designed for bacterial expression with an N-terminal

hexahistidine tag followed by MBP and TEV protease recognition site (ENLYFQG) and C-terminal

Avi-tag (LNDIFEAQKIEWHE) for enzymatic biotinylation. Based on the structural information, we

designed the 5D-binding region, the delivery and receptor-binding domain (DRBD) of TcdB for

expression using residues from amino acid position 1092 to position 1433 of TcdB. VHHs 7F and

E3 bind regions on the glucosyltransferase domain (GTD). A fragment containing this region was

designed to include amino acid positions 2 through amino acid 543 of TcdB. The constructs

were subcloned into a modified pET28 b(+) vector where the Kanamycin bacterial resistance

gene was replaced with Ampicillin bacterial resistance gene. Constructs were sequence verified

by Sanger sequencing (Genewiz). Sequence verified plasmids were transformed into BL21(DE3)

Escherichia coli cells (New England Biolabs) and plated on LB medium supplemented with 100

μg mL^{-1} ampicillin. Protein was expressed using autoinduction protocol[64]. Briefly, 1 L of Super

Broth media containing appropriate selection drug was inoculated with overnight cultures from

individual transformants, shaken at 37 °C for 8 hours followed by 16 °C for 24 hours. Expression

cultures were pelleted by centrifugation and stored at −20 °C until purification. Frozen cell

pellets were thawed on ice and resuspended in 100 mL of 50 mM Tris pH 8.0, 300 mM NaCl

supplemented with pierce protease inhibitor tablets EDTA free (Thermo Fisher Scientific cat. no.

A32965) and 1 mM phenylmethylsulfonyl fluoride (PMSF) (Thermo Fisher Scientific cat. no.

36978). Cells were lysed by high pressure homogenization (HPH) (Microfluidics LM20) at 12,000

psi. The lysates were clarified by centrifugation at 18,500*g* at 4 °C for 30 minutes using tabletop

centrifuge (Eppendorf Centrifuge 5810R). Soluble protein was separated from cell debris and

the supernatant was passed through 5 mL HisTrap columns (Cytiva, cat. no. 17524802)

equilibrated with 50mM Tris pH 8.0, 300 mM NaCl and 20 mM imidazole (Thermo Fisher

Scientific, cat. no. O3196-500). His-tagged protein bound HisTrap columns were washed with 10

column volumes (CV) wash buffer (50 mM Tris, pH 8.0, 300 mM NaCl and 20 mM imidazole).

Bound his-tagged protein were eluted using gradient flow at slope of 10% - 100% elution buffer

(50 mM Tris pH 8.0, 300 mM NaCl and 250 mM imidazole) in 5 CV. Fractions containing the

eluted protein were assessed by SDS-PAGE gel. SDS-PAGE analyses were performed by

incubating 2 μL of purified protein elution in NuPAGE LDS Sample Buffer (Thermo Fisher

Scientific) at 70˚C for 10 minutes. Bis-Tris NuPAGE 4 to 12% gels were placed into gel box filled

with NuPAGE MES SDS Buffer. Samples were loaded and gels run at constant volt of 170 mV for

40 minutes, followed by staining using Coomassie Blue gel stain. Fractions were pooled based

on level of purity, concentrated, and buffer exchanged into PBS. The samples were then filtered

through a 0.2 μm filter and run over a size-exclusion column on AKTA Pure FPLC-based size-

exclusion chromatography HiLoad 16/600 Superdex 200 pg column (Cytiva, cat. no. 28989335)

equilibrated in PBS. Purity and final analysis were performed using SDS-PAGE gel where 3 μg

purified protein samples were incubated at 70 °C in LDS loading buffer with or without 20 mM

dithiothreitol (DTT) reducing agent (Thermo Fisher Scientific, cat. no. R0862). Protein samples

were separated by size on NuPAGE Bis-Tris gels as described above. Purified samples were

biotinylated using BirA biotin ligase (Avidity Inc.) at 1 μg of BirA enzyme per 10 nM of protein.

Protein purification from spirulina biomass

The TcdB-binding proteins developed from VHHs 5D (pp1005, batch no. 145), E3 (pp1006, batch

no. 147), and 7F (pp1007, batch no. 147), and full-length lysin PlyCD (pp1092, batch no. 025)

were expressed in spirulina and purified from spray-dried biomass using Ni-NTA affinity and

size-exclusion chromatography 2-step purification protocol. Dried spirulina biomass from

strains expressing heterologous proteins were rehydrated at 30 mL/g of biomass in phosphate

buffer saline lysis buffer supplemented with protease inhibitor mix of 1 mM

phenylmethylsulfonyl fluoride (PMSF), (Thermo Fisher Scientific cat. no. 36978) and Pierce

Protease Inhibitor Tablets, EDTA-free, (Thermo Fisher Scientific cat no A32965). Rehydrated and

homogenous biomass resuspension were lysed using a high-pressure homogenizer

(Microfluidics LM20) at 12,000 psi. The lysates were clarified by centrifugation at 18500g on

tabletop Centrifuge (Eppendorf Centrifuge 5810R) for 30 minutes. Supernatant were clarified

by filtration using a coffee filter followed by 0.22 μm membrane filter. Recombinant proteins

were purified from the clarified lysate by affinity chromatography as describe above using 5 mL

HisTrap columns (Cytiva, cat. no. 17524802). Hexa-Histidine tagged and HisTrap-purified

samples were further refined using AKTA Pure FPLC-based size-exclusion chromatography on

HiLoad 16/600 Superdex 200 pg column (Cytiva, cat. no. 28989335). Protein purity and sizing

were assayed by size-exclusion chromatography on analytical Superdex 200 increase 10/300

(Cytiva, cat. no. 28990944) where 100 μL of purified protein at ~1 mg/mL was run in PBS.

Purified protein samples were also assessed on SDS-PAGE under reducing and non-reducing

conditions. 3 μg protein samples were incubated at 70 °C in LDS loading buffer with or without

20 mM dithiothreitol (DTT) reducing agent (Thermo Fisher Scientific, cat. no. R0862). Proteins

were determined to be >90% pure and exhibited the expected molecular weight size

distribution.

Native gel analysis

To assess the importance of disulfide bonds in pp1005, pp1006, and pp1007 dimer formation,

we compared protein migration in native and SDS-PAGE gel under reduced and non-reduced

conditions. Native gel analysis was performed following the manufacturer's recommendation

(Life Technologies) where 1 μg of purified protein was mixed with Native Page Sample Buffer

supplemented with or without 20 mM DTT. Native page anode buffer was prepared by mixing

50 mL NativePage 20X running buffer and 950 mL single distilled water. Similarly, 1 x

NativePage Dark Blue Cathode buffer was prepared by mixing 50 mL NativePage 20 x Running

buffer with 50 mL NativePage 20X Cathode Additive and 900 mL single distilled water. Gel

boxes were set up on ice and samples were loaded prior to adding running buffer. Inner

chamber of gel box was filled with NativePage Dark Blue Cathode buffer while the outer

chamber was filled with NativePage Anode buffer. Gels were run at a constant 150 V for 2.5

hours. Comparable sample preparations were run in SDS-PAGE gel as described above.

Assessment of VHH binding to TcdB fragments with bio-layer interferometry (BLI)

BLI measurements were performed on the Octet RED96 system (ForteBio) using High Precision

Streptavidin (SAX) Biosensors (ForteBio). Biosensors were hydrated with phosphate buffer,

pH 7.4 at room temperature for 10 minutes on Microplate 96-well flat-bottom (Greiner,

655209). All kinetics experiments were performed at 30 °C temperature with 500 rpm agitation

in the kinetics module. Biosensors were dipped into PBS containing wells for 60 seconds prior to

antigen loading. Biosensors were loaded with enzymatically biotinylated TcdB antigen

fragments at 10 µg/mL in phosphate buffer, pH 7.4 for 200 seconds to achieve ~0.6 – 1 nm

response. Loading was quenched by incubating biosensors in 50 µM Biocytin (Sigma Aldrich,

576-19-2) for 60 seconds. Baseline were established by incubating antigen loaded biosensors in

kinetics buffer (PBS + 0.02 % Tween 20, 0.1 % BSA, 0.05 % sodium azide) for 300 seconds. After

stablishing baseline, the rate of association was measured by incubating antigen loaded

biosensor tips into 3-fold dilution series of monomeric (pp1239, pp1238, and pp1240

respectively) or dimeric (pp1005, pp1006, and pp1007 respectively) 5D, E3, and 7F constructs.

The concentration series used to determine binding kinetics of pp1238 were 93.6 nM, 31.2 nM,

10.4 nM, 3.47 nM, 1.16 nM, and 0.36 nM. Concentrations used to determine binding kinetics of

pp1239 were 93.6 nM, 31.2 nM, 10.4 nM, 3.47 nM, 0.38 nM, and 0.38 nM. Concentrations used

for pp1240 were 92.2 nM, 30.7 nM, 10.2 nM, 3.41 nM, 1.14 nM and 0.38 nM. pp1005 was

assayed at 16.2 nM, 5.39 nM, 1.8 nM, 0.60 nM, 0.20 nM and 0.07 nM. Concentrations used for

pp1006 were 16.8 nM, 5.6 nM, 1.87 nM, 0.62 nM, 0.21 nM, and 0.07 nM. pp1007 were assayed

at concentrations of 84.9 nM, 28.3 nM, 9.43 nM, 3.14 nM, 1.05 nM, and 0.35 nM. Analyte

bound biosensors were dipped into kinetics buffer for 300 - 600 seconds to measure rate of

dissociation. Kinetic analysis was performed using the HT 11.1.1.39 Data Analysis module

(ForteBio). Results were double referenced. Both association and dissociation steps were used

in 1:1 binding model global data fitting model.

Toxin-neutralization assay

Toxin-neutralization assays using TcdB1.1, TcdB2.1, TcdB3.1 and TcdB5.1 were performed using

the Agilent xCELLigence RTCA MP. This system measures cell adherence via biosensors in the

bottom of a 96-well plate by applying a small electric current across microelectrodes. A unitless

measurement called cell index (CI) was then calculated based on changes in electrode

impedance. Because adherent cells increase electrode impedance, toxin induced cell rounding

can be determined from changes in CI.

Agilent 96-well E-plates were seeded with 100 µl/well of 1×10^5 cells/mL in DMEM culture

medium with 10% FBS and 1% Pen/Strep and allowed to rest for 30 minutes before going into

an xCELLigence real-time cell analyzer in a Caron CO_2 37 °C incubator to run overnight. CI was

monitored every 15 minutes for 20 to 24 hours. TcdB was serially diluted in culture medium in a

96-well set up plate. The VHH homodimer, lysin, and bezlotoxumab samples were diluted in

medium and then added to the serially diluted toxin. The TcdB/protein mixture was incubated

at 37 °C for 1 hour. After 1 hour, media was carefully removed from the cells seeded onto the E-

plates and 100 µL of the VHH homodimer and toxin mixture was added to the cells in each well.

CI was monitored every 15 minutes for at least 24 hours. Each run included a TcdB only

standard curve as a positive control and cells only as a negative control. VHH homodimer was

also tested in the absence of TcdB to assess any effects of VHH alone on CI. At the 24-hour time

point after treatment, the CI measurements for each well were normalized by setting the CI to

1 at the last reading taken before treatment was added. The normalized cell index (nCI) was

plotted using Prism and a nonlinear regression analysis was performed to calculate the IC_{50} of

TcdB under treatments.

TcdB's effect on cell adhesion and inhibition of TcdB by anti-TcdB VHH homodimers

Cell adhesion assay data were imported, normalized, and analyzed using R (version 4.1.1 The R

Foundation for Statistical Computing). Nonlinear regression was performed using the

minpack.lm routine in the *nlsLM* package, which implements the Levenberg-Marquardt

algorithm to perform bounded minimization[65].

Lysin activity assay

The lytic activity of pp1092 against *Bacillus subtilis* SL4 was assessed based upon previously

described methods with modifications to adapt the assay to a 96-well plate format[42]. Mid-log

phase grown SL4 were grown in BHI (37 °C, 250 rpm) harvested and washed twice in PBS by

centrifugation (room temperature, 6,000g, 10 minutes). Bacteria were then combined with 1ug

digested or undigested pp1092 in a final volume of 200 μL (OD = 1.0) and then turbidity

reduction was measured over time (37 °C, shaking). Each unknown was fit along with 5 3-fold

serial dilutions of undigested standards using a modified 5-parameter nonlinear logistic model.

The time of 50% OD_{600}, TC_{50}, was estimated as a 4-parameter nonlinear logistic model with lysin

concentration as the independent variable. For the standards, this concentration is known. For

the unknown sample, this value is estimated. All results were expressed as the mean ± standard

deviation as calculated by Excel software (v16.48, Renton, WA). Standard curves were

determined by linear regression analysis by Excel software (v16.48, Renton, WA). EC_{50} values

were calculated using GraphPad Prism (version number, GraphPad Software Inc, San Diego, CA).

For comparison of spirulina-expressed and *E. coli*-expressed PlyCD, *C. difficile* strain ATCC 43255

was grown to mid-log phase (optical density at 600 nm (OD_{600nm}) = 0.5) at 37 °C under

anaerobic conditions in reduced Brain Heart Infusion Medium (BD Biosciences) supplemented

with 0.5% (wt/vol) yeast extract and 0.1% (wt/vol) L-cysteine. The bacteria were harvested at

4,000 rpm for 15 minutes, washed and resuspended to an OD_{600nm} of 2.0 in 50 mM sodium

phosphate, pH 7.0. Using a 96-well nontreated polystyrene microtiter plate (Corning), 100 µL of

buffer with or without (untreated controls) lysin at initial concentrations ranging from 4 to 256

µg/mL were added to individual wells. Following the addition of 100 µL bacteria, the OD_{600nm} of

each sample was measured every 60 seconds for a total of 15 minutes at 37°C using a

SpectraMax M5 microplate reader (Molecular Devices). All data represents the average of

triplicate experiments.

VHH homodimer protease digestion for ELISA and neutralization assays

In vitro protease digestions of anti-*C. difficile* VHH homodimers were prepared in a matrix of

soluble lysate from wild-type spirulina (SP003) in Bis-Tris buffer pH 6.0 and treated with trypsin,

chymotrypsin, or elastase. VHH homodimers in PBS at were diluted in SP3 lysate, Bis-Tris buffer,

and protease so that the final concentrations were VHH homodimer, 20 mg/mL of SP3 lysate,

and a "fed" or "fasted" amount of protease. This level of SP3 lysate and VHH homodimer

simulates a expression level of 1% in spirulina (by dry weight). For trypsin and chymotrypsin,

the "fed" and "fasted" levels of protease in the digestion reactions were 0.1 and 0.01 mg/mL,

while for elastase, 5 and 0.5 µg/mL were used respectively. Samples were incubated at 37 °C for

1 hour with shaking at 900 rpm in an Eppendorf ThermoMixer. At the end of the incubation,

reactions were quenched with an equal volume of stop buffer containing 2 mM PMSF and 2x

Pierce protease inhibitor minitablet (Thermo Fisher) in PBS; elastase digests were additionally

neutralized with a 1 mM elastase inhibitor III. Samples were kept on ice until preparation for

ELISA or cell rounding assay. The final concentration of each VHH homodimer after protease

neutralization was 0.1 mg/mL.

VHH homodimer quantitation by ELISA

Digested VHH homodimers were analyzed for binding activity via ELISA using TcdB. Thermo

Scientific Nunc ELISA plates were coated with 100 µL/well of inactive TcdB at 2 µg/mL in CBC

binding buffer and incubated overnight at 4 °C. Plates were washed 3 times with PBS

supplemented with 0.05% Tween-20 (PBS-tw), then blocked for 1 hour with 300 µL/well of

block (PB-tw supplemented with 5% BLOTTO). Dilutions of undigested and digested VHH

homodimer were prepared in a low-binding 96-well plate by serially diluting with block. 6

dilutions of each undigested VHH homodimer were prepared for a standard curve. Two

dilutions of each protease digested sample were prepared for quantification by interpolation

from the standard curve. A 100 µL sample of each VHH homodimer was added to ELISA plate

wells and incubated for 1 hour at room temperature on a plate shaker. The plates were washed

as before. An HRP-conjugated anti-camelid VHH antibody cocktail (Genscript) was diluted

50,000-fold in block and 100 µL/well was added to the ELISA plates for 50 minutes with shaking.

Plates were washed 2 times with PBS-tw and 1 time with PBS. Finally, 100 µL/well of TMB Ultra

ELISA substrate (Thermo Fisher) was added to each well. The plates were developed in the dark

at room temperature on a plate shaker for 10 minutes before being quenched with 50 µL of 2.5

M sulfuric acid, and the absorbance was read at 450 nm with a Spectra Max M5 plate reader.

All digestion reactions were in duplicate, in 3 independent experiments.

Calculation of VHH homodimer protease resistance

The concentration of VHH homodimer in the protease-digested samples was calculated by

interpolating the concentration from standard curves generated with undigested VHH

homodimer. The percentage of remaining VHH homodimer activity after digestion was

calculated by dividing the interpolated value by the amount present if no VHH homodimer were

digested. Each replicate was calculated separately, and the average and standard deviation

were determined across all replicates.

Toxin neutralization assay of digested VHH homodimer s

Chymotrypsin-digested pp1005, pp1006, and pp1007 were assessed for toxin-neutralizing

activity as described. The final concentration of pp1005, pp1006, and pp1007 was 0.05 nM,

5 nM, and 5 nM respectively and the TcdB was 1:5 serially diluted starting at 250 pM.

Lysin digestion conditions

In digestion buffer (20 mM Tris, 150 mM NaCl, 3 mM $CaCl_2$ at pH 6.4) 0.2 µg/µL protein was

mixed with proteases trypsin (Promega sequencing grade), chymotrypsin (BD), and elastase

(BD) at 0.1 µg/µL or 0.01 µg/µL and incubated for 1 hour at 37 °C (200 rpm shaking). Reactions

were immediately quenched by addition of 1 mM PMSF and Protease Inhibitor Mini Tablets,

EDTA-free (Thermo Scientific, Pierce). For samples with elastase, 1 mM elastase inhibitor III

(Milipore Sigma) was added as well. To visualize digestion, 2 μg of digested and undigested

proteins were displayed a gradient BOLT-PAGE gel at 200 V for 20 minutes and then stained

with Coomassie.

Densitometry

Pixel intensity of Coomassie stained proteins were measured using ImageJ software,

normalizing all bands to the undigested band for each condition after subtracting background

pixel intensity of an identically sized bounding box.

Preparation of *C. difficile* spores for mouse study

C. difficile strain M7404, a ribotype 027 clinical isolate from Canada, was cultured on HIS-T agar

(3.7 % (w/v) heart infusion (HI) (Oxoid), 0.5 % (w/v) yeast extract, 1.5 % (w/v) agar

supplemented with 0.375 % (w/v) glucose, 0.1 % (w/v) L-cysteine and 0.1 % (w/v) sodium

taurocholate (New Zealand Pharmaceuticals)) in an anaerobic chamber (Don Whitley Scientific)

with an atmosphere of 10 % H2, 10 % CO_2 and 80 % N2 at 37 °C. To prepare spores, strains were

sub-cultured on HIS-T agar and inoculated into 500 mL tryptone yeast (TY) broth (3.0%

tryptone, 2.0% yeast extract, and 0.1% sodium thioglycolate) and grown anaerobically at 37 °C

for 10 days. Spores were harvested by centrifugation at 10,000g for 20 minutes at 4 °C and

washed 5 times with chilled distilled water prior to heat-shocking at 65 °C for 20 minutes and

quantified by plating onto HIS-T agar. Spores were adjusted to 10^6 CFU/mL in PBS containing

0.05 % Tween-80 in preparation for mouse infections.

Mouse study design

Animal handling and experimentation was performed in accordance with Victorian State

Government regulations and approved by the Monash University Animal Ethics Committee

(Monash University AEC no. 21731). Male C57BL/6J mice, 6-7 weeks old, were assigned to

either a spirulina therapeutic treatment group, positive control (vancomycin-treated), or

negative control (wild-type spirulina-treated) group and caged in groups of 5. To induce

susceptibility to *C. difficile*, the animals were pre-treated with an antibiotic cocktail in the

drinking water containing kanamycin (0.4 mg/mL), gentamicin (0.035 mg/mL), colistin (850

U/mL), metronidazole (0.215 mg/mL), vancomycin (0.045 mg/mL), and cefaclor (0.3 mg/mL) for

7 days, followed by 4 days of cefaclor (0.3 mg/mL) alone. Starting 1 day prior to infection (day -

1), the animals were orally gavaged with 200 µL of treatment once daily for the first two days

(peak of infection) and then 100 µL once daily for the remainder of the study. Mice were

infected with 10^5 spores/mouse by oral gavage.

The spirulina treatments and controls were resuspended in PBS (5% w/v) by mixing up and

down using an 18G blunt needle and 1 mL syringe for 2.5 minutes and administered within 10

minutes of preparation. Vancomycin was diluted to 6.25 mg/mL prior to administration. On the

day of *C. difficile* infection (day 0), antibiotic pre-treatment was ceased, the mice were switched

to plain water, and infection was initiated 3 hours after the treatment dose. Mice were

monitored daily for signs of infection, including weight loss, and were euthanized upon losing

15% body weight, if they reached pre-defined clinical endpoints as required by ethics, or at the

end of the study (day 4). As a surrogate assessment of colonization, fecal samples collected at

24 hours post-infection and then again at the time of euthanasia were analyzed to enumerate

the shedding of *C. difficile* spores. Subjects treated with a spirulina therapeutic cocktail were

compared to the cohort administered wild-type spirulina.

Quantification of spore shedding from mouse feces

Fecal pellets were collected 24 hours post-infection and at time of euthanasia, resuspended in

PBS (100 mg/mL), heat shocked (30 minutes, 65 °C) and plated for spore enumeration onto HIS

agar supplemented with 0.1% (w/v) L-cysteine, 0.1% (w/v) sodium taurocholate, 0.375% (w/v)

glucose, 250 μg/mL D-cycloserine, 8 μg/mL cefoxitin, and 10 μg/mL erythromycin.

Analysis of colonization data was performed by fitting linear models of $\log_{10}$ cfu/g data per

landmark time with treatment as a (potential) explanatory factor. A coefficient for wild-type

spirulina describes the mean $\log_{10}$ cfu/g for that treatment, and coefficients for each other

treatment describes the difference in mean $\log_{10}$ cfu/g for that treatment relative to wild-type

spirulina. Standard errors of estimates were reported, along with t-statistics and 2-tailed p-

value corresponding to the t-statistic with $\alpha=0.05$.

Analysis of clinical outcomes—mouse study

Analysis of maximum weight loss data was performed by fitting linear models of maximum

weight loss data with treatment as a (potential) explanatory factor. A coefficient for wild-type

spirulina describes the mean maximum weight loss for that treatment, and coefficients for each

other treatment describes the difference in mean loss for that treatment relative to wild-type

spirulina. Standard errors of estimates were reported, along with t-statistics and 2-tailed p-

value corresponding to the t-statistic with $\alpha=0.05$.

Analysis of survival data was performed on tabulated number at risk and number of events at

each time. Data was compared for each treatment to wild-type spirulina results and a P value

was computed using the log-rank statistic for each comparison.

Hamster study challenge agent preparation and confirmation

C. difficile strain R20291 (NAP1/BI/027) spore stock was titered on TCCFA plates after a 30-

minute incubation at 65 °C and determined to be 5 x 10^5 cfu/mL in a volume of 5 mL of PBS.

This spore stock was diluted to 15,000 spores/50 µL and further diluted in an equal volume of

sterile PBS containing 5% sucrose to facilitate administration of 15,000 spores in 100 µL per

animal.

To verify challenge doses, 100 µL of inoculum were serially diluted (10-fold dilutions) and plated

onto TCCFA plates. After incubation of the plates under anaerobic conditions at 37 °C for 48

hours, the number of colonies were enumerated, and the doses calculated based on these

numbers. The inoculum was measured to be 15,000 cfu.

Hamster study design

Hamsters (Envigo, Indianapolis, IN) were visibly checked upon arrival and allowed to acclimate

to their new environment for 2 days. Each animal was given a physical exam and randomly

assigned (adjusting only to distribute body weight evenly) to 1 group. They were fed *ad libitum*

with sterilized 2018S Teklad Global 18% Rodent Diet (Envigo, Indianapolis, IN) and sterile water

provided in standard bottles with sippers during the study. To sensitize the animals to *C. difficile*

infection, clindamycin, 15 mg/mL in saline, was administered intraperitoneally at 30 mg/kg,

once a day, for 3 consecutive days. During clindamycin treatment, cage bottoms were changed

daily for 3 consecutive days beginning on the first day to prevent bacterial recolonization from

coprophagy.

ACKNOWLEDGEMENTS

We thank Charles Shoemaker, George McDonald, and Fred Cross for help, discussion, and

advice. We thank Meagan James and other members of the Lyras laboratory for help with the

mouse studies. For the hamster challenge studies Lumen Bioscience has utilized the non-clinical

and pre-clinical services program offered by the National Institute of Allergy and Infectious

Diseases, especially support from Sangun Lee, Christian Gonzalez and Ryan Ranallo.

REFERENCES
748

1. Knight, D. R., Elliott, B., Chang, B. J., Perkins, T. T. & Riley, T. V. Diversity and Evolution in the
Genome of *Clostridium difficile*. *Clinical Microbiology Reviews* 28, 721–741 (2015).

2. Balsells, E. *et al.* Global burden of *Clostridium difficile* infections: a systematic review and
meta-analysis. *Journal of Global Health* 9, 010407 (2019).

3. Guh, A. Y. *et al.* Trends in U.S. Burden of *Clostridioides difficile* Infection and Outcomes. *N.*
Engl. J. Med. 382, 1320–1330 (2020).

4. Salavert, M. *et al.* Cost-effectiveness analysis of bezlotoxumab added to standard of care
versus standard of care alone for the prevention of recurrent *Clostridium difficile* Infection in
High-Risk Patients in Spain *Adv. Ther* (2018) doi:doi.org/10.1007/s12325-018-0813-y.

5. Seal, D. *et al.* Treatment of relapsing *Clostridium difficile* diarrhoea by administration of a
non-toxigenic strain. *Eur J Clin Microbiol* 6, 51–53 (1987).

6. Kuehne, S. A. *et al.* The role of toxin A and toxin B in *Clostridium difficile* infection. *Nature*
467, 711–713 (2010).

7. Lyras, D. *et al.* Toxin B is essential for virulence of *Clostridium difficile*. *Nature* 458, 1176–
1179 (2009).

8. Chen, K. *et al.* A probiotic yeast-based immunotherapy against *Clostridioides difficile*
infection. *Sci Transl Med* 12, eaax4905 (2020).

9. Tao, L. *et al.* Frizzled proteins are colonic epithelial receptors for *C. difficile* toxin B. *Nature*
538, 350–355 (2016).

10. Henkel, D. *et al.* Receptor Binding Domains of TcdB from *Clostridioides difficile* for
Chondroitin Sulfate Proteoglycan-4 and Frizzled Proteins Are Functionally Independent and
Additive. *Toxins (Basel)* 12, 736 (2020).

11. Qa'Dan, M., Spyres, L. M. & Ballard, J. D. pH-induced conformational changes in *Clostridium*
difficile toxin B. *Infection and Immunity* 68, 2470–2474 (2000).

12. Reineke, J. *et al.* Autocatalytic cleavage of *Clostridium difficile* toxin B. *Nature* 446, 415–419
(2007).

13. Sehr, P. *et al.* Glucosylation and ADP ribosylation of rho proteins: effects on nucleotide
binding, GTPase activity, and effector coupling. *Biochemistry* 37, 5296–5304 (1998).

14. Just, I. *et al.* Glucosylation of Rho proteins by *Clostridium difficile* toxin B. *Nature* 375, 500–
503 (1995).

15. Aelst, L. V. & Symons, M. Role of Rho family GTPases in epithelial morphogenesis. *Genes &*
Development 16, 1032–1054 (2002).

16. Schlegel, N., Meir, M., Spindler, V., Germer, C.-T. & Waschke, J. Differential role of Rho
GTPases in intestinal epithelial barrier regulation in vitro. *J Cell Physiol* 226, 1196–1203 (2011).

17. Pradhan, R., Ngo, P. A., Martínez-Sánchez, L. D., Neurath, M. F. & López-Posadas, R. Rho
GTPases as Key Molecular Players within Intestinal Mucosa and GI Diseases. *Cells* 10, 66 (2021).

18. Carter, G. P. *et al.* Defining the Roles of TcdA and TcdB in Localized Gastrointestinal Disease,
Systemic Organ Damage, and the Host Response during *Clostridium difficile* Infections. *mBio* 6,
e00551 (2015).

19. Mileto, S. J. *et al.* Clostridioides difficile infection damages colonic stem cells via TcdB,
impairing epithelial repair and recovery from disease. *Proceedings of the National Academy of*
Sciences 117, 8064–8073 (2020).

20. Bella, S. D., Ascenzi, P., Siarakas, S., Petrosillo, N. & Masi, A. di. *Clostridium difficile* Toxins A
and B: Insights into Pathogenic Properties and Extraintestinal Effects. *Toxins (Basel)* 8, 134–25
(2016).

21. Hutton, M. L. *et al.* Bovine antibodies targeting primary and recurrent *Clostridium difficile*
disease are a potent antibiotic alternative. *Sci. Rep.* 7, 16020–13 (2017).

22. Steele, J., Mukherjee, J., Parry, N. & Tzipori, S. Antibody Against TcdB, but Not TcdA,
Prevents Development of Gastrointestinal and Systemic *Clostridium difficile* Disease. *J. Infect.*
Dis. 207, 323–330 (2012).

23. Wilcox, M. H. *et al.* Bezlotoxumab for Prevention of Recurrent *Clostridium difficile* Infection.
N. Engl. J. Med. 376, 305–317 (2017).

24. Dieterle, M. G., Rao, K. & Young, V. B. Novel therapies and preventative strategies for
primary and recurrent *Clostridium difficile* infections. *Annals of the New York Academy of*
Sciences 1435, 110–138 (2019).

25. Song, J. H. & Kim, Y.-S. Recurrent *Clostridium difficile* Infection: Risk Factors, Treatment, and
Prevention. *Gut Liver* 13, 16–24 (2019).

26. Singh, T. *et al.* Updates in Treatment of Recurrent *Clostridium difficile* Infection. *J Clin Med*
Res 11, 465–471 (2019).

27. Pruss, K. M. & Sonnenburg, J. L. *C. difficile* exploits a host metabolite produced during toxin-
mediated disease. *Nature* 593, 261–265 (2021).

28. Yang, Z. *et al.* A novel multivalent, single-domain antibody targeting TcdA and TcdB prevents
fulminant *Clostridium difficile* infection in mice. *J. Infect. Dis.* 210, 964–972 (2014).

29. Schmidt, D. J. *et al.* A Tetraspecific VHH-Based Neutralizing Antibody Modifies Disease
Outcome in Three Animal Models of Clostridium difficile Infection. *Clinical and Vaccine*
Immunology 23, 774–784 (2016).

30. Roberts, A. K. *et al.* A Novel, Orally Delivered Antibody Therapy and Its Potential to Prevent
Clostridioides difficile Infection in Pre-clinical Models. *Front Microbiol* 11, 578903 (2020).

31. Babcock, G. J. *et al.* Human Monoclonal Antibodies Directed against Toxins A and B Prevent
Clostridium difficile-Induced Mortality in Hamsters. *Infection and Immunity* 74, 6339–6347
(2006).

32. Sponseller, J. K. *et al.* Hyperimmune Bovine Colostrum as a Novel Therapy to Combat
Clostridium difficile Infection. *J. Infect. Dis.* 5, 549–8 (2014).

33. Mansfield, M. J. *et al.* Phylogenomics of 8,839 *Clostridioides difficile* genomes reveals
recombination-driven evolution and diversification of toxin A and B. *PLoS Pathog* 16, e1009181
(2020).

34. Bartlett, J. G. Bezlotoxumab — A New Agent for *Clostridium difficile* Infection. *N. Engl. J.*
Med. 376, 381–382 (2017).

35. Dissel, J. T. van *et al.* Bovine antibody-enriched whey to aid in the prevention of a relapse of
Clostridium difficile-associated diarrhoea: preclinical and preliminary clinical data. *J Med*
Microbiol 54, 197–205 (2005).

36. Numan, S. C., Veldkamp, P., Kuijper, E. J., Berg, R. J. van den & Dissel, J. T. van. Clostridium
difficile-associated diarrhoea: bovine anti-*Clostridium difficile* whey protein to help aid the
prevention of relapses. *Gut* 56, 888 (2007).

37. Mattila, E. *et al.* A randomized, double-blind study comparing *Clostridium difficile* immune
whey and metronidazole for recurrent Clostridium difficile-associated diarrhoea: Efficacy and
safety data of a prematurely interrupted trial. *Scand J Infect Dis* 40, 702–708 (2009).

38. McGovern, B. H. *et al.* SER-109, an Investigational Microbiome Drug to Reduce Recurrence
after *Clostridioides difficile* infection: Lessons Learned from a Phase 2 Trial. *Clin Infect Dis* 313,
1719 (2020).

39. Ghose, C. & Kelly, C. P. The Prospect for Vaccines to Prevent *Clostridium difficile* Infection.
Infectious Disease Clinics of North America 29, 145–162 (2015).

40. Bruyn, G. de *et al.* Safety, immunogenicity, and efficacy of a *Clostridioides difficile* toxoid
vaccine candidate: a phase 3 multicentre, observer-blind, randomised, controlled trial. *The*
Lancet Infectious Diseases 21, 252–262 (2021).

41. Chen, P. *et al.* Structure of the full-length Clostridium difficile toxin B. *Nat Struct Mol Biol* 26,
712–719 (2019).

42. Wang, Q., Euler, C. W., Delaune, A. & Fischetti, V. A. Using a Novel Lysin To Help Control
Clostridium difficile Infections. *Antimicrobial Agents and Chemotherapy* 59, 7447–7457 (2015).

43. Fischetti, V. A. Bacteriophage lysins as effective antibacterials. *Current Opinion in*
Microbiology 11, 393–400 (2008).

44. Peng, Z. *et al.* A Novel Bacteriophage Lysin-Human Defensin Fusion Protein Is Effective in
Treatment of *Clostridioides difficile* Infection in Mice. *Front. Microbio.* 9, 3234 (2018).

45. Jester, B. *et al.* Expression and manufacturing of protein therapeutics in spirulina. *bioRxiv*
2021.01.25.427910 (2021) doi:10.1101/2021.01.25.427910.

46. Raran-Kurussi, S. & Waugh, D. S. The Ability to Enhance the Solubility of Its Fusion Partners
Is an Intrinsic Property of Maltose-Binding Protein but Their Folding Is Either Spontaneous or
Chaperone-Mediated. *Plos One* 7, e49589 (2012).

47. Hanoune, J. & Defer, N. Regulation and role of adenylyl cyclase isoforms. *Annu. Rev.*
Pharmacol. Toxicol. 41, 145–174 (2001).

48. Burgers, P. P. *et al.* Structure of smAKAP and its regulation by PKA-mediated
phosphorylation. *FEBS J* 283, 2132–2148 (2016).

49. Ryder, A. B. *et al.* Assessment of *Clostridium difficile* Infections by Quantitative Detection of
tcdB Toxin by Use of a Real-Time Cell Analysis System. *J Clin Microbiol* 48, 4129–4134 (2010).

50. Pollock, N. R. Ultrasensitive Detection and Quantification of Toxins for Optimized Diagnosis
of *Clostridium difficile* Infection. *Journal of Clinical Microbiology* 54, 259–264 (2016).

51. Shen, E. *et al.* Subtyping analysis reveals new variants and accelerated evolution of
Clostridioides difficile toxin B. *Commun Biology* 3, 347 (2020).

52. Alonso, C. D. *et al.* Ultrasensitive and quantitative toxin measurement correlates with
baseline severity, severe outcomes, and recurrence among hospitalized patients with
Clostridioides difficile infection. *Clin Infect Dis* ciab826- (2021) doi:10.1093/cid/ciab826.

53. Tickler, I. A. *et al.* Changes in molecular epidemiology and antimicrobial resistance profiles
of *Clostridioides (Clostridium) difficile* strains in the United States between 2011 and 2017.
Anaerobe 60, 102050 (2019).

54. Li, Z. *et al.* Ribotype Classification of *Clostridioides difficile* Isolates Is Not Predictive of the
Amino Acid Sequence Diversity of the Toxin Virulence Factors TcdA and TcdB. *Front Microbiol*
11, 1310 (2020).

55. Chou, T. C. & Talaly, P. A simple generalized equation for the analysis of multiple inhibitions
of Michaelis-Menten kinetic systems. *J. Biol. Chem.* 252, 6438–6442 (1977).

56. Einav, T. & Bloom, J. D. When two are better than one: Modeling the mechanisms of
antibody mixtures. *Plos Comput Biol* 16, e1007830 (2020).

57. Gilchuk, P. *et al.* Analysis of a Therapeutic Antibody Cocktail Reveals Determinants for
Cooperative and Broad Ebolavirus Neutralization. *Immunity* 52, 388-403.e12 (2020).

58. Mulangu, S. *et al.* A Randomized, Controlled Trial of Ebola Virus Disease Therapeutics. *New*
Engl J Med 381, 2293–2303 (2019).

59. Weinreich, D. M. *et al.* REGN-COV2, a Neutralizing Antibody Cocktail, in Outpatients with
Covid-19. *New Engl J Medicine* 384, NEJMoa2035002 (2020).

60. Shah, D. K. & Betts, A. M. Antibody biodistribution coefficients: inferring tissue
concentrations of monoclonal antibodies based on the plasma concentrations in several
preclinical species and human. *mAbs* 5, 297–305 (2013).

61. Mudie, D. M. *et al.* Quantification of Gastrointestinal Liquid Volumes and Distribution
Following a 240 mL Dose of Water in the Fasted State. *Mol. Pharmaceutics* 11, 3039–3047
(2014).

62. Schiller, C. *et al.* Intestinal fluid volumes and transit of dosage forms as assessed by
magnetic resonance imaging. *Aliment Pharmacol Ther* 22, 971–979 (2005).

63. Gerding, D. N., Johnson, S., Rupnik, M. & Aktories, K. *Clostridium difficile* binary toxin CDT.
Gut Microbes 5, 15–27 (2014).

64. Studier, F. W. Protein production by auto-induction in high-density shaking cultures. *Protein*
Expres Purif 41, 207–234 (2005).

65. Moré, J. J. Numerical Analysis. *Lect Notes Math* 105–116 (1978) doi:10.1007/bfb0067700.

66. *Akaike information criterion statistics*. vol. 29 (reidel, 1987).

genomically integrated gene-of-interest

Figure 1. Spirulina strain engineering and characterization. A) Diagram of modified anti-
C. difficile TdcB VHHs 5D, E3, 7F, PlyCD, PlyCD$_{1-174}$, and 5HVZ-mediated dimer formation. B) A
schematic diagram of the genomic integration locus (WP_006618409.1). PCR was performed
with indicated primers to analyze the segregation and specificity of integration for each strain.
PCR products with primers MP1/MP2 showed bands of predicted sizes in all strains, and
absence of the parental alleles, suggesting the segregation to homozygosity was complete. The
primer pairs for amplification of the left homology arm (LHA) and right homology arm (RHA)
include one priming site (MP3 and MP6) that is only present in the spirulina genome at the
target locus and absent from the donor plasmid. PCR amplification of the LHA and RHA with
MP3/MP4 and MP5/MP6, respectively, demonstrated that genomic integration was site-
specific. The DNA sequences from all PCR products were verified to match expectations.
C) Copy numbers of the genes of interest (GOIs) were determined with next-generation

914   sequencing by comparing reads of the GOIs to ten randomly selected native spirulina genes.

915   Dotted line indicates range in which the relative copy numbers were considered equivalent to

916   one.

918
919
Figure 2. Characterization of spirulina-expressed anti-TcdB VHHs and lysins. A) Chromatograms
of pp1005, pp1006, pp1007, pp1092, and pp1093 extracted from spirulina biomass and
detected under reducing conditions using a capillary electrophoresis-based immunoassay.
Soluble protein, as a percentage of washed dry weight, is indicated on each graph. B) Spirulina
expressed and purified pp1005, pp1006, and pp1007 in aqueous solution (PBS, pH 7.4) assayed
on a Superdex 200 10/300 GL FPLC column. Grey peaks correspond to molecular size standards.
The monomeric and dimeric portions of the VHHs are indicated. Disulfide bond formation
between monomers was assessed using denaturing gel electrophoresis under non-reducing and
reducing (50 mM DTT) conditions. C) Native gel electrophoresis analysis of pp1005 (1, 4),
pp1006 (2, 5), and pp1007 (3, 6) under non-reducing or reducing (50 mM DTT) conditions. In
the presence of reducing agent the dimeric state of the VHHs was mostly preserved.

B

Sample ID	K_D (nM)	k_a (1/Ms)	k_d (1/s)
pp1239 (M)	2.56 (0.02)	1.64E+05 (0.12)	4.20E-04 (0.13)
pp1005 (D)	0.002 (0)	1.16E+05 (0.11)	2.59E-07 (0)
pp1238 (M)	4.85 (0.06)	1.45E+05 (0.02)	7.02E-04 (0.02)
pp1006 (D)	0.03 (0)	7.48E+04 (1.68)	2.57E-06 (0)
pp1240 (M)	2.09 (0.06)	1.26E+05 (0.04)	2.63E-04 (0.03)
pp1007 (D)	0.58 (0.07)	1.54E+05 (0.17)	8.97E-05 (0.30)

931
932
Figure 3. Binding affinities and toxin neutralizing activity of TcdB VHHs. A) BLI binding
sensorgrams for analysis of binding kinetic for anti-TcdB VHHs as monomers (pp1239, pp1238,
and pp1240) and dimers (pp1005, pp1006, and pp1007). B) The rate constants k_a and k_d were
calculated from concentration-dependent responses. The apparent equilibrium binding
constant (K_D) for each was determined from the rate of association (k_a), and rate of dissociation
(k_d). Values of K_D, k_a and k_d are indicated in the table (M, monomer; D, dimer).

945
946
Figure 4. Comparative neutralization of the five major TcdB types by LMN-201 and
bezlotoxumab. A) Neutralizing activity of pp1005, pp1006, pp1007, and the triple cocktail on
toxin types 1.1, 2.1, 3.1 and 5.1. 1.5 nM of each dimeric VHH or the cocktail of all three dimeric
VHHs (0.5 nM of each) was tested against serially diluted TcdB with Vero cells on the RTCA
system. As comparison, 3 nM of bezlotoxumab was similarly evaluated. To rule out
interference, the effect of 1 nM lysin on VHH cocktail efficacy was assessed. B) IC$_{50}$ of TcdB
under treatment was calculated from the data shown in A and compared.

954
955
Figure 5. Characterization of spirulina-expressed anti-*C. difficile* lysin. A) A chromatogram of
pp1092 after total protein extraction (black) and soluble protein extraction (red). More than
75% of the lysin protein is soluble. B) Spirulina-expressed, purified lysins, pp1092 and pp1093,
in aqueous solution (1xPBS, pH 7.4) evaluated by size-exclusion chromatography (Superdex 200
10/300 GL FPLC column). Purified protein samples were also assessed using denaturing gel
electrophoresis under non-reducing and reducing (50 mM DTT) conditions. C) Activity of
spirulina extracts (wild-type (WT) expressing no lysin, SP1286 expressing pp1093, and SP1287
expressing pp1092) prepared at 2 mg/mL were compared by measuring the time to reduce
turbidity of living *B. subtilis* SL4 cultures by 50%.
965

966
967
Figure 6. Stability of anti-TcdB VHHs to gastrointestinal proteases. A) The percentage of TcdB-
binding activity of pp1005, pp1006, and pp1007 measured by ELISA after digestion with trypsin
(T), chymotrypsin (C), and elastase (E). Two concentrations of each protease were tested to
simulate either "fed" (higher concentration) or "fasted" (lower concentration) conditions. Data
represent the average of six independent replicates, the error bars representing the standard
deviation. B) TcdB-neutralizing activity of protease digested pp1005, pp1006, and pp1007. VHH
homodimers treated with chymotrypsin for either 10 or 60 minutes were compared in a cell-
rounding assay with the corresponding undigested protein. Each was diluted to a fixed
concentration (pp1005 was 0.05 nM, pp1006 was 5 nM, and pp1007 was 5 nM) and incubated
with serially diluted TcdB-10463. Each mixture was assayed for its effect on cell adhesion. After
digestion all exhibited neutralizing activities similar to undigested controls.
979

980
981
Figure 7. Stability of anti-*C. difficile* lysin to gastrointestinal proteases. A) Full-length lysin
(pp1093, 37 kDa), its catalytic subdomain (pp1092, 25 kDa), and a protease sensitive VHH
(pp555, 55 kDa) were digested with trypsin (T), chymotrypsin (C), and elastase (E) at pH 6.4 for
1 hour at 37 °C, quenched with protease inhibitors, the products separated by size using PAGE,
and peptides visualized by Coomassie staining. Mock digestions without target proteins were
performed to visualize the contribution by the proteases and quenching reagents. B) The image
in A (representative of six independent digestions) was quantified by densitometry with error
expressed as SEM. C) Activities of pp1092 digests were measured by determining the time to
reduce the turbidity of a culture of live *B. subtilis* SL4 by 50%. This value was then plotted
against a standard curve of undigested reference lysin to determine the activities of the
digested samples compared to undigested reference lysin. The dotted line represents the
activity of lysin when undigested. Error bars represent the standard error calculated from three
technical replicates.

995
996
Figure 8. Mouse challenge study. Shown are A) fecal spore counts 24 hours after infection and
at the time of euthanasia; B) maximum weight loss for individuals in each group; and C) Kaplan-
Meier representation of survival. Individuals were euthanized upon losing ≥15% of their initial
weight. Individuals in panels A and B are represented as a point and the group median,
interquartile range (IQR) and 1.5*IQR as boxplots. A) Log_{10} fecal shedding data were fit with a
linear model with group as a predictor per time point. Treatment group is a significant predictor
($p<0.05$) of shedding relative to the wild-type spirulina group (control/intercept) for Cocktail 1
and vancomycin. Cocktail 2 fails significance testing ($p\sim0.1$ and $p\sim0.25$). Statistic reports the t
ratio based on a Student t reference distribution, and p-value reports the two-tailed p-value
corresponding to the t ratio based on the Student t reference distribution. B) Maximum animal

weight loss was fit with a linear model with treatment as predictors of weight loss. Treatment
group is a significant predictor (p<0.05) of animal weight loss relative to the wild-type spirulina
group (control). C) Kaplan-Meier survival curves for mice infected with 10^5 *C. difficile* spores and
treated with vancomycin (orange, n=10), wild-type spirulina (black, n=5), spirulina therapeutic
Cocktail 1 (teal, n=5) or Cocktail 2 (blue, n=7). Survival rates were compared to treatment with
wild-type spirulina by log-rank test. Significance was observed with vancomycin treatment
(p=0.0068), borderline significance was observed with spirulina Cocktail 1 (p=0.080) and
Cocktail 2 (p=0.14) treatment.
1015

1016
1017
Figure 9. Hamster challenge trial. A) The Lumen therapeutic prolonged survival times when
compared to the untreated and group administered wild-type spirulina. Clindamycin-treated
Golden Syrian hamsters were administered phosphate-buffered saline, wild-type spirulina, or
the Lumen therapeutic two times a day starting one day before infection with *C. difficile*
(15,000 spores) and continuing until they reached their humane endpoints or the end of the
study (14 days post-infection). The mean survival times were compared using a Mann-Whitney
test with continuity correction, 90% confidence level, one-sided test. B) Lumen therapeutic
decreased *C. difficile* cecal colonization. Cecal contents were collected after animals reached
their humane endpoints or the termination date. *C. difficile* colony counts were quantified by
comparing RT-PCR measurements of extracted DNA from these samples as compared to a
standard curve generated by DNA extracted from vegetative *C. difficile* cells. Fewer *C. difficile*
cells colonized the cecum in hamsters treated with the Lumen therapeutic than in wild-type
spirulina-treated or untreated groups. Wilcoxon rank sum test (Mann-Whiney test) with
continuity correction, 90% confidence level, one-sided test.
1032

Parameter	Estimate	Std. error	p-value	2.5% CI	97.5% CI
TcdB IC_{50} (fM)	1.268	6.57E-05	2.24E-52	0.001138	0.001397
$Hill^{TcdB}$	1.495	0.07091	1.35E-58	1.355	1.634
EC_{50}^{pp1005} (pM)	21.44	0.002423	1.28E-16	0.01667	0.02622
$Hill^{pp1005}$ (-)	1.813	0.1245	1.12E-35	1.567	2.058
E_{max}^{pp1005} (fold)	45.25	3.076	3.20E-36	39.2	51.31
EC_{50}^{pp1006} (pM)	54.13	0.003761	4.25E-35	0.04673	0.06154
EC_{50}^{pp1007} (pM)	69.26	0.004852	1.11E-34	0.05971	0.07882
Bottom (assay units)	-0.01326	0.006172	0.0326	-0.02541	-0.001107
Top (assay units)	0.9289	0.01145	3.39E-189	0.9064	0.9515

1033
1034
Table 1. EC_{50} values for pp1005, pp1006, and pp1007. *Independent* model estimates from
analysis of the overall data set. TcdB1 acts as an inhibitor of cellular adhesion with femtomolar
potency (IC_{50}, H). pp1005, pp1006, and pp1007 act to inhibit TcdB1 activity, shifting TcdB1
potency 2-fold at 10's of picomolar potency (EC_{50}^{pp1005}, EC_{50}^{pp1006}, EC_{50}^{pp1007}). pp1005 activity
saturates at around 50-fold (E_{max}^{pp1005}, $Hill^{pp1005}$) while the somewhat less potent pp1006 and
pp1007 do not saturate.