The first sheep graph-based pan-genome reveals the spectrum of structural variations and their effects on tail phenotypes

Ran Li[1#], Mian Gong[1#], Xinmiao Zhang[1#], Fei Wang[1#], Zhenyu Liu[1#], Lei Zhang[1#],

Mengsi Xu[2], Yunfeng Zhang[2], Xuelei Dai[1], Zhuangbiao Zhang[1], Wenwen Fang[1], Yuta

Yang[1], Huanhuan Zhang[1], Weiwei Fu[1], Chunna Cao[1], Peng Yang[2], Zeinab Amiri

Ghanatsaman[3], Niloufar Jafarpour Negari[4], Hojjat Asadollahpour Nanaei[1,4],

Xiangpeng Yue[5], Yuxuan Song[1], Xianyong Lan[1], Weidong Deng[6], Xihong Wang[1],

Ruidong Xiang[7], Eveline M. Ibeagha-Awemu[8], Pat (J.S.) Heslop-Harrison[9], Johannes

A. Lenstra[10], Shangquan Gan[2]*, Yu Jiang[1]*

[1]Key Laboratory of Animal Genetics, Breeding and Reproduction of Shaanxi

Province, College of Animal Science and Technology, Center for Functional

Genomics, Institute of Future Agriculture, Northwest A&F University, Yangling,

Shaanxi, 712100, China

[2]State Key Laboratory of Sheep Genetic Improvement and Healthy Production,

Xinjiang Academy of Agricultural and Reclamation Sciences, Shihezi, Xinjiang

832000, China

[3]Department of Animal Science, Fars Agricultural and Natural Resources Research

and Education Center, Agricultural Research, Education & Extension Organization

(AREEO), Shiraz, Iran

[4]Department of Animal Science, Shahid Bahonar University of Kerman, Kerman, Iran

[5]State Key Laboratory of Grassland Agro-ecosystems, Key Laboratory of Grassland

Livestock Industry Innovation, Ministry of Agriculture and Rural Affairs, Engineering

Research Center of Grassland Industry, Ministry of Education, College of Pastoral

Agriculture Science and Technology, Lanzhou University, Lanzhou, 730020, China

[6] Faculty of Animal Science and Technology, Yunnan Agricultural University,

Kunming, 650201, China

[7]Faculty of Veterinary & Agricultural Science, The University of Melbourne,

Parkville, 3052, VIC, Australia

[8]Agriculture and Agri-Food Canada, Sherbrooke Research and Development Centre,

Sherbrooke, QC J1M 0C8, Canada

[9]Department of Genetics and Genome Biology, University of Leicester, Leicester, UK

[10]Faculty of Veterinary Medicine, Utrecht University, Utrecht, The Netherlands

[#]These authors contributed equally.

*To whom correspondence should be addressed.

Email: yu.jiang@nwafu.edu.cn; shangquangan@163.com

Abstract

Structural variations (SVs) are a major contributor to genetic diversity and phenotypic

variations, but their prevalence and functions in domestic animals are largely

unexplored. Here, we assembled 26 haplotype-resolved genome assemblies from 13

genetically diverse sheep using PacBio HiFi sequencing. We constructed a graph-

based ovine pan-genome and discovered 142,422 biallelic insertions and deletions,

7,028 divergent alleles and 13,419 multiallelic variations. We then used a graph-based

approach to genotype the biallelic SVs in 684 individuals from 45 domestic breeds

and two wild species. Integration with RNA-seq data allows to identify candidate

expression-associated SVs. We demonstrate a direct link of SVs and phenotypes by

localizing the putative causative insertion in *HOXB13* gene responsible for the long-

tail trait and identifying multiple large SVs associated with the fat-tail. Beyond

generating a benchmark resource for ovine structural variants, our study highlights

that animal genetic research will greatly benefit from using a pan-genome graph

rather than a single reference genome.

Keywords: sheep pan-genome, structural variations, *de novo* assembly, tail

morphology, *HOXB13*

Introduction

Structural variations (SVs) range from 50 base pairs (bp) to over megabases (Mb) in

size and are a major source of genetic variation. Compared to single-nucleotide

polymorphisms (SNPs), SVs could cause larger-scale genomic perturbations in genes

and regulatory regions and are therefore more likely to affect gene expression and

phenotypes[1, 2].

However, SV detection has been most challenging[3, 4]. Identification based on

short reads (≤ 150 bp) has a poor sensitivity and a high false discovery rate[5]. The long

reads of >10 kb by PacBio SMRT sequencing and Oxford Nanopore sequencing

greatly facilitated the efficacy of SV discovery by directly spanning the breakpoints[6, 7,

8, 9, 10, 11]. However, a sequencing error rate of 5-15% of the conventional long reads

impede downstream analysis and is particularly problematic for analysis of highly

repetitive regions. Another main restraint in efficient SV detection lines in the

reference genome model. The pan-genome graph model has been shown to much

superior to the linear reference assembly in representing the genome diversity and

thereby facilitating analysis of large genomic variations[12, 13, 14, 15, 16].

The novel PacBio circular consensus long-read sequencing (CCS) offers high

fidelity (HiFi) reads of 15 to 25 kb with >99.9% base accuracy[17], which now is

revolutionizing the *de novo* assembly and SV detection. It is expected to foster the

development of pan-genome applications by delivering haplotype-resolved assemblies

to depict the sequence diversity of the diploid genome[18]. With the availability of

population-scale genome assemblies, the pan-genomic reference models will allow to

determine faithfully the structure of complex regions for reliable SV detection[7, 12, 19].

Sheep are among the first domesticated livestock with economic and cultural

importance. Previous studies have shown that SVs determine the white coat color[20],

fleece variations[21] and polledness[22] in sheep – characters that have been under

selection for millennia. Nevertheless, a population-scale characterization of genetic

variation characterized by SVs remains to be explored in sheep as in most farm

animal species. In this study, we generated reference-grade *de novo* assemblies for 13

representative sheep breeds, together with two sets of phased assemblies for each

individual taking advantage of the Pacbio HiFi sequencing. These datasets allow us to

construct a graph-based ovine pan-genome and to explore the SV landscape with

unprecedented accuracy and sensitivity. We further analyzed the SV spectrum in 684

domestic and wild sheep from short-reads sequencing and highlighted SVs as largely

unexplored contributors to variations in phenotypes and gene expression.

Results

High-quality *de novo* assemblies of 13 representative sheep

Domestic sheep could be generally divided into at least four genetic groups including

European, Asian, Middle East and Africa with a separate position of the crossbred

Dorper sheep[23, 24, 25]. We selected 13 representative sheep (Fig. 1a) from seven

European, four East Asian, one Middle Eastern (Kermani from Iran) and one African

(White Dorper) breeds. The European breeds include meat (Suffolk, Dorset, Texel,

Charollais), wool (Merino, Romney) as well as dairy (East Friesian) sheep, whereas

the East Asian accessions represent the three lineages of Northern China (Ujumqin,

Kazak), Qinghai-Tibetan Plateau (Tibetan) and Yunnan-Kweichow Plateau (Yunnan)[24].

A SNP analysis showed that the 13 sheep encompass 95.5% of the SNPs with minor

allele frequency (MAF) >0.05 among the studied 643 domestic sheep worldwide.

The 13 sheep were sequenced using PacBio HiFi sequencing at an average

coverage of 20.9×, yielding highly accurate long reads with a per-base

accuracy >99.9%. We used the Hifiasm assembler[18] for *de novo* assembly, which

generated the primary contigs as well as two sets of phased contig assemblies from

HiFi reads. We ultimately assembled 13 primary collapsed assemblies with an average

length of ~2.86 Gb (Table 1). The average contig number were 1582 with contig N50

lengths of 68.2 Mb. The most complete assembly (FRI1) achieved a contig N50 of

85.1 Mb, which is much higher than for the sheep reference genome (ARS-

UI_Ramb_v2.0). The HiFi sequencing depth is not correlated with assembly length

(r_s=-0.04, Spearman rank correlation), but with contig N50 length (r_s =0.93) and

negatively with contig N50 number (r_s =-0.67). We further ordered and positioned

~93.6% of the contigs from each of the 13 collapsed assemblies into chromosomes

based on their synteny to the reference genome. The resulting chromosome-level

collapsed assemblies for the 13 representative breeds may very well serve as a

reference panel for future research in sheep genetics and genomics. In addition, the

haplotype phased assemblies achieved an average contig N50 length of 14.5 Mb,

which captures per diploid individual almost all sequence variants and was used for

the subsequent construction of the multiassembly graph.

On the basis of Benchmarking Universal Single-Copy Orthologs (BUSCO)

analysis, the collapsed assemblies displayed an average completeness of 94.1%

(93.8%-94.4%, Table 1). BUSCO scores for the partially phased assemblies were

lower but still consistently >90% complete. To measure the quality value (QV) of the

assemblies, the 13 sheep were sequenced to an average of 30× depth Illumina short-

reads, which showed for all the collapsed and phased assemblies except WDU1

higher values than obtained for the sheep reference genome assembly (QV=42.5).

Construction of sheep graph-based pan-genome and SV detection

To build a comprehensive sheep graph-based pan-genome, we used the reference

genome (ARS-UI_Ramb_v2.0) as the backbone. The 26 haplotype-resolved

assemblies from the 13 individuals were individually added to the graph to encompass

the sequence variations more comprehensively than the primary contigs. A multigraph

genome was generated using minigraph which is a state-of-the-art graph constructor

that especially is suitable for discovering complex variations[26, 27].

The resulting multiassembly graph spans 2.75 Gb and includes 167,565

nonreference nodes of 137.7 Mb. Based on the presence or absence of each node

among the reference genome and the 26 haplotype-resolved assemblies, we divided

the pan-genome into three categories (Fig. 1b): core (nodes present in all assemblies,

69.6%), flexible (nodes present in at least two assemblies, 29.0%) and private (nodes

present in one assembly, 1.4%). The variations were most associated with LINEs

followed by VNTR and mixed interspersed repeats (Fig. 1c).

The genome graph represents SVs as bubbles that are located between a single

source and a single sink node coming from the reference assembly. We classified the

SVs into three categories: (1) biallelic insertions and deletions; (2) divergent alleles

with a non-reference node; (3) multiallelic variations with more than one non-

reference path (Fig. 1d and 1e, Supplementary Data 1 and 2). The genome graph

revealed 71,401 biallelic insertions and deletions, 7,028 divergent alleles and 13,419

multiallelic variations. However, despite advantage in recovering divergent alleles and

multiallelc variations, the genome graph constructed by minigraph is not well adapted

for the discovery of small SVs[18] and thus we excluded those below 100 bp. As a

complementary approach to detect insertions and deletions, we used pbsv

(https://github.com/PacificBiosciences/pbsv) to jointly call among the 13 individuals

SVs using the PacBio HiFi reads, yielding 142,279 insertions/deletions. We observed

much more insertions than deletions, contrasting with results based on short-read

sequencing which preferentially identifies deletions[28]. The pbsv call set covers 97.5%

of the insertions and deletions found by minigraph. Therefore, we merged the pbsv

call set and the large variants (>5 kb) specifically found by minigraph, resulting in

142,422 insertions and deletions, which together with the identified divergent alleles

and multiallelic variations constituted a final comprehensive catalogue of SVs in the

sheep genome (Fig. 1d and 1e, Supplementary Data 3).

The number of insertions/deletions rapidly decreases with length, consistent with

reports in humans using long-read sequencing[6, 7, 29]. Noticeable peaks at 150 bp

correspond to SINES, 1,300 bp for LTRs, as well as 7,750 bp for LINEs (Fig. 1d). The

median length of divergent alleles (726 bp) and multiallelic loci (575 bp) are longer

than that of biallelic variations (230 bp).

The insertion sequences were predicted to harbor 588 genes whereas the deletions

potentially affect the exonic region of 1920 genes by containing exons. The

nonreference sequences of the divergent alleles contained 31 genes. We revealed the

full sequence of one divergent locus containing *MYADM-like* repeat (Fig. 1f), which

was previously reported to be associated with mean corpuscular hemoglobin

concentration traits and weight of lamb weaned but the sequence was unknown[30]. We

also noticed several other large divergent alleles that likely to correspond to

phenotypic variants (Supplementary Fig. 1). The non-reference alleles of the

multiallelic loci were predicted to harbor 146 genes. One of the largest multiallelic

region represented a deletion of 105.6 kb which is nested with an insertion of 5.5 kb

(Fig. 1g). The deletion likely results in the loss of the intelectin genes, that displays

extremally high expression in lung of sheep and potentially plays a role in mucosal

response in allergy[31]. Additional examples of large multiallelic loci that resulted in

gene gain/loss, included *HBB*, *ASIP*, *UDP-glucuronosyltransferase 2B18-like*,

olfactory receptor (*LOC101115700*), etc. (Supplementary Fig. 2).

The SVs are underrepresented in untranslated region (UTR), coding sequences

(CDS), and regulatory elements (H3K4me3, H3K27ac), suggesting strong negative

selection that acted against them (Supplementary Fig. 3). We also reported 201 SV

hotspots sites covering 18.2 Mb of the autosomes (Supplementary Fig. 4). The SVs

were significantly clustered at the terminal 5 Mb region of each chromosome with

4.1-fold enrichment (P value<0.01, permutation test) (Supplementary Fig. 5).

Frequency spectrum of SVs in the sheep populations

In order to better understand the frequency spectrum of the SVs within sheep, we used

the graph genotyping software Paragraph[32] to genotype the biallelic insertions and

deletions in autosomes in a panel of 684 resequenced samples from 45 domestic

breeds or populations and two wild ovine species. The samples included 643 domestic

sheep from Europe, East Asia, Middle East and Africa, 33 Asiatic mouflons and 8

argali with an average coverage of 18.1× (Fig. 2a, see Methods for the assignment of

the breeds and populations. Paragraph represent SVs as diverging paths in a graph and

achieves a high rate of genotyping accuracy[33].

To ensure a high-quality set of autosomal SVs for population genetics analysis,

we excluded 17,330 SVs that failed to be genotyped in >90% of the samples and

another 2,011 that by an excess of heterozygotes deviated from Hardy Weinberg

equilibrium, leaving 116,776 (85.6%) reliably genotyped SVs for downstream

analysis. The frequency of the SVs was provided in Supplementary Data 4. We also

estimated the genotyping accuracy of the remaining SVs by measuring the Mendelian

errors, which represent genotypes that are found in the offspring but could not be

inherited form either parent. The Mendelian error rate was 3.8% on average in the 11

trios of 29 individuals (each ~30× coverage) used in our pervious study[34].

PCA analysis using either insertions or deletions displays similar population

structure and shows a clear separation of the four genetic groups (Supplementary Fig.

6), agreeing with the clusters defined by SNPs. Therefore, the SV catalogue

reproduces the genetic diversity of the worldwide sheep population.

The level of linkage disequilibrium (LD) between SVs and SNPs could reflect the

extent of SVs representing independent and undiscovered genetic variations relative

to previous evolutionary studies. We estimated how frequently the SVs were linked to

the flanking SNPs (Fig. 2b). Around 39.8% SVs link strongly with surrounding SNPs

(rank>200), the characteristic of which could be detected by the surrounding linked

SNPs. It is worth noting that there are still 39.2% SVs display low levels of LD with

nearby SNPs (rank<100), implying variants that are poorly tagged by easily

genotyped SNP markers.

In order to infer the allele state (ancestral or derived) for each SV, we additionally

genotyped one snow sheep (*Ovis nivicola*) and one bighorn sheep (*Ovis canadensis*)

and, together with one argali (*Ovis ammon*) as outgroup, and identified the state for

115,089 SVs and a variable allele frequency of derived alleles (DAF) in domestic

sheep (Fig. 2c). About 28% of the derived alleles occurs only in domestic sheep (Fig.

2d) and may have reached a high frequency by domestication and improvement. The

proportion of *de novo* SVs may be overestimated due to the relatively limited number

of wild sheep but this does not explain a high geographic differentiation of 592 high-

frequency *de novo* SVs (DAF≥0.5 in at least one geographic group (Fig. 2e). Those

SVs were found to be enriched in KEGG pathways of Ras signaling and long-term

depression (adjusted *P* value<0.05). For example, we found in all regions a *de novo*

3.4 kb insertion in the first intron of *GRID2* (Fig. 2f), which plays a key role in

synaptogenesis and neurodevelopment[35]. A 1.4 kb *de novo* insertion prevalent in

European sheep was found at the 3' UTR of *IRF2BP2* (Fig. 2g), which was reported

as the causative mutation for the wool coat[21].

Population-stratified SVs and their potential effect on gene expression

After initial domestication at the Fertile Crescent, domestic sheep dispersed

worldwide and adapted to a wide range of natural and artificial selection. In this

context, we detect putative selection signals by dissecting population-stratified SVs

among the selected 33 out of the 45 breeds or populations to ensure a similar number

of breeds or populations for European and East Asian sheep. For the differentiation

between these groups, we used a modified calculation of global F_{ST} (see Methods).

Because this strategy tends to discover common selection signals present in multiple

breeds, the population-stratified SVs differentiating Europe and East Asian breeds

were preferentially detected since they are in our panel the predominant breeds.

We identified 865 population-stratified SVs with the top 1% global F_{ST} values

(Supplementary Data 5), 351 of which intersect with genic regions and 324 are within

100 kb of gene flanking region (Fig. 3a). The majority of them (698, 80.7%) were

confirmed by the Ohana approach[36] with which we detected selection signals in

multiple populations by varying the number of ancestral admixture components. The

related genes are associated with diverse phenotypic and production traits, including

wool type (*IRF2BP2*, *FGF7*)[21, 37], horn type (*RXFP2*)[38], fat deposition (*BMP2*,

PDGFD, *PDGFA*)[23, 39], and coat color (*KIT*)[40], etc. Notably, the highest stratified SV

(F_{ST}=0.78) is the 1.4 kb insertion downstream of *IRFBP2* (Fig. 2g), differentiating

European and Asian breeds. Functional annotation of the *top* F_{ST} signals reveals

enrichment in several KEGG pathways, such as those involved in signaling pathways

(Rap1, MAPK) and parasitic disease (African trypanosomiasis, amoebiasis)

(Supplementary Fig 7).

SVs may impact the gene expression of neighboring genes via the cis-regulatory

elements[10, 41, 42]. To explore SV-associated expression changes, we used 112 RNA-seq

data from 8 tissues of 14 hybrids of Texel×Kazak sheep as well as WGS data of the

same sheep[34, 43] (see Methods). This unique combination of data enabled us to identify

the candidate expression-associated SVs by allelic-specific expression (ASE)

mapping. By ASE mapping, we found 1,141 heterozygous SVs that would putatively

act as cis-regulatory elements and thus cause allelic-specific expression in associated

genes (Fig. 3b and 3c, Supplementary Data 6). These expression-associated SVs were

significantly enriched in generic regions (P value<2.2e-16, permutation test), and 34

overlapped with population-stratified SVs which is greater than random chances (P

value<1.1e-10, Chi-Square test). We additionally performed SV-eQTL mapping and

found 260 SVs with genotypes that correlate with expression levels of a nearby gene

(Supplementary Data 7).

SVs as strong candidates affecting sheep tail morphology

The tail morphology of sheep is a key genetic trait that is related to domestication,

adaptation, and animal welfare[44]. The wild ancestors (Asiatic mouflon) have a short

thin tail whereas four domestic tail types include in addition to the primitive type the

long thin-tailed, the short fat-tailed (including fat-rumped) and the long fat-tailed

types. Although several genes have been found to be associated with these traits[23, 45,

46], the underlying causative mutations are still unknown. However, the long and fat

tails are derived traits, indicating that the causative mutations are likely to be *de novo*

variants.

To identify SVs associated with tail length, we compared the long and short tailed

breeds of (1) the Asian long fat-tailed breeds versus Asian short fat-tailed breeds from

Asian breeds and (2) the European long thin-tailed breeds versus the North European

short thin-tailed breeds, respectively. Notably, in both independent comparisons the

most differentiated SVs point to an insertion of 169 bp (chr11:37525005) close to the

5' UTR of *HOXB13* (Fig. 4a). This resides in a specific haplotype exclusively found

in long tailed sheep, where the insertion is prevalent in long-tailed breeds and rare in

short-tailed breeds (Fig. 4b and 4c). *HOXB13* is mainly expressed in the embryo and

colon of sheep and cattle[47]. We found the expression of *HOXB13* in colons of the

short-tailed Texel×long-tailed Scottish Blackface individuals presumably carrying at

least one copy of the insertion[48], and in colons of individuals with short tail

presumably without the insertion (Texel sheep and small-tail sheep)[49, 50]. The insertion

was only expressed in the colons of the Texel×Scottish Blackface but not in Texel

sheep and small-tail sheep (Fig. 4d and Supplementary Fig. 8).

In order to verify the effect of this insertion on tail length, we evaluated the

association between the insertion and tail length in a population of 201 sheep of the

same age (nine months old). The three genotypes including homologous insertion (II),

heterozygote (ID) and homologous deletion (DD) were directly distinguished through

PCR amplification and agarose gel electrophoresis (Fig. 4e). Despite lacking the DD

individuals in the studied population (see Methods), the tail length of the homologues

(II) was found to be significantly longer than that of the heterozygotes (ID) by 2.77

cm on average (Mann-Whiney U test, P value<0.0001) (Fig. 4f). To validate the

function of the insertion, we performed luciferase reporter assays. Relative to the

wildtype (WT), the mutant with the insertion decreased luciferase activity by 10-fold

(Fig. 4g), indicating that the insertion reduces the transcriptional level of *HOXB13*.

The gene has been previously identified as a repressor of caudal vertebra

development[51, 52]. Therefore, the insertion is likely to increase the tail length of sheep

by depressing the expression of *HOXB13*.

To identify candidate SVs related to the fat tail, we compared the genomes of

eight fat-tail breeds (SFK, KAZ, HUS, UJM, LTH, TAN, BSB, SSS) and ten thin-tail

breeds (FIN, DEM, EFR, GTL, SEL, SFK, SOG, POD, TIB, YBB). We applied the

population branch statistic (PBS) to identify positive selection in fat-tail sheep using

Asiatic mouflon as an outgroup. The two most differentiated regions are identified by

either SV or SNPs (Fig. 5a and 5b), one located at the intergenic region between

BMP2 and *HAO1* (hereafter referred to as IBH region) and the other in *PDGFD*.

Notably, the most stratified SVs in these two regions are among the top selective

variants indicated by both SVs and SNPs. The most differentiated SV cluster are

located at IBH region which has also reported as a candidate selective region affecting

fat tail trait[53, 54]. Notably, six highly differentiated SVs including one *de novo* SV were

identified in the IBH selective region (Fig. 5c), of which the largest one is a 7728 bp

insertion. *PDGFD* has been recognized as the most plausible candidate gene for the

fat-tail phenotype in recent studies[23, 55]. We found three highly differentiated SVs

residing in the selective region of *PDGFD*, of which the largest event is a 867 bp

insertion. The three selective variants are all domestic *de novo* SVs. Therefore, our

results suggest that both regions are associated with the fat-tail phenotype and the SVs

identified within these regions are the most plausible candidate causative mutations.

Discussion

The genome-wide analysis of genomic structural variations in diverse breeds is

critical to understanding the full repertoire of genetic diversity and pinpointing their

associations with phenotypic traits. However, previous resequencing studies mainly

rely on mapping short reads to a linear reference genome, precluding the discovery of

regions with pronounced structural variations[4]. The graph-based pan-genome model

largely reshapes the methodology to assess the whole landscape of sequence diversity,

and is likely to transform the genomic research field. To build a high-quality ovine

pan-genome, we utilized the PacBio HiFi sequencing to generate a panel of reference-

level assemblies and 26 phased haplotypes for 13 representative sheep breeds,

providing a solid foundation for in-depth functional genomic studies. The long and

accurate HiFi reads guarantee the high-standard of our pan-genome through

generation of high-quality *de novo* assemblies, preservation of the two haplotypes and

precise assignment of SV' breakpoints. The development of this ovine pan-genome

will contribute to shifting in animal genomics research from a linear reference

genome to a pan-genomic model comprising multiple reference-quality representative

genomes.

The application of graph-based pan-genome has its clear opportunities and

obstacles. The pan-genomic approaches are in practice not so straightforward as the

linear reference genome and the methodologies and implementations are still in active

development[12, 14, 56, 57]. For example, minigraph is one of the most promising graph

constructors, but has limitations in performing base alignment, generating relatively

high false positives for small SVs[26]. Nevertheless, a shift to the graph-based pan-

genomes model will enhance the opportunities to systematically investigate

previously inaccessible genomic variations. In our study, we found that the minigraph

performed very well in discovering divergent alleles and multiallelic variations, which

accounted for 27.4% of the total variations in length but are previously neglected.

Combining the graph-based pan-genome with read-based calling yielded a reference

panel of high-quality SVs for downstream functional genomic studies.

We also demonstrated how these pan-genome datasets can be utilized to link SVs

to phenotypes by genotyping and assessing the SV landscape in a large population

from short-read sequencing. We genotype SVs using the graph-based Paragraph,

which enables efficient encoding of structural variation into the pan-genome[32]. The

population-scale study allows us to trace their responses to natural and artificial

selection. Taking the tail morphology traits as an example, we identified a plausible

causative mutation in *HOXB13* for the long-tail trait, which illustrates the key roles of

SVs underlying phenotypic changes. We further reported segregating SVs in fat-tailed

breeds. The observation that *de novo* SVs showed higher or comparative selection

signals than surrounding SNPs warrants further investigation.

In conclusion, the pan-genome representation with an influx of high-quality

assemblies will revolutionize our understanding of SV prevalence and their biological

significance, giving a new dimension to the nature and extent of genomic variation,

beyond SNPs, impacting traits.

Methods

Sample collection and PacBio sequencing

Samples were selected to for an optimal representation of the global sheep diversities.

The sampling information were provided in Supplementary Table 1. High-quality

genomic DNA (gDNA) was extracted from fresh blood samples as previously

described[58] and assessed for purity and quantity using Nanodrop 1000 (Thermo Fisher

Scientific, CA, USA) and Qubit (Thermo Fisher Scientific, CA, USA) assays. Library

with an average insert size of ~15 kb was generated using the SMRTbell Express

Template Prep Kit 2.0 (Pacific Biosciences, CA, USA) and fractionated on the

SageELF (Sage Science, Beverly, MA, USA) into narrow library fractions. Libraries

were then sequenced on 2-3 SMRT Cells 8M on a Sequel II instrument (Pacific

Biosciences, CA, USA) using 30-hour movie times at Annoroad Gene Technology

Co., Ltd (Beijing, China). Raw data was processed using the CCS algorithm (version

6.0.0, parameters: --minPasses 3 -- minPredictedAccuracy 0.99 –maxLength 21000)

to generate highly accurate HiFi reads.

De novo **assembly using HiFi reads and quality assessment**

Hifiasm v0.15.3-r339 was used to generate the assembly from HiFi CCS reads using

default parameters[18]. Hifiasm yields one primary contig assembly and two sets of

partially phased contig assemblies. The primary contig assembly for each breed was

scaffolded to chromosome level using RagTag v2.0.1 with the reference sheep

genome assembly[59].

The genome completeness was assessed using the BUSCO program (v.3.0.2)

containing the Mammalia odb9 set of 4,104 genes[60]. The single plus duplicated

complete BUSCO gene counts are reported. The base accuracy was measured by

assembly quality value (QV) using yak (https://github.com/lh3/yak), which compares

the 31-kmers found in short-reads and the assembly sequences.

Pan-genome construction

The pan-genome construction was performed using a previously reported workflow[15]

with minor modifications. Minigraph v0.15-r426 with option -xggs was used to

generate the graph pan-genome[26]. The sheep reference genome (ARS-UI_Ramb_v2.0,

GCF_016772045.1) was used as the backbone of the multiassembly graph and the 26

partially phased contig assemblies were added successively. Only autosomal

sequences were considered in the multigraph genome, as the X and Y chromosomes

of males were underrepresented.

Identification of non-reference divergent alleles and multiallelic alleles from the

pan-genome graph

The graph generated by minigraph is composed of chains of bubbles with the

reference as the backbone. We first used gfatools v0.5-r234 which is a bubble popping

algorithm to extract bubbles from the graph (https://github.com/lh3/gfatools). Each

bubble represents a structural variation, encompassing the start and end nodes of

reference sequences as well as path traversing the start and end nodes.

Depending on the number of paths in a bubble, the structural variations were

classified into three categories: (1) biallelic variations (insertions/deletions): either the

reference path is longer (deletions) or the non-reference path is longer (insertions).

The node in the shorter path is required to be below 100 bp. (2) Divergent alleles: the

reference nodes and the non-reference nodes were both longer than 100 bp. (3)

Multiallelic alleles: one reference path and more than one nonreference path were

present. Those nonreference paths below 100 bp were excluded.

Genome annotation of non-reference sequences

We used RepeatMasker v4.0.5 (http://www.repeatmasker.org) to softmask the non-

reference sequences. Then we performed an ab initio gene structure prediction using

Augustus v3.3.3[61] and searched the protein sequence predicted by Augustus against

the local protein database using DIAMOND[62] blastp (parameters: --more-sensitive --

evalue 1e-10 --max-target-seqs 1) and kept hits with minimum coverage of 70% and

identity of 80%. The local protein database was built using DIAMOND[62] makedb

with RefSeq protein sequences from sheep, goat and cattle.

Annotation of repeats

Interspersed repeats and low complexity DNA sequences were identified using

RepeatMasker v4.0.7 (http://www.repeatmasker.org) with a combined repeat database

including Dfam v.20170127 and RepBase v20170127 with parameters: -species

Ruminantia -xsmall -s -no_is -cutoff 255 -frag 20000 -gff. Tandem repeats were

identified using Tandem Repeat Finder v4.09[63]. The identified tandem repeats as well

as those low complexity sequences identified by RepeatMasker were merged as low-

complexity regions (LCR). Furthermore, LCR with unit motif lengths ≥7 bp were

classified as VNTR and those with unit motif lengths ≤6 bp were classified as STR.

Structural variant calling

We integrated the results from graph-based pan-genome analysis and alignment-based

SV calling to generated high-confidence SV calls. For pan-genome analysis, the SVs

were generated from our multiassebly graph genome using gfatools v0.5-r234 as

described above. For alignment-based SV calling, the SVs were detected using pbsv

tools v2.6.2 (https://github.com/PacificBiosciences/pbsv). The CCS reads were

aligned to the reference genome using pbmm2 to generate the alignment BAM file.

Then the pbsv discover module was used to identify signatures of structural variation

within each aligned BAM. The tandem repeat annotation file was provided to increase

the sensitivity and recall of this step. Finally, the signatures of all the samples were

fed to the pbsv call module to jointly call SV from structural variant signatures using

the –ccs mode.

Mendelian error rate estimation

A Mendelian inheritance error represents a genotype in the offspring that cannot be

inherited from either of their paraments. The whole genome sequencing data of 11

trios were from our recently study for allelic specific expression analysis[34]. For each

trio, the Mendelian error rate is the proportion of SVs from the children that deviate

from Mendelian inherence.

SV hotspot analysis

SV hotspot regions were determined using primatR package "hotspotter" section

(bw=200000, num.trial=1000, pval<1e-8)[64]. SVs were significantly clustered at the

terminal 5 Mb region of each chromosome (*P* value<0.01, permutation test, random

457   shuffling of 1,000 times). The random shuffling of SV intervals was performed using

458   BEDTools v2.29.0 with the shuffle command[65].

Enrichment of SVs in the genome

The H3K4me3 and H3K27ac peak regions were retrieved from our Ruminant genome

database. We only used insertions and deletions for enrichment. For insertions, the

end position is the start position plus the insertion length. We first recorded the

observed number of overlaps between SVs and UTR, exon, intron, H3K4me3 and

H3K27ac peak regions, respectively. Then we counted the average number of

overlaps between randomly shuffled SVs (1,000 times) and each genomic region.

SV genotyping in whole genome sequencing data

We collected whole-genome sequencing data for 643 domestic sheep, from four

geographic regions (Europe, East Asia, Middle East and Africa) as well as wild sheep

including 33 Asiatic mouflons (*Ovis aries musimon*) and 8 argali sheep (*Ovis ammon*).

For the domestic sheep, we selected breeds with at least five individuals and defined

populations by combining breeds from the same country with less than 5 individuals.

In this way, all the domestic sheep were assigned to 45 breeds or populations

(Supplementary Data 7). We genotyped SVs in these samples using Paragraph v2.4a[32].

The maximum permitted read count for each variant was set to 20 times the average

sample depth in order to reduce runtime for repetitive regions. The resulting genotype

files in vcf format from all samples were then combined using bcftools v1.9[66].

To obtain a high-confidence call set of SVs, the resulting genotypes were filtered

based on overall genotyping rates and Hardy-Weinberg equilibrium[33]. Those variants

that were successfully genotyped in ≥90% of all samples were retained using

vcftools[67] for further analysis. We also estimated Hardy-Weinberg equilibrium for

each variant using vcftools and those with excess of heterozygotes (P value< 1×10^{-5})

were removed.

Quantifying linkage disequilibrium (LD) of SVs with surrounding SNPs

In order to measure the extent to which SVs are correlated with SNPs, we calculated

LD (r^2 values) between the polymorphic SVs (MAF>0.05) and the neighboring

polymorphic SNPs (MAF>0.05; 150 upstream and 150 downstream SNPs) in

Mouflon and domestic sheep. SNPs located on SVs were not considered. Plink v1.9

was used to calculate LD for all pairwise cases for SV-SNP and the surrounding SNP-

SNP variant pairs. According to a previous study[41], we first obtained pairwise

genotype correlations (r^2) for all SNP-SNP and SNP-SV pairs. For each of the 300

ranked surrounding positions, the number of times the SV rank was greater than the

SNP-SNP median rank was recorded. Low, middle or high LD of each SV were then

classified if the relative LD metric of SV to SNP was≤ 100, **>**100 and ≤200, and >200,

respectively, respectively.

Inferring the derived SVs and *de novo* SVs in domestic sheep

To infer the state (ancestral or derived) for each SV, we genotyped one snow sheep

(12.11× depth, ERR4161992) and a bighorn sheep (10.3× depth, SRR501898), which

together with one argali sample served as outgroups. For each biallelic SV, if the same

homozygous genotypes were detected in at least two outgroups, this state was defined

as the ancestral state while the other was the derived state. If the derived allele is only

present in domestic sheep but absent in all the wild sheep (8 Argali and 33 Asiatic

mouflons), the SV was defined as a *de novo* SV in domestic sheep.

Detecting selection signals

To detect population-stratified SVs, we employed two strategies, the global F_{ST} and

the Ohana method, which identifies extreme allele frequency differences by modelling

ancestral admixture components[36]. The global F_{ST} was calculated following a

previously described approach with modifications[68]. For each SV site, we calculated

the F_{ST} values between all populations. Instead of using the average of all F_{ST} values,

we took the average of the top 50% F_{ST} values as the global F_{ST} value for each SV.

The top 1% of the highest global F_{ST} value by this method were considered as

putative selective signals. For the Ohana method[36], we first modeled the dataset with

the provided workflow using the SNPs[33]. The resulting covariance was used as the

neutral input for scans of selection. For each specific ancestry component, Ohana

reported a likelihood ratio statistic (LRS) to quantify the likelihood of selection for

each variant. We varied the number of ancestral admixture components from K=2 to

K=5 to find selection signals.

In order to reveal selection signals in fat-tailed sheep, we used the method of

population branch statistic (PBS) to calculate and compare the Fixation index (F_{ST})

between the fat-tailed sheep, thin-tailed sheep and Asiatic mouflon, using the Asiatic

mouflon as a distantly related population. The F_{ST} value between populations with

different traits was calculated using the python scikit-allel package v1.3.2

(https://github.com/cggh/scikit-allel). We calculated the F_{ST} values of both SNP and

SV sets. F_{ST} values for SNPs were calculated with a 10 kb sliding window with a 5 kb

step.

Detecting expression-associated SVs

The RNA-seq data of 8 tissues (liver, rumen, tail fat, skin, subcutaneous fat,

semitendinosus, semimembranosus, longissimus dorsi) from 14 Texel (♂)×Kazak (♀)

sheep hybrids with accession number PRJNA485657 as well as their whole genome

sequencing data were downloaded from NCBI Sequence Read Archive[34, 43]. The raw

reads were first trimmed for adapter and low-quality sequences using Trimmomatic[69]

and the high-quality reads were aligned to the sheep reference genome using Hisat2[70].

Reads counts per gene were computed using featureCounts [71] and normalized to

counts per million (CPM) using edgeR package[72] and those with CPM>1 in at least

two samples were retained.

It is expected that the regulatory SVs could cause allelic-specific expression. We

thus performed allelic specific expression (ASE) mapping by integrating the RNA-seq

data as well as the SV genotyping information for each of the samples. The ASE

genes were identified following the pipeline described in our previous study[34]. First,

for each sample, the ASE SNPs displaying allelic imbalance at the gene body were

identified from RNA-seq data using the following parameters: (1) at least 20 total

reads; (2) allele ratio (allele reads count/total reads count)> 0.65 or <0.35; (3)

significant allelic imbalances with FDR<0.05 (Chi-squared test). Those with at least

two ASE SNPs at the gene body and found in at least three samples were defined as

candidate ASE gene. For each SV-gene pair, if the ASE gene matches with the

heterozygous state for the associated SV in at least five RNA-seq samples, we

considered the SV as plausibly expression-associated.

The SV-based eQTL analysis was performed using the R package Matrix eQTL

v.2.3 [73]. The midpoint where SV starts and ends were defined as the position of each

SV. Cis-eQTL analysis was tested within each tissue against all SVs located within the

200 kb upstream and downstream of neighboring genes. The gender and population

structure (the first three principal components from PCA analysis of the 14 individuals)

were included as covariates to account for confounding effects using an additive

linear model. Statistical significance was considered with FDR<0.05.

Analyzing the association between the 169 bp insertion with the tail length

The association analysis was conducted in a F2 population which was generated by

backcrossing East Friesian sheep (♂) with the female Hu sheep (♀) × East Friesian

sheep (♂). A total of 211 individuals of the similar age (six months old) were used.

The tail length (cm) of each individual was measured and DNA samples were

collected. Since the insertion was dominant in East Friesian sheep with allele

frequency of 0.92 but is completely absent in Hu sheep based on our resequencing

data, the F2 population that we studied only possessed two genotypes (II and ID) due

to the grading-up using East Friesian sheep. Agarose electrophoresis was used to

determine genotypes of the PCR products. Primers were designed in the flanking

region of the insertion (forward: TTTATGAGCTTCTCTCCGCCA; reverse:

AAGTGGTATAATTGCCGGGCT). PCR amplification was performed by

denaturation of 94℃ for 2 min followed by 28 cycles of 94℃ for 30 s, 62℃ for 30 s,

and 72°C for 30 s and an extension of 72°C for 2 min; and a hold at 4°C.

Luciferase reporter assay

To analyze promoter activity of the SV variation by luciferase assay, fragments with

(mutation) and without (wild type) the insertion was synthetized by Tsingke

Biotechnology Co., Ltd. (Beijing, China). Each fragment was cloned into the pGL3-

Basic vector using Sac I and Hind□ sites to assay its promoter activity on the

downstream gene.

The 293T cells in 96-well plates were transiently transfected with 200 ng Firefly

luciferase vector (pGL3) and 4 ng pRL-TK Renilla luciferase vector (Promega,

Madison, USA) using 0.5 uL Lipofectamine 3000 (Thermo Fisher Scientific, Waltham,

USA). Twenty-four hours after transfection, luciferase activities were measured with a

dual-luciferase reporter assay kit (Promega, Madison, USA). Renilla luciferase

activity was used as the transfection control for normalizing Firefly luciferase activity.

The pGL3-Basic vector without insertion was used as the reference for promoter

activity assays and the pGL3-Control vector was used as the positive control for

promoter activity.

Author contributions

Y.J. and S.Q.G. conceived and designed the experiments. R.L., X.M.Z. and Z.Y.L.

performed *de novo* assembly and SV calling; Z.Y.L., M.S.X. and Y.F.Z. analyzed the

RNA-seq and Chip-seq data; F.W., M.G. and L.Z. contributed to the ASE analysis;

R.L., Z.Y.L., W.W.Fang and X.L.D. carried out SV genotyping and population

genetics analysis; Y.T.Y. and X.Y.L. conducted the luciferase reporter assay; Z.B.Z.,

W.W.Fu, H.H.Z., C.N.C., P.Y., Z.A.G., N.J.N., H.A.N., X.P.Y., Y.X.S. and W.D.D.

collected the samples and prepared the WGS data; R.L. wrote the manuscript and Y.J.,

S.Q.G., J.A.L., P.H., E.I.A., R.D.X. and X.H.W. revised it. All authors read and

approved the final manuscript.

Acknowledgements

We thank the High-Performance Computing platform of Northwest A&F University

for providing computing resources. We would also like to thank Prof. Xiaolong Wang

for providing the nice chart of different sheep tail patterns. This study was supported

by research grants from the National Natural Science Foundation of China to Y.J.

(U21A20247, 31822052), S.Q.G. (31760662, 31860639) and R.L. (31802027),

Natural Science Basic Research Program of Shaanxi (K3030121801), and China

Postdoctoral Science Foundation to W.W.Fu (No. 2021M702690).

Competing interests

The authors declare no competing interests.

Ethics statement

Blood samples were taken by conforming with the Helsinki Declaration of 1975 (as

revised in 2008) concerning Animal Rights.

Data availability

Sequences and metadata generated in this work are publicly available. The *de novo*

assemblies, HiFi data and WGS data for the 13 individuals are available at NCBI:

WDU1 (PRJNA788705, PRJNA788704, PRJNA784371),QK1 (PRJNA788680,

PRJNA788679, PRJNA784330), ROM1 (PRJNA788688, PRJNA788687,

PRJNA784357), SFK1 (PRJNA788697, PRJNA788696, PRJNA784366), UJM1

(PRJNA788702, PRJNA788701, PRJNA784376), CHA1 (PRJNA788666,

PRJNA788665, PRJNA784396), YUN1 (PRJNA788712, PRJNA788711,

PRJNA784413), TEX1 (PRJNA788214, PRJNA788213, PRJNA780104), DOR1

(PRJNA788571, PRJNA788570, PRJNA782680), KER1 (PRJNA788676,

PRJNA788675, PRJNA783516), MER1 (PRJNA788672, PRJNA788671,

PRJNA783537), KAZ1 (PRJNA788564, PRJNA788563, PRJNA783179), FRI1

(PRJNA791176, PRJNA791175, PRJNA721520). The details are provided in

Supplementary Table 1.

References:

1.   Chiang C, *et al.* The impact of structural variation on human gene expression. *Nat*

Genet **49**, 692-699 (2017).

2.   Sudmant PH, *et al.* An integrated map of structural variation in 2,504 human

genomes. *Nature* **526**, 75-81 (2015).

3.   Mahmoud M, Gobet N, Cruz-Dávalos DI, Mounier N, Dessimoz C, Sedlazeck FJ.

Structural variant calling: the long and the short of it. *Genome Biol* **20**, 246 (2019).

4.   Ho SS, Urban AE, Mills RE. Structural variation in the sequencing era. *Nat Rev Genet*

21, 171-189 (2020).

5.   Sedlazeck FJ, Lee H, Darby CA, Schatz MC. Piercing the dark matter: bioinformatics

of long-range sequencing and mapping. *Nat Rev Genet* **19**, 329-346 (2018).

6.   Audano PA, *et al.* Characterizing the Major Structural Variant Alleles of the Human

Genome. *Cell* **176**, 663-675.e619 (2019).

7.   Ebert P, *et al.* Haplotype-resolved diverse human genomes and integrated analysis of

structural variation. *Science*, eabf7117 (2021).

8.   Qin P, *et al.* Pan-genome analysis of 33 genetically diverse rice accessions reveals

hidden genomic variations. *Cell* **184**, 3542-3558.e3516 (2021).

9.   Liu Y, *et al.* Pan-Genome of Wild and Cultivated Soybeans. *Cell* **182**, 162-176.e113

(2020).

10.   Alonge M, *et al.* Major Impacts of Widespread Structural Variation on Gene

Expression and Crop Improvement in Tomato. *Cell* **182**, 145-161.e123 (2020).

11.   Beyter D, *et al.* Long-read sequencing of 3,622 Icelanders provides insight into the

role of structural variants in human diseases and other traits. *Nat Genet* **53**, 779-786

(2021).

12. Golicz AA, Bayer PE, Bhalla PL, Batley J, Edwards D. Pangenomics Comes of Age:

From Bacteria to Plant and Animal Applications. *Trends Genet* **36**, 132-145 (2020).

13. Talenti A, *et al.* A cattle graph genome incorporating global breed diversity. *bioRxiv*,

2021.2006.2023.449389 (2021).

14. Miga KH, Wang T. The need for a human pangenome reference sequence. *Annual*

Review of Genomics and Human Genetics **22**, 81-102 (2021).

15. Crysnanto D, Leonard AS, Fang Z-H, Pausch H. Novel functional sequences

uncovered through a bovine multiassembly graph. *Proc Natl Acad Sci* **118**,

e2101056118 (2021).

16. Leonard AS, *et al.* Bovine pangenome reveals trait-associated structural variation

from diverse assembly inputs. *bioRxiv*, 2021.2011.2002.466900 (2021).

17. De Coster W, Van Broeckhoven C. Newest methods for detecting structural variations.

Trends Biotechnol **37**, 973-982 (2019).

18. Cheng H, Concepcion GT, Feng X, Zhang H, Li H. Haplotype-resolved de novo

assembly using phased assembly graphs with hifiasm. *Nat Methods* **18**, 170-175

(2021).

19. Nurk S, *et al.* HiCanu: accurate assembly of segmental duplications, satellites, and

allelic variants from high-fidelity long reads. *Genome Res* **30**, 1291-1305 (2020).

20. Norris BJ, Whan VA. A gene duplication affecting expression of the ovine ASIP gene

is responsible for white and black sheep. *Genome Res* **18**, 1282-1293 (2008).

21.   Demars J, *et al.* Genome-Wide Identification of the Mutation Underlying Fleece

Variation and Discriminating Ancestral Hairy Species from Modern Woolly Sheep. *Mol*

Biol Evol **34**, 1722-1729 (2017).

22.   Wiedemar N, Drögemüller C. A 1.8-kb insertion in the 3′-UTR of RXFP2 is associated

with polledness in sheep. *Anim Genet* **46**, 457-461 (2015).

23.   Li X, *et al.* Whole-genome resequencing of wild and domestic sheep identifies genes

associated with morphological and agronomic traits. *Nat Commun* **11**, 2815 (2020).

24.   Yang J, *et al.* Whole-genome sequencing of native sheep provides insights into rapid

adaptations to extreme environments. *Mol Biol Evol* **33**, 2576-2592 (2016).

25.   Lv F-H, *et al.* Whole-genome resequencing of worldwide wild and domestic sheep

elucidates genetic diversity, introgression and agronomically important loci. *Mol Biol*

Evol, (2021).

26.   Li H, Feng X, Chu C. The design and construction of reference pangenome graphs

with minigraph. *Genome Biol* **21**, 265 (2020).

27.   Crysnanto D, Pausch H. Bovine breed-specific augmented reference graphs facilitate

accurate sequence read mapping and unbiased variant discovery. *Genome Biol* **21**,

184 (2020).

28.   Durbin RM, *et al.* A map of human genome variation from population-scale

sequencing. *Nature* **467**, 1061-1073 (2010).

29.   Quan C, *et al.* Characterization of structural variation in Tibetans reveals new

evidence of high-altitude adaptation and introgression. *Genome Biol* **22**, 159 (2021).

30.   Gonzalez MV, *et al.* A Divergent Artiodactyl MYADM-like Repeat Is Associated with

Erythrocyte Traits and Weight of Lamb Weaned in Domestic Sheep. *PLoS One* **8**,

e74700 (2013).

31.   French AT, *et al.* The expression of intelectin in sheep goblet cells and upregulation

by interleukin-4. *Vet Immunol Immunopathol* **120**, 41-46 (2007).

32.   Chen S, *et al.* Paragraph: a graph-based structural variant genotyper for short-read

sequence data. *Genome Biol* **20**, 291 (2019).

33.   Yan SM, *et al.* Local adaptation and archaic introgression shape global diversity at

human structural variant loci. *eLife* **10**, e67615 (2021).

34.   Wang F, *et al.* Allele-specific expression and splicing provides insight into the

phenotypic differences between thin- and fat-tailed sheep breeds. *Journal of Genetics*

and Genomics, (2021).

35.   Kalkan Z, Durasi İM, Sezerman U, Atasever-Arslan B. Potential of GRID2 receptor

gene for preventing TNF-induced neurodegeneration in autism. *Neurosci Lett* **620**,

62-69 (2016).

36.   Cheng JY, Stern AJ, Racimo F, Nielsen R. Detecting Selection in Multiple

Populations by Modeling Ancestral Admixture Components. *Mol Biol Evol*, (2021).

37.   Zhu X-J, *et al.* BMP-FGF signaling axis mediates Wnt-induced epidermal stratification

in developing mammalian skin. *PLoS Genet* **10**, e1004687 (2014).

38.   Kardos M, *et al.* Whole-genome resequencing uncovers molecular signatures of

natural and sexual selection in wild bighorn sheep. *Mol Ecol* **24**, 5616-5632 (2015).

39.   Pan Z, *et al.* Rapid evolution of a retro-transposable hotspot of ovine genome

underlies the alteration of BMP2 expression and development of fat tails. *BMC*

Genomics **20**, 1-15 (2019).

40.   Nishikawa S, *et al*. In utero manipulation of coat color formation by a monoclonal anti-

c‑kit antibody: two distinct waves of c‑kit‑dependency during melanocyte

development. *The EMBO journal* **10**, 2111-2118 (1991).

41.   Yang N, *et al*. Genome assembly of a tropical maize inbred line provides insights into

structural variation and crop improvement. *Nat Genet* **51**, 1052-1059 (2019).

42.   Võsa U, *et al*. Large-scale cis- and trans-eQTL analyses identify thousands of genetic

loci and polygenic scores that regulate blood gene expression. *Nat Genet* **53**, 1300-

1310 (2021).

43.   Pan X, *et al*. Modes of genetic adaptations underlying functional innovations in the

rumen. *Science China Life Sciences* **64**, 1-21 (2021).

44.   Kalds P, Luo Q, Sun K, Zhou S, Chen Y, Wang X. Trends towards revealing the

genetic architecture of sheep tail patterning: Promising genes and investigatory

pathways. *Anim Genet* **52**, 799-812 (2021).

45.   Moioli B, Pilla F, Ciani E. Signatures of selection identify loci associated with fat tail in

sheep1. *J Anim Sci* **93**, 4660-4669 (2015).

46.   Xu S-S, *et al*. Genome-wide association analysis identifies the genetic basis of fat

deposition in the tails of sheep (Ovis aries). *Anim Genet* **48**, 560-569 (2017).

47.   Fu W, Wang R, Nanaei HA, Wang J, Hu D, Jiang Y. RGD v2.0: a major update of the

ruminant functional and evolutionary genomics database. *Nucleic Acids Res*,

gkab887 (2021).

48.   Clark EL, *et al*. A high resolution atlas of gene expression in the domestic sheep

734   (Ovis aries). *PLoS Genet* **13**, e1006997 (2017).

735   49.   Chao T, *et al.* Transcriptome Analysis of Three Sheep Intestinal Regions reveals Key

736   Pathways and Hub Regulatory Genes of Large Intestinal Lipid Metabolism. *Scientific*

737   *Reports* **7**, 5345 (2017).

738   50.   Jiang Y, *et al.* The sheep genome illuminates biology of the rumen and lipid

739   metabolism. *Science* **344**, 1168-1173 (2014).

740   51.   Economides KD, Zeltser L, Capecchi MR. HOXB13 mutations cause overgrowth of

741   caudal spinal cordand tail vertebrae. *Dev Biol* **256**, 317-330 (2003).

742   52.   Aires R, *et al.* Tail Bud Progenitor Activity Relies on a Network Comprising Gdf11,

743   Lin28, and Hox13 Genes. *Dev Cell* **48**, 383-395.e388 (2019).

744   53.   Pan Z, *et al.* Rapid evolution of a retro-transposable hotspot of ovine genome

745   underlies the alteration of BMP2 expression and development of fat tails. *BMC*

746   *Genomics* **20**, 261 (2019).

747   54.   Baazaoui I, Bedhiaf-Romdhani S, Mastrangelo S, Ciani E. Genome-wide analyses

748   reveal population structure and identify candidate genes associated with tail fatness

749   in local sheep from a semi-arid area. *Animal* **15**, 100193 (2021).

750   55.   Dong K, *et al.* Genomic analysis of worldwide sheep breeds reveals PDGFD as a

751   major target of fat-tail selection in sheep. *BMC Genomics* **21**, 1-12 (2020).

752   56.   Hickey G, *et al.* Genotyping structural variants in pangenome graphs using the vg

753   toolkit. *Genome Biol* **21**, 35 (2020).

754   57.   Sirén J, *et al.* Pangenomics enables genotyping of known structural variants in 5202

755   diverse genomes. *Science* **374**, abg8871 (2021).

58.   Li R, *et al.* A near complete genome for goat genetic and genomic research. *Genet*

Sel Evol **53**, 74 (2021).

59.   Alonge M, *et al.* RaGOO: fast and accurate reference-guided scaffolding of draft

genomes. *Genome Biol* **20**, 224 (2019).

60.   Manni M, Berkeley MR, Seppey M, Simão FA, Zdobnov EM. BUSCO Update: Novel

and Streamlined Workflows along with Broader and Deeper Phylogenetic Coverage

for Scoring of Eukaryotic, Prokaryotic, and Viral Genomes. *Mol Biol Evol* **38**, 4647-

4654 (2021).

61.   Stanke M, Keller O, Gunduz I, Hayes A, Waack S, Morgenstern B. AUGUSTUS: ab

initio prediction of alternative transcripts. *Nucleic Acids Res* **34**, W435-W439 (2006).

62.   Buchfink B, Reuter K, Drost H-G. Sensitive protein alignments at tree-of-life scale

using DIAMOND. *Nat Methods* **18**, 366-368 (2021).

63.   Benson G. Tandem repeats finder: a program to analyze DNA sequences. *Nucleic*

Acids Res **27**, 573-580 (1999).

64.   Bakker B, *et al.* Single-cell sequencing reveals karyotype heterogeneity in murine and

human malignancies. *Genome Biol* **17**, 115 (2016).

65.   Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic

features. *Bioinformatics* **26**, 841-842 (2010).

66.   Narasimhan V, Danecek P, Scally A, Xue Y, Tyler-Smith C, Durbin R. BCFtools/RoH:

a hidden Markov model approach for detecting autozygosity from next-generation

sequencing data. *Bioinformatics* **32**, 1749-1751 (2016).

67.   Danecek P, *et al.* The variant call format and VCFtools. *Bioinformatics (Oxford,*

778   *England)* **27**, 2156-2158 (2011).

779   68.   Kijas JW, *et al.* Genome-wide analysis of the world's sheep breeds reveals high

780    levels of historic mixture and strong recent selection. *PLoS Biol* **10**, e1001258 (2012).

781   69.   Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina

782    sequence data. *Bioinformatics* **30**, 2114-2120 (2014).

783   70.   Kim D, Langmead B, Salzberg SL. HISAT: a fast spliced aligner with low memory

784    requirements. *Nat Methods* **12**, 357-360 (2015).

785   71.   Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for

786    assigning sequence reads to genomic features. *Bioinformatics* **30**, 923-930 (2013).

787   72.   Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for

788    differential expression analysis of digital gene expression data. *Bioinformatics* **26**,

789    139-140 (2010).

790   73.   Shabalin AA. Matrix eQTL: ultra fast eQTL analysis via large matrix operations.

791    *Bioinformatics* **28**, 1353-1358 (2012).

Tables

 Table 1 Quality assessments of the primary and partially phased assemblies for the 13

 individuals. For each sheep, we show the values for the primary assembly and for

 both partially phased haploid assemblies.

Breeds	Abbr.	Assembly	Assembly length (Gb)	Contig N50 (Mb)	Contig number	Depth	BUSCO (%)	Base accuracy
Kazak	KAZ1	Primary	2.87	73.4	1849	22.4	93.4	46.3
		hap1/hap2	2.85/2.84	10.7/13.9		-	92.3/91.9	46.6/46.4
Kermani	KER1	Primary	2.82	80.3	1675	21.7	93.8	45.2
		hap1/hap2	2.77/2.75	15.4/15.2		-	90.7/92.2	45.4/45.6
Dorset	DOR1	Primary	2.91	92.4	1293	25.0	94.3	45.5
		hap1/hap2	2.86/2.83	32.8/39.5		-	93.4/92.7	45.6/45.6
East Friesian	FRI1	Primary	2.90	85.1	982	28.0	94.4	48.2
		hap1/hap2	2.77/2.75	62.9/53.2		-	91.3/92.8	48.4/48.5
Chiniese Merino	MER1	Primary	2.84	60.0	1773	18.7	94.2	43.3
		hap1/hap2	2.82/2.83	3.7/4.8		-	92.1/90.0	42.4/43.3
Qiaoke	QK1	Primary	2.87	75.0	1654	21.5	94.0	47.1
		hap1/hap2	2.83/2.84	13.6/9.9		-	92.6/93.4	47.6/47.2
Romney	ROM1	Primary	2.86	68.3	1552	19.3	94.4	43.2
		hap1/hap2	2.81/2.82	6.3/5.7		-	92.1/92.4	43.3/43.1
Suffolk	SFK1	Primary	2.88	65.3	1513	18.8	94.3	43.4
		hap1/hap2	2.86/2.80	5.7/4.9		-	92.4/92.5	43.2/43.3
Texel	TEX1	Primary	2.82	47.6	1835	17.9	94.4	44.6
		hap1/hap2	2.76/2.75	4.52/4.49		-	91.4/90.2	44.8/44.8
White Dorper	WDU1	Primary	2.81	17.8	2125	15.4	93.8	41.7
		hap1/hap2	2.82/2.77	1.58/1.26		-	90.2/89.8	41.9/41.8
Ujumqin	UJM1	Primary	2.87	75.6	1534	21.3	94.4	44.4
		hap1/hap2	2.86/2.84	11.5/11.7		-	93.9/93.5	44.5/44.6
Charollais	CHA1	Primary	2.86	71.5	1427	19.2	94.1	44.1
		hap1/hap2	2.82/2.80	6.4/5.1		-	91.7/92.1	44.3/44.3
Yunnan	YUN1	Primary	2.91	73.9	1293	22.0	94.1	46.9
		hap1/hap2	2.89/2.85	18.7/13.1		-	93.6/93.4	46.7/46.6

 Figure legends

Fig. 1 Ovine pan-genome graph construction. (a) PCA plot showing the representation

of the genetic diversity of domestic sheep by the 13 individuals used for HiFi

sequencing and pan-genome construction. (b) The pan-genome size changes with

increase in the number of genome assemblies. (c) Repeat annotation of the structural

variations stratified by SV types. (d) Length distribution of different type of SVs. (e)

Diagram and number of different SV types. The red bold lines indicate the reference

sequence and the blue bold lines represent the non-reference. (f) An example of a

divergent allele of 50,757 bp containing an MYADM-like repeat. (g) An example of a

multiallelic variation containing a large deletion of 105,650 bp which is nested with

another insertion of 5,518 bp. The red lines with arrows show the reference path and

the blue lines with arrows refer to the non-reference path.

Fig. 2 The repertoire of structural variations in sheep genome and frequency spectrum

in domestic and wild sheep. (a) Geographical distribution of the 45 breeds or

populations (see Supplementary Data 7 for the assignment of the breeds or

populations). The Dorper sheep from South Africa and the white Suffolk from

Australia are not shown on the map. (b) Histogram of the number of SV ranks (0-300)

of r^2 values that surpass the median SNP-based r^2 value for common SVs

(MAF>0.05). (c) Derived allele frequencies of the insertions and deletions. (d) The

proportion of standing and *de novo* SVs. (e) High-frequency *de novo* SVs in domestic

sheep. (f) A *de novo* insertion within *GRID2* occurs at high frequencies in all studied

populations. (g) A *de novo* insertion of a retrogene occurs at high frequencies in

European sheep.

Fig. 3 Population-stratified SVs and their effects on gene expression. (a) Genome-

wide distribution of global F_{ST} which is measured by the average value of the top 50%

F_{ST} values for each SV across 33 sheep breeds or populations. The red dots represent

population-stratified SVs that were also found by Ohana approach. (b) Examples of

expression-associated SVs identified by allelic specific expression (ASE) mapping.

Each column represents one ASE SNP in exons and each row represents one RNA-seq

sample. (c) Allele distributions of the three expression-associated SVs in diverse

breeds or populations.

Fig. 4 Selective SVs associated with the long tail trait. (a) The genome-wide Fixation

index (F_{ST}) of long fat-tailed versus short fat-tailed Asian sheep breeds and long thin-

tailed versus short thin-tailed European sheep breeds. (b) The selective haplotype in

long-tailed sheep. The 169 bp-insertion is located upstream of the 5' UTR region of

HOXB13. The haplotypes include the genotypes of the insertion and surrounding

SNPs. (c) The frequency of the insertion in wild sheep and domestic sheep. (d) RNA-

seq data shows that the expression of the insertion. The top panel corresponds to the

expression in the colon of Texel × long-tailed Scottish Blackface sheep

(ERS2107126); the bottom panel corresponds to the expression in colon of one small-

tail sheep (SRR5122597) without the insertion. (e) The electrophoresis diagrams of

PCR products for the three genotypes (homologous insertion: II, heterozygote: ID,

homologous deletion: DD). (f) The carriers and non-carriers of the insertion differ in

tail length. (g) *HOXB13* promoter activity is modulated by the 169 bp insertion.

Luciferase activity in HEK293T cells transfected with recombinant plasmids

containing the insertion or not. Experiments were repeated at least five times. The

data are presented as mean ± standard error of the normalized luciferase activity.

Fig. 5 Selective SVs associated with the fat-tail trait. (a) Population branch statistic

(PBS) values across the whole genome by comparing fat-tailed sheep to thin-tailed

sheep using the mouflon sheep as an outgroup. (b) The two most differentiated

regions between fat-tailed sheep and thin-tailed sheep. The left panel shows the

intergenic region between *BMP2* and *HAO1* while the right panel corresponds to the

region surrounding *PDGFD*. SVs with PBS value >0.8 are highlighted by blue dotted

lines. (c) The haplotypes of mouflons and domestic sheep for the two most selective

regions. Each column represents one SV and each row represent one individual. The

black reverted triangles represent *de novo* SVs.

Supplementary Files

Supplementary Table 1 Accession numbers of the *de novo* assemblies.

Supplementary Data

Supplementary Data 1 Information of divergent variations.

Supplementary Data 2 Information of multiallelic variations.

Supplementary Data 3 List of the biallelic insertions and deletions identified in this

study.

Supplementary Data 4 SV frequency in 45 sheep breeds or populations.

Supplementary Data 5 List of population-stratified SVs.

Supplementary Data 6 List of putatively expression-associated SVs identified by

ASE mapping.

Supplementary data 7 List of putatively expression-associated SVs identified by

eQTL mapping.

Supplementary Data 8 Sample list of the resequencing data.

Supplementary figures

Supplementary Fig. 1 Example of three representative divergent variations.

intersecting with IDE (a), PRP3-like (b) and MAGI2 (c), respectively.

Supplementary Fig. 2 Example of three representative multiallelic variations

affecting ASIP (a), UDP-glucuronosyltransferase 2B18-like (b) and olfactory receptor

(LOC101115700) (c), respectively.

Supplementary Fig. 3 The number of SVs intersecting functional elements compared

to randomly permuting SV locations.

Supplementary Fig. 4 An ideogram showing SV hotspots using all SVs. The total

number of SVs in each detected hotspot is shown by a scale going from blue to red.

Supplementary Fig. 5 Enrichment analysis of SVs with respect to the last 5 Mb of

each chromosome end.

Supplementary Fig. 6 PCA plot showing the clusters of domestic sheep by insertions

(a) and deletions (b).

Supplementary Fig. 7 Enriched KEGG pathways for the population stratified SVs.

Supplementary Fig. 8 RNA-seq mapping for the 169 bp insertion shows that the

insertion belongs to 5' UTR region.

a

b

KHK pbsv.INS.30596 (119 bp)

APOL3-Like pbsv.INS.38285 (51 bp)

pbsv.INS.49258 (78 bp) **DOCK6**

Allelic imbalance
Ref allele ratio
Alt allele ratio

c

a

b

chr11:37,502,704 - 37,547,639 bp

169 bp Insertion

REF HET ALT

c

Fat tail

Thin tail

Long tail

LTH   TAN   SSS   EFR   SOG   DEM   SFK

Short tail

HUS   STH   UJM   SEL   FIN   OUE   GTL

Wild

Mouflon

REF
ALT

d

chr11:37,525,005   *HOXB13*

169 bp

Long tailed

Short tailed

Depth

e

Marker   II   ID   DD

bp

bp

f

$P<0.0001$

II (n=57)   ID (n=144)

Genotype

g

$P<0.0001$

HOXB13-WT   HOXB13-MUT