The first sheep graph pan-genome reveals the spectrum of structural variations and their effects on different tail phenotypes

Ran Li[1#], Mian Gong[1#], Xinmiao Zhang[1#], Fei Wang[1#], Zhenyu Liu[1#], Lei Zhang[1#],

Mengsi Xu[2], Yunfeng Zhang[2], Xuelei Dai[1], Zhuangbiao Zhang[1], Wenwen Fang[1], Yuta

Yang[1], Huanhuan Zhang[1], Weiwei Fu[1], Chunna Cao[1], Peng Yang[2], Zeinab Amiri

Ghanatsaman[3], Niloufar Jafarpour Negari[4], Hojjat Asadollahpour Nanaei[1,4],

Xiangpeng Yue[5], Yuxuan Song[1], Xianyong Lan[1], Weidong Deng[6], Xihong Wang[1],

Ruidong Xiang[7], Eveline M. Ibeagha-Awemu[8], Pat (J.S.) Heslop-Harrison[9], Johannes

A. Lenstra[10], Shangquan Gan[2]*, Yu Jiang[1]*

[1]Key Laboratory of Animal Genetics, Breeding and Reproduction of Shaanxi

Province, College of Animal Science and Technology, Northwest A&F University,

Yangling, Shaanxi, 712100, China

[2]State Key Laboratory of Sheep Genetic Improvement and Healthy Production,

Xinjiang Academy of Agricultural and Reclamation Sciences, Shihezi, Xinjiang

832000, China

[3]Department of Animal Science, Fars Agricultural and Natural Resources Research

and Education Center, Agricultural Research, Education & Extension Organization

(AREEO), Shiraz, Iran

[4]Department of Animal Science, Shahid Bahonar University of Kerman, Kerman, Iran

[5]State Key Laboratory of Grassland Agro-ecosystems, Key Laboratory of Grassland

Livestock Industry Innovation, Ministry of Agriculture and Rural Affairs, Engineering

Research Center of Grassland Industry, Ministry of Education, College of Pastoral

Agriculture Science and Technology, Lanzhou University, Lanzhou, 730020, China

[6] Faculty of Animal Science and Technology, Yunnan Agricultural University,

Kunming, 650201, China

[7] Faculty of Veterinary & Agricultural Science, The University of Melbourne,

Parkville, 3052, VIC, Australia

[8] Agriculture and Agri-Food Canada, Sherbrooke Research and Development Centre,

Sherbrooke, QC J1M 0C8, Canada

[9] Department of Genetics and Genome Biology, University of Leicester, Leicester, UK

[10] Faculty of Veterinary Medicine, Utrecht University, Utrecht, The Netherlands

[#] These authors contributed equally.

*To whom correspondence should be addressed.

Email: yu.jiang@nwafu.edu.cn; shangquangan@163.com

Abstract

Structural variations (SVs) are a major contributor of genetic diversity and phenotypic

variations, however their prevalence and functions in domestic animals are largely

unexplored. Here, we assembled 26 haplotype-resolved genome assemblies from 13

genetically diverse sheep breeds using PacBio HiFi sequencing. We then constructed

an ovine graph pan-genome and demonstrated its advantage in discovering 142,593

biallelic SVs (Insertions and deletions), 7,028 divergent alleles and 13,419

multiallelic variations with high accuracy and sensitivity. To link the SVs to

genotypes, we genotyped the SVs in 687 resequenced individuals of domestic and

wild sheep using a graph-based approach and identified numerous population-

stratified variants, of which expression-associated SVs were detected by integrating

RNA-seq data. Taking the varying sheep tail morphology as example, we located a

putative causative insertion in *HOXB13* gene responsible for the long tail and reported

multiple large SVs associated with the fat tail. Beyond generating a benchmark

resource for ovine structural variants, our study also highlighted that the population

genetics analysis based on graph pan-genome rather than reference genome will

greatly benefit the animal genetic research.

Keywords: sheep pan-genome, structural variations, *de novo* assembly, tail
morphology

Introduction

Structural variations (SVs) are a major source of genetic variation that range from 50

base pairs (bp) to over megabases (Mb) in size. Compared to single-nucleotide

polymorphisms (SNPs), SVs could cause larger scale genomic perturbations in genes

and regulatory regions and are therefore more likely to affect gene expression and

phenotypes [1, 2].

However, SV detection has been challenging primarily due to technical

limitations [3]. The approach based on short reads (100-150 bp) causes poor

sensitivity and high false discovery rate [4]. To overcome the limitation of short reads,

the long-read sequencing such as PacBio SMRT sequencing and Oxford Nanopore

sequencing produce long reads (>10 kb) and greatly facilitated the efficacy of SV

discovery by directly spanning the breakpoints [5-9]. Despite that, the conventional

long reads suffer from high sequencing error rate (5-15%), which not only implicated

downstream analysis but also hampered discernment in highly repetitive regions. In

addition, the linear reference assembly has been shown to be not fully compatible

with the methodology of SV detection [10, 11].

The emergence of PacBio high fidelity (HiFi) reads is revolutionizing our

methodologies in *de novo* assembly and SV detection [12]. As a major leap forward in

long-read sequencing, the HiFi reads are both long（15 kb – 25 kb）and highly

accurate（>99.9% base accuracy）[13], offering base-level resolution at SV

breakpoints. Furthermore, the HiFi sequencing could deliver haplotype-resolved

assemblies to depict the sequence diversity of the diploid genome [14], which are

essential for pan-genomic studies. Compared with a linear reference genome, the pan-

genomic reference models can now maximumly utilize the information from

haplotype-resolved genome assemblies to faithfully capture complex regions and SVs

in the sequenced individual [11, 15]. Yet the superiority of HiFi sequencing in pan-

genome construction and SV detection has just revealed [6, 16, 17]. The analysis of *de*

novo assemblies at population-scale using HiFi sequencing is going to transform our

understanding of SV complexity in each species.

Sheep are among the first domesticated animals with economic and cultural

importance. Previous studies have shown that SVs determine the white coat color [18],

fleece variations [19] and horn phenotypes (polledness) [20] – characters that have

been under selection for millennia. Nevertheless, a comprehensive characterization of

genetic variation characterized by SVs remains to be explored in sheep as in most

animal species. In this study, we generated reference-grade *de novo* assemblies for 13

representative Eurasian sheep breeds, together with two sets of partially phased

assemblies for each individual taking advantage of Pacbio HiFi sequencing. These

datasets allow us to construct a graph-based ovine pangenome and explore the SV

landscape with unprecedent accuracy and sensitivity. We further analyzed the SV

spectrum in 686 domestic and wild sheep from short-reads sequencing and

highlighted SVs as largely unexplored contributors to variations in gene expression

and phenotypes.

Results

High-quality *de novo* assemblies of 13 representative sheep breeds

Domestic sheep could be generally divided into five genetic groups including

European, Asian, Middle Eastern and two African lineages (one of the Dorper sheep

and the other of the African landraces) [21]. We selected 13 genetically diverse

accessions to cover the major genetic groups, including seven European, four East

Asian, one Middle Eastern (Kermani sheep from Iran) and one African breeds (White

dorper). The seven distinct European breeds covered those for meat (Suffolk, Dorset,

Texel, Charollais), wool (Merino, Romney) and dairy (East Friesian) purposes and the

East Asian accessions covered all the three lineages of Northern China (Ujumqin,

Kazak), Qinghai-Tibetan Plateau (Tibetan) and Yunnan-Kweichow Plateau (Yunnan).

The 13 individuals displayed distinct phenotypes and a good representation especially

for the Eurasian breeds. A SNP analysis showed that they encompass 95.5% of the

SNPs with minor allele frequency (MAF) >0.05 among the studied 645 domestic

sheep worldwide.

To obtain high-quality *de novo* assemblies, the 13 accessions were sequenced

using PacBio HiFi sequencing at an average coverage of 20.9×, yielding highly

accurate long reads with per-base accuracy >99.9%. We used the Hifiasm assembler

[14] for *de novo* assembly, which could generate the primary contigs as well as two

sets of partially phased contig assemblies from HiFi reads. We ultimately assembled

13 primary collapsed assemblies with average length of 2.86 Gb (2.81 Gb- 2.92 Mb)

(Table 1). The average contig number were 1582 with contig N50 length of 68.2 Mb.

The most complete assembly (FRI1) achieved a final contig N50 of 85.1 Mb which is

much higher than the sheep reference genome (ARS-UI_Ramb_v2.0). We found that

the HiFi sequencing depth is not correlated with assembly length (r^2=-0.04, Spearman

rank correlation), but positively correlated with contig N50 length (r^2=0.93, Spearman

rank correlation) and negatively correlated with contig N50 number (r^2=-0.67,

Spearman rank correlation). We further ordered and oriented contigs of the 13

collapsed assemblies into chromosomes based on their synteny to the reference

genome. As a result, on average 93.58% of the contigs per assembly were placed into

chromosomes. The resulting chromosome-level collapsed assemblies for the 13

representative breeds will serve as a reference panel for future research in sheep

genetics and genomics. Meanwhile, the partially phased assemblies displayed an

average contig N50 length of 14.5 Mb, which could capture the variations of the

diploid and will be used for the subsequent construction of the multiassembly graph.

We assessed the completeness using Benchmarking Universal Single-Copy

Orthologs (BUSCO) analysis and the base accuracy using quality value (QV) for the

collapsed and partially phased assemblies (Table 1). The collapsed assemblies

displayed an average completeness of 94.1% (93.8%-94.4%) whereas the BUSCO

scores for the partially phased assemblies were lower but still consistently >90%

complete. To measure the QV of the assemblies, the 13 accessions were also

sequenced to an average of 30× depth Illumina short-reads. All the collapsed and

partially phased assemblies displayed higher quality value than that of the sheep

reference genome assembly (QV=42.5) except for WDU1.

Construction of sheep graph pan-genome

To build a comprehensive sheep graph pan-genome, we used the reference genome

(ARS-UI_Ramb_v2.0) as the backbone and the 26 haplotype-resolved assemblies

from the 13 individuals were augmented into the graph individually. The partially

phased assemblies, instead of the primary contigs, were used as they would

encompass more comprehensive sequence variations of a haploid genome. A

multigraph genome was generated using minigraph [22] which is one of the state-of-

art graph constructors especially for discovering complex variations. Only autosomal

sequences were considered in the multigraph genome, as the X and Y chromosomes

of males were underrepresented in the sequencing.

The resulting multiassembly graph spanned 2.75 Gb including 167,565

nonreference nodes of 137.7 Mb. Based on the presence and absence of each node

among reference genome and the 26 assemblies, we classified the graph pangenome

into three categories: core (nodes present in all assemblies), flexible (nodes present in

at least two assemblies) and private (nodes present in one assembly). The core and the

flexible genome accounted for 69.6% and 29.0% of the total size, respectively,

whereas the private part was minimal (1.4%) (Fig. 1B). The variations were most

associated with LINEs following by those with mixed interspersed repeats (Fig. 1C).

The graph genome encompassed 91,848 structural variations with length ≥100 bp,

representing bubbles that are located between a single source and a single sink node

coming from the reference assembly. The paths in a bubble below 100 bp were not

considered, since we found that the small variants contain abundant false positives by

manual inspection. On the other hand, the extended graph coordinate system proposed

by minigraph enabled us to annotate complex variations including those in highly

diverse regions. We classified the structural variations into three categories: (1)

biallelic variations (Insertions and deletions); (2) divergent alleles where a non-

reference node of ≥100 bp from a single path were found; (3) multiallelic variations

where more than one non-reference paths were present (Fig. 1D and Supplementary

data 1 and 2). Notably, from the multiassembly graph, we identified 7028 divergent

loci adding up to 11.7 Mb and another 13，419 multiallelic loci of 33.3 Mb. The

median length of divergent alleles (726 bp) and multiallelic loci (575 bp) were longer

than that of biallelic variations (452 bp).

The major advantage of the multiassembly graph does not lie in the efficacy in

discovering regular biallelic variations (insertions/deletions); instead, it is most

efficient in recovering complex SVs, i.e., the divergent and multiallelic variations.

These complex SV could be an import type of structural variations that can be trait-

associated but are generally neglected by most SV callers. The divergent variations

intersected with 1797 genes with the non-reference alleles predicted to harbor 27

genes. A divergent locus containing MYADM-like repeat was previously reported to

be associated with mean corpuscular hemoglobin concentration traits and weight of

lamb weaned [23], but the full sequence of the divergent allele was not resolved. We

found this divergent allele to be 50.7 kb long and was presented in 14/26 of the

partially phased haplotypes (Fig. 1E). We also noticed several large divergent alleles

that affected genic region of insulin degrading enzyme (IDE), Membrane Associated

Guanylate Kinase, WW And PDZ Domain Containing 2 (MAGI2), placental

prolactin-related protein 3-like (Supplementary fig. 1). The multiallelic loci

intersected 5092 genes and the non-reference alleles themselves were predicted to

harbor 161 genes. One of the largest multiallelic region represented a deletion of

105.6 kb which is nested with an insertion of 5.5 kb (Fig. 1F). The deletion likely

result in the loss of intelectin gene, that displays specific expression in lung of sheep

and potentially plays a role in mucosal response in allergy [24]. Another loci (45.6 kb)

harboring beta hemoglobin genes was previously reported as divergent allele and

associated with tolerance of anemia [25], but it actually also belonged to multiallelic

variation (Supplementary fig. 2). We provided additional examples of multiallelic loci

that would result in gene gain/loss, such as ASIP, UDP-glucuronosyltransferase 2B18-

like, olfactory receptor (LOC101115700), etc. (Supplementary fig. 3).

A comprehensive catalog and frequency spectrum of SVs in sheep population

Despite advantages in discovering divergent alleles and multiallelic variations, the

graph genome generated by minigraph does not involve base alignment, rendering

accurate breakpoint assignments for many structural variations. In addition, minigraph

is not well adapted for discovering small SVs, i.e. SVs below 100 bp [14]. We

therefore used pbsv (https://github.com/PacificBiosciences/pbsv), a read based SV

caller, to jointly call SVs among the 13 individuals. As a complementing strategy to

the pan-genome, the reads-based approach could determine the precise breakpoints

and included SVs below 100 bp. Aligning PacBio HiFi reads to the reference genome

generated 147,832 SVs including 142,450 insertions/deletions, 516 inversions and

4,866 duplications (Supplementary data 3). We observed much more insertions than

deletions, which contrasts with results based on short read sequencing where deletions

are typically more frequent and easier to identify [26]. The pbsv call set covers 94.8%

of the biallelic variations from minigraph; we therefore merged the pbsv call set and

the large variants (>5 kb) with from minigraph that did not overlap with pbsv call set,

resulting a final comprehensive catalogue of 147,975 SVs (insertions and deletions) in

sheep genome. It should be noted that the SV catalogue include the 516 inversions,

4,866 duplications 7,028 divergent alleles and 13,419 multiallelic variations, which

however were not subjected to the subsequent population-level genotyping using

short-reads data.

The number of variants rapidly decreases with length, consistent with the reports

in humans using long-read sequencing [5, 6, 27]. We observed noticeable peaks at 150

bp corresponding to SINES, 1,350 bp and 1,600 bp corresponding to LTR, as well as

7,750 bp corresponding to LINES (Fig 2A). The SVs were most underrepresented in

untranslated region (UTR), coding sequences (CDS), and regulatory elements

(H3K4me3, H3K27ac), suggesting strong negative selection that acted against them

(Supplementary fig. 4). We also reported 201 SV hotspots sites covering 18.2 Mb of

the autosomes (Supplementary fig. 5). The SVs were significantly clustered at the

terminal 5 Mb region of each chromosome with 4.1-fold enrichment (p value<0.01,

permutation test) (Supplementary fig. 6).

In order to better understand the frequency spectrum of the SVs and identify

those under selection, we used the graph genotyping software Paragraph [28] to

genotype the variants in a panel of 687 resequenced samples. Only the insertions and

deletions from autosomes were genotyped. Paragraph represent SVs as diverging

paths in a graph and could achieve high rate of genotyping accuracy [29]. The

samples included 644 domestic sheep from Europe, East Asia Middle East and Africa ,

33 Asiatic mouflons and 8 argali with an average coverage of 18.1$\times$ (Fig. 2B).

To ensure a high-quality set of genotyped SVs for population genetics analysis,

we filtered genotyping results based on genotyping rates and adherence to Hardy

Weinberg equilibrium [29]. We excluded 12,705 SVs that failed to be genotyped

in >90% of the samples and another 2011 displaying deviation from Hardy Weinberg

equilibrium by displaying excess of heterozygotes, leaving 121,258 (89.2%) reliably

genotyped SVs for downstream analysis (Supplementary data 4). We excluded the

SVs not because that they are unreliable but because they cannot be correctly

genotyped probably due to the limitation of short reads. We also estimated the

genotyping accuracy of the remaining SVs by measuring the Mendelian errors, which

represent genotypes that are found in the offspring but could not be inherited form

either parent. The Mendelian error rate was 3.8% on average in the 11 trios of 29

individuals (each ~30x), suggesting high accuracy of the genotyping results.

PCA analysis using either insertions or deletions displays similar population

structure and shows clear separation of the four groups, and in many cases, even

individual breeds could be distinguished (Supplementary fig. 7), agreeing with the

clusters defined by SNPs. Therefore, the SV catalogue represented a broad diversity

of worldwide sheep population.

The level of linkage disequilibrium (LD) between SVs and SNPs could reflect the

extent of SVs representing independent and undiscovered genetic variations relative

to previous evolutionary studies. We estimated how frequently the SVs were linked to

the flanking SNPs. About 41.8% SVs link strongly with surrounding SNPs

(rank>200), the characteristic of which could be detected by the surrounding linked

SNPs. It is worth noting that there are still 37.8% SVs display low levels of LD with

nearby SNPs (rank < 100), implying variants that are poorly tagged by easily

genotyped SNP markers (Supplementary fig. 8).

In order to infer the allele state (ancestral or derived) for each SV, we additionally

genotyped one snow sheep and one bighorn sheep and, together with one argali as

outgroup, and could identify the state for 115,089 SVs. Based on the allele frequency

of derived alleles (DAF) in domestic sheep, the variants were classified into four

categories: shared (DAF $\geq$ 0.95), major (0.95 > DAF $\geq$ 0.5), polymorphic (0.5 > DAF

$\geq$ 0.05) and rare (DAF < 0.05) SVs (Fig. 2C).

We next dissected *de novo* SVs in particular those arising to high frequency in the

domestication and improvement process. The *de novo* SVs accounted for 28% of all

the SVs (Fig. 2D). It should be noted that the proportion of *de novo* SVs would be

overestimated due to the relative limited number of wild sheep. Among them, we

found 592 high-frequency *de novo* SVs (DAF $\geq$ 0.5 in at least one geographic group)

(Fig. 2E). The *de novo* SVs reaching high frequency in all domestic sheep might be

associated with domestication. Those SVs were found to be enriched in KEGG

pathways of Ras signaling and long-term depression (adjusted p value < 0.05). For

example, a *de novo* mutation of 3.4 kb insertion dominating in domestic sheep were

identified in the first intron of *GRID2* (Fig. 2F). The gene plays a key role in

synaptogenesis and neurodevelopment [30]. Those that rapidly arose in certain

geographic regions might be involved in the emergence of some domesticated traits. A

1.4 kb *de novo* insertion prevalent in European sheep was found at the 3' UTR of

IRF2BP2 gene (Fig. 2G), which was reported as the causative mutation for the woolly

hair [19].

Population-stratified SVs and their potential effect on gene expression

After initial domestication at the Fertile Crescent, domestic sheep dispersed

worldwide and adapted to a wide range of natural and artificial selection. In this

context, we detected putative selection signals by dissecting population-stratified SVs

among the 33 breeds/populations. The population differentiation was measured

among each breed versus all others breeds using a modified calculation of global Fst

(see methods). It should be noted that this strategy tends to discover common

selection signals present in multiple breeds. Therefore, the population-stratified SVs

between Europe and East Asian breeds are preferentially detected since they are the

predominant breeds in the comparison.

We identified 865 population-stratified SVs with the top 1% global Fst values

(Supplementary data 4), of which 351 intersected with genic regions and 324 were

within 100 kb of gene flanking region (Fig. 3A). The majority of them (698, 80.7%)

were confirmed by the Ohana approach with which we detected selection in multiple

populations by varying the number of ancestral admixture components. The related

genes were associated with diverse phenotypic and production traits, including wool

type (IRF2BP2, FGF7) [19, 31], tail morphology (HOXB13) [32], horn type (RXFP2)

[33], ear type (MSRB3) [34], fat deposition (BMP2, PDGFD, PDGFA) [35], coat

color (KIT) [36], etc. Notably, the highest stratified SV (Fst=0.78) was the insertion

as shown in Fig. 2G, representing the most differentiated SV among European and

Asian breeds. The next two highest signals affecting HOXB13 and BMP2 were both

associated with tail morphology which will be described in following sections.

Functional annotation of the putatively affected genes revealed enrichment in KEGG

pathways involved in important signaling pathways (Rap1 signaling pathway, MAPK

signaling pathway), parasitic disease (African trypanosomiasis, amoebiasis), etc.

(Supplementary fig 9).

SVs could impact the gene expression of neighboring genes by altering the

composition or location of cis-regulatory elements [9, 37]. To explore SV-associated

expression changes, we leveraged transcriptomic data from 13 hybrids of Texel×

Kazak sheep [38, 39], for each of which RNA-seq data for 8 tissues were available.

Additionally, we obtained the WGS data for each of the individual as well as their

parents. With these datasets, we were able to identify the candidate expression-

associated SVs using two complementary strategies by either SV-eQTL mapping or

allelic-specific expression mapping (see methods). For SV-eQTL mapping, we

defined a list of SV-gene pairs based on their location and correlate the genotype of

each SV with different expression levels of associated gene. 705 SVs were found to

be associated with gene expression change in at least two tissues (adjusted p

value<0.05, two-sided Mann-Whitney U test) (Fig. 3B). For allelic-specific

expression mapping, we expected that heterozygous SVs could act as cis-regulatory

elements and cause allelic-specific expression in associated genes. In this context,

1108 SVs were found to be associated with allelic-specific expression of nearby genes

(Fig. 3C, 3D). Therefore, based on the two approaches, we identified 1794 SVs which

potentially affect gene expression (Supplementary data 5). Notably, 43 of the

expression-associated SVs were among the population-stratified SVs, influencing

expression of genes involved in immunity (APOL3-like, LOC114118416) [40-42],

lipid and fructose metabolism (SEC31A, KHK) [43, 44], osteoblast differentiation

(BMP2, METTL7A) [45, 46], and development (DOCK6, DIP2B) [47, 48], etc.

SVs as strong candidates affecting sheep tail morphology

The sheep tail is a key genetic trait that is related to domestication, adaptation, and

animal welfare. The wild ancestors (Asiatic mouflon) have a thin and short tail

whereas the domestic sheep developed four tail types including the primitive short

thin-tailed type and the derived short fat-tailed (including fat-rumped), the long fat-

tailed and long thin-tailed types. The underlying causative mutations are still

unknown, but the long tail and the fat tail traits are apparently derived traits that arose

in subsequent improvement process indicating that the causative mutations are likely

to be *de novo* variants in domestic sheep.

Since the long tail is the derived trait, the causative SV is expected to be under

selection in long tailed sheep breeds. To identify SVs associated with tail length, we

performed comparisons within the fat-tailed type (the long fat-tailed breeds versus the

short fat-tailed breeds) from Asian breeds and the thin-tailed type (the European long

thin-tailed breeds versus the North European short thin-tailed breeds), respectively.

Notably the most differentiated SV from the two independent comparisons both point

to an insertion of 169 bp (chr11:37525005) close to the 5' UTR of HOXB13 gene (Fig.

4A). Residing in a specific haplotype exclusively found in long tailed sheep, the

insertion is prevalent in long-tailed breeds whereas rare in short-tailed breeds (Fig. 4B

and 4C). This gene is mainly expressed in the embryo and colon of sheep and cattle

by searching the Ruminant genome database [49]. We luckily found the expression of

HOXB13 in colons of the Texel x Scottish Blackface individuals presumably carrying

at least one copy of the insertion [50], and individuals with short tail presumably

absence of the insertion (Texel sheep and small-tail sheep) [51, 52]. As expected, the

insertion was only expressed in the colons of the Texel x Scottish Blackface with no

expression in Texel sheep and small-tail sheep, indicating that it belongs to the 5'

UTR region of HOXB13 gene (Fig. 4D and Supplementary fig 10).

In order to verify the effect of this insertion on tail length, we evaluated the

association between the insertion and tail length in a population of 201 sheep of the

same age (nine months old). The three genotypes including homologous insertion (II),

heterozygote (ID) and homologous deletion (DD) could be directly distinguished

using agarose gel electrophoresis of PCR products (Fig. 4E). The tail length of the

homologues (II) found to be significantly longer than that of the heterozygotes (ID)

by 2.77 cm on average (Mann-Whiney U test, $p<0.0001$) (Fig. 4F). To validate the

function of the insertion, we performed luciferase reporter assays. Relative to the

wildtype (WT), the mutant with the insertion dramatically decreased luciferase

activity by 10-fold (Fig. 4G), indicating that the insertion could reduce transcription

level of HOXB13. HOXB13 has been previously identified as a repressor of caudal

vertebra development [32]. Therefore, the insertion is likely to increase the tail length

of sheep by depressing the expression of HOXB13.

To identify candidate SVs related with fat tail, we compared the genomes of eight

fat tail breeds (SFK, KAZ, HUS, UJM, LTH, TAN, BSB, SSS) and ten thin tail breeds

(FIN, DEM, EFR, GTL, SEL, SFK, SOG, POD, TIB, YBB). We applied the

population branch statistic (PBS) to identify positive selection in fat tail sheep using

Asiatic mouflon as outgroup. Two most differentiated region were identified by either

SV or SNPs (Fig. 5A, 5B), one located at the intergenic region between BMP2 and

HAO1 gene (hereafter referred to as IBH region) and the other in PDGFD gene.

Notably, the most stratified SVs in these two regions were among the top selective

variants of both SVs and SNPs. The most differentiated SV cluster were located at

IBH region which has also reported as a candidate selective region affecting fat tail

trait [53, 54]. Notably, six highly differentiated SVs including one *de novo* SV were

identified in the selective region (Fig. 5C), of which the largest one is a 7728 bp

insertion. PDGFD is recognized as the most plausible candidate gene for the fat-tail

phenotype in recent studies [21, 55]. We found three highly differentiated SVs

residing in the selective region of PDGFD, of which the largest event is a 867 bp

insertion. The three selective variants were all *de novo* SVs that were not present in

Asiatic mouflon but prevalent in fat-tailed sheep. Therefore, our result suggested that

both of these regions were likely to be involved in the fat tail phenotype and the SVs

that we identified should be prioritized when searching for the causative mutations.

Discussion

The genome-sed analysis of genomic structural variations in diverse breeds is critical

to understanding the full repertoire of genetic diversity and pinpointing their

associations with phenotypic traits. However, previous resequencing studies mainly

rely on mapping short reads to a linear reference genome to detect genetic variations,

precluding the discovery of regions with pronounced structural variations [56]. The

graph-based pan-genome model presents a new approach to assess the whole

landscape of sequence diversity, and is likely to become the new guide for future

genomics research. To build a high-quality ovine pan-genome, we utilized the PacBio

HiFi sequencing to generate a panel of reference-level assemblies and partially phased

haplotypes for 13 representative sheep breeds, providing a solid foundation for in-

depth functional genomic studies. The long and accurate HiFi reads guarantee the

high-standard of our pan-genome through generation of high-quality *de novo*

assemblies, preservation of the two haplotypes and precise assignment of SVs'

breakpoints. The development of this ovine pan-genome will be a remarkable step

toward shifting in animal genomics research from a linear reference genome to a pan-

genomic model comprising of multiple reference-quality representative genomes.

Application of graph-based pangenome has its clear opportunities and obstacles.

The pan-genomic approaches are not so straightforward in practice as the linear

reference genome and are still not fully developed in the methodologies and

implementations [11, 57]. For example, despite becoming one of the most promising

graph constructors, the minigraph that we applied has limitations in performing base

alignment, presenting relative high positives for small structural variants [22].

Nevertheless, a shift to the graph-based pan-genomes model will provide researchers

with enhanced opportunities to systematically investigate previously inaccessible

genomic variations. In our study, we found that the minigraph shows great

performance in discovering divergent alleles and multiallelic variations，which

account for 27.4% of total variations in length but are mostly neglected in previous

studies. We further applied the read-based approach to resolve small SVs. The two

complementary approaches of graph-based pan-genome and read-based calling

together ensures a reference panel of high-quality SVs for downstream functional

genomic studies.

The pangenome representation with an influx of high-quality assemblies will

revolutionize our understanding of SV prevalence and their biological significance,

giving a new dimension to the nature and extent of genomic variation, beyond SNPs,

impacting traits. We demonstrated how this pangenome datasets can be utilized to link

SVs to phenotypes by genotyping and assessing the SV landscape in a large

population from short-read sequencing. We applied Paragraph to genotype SVs,

which is a graph-based genotyper and enables efficient encoding of structural

variation into the pangenome and genotyping of those variants [28]. The population-

scale study allows us to trace their responses to natural and artificial selection. Taking

the tail morphology evolution as an example, we identified the putative causative

mutation in HOXB13 gene that causes long tail, highlighting the keys roles of SVs

underlying phenotypic changes. We further reported segregating SVs in fat tail breeds

and in particular *de novo* SVs that showed higher or comparative selection signals

than surrounding SNPs, which warrant further investigation.

Methods

Sample collection and PacBio sequencing

Samples were selected to maximumly represent world sheep diversities, especially

Eurasian breeds. Fresh blood samples were collected for each individual. High-quality

genomic DNA (gDNA) was extracted as previously described [58] and assessed for

purity and quantity using Nanodrop 1000 (Thermo Fisher) and Qubit (Thermo Fisher)

assays. Library with an average insert size of ~15 kb was generated using the

SMRTbell Express Template Prep Kit 2.0 (PacBio) and fractionated on the SageELF

(Sage Science, Beverly, MA) into narrow library fractions. Libraries were then

sequenced on 2-3 SMRT Cells 8M on a Sequel II instrument using Sequel II

instrument (Pacbio) using 30-hour movie times. Raw data was processed using the

CCS algorithm (version 6.0.0, parameters: --minPasses 3 -- minPredictedAccuracy

0.99 –maxLength 21000) to generate highly accurate HiFi reads.

De novo **assembly using HiFi reads and quality assessment**

Hifiasm v0.15.3-r339 was used to generate the assembly from HiFi CCS reads using

default parameters [14]. Hifiasm yields one primary contig assembly and two sets of

partially phased contig assemblies. The primary contig assembly for each breed was

scaffolded to chromosome level using RagTag v2.0.1 with the reference sheep

genome assembly [59].

The genome completeness was assessed using the BUSCO program (v.3.0.2)

containing the Mammalia odb9 gene set (4104 BUSCO genes) [60]. The single plus

duplicated complete BUSCO gene counts are reported. The base accuracy was

measured by assembly quality value (QV) using yak (https://github.com/lh3/yak),

which compares the 31-kmers found in short-reads and the assembly sequences.

Pan-genome construction

The pan-genome construction was performed using a previously reported workflow

[61] with minor modifications. Minigraph v0.15-r426 with option -xggs was used to

generate the graph pan-genome [22]. The sheep reference genome (ARS-

UI_Ramb_v2.0, GCF_016772045.1) was used as the backbone of the multiassembly

graph and the 26 partially phased contig assemblies were added successively.

Identification of non-reference divergent alleles and multiallelic alleles from the

graph pan-genome

The resulting graph by minigraph is composed of chains of bubbles with the reference

as the backbone. We first used gfatools v0.5-r234 which is a bubble popping

algorithm to extract bubbles from the graph (https://github.com/lh3/gfatools). Each

bubble represents a structural variation, encompassing the start and end nodes of

reference sequences as well as path traversing the start and end nodes.

Depending on the number of paths in a bubble, the structural variations were

classified into three categories: (1) biallelic variations (Insertions/Deletions): either

the reference path is longer (deletions) or non-reference path is longer (insertions).

The node in the shorter path is required to be below 100 bp. (2) Divergent alleles:

biallelic variations where the reference nodes and the non-reference nodes were both

longer than 100 bp. (3) Multiallelic alleles: multiallelic variations where one reference

path and more than one nonreference path were present. Those nonreference paths

below 100 bp were excluded.

Genome annotation of non-reference sequences

The nodes in a bubble that are not derived from the reference path were referred to as

non-reference alleles. The genes within non-reference sequences were annotated using

a homology-based approach. Protein sequences from different species including cattle

(Bos taurus), goat (Capra hircus), sheep (Ovis aries) and pig (Sus scrofa) were

downloaded from NCBI and processed with Exonerate v2.2.0 [62] to find the best hit

per protein sequence. EvidenceModeler v1.1.1 [63] was used to integrate all evidence

into a non-redundant gene annotation.

Annotation of repeats

Interspersed repeats and low complexity DNA sequences were identified using

RepeatMasker v4.0.7 (http://www.repeatmasker.org) with a combined repeat database

including Dfam v.20170127 and RepBase v20170127 with parameters: -species

Ruminantia -xsmall -s -no_is -cutoff 255 -frag 20000 -gff. Tandem repeats were

identified using Tandem repeat finder [64]. The identified tandem repeats as well as

those low complexity sequences identified by RepeatMakser were merged as low-

complexity regions (LCR). Of the LCR, regions of tandem repeat with the unit motif

length ≥7 bp was classified as VNTR and those with the unit motif length ≤6 bp were

classified as STR. The remaining LCR were classified as Other-LCR.

SV calling

We integrated the results from pan-genome analysis and alignment-based SV calling

to generated a high-confidence SV calls. For pan-genome analysis, the SVs were

generated from our multiassebly graph genome using gfatools v0.5-r234 as we

described in previous section. For alignment-based SV calling, the SVs were detected

using pbsv tools v2.6.2 (https://github.com/PacificBiosciences/pbsv). The CCS reads

were aligned to the reference genome using pbmm2 to generate the alignment BAM

file. Then the pbsv discover module was used to identify signatures of structural

variation within each aligned BAM. The tandem repeat annotation file was provided

to increase sensitivity and recall of this step. Then the signatures of all the samples

were fed to pbsv call module to jointly call SV from structural variant signatures

using the –ccs mode.

Mendelian error rate estimation

A Mendelian inheritance error represents a genotype in the offsprint that cannot be

inherited from either of their paraments. The whole genome sequencing data of 11

trios were from our recently study for allelic specific expression analysis [38]. For

each trio, the Mendelian error rate is the proportion of SVs from the children that

deviate from Mendelian inherence.

SV hotspot analysis

SV hotspot regions were determined using primatR package "hotspotter" section

(bw=200000, num.trial=1000, pval<1e-8) [65]. SVs were significantly clustered at the

terminal 5 Mb region of each chromosome (pvalue < 0.01, permutation test, random

shuffling of 1000 times). The random shuffling of SV intervals were performed using

Bedtools v2.29.0 with the shuffle command [66].

Enrichment of SVs in the genome

The H3K4me3 and H3K27ac peak regions were retrieved from our Ruminant genome

database. We only used insertions and deletions for enrichment. For insertions, the

end position is the start position plus the insertion length. We first recorded the

observed number of overlaps between SVs and UTR, exon, intron, H3K4me3 and

H3K27ac peak regions, respectively. Then we counted the average number of

overlaps between randomly shuffled SVs (1000 times) and each genomic region.

SV genotyping in resequenced samples

We collected whole genome sequencing data for 645 domestic sheep, from four

geographic regions (Europe, East Asia, Middle east and Africa) as well as wild sheep

including 33 Asiatic mouflons (*Ovis aries musimon*) and 8 argali sheep (*Ovis ammon*).

For the domestic sheep, those breeds with at least five individuals remain unchanged

whereas those samples from one country were grouped into one population if each of

the breed is composed of less than 5 individuals. In this way, all the domestic sheep

were assigned to 33 breeds/populations (Supplementary data 6). We genotyped SVs in

these samples using Paragraph v2.4a [28]. The maximum permitted read count for

each variant were set to 20 times the average sample depth in order to reduce runtime

for repetitive regions. The resulting genotype files in vcf format from all samples

were then combined using bcftools v1.9 [67].

To obtain a high-confidence call set of SVs, the resulting genotypes were filtered

based on overall genotyping rates and Hardy-Weinberg equilibrium [29]. Those

variants that were successfully genotyped in ≥10% of all samples were removed using

vcftools [68]. We also estimated Hardy-Weinberg equilibrium for each variant using

vcftools and those shows excess of heterozygotes (pvalue< 1×10-5) were removed.

Quantifying linkage disequilibrium (LD) of SVs with surrounding SNPs

In order to measure the extent to which SVs are correlated with SNPs, we calculated

LD (r^2 values) between the polymorphic SVs (MAF > 0.05) and the neighboring

polymorphic SNPs (MAF > 0.05; 150 upstream and 150 downstream SNPs) in

Mouflon and domestic sheep. SNPs located on SVs were not considered. Plink v1.9

was used to calculate LD for all pairwise cases for SV-SNP and the surrounding SNP-

SNP variant pairs. We then calculated degree of LD between each SV and flanking

SNPs following a previous study [37]. We first obtained pairwise genotype

correlations (r^2) for all SNP-SNP and SNP-SV pairs. For each of the 300 ranked

surrounding positions, the number of times the SV rank was greater than the SNP-

SNP median rank was recorded as a relative LD metric of SV to SNP. Low, middle or

high LD of each SV were then classified if the number of exceed was less than 100,

between 100 and 200, and greater than 200, respectively.

Inferring the derived SVs and *de novo* SVs in domestic sheep

To infer the state (ancestral or derived) for each SV, we additionally genotyped one

snow sheep (12.11×, ERR4161992) and one bighorn sheep (10.3×, SRR501898). In

this way, we used three wild sheep samples from one argali, one snow sheep, one

bighorn sheep as outgroups. For each biallelic SV, if the same homozygous genotypes

were detected in at least two outgroups, this state is defined as the ancestral state

while the other is the derived state. If the derived allele is only present in domestic

sheep but absent in all the wild sheep (8 Argali and 33 Asiatic mouflons), the SV was

defined as a *de novo* SV in domestic sheep.

Detecting selection signals

In order to reveal selection signals in fat tail sheep, we used the method of population

branch statistic (PBS) to calculate and compare the Fixation index (Fst) between the

fat tail sheep, thin tail sheep and Asiatic mouflon, using the Asiatic mouflon as a

distantly related population. The Fst value between populations with different traits

was calculated using the python scikit-allel package v1.3.2

(https://github.com/cggh/scikit-allel). We calculated the Fst values of both SNP and

SV sets. Fst values for SNPs were calculated with a 10 kb sliding window with a 5 kb

step.

To scan positive selected loci between different breeds, we employed two

strategies, including the global Fst and the Ohana method which identify extreme

allele frequency differences by modelling ancestral admixture components. The

global Fst was calculated following previously described approach with modifications

[69]. For each SV site, we calculated the Fst values between all populations. Instead

of using the average of all Fst values, we took the average of the top 50% Fst values

as the global FST value for each SV. The top 1% of the highest global Fst value by

this method were considered as putative selective signals. For Ohana method, we first

modeled the dataset with the provided workflow using the SNPs [29]. The resulting

covariance was used as the neutral input for scans of selection. For each specific

ancestry component, Ohana reported a likelihood ratio statistic (LRS) to quantify the

likelihood of selection for each variant. We varied the number of ancestral admixture

components from K=2 to K=5 to find selection signals.

Detecting expression-associated SVs

The RNA-seq data were downloaded from NCBI SRA database (accession number

PRJNA485657) from 14 Texel (♂)×Kazakh (♀) sheep hybrids from 8 tissues

including liver, rumen, tail fat, skin, subcutaneous fat, semitendinosus,

semimembranosus, longissimus dorsi [38, 39]. The raw reads were first trimmed for

adapter and low-quality sequences using Trimmomatic [70] and the high-quality reads

were aligned to sheep reference genome using Hisat2 [71]. Reads counts per gene

were computed using featureCounts [72] and normalized to counts per million (CPM)

using edgeR package [73] and those with CPM > 1 in at least two samples were

retained.

To identify expression-associated SVs, we first performed SV-eQTL mapping by

measuring the association between SV genotypes and expression level of nearby

genes. We defined a list of SV-gene pairs based on the distance between SV and

nearby genes within 200 kb. Only the SVs with presence and absence in at least 3

samples, respectively, were considered. For each of the SV-gene pairs, the expression

difference of each gene was compared between those with and without SV using a

two-sided Mann-Whitney U test. A P-value threshold of < 0.05 was applied to identify

candidate SV-eQTL. The candidate expression-associated SV was required to be

repeatedly found in at least two tissues.

It is expected that the regulatory SVs could cause allelic specific expression. We

also performed allelic specific expression (ASE) mapping by integrating the RNA-seq

data as well as the SV genotyping information for each of the samples. The ASE

genes were identified following the pipeline described in our previous study [38].

First, for each sample, the ASE SNPs displaying allelic imbalance at the gene body

were identified from RNA-seq data using the following parameters: (1) at least 20

total reads; (2) allele ratio (allele reads count / total reads count)> 0.65 or <0.35; (3)

significant allelic imbalances with FDR<0.05. Those with at least two ASE SNPs at

the gene body and were found in at least three samples were defined as candidate ASE

gene. For each SV-gene pair, if the ASE gene matches with the heterozygous state for

the associated SV in at least five RNA-seq samples, we considered the SV as

putatively expression-associated.

Analyzing the association between the 169-bp insertion with tail length

The association analysis was conducted in a F2 population which was generated by

backcrossing East Friesian sheep (♂) with the female Hu sheep (♀) x East Friesian

sheep (♂). We selected 211 individuals at the same age (six months old) to measure

the tail length and collect DNA samples for each individual. Since the insertion was

dominant in East Friesian sheep with allele frequency of 0.92 but is completely absent

in Hu sheep based on our resequencing data, the F2 population that we studied only

possessed two genotypes (II and ID) due to the grading-up using East Friesian sheep.

Agarose electrophoresis was used to determine genotypes of the PCR products.

Primers were designed in the flanking region of the insertion (forward:

TTTATGAGCTTCTCTCCGCCA; reverse: AAGTGGTATAATTGCCGGGCT). PCR

amplification was performed with 94°C for 2 min; 94°C for 30 s, 62°C for 30 s, and

72°C for 30 s for 28 cycles; 72°C for 2 min; and a hold at 4°C.

Luciferase reporter assay

To analyze promoter activity of the SV variation by luciferase assay, fragments with

(mutation) and without (wild type) the insertion was synthetized by Tsingke

Biotechnology Co., Ltd. (Beijing, China). Each fragment was cloned into the pGL3-

Basic vector using Sac I and Hind□ sites to assay its promoter activity on the

downstream gene.

293T cells in 96-well plates were transiently transfected with 200 ng Firefly

luciferase vector (pGL3) and 4 ng pRL-TK Renilla luciferase vector (Promega,

Madison, USA) using 0.5 uL Lipofectamine 3000 (Thermo Fisher Scientific, Waltham,

USA). Twenty-four hours after transfection, luciferase activities were measured with a

dual-luciferase reporter assay kit (Promega, Madison, USA). Renilla luciferase

activity was used as the transfection control for normalizing Firefly luciferase activity.

Empty pGL3-Basic vector was used as the reference for promoter activity assays.

Empty pGL3-Control vector (Control) was used as the positive control for promoter

activity. The data are presented as means with standard deviation.

660 Authors' contributions

Y.J. and S.Q.G. conceived and designed the experiments. R.L., X.M.Z. and Z.Y.L.

performed *de novo* assembly and SV calling; Z.Y.L., M.S.X. and Y.F.Z. analyzed the

RNA-seq and Chip-seq data; F.W., M.G. and L.Z. performed the ASE analysis; R.L.,

Z.Y.L., W.W.Fang and X.L.D. carried out SV genotyping and population genetics

analysis; Y.T.Y. and X.Y.L. conducted the luciferase reporter assay; Z.B.Z., W.W.Fu,

H.H.Z., C.N.C., P.Y., Z.A.G., N.J.N., H.A.N., X.P.Y., Y.X.S. and W.D.D. collected the

samples and prepared the WGS data; R.L. wrote the manuscript and Y.J., S.Q.G.,

J.A.L., P.H., E.I.A., R.D.X. and X.H.W. revised it. All authors read and approved the

final manuscript.

670 Acknowledgements

671 We thank the High-Performance Computing platform of Northwest A&F University

672 for providing computing resources. We would also like to thank Prof. Xiaolong Wang

673 for providing the nice chart of different sheep tail patterns.

674 Competing interests

675 The authors declare no competing interests.

676 Ethics statement

Blood samples were taken by conforming with the Helsinki Declaration of 1975 (as

revised in 2008) concerning Animal Rights.

Data availability

Sequences and metadata generated in this work are publicly available. The *de novo*

assemblies, HiFi data and WGS data for the 13 individuals were available at NCBI:

WDU1 (PRJNA788705, PRJNA788704, PRJNA784371),QK1 (PRJNA788680,

PRJNA788679, PRJNA784330), ROM1 (PRJNA788688, PRJNA788687,

PRJNA784357), SFK1 (PRJNA788697, PRJNA788696, PRJNA784366), UJM1

(PRJNA788702, PRJNA788701, PRJNA784376), CHA1 (PRJNA788666,

PRJNA788665, PRJNA784396), YUN1 (PRJNA788712, PRJNA788711,

PRJNA784413), TEX1 (PRJNA788214, PRJNA788213, PRJNA780104), DOR1

(PRJNA788571, PRJNA788570, PRJNA782680), KER1 (PRJNA788676,

PRJNA788675, PRJNA783516), MER1 (PRJNA788672, PRJNA788671,

PRJNA783537), KAZ1 (PRJNA788564, PRJNA788563, PRJNA783179), FRI1

(PRJNA791176, PRJNA791175, PRJNA721520). The details are provided in

Supplementary table 1.

Funding

This study was supported by research grants from the National Natural Science

Foundation of China to Y.J. (U21A20247, 31822052), S.Q.G. (31760662, 31860639)

and R.L. (31802027), China Postdoctoral Science Foundation to W.W.Fu (No.

2021M702690).

703 **Tables**

704 **Table 1** Quality assessments of the primary and partially phased assemblies for the 13

705 individuals.

Breeds	Abbr.	Assembly	Assembly length (Gb)	Contig N50 (Mb)	Contig number	Depth	BUSCO (%)	Base accuracy
Kazak	KAZ1	Primary	2.87	73.4	1849	22.4	93.4	46.3
		hap1/hap2	2.85/2.84	10.7/13.9		-	92.3/91.9	46.6/46.4

Kermani	KER1	Primary	2.82	80.3	1675	21.7	93.8	45.2
		hap1/hap2	2.77/2.75	15.4/15.2		-	90.7/92.2	45.4/45.6
Dorset	DOR1	Primary	2.91	92.4	1293	25.0	94.3	45.5
		hap1/hap2	2.86/2.83	32.8/39.5		-	93.4/92.7	45.6/45.6
East Friesian	FRI1	Primary	2.90	85.1	982	28.0	94.4	48.2
		hap1/hap2	2.77/2.75	62.9/53.2		-	91.3/92.8	48.4/48.5
Chiniese Merino	MER1	Primary	2.84	60.0	1773	18.7	94.2	43.3
		hap1/hap2	2.82/2.83	3.7/4.8		-	92.1/90.0	42.4/43.3
Qiaoke	QK1	Primary	2.87	75.0	1654	21.5	94.0	47.1
		hap1/hap2	2.83/2.84	13.6/9.9		-	92.6/93.4	47.6/47.2
Romney	ROM1	Primary	2.86	68.3	1552	19.3	94.4	43.2
		hap1/hap2	2.81/2.82	6.3/5.7		-	92.1/92.4	43.3/43.1
Suffolk	SFK1	Primary	2.88	65.3	1513	18.8	94.3	43.4
		hap1/hap2	2.86/2.80	5.7/4.9		-	92.4/92.5	43.2/43.3
Texel	TEX1	Primary	2.82	47.6	1835	17.9	94.4	44.6
		hap1/hap2	2.76/2.75	4.52/4.49		-	91.4/90.2	44.8/44.8
White Dorper	WDU1	Primary	2.81	17.8	2125	15.4	93.8	41.7
		hap1/hap2	2.82/2.77	1.58/1.26		-	90.2/89.8	41.9/41.8
Ujumqin	UJM1	Primary	2.87	75.6	1534	21.3	94.4	44.4
		hap1/hap2	2.86/2.84	11.5/11.7		-	93.9/93.5	44.5/44.6
Charollais	CHA1	Primary	2.86	71.5	1427	19.2	94.1	44.1
		hap1/hap2	2.82/2.80	6.4/5.1		-	91.7/92.1	44.3/44.3
Yunnan	YUN1	Primary	2.91	73.9	1293	22.0	94.1	46.9
		hap1/hap2	2.89/2.85	18.7/13.1		-	93.6/93.4	46.7/46.6

Figure legends

Fig. 1 Ovine graph pan-genome construction. (A) PCA plot shows the representation of the genetic diversity of domestic sheep by the 13 individuals used for HiFi sequencing and pan-genome construction. (B) The pan-genome size changes with the increase in number of genome assemblies. (C) Repeat annotation of the structural variations stratified by SV types. (D) The ovine graph pan-genome reveals diverse SV types. (E) An example of divergent allele of 50757 bp containing a MYADM-like repeat. (F) An example of multiallelic variations containing a large deletion of

721 105,650 bp which is nested with another insertion of 5,518 bp.

722 **Fig. 2** The repertoire of structural variations in sheep genome and frequency spectrum

723 in domestic and wild sheep. (A) Length distribution of insertions and deletions. (B)

724 Geographical distribution of the 645 domestic sheep and 33 Asiatic mouflons. Due to

725 space limitation, dorper sheep from South Africa is not shown. (C) Derived allele

726 frequency of the insertions and deletions. (D) Proportion of standing and de novo SVs.

727 (E) High-frequency de novo SVs in domestic sheep. (F) A de novo insertion within

728 GRID2 gene rises to high frequency in all geographical populations. (G) A de novo

729 insertion of a retrogene rises to high frequency in European sheep.

730 **Fig. 3** Population-stratified SVs and their effects on gene expression. (A) Genome-

731 wide distribution of global Fst which is measured by the average value of top 50% Fst

732 values for each SV across 33 sheep breeds. (B) Examples of expression-associated

733 SVs identified by SV-eQTL mapping. (C) Examples of expression-associated SVs

734 identified by allelic specific expression (ASE) mapping. Each column represents one

735 ASE SNP in exons and each row represent one RNA-seq sample. (D) Frequency

736 distribution of the six expression-associated SVs in 33 breeds/populations. CNM,

737 Chinese Merino; DEM, German merino; EFR, East Friesian Dairy; ESO, Spain; FIN,

738 Finnsheep; GTL, Gotland; NOW, Norway; OUE, Ouessant; POD, Poll Dorset; RUS,

739 Russia; SEL, Shetland; SFK, Suffolk; SOG, Solognote; WSF, Suffolk White; ALT,

740 Altay; BSB, Bashibai; BYK, Bayinbuluke; CLB, Cele Black; DUL, Duolang; HUS,

741 Hu; KAZ, Kazakh; LTH, Large-tailed Han; SSS, Sishui Fur; STH, Small-tailed Han;

742 TAN, Tan; UJM, Ujumqin; WAG, Waggir; WDS, Wadi; OLA, Oula; TIB, Tibetan;

743 YBB, Yunnan; AFS, Afshari; MDE, Middle East; CND, Dorper; MAOA, Morocco;

744 MOF, Mouflon.

745 **Fig. 4** Selective SVs associated with long tail trait. (A) The genome-wide Fixation

746 index (Fst) of long fat-tailed sheep versus short fat-tailed sheep from Asia breeds and

747 long thin-tailed sheep versus short thin-tailed sheep from Europe breeds. (B) The

748 selective haplotype in long-tailed sheep. The 169bp-insertion is close to the 5' UTR

749 region of HOXB13 gene. The haplotypes include the genotypes of the insertion and

750 surrounding SNPs. (C) The frequency of the insertion in wild sheep and domestic

sheep. LTH, Large-tailed Han; TAN, Tan; SSS, Sishui Fur; EFR, East Friesian Dairy;

SOG, Solognote; DEM, German merino; SFK, Suffolk; HUS, Hu; STH, Small-tailed

Han; UJM, Ujumqin; SEL, Shetland; FIN, Finnsheep; OUE, Ouessant; GTL, Gotland.

(D) RNA-seq data shows that the insertion belongs to the 5' UTR region. The top

panel corresponds to the expression in colon of Texel x Scottish Blackface sheep

(ERS2107126) presumably carrying at least one copy of the insertion; the bottom

panel corresponds to the expression in colon of one small-tail sheep (SRR5122597)

presumably absence of the insertion. (E) The electrophoresis diagrams of PCR

products for the three genotypes (homologous insertion: II, heterozygote: ID,

homologous deletion: DD). (F)The carriers and non-carriers of the insertion differs in

tail length. (G) HOXB13 promoter activity modulated by the 169 bp insertion.

Luciferase activity in HEK293T cells transfected with recombinant plasmids

containing the insertion or not. Each experiment was repeated at least five times. The

data were the mean ± standard error of the normalized luciferase activity. **P-

value<0.01.

Fig. 5 Selective SVs associated with fat tail trait. (A) Population branch statistic (PBS)

values across the whole genome by comparing fat tailed sheep to thin tailed sheep

using the mouflon sheep as outgroup. (B) The two most differentiated regions

between fat tailed sheep and thin tailed sheep. The left panel corresponds to the

intergenic region between BMP2 and HAO1 gene while the right panel corresponds to

the region surrounding PDGFD gene. Those SVs with PBS value >0.8 were

highlighted. (C) The haplotypes of mouflons and domestic sheep for the two most

selective regions. Each column represents one SV and each row represent one

individual. The black reverted triangles represent *de novo* SVs.

Supplementary files

Supplementary table 1 Accession numbers of the *de novo* assemblies.

Supplementary data

Supplementary data 1 Information of divergent variations

Supplementary data 2 Information of multiallelic variations

Supplementary data 3. List of the structural variations including insertion, deletion,

inversion and duplications

Supplementary data 4 SV frequency in mouflon sheep and domestic sheep

Supplementary data 5 List of putatively expression associated SVs

Supplementary data 6 Sample list of the resequencing data

Supplementary figures

Supplementary fig. 1 Example of three representative divergent variations

intersecting with IDE (a), PRP3-like (b) and MAGI2 (c), respectively.

Supplementary fig. 2 The multiallelic region harboring beta hemoglobin genes.

Supplementary fig. 3 Example of three representative multiallelic variations

affecting ASIP (a), UDP-glucuronosyltransferase 2B18-like (b) and olfactory receptor

(LOC101115700) (c), respectively.

Supplementary fig. 4 The number of SVs intersecting functional elements compared

to randomly permuting SV locations.

Supplementary fig. 5 An ideogram showing SV hotspots using all SVs. The total

number of SVs in each detected hotspot is shown by a scale going from blue to red.

Supplementary fig. 6 Enrichment analysis of SVs with respect to the last 5 Mb of

each chromosome end.

Supplementary fig. 7 PCA plot showing the clusters of domestic sheep by insertions

(a) and deletions (b).

Supplementary fig. 8 Histogram of the number of SV ranks (0-300) of r2 values that

surpass the median SNP-based r2 value for common SVs (MAF > 0.05).

Supplementary fig. 9 Enriched KEGG pathways for the population-stratified SVs.

Supplementary fig. 10 RNA-seq mapping for the 169 bp insertion shows that the
insertion belongs to 5' UTR region.

References:

[1] Chiang C, Scott A J, Davis J R, Tsang E K, Li X, Kim Y, et al. The impact of

structural variation on human gene expression. Nat Genet. 2017;49:692-9.

[2] Sudmant P H, Rausch T, Gardner E J, Handsaker R E, Abyzov A, Huddleston J, et

al. An integrated map of structural variation in 2,504 human genomes. Nature.

2015;526:75-81.

[3] Mahmoud M, Gobet N, Cruz-Dávalos D I, Mounier N, Dessimoz C, Sedlazeck F

J. Structural variant calling: the long and the short of it. Genome Biol. 2019;20:246.

[4] Sedlazeck F J, Lee H, Darby C A, Schatz M C. Piercing the dark matter:

bioinformatics of long-range sequencing and mapping. Nat Rev Genet. 2018;19:329-

46.

[5] Audano P A, Sulovari A, Graves-Lindsay T A, Cantsilieris S, Sorensen M, Welch

A E, et al. Characterizing the Major Structural Variant Alleles of the Human Genome.

Cell. 2019;176:663-75.e19.

[6] Ebert P, Audano P A, Zhu Q, Rodriguez-Martin B, Porubsky D, Bonder M J, et al.

Haplotype-resolved diverse human genomes and integrated analysis of structural

variation. Science. 2021:eabf7117.

[7] Qin P, Lu H, Du H, Wang H, Chen W, Chen Z, et al. Pan-genome analysis of 33

genetically diverse rice accessions reveals hidden genomic variations. Cell. 2021.

[8] Liu Y, Du H, Li P, Shen Y, Peng H, Liu S, et al. Pan-Genome of Wild and

Cultivated Soybeans. Cell. 2020;182:162-76.e13.

[9] Alonge M, Wang X, Benoit M, Soyk S, Pereira L, Zhang L, et al. Major Impacts

of Widespread Structural Variation on Gene Expression and Crop Improvement in

Tomato. Cell. 2020;182:145-61.e23.

[10]Tuzun E, Sharp A J, Bailey J A, Kaul R, Morrison V A, Pertz L M, et al. Fine-

scale structural variation of the human genome. Nat Genet. 2005;37:727-32.

[11]Golicz A A, Bayer P E, Bhalla P L, Batley J, Edwards D. Pangenomics Comes of

Age: From Bacteria to Plant and Animal Applications. Trends Genet. 2020;36:132-45.

[12]De Coster W, Van Broeckhoven C. Newest methods for detecting structural

variations. Trends Biotechnol. 2019;37:973-82.

[13] Wenger A M, Peluso P, Rowell W J, Chang P-C, Hall R J, Concepcion G T, et al.

Accurate circular consensus long-read sequencing improves variant detection and

assembly of a human genome. Nat Biotechnol. 2019;37:1155-62.

[14] Cheng H, Concepcion G T, Feng X, Zhang H, Li H. Haplotype-resolved de novo

assembly using phased assembly graphs with hifiasm. Nat Methods. 2021;18:170-5.

[15] Nurk S, Walenz B P, Rhie A, Vollger M R, Logsdon G A, Grothe R, et al. HiCanu:

accurate assembly of segmental duplications, satellites, and allelic variants from high-

fidelity long reads. Genome Res. 2020;30:1291-305.

[16] Talenti A, Powell J, Hemmink J D, Cook E A J, Wragg D, Jayaraman S, et al. A

cattle graph genome incorporating global breed diversity. bioRxiv.

2021:2021.06.23.449389.

[17] Leonard A S, Crysnanto D, Fang Z-h, Heaton M P, Vander Ley B L, Herrera C, et

al. Bovine pangenome reveals trait-associated structural variation from diverse

assembly inputs. bioRxiv. 2021:2021.11.02.466900.

[18] Norris B J, Whan V A. A gene duplication affecting expression of the ovine ASIP

gene is responsible for white and black sheep. Genome Res. 2008;18:1282-93.

[19] Demars J, Cano M, Drouilhet L, Plisson-Petit F, Bardou P, Fabre S, et al.

Genome-Wide Identification of the Mutation Underlying Fleece Variation and

Discriminating Ancestral Hairy Species from Modern Woolly Sheep. Mol Biol Evol.

2017;34:1722-9.

[20] Wiedemar N, Drögemüller C. A 1.8-kb insertion in the 3ʹ-UTR of RXFP2 is

associated with polledness in sheep. Anim Genet. 2015;46:457-61.

[21] Li X, Yang J, Shen M, Xie X-L, Liu G-J, Xu Y-X, et al. Whole-genome

resequencing of wild and domestic sheep identifies genes associated with

morphological and agronomic traits. Nat Commun. 2020;11:2815.

[22] Li H, Feng X, Chu C. The design and construction of reference pangenome

graphs with minigraph. Genome Biol. 2020;21:265.

[23] Gonzalez M V, Mousel M R, Herndon D R, Jiang Y, Dalrymple B P, Reynolds J

O, et al. A Divergent Artiodactyl MYADM-like Repeat Is Associated with Erythrocyte

Traits and Weight of Lamb Weaned in Domestic Sheep. PLOS ONE. 2013;8:e74700.

[24]French A T, Bethune J A, Knight P A, McNeilly T N, Wattegedera S, Rhind S, et

al. The expression of intelectin in sheep goblet cells and upregulation by interleukin-4.

Vet Immunol Immunopathol. 2007;120:41-6.

[25]Jiang Y, Wang X, Kijas J W, Dalrymple B P. Beta-globin gene evolution in the

ruminants: evidence for an ancient origin of sheep haplotype B. Anim Genet.

2015;46:506-14.

[26]Durbin R M, Altshuler D, Durbin R M, Abecasis G R, Bentley D R, Chakravarti

A, et al. A map of human genome variation from population-scale sequencing. Nature.

2010;467:1061-73.

[27]Quan C, Li Y, Liu X, Wang Y, Ping J, Lu Y, et al. Characterization of structural

variation in Tibetans reveals new evidence of high-altitude adaptation and

introgression. Genome Biol. 2021;22:159.

[28]Chen S, Krusche P, Dolzhenko E, Sherman R M, Petrovski R, Schlesinger F, et al.

Paragraph: a graph-based structural variant genotyper for short-read sequence data.

Genome Biol. 2019;20:291.

[29]Yan S M, Sherman R M, Taylor D J, Nair D R, Bortvin A N, Schatz M C, et al.

Local adaptation and archaic introgression shape global diversity at human structural

variant loci. eLife. 2021;10:e67615.

[30]Kalkan Z, Durasi İ M, Sezerman U, Atasever-Arslan B. Potential of GRID2

receptor gene for preventing TNF-induced neurodegeneration in autism. Neurosci Lett.

2016;620:62-9.

[31]Zhu X-J, Liu Y, Dai Z-M, Zhang X, Yang X, Li Y, et al. BMP-FGF signaling axis

mediates Wnt-induced epidermal stratification in developing mammalian skin. PLoS

Genet. 2014;10:e1004687.

[32]Economides K D, Zeltser L, Capecchi M R. HOXB13 mutations cause

overgrowth of caudal spinal cordand tail vertebrae. Dev Biol. 2003;256:317-30.

[33]Kardos M, Luikart G, Bunch R, Dewey S, Edwards W, McWilliam S, et al.

Whole genome resequencing uncovers molecular signatures of natural and sexual

selection in wild bighorn sheep. Mol Ecol. 2015;24:5616-32.

[34]Chen C, Liu C, Xiong X, Fang S, Yang H, Zhang Z, et al. Copy number variation

in the MSRB3 gene enlarges porcine ear size through a mechanism involving miR-584-5p. Genet Sel Evol. 2018;50:1-18.

[35]Pan Z, Li S, Liu Q, Wang Z, Zhou Z, Di R, et al. Rapid evolution of a retro-transposable hotspot of ovine genome underlies the alteration of BMP2 expression and development of fat tails. BMC Genomics. 2019;20:1-15.

[36]Nishikawa S, Kusakabe M, Yoshinaga K, Ogawa M, Hayashi S-l, Kunisada T, et al. In utero manipulation of coat color formation by a monoclonal anti□c□kit antibody: two distinct waves of c□kit□dependency during melanocyte development. The EMBO journal. 1991;10:2111-8.

[37]Yang N, Liu J, Gao Q, Gui S, Chen L, Yang L, et al. Genome assembly of a tropical maize inbred line provides insights into structural variation and crop improvement. Nat Genet. 2019;51:1052-9.

[38]Wang F, Shao J, He S, Guo Y, Pan X, Wang Y, et al. Allele-specific expression and splicing provides insight into the phenotypic differences between thin- and fat-tailed sheep breeds. Journal of Genetics and Genomics. 2021.

[39]Pan X, Cai Y, Li Z, Chen X, Heller R, Wang N, et al. Modes of genetic adaptations underlying functional innovations in the rumen. Sci China Life Sci. 2021;64:1-21.

[40]Nanjappa S G, Mudalagiriyappa S, Fites J S, Suresh M, Klein B S. CBLB Constrains Inactivated Vaccine–Induced CD8+ T Cell Responses and Immunity against Lethal Fungal Pneumonia. The Journal of Immunology. 2018;201:1717-26.

[41]Gaudet R G, Zhu S, Halder A, Kim B-H, Bradfield C J, Huang S, et al. A human apolipoprotein L with detergent-like activity kills intracellular pathogens. Science. 2021;373:eabf8113.

[42]Lascano J, Uchil P D, Mothes W, Luban J. TRIM5 retroviral restriction activity correlates with the ability to induce innate immune signaling. J Virol. 2015;90:308-16.

[43]Iwaki A, Moriwaki K, Sobajima T, Taniguchi M, Yoshimura S i, Kunii M, et al. Loss of Rab6a in the small intestine causes lipid accumulation and epithelial cell death from lactation. The FASEB Journal. 2020;34:9450-65.

[44]Han J, Li E, Chen L, Zhang Y, Wei F, Liu J, et al. The CREB coactivator CRTC2

controls hepatic lipid metabolism by regulating SREBP1. Nature. 2015;524:243-6.

[45]Matsubara T, Kida K, Yamaguchi A, Hata K, Ichida F, Meguro H, et al. BMP2

Regulates Osterix through Msx2 and Runx2 during Osteoblast Differentiation*. J Biol

Chem. 2008;283:29119-25.

[46]Lee E, Kim J Y, Kim T K, Park S Y, Im G I. Methyltransferase-like protein 7A

(METTL7A) promotes cell survival and osteogenic differentiation under metabolic

stress. Cell Death Discov. 2021;7:154.

[47]Sukalo M, Tilsen F, Kayserili H, Müller D, Tüysüz B, Ruddy D M, et al. DOCK6

mutations are responsible for a distinct autosomal-recessive variant of Adams-Oliver

syndrome associated with brain and eye anomalies. Hum Mutat. 2015;36:593-8.

[48]Winnepenninckx B, Debacker K, Ramsay J, Smeets D, Smits A, FitzPatrick D R,

et al. CGG-repeat expansion in the DIP2B gene is associated with the fragile site

FRA12A on chromosome 12q13.1. Am J Hum Genet. 2007;80:221-31.

[49]Fu W, Wang R, Nanaei H A, Wang J, Hu D, Jiang Y. RGD v2.0: a major update of

the ruminant functional and evolutionary genomics database. Nucleic Acids Res. 2021.

[50]Clark E L, Bush S J, McCulloch M E B, Farquhar I L, Young R, Lefevre L, et al.

A high resolution atlas of gene expression in the domestic sheep (Ovis aries). PLoS

Genet. 2017;13:e1006997.

[51]Chao T, Wang G, Ji Z, Liu Z, Hou L, Wang J, et al. Transcriptome Analysis of

Three Sheep Intestinal Regions reveals Key Pathways and Hub Regulatory Genes of

Large Intestinal Lipid Metabolism. Scientific Reports. 2017;7:5345.

[52]Jiang Y, Xie M, Chen W, Talbot R, Maddox J F, Faraut T, et al. The sheep genome

illuminates biology of the rumen and lipid metabolism. Science. 2014;344:1168-73.

[53]Pan Z, Li S, Liu Q, Wang Z, Zhou Z, Di R, et al. Rapid evolution of a retro-

transposable hotspot of ovine genome underlies the alteration of BMP2 expression

and development of fat tails. BMC Genomics. 2019;20:261.

[54]Baazaoui I, Bedhiaf-Romdhani S, Mastrangelo S, Ciani E. Genome-wide

analyses reveal population structure and identify candidate genes associated with tail

fatness in local sheep from a semi-arid area. Animal. 2021;15:100193.

[55]Dong K, Yang M, Han J, Ma Q, Han J, Song Z, et al. Genomic analysis of

worldwide sheep breeds reveals PDGFD as a major target of fat-tail selection in sheep.

BMC Genomics. 2020;21:1-12.

[56]Ho S S, Urban A E, Mills R E. Structural variation in the sequencing era. Nat Rev

Genet. 2020;21:171-89.

[57]Miga K H, Wang T. The Need for a Human Pangenome Reference Sequence.

Annual Review of Genomics and Human Genetics. 2021;22:81-102.

[58]Li R, Yang P, Dai X, Asadollahpour Nanaei H, Fang W, Yang Z, et al. A near

complete genome for goat genetic and genomic research. Genet Sel Evol. 2021;53:74.

[59]Alonge M, Soyk S, Ramakrishnan S, Wang X, Goodwin S, Sedlazeck F J, et al.

RaGOO: fast and accurate reference-guided scaffolding of draft genomes. Genome

Biol. 2019;20:224.

[60]Manni M, Berkeley M R, Seppey M, Simão F A, Zdobnov E M. BUSCO Update:

Novel and Streamlined Workflows along with Broader and Deeper Phylogenetic

Coverage for Scoring of Eukaryotic, Prokaryotic, and Viral Genomes. Mol Biol Evol.

2021;38:4647-54.

[61]Crysnanto D, Leonard A S, Fang Z-H, Pausch H. Novel functional sequences

uncovered through a bovine multiassembly graph. Proc Natl Acad Sci.

2021;118:e2101056118.

[62]Keilwagen J, Wenk M, Erickson J L, Schattat M H, Grau J, Hartung F. Using

intron position conservation for homology-based gene prediction. Nucleic Acids Res.

2016;44:e89.

[63]Haas B J, Salzberg S L, Zhu W, Pertea M, Allen J E, Orvis J, et al. Automated

eukaryotic gene structure annotation using EVidenceModeler and the Program to

Assemble Spliced Alignments. Genome Biol. 2008;9:R7.

[64]Benson G. Tandem repeats finder: a program to analyze DNA sequences. Nucleic

Acids Res. 1999;27:573-80.

[65]Bakker B, Taudt A, Belderbos M E, Porubsky D, Spierings D C J, de Jong T V, et

al. Single-cell sequencing reveals karyotype heterogeneity in murine and human

malignancies. Genome Biol. 2016;17:115.

[66]Quinlan A R, Hall I M. BEDTools: a flexible suite of utilities for comparing

genomic features. Bioinformatics. 2010;26:841-2.

[67]Narasimhan V, Danecek P, Scally A, Xue Y, Tyler-Smith C, Durbin R.

BCFtools/RoH: a hidden Markov model approach for detecting autozygosity from

next-generation sequencing data. Bioinformatics. 2016;32:1749-51.

[68]Danecek P, Auton A, Abecasis G, Albers C A, Banks E, DePristo M A, et al. The

variant call format and VCFtools. Bioinformatics (Oxford, England). 2011;27:2156-8.

[69]Kijas J W, Lenstra J A, Hayes B, Boitard S, Neto L R P, San Cristobal M, et al.

Genome-wide analysis of the world's sheep breeds reveals high levels of historic

mixture and strong recent selection. PLoS Biol. 2012;10:e1001258.

[70]Bolger A M, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina

sequence data. Bioinformatics. 2014;30:2114-20.

[71]Kim D, Langmead B, Salzberg S L. HISAT: a fast spliced aligner with low

memory requirements. Nat Methods. 2015;12:357-60.

[72]Liao Y, Smyth G K, Shi W. featureCounts: an efficient general purpose program

for assigning sequence reads to genomic features. Bioinformatics. 2013;30:923-30.

[73]Robinson M D, McCarthy D J, Smyth G K. edgeR: a Bioconductor package for

differential expression analysis of digital gene expression data. Bioinformatics.

2010;26:139-40.

1008

A

Long fat-tailed vs. Short fat-tailed

Long thin-tailed vs. Short thin-tailed

● SNP △ SV

B

REF HET ALT

chr11:37,502,704 - 37,547,639 bp

169 bp Insertion

Mouflon

C

Fat tail | Thin tail

Long tail

LTH TAN SSS EFR SOG DEM SFK

Short tail

HUS STH UJM SEL FIN OUE GTL

Wild

Mouflon

● REF ● ALT

D

chr11:37,525,005 HOXB13

169 bp

Long tailed Depth 400 0

Short tailed Depth 70 0

E

Marker II ID DD

bp
600
500
412
400 412
300
243
200 243
100

F

p<0.0001

Tail length (cm)

II (n=57) ID (n=144)

Genotype

G

**

Relative luciferase activity

HOXB13-WT HOXB13-MUT

A

PBS value

SNP ● SV △

Chromosome

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26

B

PBS value

Chr13

48400 Kb 48800 Kb 49200 Kb 49600 Kb 50000 Kb 50400 Kb

PBS value

Chr15

3300 Kb 3500 Kb 3700 Kb 3900 Kb 4100 Kb 4300 Kb

BMP2 *LOC101117953* *LOC101118207* *LOC101110166* *LOC101110438* *HAO1*

INS (638 bp) INS (295 bp) INS (7,728 bp) INS (947 bp)
INS (2,917 bp) DEL (73 bp)

LOC106991615 *PDGFD*

INS (867 bp) DEL (131 bp) DEL (157 bp)

Insertion
Deletion

C

Mouflon

Chr13:48,400,000 bp-50,400,000 bp

Chr15:3,300,000 bp-4,300,000 bp