Continental-scale Hyperspectral tree species classification in the National Ecological Observatory Network

Sergio Marconi[1] , Ben. G. Weinstein[1], Sheng Zou[3], Stephanie A. Bohlman[2], Alina Zare[3], Aditya Singh[4], Dylan Stewart[3], Ira Harmon[5], Ashley Steinkraus[1], Ethan P. White[1]

[1]Department of Wildlife Ecology and Conservation, University of Florida, Gainesville, Florida, USA

[2]School of Forest, Fisheries and Geomatics Sciences, University of Florida, Gainesville, Florida, USA

[3]Department of Electrical and Computer Engineering, University of Florida, Gainesville, Florida, USA

[4]Department of Agricultural & Biological Engineering, University of Florida, Gainesville, FL, USA

[5]Department of Computer and Information Sciences and Engineering, University of Florida, Gainesville, FL, USA

Abstract

Advances in remote sensing imagery and computer vision applications unlock the potential for developing algorithms to classify individual trees from remote sensing at unprecedented scales. However, most approaches to date focus on site-specific applications and a small number of taxonomic groups. This limitation makes it hard to evaluate whether these approaches generalize well across broader geographic areas and ecosystems. Leveraging field surveys and hyperspectral remote sensing data from the National Ecological Observatory Network (NEON), we developed a continental extent model for tree species classification that can be applied to the entire network including a wide range of US terrestrial ecosystems. We compared the performance of the generalized approach to models trained at each individual site, evaluating advantages and challenges posed by training species classifiers at the US scale. We evaluated the effect of geography, environmental, and ecological conditions on the accuracy and precision of species predictions.

On average, the general model resulted in good overall classification accuracy (micro-F1 score), with better accuracy than site-specific classifiers (average individual tree level accuracy of 0.77 for the general model and 0.72 for site-specific models). Aggregating species to the genus-level increased accuracy to 0.83. Regions with more species exhibited lower classification accuracy. Trees were more likely to be confused with congeneric and co-occurring species and confusion was highest for trees with structural damage and in complex closed-canopy forests. The model produced accurate estimates of uncertainty, correctly identifying trees where confusion was likely. Using only data from NEON this single integrated classifier can make predictions for 20% of all tree species found in forest ecosystems across the US, suggesting the potential for broad scale general models for species classification from hyperspectral imaging.

1. Introduction

Forest ecosystems play a central role in essential services like assessment of natural resources and biodiversity conservation at national scales (Wiens, 2016; Pecl et al., 2017), but understanding this role at the large range of scales over which ecological processes operate can be challenging. A common strategy to monitor biodiversity and biomass of forests at broad scales is to use field surveys of plots as a sample for landscape level characteristics (USDA Forest Service, 2001, Lawrence et al. 2010). Data collection within survey plots requires extensive effort, limiting even the most extensive national forest inventories to several thousand permanent plots sampled every few years (White et al., 2016), which can be too sparse for investigating the effects of local and landscape ecological properties on forest structure and diversity (Tomppo et al., 2008). Remote sensing can help bridge this gap between local and regional scales by providing individual tree level data at scales beyond what is feasible for traditional plot-level inventories (Anderson, 2018). Models linking remotely sensed imagery to field surveys can identify the location and species identity of individual trees (Henrys & Jarvis, 2019), alleviating the challenge of inferring local patterns from sparsely sampled data (Ayrey et al., 2019, Bastin et al., 2019, Kandare et al., 2017) for understanding species distributions and abundances.

Numerous approaches have been developed for pixel- or canopy-scale species-level classification using hyperspectral remote sensing based on exploiting differences in foliar properties and canopy structure (Mayra et al, 2021, Belgiu & Dragut, 2016, Ballanti et al., 2016, Ab Majid et al., 2016). In general, however, most approaches are site- and/or ecosystem-specific (Fassnacht et al., 2016). Site and/or ecosystem-specific techniques have the potential to limit the generality of algorithms and make it difficult to: 1) conduct research at regional to continental scales due to the lack of general models that can be applied across ecosystems; 2) identify rare or uncommon species due to limited data for training models, which often results in

studies focusing on a limited subset of common species; and 3) accurately apply the model

beyond the scale of the associated field data. Furthermore, training data from single site studies

often lack the full range of variation in spectral characteristics that can occur for each species

due to intraspecific variation and variability from the remote sensing acquisition process.

Developing species classification approaches that combine data from across broad geographic

extents has the potential to address these limitations and increase the utility of remote sensing

models in duplicated effort across sites and projects.

Developing species classification models that span geographic areas, forest types and

species pools poses a novel set of challenges. First, it requires building a library of coregistered

field and remote sensing data that includes data from multiple sites and ecoregions for training

and testing algorithms. Second, increasing the geographic extent of species classification risks

confusing species with similar spectral properties that do not overlap in their geographic

distributions. Third, combining data from multiple sites may introduce variation in spectral

reflectance due to differences in phenology (which affect leaf greenness) and environmentally

driven intraspecific variation, which affect leaf biochemistry (Sims & Gamon, 2002). Finally,

aggregating remote sensing data from multiple flights, sensors, and sites may increase variation

in spectral signatures due to complex sources of spatial and temporal variation that are linked,

but not limited to, acquisition dates, solar angles, ecosystems types and variation in sensor

calibrations (Pax-Lenney et al., 2001). Therefore, while there are many potential benefits to

more broad scale models for species classification, it is unclear how they will perform compared

to models developed for specific ecosystems.

Here we leverage newly available data from the National Ecological Observatory

Network (NEON) to develop a continental level model for tree species classification that can be

applied to the entire network and compare its performance to the traditional approach of building

individual models for each site. We used NEON remote sensing and field data on individual

trees at 30 terrestrial sites from Puerto Rico to Alaska, covering a wide range of ecoregions and

biomes across the United States (US). Several studies have developed species classification

models for NEON data, but all these studies focused on individual NEON sites (Scholl et al.,

2020, Fricker et al., 2019, Marrs & Ni-Meister, 2019, Marconi et al., 2020), or 2-3 sites in the

same region (Rana et al. 20XX, Graves et al. 2021), and a small number of species within each

site. Even outside of NEON, species classification efforts from airborne remote sensing focus on

site-specific algorithms, making it hard to assess how well these approaches transfer to different

regions (but see Rana et al., 2018). We build on this work to develop a general model that can

be applied across the entire National Ecological Observatory Network by connecting field-

identified tree stems to hyperspectral images then using an ensemble of machine learning

algorithms to identify species. This ensemble-based approach allows for leveraging the

strengths of different machine learning classifiers and provides effective ways to estimate the

uncertainty of predictions. Using this model, we (1) assess whether a general model approach

improves performance compared to separate models for each site, (2) determine the

importance of reflectance, geography, environmental and ecological conditions on the accuracy

and precision of species predictions; (3) evaluate the uncertainty in predictions; and (4) discuss

the potential for this general model to be used for ecological applications from monitoring

populations of individual species to understanding patterns of biodiversity.

2. Methods

2.1. Field Data

Vegetation structure field data (https://data.neonscience.org/data-

products/DP1.10098.001) were collected by the NEON terrestrial observatory system (TOS)

between 2015 and 2019 (Table S.1). This dataset, sampled from 400 m^2 plots distributed across

the landscape of each NEON site, includes information about individual trees' geolocation and

properties such as species identity, health status, canopy position, crown diameter, and tree

height. Vegetation structure plot locations are located randomly across the sites stratified by

vegetation type within each site with the aim of capturing landscape level biodiversity and

structural features at each site. Each subplot is assigned to an ecosystem type extracted from

the National Land Cover Dataset. For this study we used data from 30 of the 41 NEON sites

with partial to complete forest cover, encompassing 17 out of 18 ecoclimatic domains in the US

(Figure S.1). Data from the other NEON sites could not be used because field data was either

missing information about tree stem positions or the remote sensing imagery contained gaps in

the hyperspectral or missing information about the sensor angle at the time of data collection.

We only included individual stem data that met the following criteria: (a) the stem had a species

label assigned to it, (b) it was marked as "alive" and "tree" in the NEON field inventory, and (c) it

belonged to a species with more than 5 entries for the entire dataset. We also did not use stems

designated in the NEON vegetation structure data as fully shaded, shrubs or sapling, as these

stems are most likely not visible in the remote sensing imagery and would therefore be

erroneously paired with pixels belonging to species from neighboring overstory crowns. The

final dataset used for species classification consisted of 5867 individual trees of 80 species.

2.2. Remote sensing data

For this study we used the hyperspectral L3 data from the NEON Airborne Observatory

Platform (NEON, 2021). This data consists of 1km^2 tiles with 426 channels recording reflectance

in 5nm bands from 350 to 2450nm. Reflectance data was already atmospherically corrected

using the ATCOR-4 approach (Krause et al., 2011). Pixel resolution is 1m^2. We applied

bidirectional reflectance distribution function (BRDF) correction, topographic correction,

and L2 normalization to reduce the effect of peripheral light and non-Lambertian scattering with

the goal of minimizing variation in reflectance ascribable to flight path and airplane position

(Marconi et al., 2020). For all tiles (n = 4500), we used the same general parameterization to

define the BRDF kernel. We also dropped bands in the water absorption regions of the spectra

as well as the spectrometer's peripheral bands to reduce the effects of noise and artifacts.

Thus, the hyperspectral data was reduced to a total of 350 channels. In the models, we included

terrain elevation ($1m^2$ spatial resolution) along with the hyperspectral data because of

elevation's potential information in discriminating species within landscapes (Strahler et al.,

1978, Scholl et al., 2020). This data was derived from a LiDAR sensor mounted along with the

hyperspectral sensor on the aircraft, which was converted into a 1m spatial resolution raster,

and appended to the hyperspectral data as an additional band.

We assigned each individual tree from the filtered field dataset to a square clip of 16

pixels (4m crown diameter), centered around the stem's GPS coordinates. This threshold was

selected because it is smaller than more than 95% of individual tree crowns diameter measured

from the NEON vegetation structure dataset. We adopted this strategy to reduce the number of

mislabeled pixels at the edges of the crown that belong to neighboring trees, especially in dense

closed canopies. To remove shaded and non-vegetation pixels from these clips, we removed all

pixels with NDVI < 0.5 and low reflectance in the NIR (reflectance at 825nm < 0.2). Since stem

positions often do not match precisely with the center of the tree crown in the canopy, pixels will

sometimes be assigned to the wrong label (hereafter referred to as pixel pollution). To reduce

pixel pollution, we filtered out pixels that were much shorter than the maximum height of the

crown. These pixels are less likely to belong to the sunlit portion of the target crown or may

even measure the reflectance from neighboring shorter tree crowns, or the understory within a

gap in the target crown. We filtered out pixels that were ≥5m below the top height of the tree as

determined by the maximum height of the tree from the LiDAR data in the 16-pixel clip. Finally,

we removed stems with field GPS locations that fell within 3 meters of one another where the

stems belonged to different taxa to decrease the chance of confusing closely neighboring, and

potentially intermixed tree crowns of different species. After all these steps, the final, filtered

dataset used ~50,000 out of 200,000 initial pixels and 7464 out of ~21,000 crowns in the original

vegetation structure dataset.

Due to the large number of bands in hyperspectral data it is necessary to reduce the

number of features used in classifiers to reduce the number of correlated bands and limit the

potential for overfitting (Li et al., 2011). We reduced the 356 hyperspectral bands to 45 spectral

features using a hierarchical dimensionality reduction, consisting of clustering bands with similar

standardized distributions according to Kullback-Leibler divergence (KLD) (Zare et al., 2019).

We chose this approach because it is well suited as a dimensionality reduction tool for data in

the form of probability densities like hyperspectral reflectance spectra (Delicado, 2011) and

allows for a direct identification of the most informative spectral regions. For this study we used

15 spectral regions, and we used the minimum, mean and maximum of the reflectance in each

region as the hyperspectral features for the species classifier, resulting in 45 spectral predictors.

2.3. Site effects

To provide the model with information on site location that could be useful to reduce

confusion across species that do not co-occur in a site but are characterized by similar spectral

signatures, we included the latitude and longitude of the centroid of each site in the model. This

approach incorporates information on the proximity of different sites and can be readily

generalized to use outside of NEON. To help control for potential differences resulting from

variation in sensor calibration of the specific flight missions, which would be specific to each

site, we added a "site identifier" to the remote sensing features in the model. The site identifier

was calculated using a Leave-One-Out regression encoding strategy to transform site

categories into real positive numbers (https://contrib.scikit-learn.org/category_encoders; Wright

& König, 2019). The advantage of this approach over the commonly used one-hot-encoder (ref)

is that it compresses the information into a single feature (rather than one feature per site),

which is avoids undesired sparsity and potential overfitting due to a large number of encoded

classes (30 in this study) (Rodriguez et al., 2018). We used data in the training set to fit the

encoder and assigned its average value to each site category. The final model input for the

general model was hyperspectral features, elevation, latitude and longitude and site. For the

site-specific models, only spectral features and elevation were used.

2.4. Species classification

For both the general and site-specific models, we performed species classification at the

pixel level using an ensemble of five classifiers (Figure S.2): (1) a random forest classifier

(Belgiu & Dragut, 2016), (2) a k-nearest neighbors classifier (Laaksonen & Oja, 1996), (3) a

histogram gradient boosting classifier (Guryanov, 2019), (4) a fully connected multilayer

perceptron (Pacifico et al., 2018), and (5) a bagging classifier with support vector machine as

base estimators, using tools from the scikit-learn python package (Pedrosa et al., 2011). Details

for each classifier can be found in supplementary materials (Supplement 1: classifiers).

Ensemble-based approaches generally provide better performance and limit overfitting

compared to using one classifier alone (Knauer et al., 2019). We chose the individual classifiers

to ensemble because they have been shown to perform well for species classification on NEON

data (Marconi et al., 2019). All predictors were normalized for model fitting by subtracting the

mean and dividing by the standard deviation (i.e., setting the mean to zero and the standard

deviation to 1). Parameters for all models and the ensemble were extracted by performing cross

validation.

We used entropy loss to measure the quality of tree-splits for random forests, categorical

cross-entropy as the loss function for the histogram-gradient boosting, a radial basis function

kernel to allow for a non-linear decision surface for the support vector classifiers, and the

Manhattan distance for calculating the distance between k-nearest neighbors in the KNN

classifier. We stacked these five pixel-based models by using the probability vectors produced

by each classifier as features for a meta-ensemble elastic net logistic model (Tang et al., 2015,

Hui & Hastie, 2005). We chose this approach because logistic classifiers are easily interpretable

and use maximum likelihood to obtain estimates of the coefficients, returning as a result

confidence scores that match the probability of a label-match and not just the single best

predicted classification (Maddala, 1986), which is fundamental for assigning each prediction to

a reliable uncertainty score. Pairing predictions to robust estimates of uncertainty is fundamental

to increase the utility of remote sensing tree surveys for ecological analysis because it allows for

(1) selecting trees and areas that meet or exceed minimum confidence in the derived measures

for being used for scientific analyses, and (2) allows for cascading the uncertainty in predictions

onto the results of analyses downstream (Dietze, 2017). Training the logistic meta-ensemble on

calibrated scores from sub-classifiers offers an advantage over other modern algorithms, whose

estimates of uncertainty do not match true probabilities and are not well calibrated to the output

of interest (Guo et al., 2017, Mukhati et al. 2020).

One of the main challenges of species classification algorithms is the imbalance

between number of individual samples for rare and common species, which can cause models

to overfit to highly abundant classes. In our data set, the number of pixels per species ranged

from 44-28000 and the number of individual trees per species ranged from 5-1000. We used

SMOTETomek technique (Batista et al., 2003) to reduce the effects of species class

imbalances in the training set. SMOTETomek consists of a combination of under and

oversampling which resulted in roughly 1000 spectral signatures (pixels) per species. First we

undersampled pixels from the most abundant species using Tomek links, which removes noisy

and borderline pixels (Tomek, 1976). Then, we used a SMOTE oversampling approach (Chawla

et al., 2002) to create non-identical synthetic pixels for any species with fewer pixels than the

majority class, thus balancing each class to roughly 1000 pixels each. No over-undersampling

was applied to the test data. Because of the stratified design of the train-test split, most species

and sites had a number of test trees proportional to their frequency in the original dataset. We

also used the same train-test split to repeat the entire analysis once for each NEON site by

building and testing site-specific models built using only data from each particular site.

2.5.   Evaluation

We evaluated the performance of the models by training the model on 80% of the data

and evaluating its performance on the remaining 20%. Since spatial autocorrelation across train

and test data can lead to optimistic bias in classification (Millard & Richardson 2015), we placed

all individuals within a plot together into either the training or testing data sets. A series of

randomizations of the plots were performed to create a 80:20 split of individuals that optimized

the number of species in the train and test data sets. For each randomization, we calculated the

total number of species in the test set and repeated this random operation until we found the

split which maintained the highest number of species from the original data in both train and test

set. For the general model, the training data set contained 5867 individuals of 80 species and

the test data set 1599 individuals of 75 species. The resulting data represents 56% of the total

tree species in the original unfiltered vegetation structure dataset and these species account for

an average of 89% of individuals per site (Figure S.3).

Predictions for the species class of each individual in the test set were made using a 4x4

clip centered on the location of the test stem. Model performance was then evaluated using

overall accuracy, individual tree level (micro) and average species-level (macro) F1 scores

(hereafter referred to as individual-level and species-level accuracy respectively). The F1 score

combines precision and recall to provide a general measure of the overall accuracy of the

species classification, allowing for direct comparison between models using a single metric

(Chinchor, 1992). For each site, F1 scores for the general model were compared to those

produced by equivalent single site models to determine how the general model performed

relative to the traditional single site approach.

To understand the performance of the general model in different ecological contexts, we

evaluated how performance varied across the United States, how performance correlated with

the number of species being predicted at the NEON site, and which components of the model

(site effects, elevation, geographic location, and hyperspectral reflectance) were most important

for prediction. We identified the relative importance of site, geolocation, elevation and

reflectance by performing bootstrap features importance (Breiman, 2001). This approach is

based on evaluating how the overall accuracy is affected by each individual feature. At every

iteration, one feature is selected and the values are randomly shuffled among the samples,

effectively removing the information held in it. The accuracy is recorded with the shuffled feature

to determine the loss of performance compared to the unshuffled data. We used the same

approach to quantify the relative importance of the 15 spectral regions in which we grouped the

hyperspectral data.

We also evaluated the characteristics of trees and forests associated with the most

confusion between species (i.e., misclassification) based on forest type (using the National Land

Cover Database; Homer et al., 2001) and information from the NEON field data on canopy

position, tree status, and growth form from the NEON field data. We also assessed spatial

structure in confusion by determining, for every misclassified tree, whether the species it was

incorrectly classified as also occurred in the same NEON field plot. Finally, since confusion

commonly occurred within genera we also evaluated model performance for predicting genus

instead of species.

2.6. Prediction

We generated predictions for individual trees at the landscape scale (~350 km^2) by

integrating our approach with individual tree detections from previous work (Weinstein et al.,

2021). The Weinstein et al. (2021) dataset consists of 100 million individual tree crowns from 37

NEON sites identified using a retinanet neural network object detector and represented by

quadrangular polygons (i.e. bounding boxes) roughly representing the surface of the sunlit

portion of the crown. For consistency between the approach used for training and testing the

model (16-pixel clips), we extracted the pixels from the centroid of each estimated bounding

box. First, we extracted a 4x4 square window of pixels around the centroid of each detection.

For bounding boxes smaller than 16m^2, we dropped the pixels falling outside the bounding

boxes. Second, we filtered vegetation pixels from the background using the same procedure as

applied to the training/test data set. We finally selected all pixels with uncertainty scores > 0.5 to

be used to make predictions at the level of individual trees. We assigned each tree to a species

class by averaging the probability vectors (i.e., probability that the pixel is assigned to any of the

80 classes) of each pixel in the crown and selecting the species with the highest average

probability. We assigned each individual-tree prediction an uncertainty score consisting of the

average pixel probability, which ranged from 0-1.

3. Results

The general (cross-site) model yielded more accurate species classifications (larger F1

scores) than site-level models for 17 (species-level F1) and 18 (individual-level F1) of the 30

sites and identical accuracies for 8 (species-level F1) and 10 (individual-level F1) additional

sites. There were only two sites that showed better site-level species-level and individual-level

F1 scores: Blandy Experimental Farm, Washington (BLAN) and Rocky Mountain National Park,

Colorado (RMNP) (Figure 1, Figure S. 4, Figure S. 5). On average, the general model resulted

in increases in the accuracy of individual tree level classification (increases in individual-level

F1) from 0.72 to 0.77 and species-average accuracy (increases in species-level F1) from 0.46

to 0.54. Accuracy of the ensemble was higher than its sub-models trained singularly (raw

outputs available at https://doi.org/10.5281/zenodo.5796142), which is consistent with the

general observation that ensemble-based approaches produce more accurate predictions

(Healey et al., 2018). Since the general ensemble model proved to be the best performing

approach in this study, we focus primarily on it from this point forward.

Our results show a link between classification accuracy and ecological properties such

as ecosystem type, tree health, and growth form (Figure 2, Figure S.6). Damaged trees,

including broken boles and other types of damage (but not diseased trees), exhibited higher

rates of misclassification than healthy crowns (Figure 2), with broken boles exhibiting a >50%

misclassification rate. The general model performed best in shrub scrub and evergreen forests

and worst in wetlands (~32% misclassification rate), with deciduous forests falling in between.

Average classification accuracy was higher in eastern forests compared to western forests

(Figure 3a), and was significantly correlated with the number of species within the site (Figure

3b,d). The algorithm generally underperformed in the Prairie Peninsula and Central and

Southern Plains ecoregions which are characterized by patches of closed forest at the edges of

prairies or farmland (Figure 3a, Figure S.6). These results align with previous work in showing

that classification from remote sensing is more challenging for more complex canopies,

overlapping crowns, and coexisting species with similar life history and spectral properties

(Heinzel & Koch 2016, Bioucas-Dias, 2013).

Roughly 80% of the information used by the algorithm for classifying species was from

the hyperspectral reflectance (Figure 4). Important information was present across the entire

spectrum, but our results show that each spectral region is characterized by groups of more

informative bands. Specifically, the most important band groupings are the blue and green

(0.450 to 0.550 nm) in the visible region, the red-edge in the near infrared (0.62 to 0.85), 1.15 to

1.27 nm in SWIR1 and 1.62 to 1.68 nm in SWIR2. The site's coordinates, which represented the

geographic location of sites, explained 11% of total variation and were the second most

important variable (Figure 4). Elevation, a proxy of potential local changes in the environment

within each site, accounted for another 4%. The site effect, a proxy of other site level ancillary

information (e.g. sensor calibration, flight and atmospheric conditions), only accounted for 3% of

the total explained variance.

Comparing misclassification among species shows there is greater confusion for rare

species, congenerics, and species that co-occur within NEON field plots, and that model-

estimated uncertainty accurately reflects confidence in the model prediction. All species

performing poorly (F1 < 0.5) belonged to taxa with low sample sizes (less than 50 trees for

training) (Figure S6, Figure S7). In general, most of the confusion was among species co-

occurring within plot (74%) and site (93%). A large amount of confusion also occurred among

congeneric species (~27% of total misclassifications), mostly within pines, poplars, oaks and

maples, which make up 57% of the test dataset (Figure S.9). Oaks, pines and poplars in

particular accounted for ~87% of the total within genus confusion, and most misclassifications

had confidence scores >0.8. Aggregating predictions at the genus level improved the overall

accuracy by 6% (individual-level F1 accuracy of 83%), confirming that part of the confusion is

embedded in physiological similarities across taxonomically related trees. The model produced

reliable estimates of uncertainty for all species regardless of the accuracy. Uncertainty scores

matched closely with the probability of correct classification (R^2 =of 0.89, Figure 5).

4. Discussion

Using a single general model that integrated data across a continental scale resulted in

more accurate classification of tree species identity from remote sensing data than building

separate models for individual forest sites. The more accurate classification occurred despite

the continental data set containing many different forest types, structures and species

compositions across 30 sites. This suggests that the benefits of increasing the number of

samples for less common species and more fully characterizing within-species variance

outweighs the costs associated with including species that do not overlap geographically and

including components of within-species variance not observed at individual sites (Figure 1). To

our knowledge this is the first study to attempt to develop a generalized model across dozens of

sites and the success of the general model here suggests that developing generalized

algorithms offers a potential step forward in species classification from remote sensing more

broadly. Our generalized approach leveraged the information from multiple locations and survey

efforts, increasing the number of individuals from rarely sampled or highly variable classes and

allowing models to learn more broadly about how to distinguish species in the taxonomic group

of interest. In addition to yielding improved predictions, generalized cross-site approaches can

potentially generate predictions for a wide range of ecosystems, including those with limited or

no training data, allowing other studies to leverage the same shared model and thereby

facilitating large-scale ecological research (Weinstein et al. 2021).

By virtue of classification of the most common tree species occurring in the canopy, the

results of this model are potentially useful for several ecological applications, such as mapping

biomass and modeling carbon, energy and water flux. Our model captured species making up

~80% of the individual trees in the upper canopy, although this varied among sites.

Furthermore, given the stratified sampling of the NEON vegetation structure data used to

develop the generalized model, the model is likely to capture the major vegetation types and

most common species at each site. Canopy trees, which are visible from optical remote sensing

devices, contribute the majority of biomass in forests (Lutz et al., 2012). Because they form the

interface between the atmosphere and land surface, the canopy layer also is particularly

important for water and energy flux (Paul-Limogens et al., 2017). Because carbon storage,

water and energy flux can vary among species (Wright et al., 2006), the ability to map the

location and coverage of canopy species is important for assessing these important ecosystem

characteristics. Other ecological applications, such as assessing total forest species richness,

and quantifying tree regeneration, cannot be addressed using the model because our model

could not classify rare canopy species or understory individuals,

One of the main challenges in developing models that generalize well across the

continent is overcoming differences across sites in factors including seasonality, background,

and sensor calibration (Hesketh & Sanchez-Azofeifa, 2012, Clark et al., 2005, Pu, 2021). To

quantify the sensitivity of the algorithm to this ancillary information, we evaluated the relative

importance of the site-effect features compared to reflectance, geography and elevation. Our

results showed that the relative importance of the site-effect is marginal and accounts for less

than 3% of the total information captured (Figure 4). This suggests that the spectral signal from

NEON data is comparable across different flights and that flight-specific noise can be minimized

using BRDF corrections and vector normalization to limit the impact on the accuracy of

generalized algorithms. This is due in part to NEON data being highly standardized and using

the same image pre-processing protocol across the entire network (Kampe et al., 2014). NEON

remote sensing data is also collected at the peak of vegetation productivity for each site,

reducing the confounding effect of different phenological stages for species occurring at multiple

sites (Gartner et al., 2016). Expanding large scale surveys outside the NEON network would

require integrating information from less standardized sources, raising new challenges related to

fusion of sensors that are not cross-calibrated and images collected in different seasons (Brook

& Ben-Dor, 2015, Zou et al., 2018). Further investigation is therefore fundamental to evaluating

whether our findings apply to applications that involve integrating multiple sensors, missions, or

resolutions.

One of the advantages to broad scale general models is that they allow assessment of

how different ecological and environmental conditions influence the quality of the species

classification. Understanding variation in model performance across space, forest types, and

taxa is fundamental to better understanding where and when these models can be applied and

improvement of large-scale surveys from remote sensing. In our analysis, eastern forests

showed lower accuracy compared to western ecosystems. We believe this is at least partly

because eastern ecosystems are characterized by the higher species diversity of canopy trees

as well as crown geometry that makes aligning stems to crowns more difficult compared to

western conifer stands (Figure 2, 3, S.6). Higher species diversity in eastern forests (mean

species per site ~15) compared to western forests (mean species diversity per site ~4),

inherently makes classification tasks more challenging due to larger numbers of classes

typically resulting in lower accuracy predictions (Takahashi et al., 2020). Continuous closed

canopies also increase the likelihood pixels selected in a window centered on the stem will be

from neighboring tree crowns. This is due to the difficulty of obtaining accurate GPS points of

stems in closed canopy (Rodriguez-Perez et al., 2007), as well as the increased likelihood of

sunlit portions of the crown being displaced from the stem location in continuous broadleaf

forests (Strigul et al., 2008). This is a common problem, since field surveys often provide only

the geographic coordinates of tree stems and lack information about crown position or size,

making it impossible to correctly align crown borders with species labels. For example, pixel

mislabeling may be one of the reasons why our classifier was weaker at sites in the Great

Plains region (e.g., the NEON sites of Lyndon B. Johnson National Grasslands, CLBJ and

University of Kansas Field Station, UKFS), where patches of grasslands alternate with dense

forests characterized by multiple oak species forming a complex mosaic of crowns that may

have shifted their stem locations. In contrast, conifers in western US forests tend to be

dominated by species characterized by apical dominance (e.g. aspens and firs) with crowns

centered directly above the main stem reducing pixel mislabeling and improving classification.

Finally, savannas, such as the San Joaquin Experimental Range (SJER), characterized by

isolated trees of few species (mostly broadleaves) may be less likely to suffer from confounding

effects like crown displacement and stem-crown misalignment, making them less prone to

spectral mixing or potential pixel-mislabeling (Heinzel & Koch, 2012). The most challenging

ecosystem type in our analysis, wetlands, combines all of these challenges. Species like

Carpinus caroliniana and *Betula papyrifera*, found often in plots from wetland ecosystems, were

among the species with the worst classification accuracy, partly because they are generally

smaller trees that can occur in the understory, grow in closed canopies in the overstory (e.g. an

average dbh ~16.5cm and average height of ~10m), and often include limited training samples

because they are found in riparian ecosystems which make up a small fraction of the

landscapes from the NEON sites included in this study (less than 50 individuals per species).

Because of these challenges, the accuracy of the species predictions need to be assessed

depending on the site and ecosystem types within sites to ensure it is sufficient for the intended

ecological application.

Because species predictions from remote sensing are imperfect, it is important that

classification models produce robust estimates of uncertainty to allow this uncertainty to be

propagated through ecological analyses and considered during decision making. This is

particularly important when generating large numbers of predictions at large scales, because

this will result in including species located in undersampled areas and challenging ecosystems

as well as species that are difficult to classify due to rarity or similarity to other closely related

species. Our results confirmed that stacking scores from different classifiers using a logistic

regression produces accurate estimation of classification uncertainty (Figure 6).

While our generalized approach resulted in significant improvements over site-level

models, it is important to recognize that the accuracy of this approach is still insufficient for

ecological analyses contingent on rare or untrained species. For example, biodiversity patterns

are often driven by rare species (Leitao et al., 2016, Mouilllot et al., 2013), which are the most

challenging taxa for species classification, especially species so rare that they cannot be

included in the model due to data limitations (n < 5 individual trees in this study). Extrapolating

outside of NEON sites, a goal for general models, would also result in the presence of additional

species missing from the field dataset, restricting the range of ecological analyses to species

sampled within the footprint of NEON sites. Some of these limitations may be mitigated by

classifying trees into higher level taxonomic levels. In this study we observed that misclassified

trees were generally limited to species in the same genus and species co-occurring in the same

plot (Figure S.9, Figure S.10). Oaks, pines and poplars in particular accounted for ~87% of the

total within genus confusion. One possible driver of confusion among oak species is their similar

physiological and spectral characteristics (Figure S.11). Some co-occurring oaks species like

Quercus alba and *Quercus stellata* can also cross-breed and therefore be physiologically very

similar (Hardin, 1975), making them particularly hard to distinguish from imagery. For these

reasons, most of the cases leading to misclassification resulted from within-genus confusion.

This implies that uncertainty can be significantly reduced by aggregating predictions to the

genus level, offering a more robust solution for large scale ecological applications that can be

successfully addressed by accurately classifying trees at the level of genus, families or plant

functional type. For example, earth system models use plant functional types as the taxonomic

unit for quantifying carbon dynamics at continental to global scale (Lawrence et al., 2019), large

scale fire risk assessment and management can be achieved by using genus level surveys of

the most dominant taxa (Ma et al., 2021), and patterns of forest biomass largely depending on

which taxa dominate the ecosystem (Cheng et al., 2018). An increase in taxonomic level also

reduces issues with applying general models beyond the training data (e.g., outside of NEON

sites) because it is much more likely that all genera or families have been sampled in the

training data.

Building generalized algorithms provides an approach to overcome the significant field

data limitations present in most remote sensing tasks in ecology, by allowing pooling data from

ever growing sources of spatially explicit field surveys and high-resolution remote sensing

imagery. Our results showed that by integrating field surveys from dozens of NEON sites, it is

possible to produce a general model that provides improvements over single-site models for

species classification, with good estimates of uncertainty, and the ability to increase accuracy

further by aggregating predictions at the genus level. This general approach also unlocks the

potential for making predictions outside of NEON sites. The ultimate goal is to develop general

models that can be used anywhere in the region of interest (in our case the United States).

Using only NEON data, we successfully built a single integrated classifier that includes 20% of

all tree species found in forest ecosystems across the US (n=80 out of 396 surveyed by the

United States Forest Inventory and Analysis project; appendix F; Woudenberg, et al., 2010).

Beyond NEON, more and more openly available field and hyperspectral datasets are being

released from aerial (airborne and UAV) and satellite hyperspectral missions worldwide (Cook

et al., 2013, Vangi et al., 2021, Claverie et al., 2018). Our results show that instead of training

hundreds of separate models for local applications, there is the potential for integrating field and

remote sensing collections from multiple locations and sources to build general models with

improved accuracy for a broader range of landscapes and geographic locations. Further

integration with more field and remote sensing datasets could potentially provide remote

sensing based surveys of hundreds of millions of trees, making it possible to investigate the

properties of ecosystems from local to continental scales.

5. Conclusions

Remote sensing is facing a revolution in data quality and methods accuracy, making it a good candidate for developing applications to survey species assembly and forest properties at scale. Leveraging data collected from NEON across the US, we demonstrated that building continental scale algorithms for generalized species classification offers several advantages over the more traditional site level applications. Despite being very high for dominant taxa, accuracy in predictions for less represented species can be taunted by limitations in field-to-image misalignment, the number of species and individuals from rarely sampled taxa, making surveys from remote sensing unsuited to date for analyzing patterns in species alpha diversity at scale. Yet, building generalized algorithms is a fundamental cornerstone to overcome these limitations, because it allows for pooling from ever growing sources of geo-explicit field surveys and high resolution remote sensing imagery. Our results showed that by integrating field surveys with NEON airborne data, it is possible already to generate highly accurate predictions at the genus level and overall good estimates of uncertainty for individual trees. This allows for generating surveys of hundreds of millions of individual crowns across the continent, unlocking the potential for investigating large scale ecological applications focusing on the sun-exposed part of the canopy, dominant species, genuses or functional types.

Acknowledgements

This work was supported by the Gordon and Betty Moore Foundation's Data-Driven Discovery Initiative through grant GBMF4563 to E. P. White and by the National Science Foundation through grant 1926542 to E. P. White, S. A. Bohlman, A. Zare, D. Z. Wang, and A. Singh; by the NSF Dimension of Biodiversity program grant (DEB-1442280) and USDA/NIFA McIntire-Stennis program (FLA-FOR-005470) to S. A. Bohlman; by the University of Florida

Biodiversity Institute (UFBI) and Informatics Institute (UFII) Graduate Fellowship to Sergio

Marconi. There was no additional external funding received for this study.

All data can be found in the following Zenodo archive:

https://doi.org/10.5281/zenodo.5796143. All code for data preprocessing, model training and

testing and analyses can be found in the following GitHub repo:

https://github.com/MarconiS/Continental-scale-Hyperspectral-tree-species-classification-in-the-

National-Ecological-Observatory-N

References

1. Ab Majid, Ibtisam, Zulkiflee Abd Latif, and Nor Aizam Adnan. "Tree species classification

using worldview-3 data." 2016 7th IEEE Control and System Graduate Research

Colloquium (ICSGRC). IEEE, 2016.

2. Alina Zare, Susan Meerdink, Yutai Zhou, Caleb Robey, Ron Fick, John Henning, & Paul

Gader. (2019, April 12). GatorSense/hsi_toolkit_py: Initial Release (Version v1.0).

Zenodo. http://doi.org/10.5281/zenodo.2638117

3. Anderson-Teixeira, K.J., Davies, S.J., Bennett, A.C., Gonzalez-Akre, E.B., Muller-

Landau, H.C., Joseph Wright, S., Abu Salim, K., Almeyda Zambrano, A.M., Alonso, A.,

Baltzer, J.L. and Basset, Y., 2015. CTFS-ForestGEO: a worldwide network monitoring

forests in an era of global change. Global change biology, 21(2), pp.528-549.

4. Anderson, C.B. "Biodiversity monitoring, earth observations and the ecology of scale."

Ecology letters 21.10 (2018): 1572-1585.

5. Ayrey, E., et al. "Synthesizing Disparate LiDAR and Satellite Datasets through Deep

Learning to Generate Wall-to-Wall Regional Forest Inventories." BioRxiv (2019): 580514.

6. Ballanti, Laurel, et al. "Tree species classification using hyperspectral imagery: A

comparison of two classifiers." Remote Sensing 8.6 (2016): 445.

7. Bastin, J.F., et al. "The global tree restoration potential." Science 365.6448 (2019): 76-79.

8. Belgiu, Mariana, and Lucian Drăguţ. "Random forest in remote sensing: A review of applications and future directions." ISPRS journal of photogrammetry and remote sensing 114 (2016): 24-31.

9. Bioucas-Dias, José M., et al. "Hyperspectral remote sensing data analysis and future challenges." IEEE Geoscience and remote sensing magazine 1.2 (2013): 6-36.

10. Breiman, Leo. "Random forests." Machine learning 45.1 (2001): 5-32.

11. Brook, Anna, and Eyal Ben-Dor. "Supervised vicarious calibration (SVC) of multi-source hyperspectral remote-sensing data." Remote Sensing 7.5 (2015): 6196-6223.

12. Chawla, Nitesh V., et al. "SMOTE: synthetic minority over-sampling technique." Journal of artificial intelligence research 16 (2002): 321-357.

13. Cheng, Yanxia, et al. "Biomass-dominant species shape the productivity-diversity relationship in two temperate forests." Annals of Forest Science 75.4 (2018): 1-9.

14. Chinchor, N. "MUC-4 evaluation metrics in Proc. of the Fourth Message Understanding Conference 22–29." (1992).

15. Claverie, Martin, et al. "The Harmonized Landsat and Sentinel-2 surface reflectance data set." Remote Sensing of Environment 219 (2018): 145-161.

16. Cook, Bruce D., et al. "NASA Goddard's LiDAR, hyperspectral and thermal (G-LiHT) airborne imager." Remote Sensing 5.8 (2013): 4045-4066

17. Delicado, Pedro. "Dimensionality reduction when data are density functions." Computational Statistics & Data Analysis 55.1 (2011): 401-420.

18. Dietze, Michael C. "Prediction in ecology: A first-principles framework." Ecological Applications 27.7 (2017): 2048-2060.

19. Fassnacht, Fabian Ewald, et al. "Review of studies on tree species classification from remotely sensed data." Remote Sensing of Environment 186 (2016): 64-87.

20. Fricker, G. A., Ventura, J. D., Wolf, J. A., North, M. P., Davis, F. W., & Franklin, J.

(2019). A convolutional neural network classifier identifies tree species in mixed-conifer

forest from hyperspectral imagery. Remote Sensing, 11(19), 2326.

21. G. Batista, B. Bazzan, M. Monard, "Balancing Training Data for Automated Annotation of

Keywords: a Case Study," In WOB, 10-18, 2003.

22. Gärtner, Philipp, Michael Förster, and Birgit Kleinschmit. "The benefit of synthetically

generated RapidEye and Landsat 8 data fusion time series for riparian forest

disturbance monitoring." Remote Sensing of Environment 177 (2016): 237-247.

23. Guo, Chuan, et al. "On calibration of modern neural networks." International Conference

on Machine Learning. PMLR, 2017.

24. Guryanov, Aleksei. "Histogram-based algorithm for building gradient boosting ensembles

of piecewise linear decision trees." International Conference on Analysis of Images,

Social Networks and Texts. Springer, Cham, 2019.

25. Hardin, James W. "Hybridization and introgression in Quercus alba." Journal of the

Arnold Arboretum 56.3 (1975): 336-363.

26. Healey, Sean P., et al. "Mapping forest change using stacked generalization: An

ensemble approach." Remote Sensing of Environment 204 (2018): 717-728.

27. Heinzel, Johannes, and Barbara Koch. "Investigating multiple data sources for tree

species classification in temperate forest and use for single tree delineation."

International Journal of Applied Earth Observation and Geoinformation 18 (2012): 101-

110.

28. Henrys, Peter A., and Susan G. Jarvis. "Integration of ground survey and remote

sensing derived data: Producing robust indicators of habitat extent and condition."

Ecology and evolution 9.14 (2019): 8104-8112.

29. Hesketh, Michael, and G. Arturo Sánchez-Azofeifa. "The effect of seasonal spectral

variation on species classification in the Panamanian tropical forest." Remote Sensing of

Environment118 (2012): 73-82.

30. Homer, Collin, et al. "Development of a 2001 national land-cover database for the United

States." Photogrammetric Engineering & Remote Sensing 70.7 (2004): 829-840.

31. Kampe, T., et al. "NEON imaging spectrometer geolocation processing algorithm

theoretical basis document." NEON Doc.# 001290 Rev A (2014).

32. Kandare, K., et al. "Prediction of species-specific volume using different inventory

approaches by fusing airborne laser scanning and hyperspectral data." Remote Sensing

9.5 (2017): 400.

33. Knauer, Uwe, et al. "Tree species classification based on hybrid ensembles of a

convolutional neural network (CNN) and random forest classifiers." Remote Sensing

11.23 (2019): 2788.

34. Krause, Keith S., et al. "Early algorithm development efforts for the National Ecological

Observatory Network Airborne Observation Platform imaging spectrometer and

waveform lidar instruments." Imaging Spectrometry XVI. Vol. 8158. International Society

for Optics and Photonics, 2011.

35. Laaksonen, Jorma, and Erkki Oja. "Classification with learning k-nearest neighbors."

Proceedings of International Conference on Neural Networks (ICNN'96). Vol. 3. IEEE,

1996.

36. Lawrence, David M., et al. "The Community Land Model version 5: Description of new

features, benchmarking, and impact of forcing uncertainty." Journal of Advances in

Modeling Earth Systems 11.12 (2019): 4245-4287.

37. Lawrence, M., et al. "Comparisons of national forest inventories." National forest

inventories. Springer, Dordrecht, 2010. 19-32.

638. Leitão RP, Zuanon J, Villéger S, Williams SE, Baraloto C, Fortunel C, Mendonça FP, Mouillot D. 2016 Rare species contribute disproportionately to the functional structure of species assemblages. Proc. R. Soc. B 283: 20160084. http://dx.doi.org/10.1098/rspb.2016.0084

39. Li, Wei, et al. "Locality-preserving dimensionality reduction and classification for hyperspectral image analysis." IEEE Transactions on Geoscience and Remote Sensing 50.4 (2011): 1185-1198.

40. Loizzo, R., et al. "PRISMA: The Italian hyperspectral mission." IGARSS 2018-2018 IEEE International Geoscience and Remote Sensing Symposium. IEEE, 2018.

41. Lutz, James A., et al. "Ecological importance of large-diameter trees in a temperate mixed-conifer forest." PloS one 7.5 (2012): e36131.

42. M. L. Clark, D. A. Roberts, and D. B. Clark, "Hyperspectral discrimination of tropical rain forest tree species at leaf to crown scales," Remote Sensing of Environment, vol. 96, no. 3–4, pp. 375–398, 2005.

43. Ma, Wu, et al. "Assessing climate change impacts on live fuel moisture and wildfire risk using a hydrodynamic vegetation model." Biogeosciences (2021).

44. Maddala, Gangadharrao S. Limited-dependent and qualitative variables in econometrics. No. 3. Cambridge university press, 1986.

45. Marconi, Sergio, et al. "Rethinking the fundamental unit of ecological remote sensing: Estimating individual level plant traits at scale." bioRxiv (2019): 556472.

46. Mäyrä, Janne, et al. "Tree species classification from airborne hyperspectral and LiDAR data using 3D convolutional neural networks." Remote Sensing of Environment 256 (2021): 112322.

47. Mouillot D, Bellwood DR, Baraloto C, Chave J, Galzin R, et al. Rare Species Support Vulnerable Functions in High-Diversity Ecosystems. PLoS Biology, 2013 DOI: 10.1371/journal.pbio.1001569

48. Mukhoti, Jishnu, et al. "Calibrating deep neural networks using focal loss." arXiv preprint

arXiv:2002.09437 (2020).

49. NEON (National Ecological Observatory Network). Spectrometer orthorectified surface

directional reflectance - mosaic, RELEASE-2021 (DP3.30006.001).

https://doi.org/10.48443/qeae-3x15. Dataset accessed from https://data.neonscience.org

on March 7, 2021

50. Pacifico, Luciano DS, Valmir Macario, and Joao FL Oliveira. "Plant classification using

artificial neural networks." 2018 International Joint Conference on Neural Networks

(IJCNN). IEEE, 2018.

51. Paul-Limoges, Eugénie, et al. "Below-canopy contributions to ecosystem CO2 fluxes in a

temperate mixed forest in Switzerland." Agricultural and Forest Meteorology 247 (2017):

582-596.

52. Pax-Lenney, Mary, et al. "Forest mapping with a generalized classifier and Landsat TM

data." Remote Sensing of Environment 77.3 (2001): 241-250.

53. Pecl, G.T., Araújo, M.B., Bell, J.D., Blanchard, J., Bonebrake, T.C., Chen, I.C., Clark,

T.D., Colwell, R.K., Danielsen, F., Evengård, B. and Falconi, L., 2017. Biodiversity

redistribution under climate change: Impacts on ecosystems and human well-being.

Science, 355(6332), p.eaai9214.

54. Pedregosa, Fabian, et al. "Scikit-learn: Machine learning in Python." the Journal of

machine Learning research 12 (2011): 2825-2830.

55. Pu, Ruiliang. "Mapping Tree Species Using Advanced Remote Sensing Technologies: A

State-of-the-Art Review and Perspective." Journal of Remote Sensing 2021 (2021).

56. Rana, Parvez, et al. "Towards a generalized method for tree species classification using

multispectral airborne laser scanning in Ontario, Canada." IGARSS 2018-2018 IEEE

International Geoscience and Remote Sensing Symposium. IEEE, 2018.

57. Rodríguez-Pérez, José R., M. F. Álvarez, and Enoc Sanz-Ablanedo. "Assessment of

low-cost GPS receiver accuracy and precision in forest environments." Journal of

Surveying Engineering 133.4 (2007): 159-167.

58. Scholl, V. M., Cattau, M. E., Joseph, M. B., & Balch, J. K. (2020). Integrating national

ecological observatory network (neon) airborne remote sensing and in-situ data for

optimal tree species classification. Remote Sensing, 12(9), 1414.

59. Sims, Daniel A., and John A. Gamon. "Relationships between leaf pigment content and

spectral reflectance across a wide range of species, leaf structures and developmental

stages." Remote sensing of environment 81.2-3 (2002): 337-354.

60. Strahler, A. H., Thomas L. Logan, and Nevin A. Bryant. "Improving forest cover

classification accuracy from Landsat by incorporating topographic information." (1978).

61. Strahler, Alan H., Thomas L. Logan, and Nevin A. Bryant. "Improving forest cover

classification accuracy from Landsat by incorporating topographic information." (1978).

62. Strigul, Nikolay, et al. "Scaling from trees to forests: tractable macroscopic equations for

forest dynamics." Ecological Monographs 78.4 (2008): 523-545.

63. Takahashi Miyoshi, Gabriela, et al. "Evaluation of hyperspectral multitemporal

information to improve tree species identification in the highly diverse atlantic forest."

Remote Sensing 12.2 (2020): 244.

64. Tang, J., S. Alelyani, and H. Liu. "Data Classification: Algorithms and Applications." Data

Mining and Knowledge Discovery Series, CRC Press (2015): pp. 498-500.

65. Tavares, P. A., Beltrão, N. E. S., Guimarães, U. S., & Teodoro, A. C. (2019). Integration

of sentinel-1 and sentinel-2 for classification and LULC mapping in the urban area of

Belém, eastern Brazilian Amazon. Sensors, 19(5), 1140.

66. Tomek, Ivan. "Two modifications of CNN." (1976).

67. Tomppo, E., et al. "Combining national forest inventory field plots and remote sensing

data for forest databases." Remote Sensing of Environment 112.5 (2008): 1982-1999.

68. USDA Forest Service, 2001. Forest Inventory and Analysis National Core Field Guide,

Volume I: Field Data Collection Procedures For Phase 2 Plots, Version 1.5. US

Department of Agriculture, Forest Service, Washington, DC.

69. Vangi, Elia, et al. "The new hyperspectral satellite PRISMA: Imagery for forest types

discrimination." Sensors 21.4 (2021): 1182.

70. Weinstein, B.G., Gardner, L., Saccomanno, V., Steinkraus, A., Ortega, A., Brush, K.,

Yenni, G., McKellar, A.E., Converse, R., Lippitt, C. and Wegmann, A., 2021. A general

deep learning model for bird detection in high resolution airborne imagery. bioRxiv.

https://doi.org/10.1101/2021.08.05.455311

71. White, J. C., et al. "Remote sensing technologies for enhancing forest inventories: A

review." Canadian Journal of Remote Sensing 42.5 (2016): 619-641.

72. Wiens, J.J., 2016. Climate-related local extinctions are already widespread among plant

and animal species. PLoS biology, 14(12).

73. Woudenberg, S. W., et al. "The Forest Inventory and Analysis Database: Database

description and users manual version 4.0 for Phase 2." Gen. Tech. Rep. RMRS-GTR-

245. Fort Collins, CO: US Department of Agriculture, Forest Service, Rocky Mountain

Research Station. 336 p. 245 (2010).

74. Wright, Marvin N., and Inke R. König. "Splitting on categorical predictors in random

forests." PeerJ 7 (2019): e6339.

75. Wright, Ian J., et al. "Cross-species patterns in the coordination between leaf and stem

traits, and their implications for plant hydraulics." Physiologia Plantarum 127.3 (2006):

445-456.

76. Xi, Zhouxin, et al. "See the forest and the trees: Effective machine and deep learning

algorithms for wood filtering and tree species classification from terrestrial laser

scanning." ISPRS Journal of Photogrammetry and Remote Sensing 168 (2020): 1-16.

77. Zou, Hui, and Trevor Hastie. "Regularization and variable selection via the elastic net."

Journal of the royal statistical society: series B (statistical methodology) 67.2 (2005):

301-320.

Figure 1. Performance of generalized vs site-specific classification models for each NEON site.

Positive values are sites for which the generalized model performed better than site-level.

Negative values are sites for which the generalized model performed worse compared to site-

level. Blue bars represent species-level F1 score, yellow bars individual-level F1. Numbers

separated by (:) on top of each site name represent the total number of species in the training

and test set specific for each site. Numbers separated by (/) in BLAN represent the different

number of species in the test set for the site specific and general models.

764   Figure 2. Fraction of misclassified trees across ecosystem types (blue), growth form (yellow),

765   canopy position (red) and health status (grey). Numbers above the x-axis labels are the number

766   of trees in each category.

Figure 3. Variation in accuracy of the generalized algorithm across the US. (a) map of average

individual-level F1 scores for each ecological domain. Dots represent the location of each

NEON site. Blue polygons represent the Prairie Peninsula and Central-Southern Plains. (b)

Relationship between individual-level F1 and number of species in the training dataset for each

site.The blue line is the loess smoother relationship over the 29 sites. (c) Kernel density

estimate of the distribution of individual-level F1 scores (averages per site). (d) Relationship

between species-level F1 and number of trained species found in site; orange line is the loess

smoother relationship over the 29 sites. (e) Kernel density estimate of the distribution of

species-level F1 scores (averages per site).

Figure 4: Features importance calculated from the permutation feature importance procedures

described in Breiman, 2001, on the meta-ensemble model. (a) Relative contribution of different

feature types.reflectance (grey), site coordinates (yellow), elevation (blue) and site effect

(red).(b) Relative importance of each Kullback-Leibler group used for dimensionality reduction of

reflectance. Blue bars represent the reflectance for the average minimum, mean and maximum

band in the specific KL group. Bar width represents the range of bands covered by the specific

KL group. Color intensity represents the relative importance of the specific KL group for the

classifier (from light blue being of little importance, to dark blue being highly important). Black

lines represent the reflectance of a randomly selected pixel to illustrate a typical vegetation

reflectance pattern. Reflectance was normalized using L2 normalization.

Figure 5. Evaluation of model confidence score (the probability of assigning the correct label to

a prediction) as a measure of uncertainty. Confidence score was binned into 34 equal-width

bins (each bin representing an interval of 0.03). Bin centers were plotted against the fraction of

trees in that confidence score bin that were correctly classified. The blue line shows the fitted

linear relationship between the confidence score and the proportion of correctly classified trees.

The black line is the 1:1 line.

Figure 6. Example of species classification maps for all individual trees at the Bartlett

Experimental Forest (BART) NEON site in New Hampshire. Species in legend include six of the

most abundant taxa predicted at the site. Individual crown boundaries were estimated using

predictions from Weinstein et al., 2021. The background is gray scale imagery of the site, so the

gray areas on the right hand panels are areas without any trees including roads and other open

areas.

Supplement 1: parameterization of classifiers and meta ensemble

KNN classifier was trained using 20 neighboring points, with distance weighted by the

inverse of their Manhattan distance. The Random Forest classifier was trained using 300 trees

with up to 7 features (square root of the total predictors) considered for better split, validated

using cross-entropy loss function on out-of-bag samples. The gradient boosting classifier was

trained using 1000 maximum iterations, a learning rate of 0.01, max depth of 25 and 0.5 L2

regularization. Loss was calculated using categorical cross-entropy on out-of-bag samples.

Multi-layer Perceptron classifier was trained for 1200 max iterations, using relu

activation, 1 hidden layer, weight optimization through adam booster with exponential

decay rate of 0.9. The Bagging Classifier was trained using 10 support vector machine

classifiers as base estimators. We used loose regularization (C = 1000), RBF kernel, and

5-fold cross validation to calibrate probability estimates. The meta ensemble was trained

using probability vectors produced by each weak classifier. We used a regularized logistic

regression (elasticnet), with 0.5 L1 to L2 penalty ratio. We used a saga solver to optimize the

loss function.

Supplementary Figures

Figure S1 Geographic distribution of NEON sites included for this study. Colored regions

represent ecological regions defined by NEON (https://www.neonscience.org/field-sites/about-

field-sites). A description of each site and their ecological domain can be found in the

Supplementary Table 1.

Figure S. 2: Flowchart of the species classification pipeline developed for this study

Figure S.3 For each site, fraction of all species (red) and their relative frequency in number of

trees (blue) covered by the species included in the analysis, compared to the total number of

species (red) and individual trees (blue) included in the NEON vegetation structure dataset.

Numbers separated by (/) above each site name represent the total number of species in the

original dataset and in the filtered data respectively, specific for each site. Trees in the canopy

(blue bars) were determined by canopy position data in the vegetation structure data where

trees in the canopy were designated as "Full sun", "Mostly shaded", "Partially shaded", "Open

growth", or non classified ("NA")..

Figure S.4 Density functions of the difference in ΔF1 scores between the generalized and each

single-site algorithm for species-level F1 (yellow) and individual-level F1 (blue). Positive ΔF1

values (17 out of 29 sites) represent sites where the generalized algorithm outperformed its site-

specific counterpart. Dashed vertical lines represent the average ΔF1 across sites (species-

level F1 = 0.09, individual-level F1 = 0.05).

Supplement S.5. Difference in accuracy between the general and site-specific approaches for

each species-site combination. Negative values (red) represent taxa whose accuracy is higher

in the general approach. Blue values represent taxa whose accuracy is higher in the site-

specific approach. White values where accuracy was similar for the general and site-specific

approaches. Grey are species that do not occur at the site. Sites are sorted by geographic

similarity.

Figure S.6 Example of 400m2 plots for 6 sites from western (left panel), central (center panel)

and eastern US (right panel). Dots represent field stem data collected from NEON vegetation

structure. Different dot colors represent different species. Only stems that have been filtered to

include only stems that are likely to be in the canopy. From top left to bottom right sites

acronyms are Abby Road (ABBY), Delta Junction (DEJU), Harvard Forest (HARV), San Joaquin

Experimental Range (SJER), Blandy Experimental Farm (BLAN), Lenoir Landing (LENO),

Figure S.7. Precision, Recall and F1for each individual species included in the dataset. Results

can be found in the supplement table in a separate supplementary file.

892 Figure S.8. The relationship between individual species F1 scores and number of individual

893 trees available for training for that species. The blue line shows a fitted relationship using local

894 polynomial regression fitting (loess) and the grey region shows the 95% confidence interval

895 around that relationship.

904 Figure S.9. Confidence score P(x) for those taxa each species was most confused with. Taxa

905 with a P(x) lower than 0.02 were not included.

Figure S.10. Distribution of species across sites.

Figure S.11 Example of spectral signature overlap between often confused congeneric species.

Lines represent the average spectra, shaded areas represent the standard deviation for all

pixels extracted for that particular species. The first species in legend is in blue, the second in

grey. The X-axis is the band number from brdf corrected hyperspectral image. Couples of

species are: (a) *Quercus marilandica* and *Quercus stellata*, (b) *Quercus alba* and *Quercus*

rubra, (c) *Pinus palustris* and *Pinus elliottii*, (d) *Abies magnifica* and *Abies lowiana*.

Site Name	Ecological Domain	Domain Name	State	Geolocation (Lat-Lon)
Abby Road NEON (ABBY)	D16	Pacific Northwest	Washington	45.76
				-122.33
Bartlett Experimental Forest NEON (BART)	D01	Northeast	New Hampshire	44.06
				-71.29
Blandy Experimental Farm NEON (BLAN)	D02	Mid Atlantic	Virginia	39.03
				-78.04
Caribou-Poker Creeks Research Watershed NEON (BONA)	D19	Taiga	Alaska	65.15
				-147.5
Dead Lake NEON (DELA)	D08	Ozarks Complex	Alabama	32.54
				-87.8
Delta Junction NEON (DEJU)	D19	Taiga	Alaska	63.88
				-145.75
Disney Wilderness Preserve NEON (DSNY)	D03	Southeast	Florida	28.13
				-81.44
Guanica Forest NEON (GUAN)	D04	Atlantic Neotropical	Puerto Rico	17.97
				-66.87
Harvard Forest & Quabbin	D01		Massachusetts	42.54

Watershed NEON (HARV)		Northeast		-72.17
KU Field Station NEON (UKFS)	D06	Prairie Peninsula	Kansas	39.04
				-95.19
Konza Prairie Biological Station NEON (KONZ)	D06	Prairie Peninsula	Kansas	39.1
				-96.56
Lenoir Landing NEON (LENO)	D08	Ozarks Complex	Alabama	31.85
				-88.16
Lyndon B. Johnson National Grassland NEON (CLBJ)	D11	Southern Plains	Texas	33.4
				-97.57
Moab NEON (MOAB)	D13	Southern Rockies / Colorado Plateau	Utah	38.25
				-109.39
Mountain Lake Biological Station NEON (MLBS)	D07	Appalachians / Cumberland Plateau	Virginia	37.38
				-80.52
Niwot Ridge NEON (NIWO)	D13	Southern Rockies / Colorado	Colorado	40.05
				-105.58

		Plateau		
Ordway-Swisher Biological Station NEON (OSBS)	D03	Southeast	Florida	29.69
				-81.99
Rocky Mountains NEON (RMNP)	D10	Central Plains	Colorado	40.28
				-105.55
San Joaquin Experimental Range NEON (SJER)	D17	Pacific Southwest	California	37.11
				-119.73
Smithsonian Conservation Biology Institute NEON (SCBI)	D02	Mid Atlantic	Virginia	38.89
				-78.14
Smithsonian Environmental Research Center NEON (SERC)	D02	Mid Atlantic	Maryland	38.89
				-76.56
Steigerwaldt-Chequamegon NEON (STEI)	D05	Great Lakes	Wisconsin	45.51
				-89.59
Talladega National Forest NEON (TALL)	D08	Ozarks Complex	Alabama	32.95
				-87.39
The Jones Center At Ichauway NEON (JERC)	D03	Southeast	Georgia	31.19
				-84.47
Treehaven NEON (TREE)	D05	Great Lakes	Wisconsin	45.49
				-89.59

				46.23
University of Notre Dame Environmental Research Center NEON (UNDE)	D05	Great Lakes	Michigan	-89.54
Wind River Experimental Forest NEON (WREF)	D16	Pacific Northwest	Washington	45.82
				-121.95
Yellowstone National Park NEON (YELL)	D12	Northern Rockies	Wyoming	44.95
				-110.54

923   Table S.1 Description of NEON sites and ecological domains used in this study.

924