Dead infant carrying by chimpanzee mothers in the Budongo Forest

Adrian Soldati[1,2,3], Pawel Fedurek[3,4,5], Catherine Crockford[4], Sam Adue[3], John Walter

Akankwasa[3], Caroline Asiimwe[3], Jackson Asua[3], Gideon Atayo[3], Boscou Chandia[3], Elodie

Freymann[3,6], Caroline Fryns[2,3], Geresomu Muhumuza[3], Derry Taylor[1,3], Klaus

Zuberbühler[1,2,3], Catherine Hobaiter[1,3]

Affiliations:

[1] School of Psychology and Neuroscience, University of St Andrews, St Andrews, UK

[2] Department of Comparative Cognition, Institute of Biology, University of Neuchâtel,

Neuchâtel, Switzerland

[3] Budongo Conservation Field Station, Masindi, Uganda

[4] Department of Primatology, Max Planck Institute for Evolutionary Anthropology, Leipzig,

Germany

[5] Division of Psychology, Faculty of Natural Sciences, University of Stirling, Stirling, UK

[6] Department of Anthropology and Museum Ethnography, University of Oxford, Oxford, UK

Email of corresponding author: as391@st-andrews.ac.uk

ORCID of Adrian Soldati: 0000-0002-3618-5022

ORCID of Catherine Hobaiter: 0000-0002-3893-0524

ORCID of Klaus Zuberbühler: 000-0001-8378-088X

ORCID of Elodie Freymann: 0000-0002-5484-8643

ORCID of Caroline Fryns: 0000-0001-7686-7274

ORCID of Derry Taylor: 0000-0002-8559-7154

Abstract

It has been suggested that non-human primates (hereafter primates) can respond to deceased

conspecifics in ways that suggest they experience psychological states not unlike humans,

which would indicate they exhibit some notion of death. Here, we report long-term

demographic data from two East African chimpanzee groups. During a combined 40-year

observation period we recorded 191 births of which around a third died in infancy, mostly

within the first year. We documented the post-mortem behaviour of the mothers and found

that Budongo chimpanzee mothers routinely carried deceased infants after their death,

usually until the body started to decompose after a few days. However, we also observed

three cases of extended carrying lasting for more than 2-weeks, one of which was followed

by the unusual extended carrying of a substitute object and another which lasted three

months. In each case, the corpses mummified. We discuss these data in view of functional

hypotheses of dead infant carrying in primates and regarding the potential proximate

mechanisms involved in this behaviour. We suggest that chimpanzees may exhibit

psychological processes related to death of conspecifics similar to human grieving.

Introduction

Primate thanatology, the study of the behaviour and underlying physiological and

psychological factors associated with dead or dying individuals in non-human primates

(hereafter primates), continues to raise important questions about human uniqueness

(Anderson 2016, 2018; Anderson et al. 2018). Humans experiencing the loss of socially close

individuals experience significant psychological trauma with long-term physiological effects,

including symptoms of post-traumatic stress disorder, anxiety and depression, and increased

used of health-care resources (Figley et al. 1997; Parkes et al. 1997; Lannen et al. 2008).

Given our biological and social similarities, other primate species – in particular other great

apes – may experience similar cognitive and physiological changes. Cross-species

comparisons, especially in primates, are often used to reveal past evolutionary trajectories of

the hominid lineage. Primate behaviour and physiology in relation to deceased individuals

provide valuable data to better understand the origins of why humans are so powerfully

affected by death. More generally, primate responses to death may provide further insight

into other aspects of animal cognition, such as animacy or the perception of time (Gonçalves

and Carvalho 2019).

Because death is unpredictable and rarely observed in wild primate communities, the

available datasets are usually anecdotal and descriptions of events are often incomplete

(Watson and Matsuzawa 2018; Ramsay and Teichroeb 2019). Nevertheless, an increasing

number of primate groups have been habituated to human observers (Kappeler et al. 2012),

which has led to more frequent reporting and more systematic efforts to extract patterns of

behaviour in responses to death (Anderson 2020). These observations have led to claims that

primates respond to death in ways that are similar to humans, by producing strong emotional,

social, and exploratory responses (Watson and Matsuzawa 2018; Gonçalves and Carvalho

2019). Among non-human animals, the emphasis on primates may result from easier

detection, given the relatively large number of long-term studies, and observations from

corvid, elephant, and cetacean species suggest that some understanding of death may be

widespread among a wide range of long-lived highly-social species (Gonçalves and Biro

2018; Bercovitch 2020).

Here, we focus on a particularly remarkable behaviour seen in many primates, dead infant

carrying by mothers, which in chimpanzees typically occurs for a period of up to three days

(Gonçalves and Carvalho 2019). Dead infant carrying (also referred as infant corpse carrying)

is the most frequently reported thanatological behaviour and shows substantial variation in

accompanying behaviours across primate species (Fernández-Fueyo et al. 2021), as a result, it

offers rich data against which to explore possible explanatory factors. This behaviour has not

only been reported in chimpanzees (Matsuzawa 1997; Hosaka et al. 2000; Kooriyama 2009;

Biro et al. 2010) but also in bonobos (Fowler and Hohmann 2010; Tokuyama et al. 2017),

gorillas (Warren and Williamson 2004; Masi 2020), chacma baboons (Carter et al. 2020), red

colobus (Georgiev et al. 2019), geladas (Fashing et al. 2011), ring-tailed lemurs (Nakamichi

et al. 1996), bonnet and lion-tailed macaques (Das et al. 2019), Japanese macaques

(Sugiyama et al. 2009; Takeshita et al. 2020), marmosets (Thompson et al. 2020), and vervet

monkeys (Botting and van de Waal 2020). Several hypotheses have been put forward to

explain the function of and motivation for this behaviour (Table 2; for a review see: Watson

and Matsuzawa 2018; Gonçalves and Carvalho 2019).

Within an increasingly large field of hypotheses – some of which are mutually non-exclusive

– there are some natural groupings based on similarity in the suggested mechanisms or

explanations. A first group of hypotheses presumes that primate mothers are unable to

understand the ramifications of death and their behavioural responses are simply side effects

of evolved physiological mechanisms. Specifically, the 'unawareness hypothesis' states that

mothers are unable to discriminate between temporarily unresponsive and irreversibly

deceased individuals and simply try to avoid the costs of premature abandonment (Hrdy

1999). The hypothesis predicts that mothers are unable to understand their infants' state and

continue providing maternal care (e.g. grooming) and try to elicit responsiveness (e.g.

poking, smelling). With increasing time, the ambiguity will dissolve due to the accelerating

decomposition of the corpse, which is usually mediated by local climate. A second prediction

therefore is that dry and particularly hot or cold conditions favouring mummification should

favour prolonged carrying of dead infants (see also 'climate hypothesis': Matsuzawa 1997;

Fashing et al. 2011). The 'post-parturient condition' hypothesis similarly proposes that the

maternal physiological conditions associated with pregnancy and birth favour persistent care

of the dead infant as long as the mother is lactating or until resumption of ovulation, for

example indicated by sexual swelling (Kaplan 1973). Here, the younger the infant the longer

the expected carrying period.

A second kind of hypotheses assumes that primate mothers can have a notion of death,

provided they have relevant personal experience. Specifically, the 'learning about death'

hypothesis states that chimpanzees do not intuitively understand death but can acquire the

notion by learning to attend to the relevant cues (Cronin et al. 2011). Here, the predication is

that experienced mothers will be aware of the irreversible change and, unlike first-time

mothers, should have shorter dead-infant carrying periods. An alternative hypothesis, the

'grief-management hypothesis', also assumes that chimpanzees possess a notion of death, and

suggests dead-infant carrying represents a consoling strategy to cope with grief. This

hypothesis predicts that mothers who are able to carry their dead infants experience lower

levels of 'stress' hormones (i.e. glucocorticoids) than mothers who could not or did not carry

them (Nicolson 1991). A recent study by Girard-Buttoz and colleagues (Girard-Buttoz et al.

2021) reported elevated cortisol levels in infant chimpanzees who lost their mothers,

supporting the notion that disruption of the mother-infant bond leads to elevated stress.

Similarly, female baboons experience high levels of glucocorticoids when losing an ally to

predation and in periods of infanticidal attacks (Engh et al. 2006a, b). Dead infant carrying,

therefore, could be a strategy to cope with stress associated with infant loss.

A third group of hypotheses are agnostic about whether chimpanzees possess a notion of

death but proposes various adaptive mechanisms that favour post-mortem mothering

behaviour. First, the 'learning-to-mother hypothesis' states that dead-infant carrying

improves maternal skills (Warren and Williamson 2004), predicting that the behaviour should

mainly be observed in inexperienced primiparous females. Second, the 'maternal-bond

strength hypothesis' predicts that mothers with older infants share a stronger bond and are

thus more likely to show extensive carrying as compared to mothers with younger infants

(Watson and Matsuzawa 2018). On the other hand, the 'hormonal hypothesis' states that,

after giving birth, the endocrine system of the mother releases hormones (e.g. oxytocin) that

stimulate maternal behaviours (Biro et al. 2010). Here, persistent carrying would be the result

of a strong mother-infant bond and the younger the infant at time of death the longer the

carrying period.

The existence of multiple – sometime contrasting – hypotheses is likely a reflection of the

small and highly variable data available and the diversity of potential drivers of this

behaviour across different species. A recent systematic study on primates (n=18 species)

showed that duration of infant carrying is affected by the age of the mother, with older

mothers carrying for longer periods (Das et al. 2019), in contrast with previous findings by

Sugiyama and colleagues who found no effect of age (Sugiyama et al., 2009). Infants that

died of sickness were carried for longer than those who were stillborn or victims of

infanticide, while the age of the infant did not influence duration of carrying (Das et al.

2019). Although dead infant carrying is a shared but rare behaviour in primates, carrying

duration seems to be the longest in great apes (Fernández-Fueyo et al. 2021), particularly in

chimpanzees (Das et al. 2019). Wild chimpanzees have higher levels of infant mortality

compared to captive conspecifics (Courtenay and Santow 1989; Hill et al. 2001), which

provides increased opportunities to observe mothers' responses to their infants' death

(Watson and Matsuzawa 2018). Chimpanzee mothers typically carry their dead infants for a

few days, though a recent analysis of the largest chimpanzee dataset (n=33 cases) did not

provide clear support for any of the previous hypotheses (Lonsdorf et al. 2020). Despite this,

it was proposed that the 'unawareness hypothesis' is unlikely because of the presence of

atypical carrying postures and sensory cues after death compared to those displayed towards

alive infants (Lonsdorf et al. 2020). More recently, a study using the largest primate database

to date (n=409 cases and n=50 species) found that dead infant carrying was more likely to

occur when the cause of death was non-traumatic and when mothers were younger

(Fernández-Fueyo et al. 2021). In addition, the authors showed that carrying duration was

greater for younger infants, supporting the importance of the maternal bond. Here, we revisit

the discussion on the potential reasons behind dead infant carrying by female chimpanzees

with a 40-year dataset of two study groups of the Budongo Forest, Uganda, including three

detailed observations of extended dead infant carrying by two different females.

Methods

Study site and subjects

The Budongo Forest Reserve is a semi-deciduous tropical rain forest located along the

western Rift Valley in Uganda. This reserve is made of 793 km^2 of protected forest and

grassland, including 482 km^2 of continuous forest cover (Eggeling 1947) . The reserve

contains a population of approximated 600 East African chimpanzees (*Pan troglodytes*

schweinfurthii). Our observations took place in two adjacent communities, Sonso and

Waibira, studied regularly by researchers and followed on a daily basis by field assistants

since 1990 and 2011 respectively (Reynolds 2005; Samuni et al. 2014).

At the time of the events, the Waibira community contained an estimated 120 individuals, 96

of which could be individually recognised. Individuals involved in the first event were Ketie

(KET), a 20-year old adult primiparous female (estimated birth 1998) and her 2-year old

female infant Karyo (KYO) born in December 2015. The Sonso community contained 65

named individuals, in addition to three unnamed females in the process of immigrating. The

individuals involved in the event were Upesi (UP), a 21-year old parous female (estimated

birth 1999) and a) in the second event, her recently born unnamed unsexed infant UP3, born

mid-September 2020 and b) in the third event, her 4th born unnamed unsexed infant UP4,

estimated birth 7th August 2021. Her first two infants (born in 2017 and 2018) were victims

of within-community infanticide before reaching a month old (see Leroux et al. 2021 for one

reported case).

In this paragraph we consider the scope for bias in our study subjects by using the

STRANGE framework to report potential sampling biases in our study (Webster and Rutz

2020; Rutz and Webster 2021). The Sonso community are of typical size whereas the

Waibira community are particularly large as compared to that of other chimpanzees (in a

recent comparison of 18 groups across three subspecies: *P.t. schweinfurthii, P.t. troglodytes,*

P.t. verus; communities range from 7-144 individuals with a mean 42; within these data the

East African sub-species (*P.t. schweinfurthii*) range is 18-144 with a median 49; Wilson et al.

2014). Sonso have a typical female-basied sex ratio among mature individuals (M:F; 1:1.7),

whereas the Waibira community have more unusual evenly-balanced sex ratio among mature

individuals (M:F; 1:1.2; mean among 9 *P.t. schweinfurthii* communities 1:1.7; Wilson et al.

2014). Of relevance to sampling biases in infant mortality and opportunities to carry dead

infants, the Sonso community experience high levels of infanticide (Lowe et al. 2019, 2020).

A medium altitude rainforest (~1100m) with significant annual rainfall (~1500mm per year)

the area is slightly more seasonal than true rainforest with a distinct dry-season during

December-March and a drier season during June-August (Newton-Fisher 1999).

Ethical note

Data collection was observational and adhered to the International Primatological Society's

Code of Best Practice for Field Primatology (MacKinnon et al. 2014). All applicable

international, national, and institutional guidelines for the care of animals were followed.

Research was conducted under approval by the Uganda Wildlife Authority and the Uganda

National Council for Science and Technology. All work was in accordance with the ethical

standards of the Budongo Conservation Field Station at which the study was conducted.

Data collection

Researchers and a team of field assistants followed chimpanzees daily (Waibira: from 06:00

to 18:00; Sonso: from 07:00 to 16:30). Long-term data collection included focal individual

activity and party composition taken on a 15-min scan basis. In addition, all unusual events or

otherwise remarkable behaviour were recorded in detail in logbooks for each community,

including births, deaths, and associated descriptions of behaviour (Sonso: since 1993;

Waibira: since 2011).

In addition to long-term records, AS, PF, EF, CF, DT and CH together with field-assistants

SA, JA, GA, BC, and GM of the Budongo Conservation Field Station observed the extended

carry events we report. KET and UP are typically comfortable with the presence of human

observers; however, following the death of KETs infant we avoided selecting her as a focal

individual because we observed apparent signs of greater than typical arousal (e.g. self-

scratching and vigilance) in her interactions with other chimpanzees and we did not want our

extended presence to further impact these. Observations of her behaviour were taken on an ad

libitum basis whenever she joined the party of chimpanzees that included a focal individual,

but we made an effort to locate and observe her for a brief period of time each day to obtain

regular updates on her and her infant's state of decomposition. During the births and deaths

of UP's infants, regular research practices had been adjusted due to the Covid19 pandemic. In

2020 activities were restricted to shorter hours of observation (7:30 to 13:00) and limited to

CH and the permanent field staff, who focused primarily on health monitoring of the

chimpanzees during this period; in late 2021, at the time of UP's second extended infant

carry, restricted research activities had resumed. Researchers and field staff opportunistically

noted any unusual behaviour exhibited. Particular attention was given to how the bodies, and

in one event a potential substitute object, were transported, the response of nearby individuals

to the mother or the carcass, the interactions of the mother with the corpse, and the state of

the corpse. We were not able to collect physiological samples from either corpse to perform

laboratory analyses on the cause of death, nor we were able to retrieve either body for

autopsy.

Results

Over a combined 40-year period of observations (30 years Sonso, 10 years Waibira) a total of

191 births were recorded. Of these: 68 (36%) died in infancy ($\leq$ 5 years) offering

opportunities for their mothers to carry the infant's corpse post-mortem. We found no

evidence for seasonality as deaths (with a confirmed observation month, n=59) occurred

throughout the year (Jan n=5; Feb n=2; Mar n=2; April n=3; May n=1; Jun n=3; Jul n=9; Aug

n=6; Sep n=10; Oct n=6; Nov n=9; Dec n=3). Of the n=68 infant deaths, we excluded n=3

that died together with their mothers and n=12 because they were partially dismembered or

cannibalised during infanticides. Of the remaining n=53 cases, we were able to estimate the

infant's age at death (±1 month) in n=46 cases (87%). The majority (n=25; 54%) died within

the first month, n=17 (37%) at 1-month to 1-year old; n=3 (7%) at 1- to 3-years old, and n=1

(2%) at 3- to 5-years old.

We observed n=12 carries of dead infants by their mothers (Table 1), 23% of observed

opportunities (n=53). To be included as a case of dead infant carrying we required that the

mother be seen with the infant the day after death was estimated to have occurred. In 9 cases

the minimum carry length observed was 1-3 days, in 3 cases we observed a longer minimum

carry of n=18, n=56, and n= 89 days. These are described in more detail below. The 12

carries occurred in both primiparous (n=1) and multi-parous females (n=11), including a 7[th]

born infant. However, these observations are likely an under-estimate of the frequency of

dead infant carrying behaviour in Budongo mothers. In n=36 the mother reappeared alone

and could have carried for an unknown period prior to this. In total there were n=29 cases

where the mother and dead infant were seen together, of these n=12 included a death with the

mother or infanticide with cannibalism. Of the remaining 17 cases, n=12 showed carrying of

the dead-infant, a rate of 71%. Of the five cases where the mother was observed with the

dead infant but did not carry it, all were infanticides.

Table 1 Carrying of dead infants by Budongo chimpanzee mothers: mother-infant dyads

(with the mother first), mother's age, parity (indicated as multi-parous (multi) or primi-

parous (primi)), infant's age, cause of death, duration of carrying

Mother-infant	Age mother (years)	Parity	Age infant (months)	Cause of death	Minimum carrying duration (days)
KG-KG2	21 ±3	multi	0.03	still birth (suspected)	2
ML-ML2	26 ±3	multi	0.03	still birth (suspected)	1

JN-JN2	21 ±1	multi	0.5	unknown	1
ZM-ZM6	41 ±5	multi	0.4	unknown	1
JN-JN4	28 ±1	multi	0.25	infanticide	3
KU-KU5	34 ±3	multi	0.1	infanticide (suspected)	2
KL-KL7	34 ±1	multi	0.12	infanticide	2
ML-ML5	39 ±5	multi	0.06	infanticide (suspected)	3
KET-KYO	20 ±1	primi	25	respiratory infection (inferred)	18
KU-KU7	40 ±3	multi	0.15	unknown	2
UP-UP3	21 ±1	multi	0.25	unknown	56
UP-UP4	22 ±1	multi	0.25	unknown	89

Extended dead infant carrying

A detailed description and videos of the observations are available in the Supplementary

Materials. Here we provide a summary of the key information..

Observation 1: KET, extended dead infant carrying in Waibira

KET's first born infant KYO was last seen alive on the 6[th] of January 2018, aged 25 months.

On the 7th of January 2018 KET was observed carrying KYO who appeared lifeless. Other

chimpanzees were present and were apparently aware of her arrival with the infant but

showed no atypical reactions to KET or the corpse. The likely cause of death was inferred to

be respiratory infection. During the first day KET was observed scratching herself repeatedly

before approaching a water area and when sitting close to a sub-adult male. These scratches

appeared to be a sign of arousal (fast and repeated, and not accompanied by grooming or

response waiting). On several occasions she moved her hand over the dead body apparently

to chase away the flies. Other than this, during the entire 18-day period, she was never

observed to provide any direct maternal care (grooming, inspecting, touching, or peering)

other than carrying, and regularly left the body at short distances (up to 4m) without visually monitoring it. She did not stop others from approaching herself or the dead infant. When moving or feeding in a tree, the dead body was usually (15 out of 16 observations) placed in her right leg pocket, when on the ground she carried the body in her hand or arm (see Online Resources). On one occasion a nulliparous young adult female (MON) was observed to briefly carry the corpse in one hand while KET followed her. Across the 18-days KYO's corpse decomposed, initially increasing in smell. By the 4[th] day, no hair remained on the body. By the 9th day, the body looked "dried". By the 10[th] day the pungent smell and number of flies decreased. It is likely that at this stage the body was completely mummified. No other chimpanzees responded noticeably to either the smell or the flies. On the last day of observed carrying, the body was still intact with only eyes missing and one deformed ear.

Fig. 1 KET holding KYO's mummified body while sitting on a tree (picture taken by AS on the 21.01.2018)

Observation 2: UP, extended dead infant and substitute object carrying in Sonso

UP was first seen on the 25th September 2020 with an apparently recently dead infant (UP3), estimated to be 1-week old. Her two previous infants were killed by intra-community infanticide when under 2-weeks old. While some immature individuals (< 10 years old) inspected the carcass, no others did, and an adult male showed no interest even while grooming UP. The infant's corpse had started to dry out, but had a noticeable smell and flies, and was assumed to have died several days earlier. UP was observed on the 4th and 26th of October and the 8th and 19th of November, carrying the corpse on all occasions. She held it in her hand when on the ground and moved it to a leg pocket when climbing or moving in trees. By the 8th November the corpse appeared fully mummified. UP was last seen with the corpse on the 19th November, a minimum carry duration of 56-days (although likely several days longer given corpse appeared partially dry on first observation). She was next seen on the 23rd November. When approaching a water hole at the base of a tree, she carried a twig ~50cm long in her mouth. As she reached the hole she transferred the twig to her hand and left leg pocket, drank, and then returned it to her mouth (see Online Resources). She continued to carry the twig throughout the morning, including while climbing large trees, and when patrolling with the group over several kilometres. She was seen on the 24th November, and 3rd and 4th December, and was again observed to be carrying a similar twig consistently. She was not seen to put it down on the ground. She was seen briefly on the 5th December, but it was not clear if she had a twig with her, and when she was next observed in the new year (30th January) she no longer carried anything.

Observation 3: UP, 2nd extended dead infant carrying in Sonso

UP was first seen on the 28th August 2021 with another dead infant (UP4), estimated to be 1-week old. The infant's corpse had started to dry out but had a strong smell and on the 30th of

August flies could be seen hovering around it. Given that the corpse still had a strong smell

but was already partially dried, it was assumed to have died at least one week prior. The

cause of death was unclear; however, UP was observed with wounds on her head and on her

left arm. UP was observed using three main carrying styles when traveling on the ground or

moving in trees. She either carried the corpse in one hand (typically left one), in the mouth,

or in one leg pocket (typically left one) (see Online Resources). When resting, she placed the

corpse on her lap, in a leg pocket, held it in one hand, or placed it on the ground close to her.

UP was not observed providing direct maternal care (e.g. grooming, inspecting, or peering) to

the corpse, though she was observed moving her hand around the dead body to chase away

flies on several occasions. On one instance UP was victim of aggression from other females

during which she dropped the corpse, and then followed the group when traveling and left the

corpse behind. Soon after, she was observed returning to retrieve the corpse. We observed a

juvenile male orphan (KJ) following her and peering close to the corpse on a few occasions.

No other individual was observed taking interest in or showing response to the corpse.

Throughout the observation period, UP was often seen in large groups and regularly

socialising with adult males (e.g. grooming). On this occasion there were no observations of

object carrying. UP was seen carrying the corpse on: the 28th, 30th and 31st of August; the 2nd,

6th, 8th, 9th, 11th, 13th, 14th, 16th, 18th-23rd,24th, 25th and 29th of September; the 7th, 12th, 13th,

15th-21st, 26th, 28th of October; and the 3rd, 10th, 14th and 17th of November. On the 18th of

November she was seen without the corpse and had resumed her sexual cycle (with visible

sexual swelling) for the first time since the last pregnancy.

Fig. 2 UP carrying UP4's mummified body in her hand while traveling on the ground

(picture taken by EF on the 30.08.2021)

Discussion

Human responses to deceased conspecifics are remarkable. Despite the fact that dead

individuals have lost all biological importance in terms of kinship, reproduction, or

cooperation, humans do not immediately abandon their dead but exhibit a plethora of post-

mortem social behaviour towards them, including social gatherings, physical tending, and

various burial practices (Parkes et al. 1997). These activities can last for days but are

typically terminated by the onset of physical decay, though certain cultures continue to

interact with the deceased afterwards (e.g. Hollan 1995). There is archaeological evidence

that tending the dead evolved before modern humans (Martinón-Torres et al. 2021), with

burials now documented in *Homo sapiens neanderthalensis* (Rendu et al. 2014) and *Homo*

naledi (Dirks et al. 2015), and interpreted as an indication of some abstract notion of death

and understanding of irreversible loss (Pettitt 2018).

What do great apes experience, cognitively and emotionally, when losing a socially close

individual? In this study we compiled long-term data on chimpanzee mothers that had lost

their infants but continued carrying them for days, raising questions about evolutionary

continuity in the function of this behaviour and the cognition of understanding death. Dead-

infant carrying was practiced by both parous and primiparous mothers with both new-born

and older infants. It occurred in at least a fifth, but more likely in almost three quarters of

cases (if we consider only those cases in which we were able to observe the mother

immediately after the infant's death). As compared to other chimpanzee communities even

this value may be an under-estimate of the frequency with which bereaved female

chimpanzees carry their infants. The Sonso community, in which we made most of our

observations, experiences periodic high levels of infanticide (Lowe et al. 2020). These

infanticide cases are often accompanied by some level of cannibalism or dismemberment,

and/or the infant is taken from the mother (Lowe et al. 2019), which may limit or shorten

mothers' opportunities to carry (Gonçalves and Carvalho 2019; Fernández-Fueyo et al.

2021).

We found no seasonal or infant age effect on dead infant carrying. We also reported three

prolonged cases of extended infant carrying. Our observations suggest that these mothers,

despite the evidence of irreversible loss including absence of any resemblance to living

infants, continued to experience a strong attachment to their deceased infants. Neither female

had any other living offspring and one, after eventually abandoning her dead infant after 56

days, replaced the corpse with a twig, which she carried for at least another two weeks. In our

cases dead-infant carrying was not accompanied by other forms of maternal care, such as

grooming or other forms of maternal attention or interactions, which sometimes occurs very

early after an infant's death (Matsuzawa 1997; Biro 2011), suggesting that the two mothers

had become aware of the biological facts. Both mothers were forced to use atypical modes of

infant carrying, including mouth carrying, more typically used for objects (Lonsdorf et al.

2020), as live infants cling to the back or belly of their mothers during travel. Also relevant is

that, unlike other groups of animals, chimpanzees do not often cannibalise their own dead

infants (but see Fedurek et al. 2020 for two exceptions). Overall, these data suggest that the

'unawareness hypothesis' is an unlikely explanation of chimpanzee behaviour towards dead

conspecifics.

As neither mother appeared to inspect or interact with the infant beyond carrying, our

observations do not support the 'learning about death' hypothesis, although maternal social

behaviour has been seen in other chimpanzee mothers during the initial stages after death

(Matsuzawa 1997; Biro et al. 2010). Similarly, we did not find general support for the

'learning to mother hypothesis'. While both KET and UP were inexperienced mothers

(primiparous or parous but all offspring killed at under a month old),10 of the 11 mothers

who carried their dead infants were parous and two of these cases were with 6^{th} and 7^{th} born

infants. Cases of extended carrying by parous mothers in other groups also suggest limited

support the 'learning to mother hypothesis' (Matsuzawa 1997; Biro et al. 2010; Biro 2011;

Lonsdorf et al. 2020). However, in line with the fact that younger mothers carry dead infants

for longer (Fernández-Fueyo et al. 2021), our observations would fit a suggestion that rare

instances of particularly extended carrying across several months might occur when mothers

have limited mothering experience.

Because we did not collect any hormonal data to assess the levels of stress associated with

this event, we were unable to evaluate the 'grief-management hypothesis' or the 'hormonal

hypothesis'. However, the fact that elevated glucocorticoid levels have been reported in

chimpanzees and baboons following the death of a socially important group member strongly

suggests that these would also be present for chimpanzee mothers in these instances given

that the bond they share with their infants is among their most significant, with typically 3-5

years before weaning, and often up to 10-years of daily close contact (Pusey 1983; Lonsdorf

and Ross 2012; Stanton et al. 2017).

Our observations provide more mixed support for other hypotheses. One observation took

place during the peak of the driest season of the year, whereas the other two took place

during the wettest season of the year. Thus, rather than mummification being the result of

favourable climactic conditions, it is possible mummification was observed because the

extended carrying durations allowed for it. Recent explorations of several large datasets also

found no support for the 'climate hypothesis' (Das et al. 2019; Lonsdorf et al. 2020;

Fernández-Fueyo et al. 2021). The evidence for the 'post-parturient condition hypothesis' is

similarly mixed; of the three extended carries two infants were new-born, while the other was

2-years old. The longest carry reported was for a new-born; however, other new-born infants

were carried for short periods of just a few days. One of the three extended carries terminated

with the resumption of the mother's reproductive cycles. Our data do not provide

straightforward support for the 'hormonal hypothesis' – the extended carrying by KET of her

2-year old would fit this pattern; however, the extended carrying by UP of her two new-borns

do not. Other observations of non-mothers carrying dead infants suggest that the hormonal

hypothesis cannot explain observed extended carrying alone (Watson and Matsuzawa 2018;

Masi 2020). However, hormonal data (that could potentially also show abnormal hormonal

production in either of the two females) are needed to investigate this hypothesis effectively.

While the same argument may apply to the 'maternal-bond strength hypothesis', with KYO's

case providing support and UP's cases providing counterevidence, it may be worth noting

that these were UP's third and fourth infants in a five-year period, the first two having been

killed at under a month old in within-community infanticides (one suspected, one confirmed).

As a result, it may be difficult to assess the nature of her bond with these infants. Her

apparent use of a 'substitute' in her carrying of a twig together with the even more prolonged

second carry suggest she had a particularly strong motivation to carry. These observations,

combined with the fact that all recorded instances of carrying in our dataset concern infants

who died before weaning age, indicate that maternal behaviours, which are not limited to

maternal care, likely play an important role (Fernández-Fueyo et al. 2021).

Table 2 Hypotheses, predictions, and supporting evidence from the present study for dead

infant carrying

Hypothesis	Prediction	Study support
Unawareness	Dead infants are treated as alive.	Unlikely
Post-parturition	Younger infants are carried for longer.	Mixed
Learning about death	Mothers inspect and check state of infant.	Unlikely
Grief-management	Stress levels are lower in mothers carrying dead infants.	Not tested
Learning-to-mother	Primiparous mothers carry dead infants more often/for longer.	Unlikely
Maternal-bond strength	Older infants are carried for longer.	Mixed
Hormonal	Strongly bonded and older infants are carried for longer.	Not tested-Mixed

Within the detailed observations of longer carries, there were two examples of particularly

rare behaviour: KET allowing MON to carry KYO, and UP's stick carrying. Carrying the

infant of others over periods of time longer than one day is an extremely unusual behaviour

in primates, though it has been observed on two occasions in Budongo: adult males snatched

new born infants and carried them still alive for at least two days (in one case the male

continued to carry the infant for a further two days after its death; Notman and Munn 2003;

unpublished long-term data) and was suspected in another case where a daughter was

observed carrying her mother's new infant for several days without the mother being present

(unpublished long-term data). It is possible that KET tolerated MON's behaviour because of

a close bond the two shared, another explanation is that her own bond with the infant's body

apparently had decreased by the 13th day.

We are not aware of any other reports of primate mothers carrying substitute objects

following their infant's death and we are cautious about interpreting this observation.

Chimpanzees may carry objects, including twigs, for many reasons; however, a number of

features suggest that this was related to UP's infant carrying. Prior to the incident, neither UP

nor any other adult Budongo chimpanzees had ever been observed to carry non-food objects

between locations. They are notoriously non-stick-tool users (Whiten et al. 1999; Gruber et

al. 2011), with a substantial literature suggesting that this is both absent, and robust to

scaffolding of its acquisition (Gruber et al. 2009; Gruber 2016). Chimpanzees have been

reported at several sites to engage in 'doll' play, where substitute objects, including logs and

sticks, are carried as if they were a young infant (Matsuzawa 1997; Kahlenberg and

Wrangham 2010). This behaviour typically peaks in juveniles and is more frequent in

females and while it is observed in some adult females it ceased once they became mothers

(Kahlenberg and Wrangham 2010). The description of log doll use in Bossou is of particular

interest here, as it was carried by a juvenile female during the period that her mother was

carrying her sick infant sister, who subsequently died and whose body was also then carried

(Matsuzawa 1997). UP's behaviour was observed multiple times over several weeks. Unlike

the descriptions of other 'dolls' she was not seen to play with or interact with the object,

treating it instead in the same way as she had her infant's corpse. Thus, in addition to object-

carrying being associated with the absence of infants in nulliparous mothers, it may also be

associated with the loss of an infant in bereaved mothers. In humans, the use of transitional

objects has been suggested to function as a coping mechanism for grief following a

bereavement (Lister et al. 2008).

To sum up, our observations are consistent with previous observations that chimpanzee

mothers respond to the death of their infants with carrying behaviour across communities.

Furthermore, our observations support the argument that these mothers act as if they are

aware of the loss but are continuing to display a strong attachment to the bodies of their

infants and may be affected by psychological processes akin to human grieving.

Nevertheless, more detailed hormonal data are needed for a test of this potential mechanism.

A combination of ecological conditions favouring mummification, and social factors, such as

the strong bond shared between mothers and their infants, may explain the three particularly

extended carries by Budongo chimpanzees. While we did not observe other indications of

maternal care in these cases, we are cautious about interpreting this as a wider absence in

Budongo mothers. Mothers' pattern of behavioural responses to death may be individually

specific and nuanced, resulting from a combination of physical, ecological, and psychological

factors, and many more observations are needed to generalise at the population or species

level. Our interpretations are limited by the small dataset and the large number of competing

and non-mutually exclusive hypotheses, and we encourage researchers and long-term field

sites to continue to report the rare behaviours observed in different populations, for example

by contributing to open-access databases such as 'ThanatoBase'

(http://thanatobase.mystrikingly.com), to allow a richer exploration and more robust

hypothesis testing of non-human primates' reaction to death.

Acknowledgments

We are grateful to all field assistants of the Sonso and Waibira communities who provided

essential assistance throughout the observation period. We are thankful to the management,

staff, and researchers of the Budongo Conservation Field Station for their support and thank

the project's founder Vernon Reynolds. We thank the Royal Zoological Society of Scotland

for their long-term financial support to the field station. We thank the Uganda Wildlife

Authority and the Uganda National Council for Science and Technology for permission to

work in Uganda. AS's fieldwork was supported by the European Research Council project

grant to CC (grant agreement number: 679787). CH is supported by funding from the

European Union's 8th Framework Programme, Horizon 2020 (grant agreement number:

802719).

References

Anderson JR (2016) Comparative thanatology. Curr Biol 26:R553–R556.

https://doi.org/10.1016/j.cub.2015.11.010

Anderson JR (2018) Chimpanzees and death. Phil Trans R Soc B 373:20170257.

https://doi.org/10.1098/rstb.2017.0257

Anderson JR (2020) Responses to death and dying: primates and other mammals. Primates

61:1–7. https://doi.org/10.1007/s10329-019-00786-1

Anderson JR, Biro D, Pettitt P (2018) Evolutionary thanatology. Phil Trans R Soc B

373:20170262. https://doi.org/10.1098/rstb.2017.0262

Bercovitch FB (2020) A comparative perspective on the evolution of mammalian reactions to

dead conspecifics. Primates 61:21–28. https://doi.org/10.1007/s10329-019-00722-3

Biro D (2011) Chimpanzee Mothers Carry the Mummified Remains of Their Dead Infants:

Three Case Reports from Bossou. In: Matsuzawa T, Humle T, Sugiyama Y (eds) The

Chimpanzees of Bossou and Nimba. Springer Japan, Tokyo, pp 241–250

Biro D, Humle T, Koops K, et al (2010) Chimpanzee mothers at Bossou, Guinea carry the

mummified remains of their dead infants. Curr Biol 20:R351–R352.

https://doi.org/10.1016/j.cub.2010.02.031

Botting J, van de Waal E (2020) Reactions to infant death by wild vervet monkeys

(Chlorocebus pygerythrus) in KwaZulu-Natal, South Africa: prolonged carrying, non-

mother carrying, and partial maternal cannibalism. Primates 61:751–756.

https://doi.org/10.1007/s10329-020-00851-0

Carter AJ, Baniel A, Cowlishaw G, Huchard E (2020) Baboon thanatology: responses of

filial and non-filial group members to infants' corpses. R Soc open sci 7:192206.

https://doi.org/10.1098/rsos.192206

Courtenay J, Santow G (1989) Mortality of wild and captive chimpanzees. Folia Primatol

52:167–177

Cronin KA, van Leeuwen EJC, Mulenga IC, Bodamer MD (2011) Behavioral response of a

chimpanzee mother toward her dead infant. Am J Primatol 73:415–421.

https://doi.org/10.1002/ajp.20927

Das S, Erinjery JJ, Desai N, et al (2019) Deceased-infant carrying in nonhuman anthropoids:

Insights from systematic analysis and case studies of bonnet macaques (Macaca radiata)

and lion-tailed macaques (Macaca silenus). J Comp Psych 133:156–170.

https://doi.org/10.1037/com0000140

Dirks PH, Berger LR, Roberts EM, et al (2015) Geological and taphonomic context for the

new hominin species Homo naledi from the Dinaledi Chamber, South Africa. Elife

4:e09561

Eggeling WJ (1947) Observations on the Ecology of the Budongo Rain Forest, Uganda. J

Ecol 34:20–87. https://doi.org/10.2307/2256760

Engh AL, Beehner JC, Bergman TJ, et al (2006a) Behavioural and hormonal responses to

predation in female chacma baboons (Papio hamadryas ursinus). Proc R Soc B 273:707–

712. https://doi.org/10.1098/rspb.2005.3378

Engh AL, Beehner JC, Bergman TJ, et al (2006b) Female hierarchy instability, male

immigration and infanticide increase glucocorticoid levels in female chacma baboons.

Anim Behav 71:1227–1237. https://doi.org/10.1016/j.anbehav.2005.11.009

Fashing PJ, Nguyen N, Barry TS, et al (2011) Death among geladas (Theropithecus gelada):

a broader perspective on mummified infants and primate thanatology. Am J Primatol

73:405–409. https://doi.org/10.1002/ajp.20902

Fedurek P, Tkaczynski P, Asiimwe C, et al (2020) Maternal cannibalism in two populations

of wild chimpanzees. Primates 61:181–187. https://doi.org/10.1007/s10329-019-00765-6

Fernández-Fueyo E, Sugiyama Y, Matsui T, Carter AJ (2021) Why do some primate mothers

carry their infant's corpse? A cross-species comparative study. Proc R Soc B 288:1–10.

https://doi.org/10.1098/rspb.2021.0590

Figley CR, Bride BE, Mazza N (1997) Death and trauma: The traumatology of grieving.

Taylor & Francis

Fowler A, Hohmann G (2010) Cannibalism in wild bonobos (Pan paniscus) at Lui Kotale.

Am J Primatol 72:509–514. https://doi.org/10.1002/ajp.20802

Georgiev AV, Melvin ZE, Warketin A-S, et al (2019) Two Cases of Dead-Infant Carrying by

Female Zanzibar Red Colobus (Piliocolobus kirkii) at Jozani-Chwaka Bay National Park,

Zanzibar. Afr Primates 13:57–60

Girard-Buttoz C, Tkaczynski PJ, Samuni L, et al (2021) Early maternal loss leads to short-

but not long-term effects on diurnal cortisol slopes in wild chimpanzees. Elife 10:e64134.

https://doi.org/10.7554/eLife.64134

Gonçalves A, Biro D (2018) Comparative thanatology, an integrative approach: exploring

sensory/cognitive aspects of death recognition in vertebrates and invertebrates. Phil Trans

R Soc B 373:20170263. https://doi.org/10.1098/rstb.2017.0263

Gonçalves A, Carvalho S (2019) Death among primates: a critical review of non-human

primate interactions towards their dead and dying. Biol Rev 94:1502–1529.

https://doi.org/10.1111/brv.12512

Gruber T (2016) Great Apes Do Not Learn Novel Tool Use Easily: Conservatism, Functional

Fixedness, or Cultural Influence? Int J Primatol 37:296–316.

https://doi.org/10.1007/s10764-016-9902-4

Gruber T, Muller MN, Reynolds V, et al (2011) Community-specific evaluation of tool

affordances in wild chimpanzees. Sci Rep 1:128. https://doi.org/10.1038/srep00128

Gruber T, Muller MN, Strimling P, et al (2009) Wild Chimpanzees Rely on Cultural

Knowledge to Solve an Experimental Honey Acquisition Task. Curr Biol 19:1806–1810.

https://doi.org/10.1016/j.cub.2009.08.060

Hill K, Boesch C, Goodall J, et al (2001) Mortality rates among wild chimpanzees. J Hum

Evol 40:437–450. https://doi.org/10.1006/jhev.2001.0469

Hollan D (1995) To the Afterworld and Back: Mourning and Dreams of the Dead among the

Toraja. Ethos 23:424–436. https://doi.org/10.1525/eth.1995.23.4.02a00030

Hosaka K, Matsumoto-Oda A, Huffman MA, Kawanaka K (2000) Reactions to dead bodies

of conspecifics by wild chimpanzees in the Mahale Mountains, Tanzania. Primate Res

16:1–15

Hrdy SB (1999) Mother Nature: Ruthless Competitive Nurturing and Tender: Maternal

Instincts & the Shaping of the Species

Kahlenberg SM, Wrangham RW (2010) Sex differences in chimpanzees' use of sticks as play

objects resemble those of children. Curr Biol 20:R1067–R1068.

https://doi.org/10.1016/j.cub.2010.11.024

Kaplan J (1973) Responses of mother squirrel monkeys to dead infants. Primates 14:89–91.

https://doi.org/10.1007/BF01730518

Kappeler PM, van Schaik CP, Watts DP (2012) The Values and Challenges of Long-Term

Field Studies. In: Kappeler PM, Watts DP (eds) Long-Term Field Studies of Primates.

Springer Berlin Heidelberg, Berlin, Heidelberg, pp 3–18

Kooriyama T (2009) The death of a newborn chimpanzee at Mahale: reactions of its mother

and other individuals to the body. Pan Afr News 16:19–21

Lannen PK, Wolfe J, Prigerson HG, et al (2008) Unresolved Grief in a National Sample of

Bereaved Parents: Impaired Mental and Physical Health 4 to 9 Years Later. JCO

26:5870–5876. https://doi.org/10.1200/JCO.2007.14.6738

Leroux M, Monday G, Chandia B, et al (2021) First observation of a chimpanzee with

albinism in the wild: Social interactions and subsequent infanticide. Am J Primatol.

https://doi.org/10.1002/ajp.23305

Lister S, Pushkar D, Connolly K (2008) Current bereavement theory: Implications for art

therapy practice. Arts Psychother 35:245–250. https://doi.org/10.1016/j.aip.2008.06.006

Lonsdorf EV, Ross SR (2012) Socialization and Development of Behavior. In: The Evolution

of Primate Societies. University of Chicago Press, pp 245–268

Lonsdorf EV, Wilson ML, Boehm E, et al (2020) Why chimpanzees carry dead infants: an

empirical assessment of existing hypotheses. R Soc open sci 7:200931.

https://doi.org/10.1098/rsos.200931

Lowe AE, Hobaiter C, Asiimwe C, et al (2020) Intra-community infanticide in wild, eastern

chimpanzees: a 24-year review. Primates 61:69–82. https://doi.org/10.1007/s10329-019-

00730-3

Lowe AE, Hobaiter C, Newton-Fisher NE (2019) Countering infanticide: Chimpanzee

mothers are sensitive to the relative risks posed by males on differing rank trajectories.

Am J Phys Anthropol 168:3–9. https://doi.org/10.1002/ajpa.23723

MacKinnon KC, Riley EP, Garber PA, et al (2014) Code of Best Practices for Field

Primatology. https://doi.org/10.13140/2.1.2889.1847

Martinón-Torres M, d'Errico F, Santos E, et al (2021) Earliest known human burial in Africa.

Nature 593:95–100. https://doi.org/10.1038/s41586-021-03457-8

Masi S (2020) Reaction to allospecific death and to an unanimated gorilla infant in wild

western gorillas: insights into death recognition and prolonged maternal carrying.

Primates 61:83–92. https://doi.org/10.1007/s10329-019-00745-w

Matsuzawa T (1997) The death of an infant chimpanzee at Bossou, Guinea. Pan Afr News

4:4–6. https://doi.org/10.5134/143350

Nakamichi M, Koyama N, Jolly A (1996) Maternal responses to dead and dying infants in

wild troops of ring-tailed lemurs at the Berenty Reserve, Madagascar. Int J Primatol

17:505–523. https://doi.org/10.1007/BF02735189

Newton-Fisher NE (1999) The diet of chimpanzees in the Budongo Forest Reserve, Uganda:

Diet of Budongo chimpanzees. Afr J Ecol 37:344–354. https://doi.org/10.1046/j.1365-

2028.1999.00186.x

Nicolson NA (1991) Maternal behavior in human and nonhuman primates. In: Understanding

behavior: What primate studies tell us about human behavior. Oxford Univ. Press NY, pp

17–50

Notman H, Munn J (2003) A Case of Infant Carrying by an Adult Male Chimpanzee in the

Budongo Forest. Pan Afr News 10:7–9

Parkes CM, Langani P, Young B, Speck PW (1997) Death and bereavement across cultures.

London: Routledge. Palliat Med 11:427–427

Pettitt P (2018) Hominin evolutionary thanatology from the mortuary to funerary realm: the

palaeoanthropological bridge between chemistry and culture. Phil Trans R Soc B

373:20180212. https://doi.org/10.1098/rstb.2018.0212

Pusey AE (1983) Mother-offspring relationships in chimpanzees after weaning. Anim Behav

31:363–377. https://doi.org/10.1016/S0003-3472(83)80055-4

Ramsay MS, Teichroeb JA (2019) Anecdotes in Primatology: Temporal Trends,

Anthropocentrism, and Hierarchies of Knowledge. Am Anthropol 121:680–693.

https://doi.org/10.1111/aman.13295

Rendu W, Beauval C, Crevecoeur I, et al (2014) Evidence supporting an intentional

Neandertal burial at La Chapelle-aux-Saints. Proc Natl Acad Sci USA 111:81–86.

https://doi.org/10.1073/pnas.1316780110

Reynolds V (2005) The chimpanzees of the Budongo forest: Ecology, behaviour and

conservation. OUP Oxford

Rutz C, Webster MM (2021) *Ethology* adopts the STRANGE framework for animal

behaviour research, to improve reporting standards. Ethology 127:99–101.

https://doi.org/10.1111/eth.13118

Samuni L, Mundry R, Terkel J, et al (2014) Socially learned habituation to human observers

in wild chimpanzees. Anim Cogn 17:997–1005. https://doi.org/10.1007/s10071-014-

0731-6

Stanton MA, Lonsdorf EV, Pusey AE, Murray CM (2017) Do juveniles help or hinder?

Influence of juvenile offspring on maternal behavior and reproductive outcomes in wild

chimpanzees (Pan troglodytes). J Hum Evol 111:152–162.

https://doi.org/10.1016/j.jhevol.2017.07.012

Sugiyama Y, Kurita H, Matsui T, et al (2009) Carrying of dead infants by Japanese macaque

(Macaca fuscata) mothers. AS 117:113–119. https://doi.org/10.1537/ase.080919

Takeshita RSC, Huffman MA, Kinoshita K, Bercovitch FB (2020) Changes in social

behavior and fecal glucocorticoids in a Japanese macaque (Macaca fuscata) carrying her

dead infant. Primates 61:35–40. https://doi.org/10.1007/s10329-019-00753-w

Thompson CL, Hrit R, Melo LCO, et al (2020) Callitrichid responses to dead and dying

infants: the effects of paternal bonding and cause of death. Primates 61:707–716.

https://doi.org/10.1007/s10329-020-00824-3

Tokuyama N, Moore DL, Graham KE, et al (2017) Cases of maternal cannibalism in wild

bonobos (Pan paniscus) from two different field sites, Wamba and Kokolopori,

Democratic Republic of the Congo. Primates 58:7–12. https://doi.org/10.1007/s10329-

016-0582-7

Warren Y, Williamson EA (2004) Transport of dead infant mountain gorillas by mothers and

unrelated females. Zoo Biol 23:375–378. https://doi.org/10.1002/zoo.20001

Watson CFI, Matsuzawa T (2018) Behaviour of nonhuman primate mothers toward their

dead infants: uncovering mechanisms. Phil Trans R Soc B 373:20170261.

https://doi.org/10.1098/rstb.2017.0261

Webster MM, Rutz C (2020) How STRANGE are your study animals? Nature 582:337–340.

https://doi.org/10.1038/d41586-020-01751-5

Whiten A, Goodall J, McGrew WC, et al (1999) Cultures in chimpanzees. Nature 399:682–

685. https://doi.org/10.1038/21415

Wilson ML, Boesch C, Fruth B, et al (2014) Lethal aggression in Pan is better explained by

adaptive strategies than human impacts. Nature 513:414–417.

https://doi.org/10.1038/nature13727