Dead infant carrying by chimpanzee mothers in the Budongo Forest

Adrian Soldati[1,2,3], Pawel Fedurek[3,4,5], Catherine Crockford[4], Sam Adue[3], John Walter

Akankwasa[3], Caroline Asiimwe[3], Jackson Asua[3], Gideon Atayo[3], Boscou Chandia[3], Elodie

Freymann[3,6], Caroline Fryns[2,3], Geresomu Muhumuza[3], Derry Taylor[2,3], Klaus

Zuberbühler[1,2,3], Catherine Hobaiter[1,3]

Affiliations:

[1] School of Psychology and Neuroscience, University of St Andrews, St Andrews, UK

[2] Department of Comparative Cognition, Institute of Biology, University of Neuchâtel,

Neuchâtel, Switzerland

[3] Budongo Conservation Field Station, Masindi, Uganda

[4] Department of Primatology, Max Planck Institute for Evolutionary Anthropology, Leipzig,

Germany

[5] Division of Psychology, Faculty of Natural Sciences, University of Stirling, Stirling, UK

[6] Department of Anthropology and Museum Ethnography, University of Oxford, Oxford, UK

Email of corresponding author: as391@st-andrews.ac.uk

ABSTRACT

It has been suggested that non-human primates can respond to deceased conspecifics in ways

that suggest they experience psychological states not unlike humans, some of which could

indicate they exhibit a notion of death. Here, we report long-term demographic data from two

East African chimpanzee groups. During a combined 40-year observation period we recorded

191 births of which 68 died in infancy, mostly within the first year. We documented the post-

mortem behaviour of the mothers and found that Budongo chimpanzee mothers carried

infants after their death on nine occasions, usually until the body started to decompose after a

few days. However, we also observed three cases of extended carrying lasting for more than

two weeks, one of which was followed by the unusual extended carrying of an object and

another which lasted three months. In each case, the corpses mummified. In addition, we

report four instances of recurring dead infant carrying by mothers, three of whom carried the

corpse for longer during the second instance. We discuss these observations in view of

functional hypotheses of dead infant carrying in primates and the potential proximate

mechanisms involved in this behaviour.

INTRODUCTION

Primate thanatology, the study of the behaviour and underlying physiological and

psychological factors associated with dead or dying individuals in non-human primates

(hereafter primates), continues to raise important questions about human uniqueness

(Anderson 2011, 2018; Anderson et al. 2018). Humans experiencing the loss of socially close

individuals experience significant psychological trauma with long-term physiological effects,

including symptoms of post-traumatic stress disorder, anxiety and depression, and increased

used of health-care resources (Figley et al. 1997; Parkes et al. 1997; Lannen et al. 2008).

Given our biological and social similarities, other primate species – in particular other great

apes – may experience similar cognitive and physiological changes. Cross-species

comparisons, especially in primates, are often used to reveal past evolutionary trajectories of

the hominid lineage. Primate behaviour and physiology in relation to deceased individuals

provide valuable data to better understand the origins of why humans are so powerfully

affected by death. More generally, primate responses to death may provide further insight

into other aspects of animal cognition, such as animacy or the perception of time (Gonçalves

and Carvalho 2019).

Because death is unpredictable and rarely observed in wild primates, the available datasets

are usually anecdotal, and descriptions of events are often incomplete (Watson and

Matsuzawa 2018; Ramsay and Teichroeb 2019). Nevertheless, an increasing number of

primate groups have been habituated to human observers (Kappeler et al. 2012), which has

led to more frequent reporting and more systematic efforts to extract patterns of behaviour in

responses to death (Anderson 2020). These observations have led to claims that primates

respond to death in ways that are similar to humans, by producing strong emotional, social,

and exploratory responses (Watson and Matsuzawa 2018; Gonçalves and Carvalho 2019).

Among non-human animals, the emphasis on primates may result from easier detection,

given the relatively large number of long-term studies, and observations from corvid,

elephant, and cetacean species, which suggest that attentive responses towards dead

conspecifics may be widespread among a wide range of long-lived highly-social species

(Reggente et al. 2016; Gonçalves and Biro 2018; Bercovitch 2020).

Here, we focus on a particularly remarkable behaviour seen in many primates, dead infant

carrying by mothers, which in chimpanzees typically occurs for a period of up to three days

(Gonçalves and Carvalho 2019). Dead infant carrying (also referred as infant corpse carrying)

is the most frequently reported thanatological behaviour and shows substantial variation in

how it is expressed across primate species (Fernández-Fueyo et al. 2021; Watson and

Matsuzawa 2018), as a result, it offers rich data against which to explore possible explanatory

factors. This behaviour has not only been reported in chimpanzees (Matsuzawa 1997; Hosaka

et al. 2000; Kooriyama 2009; Biro et al. 2010) but also in bonobos (Fowler and Hohmann

2010; Tokuyama et al. 2017), gorillas (Warren and Williamson 2004; Masi 2020), chacma

baboons (Carter et al. 2020), red colobus (Georgiev et al. 2019), geladas (Fashing et al.

2011), bonnet and lion-tailed macaques (Das et al. 2019), Japanese macaques (Sugiyama et

al. 2009; Takeshita et al. 2020), and vervet monkeys (Botting and van de Waal 2020) among

others (Fernández-Fueyo et al. 2021), while failed apparent attempts at carrying have been

observed in ring-tailed lemurs (Nakamichi et al. 1996) and marmosets (Thompson et al.

2020). Several hypotheses have been put forward in an attempt to explain the function of and

motivation for this puzzling behaviour (for a review see: Watson and Matsuzawa 2018;

Gonçalves and Carvalho 2019). We summarise six of the main current hypotheses from the

literature to provide a framework for evaluating the potential functions and mechanisms of

new observations and discuss their implications.

A first group of hypotheses presumes that primate mothers are unable to understand the

ramifications of death and their behavioural responses are simply side effects of evolved

physiological mechanisms. Specifically, the 'unawareness hypothesis' states that mothers are

unable to discriminate between temporarily unresponsive and irreversibly deceased

individuals and simply try to avoid the costs of premature abandonment (Hrdy 1999). The

hypothesis predicts that mothers are unable to understand their infants' state and continue

providing maternal care (e.g., grooming) and try to elicit responsiveness (e.g., poking,

smelling). With increasing time, the ambiguity will dissolve due to the accelerating

decomposition of the corpse, which can be mediated by local climate. A second prediction

therefore is that dry and particularly hot or cold conditions favouring mummification should

also favour prolonged carrying of dead infants (see also 'climate hypothesis': Matsuzawa

1997; Fashing et al. 2011). The 'post-parturient condition' hypothesis (also referred as

'hormonal', see Gonçalves and Carvalho 2019) proposes that the maternal physiological

conditions associated with pregnancy and birth favour persistent care of the dead infant as

long as the mother is lactating or until resumption of ovulation (Biro et al. 2010; Kaplan

1973). After giving birth, the endocrine system of the mother releases hormones (e.g.,

oxytocin) that stimulate maternal behaviours (Keverne 1988; Bercovitch 2020). Here, the

prediction is that when a mother loses her infant close to its birth (and the associated

hormonal changes), they are more likely to continue to invest in infant care, increasing the

likelihood and perhaps duration of any post-mortem carrying behaviour.

A second kind of hypotheses assumes that primate mothers can have a notion of death,

provided they have relevant personal experience. Specifically, the 'learning about death'

hypothesis states that chimpanzees do not intuitively understand death but can acquire the

notion by learning to attend to the relevant cues (Cronin et al. 2011). Here, the predication is

that younger or inexperienced mothers are more likely to carry dead-infants than older or

experienced mothers to learn about death (Watson and Matsuzawa 2018). While females

could learn by handling the corpses of the offspring of others, another possible source of

information comes from experiencing multiple deaths of their own offspring. In addition,

experienced mothers might also be more aware of the irreversible change of death due to

having interacted with more living infants and/or for longer periods of time. Overall, if

mothers' reason for carrying is due to a lack of experience or understanding of what a dead

infant is like, they should be less likely to carry or carry for shorter periods dead-infants in

subsequent cases. An alternative hypothesis, the 'grief-management hypothesis', also

assumes that chimpanzees possess a notion of death, and suggests dead-infant carrying

represents a consoling strategy to cope with grief. This hypothesis predicts that mothers who

are able to carry their dead infants experience lower levels of 'stress' hormones (i.e.,

glucocorticoids) than mothers who did not carry them (Nicolson 1991), which could be a

strategy to cope with stress associated with infant loss (e.g., Takeshita et al. 2020).

A third group of hypotheses are agnostic about whether chimpanzees possess a notion of

death but proposes various adaptive mechanisms that favour post-mortem mothering

behaviour. First, the 'learning-to-mother hypothesis' states that dead-infant carrying

improves maternal skills (Warren and Williamson 2004), predicting that the behaviour should

mainly be observed in inexperienced primiparous females. Second, the 'maternal-bond

strength hypothesis' predicts that mothers with older infants share a stronger bond, due to

more extensive association and interactions between them, and are thus more likely to show

extensive carrying as compared to mothers with younger infants (Watson and Matsuzawa

2018).

The existence of multiple – sometime contrasting – hypotheses is likely a reflection of the

small and highly variable data available and the diversity of potential drivers of this

behaviour across different species. A recent systematic study on 18 primate species (n=48

cases) showed that duration of infant carrying is affected by the age of the mother, with older

mothers carrying for longer periods (Das et al. 2019). However, some species do not fit this

pattern, with previous findings by Sugiyama and colleagues (2009) detecting no effect of age

in a large study on Japanese macaques. Infants that died of sickness were carried for longer

than those who were stillborn or victims of infanticide, while the age of the infant did not

influence duration of carrying (Das et al. 2019). More recently, a study using the largest

primate database to date (n=50 species and n=409 cases) found that dead infant carrying was

more likely to occur when the cause of death was non-traumatic, when mothers were

younger, and that older infants are carried for shorter periods (Fernández-Fueyo et al. 2021).

While dead infant carrying is a shared behaviour in primates, the frequency of occurrence,

duration of carrying, and ease of observation, vary considerably between species and

individuals. Nevertheless, given the observations currently available, carrying duration seems

to be the longest in great apes, particularly in chimpanzees (Das et al. 2019). Wild

chimpanzees have higher levels of infant mortality compared to captive conspecifics

(Courtenay and Santow 1989; Hill et al. 2001), which provides increased opportunities to

observe mothers' responses to their infants' death (Watson and Matsuzawa 2018).

Chimpanzee mothers typically carry their dead infants for a few days, though a recent

analysis of the largest chimpanzee dataset (n=33 cases) did not provide clear support for any

of the previous hypotheses (Lonsdorf et al. 2020). Lonsdorf and colleagues (2020) proposed

that the 'unawareness hypothesis' is unlikely because of the presence of atypical carrying

postures and sensory cues after death as compared to those displayed towards living infants.

However, while a decomposing body may be perceived differently from a living one through

several sensory modalities (Gonçalves and Biro 2018), chimpanzee mothers may also be

responding to an inability of their infant to grasp, rather than a recognition of its status as

living or dead. Here, we report observations of dead infant carrying by female chimpanzees

using a 40-year dataset of two study groups of the Budongo Forest, Uganda, including three

detailed observations of extended dead infant carrying by two different females.

METHODS

Study site and subjects

The Budongo Forest Reserve is a semi-deciduous tropical rain forest located along the

Western Rift Valley in Uganda. This reserve is made of 793 km^2 of protected forest and

grassland, including 482 km^2 of continuous forest cover (Eggeling 1947) . The reserve

contains a population of approximated 600 East African chimpanzees (*Pan troglodytes*

schweinfurthii). Our observations took place in two adjacent communities, Sonso and

Waibira, studied regularly by researchers and followed on a daily basis by field assistants

since 1990 and 2011 respectively, contributing a combined 43-years of long-term

observational records (Reynolds 2005; Samuni et al. 2014).

At the time of the events, the Waibira community contained an estimated 120 individuals, 96

of which could be individually recognised. Individuals involved in the first event were Ketie

(KET), a 20-year old adult primiparous female (estimated birth 1998) and her 2-year old

female infant Karyo (KYO) born in December 2015. The Sonso community contained 65

named individuals, in addition to three unnamed females in the process of immigrating. The

individuals involved in the event were Upesi (UP), a 21-year old parous female (estimated

birth 1999) and a) in the second event, her recently born unnamed unsexed infant UP3, born

mid-September 2020 and b) in the third event, her 4th born unnamed unsexed infant UP4,

estimated birth 7th August 2021. Her first two infants (born in 2017 and 2018) were victims

of within-community infanticide before reaching a month old (see Leroux et al. 2021 for one

reported case).

We considered the scope for bias in our study subjects by using the STRANGE framework to

report potential sampling biases in our study (Webster and Rutz 2020; Rutz and Webster

2021). The Sonso community are of typical size whereas the Waibira community are

particularly large as compared to that of other chimpanzees (in a recent comparison of 18

groups across three subspecies: *P.t. schweinfurthii, P.t. troglodytes, P.t. verus*; communities

range from 7-144 individuals with a mean 42; within these data the East African sub-species

(*P.t. schweinfurthii*) range is 18-144 with a median 49; Wilson et al. 2014). Sonso have a

typical female-biased sex ratio among mature individuals (M:F; 1:1.7), whereas the Waibira

community have more unusual evenly-balanced sex ratio among mature individuals (M:F;

1:1.2; mean among 9 *P.t. schweinfurthii* communities 1:1.7; Wilson et al. 2014). Of relevance

to sampling biases in infant mortality and opportunities to carry dead infants, the Sonso

community experience high levels of infanticide (Lowe et al. 2019, 2020). A medium altitude

rainforest (~1100m) with significant annual rainfall (~1500mm per year) the area is slightly

more seasonal than true rainforest with a distinct dry-season during December-March and a

drier season during June-August (Newton-Fisher 1999).

Observations of wild chimpanzees in dense secondary forest are often challenging and

individuals in highly fission-fusion East African chimpanzee communities may not be

observed for days or weeks (Badihi et al. Preprint). In addition, we often choose not to

follow, or only to follow at an extended distance, individuals who are experiencing high-

stress events, such as maternal loss. We considered a mother to be carrying a dead infant if

they were seen to be doing so on the day following their infant's death. In doing so, we may

underestimate occurrences of shorter carries (i.e., under a day), or the total duration of

carries. Similarly, as we rarely observed the moment that the mother stopped carrying the

dead infant, our estimation of duration may be particularly conservative. However, given the

challenges in discriminating the precise moment of infant death from, for example:

unconsciousness, we felt that a conservative approach was appropriate.

Ethical note

Data collection was observational and adhered to the International Primatological Society's

Code of Best Practice for Field Primatology (MacKinnon et al. 2014). All applicable

international, national, and institutional guidelines for the care of animals were followed.

Research was conducted under approval by the Uganda Wildlife Authority and the Uganda

National Council for Science and Technology. All work was in accordance with the ethical

standards of the Budongo Conservation Field Station at which the study was conducted.

Data collection

Researchers and a team of field assistants followed chimpanzees daily (Waibira: from 06:00

to 18:00; Sonso: from 07:00 to 16:30). Long-term data collection included continuous focal

individual activity and party composition taken on a 15-min scan basis. In addition, all

unusual events or otherwise remarkable behaviour were recorded in detail in logbooks for

each community, including births, deaths, and associated descriptions of behaviour (Sonso:

since 1993; Waibira: since 2011).

In addition to long-term records, AS, PF, EF, CF, DT and CH together with field-assistants

SA, JA, GA, BC, and GM of the Budongo Conservation Field Station observed the extended

carry events we report. KET and UP are typically comfortable with the presence of human

observers; however, following the death of KETs infant we avoided selecting her as a focal

individual because we observed behavioural cues known (or assumed) to be associated with

greater than typical arousal (e.g., self-scratching and vigilance) in her interactions with other

chimpanzees and we did not want our extended presence to further impact these.

Observations of her behaviour were taken on an ad libitum basis whenever she joined the

party of chimpanzees that included a focal individual, but we made an effort to locate and

observe her for a brief period of time each day to obtain regular updates on her and her

infant's state of decomposition. During the births and deaths of UP's infants, regular research

practices had been adjusted due to the Covid19 pandemic. In 2020 activities were restricted

to shorter hours of observation (7:30 to 13:00) and limited to CH and the permanent field

staff, who focused primarily on health monitoring of the chimpanzees during this period; in

late 2021, at the time of UP's second extended infant carry, restricted research activities had

resumed. Researchers and field staff opportunistically noted any unusual behaviour exhibited.

Particular attention was given to how the bodies, and in one event a potential substitute

object, were transported, the response of nearby individuals to the mother or the carcass, the

interactions of the mother with the corpse, and the state of the corpse. We were not able to

collect physiological samples from either corpse to perform laboratory analyses on the cause

of death, nor we were able to retrieve either body for autopsy.

RESULTS

Over a combined 40-year period of observations (30 years Sonso, 10 years Waibira) a total of

191 births were recorded. Of these: 68 (36%) died in infancy (≤ 5 years) offering

opportunities for their mothers to carry the infant's corpse post-mortem. We found no

evidence for seasonality as deaths (with a confirmed observation month, n=59) occurred

throughout the year (Jan n=5; Feb n=2; Mar n=2; Apr n=3; May n=1; Jun n=3; Jul n=9; Aug

n=6; Sep n=10; Oct n=6; Nov n=9; Dec n=3). Of the 68 infant deaths, we excluded three that

died together with their mothers and 12 because they were partially dismembered or

cannibalised during infanticides. Of the remaining 53 cases, for 46 (87%) we were able to

estimate the infant's age at death (±1 month). The majority (n=25; 54%) died within the first

month, 17 (37%) at 1-month to 1-year old; 3 (7%) at 1- to 3-years old, and one (2%) at 3- to

5-years old.

We observed 12 carries of dead infants by their mothers (Table 1), 23% of observed

opportunities (n=53). To be included as a case of dead infant carrying we required that the

mother be seen with the infant the day after death was estimated to have occurred. In nine

cases the minimum carry length observed was 1-3 days, in three cases we observed a longer

minimum carry of n=18, n=56, and n= 89 days. These are described in more detail below.

The 12 carries occurred in both primiparous (n=1) and multi-parous females (n=11),

including a 7[th] born infant. However, these observations are likely an under-estimate of the

frequency of dead infant carrying behaviour in Budongo mothers. In 36 the mother

reappeared alone and could have carried for an unknown period prior to this. In total there

were 29 cases where the mother and dead infant were seen together, of these 12 included a

death with the mother or infanticide with cannibalism. Of the remaining 17 cases, 12 showed

carrying of the dead-infant, a rate of 71%. Of the five cases where the mother was observed

with the dead infant but did not carry it, all were infanticides (without cannibalism). Finally,

four out of eight mothers (ML, JN, KU, UP) were observed carrying the dead bodies of two

of their infants (Table 1), one carried for the same duration and three carried for longer

durations on the second occasion.

Table 1 Carrying of dead infants by Budongo chimpanzee mothers: mother-infant dyads

(with the mother first), mother's age, parity (indicated as multi-parous (multi) or primi-

parous (primi)), infant's age, cause of death, duration of carrying, community (Sonso or

Waibira), and date of infant death.

Mother-infant	Age mother (years)	Parity	Age infant (months)	Cause of death	Minimum carrying duration (days)	Community	Date of infant death
KG-KG2	21 ±3	multi	0.03	still birth (suspected)	2	Sonso	09.09.1997
ML-ML2	26 ±3	multi	0.03	still birth (suspected)	1	Sonso	08.07.2002
JN-JN2	21 ±1	multi	0.5	unknown	1	Sonso	24.01.2005
ZM-ZM6	41 ±5	multi	0.4	unknown	1	Sonso	19.03.2009
JN-JN4	28 ±1	multi	0.25	infanticide	3	Sonso	10.11.2012
KU-KU5	34 ±3	multi	0.1	infanticide (suspected)	2	Sonso	18.02.2013

KL-KL7	34 ±1	multi	0.12	infanticide	2	Sonso	30.07.2013
ML-ML5	39 ±5	multi	0.06	infanticide (suspected)	3	Sonso	17.11.2014
KET-KYO	20 ±1	primi	25	respiratory infection (inferred)	18	Waibira	06.01.2018
KU-KU7	40 ±3	multi	0.15	unknown	2	Sonso	26.11.2019
UP-UP3	21 ±1	multi	0.25	unknown	56	Sonso	23.09.2020
UP-UP4	22 ±1	multi	0.25	unknown	89	Sonso	21.08.2021

We found no evidence for a clear effect of seasonality on the onset of carrying behaviour, but

our sample size is small (see Fig. 1). The Budongo Forest shows two marked periods of

intense rainfall (March – May, September – November), an intense drier season (December –

February) and a light drier season (June – August). Carrying of dead infants was as likely to

occur in the Wet (n=6) or Dry (n=6) seasons ($X^2 = 0.00$, p=1.000).

Fig. 1 Dead infant carrying occurrences for each mother-infant pairs relative to the average

daily rainfall (data extracted from Budongo Conservation Field Station long-term records

1993-2018). Single red dots represent occurrences of carrying that lasted between 1 and 3

days. Red dots connected with a line represent the approximated duration of extended

carrying, with the dots representing the start and end of carrying.

We found a possible effect of infant age. Our sample size of older infants was very small: of

the 46 opportunities to carry described above, we only observed four infants who died

between 1-5 years old, and a single case (25%) of carrying (aged 2-years, described below as

the extended carry by KET of KYO). We found a similar frequency of carries (n=11), given

opportunities to carry (n=42) of infants under one year old. However, within this larger

sample of younger infants aged 1-12 months, we found that only very young infants (aged

under 1-month) were carried. When considering the number of opportunities to carry, infants

aged under 1-month (11 of 25 cases, 44%) were more likely to be carried than infants aged 1-

12months (0 of 17 cases; Fisher's exact test p=0.010).

Extended dead infant carrying

A detailed description and videos of the observations are available in the Supplementary

Materials. Here we provide a summary of the key information.

Observation 1: KET, extended dead infant carrying in Waibira

KET's first born infant KYO was last seen alive on the 6th January 2018, aged 25 months. On

the 7th January 2018 KET was observed carrying KYO who appeared lifeless. Other

chimpanzees were present and were apparently aware of her arrival with the infant but

showed no atypical reactions to KET or to the corpse. Based on the fact that most

chimpanzees exhibited signs of respiratory infection during that period and given the infant

had shown no other sign of illness, the likely cause of death was inferred to be respiratory

infection. During the first day KET was observed scratching herself repeatedly before

approaching a water area and when sitting close to a sub-adult male. These scratches

appeared to be a sign of arousal (fast and repeated, and not accompanied by grooming or

response waiting). On several occasions she moved her hand over the dead body apparently

to chase away the flies. Other than this, during the entire 18-day period, she was never

observed to provide any direct maternal care (grooming, inspecting, touching, or peering)

other than carrying, and regularly left the body at short distances (up to 5m) without visually

monitoring it. She did not stop others from approaching herself or the dead infant. When

moving or feeding in a tree, the dead body was usually (15 out of 16 observations) placed in

her right leg pocket, when on the ground she carried the body in her hand or arm (see Online

Resources). On one occasion a nulliparous young adult female (MON) was observed to

briefly carry the corpse in one hand while KET followed her. Across the 18-days KYO's

corpse decomposed, initially increasing in smell. By the 4^{th} day, no hair remained on the

body. By the 9th day, the body looked "dried". By the 10^{th} day the pungent smell and number

of flies decreased. It is likely that at this stage the body was completely mummified. No other

chimpanzees responded noticeably to either the smell or the flies. On the last day of observed

carrying, the body was still intact with only eyes missing and one deformed ear.

Fig. 2 KET holding KYO's mummified body while sitting on a tree (picture taken by

"anonymous author" on the 21.01.2018)

Observation 2: UP, extended dead infant and substitute object carrying in Sonso

UP was first seen on the 25th September 2020 with an apparently recently dead infant (UP3),

estimated to be 1-week old. Her two previous infants were killed by intra-community

infanticide when under 2-weeks old. While some immature individuals (< 10 years old)

inspected the carcass, no others did, and an adult male showed no interest even while

grooming UP. The infant's corpse had started to dry out, but had a noticeable smell and flies,

and was assumed to have died several days earlier. UP was observed on the 4th and 26th

October and the 8th and 19th November, carrying the corpse on all occasions. She held it in

her hand when on the ground and moved it to a leg pocket when climbing or moving in trees.

By the 8th November the corpse appeared fully mummified. UP was last seen with the corpse

on the 19th November, a minimum carry duration of 56-days (although likely several days

longer given corpse appeared partially dry on first observation). She was next seen on the

23rd November. When approaching a water hole at the base of a tree, she carried a twig

~50cm long in her mouth. As she reached the hole she transferred the twig to her hand and

left leg pocket, drank, and then returned it to her mouth (see Online Resources). She

continued to carry the twig throughout the morning, including while climbing large trees, and

when patrolling with the group over several kilometres. She was seen on the 24[th] November,

and 3[rd] and 4[th] December, and was again observed to be carrying a similar twig consistently.

She was not seen to put it down on the ground. She was seen briefly on the 5[th] December, but

it was not clear if she had a twig with her, and when she was next observed in the new year

(30[th] January) she no longer carried anything.

Observation 3: UP, 2[nd] extended dead infant carrying in Sonso

UP was first seen on the 28[th] August 2021 with another dead infant (UP4), estimated to be 1-

week old. While we did not see her with a live infant, and it is thus at least possible that the

infant was not hers, we believe this to be very unlikely. UP was last observed in maximal

oestrus approximately 8 months prior to being observed with the dead infant, and all other

Sonso females were either confirmed to be pregnant, had unweaned infants at the time, had

been observed cycling regularly in the months prior to the observation, or were considered to

be post-reproductive. Finally, we know of no other reports of extended carrying of non-kin

dead infants. The infant's corpse had started to dry out but had a strong smell and on the 30[th]

August flies could be seen hovering around it. Given that the corpse still had a strong smell

but was already partially dried, it was assumed to have died at least one week prior. The

cause of death was unclear; however, UP was observed with wounds on her head and on her

left arm. UP was observed using three main carrying styles when traveling on the ground or

moving in trees. She either carried the corpse in one hand (typically left one), in the mouth,

or in one leg pocket (typically left one) (see Online Resources). When resting, she placed the

corpse on her lap, in a leg pocket, held it in one hand, or placed it on the ground close to her.

UP was not observed providing direct maternal care (e.g., grooming, inspecting, or peering)

to the corpse, though she was observed moving her hand around the dead body to chase away

flies on several occasions. On one instance UP was victim of aggression from other females

during which she dropped the corpse, and then followed the group when traveling and left the

corpse behind. Soon after, she was observed returning to retrieve the corpse. We observed a

juvenile male orphan (KJ) following her and peering close to the corpse on a few occasions.

No other individual was observed taking interest in or showing response to the corpse.

Throughout the observation period, UP was often seen in large groups and regularly

socialising with adult males (e.g., grooming). On this occasion there were no observations of

object carrying. UP was seen carrying the corpse on: the 28th, 30th and 31st August; the 2nd,

6th, 8th, 9th, 11th, 13th, 14th, 16th, 18th-23rd, 24th, 25th and 29th September; the 7th, 12th, 13th, 15th-

21st, 26th, 28th October; and the 3rd, 10th, 14th and 17th November. On the 18th November she

was seen without the corpse and had resumed her sexual cycle (with visible sexual swelling)

for the first time since the last pregnancy.

Fig. 3 UP carrying UP4's mummified body in her hand while traveling on the ground

(picture taken by "anonymous author" on the 30.08.2021)

DISCUSSION

Human responses to deceased conspecifics are remarkable. Despite the fact that dead

individuals have lost all biological importance in terms of kinship, reproduction, or

cooperation, humans do not immediately abandon their dead but exhibit a plethora of post-

mortem social behaviour towards them, including social gatherings, physical tending, and

various burial practices (Parkes et al. 1997). These activities can last for days or weeks but

are typically terminated by the onset of physical decay, though certain cultures continue to

interact with the deceased for long after (e.g., Hollan 1995). There is archaeological evidence

that tending the dead evolved before modern humans (Martinón-Torres et al. 2021), with

mortuary behaviour documented in *Homo sapiens neanderthalensis* (Rendu et al. 2014) and

Homo naledi (Dirks et al. 2015), having been interpreted as an indication of some abstract

notion of death and understanding of irreversible loss (Pettitt 2018).

What are the behavioural responses of great apes when losing a socially close individual? In

this study we compiled over 40-years of long-term data on chimpanzee mothers from the

Budongo forest that had lost their infants but continued carrying them for days, raising

questions about evolutionary continuity in the function of this behaviour and the cognition of

understanding death. Dead-infant carrying was practiced by both parous and primiparous

mothers with both new-born and older infants. It occurred in at least a fifth of the cases if we

consider only those in which we were able to observe the mother immediately after the

infant's death. However, this value is very likely an under-estimate of the frequency with

which bereaved Budongo female chimpanzees carry their infants. The Sonso community, in

which we made most of our observations, experiences periodic high levels of infanticide

(Lowe et al. 2020). These infanticide cases are often accompanied by some level of

cannibalism or dismemberment, and/or the infant is taken from the mother (Lowe et al.

2019), which may limit or shorten mothers' opportunities to carry (Fernández-Fueyo et al.

2021; Gonçalves and Carvalho 2019). In many cases the mother was simply not seen in the

days following her infant's death, in the limited number of cases where the mother was seen

in the days following the infant's death, and where the death was not an infanticide with

cannibalism of the corpse, almost three quarters of cases involved carrying of the dead infant.

We found no seasonal effect on dead infant carrying, but we do show a possible age effect, in

which – amongst infants who died at under 1-year old – very young infants were more likely

to be carried. Our observations also support the suggestion that infants may be less likely to

be carried following traumatic death (Fernández-Fueyo et al. 2021). Most cases of infant

carrying were relatively short (minimum confirmed carry length of a few days); however, we

also reported three prolonged cases of extended infant carrying. Our observations suggest that

these mothers, despite the evidence of irreversible loss including absence of any resemblance

to living infants, continued to experience a strong attachment to their deceased infants.

Neither female had any other living offspring and one, after eventually abandoning her dead

infant after 56 days, carried an object: a twig, for at least another two weeks. In the three

cases of prolonged carrying that we report, dead-infant carrying was not accompanied by

other forms of maternal care, such as grooming or other forms of maternal attention or

interactions, which sometimes occurs very early after an infant's death (Matsuzawa 1997;

Biro 2011), suggesting that the two mothers had become aware of the biological facts. Both

mothers were forced to use atypical modes of infant carrying, including mouth carrying, more

typically used for objects (Lonsdorf et al. 2020), as live infants cling to the back or belly of

their mothers during travel. Also relevant is that, unlike other groups of animals,

chimpanzees do not often cannibalise their own dead infants (but see Fedurek et al. 2020 for

two exceptions). Overall, these data suggest that the 'unawareness hypothesis' is unable to

fully explain chimpanzee behaviour towards dead conspecifics.

As neither mother appeared to inspect or interact with the infant beyond carrying, our

observations do not support the 'learning about death' hypothesis, although maternal social

behaviour has been seen in other chimpanzee mothers during the initial stages after death

(Matsuzawa 1997; Biro et al. 2010). Further support against the 'learning about death'

hypothesis is provided by the mothers who carried their dead infants on more than on

occasion – of the four mothers, three carried for longer on the second occasion (one for the

same amount of time). Thus, a lack of knowledge about dead infants cannot fully explain

their reason for carrying. This pattern appears repeated in other studies: we extracted the data

from Lonsdorf et al. (2020, n=6), Biro (2011, n=1), and Hanamura et al. (2015, n=2). Six of

the nine cases described were longer on subsequent carries. Recurring and prolonged carrying

behaviours seem to indicate that mothers are not unaware of death and instead their responses

are affected by other factors.

While both KET and UP were inexperienced mothers (primiparous and parous but all

offspring killed at under a month old respectively), 10 of the 11 mothers who carried their

dead infants were parous and two of these cases were with 6^{th} and 7^{th} born infants. Cases of

extended carrying by parous mothers in other groups also suggest limited support for the

'learning to mother hypothesis' (Matsuzawa 1997; Biro et al. 2010; Biro 2011; Lonsdorf et

al. 2020). However, in line with the fact that younger primate mothers are more likely to

carry dead infants (Fernández-Fueyo et al. 2021), our three observations support the

suggestion that rare instances of particularly extended carrying across several months might

be more frequent in young mothers. Nevertheless, repeated prolonged carries (as seen in UP)

do not support the suggestion that this is due to inexperience and, given that no clear pattern

emerges when considering observations across sites (e.g., Hanamura et al. 2015; Lonsdorf et

al. 2020) and that sample sizes of chimpanzee mothers remain very small, these rare

instances could also reflect individual differences.

Because we did not collect any hormonal data to assess the levels of stress associated with

dead infant carrying, we were unable to evaluate our observations under the 'grief-

management hypothesis'. However, a recent study by Girard-Buttoz and colleagues (2021)

reported elevated cortisol levels in infant chimpanzees who lost their mothers, supporting the

notion that disruption of the mother-infant bond leads to elevated stress. Similarly, female

baboons experience high levels of glucocorticoids when losing an ally to predation and in

periods of infanticidal attacks (Engh et al. 2006a, b). Given the bond chimpanzee mothers

share with their infants is among their most significant, with typically 3-5 years before

weaning, and often up to 10-years of daily close contact (Pusey 1983; Lonsdorf and Ross

2012; Stanton et al. 2017), we expect chimpanzee mothers to experience elevated stress

levels following the death of their infant.

One of the three prolonged carries took place during the peak of the driest season of the year,

whereas the other two took place during the wettest season of the year, and we found no

effect of seasonality in our wider data. Thus, rather than mummification being the result of

favourable climactic conditions, it is possible mummification was observed because the

extended carrying durations allowed for it (see also Biro 2011). Recent explorations of

several large datasets also found no support for a 'climate hypothesis' (Das et al. 2019;

Lonsdorf et al. 2020; Fernández-Fueyo et al. 2021). Of the three extended carries two infants

were new-born, while the other was 2-years old. The longest carry reported was for a new-

born; however, other new-born infants were carried for short periods of just a few days. One

of the three extended carries terminated with the resumption of the mother's reproductive

cycles. Our observation of carrying being more likely in very young infants (under 1-month)

than in infants aged 1-12months, as well as the extended carrying by UP of her two young

new-borns would fit the pattern proposed for the 'post parturient hypothesis'; however, the

extended carrying by KET of her 2-year old infant does not. Other observations of non-

mothers carrying dead infants suggest that while post parturient effects may contribute to this

behaviour, they cannot explain extended carrying alone (Watson and Matsuzawa 2018; Masi

2020). However, hormonal data are needed to investigate this hypothesis effectively. While

the same argument may apply to the 'maternal-bond strength hypothesis', with KET's case

providing support and UP's cases providing counterevidence, it may be worth noting that

these were UP's third and fourth infants in a five-year period, the first two having been killed

at under a month old in within-community infanticides (one suspected, one confirmed). As a

result, it may be difficult to assess the nature of her bond with these infants. Her possible use

of a 'substitute object' in her carrying of a twig together with the even more prolonged

second carry suggest she had a particularly strong motivation to carry. These observations,

combined with the fact that all recorded instances of carrying in our dataset concern infants

who died before weaning age, seem to indicate that maternal behaviours, which are not

limited to maternal care, and the bond between the mother and the offspring likely play an

important role in dead infant carrying (Fernández-Fueyo et al. 2021).

Table 2 Hypotheses, predictions, and supporting evidence from the present study for dead

infant carrying

Hypothesis	Prediction	Study support

Unawareness	Dead infants are treated as alive.	Unlikely
Post-parturition	Young infants are carried for longer.	Mixed
Learning about death	Mothers inspect and check state of infant.	Unlikely
Grief-management	Stress levels are lower in mothers carrying dead infants.	Not tested
Learning-to-mother	Primiparous mothers carry dead infants more often/for longer.	Unlikely
Maternal-bond strength	Strongly bonded and intermediate/old infants are carried for longer.	Mixed

Within the detailed observations of longer carries, there were two examples of particularly

rare behaviour: KET allowing MON to carry KYO, and UP's stick carrying. Carrying the

infant of others over periods of time longer than one day is an extremely unusual behaviour

in primates, though it has been observed on two occasions in Budongo: adult males snatched

new born infants and carried them still alive for at least two days (in one case the male

continued to carry the infant for a further two days after its death; Notman and Munn 2003;

unpublished long-term data) and was suspected in another case where a daughter was

observed carrying her mother's new infant for several days without the mother being present

(unpublished long-term data). It is possible that KET tolerated MON's behaviour because

they may share a close bond, but another explanation is that her own bond with the infant's

body apparently had decreased by the 13[th] day.

We are not aware of any other reports of primate mothers carrying substitute objects

following their infant's death and we are cautious about interpreting this unique and unusual

observation. Chimpanzees may carry objects, including twigs, for many reasons; however, a

number of features suggest that this was related to UP's infant carrying. Prior to the incident,

neither UP nor any other adult Budongo chimpanzees had ever been observed to carry non-

food objects between locations. They are notoriously non-stick-tool users (Whiten et al.

1999; Gruber et al. 2011), with a substantial literature suggesting that this is both absent, and

robust to scaffolding of its acquisition (Gruber et al. 2009; Gruber 2016). Chimpanzees have

been reported at several sites to engage in 'doll' play, where substitute objects, including logs

and sticks, are carried as if they were a young infant (Matsuzawa 1997; Kahlenberg and

Wrangham 2010). This behaviour typically peaks in juveniles and is more frequent in

females and while it is observed in some adult females it ceased once they became mothers

(Kahlenberg and Wrangham 2010). The description of log doll use in Bossou is of particular

interest here, as it was carried by a juvenile female during the period that her mother was

carrying her sick infant sister, who subsequently died and whose body was also then carried

(Matsuzawa 1997). UP's behaviour was observed multiple times over several weeks. Unlike

the descriptions of other 'dolls' she was not seen to play with or interact with the object,

treating it instead in the same way as she had her infant's corpse. Thus, in addition to object-

carrying being associated with the absence of infants in nulliparous mothers, it may also be

associated with the loss of an infant in bereaved chimpanzee mothers. In humans, the use of

transitional objects has been suggested to function as a coping mechanism for grief following

a bereavement (Graham et al. 1987; Lister et al. 2008). A similar suggestion has been made

for beluga whales where both wild (Smith and Sleno 1986) and captive (Kilborn 1994),

individuals have been seen to carry inanimate objects, apparently as 'surrogates'. The captive

whale carried a buoy followed the the removal of her dead calf immediately after birth

(Kilborn 1994), and in the wild observations included carrying of planks and netting (Smith

and Sleno 1986). Further observations are necessary to validate the use of object carrying

following the death of an infant as a coping mechanism in primates,

To sum up, our observations are consistent with previous observations that chimpanzee

mothers respond to the death of their infants with carrying behaviour across communities.

Furthermore, our observations support the argument that these mothers act as if they are

aware of the loss but are continuing to display a strong attachment to the bodies of their

infants and may be affected by psychological processes akin to human grieving.

Nevertheless, more detailed hormonal data are needed for a test of this potential mechanism.

A combination of ecological conditions favouring mummification, and social factors, such as

the strong bond shared between mothers and their infants, may explain the three particularly

extended carries by Budongo chimpanzees. While we did not observe other indications of

maternal care in these cases, we are cautious about interpreting this as a wider absence in

Budongo mothers. Mothers' pattern of behavioural responses to death may be individually

specific and nuanced, resulting from a combination of physical, ecological, and psychological

factors, and more observations are needed to generalise at the population or species level.

Our interpretations are limited by the small number of observations and the multitude of

possible influential factors to consider. We encourage researchers and long-term field sites to

continue to report the rare behaviours observed in different populations, for example by

contributing to open-access databases such as 'ThanatoBase'

(http://thanatobase.mystrikingly.com), to allow a richer exploration and more robust

hypothesis testing of non-human primates' reaction to death through data sharing and

collaborations across sites.

ACKNOWLEDGMENTS

We are grateful to all field assistants of the Sonso and Waibira communities who provided

essential assistance throughout the observation period. We are thankful to the management,

staff, and researchers of the Budongo Conservation Field Station for their support and thank

the project's founder Vernon Reynolds. We thank the Royal Zoological Society of Scotland

for their long-term financial support to the field station. We thank the Uganda Wildlife

Authority and the Uganda National Council for Science and Technology for permission to

work in Uganda. We also thank André Gonçalves and Elisa Fernández-Fueyo for helpful

comments on a previous version of the draft. AS's fieldwork was supported by the

European Research Council project grant to CC (grant agreement number: 679787). CH is

supported by funding from the European Union's 8th Framework Programme, Horizon 2020

(grant agreement number: 802719).

REFERENCES

Anderson JR (2011) A primatological perspective on death. Am J Primatol 73:410–414.

https://doi.org/10.1002/ajp.20922

Anderson JR (2018) Chimpanzees and death. Phil Trans R Soc B 373:20170257.

https://doi.org/10.1098/rstb.2017.0257

Anderson JR (2020) Responses to death and dying: primates and other mammals. Primates

61:1–7. https://doi.org/10.1007/s10329-019-00786-1

Anderson JR, Biro D, Pettitt P (2018) Evolutionary thanatology. Phil Trans R Soc B

373:20170262. https://doi.org/10.1098/rstb.2017.0262

Badihi G, Bodden K, Zuberbühler K, et al (2021) Flexibility in the social structure of male

chimpanzees (Pan troglodytes schweinfurthii) in the Budongo Forest, Uganda (preprint).

https://doi.org/10.1101/2021.12.11.472209

Bercovitch FB (2020) A comparative perspective on the evolution of mammalian reactions to

dead conspecifics. Primates 61:21–28. https://doi.org/10.1007/s10329-019-00722-3

Biro D (2011) Chimpanzee mothers carry the mummified remains of their dead infants: three

case reports from Bossou. In: Matsuzawa T, Humle T, Sugiyama Y (eds) The

chimpanzees of Bossou and Nimba. Springer Japan, Tokyo, pp 241–250

Biro D, Humle T, Koops K, et al (2010) Chimpanzee mothers at Bossou, Guinea carry the

mummified remains of their dead infants. Curr Biol 20:R351–R352.

https://doi.org/10.1016/j.cub.2010.02.031

Botting J, van de Waal E (2020) Reactions to infant death by wild vervet monkeys

(Chlorocebus pygerythrus) in KwaZulu-Natal, South Africa: prolonged carrying, non-

mother carrying, and partial maternal cannibalism. Primates 61:751–756.

https://doi.org/10.1007/s10329-020-00851-0

Carter AJ, Baniel A, Cowlishaw G, Huchard E (2020) Baboon thanatology: responses of

filial and non-filial group members to infants' corpses. R Soc open sci 7:192206.

https://doi.org/10.1098/rsos.192206

Courtenay J, Santow G (1989) Mortality of wild and captive chimpanzees. Folia Primatol

52:167–177

Cronin KA, van Leeuwen EJC, Mulenga IC, Bodamer MD (2011) Behavioral response of a

chimpanzee mother toward her dead infant. Am J Primatol 73:415–421.

https://doi.org/10.1002/ajp.20927

Das S, Erinjery JJ, Desai N, et al (2019) Deceased-infant carrying in nonhuman anthropoids:

Insights from systematic analysis and case studies of bonnet macaques (Macaca radiata)

and lion-tailed macaques (Macaca silenus). J Comp Psych 133:156–170.

https://doi.org/10.1037/com0000140

Dirks PH, Berger LR, Roberts EM, et al (2015) Geological and taphonomic context for the

new hominin species Homo naledi from the Dinaledi Chamber, South Africa. Elife

4:e09561

Eggeling WJ (1947) Observations on the ecology of the Budongo rain forest, Uganda. J Ecol

34:20–87. https://doi.org/10.2307/2256760

Engh AL, Beehner JC, Bergman TJ, et al (2006a) Behavioural and hormonal responses to

predation in female chacma baboons (Papio hamadryas ursinus). Proc R Soc B 273:707–

712. https://doi.org/10.1098/rspb.2005.3378

Engh AL, Beehner JC, Bergman TJ, et al (2006b) Female hierarchy instability, male

immigration and infanticide increase glucocorticoid levels in female chacma baboons.

Anim Behav 71:1227–1237. https://doi.org/10.1016/j.anbehav.2005.11.009

Fashing PJ, Nguyen N, Barry TS, et al (2011) Death among geladas (Theropithecus gelada):

a broader perspective on mummified infants and primate thanatology. Am J Primatol

73:405–409. https://doi.org/10.1002/ajp.20902

Fedurek P, Tkaczynski P, Asiimwe C, et al (2020) Maternal cannibalism in two populations

of wild chimpanzees. Primates 61:181–187. https://doi.org/10.1007/s10329-019-00765-6

Fernández-Fueyo E, Sugiyama Y, Matsui T, Carter AJ (2021) Why do some primate mothers

carry their infant's corpse? A cross-species comparative study. Proc R Soc B 288:1–10.

https://doi.org/10.1098/rspb.2021.0590

Figley CR, Bride BE, Mazza N (1997) Death and trauma: The traumatology of grieving.

Taylor & Francis, Philadelphia

Fowler A, Hohmann G (2010) Cannibalism in wild bonobos (Pan paniscus) at Lui Kotale.

Am J Primatol 72:509–514. https://doi.org/10.1002/ajp.20802

Georgiev AV, Melvin ZE, Warketin A-S, et al (2019) Two cases of dead-infant carrying by

female zanzibar red colobus (Piliocolobus kirkii) at Jozani-Chwaka Bay National Park,

Zanzibar. Afr Primates 13:57–60

Girard-Buttoz C, Tkaczynski PJ, Samuni L, et al (2021) Early maternal loss leads to short-

but not long-term effects on diurnal cortisol slopes in wild chimpanzees. Elife 10:e64134.

https://doi.org/10.7554/eLife.64134

Gonçalves A, Biro D (2018) Comparative thanatology, an integrative approach: exploring

sensory/cognitive aspects of death recognition in vertebrates and invertebrates. Phil Trans

R Soc B 373:20170263. https://doi.org/10.1098/rstb.2017.0263

Gonçalves A, Carvalho S (2019) Death among primates: a critical review of non-human

primate interactions towards their dead and dying. Biol Rev 94:1502–1529.

https://doi.org/10.1111/brv.12512

Graham MA, Thompson SC, Estrada M, Yonekura ML (1987) Factors affecting

psychological adjustment to a fetal death. Am J Obstet Gynecol 157:254–257.

https://doi.org/10.1016/S0002-9378(87)80145-X

Gruber T (2016) Great apes do not learn novel tool use easily: conservatism, functional

fixedness, or cultural influence? Int J Primatol 37:296–316.

https://doi.org/10.1007/s10764-016-9902-4

Gruber T, Muller MN, Reynolds V, et al (2011) Community-specific evaluation of tool

affordances in wild chimpanzees. Sci Rep 1:128. https://doi.org/10.1038/srep00128

Gruber T, Muller MN, Strimling P, et al (2009) Wild chimpanzees rely on cultural

knowledge to solve an experimental honey acquisition task. Curr Biol 19:1806–1810.

https://doi.org/10.1016/j.cub.2009.08.060

Hanamura S, Kooriyama T, Hosaka K (2015) Diseases and deaths: variety and impact on

social life. In: Hosaka K, Zamma K, Nakamura M, Itoh N (eds) Mahale Chimpanzees: 50

Years of Research. Cambridge University Press, Cambridge, pp 354–371

Hill K, Boesch C, Goodall J, et al (2001) Mortality rates among wild chimpanzees. J Hum

Evol 40:437–450. https://doi.org/10.1006/jhev.2001.0469

Hollan D (1995) To the afterworld and back: mourning and dreams of the dead among the

Toraja. Ethos 23:424–436. https://doi.org/10.1525/eth.1995.23.4.02a00030

Hosaka K, Matsumoto-Oda A, Huffman MA, Kawanaka K (2000) Reactions to dead bodies

of conspecifics by wild chimpanzees in the Mahale Mountains, Tanzania. Primate Res

16:1–15 (in Japanese)

Hrdy SB (1999) Mother nature: ruthless competitive nurturing and tender: maternal instincts

& the shaping of the species. Chatto and Windus, London

Kahlenberg SM, Wrangham RW (2010) Sex differences in chimpanzees' use of sticks as play

objects resemble those of children. Curr Biol 20:R1067–R1068.

https://doi.org/10.1016/j.cub.2010.11.024

Kaplan J (1973) Responses of mother squirrel monkeys to dead infants. Primates 14:89–91.

https://doi.org/10.1007/BF01730518

Kappeler PM, van Schaik CP, Watts DP (2012) The values and challenges of long-term field

studies. In: Kappeler PM, Watts DP (eds) Long-term Field studies of primates. Springer

Berlin Heidelberg, Berlin, pp 3–18

Keverne EB (1988) Central mechanisms underlying the neural and neuroendocrine

determinants of maternal behaviour. Psychoneuroendocrinology 13:127–141.

https://doi.org/10.1016/0306-4530(88)90010-8

Kilborn SS (1994) Object carrying in a captive beluga whale (Delphinapterus leucas) as

possible surrogate behavior. Mar Mamm Sci 10:496–501. https://doi.org/10.1111/j.1748-

7692.1994.tb00510.x

Kooriyama T (2009) The death of a newborn chimpanzee at Mahale: reactions of its mother

and other individuals to the body. Pan Afr News 16:19–21

Lannen PK, Wolfe J, Prigerson HG, et al (2008) Unresolved grief in a national sample of

bereaved parents: impaired mental and physical health 4 to 9 years later. J Clin Oncol

26:5870–5876. https://doi.org/10.1200/JCO.2007.14.6738

Leroux M, Monday G, Chandia B, et al (2021) First observation of a chimpanzee with

albinism in the wild: Social interactions and subsequent infanticide. Am J Primatol.

https://doi.org/10.1002/ajp.23305

Lister S, Pushkar D, Connolly K (2008) Current bereavement theory: Implications for art

therapy practice. Arts Psychother 35:245–250. https://doi.org/10.1016/j.aip.2008.06.006

Lonsdorf EV, Ross SR (2012) Socialization and development of behavior. In: The evolution

of primate societies. University of Chicago Press, Chicago, pp 245–268

Lonsdorf EV, Wilson ML, Boehm E, et al (2020) Why chimpanzees carry dead infants: an

empirical assessment of existing hypotheses. R Soc open sci 7:200931.

https://doi.org/10.1098/rsos.200931

Lowe AE, Hobaiter C, Asiimwe C, et al (2020) Intra-community infanticide in wild, eastern

chimpanzees: a 24-year review. Primates 61:69–82. https://doi.org/10.1007/s10329-019-

00730-3

Lowe AE, Hobaiter C, Newton-Fisher NE (2019) Countering infanticide: Chimpanzee

mothers are sensitive to the relative risks posed by males on differing rank trajectories.

Am J Phys Anthropol 168:3–9. https://doi.org/10.1002/ajpa.23723

MacKinnon KC, Riley EP, Garber PA, et al (2014) Code of Best Practices for Field

Primatology. https://doi.org/10.13140/2.1.2889.1847

Martinón-Torres M, d'Errico F, Santos E, et al (2021) Earliest known human burial in Africa.

Nature 593:95–100. https://doi.org/10.1038/s41586-021-03457-8

Masi S (2020) Reaction to allospecific death and to an unanimated gorilla infant in wild

western gorillas: insights into death recognition and prolonged maternal carrying.

Primates 61:83–92. https://doi.org/10.1007/s10329-019-00745-w

Matsuzawa T (1997) The death of an infant chimpanzee at Bossou, Guinea. Pan Afr News

4:4–6. https://doi.org/10.5134/143350

Nakamichi M, Koyama N, Jolly A (1996) Maternal responses to dead and dying infants in

wild troops of ring-tailed lemurs at the Berenty Reserve, Madagascar. Int J Primatol

17:505–523. https://doi.org/10.1007/BF02735189

Newton-Fisher NE (1999) The diet of chimpanzees in the Budongo Forest Reserve, Uganda.

Afr J Ecol 37:344–354. https://doi.org/10.1046/j.1365-2028.1999.00186.x

Nicolson NA (1991) Maternal behavior in human and nonhuman primates. In: Loy JD, and

Peters, CB (eds) Understanding behavior: What primate studies tell us about human

behavior. Oxford Univ. Press, New York, pp 17–50

Notman H, Munn J (2003) A case of infant carrying by an adult male chimpanzee in the

Budongo forest. Pan Afr News 10:7–9

Parkes CM, Langani P, Young B, Speck PW (1997) Death and bereavement across cultures.

Palliat Med 11:427–427

Pettitt P (2018) Hominin evolutionary thanatology from the mortuary to funerary realm: the

palaeoanthropological bridge between chemistry and culture. Phil Trans R Soc B

373:20180212. https://doi.org/10.1098/rstb.2018.0212

Pusey AE (1983) Mother-offspring relationships in chimpanzees after weaning. Anim Behav

31:363–377. https://doi.org/10.1016/S0003-3472(83)80055-4

Ramsay MS, Teichroeb JA (2019) Anecdotes in primatology: temporal trends,

anthropocentrism, and hierarchies of knowledge. Am Anthropol 121:680–693.

https://doi.org/10.1111/aman.13295

Reggente MAL, Alves F, Nicolau C, et al (2016) Nurturant behavior toward dead

conspecifics in free-ranging mammals: new records for odontocetes and a general

review. J Mammal 97:1428–1434. https://doi.org/10.1093/jmammal/gyw089

Rendu W, Beauval C, Crevecoeur I, et al (2014) Evidence supporting an intentional

Neandertal burial at La Chapelle-aux-Saints. Proc Natl Acad Sci USA 111:81–86.

https://doi.org/10.1073/pnas.1316780110

Reynolds V (2005) The chimpanzees of the Budongo forest: Ecology, behaviour and

conservation. Oxford Univ. Press, Oxford

Rutz C, Webster MM (2021) *Ethology* adopts the STRANGE framework for animal

behaviour research, to improve reporting standards. Ethology 127:99–101.

https://doi.org/10.1111/eth.13118

Samuni L, Mundry R, Terkel J, et al (2014) Socially learned habituation to human observers

in wild chimpanzees. Anim Cogn 17:997–1005. https://doi.org/10.1007/s10071-014-

0731-6

Smith TG, Sleno GA (1986) Do white whales, Delphinapterus leucas, carry surrogates in

response to early loss of their young? Can J Zool 64:1581–1582.

https://doi.org/10.1139/z86-237

Stanton MA, Lonsdorf EV, Pusey AE, Murray CM (2017) Do juveniles help or hinder?

Influence of juvenile offspring on maternal behavior and reproductive outcomes in wild

chimpanzees (Pan troglodytes). J Hum Evol 111:152–162.

https://doi.org/10.1016/j.jhevol.2017.07.012

Sugiyama Y, Kurita H, Matsui T, et al (2009) Carrying of dead infants by Japanese macaque

(Macaca fuscata) mothers. Anthropol Sci 117:113–119.

https://doi.org/10.1537/ase.080919

Takeshita RSC, Huffman MA, Kinoshita K, Bercovitch FB (2020) Changes in social

behavior and fecal glucocorticoids in a Japanese macaque (Macaca fuscata) carrying her

dead infant. Primates 61:35–40. https://doi.org/10.1007/s10329-019-00753-w

Thompson CL, Hrit R, Melo LCO, et al (2020) Callitrichid responses to dead and dying

infants: the effects of paternal bonding and cause of death. Primates 61:707–716.

https://doi.org/10.1007/s10329-020-00824-3

Tokuyama N, Moore DL, Graham KE, et al (2017) Cases of maternal cannibalism in wild

bonobos (Pan paniscus) from two different field sites, Wamba and Kokolopori,

Democratic Republic of the Congo. Primates 58:7–12. https://doi.org/10.1007/s10329-

016-0582-7

Warren Y, Williamson EA (2004) Transport of dead infant mountain gorillas by mothers and

unrelated females. Zoo Biol 23:375–378. https://doi.org/10.1002/zoo.20001

Watson CFI, Matsuzawa T (2018) Behaviour of nonhuman primate mothers toward their

dead infants: uncovering mechanisms. Phil Trans R Soc B 373:20170261.

https://doi.org/10.1098/rstb.2017.0261

Webster MM, Rutz C (2020) How STRANGE are your study animals? Nature 582:337–340.

https://doi.org/10.1038/d41586-020-01751-5

Whiten A, Goodall J, McGrew WC, et al (1999) Cultures in chimpanzees. Nature 399:682–

685. https://doi.org/10.1038/21415

Wilson ML, Boesch C, Fruth B, et al (2014) Lethal aggression in Pan is better explained by

adaptive strategies than human impacts. Nature 513:414–417.

https://doi.org/10.1038/nature13727