Prenatal thyroid hormones accelerate postnatal growth and telomere shortening in wild great tits

Bin-Yan Hsu[1], Nina Cossin-Sevrin[1,2], Antoine Stier[1,3*], Suvi Ruuskanen[1,4*$]

[1] Department of Biology, University of Turku, Finland

[2] Université de Strasbourg, Centre National de la Recherche Scientifique, Institut Pluridisciplinaire Hubert Curien, UMR 7178, 67087 Strasbourg, France

[3] Univ Lyon, Université Claude Bernard Lyon 1, CNRS, ENTPE, UMR 5023 LEHNA, F-69622, Villeurbanne, France

[4] Department of Biological and Environmental Science, University of Jyväskylä

*joint senior authors

[$]*Corresponding author:* suvi.k.ruuskanen@gmail.com

Running headline: Prenatal thyroid hormones shorten telomeres

Abstract

 Early-life environment is known to affect later-life health and disease, which could be mediated by the early-life programming of telomere length, a key hallmark of ageing. According to the *fetal programming of telomere biology hypothesis*, variation in prenatal exposure to hormones is likely to influence telomere length. Yet the contribution of key metabolic hormones, *i.e.* thyroid hormones (THs), has been largely ignored. We recently showed that in contrast to predictions, exposure to elevated prenatal THs increased postnatal telomere length in wild collared flycatchers, but the generality of such effect, its underlying proximate mechanisms and consequences on survival have not been investigated. We therefore conducted a comprehensive study evaluating the impact of THs on potential drivers of telomere dynamics (growth, post-natal THs, mitochondria and oxidative stress), telomere length and medium-term survival using wild great tits as a model system. While prenatal THs did not significantly affect telomere length a week after hatching (*i.e.* day 7), they influenced postnatal telomere shortening (*i.e.* shorter telomeres at day 14 and the following winter) but not apparent survival. Circulating THs, mitochondrial density or oxidative stress biomarkers were not significantly influenced, whereas TH-supplemented group showed accelerated growth, which may explain the observed delayed effect on telomeres. We discuss several alternative hypotheses that may explain the contrast with our previous findings in flycatchers. Given that shorter telomeres in early life tend to be carried until adulthood and are often associated with decreased survival prospects, the effects of prenatal THs on telomeres may have long-lasting effects on senescence.

Keywords: developmental programming, thyroid hormone, telomere, ageing, mitochondria, metabolism, oxidative stress, maternal effects

Introduction

Early-life environment has been repeatedly observed to affect adult health and survival prospects in human and non-human vertebrates (Barnes and Ozanne, 2011; Godfrey and Barker, 2001; Metcalfe and Monaghan, 2001). While the mechanisms underlying such delayed effects remained somewhat elusive (Barnes and Ozanne, 2011), the early-life programming of telomere length (i.e. the protective end caps of chromosomes) has emerged as a key candidate (Entringer et al., 2018). Telomere length is considered as a hallmark of ageing (López-Otín et al., 2013) since telomeres shorten with age, and shorter telomeres are often predictive of lower survival or lifespan in both epidemiological and experimental studies (Arbeev et al., 2020; Heidinger et al., 2012; Muñoz-Lorente et al., 2019; Wilbourn et al., 2018). The prenatal hormonal environment, such as exposure to elevated glucocorticoid levels, has been coined as an important factor influencing early-life telomere length and its associated long-term outcomes (Criscuolo et al., 2020; Haussmann et al., 2012; Marchetto et al., 2016; Parolini et al., 2019). While there has been a considerable interest in prenatal glucocorticoids (Haussmann et al. 2012; Tissier, Williams, and Criscuolo 2014; Noguera, da Silva, and Velando 2020) and to a lesser extent androgens (Parolini et al., 2019; Tissier et al., 2014) in the context of the '*fetal programming of telomere biology*' hypothesis (Entringer et al., 2018), the potential impact of prenatal thyroid hormones has been mostly ignored so far (Stier et al. 2020).

Thyroid hormones (THs) are key coordinators of development and metabolism (McNabb, 2007), which are transferred from mothers to offspring (Ruuskanen and Hsu, 2018). Variation in exposure to prenatal thyroid hormones (T3, triiodothyronine, and T4, thyroxine) could influence telomere length via several mutually non-exclusive proximate pathways: (i) Prenatal THs can stimulate growth (yet results are inconsistent across studies and species,

(Hsu et al., 2020, 2019a, 2017; Medici et al., 2013; Ruuskanen et al., 2016a; Sarraude et al.,

2020a; Sarraude et al., 2020b; Vrijkotte et al., 2017; Zhang et al., 2019), which can directly

contribute to telomere attrition through increasing cellular division (Monaghan and Ozanne,

2018; Stier et al., 2020), or indirectly accelerate telomere shortening through increasing

oxidative stress (Monaghan and Ozanne, 2018; Reichert and Stier, 2017; Smith et al., 2016).

(ii) Elevated TH levels are often associated with higher metabolic rates (Liu et al., 2006; Mullur

et al., 2014; Welcker et al., 2013) and stimulate mitochondrial aerobic metabolism (Cioffi et

al., 2013), both of which can potentially increase reactive oxygen species (ROS) and oxidative

damage (Stier, Massemin, and Criscuolo 2014), accelerating telomere shortening (Reichert

and Stier, 2017). It was recently shown that exposure to elevated prenatal TH levels can lead

to a sex-specific increase in metabolic rate and circulating TH levels shortly after hatching (rock

pigeons *Columba livia*, Hsu et al. 2017, but see Ruuskanen et al. 2016), which suggests that

prenatal hormones may program postnatal metabolism and TH-axis function. (iii) The

'metabolic telomere attrition hypothesis' (Casagrande and Hau, 2019) postulates that

telomere shortening might be adaptive by amplifying the cellular signaling of energy debt to

re-direct critical resources to immediately important processes. In this scenario, we would

expect accentuated shortening when catabolism is favored over anabolism via mTOR

inhibition (i.e. mechanistic target of rapamycin, a key regulator of cellular homeostasis)

leading to a decrease in telomere maintenance processes and an active shortening through

exonuclease action (Casagrande and Hau, 2019). Since THs can have both anabolic and

catabolic actions (Mullur et al., 2014), predictions can be made in both directions between

prenatal THs and telomere length.

From an evolutionary perspective, increased offspring metabolism or growth, diverts

resources from somatic maintenance if resources are limited. This can accelerate damage to

biomolecules and/or decrease repair/maintenance processes, and therefore accentuate

telomere shortening. Therefore, prenatal TH levels would be expected to vary in relation to

predicted environmental conditions (as observed in terms of temperature and laying order,

e.g. Ruuskanen, Groothuis, et al. 2016; Hsu, Verhagen, et al. 2019) to optimize this trade-off.

In contrast to most predictions relating prenatal THs to telomere length (see above),

we recently reported that prenatal exposure to experimentally elevated THs increased

telomere length in nestlings of a wild passerine, the collared flycatcher (*Ficedula albicollis,*

Stier et al. 2020). To better understand the potential generality of this surprising finding as

well as assess underlying mechanisms and potential carry-over effects of variation in prenatal

THs on later life-stages and survival, we conducted a more detailed study in another passerine

species, the great tit *(Parus major)*. The aim of this study was to comprehensively investigate

the influence of prenatal THs on growth, oxidative stress, plasma THs, mitochondrial density

(i.e. mitochondrial DNA copy number) and telomere dynamics as well as survival via an

experimental manipulation of prenatal THs in a wild population. We monitored offspring

multiple times during development and as juveniles a few months after fledging. Based on the

majority of prior literature, we would predict that elevated prenatal THs could lead to faster

growth, increased plasma THs, oxidative stress and mitochondrial density (higher density can

lead to higher ROS production), ultimately accelerating telomere shortening. Alternatively, if

our previous finding in the collared flycatcher reflected a general pattern, we would predict

that despite accelerating growth, elevated prenatal THs could increase telomere length. We

also predict that elevated prenatal TH could increase post-fledging survival of the juveniles

(*e.g.* both due to accelerated postnatal growth and potential beneficial effects for

thermoregulation under low autumn-winter temperatures). However, longer-term survival,

that we were not able to evaluate here, could be decreased for example due to shorter

telomeres.

Methods

The experiment and all methods we used were in accordance with all relevant guidelines and

regulations and have been approved by the Animal Experiment Board of the Administrative

Agency of South Finland (ESAVI/2902/2018) and the Environmental Center of Southwestern

Finland (license number VARELY549/2018). The experiment was conducted in 2018 in a nest

box population (314 nest-boxes distributed over seven forest plots) on the island of Ruissalo

in southwestern Finland (60° 25' N, 22° 10' E).

Field experiment

The nest boxes were monitored with five-day intervals to track egg laying. Yolk T3 and T4 (i.e.

a combined injection of the two hormones) levels were elevated in half of the nests (n = 21

TH nests and n = 21 control nests) by injection into the egg, following methods in Ruuskanen,

Darras, et al. (2016). Control nests were injected with the vehicle (0.9% NaCl) only. The TH

content (±SD) in great tit eggs in the population is 0.053 (±0.02) ng/yolk for T3 and 0.46 (± 0.16

ng/yolk) for T4 (Ruuskanen et al. unpubl data). We aimed to raise the amount of yolk TH by

2SD via injection into the egg yolk, a dose that has been recommended in relation to the

natural hormone range of the study species (Podmokla et al., 2018). This corresponded to

target doses of 0.041 ng/yolk for T3 and 0.325 ng/yolk for T4. To make sure injections would

be performed on unincubated eggs mimicking maternal hormone levels (great tits can start

incubation before the clutch is complete), injections were conducted on the day the 5th egg

was laid to all eggs in the clutch. Thereafter, injections were conducted each day for the newly

laid egg.

Hatching date and success was monitored by visiting nests daily starting before the

estimated hatching day. Nestling body mass (~0.01g) and tarsus (~0.5 mm) were measured on

day 2 (mass only), 7 and 14 post-hatching. All nestling measurements were conducted

between 8 am and 2 pm. After measurements on day 2, nestlings were nail-coded, the brood

was split and nestlings were cross-fostered with same-age nestlings in another nest of the

opposite hormone treatment. This "split-brood design" allows chicks from TH and control-

injected eggs to be raised under the same postnatal environment and is in theory more

sensitive to detect effects that may otherwise require a much larger number of nests in an

experiment of "between-brood" design (e.g. Tschirren et al. 2005). Preferentially, half of the

brood was swapped between the nests with opposite treatments whenever possible, but we

also split a brood for cross-fostering when we did not have an equal number of nests with

opposite treatments on a given day. The proportion of TH-nestlings (i.e. number of TH-

nestlings divided by the brood size) in each nest after cross-fostering was therefore recorded.

All nestlings, cross-fostered or not, were included in the statistical analyses. There was no clear

bias on the number of nestlings that were cross-fostered or stayed in the original nests with

respect to the hormone treatment (CO-nestlings: 58 stayed and 41 moved, TH-nestlings: 68

stayed and 51 moved, χ^2 = 0.006, p = 0.939). On day 7 a blood sample (ca 40µl) was taken and

snap-frozen in liquid nitrogen for molecular analyses (telomere length, mitochondrial density

and molecular sexing). On day 14 a small blood sample (10-15µl) was snap-frozen in liquid

nitrogen for oxidative stress biomarker analyses and stored at -80ºC, while the rest of the

sample (ca 40 µl) was kept on ice, and centrifuged. RBCs (15 µl) were used for mitochondria

and telomere measurements as above and plasma (15 µl) for TH analyses. After day 7

measurements half of the nest were subject to a temperature manipulation (nest temperature increased for *ca.* 2^{o}C during day 7 – day 14) for the purposes of another study. To avoid potential confounding effects, for data after day 7 and in juveniles, we only included the individuals from the nests which were not temperature-manipulated (N = 20 nests, 111 nestlings on day 7 and 99 nestlings on day 14). Yet, for day 2 and day 7 measurements (prior to the temperature manipulation), we prefer to include all data (N = 42 nests, in total 224 nestlings, 218 nestlings on d2 and 221 nestlings on d7) to make use of the larger sample size.

To study long-term effects of prenatal thyroid hormones and offspring apparent post-fledging survival, we recaptured great tits during the following autumn-winter using mist-netting. Seven feeding stations were mounted in the study plots in August (generally any given nest box had a station within 200m), and nets were positioned close to the feeding stations. The distances between adjacent feeding stations were ca. 500 m, and most birds were captured at a different feeding station than their natal forest plot. In addition, 5 birds were captured at a constant effort site > 3 km from the study plots, suggesting that all stations were potentially accessible to all birds. Circa 20 kg of peeled sunflower seeds and 2 kg of fat were provided at each station, checked and filled bi-weekly, and consumption noted. Only in a very few cases the feeders were completely empty. Mist-netting was conducted at each feeding station for 3 hours on three different days in September-October and similarly again in February, summing up to a total 126 hours of mist-netting. The time of day (morning/day/afternoon) was rotated for each site. Nets were checked every 30 minutes and mass (~0.01g), wing (~0.5 mm) were recorded for each bird. A small blood sample (40-60μl) was taken, kept on ice, centrifuged within 8 hours, and RBCs frozen at -80ºC for telomere and mitochondria density analyses.

 Thyroid hormones and oxidative stress assays

Plasma thyroid hormones, the biologically active form triiodothyronine (T3), and a

precursor thyroxine (T4) (expressed as ng/mL) were measured from 14-day-old nestlings with

nano-LC-MS/MS following Ruuskanen et al. (2019; 2018). Due to practical constraints (time,

effort and finances), we randomly selected one to four nestling per nest, n = 13 TH and n = 11

control nestlings (from 18 nests of origin and 11 nests of rearing) for analysis.

Total glutathione (tGSH), the most abundant intra-cellular antioxidant, and the ratio

between reduced and oxidized gluthatione (GSH:GSSG, an indicator of oxidative challenge)

were measured from whole-blood samples (12µl of 0.9% NaCl-diluted whole blood) with the

ThioStar® Glutathione Fluorescent Detection Kit (K005-FI, Arbor Assays, USA) (Sarraude et al.,

2020a). As a measure of oxidative damage, we assessed blood lipid peroxidation

(malonaldehyde, MDA, 50µl of 0.9% NaCl-diluted whole blood) using the TBARS-assay

following Espin et al. (2017). Both measurements had CV% < 10 and are expressed per mg of

protein (measured via BCA protein assay, ThermoFisher Scientific), following Espin et al. 2017.

We aimed to measure biomarkers of oxidative stress from 2 control or 2 TH-nestlings per nest,

yet sometimes fewer nestlings were available and some samples failed in the laboratory

analysis, yielding the final sample size as n = 33 TH and 26 control nestlings from 18 rearing

nests.

qPCR assays for relative telomere length, mtDNA copy number and molecular sexing

We extracted DNA from blood cells using a standard salt extraction alcohol

precipitation method (Aljanabi and Martinez, 1997). Extracted DNA was diluted in elution

buffer BE for DNA preservation. DNA concentration and quality (260/280 > 1.80 and 260/230

> 2.00) were checked with a ND-1000-Spectrophotometer (NanoDrop Technologies,

Wilmington, USA). DNA integrity was verified in 24 samples chosen randomly using gel

electrophoresis (50 ng of DNA, 0.8 % agarose gel at 100 mV for 60 min) and DNA staining with

Midori Green. Each sample was then diluted to a concentration of 1.2 ng.μl^{-1} for subsequent

qPCR analysis.

Relative telomere length (*rTL*) and mitochondrial DNA copy number (*mtDNAcn*, an

index of mitochondrial density) were quantified using qPCR. This technique estimates relative

telomere length by determining the ratio (T/S) of telomere repeat copy number (T) to a single

copy gene (SCG), and the relative mtDNAcn as the ratio between one mitochondrial gene and the

same single copy gene. Here, we used RAG1 as a SCG (verified as single copy using a BLAST

analysis on the great tit genome) and cytochrome oxidase subunit 2 (COI2) as a mitochondrial

gene (verified as non-duplicated in the nuclear genome using a BLAST analysis). Forward and

reverse telomere primers were 5'-CGGTTTGTTTGGGTTTGGGTTTGGGTTTGGGTTTGGGTT-3'

(Tel-1b) and 5'-GGCTTGCCTTACCCTTACCCTTACCCTTACCCTTACCCT-3' (Tel-2b) respectively.

Forward and reverse RAG1 primers were 5'-TCGGCTAAACAGAGGTGTAAA-3' and 5'-

CAGCTTGGTGCTGAGATGTAT-3', respectively. Forward and reverse COI2 primers were 5'-

CAAAGATATCGGCACCCTCTAC-3'; 5'- GCCTAGTTCTGCACGGATAAG-3', respectively. For the

qPCR assays, the reactions were performed on a 384-QuantStudio™ 12K Flex Real-Time PCR

System (Thermo Fisher), in a total volume of 12μL including 6ng of DNA, primers at a final

concentration of 300nM and 6μL of SensiFAST™ SYBR lo-ROX (Bioline). Telomere, RAG1 and

COI2 reactions were performed in triplicates on the same plates (10 plates in total); the qPCR

conditions were: 3min at 95°C, followed by 35 cycles of 10 s at 95°C, 15 s at 58°C and 10s at

72°C. A DNA sample being a pool of DNA from 10 adult individuals was used as a reference

sample and was included in triplicate on every plate. The efficiency of each amplicon was

estimated from a standard curve of the reference sample ranging from 1.5 to 24ng. The mean

reaction efficiencies were 109.1 ± 1.8% for telomere, 102.2 ± 1.6% for RAG1, 96.3 ± 1.1% for

COI2. The relative telomere length and mtDNAcn of each sample were calculated as $(1+Ef_{\text{Tel or}}$

$_{\text{COI2}})^{\Delta Cq\,\text{Tel or COI2}}/(1+Ef_{\text{RAG1}})^{\Delta CqRAG1}$; Ef being the amplicon efficiency, and ΔCq the difference in

Cq-values between the reference sample and the focal sample. Intra-plate technical

repeatabilities of telomere and *mtDNAcn* based on triplicates were 0.87 (95% C.I. [0.85-0.89])

and 0.96 (95% C.I. [0.95-0.97]) respectively. Inter-plate technical repeatabilities of telomere

and *mtDNAcn* based on one repeated plate were 0.98 (95% C.I. [0.97-0.99]) and 0.77 (95% C.I.

[0.59-0.88]) respectively.

The use of *mtDNAcn* as an index of mitochondrial density has been questioned in human

(Larsen et al., 2012), but we have previously shown good correlations between mtDNAcn and

mitochondrial respiration rates in pied flycatcher (Stier et al. 2019). Great tits have quite peculiar

telomeres, characterized notably by some ultra-long telomeres that do not seem to shorten with

age in adults (Atema et al., 2019). Since qPCR only provides an estimate of overall telomere

length, it has been suggested it could be suboptimal for this study species. Yet, relative telomere

length (*i.e.* measured using qPCR) in this species has been shown to shorten during the nestling

stage (Stier et al., 2016, 2015), to respond to environmental factors (*e.g.* hatching asynchrony:

(Salmón et al., 2016; Stier et al., 2016, 2015) and to predict adult survival (Salmón et al., 2017).

Within-individual repeatability of telomere length has recently been suggested to be an important

factor to evaluate the pertinence of telomere length data in a given study/species (Kärkkäinen et

al., n.d.), and the biological repeatability in our dataset was R = 0.66 [0.55-0.79], which is above

the average reported by qPCR studies (*i.e. R* = 0.47), and within the upper range of what has been

reported for great tits (Kärkkäinen et al. 2021)

Birds were molecularly sexed using a qPCR approach adapted from(Chang et al., 2008;

Ellegren　and　Fridolfsson,　1997).　Forward　and　reverse　sexing　primers　were　5'-

CACTACAGGGAAAACTGTAC-3' (2987F) and 5'- CCCCTTCAGGTTCTTTAAAA -3' (3112R),

respectively. qPCR reactions were performed in a total volume of 12μL including 6ng of DNA,

primers at a final concentration of 800nM and 6μL of SensiFAST™ SYBR® Lo-ROX Kit (Bioline).

qPCR conditions were: 3 min at 95°C, followed by 40 cycles of 45 s at 95°C, 60 s at 52°C and

60s at 72°C, then followed by a melting curve analysis (95°C 60s, 45°C 50s, increase to 95°C at

0.1°C/s, 95°C 30s). Samples were run in duplicates in a single plate and 6 adults of known sex

were included as positive controls. Sex was determined by looking at the dissociation curve,

with two peaks indicating the presence of a Z and W chromosome (female), and one peak

indicating the presence of only the Z chromosomes (male).

Statistical analyses

We ran several linear or generalized linear mixed models (LMMs/GLMMs) for both

nestling and juvenile data. In order to account for genetic effects and the effects of growing

environment other than prenatal hormones, we included the IDs of both the nest of origin

(i.e. where a nestling hatched) and the nest of rearing (i.e. where a nestling grew up) as

random intercepts in all suitable models. For physiological measurements, assay batch ID was

included either as random intercept or fixed factors (depending on the number of levels, see

Tables for details). All initial models included the prenatal hormone treatment (TH vs. CO),

date (continuous variable), cross-fostering (yes/no), brood size (continuous variable),

proportion of TH-nestlings (continuous variable) in a nest, and their interactions with hormone

treatment as fixed factors. Except for hormone treatment, other variables were initially

included because of their potential confounding effects on nestling growth and development

but only retained in the final models when $p < 0.15$ in order not to miss potentially significant

factors. Nevertheless, in the final models, significance level was still set at 0.05. In models of

nestling growth (i.e. body mass and tarsus length), the ID of each individual was included as a

random intercept.

To analyze nestling growth, we separated the data into the early (i.e. from day 2 to day

7) and late (i.e. from day 7 to day 14) phases. Because of the start of heating treatment at day

7, this separation allows us to make use of the larger sample size during the early phase and

focus on the nestlings from non-heated nests during the late phase. In these models, age (as

a continuous variable) and hormone × interaction were included as additional fixed factors,

and individual ID as a random intercept. In order to account for the variation in growth rate

between nests, age (continuous variable) was also treated as a random slope to interact with

both the original and rearing nests. For tarsus length, we additionally controlled for the

measurer ID as a fixed factor. Because molecular sexing was only conducted on nestlings

having DNA extracted for mtDNAcn and rTL, not all nestlings included in the growth analyses

were sexed. We therefore did not include sex in order to make use of the full data set.

Repeating our models with only the sexed nestlings still gave qualitatively the same results

and did not show sex-specific effects of prenatal THs on growth. For the juveniles captured

during autumn/winter we ran LMMs with hormone treatment, date of capture, and sex

(because of clear sexual size dimorphism at this age) as fixed factors.

For all models of physiological measurements fixed factors included prenatal hormone

treatment, nestling sex and body mass. If not showing any trends (i.e. $p < 0.15$), sex and body

mass were then removed from the final models, given the smaller sample sizes for the

physiological measurements. Natural log transformation was conducted on highly-skewed

data: mtDNAcn and three oxidative stress biomarkers to ensure normal residual distributions.

rTL and plasma THs were untransformed as residuals qualified the assumption of normality.

Because both mtDNAcn and rTL are ratios to a single copy gene, we also z-transformed the

data to allow across-study comparison (Verhulst, 2020). The ages of all juveniles were pooled

as a single age category "autumn", and age was thus included as a three-level categorical

variable (day 7, day 14, and autumn) in order to estimate the changes of mtDNAcn and rTL

between each time point.

We used GLMMs to estimate the influence of prenatal thyroid hormones on hatching

success, fledging success (i.e. pre-fledging survival), and post-fledging survival. We modelled

the outcome of each egg (i.e. hatched or not) or nestling (i.e. survived or not) using a logit link

function and specifying a binomial residual distribution. For hatching success, two nests in

which eggs never hatched (i.e. cause of failure likely unrelated to our treatment) were

excluded, giving a final sample size of 354 eggs from 42 nests. The nest ID was treated as a

random intercept, and prenatal hormone treatment and laying date of each nest as fixed

factors. For fledging success, we only included the 131 nestlings (reared in 27 nests after cross-

fostering) that successfully hatched and were subsequently reared in the nests that were not

subject to temperature manipulation in the analysis. Among the 131 nestlings, 99 successfully

fledged and were included in the analysis of post-fledging survival. According to the nestling

ringing records since 1999, only 8 out of >6200 birds ringed on the Ruissalo island were

recaptured outside the island, suggesting a very low dispersal rate. The individuals that were

never recaptured in the autumn were therefore presumed dead. The ID of rearing and original

nests was treated as a random intercept and prenatal hormone treatment as the fixed factor.

Laying date, brood size, and cross-fostering, as well as their interactions with the hormone

treatment, were initially included but only retained in the final model when p < 0.15.

In all statistical models described above, hormone-related two-way interactions were

of interest and therefore included. Following the suggestion by (Schielzeth, 2010), input

variables (except for the technical variables) were mean-centered before model fitting.

Significant interactions were further examined by post-hoc interaction analysis using the R

package *emmeans* (Lenth 2021).

All LMMs and GLMMs were conducted in the R environment 3.6.1, using the package

lme4 (Bates et al., 2015) and *lmerTest* (Kuznetsova et al., 2017). P values were determined

using the Kenward-Roger method to approximate the denominator degrees of freedom (R

package *pbkrtest*, (Halekoh and Hojsgaard, 2014), implemented within *lmerTest*) for LMMs

and by Laplace approximation for GLMMs. Model assumptions of normality and

homoscedasticity were diagnosed by visual inspection on simulated residuals using the R

package *DHARMa* (Hartig 2021). Clear violation was only observed in the models for plasma

T3 in which residuals were heteroscedastic. Nevertheless, given the fact that the model of T3

did not detect any significant effect and the general robustness of LMMs (Schielzeth et al.,

2020), this did not influence our conclusion.

Results

Early body mass growth (between day 2 and day 7 post-hatching) did not differ

between TH-supplemented and control groups (Table 1a, Fig 1a), but individuals from TH-

supplemented eggs grew faster at the later nestling stage, i.e. between day 7 and day 14 post-

hatching (Hormone × age interaction, $F_{1,20.18}$ = 5.449, p = 0.030, Table 1b, Fig 1b; body mass

gain (mean±SE) in TH-nestlings 0.899 ± 0.042 g/day, CO-nestlings 0.786±0.043 g/day).

Similarly, individuals from TH-supplemented eggs expressed marginally faster tarsus growth

rate between day 7 and day 14 post-hatching ($F_{1,31.06}$ = 3.737, p=0.062, Table 1c, TH:

0.567±0.033 mm/day, CO: 0.476±0.034 mm/day, Fig 1c). Cross-fostering was retained in the

final model of early body mass growth and suggested that fostered chicks had a lower growth

rate (-0.139±0.069 g/day) than non-fostered chicks (p = 0.046, Table 1a). When analyzing the

body mass and tarsus separately for each age, body mass was slightly smaller on day 2 and

day 7 in the TH-supplemented group compared to control, yet there was no statistically

significant difference between the groups at any age (Fig 1, Table S1). At day 7, fostered

nestlings tended to have a lower body mass by 0.347 g (SE = 0.179, p=0.053) on average, and

had significantly shorter tarsi by 0.33 mm (SE = 0.146, p=0.025), yet such differences

disappeared by day 14 (Table S1). In juveniles, there was no difference in body mass or wing

length between the treatment groups (Body mass: $F_{1,9.79}$ = 0.003, p = 0.959; wing length: $F_{1,10.34}$

= 0.011, p = 0.919).

Table 1. Linear mixed models of early and late nestling growth in response to prenatal TH-

supplementation.

(a) Early nestling body mass growth (day 2 – day 7)

Random effects:		Variance	Std. Dev.			
Individual ID (n=224)	Intercept	0.839	0.916			
Nest of origin (n=42)	Intercept	0.627	0.792			
	Age	0.005	0.070			
Nest of rearing (n=42)	Intercept	0.061	0.248			
	Age	0.007	0.083			
Residual		0.202	0.450			

Fixed factors	Estimate	SE	t	df	F	p
Intercept	8.232	0.146	56.478			
Hormone (TH)	-0.182	0.142	-1.284	1, 38.29	1.650	0.207
Age	1.730	0.019	89.716	1, 45.64	8045.995	**<0.001**
Cross-foster (yes)	-0.139	0.069	-2.008	1, 188.30	4.031	**0.046**
Hormone × Age	0.009	0.016	0.560	1. 28.84	0.314	0.580

(b) Late nestling body mass growth (day 7 – day 14)

Random effects:		Variance	Std. Dev.			
Individual ID (n=111)	Intercept	0.554	0.744			
Nest of origin (n=36)	Intercept	0.374	0.611			
	Age	0.007	0.083			
Nest of rearing (n=20)	Intercept	0.001	0.037			
	Age	0.015	0.122			
Residual		0.534	0.731			

Fixed factors	Estimate	SE	t	df	F	p
Intercept	15.525	0.141	101.416			
Hormone (TH)	-0.272	0.287	-0.947	1, 29.68	0.896	0.351
Age	0.841	0.035	24.076	1, 19.49	579.635	<0.001
Hormone × Age	0.113	0.049	2.334	1, 20.18	5.449	**0.030**

(c) Late nestling tarsus growth (day 7 – day 14)

Random effects:		Variance	Std. Dev.			
Individual ID (n=111)	Intercept	0.235	0.485			
Nest of origin (n=36)	Intercept	0.213	0.461			
	Age	0.012	0.111			
Nest of rearing (n=20)	Intercept	0.000[§]	0.000[§]			
	Age	<0.001	<0.001			
Residual		0.376	0.613			

Fixed factors	Estimate	SE	t	df	F	p
Intercept	20.660	0.164	126.251			
Hormone (TH)	-0.172	0.21	-0.818	1, 30.22	0.669	0.420
Age	0.521	0.024	22.090	1, 27.16	487.968	<0.001
Measurer 2	0.2220	0.184	1.197	2, 95.70	6.832	0.002
Measurer 3	0.586	0.173	3.381			
Hormone × Age	0.091	0.047	1.933	1, 31.06	3.737	0.062

Sample sizes:
Early growth, n = 99 control and 119 TH-supplemented individuals at day 2, 104 control and
117 TH-supplemented individuals at day 7. Please noticed that in total n = 42 nests during the
early growth, but n = 41 nests at day 2 and n = 41 nests at day 7, because we missed out the

Fig. 1. Nestling growth in prenatally TH-supplemented (blue dots) and control (red dots)

groups. (a) body mass growth during early nestling stage (day 2 to day 7 post-hatching) (b)

body mass growth late nestling stage (day 7 to day 14 post-hatching), (c) structural size (tarsus)

growth during late nestling stages. Means±SE (large dots) and scatter of the raw data (small,

semi-transparent dots) are shown. See sample sizes in the text.

Fig 2. (a) Relative mitochondrial density (z-scale) and (b) relative telomere length in prenatally

TH-supplemented (blue dots) and control groups (red dots) at different developmental stages

(day 7 and 14 post-hatching and as juveniles, ca. 3 months of age). Means±SE (large dots) and

scatter of the raw data (small, semi-transparent dots) are shown. See sample sizes in the text.

Biomarkers of intracellular oxidative status (total GSH, GSH:GSSG ratio) and damage to lipids

14 days post-hatching did not clearly differ among the treatment groups (marginal means±SE:

total GSH μmol/mg protein: TH-supplemented, 0.253±0.022, control, 0.249±0.023; GSH:GSSG

ratio: TH-supplemented, 0.045±0.008, control, 0.053±0.010; MDA nmol/mg protein: TH-

supplemented, 0.038±0.005, control, 0.037±0.005; Table S2). Plasma T3 and T4 also did not

differ among prenatally TH-supplemented and control groups 14 days post hatching (marginal

means±SE pmol/ml T3: TH-supplemented, 1.21±0.17, control, 1.23±0.19; T4: TH-

supplemented, 7.35±0.946, control, 7.39±1.02; Table S3).

Mitochondrial density decreased with age, but there were no clear differences between

prenatally TH-supplemented and control groups at any age (d7, d14 or juveniles, Fig 2a, Table

2). Post-hoc analyses indicated significant decrease between each age category (all Tukey-

adjusted p < 0.001), regardless of the hormone treatment.

Telomere length decreased with age (Table 2, Fig 2b), but the pattern was different

between TH-supplemented and control group (Table 2, Fig 2b): Early in the nestling phase the

difference between the prenatally TH-supplemented and control group approached

significance (d7, Tukey post-hoc test: TH vs control: $t_{46.2}$ = 1.804, p = 0.078). On day 14 post-

hatching, telomeres were significantly shorter (13%) in the TH group (Tukey post-hoc test, TH

vs Control: $t_{46.8}$ = 3.166, p = 0.003), and this difference was accentuated in juveniles, offspring

from TH supplemented eggs having telomeres being ca. 33% shorter than offspring from

control group (Tukey post-hoc test, TH vs Control: $t_{109.8}$ = 4.311, p < 0.0001). Further, we added

the percentage of body mass gain from day 7 to day 14 in the model to examine whether the

effect of prenatal TH supplementation was via its effect on nestling growth, but this did not

qualitatively change the results. The percentage of body mass gain did not show a clear effect

on telomere length (estimate±SE = -0.575±0.580, $F_{1,64.54}$=0.981, p=0.326). Similarly, the three

oxidative stress biomarkers also did not explain the difference in telomere shortening (all p's

> 0.31).

Hatching success did not clearly differ among prenatally TH supplemented and control individuals (TH 67.6%, control 69.1%; z =0.306, p = 0.760, Table 3a). While nestling survival to fledging was higher in prenatally TH supplemented than control nestlings (TH: 83.05% vs CO 69.44%), our model did not provide clear support for such a difference (z = 0.136, p = 0.892, Table 3b) and an analysis from egg to fledgling stage yielded qualitatively similar results (treatment p = 0.86). Fledging success was lower in the nests with larger brood sizes (z = -2.333, p = 0.020, Table 3b). Juvenile recapture rate was not significantly different across the groups (TH: 28.57% vs CO 34.00%, z=-0.993, p = 0.321, Table 3c).

413   **Table 2.** Linear mixed models of mitochondrial density and telomere length in blood cells

414   growth in response to prenatal TH-supplementation. In these models, age was as a categorical

415   factor with three levels. For the purpose of mean-centering, two dummy variables ("Age day

416   7" and "Age day 14") were created to represent the respective contrast against the juvenile

417   age.

(a) Mitochondria density

Random effects:		Variance	Std. Dev.			
Individual ID (n=78)	Intercept	0.084	0.290			
Nest of origin (n=35)	Intercept	0.041	0.203			
Nest of rearing (n=19)	Intercept	0.000[§]	0.000[§]			
DNA batch (n=8)	Intercept	0.017	0.129			
Residual		0.244	0.494			
Fixed factors	**Estimate**	**SE**	**t**	**df**	**F**	**p**
Intercept	-0.023	0.079	-0.295			
Hormone (TH)	0.021	0.135	0.155	1, 24.00	0.024	0.878
Sex	0.200	0.121	1.651	1, 73.16	2.723	0.103
Age day 7	2.202	0.121	18.134	1, 111.27	328.825	**<0.001**
Age day 14	1.225	0.123	9.968	1, 111.12	99.357	**<0.001**
Hormone × Age day 7	0.352	0.240	1.464	1, 111.78	2.142	0.146
Hormone × Age day 14	0.197	0.243	0.807	1, 111.70	0.652	0.421

(b) relative telomere length

Random effects:		Variance	Std. Dev.			
Individual ID (n=77)	Intercept	0.093	0.305			
Nest of origin (n=35)	Intercept	0.000[§]	0.000[§]			
Nest of rearing (n=19)	Intercept	0.026	0.162			
DNA batch (n=8)	Intercept	0.394	0.628			
Residual		0.513	0.716			
Fixed factors	**Estimate**	**SE**	**t**	**df**	**F**	**p**
Intercept	-0.085	0.236	-0.361			
Hormone (TH)	-0.635	0.153	-4.150	1, 16.06	17.223	**<0.001**
Sex	-0.268	0.154	-1.742	1, 65.94	3.035	0.086
Age day 7	0.904	0.174	5.197	1, 117.27	27.008	**<0.001**
Age day 14	0.654	0.175	3.728	1, 115.66	13.895	**<0.001**
Brood size at day 2	0.213	0.094	2.271	1, 10.18	5.157	**0.046**
Hormone × Age day 7	1.039	0.343	3.027	1, 120.32	9.165	**0.003**
Hormone × Age day 14	0.741	0.347	2.137	1, 119.21	4.567	**0.035**
Hormone × Brood size	-0.255	0.151	-1.681	1, 32.83	2.827	0.102

Sample sizes: n = 37 control and 41 TH-supplemented individuals at day 7; 33 control and 38 TH-supplemented individuals at day 14; 16 control and 13 TH-treatment recaptured juveniles
[§] Singular fit (i.e. insufficient data to reliably estimate the variance)

418   **Table 3.** Generalized linear mixed models of hatching success, fledging success, and post-

419   fledging survival. For fledging success and post-fledging survival, only the nestlings that were

420   reared in the nests without temperature manipulation were included. The models were fit

421   by maximum likelihood approach using Laplace approximation with binomial distribution

422   and logit link function. All input variables were mean-centered before model-fitting.

(a) Hatching success (n = 354 eggs)				
Random effects:		**Variance**	**Std. Dev.**	
Nest of origin (n=42)	Intercept	0.052	0.228	
Fixed factors	**Estimate**	**SE**	**z**	**p**
Intercept	0.781	0.123	6.339	
Hormone (TH)	0.074	0.241	0.306	0.760
(b) Fledging success (n = 131 nestlings)				
Random effects:		**Variance**	**Std. Dev.**	
Nest of origin (n=39)	Intercept	0.00[§]	0.00[§]	
Nest of rearing (n=27)	Intercept	582.60	24.14	
Fixed factors	**Estimate**	**SE**	**z**	**p**
Intercept	-0.263	3.036	-0.087	
Hormone (TH)	0.195	1.436	0.136	0.892
Brood size	-10.640	4.562	-2.333	**0.020**
(c) Post-fledging survival (n = 99 fledlings)				
Random effects:		**Variance**	**Std. Dev.**	
Nest of origin (n=34)	Intercept	0.000[§]	0.000[§]	
Nest of rearing (n=19)	Intercept	1.584	1.258	
Fixed factors	**Estimate**	**SE**	**z**	**p**
Intercept	-0.925	0.395	-2.339	
Hormone (TH)	-0.569	0.573	-0.993	0.321

[§] **Singular fit (i.e. insufficient data to reliably estimate the variance)**
Sample size:
Hatching success: n = 176 control eggs and 178 TH-eggs
Fledging success: n = 72 control nestlings and 59 TH-nestlings

Discussion

By manipulating prenatal exposure to THs in a wild passerine species, we demonstrate that an increase in prenatal THs can accelerate both postnatal growth and telomere shortening. Yet, we did not detect significant effects of elevated prenatal THs on postnatal oxidative stress levels, cellular energetics measured as mtDNA copy number, circulating TH levels, or short to medium-term survival (*i.e.* hatching success, fledging success, and apparent survival to the next autumn/winter).

Shorter telomeres in the TH-supplemented group were only detected from 14 days after hatching onwards, which seems to exclude direct effects or prenatal TH on telomere dynamics during embryonic development. Yet, this coincides with the accelerated postnatal body mass and tarsus growth observed in the TH-group (between day 7 and 14) compared to controls, and faster growth can accelerate telomere shortening either through enhanced cell division or through inducing oxidative stress (Monaghan and Ozanne, 2018). Considering the lack of impact of prenatal THs on oxidative stress reported here and in previous work on birds (Hsu et al., 2020, 2019a; Sarraude et al., 2020a; Sarraude et al., 2020b), it is not surprising the accelerated telomere shortening observed in this study could not be explained by the measured oxidative stress biomarkers. Yet, we cannot fully rule out this explanation since DNA damage was not directly assessed and oxidative stress sensitivity might vary between different biomolecules (Reichert and Stier, 2017). The effect of prenatal THs on telomere length seemed to increase with age (*i.e.* stronger effect observed in juveniles than at day 14), which could be explained by a delayed effect of fast postnatal growth (day 7 to day 14) because telomeres only shorten at the subsequent cellular division, and therefore delayed effects are likely to be observed (Monaghan and Ozanne, 2018). Nevertheless, including body mass growth between

day 7 and day 14 in the model did not explain the difference in telomere shortening. Our

hypothesis that prenatal THs could program postnatal metabolic and endocrine function (and

thus affect telomere shortening indirectly) was not supported as we found no evidence for

differences in mitochondrial density or plasma TH levels (key coordinators of metabolism)

across the treatment groups. The latter results are not fully surprising considering the limited

evidence supporting a prenatal programming of plasma THs (sex-specific effect on T4 only,

Hsu et al. 2017; no effect, Hsu et al. 2020) or mitochondrial density (no effect, Stier et al. 2020;

Hsu et al. 2020). Yet, our measurements are limited to blood and tissue-specific effects of TH

on mitochondrial density or TH-related cellular machinery (receptors and converting enzymes;

Ruuskanen et al. 2022) may occur as this has been shown for instance for glucocorticoids

(Weber et al. 2002). According to the *metabolic telomere attrition hypothesis*, telomere

shortening is likely to be increased during energy-demanding periods, and accelerated growth

under limited resources is likely carrying a metabolic cost (Casagrande and Hau, 2019).

The effects observed here on telomeres in response to increased prenatal TH levels

are in sharp contrast with our previous findings in another passerine species, the collared

flycatcher (Stier et al. 2020). Indeed, increasing prenatal THs in the collared flycatcher

increased telomere length measured very shortly after hatching (day 2), while not affecting

telomere shortening during postnatal growth (Stier et al. 2020). There are several alternative

explanations for these contrasted findings: (1) we cannot exclude that prenatal THs increased

telomere length during embryonic development in great tits since our first telomere length

measurement (day 7, *ca.* 70% of fledging body mass) was done considerably later than in

collared flycatchers (*i.e.* day 2, *ca.* 20% of fledging body mass). (2) THs differently influenced

post-natal growth in the two species: great tits from the TH-supplemented eggs were initially

slightly smaller, but grew faster in the late nestling period, whereas collared flycatchers from

the TH-supplemented eggs were bigger soon after hatching, but grew slightly slower during

the nestling stage (Hsu et al., 2019a). These contrasting growth patterns may explain, at least

partly, the contrasted findings in these two species: an increased metabolic demand for fast

postnatal growth in great tit in response to prenatal TH could accelerate telomere shortening,

while a reduced metabolic demand in collared flycatcher from TH-injected eggs could enable

telomere maintenance (Casagrande and Hau, 2019). Measurements of the mTOR signaling

pathway could shed light on the validity of this hypothesis (Casagrande and Hau, 2019). (3)

The effects of maternal signals on offspring are expected to differ based on environmental

conditions, so called context-dependent effects: For example, prenatal androgens have been

found to differentially influence offspring growth depending on season or food availability

(Groothuis et al., 2020; Muriel et al., 2015). We recently found no evidence for prenatal THs

differentially influencing growth and early-life survival depending on rearing temperature

(Hsu et al. 2020), yet the influence of resource availability was not tested. In two sister species,

collared and pied flycatchers, differential effects of prenatal THs on growth may have been

explained by differences in food resources (Sarraude et al. 2020b). (4) These two species

exhibit different life-histories, as collared flycatchers are migratory and great tits (relatively)

sedentary. Migratory species have generally higher metabolism (Jetz et al., 2008) as well as an

overall faster pace of life (Soriano-Redondo et al. 2021), and may thus present differences in

TH physiology. We may speculate that the telomere maintenance, and role of THs in the

regulation of telomere maintenance may differ across species with different life-histories.

Telomere maintenance is known to be influenced by e.g. species lifespan (Haussmann et al.,

2004) and migratory populations within a species have shorter telomeres (Bauer et al., 2016).

Yet, to our knowledge, studies have not considered migratory vs. non-migratory species in

telomere maintenance. Species-specific effects of the prenatal hormonal environment on

telomere dynamics have already been described: elevated prenatal glucocorticoid levels led

to shorter telomeres in domestic chicken and female zebra finch (Haussmann et al., 2012;

Tissier et al., 2014), but to longer telomeres in yellow-legged gull (Noguera, da Silva, and

Velando 2020). To understand if any of the hypotheses presented above is likely to explain the

contrasted effects found here and in Stier et al. (2020), more studies on the impact of prenatal

THs on telomere biology across taxa are needed.

As mentioned above, in the collared flycatcher, elevated yolk THs increased nestlings

body mass at day 2 (Hsu et al. 2019), whereas in this great tit study, elevated yolk THs only

non-significantly reduced nestling body mass at day 2 (marginal mean ± SE = -0.402 ± 0.245,

p=0.109). In both studies, the lighter group of nestlings caught up with the other group before

fledging. It therefore appears that the main difference lies on the differential effects of THs

during the peri-hatching period. Interestingly, another experiment in the same population of

great tits found that elevated yolk THs significantly shortened the time needed to hatch by 0.6

days on average (Cossin-Sevrin et al. 2022). Despite the shorter developmental time, the

nestling body mass at day 2 was also found non-significant (Cossin-Sevrin et al. 2022). Thus,

the exact peri-hatching effects of prenatal THs in altricial birds still need more information to

clarify.

To understand the selective pressures on maternal allocation and signaling, it is

important to characterize true fitness effects from development to survival and lifetime

reproductive success. In contrast to our predictions, we found no effects of elevated prenatal

TH-supplementation on apparent survival or individual condition during their first autumn/

winter. Unfortunately, longer-term effects on survival and reproductive success are very

challenging to measure in such an experimental setting in the wild, as the recruitment to first

breeding is usually very low (e.g. < 10% Radersma et al., 2015), which requires a very high

number of nests to be manipulated, something often not feasible for both logistical and ethical

reasons. Yet, long-term effects of accelerated early-life telomere shortening may lead to a

decrease in longer-term survival and lifespan (Eastwood et al., 2019; Heidinger et al., 2012).

The negative effects on later stages suggest that there need to be benefits of high prenatal

THs for the offspring or mother, potentially during early-life stages, where selection can be

strong. In this study, the elevated growth rates could increase offspring competitive abilities

early in life, yet, the effects of prenatal THs on growth seem to be highly inconsistent across

studies (Hsu et al., 2020, 2019a, 2017; Ruuskanen et al., 2016a; Sarraude et al., 2020a;

Sarraude et al., 2020b). While circulating THs are known to be associated with health and

ageing in humans and mammalian models (Bano et al. 2017, 2019; Møllehave et al. 2020), our

study is the first to show that exposure to higher prenatal thyroid hormones can lead to

accelerated cellular ageing measured through telomere length. This should stimulate further

research using both epidemiological and experimental approaches across taxa to uncover the

potential regulation of telomere biology by thyroid hormones both pre- and postnatally.

Author contribution

BYH, AS & SR designed the study and collected field data. NCS, AS & SR conducted laboratory
work. BYH analyzed the data. BYH, AS & SR wrote the manuscript, with input from NCS.
537

Acknowledgements

This study was financially supported by the Academy of Finland (#286278 to SR). NCS

acknowledges support from the EDUFI Fellowship and Maupertuis Grant. B-Y.H work was supported

by grants from the Ella and Georg Ehrnrooth Foundation and Academy of Finland (#332716). AS was

supported by a 'Turku Collegium for Science and Medicine' Fellowship and a Marie Sklodowska-Curie

Postdoctoral Fellowship (#894963) at the time of writing. We thank all field assistants, especially

Lucas Bousseau, Thomas Rosille, Hsiao-Yin Liu, and Jorma Nurmi for their great effort. All

authors declare no conflict of interest.

Data accessibility

548   All data is archived and available in Figshare (DOI: 10.6084/m9.figshare.20110409).

References

550   Aljanabi, S.M., Martinez, I., 1997. Universal and rapid salt-extraction of high quality genomic DNA for
551   PCR-based techniques. Nucleic Acids Research. https://doi.org/10.1093/nar/25.22.4692
552   Arbeev, K.G., Verhulst, S., Steenstrup, T., Kark, J.D., Bagley, O., Kooperberg, C., Reiner, A.P., Hwang,
553   S.-J., Levy, D., Fitzpatrick, A.L., Christensen, K., Yashin, A.I., Aviv, A., 2020. Association of
554   Leukocyte Telomere Length With Mortality Among Adult Participants in 3 Longitudinal
555   Studies. JAMA Netw Open 3, e200023.
556   https://doi.org/10.1001/jamanetworkopen.2020.0023
557   Atema, E., Mulder, E., van Noordwijk, A.J., Verhulst, S., 2019. Ultralong telomeres shorten with age in
558   nestling great tits but are static in adults and mask attrition of short telomeres. Mol Ecol
559   Resour 19, 648–658. https://doi.org/10.1111/1755-0998.12996
560   Bano, A., Chaker, L., Mattace-Raso, F.U.S., Terzikhan, N., Kavousi, M., Ikram, M.A., Peeters, R.P.,
561   Franco, O.H., 2019. Thyroid function and life expectancy with and without noncommunicable
562   diseases: A population-based study. PLoS Med 16, e1002957.
563   https://doi.org/10.1371/journal.pmed.1002957
564   Bano, A., Dhana, K., Chaker, L., Kavousi, M., Ikram, M.A., Mattace-Raso, F.U.S., Peeters, R.P., Franco,
565   O.H., 2017. zeta Association of Thyroid Function With Life Expectancy With and Without
566   Cardiovascular Disease The Rotterdam Study. Jama Internal Medicine.
567   https://doi.org/10.1001/jamainternmed.2017.4836
568   Barnes, S.K., Ozanne, S.E., 2011. Pathways linking the early environment to long-term health and
569   lifespan. Progress in Biophysics and Molecular Biology 106, 323–336.
570   https://doi.org/10.1016/j.pbiomolbio.2010.12.005
571   Bates, D., Machler, M., Bolker, B.M., Walker, S.C., 2015. Fitting Linear Mixed-Effects Models Using
572   lme4. Journal of Statistical Software. https://doi.org/10.18637/jss.v067.i01
573   Bauer, C.M., Heidinger, B.J., Ketterson, E.D., Greives, T.J., 2016. A migratory lifestyle is associated
574   with shorter telomeres in a songbird (Junco hyemalis). Auk. https://doi.org/10.1642/auk-16-
575   56.1
576   Casagrande, S., Hau, M., 2019. Telomere attrition: metabolic regulation and signalling function?
577   Biology Letters. https://doi.org/10.1098/rsbl.2018.0885
578   Chang, H.W., Cheng, C.A., Gu, D.L., Chang, C.C., Su, S.H., Wen, C.H., Chou, Y.C., Chou, T.C., Yao, C.T.,
579   Tsai, C.L., Cheng, C.C., 2008. High-throughput avian molecular sexing by SYBR green-based
580   real-time PCR combined with melting curve analysis. Bmc Biotechnology.
581   https://doi.org/10.1186/1472-6750-8-12

Cioffi, F., Senese, R., Lanni, A., Goglia, F., 2013. Thyroid hormones and mitochondria: With a brief
look at derivatives and analogues. Molecular and Cellular Endocrinology.
https://doi.org/10.1016/j.mce.2013.06.006
Cossin-Sevrin, N., Hsu, B.-Y., Marciau, C., Viblanc, V.A., Ruuskanen, S., Stier, A., 2022. Effect of
prenatal glucocorticoids and thyroid hormones on developmental plasticity of mitochondrial
aerobic metabolism, growth and survival: an experimental test in wild great tits. Journal of
Experimental Biology 225, jeb243414. https://doi.org/10.1242/jeb.243414
Criscuolo, F., Torres, R., Zahn, S., Williams, T.D., 2020. Telomere dynamics from hatching to sexual
maturity and maternal effects in the 'multivariate egg.' Journal of Experimental Biology 223,
jeb232496. https://doi.org/10.1242/jeb.232496
Eastwood, J.R., Hall, M.L., Teunissen, N., Kingma, S.A., Aranzamendi, N.H., Fan, M., Roast, M.,
Verhulst, S., Peters, A., 2019. Early-life telomere length predicts lifespan and lifetime
reproductive success in a wild bird. Molecular Ecology. https://doi.org/10.1111/mec.15002
Ellegren, H., Fridolfsson, A.K., 1997. Male-driven evolution of DNA sequences in birds. Nat Genet 17,
182–184. https://doi.org/10.1038/ng1097-182
Entringer, S., de Punder, K., Buss, C., Wadhwa, P.D., 2018. The fetal programming of telomere biology
hypothesis: an update. Philos Trans R Soc Lond B Biol Sci 373, 20170151.
https://doi.org/10.1098/rstb.2017.0151
Espin, S., Sanchez Virosta, P., García-Fernández, A., Eeva, T., 2017. A microplate adaptation of the
thiobarbituric acid reactive substances assay to determine lipid peroxidation fluorometrically
in small sample volumes. Revista de Toxicologia 34, In press.
Godfrey, K.M., Barker, D.J., 2001. Fetal programming and adult health. Public Health Nutr 4, 611–
624. https://doi.org/10.1079/phn2001145
Groothuis, T.G.G., Kumar, N., Hsu, B.Y., 2020. Explaining discrepancies in the study of maternal
effects: the role of context and embryo. Current Opinion in Behavioral Sciences.
https://doi.org/10.1016/j.cobeha.2020.10.006
Halekoh, U., Hojsgaard, S., 2014. Kenward-Roger Approximation and Parametric Bootstrap Methods
for Tests in Linear Mixed Models - The R Package pbkrtest. Journal of Statistical Software.
Hartig, F. (2021). DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed) Regression
Models. R package version 0.4.3. https://CRAN.R-project.org/package=DHARMa
Haussmann, M.F., Longenecker, A.S., Marchetto, N.M., Juliano, S.A., Bowden, R.M., 2012. Embryonic
exposure to corticosterone modifies the juvenile stress response, oxidative stress and
telomere length. Proceedings of the Royal Society B-Biological Sciences.
https://doi.org/10.1098/rspb.2011.1913
Haussmann, M.F., Winkler, D.W., Huntington, C.E., Nisbet, I.C.T., Vleck, C.M., 2004. Telomerase
expression is differentially regulated in birds of differing life span. Strategies for Engineered
Negligible Senescence: Why Genuine Control of Aging May Be Foreseeable.
https://doi.org/10.1196/annals.1297.029
Heidinger, B.J., Blount, J.D., Boner, W., Griffiths, K., Metcalfe, N.B., Monaghan, P., 2012. Telomere
length in early life predicts lifespan. Proceedings of the National Academy of Sciences 109,
1743–1748. https://doi.org/10.1073/pnas.1113306109
Hsu, B.Y., Dijkstra, C., Darras, V.M., de Vries, B., Groothuis, T.G.G., 2017. Maternal thyroid hormones
enhance hatching success but decrease nestling body mass in the rock pigeon (Columba
livia). General and Comparative Endocrinology. https://doi.org/10.1016/j.ygcen.2016.10.011
Hsu, B.Y., Doligez, B., Gustafsson, L., Ruuskanen, S., 2019a. Transient growth-enhancing effects of
elevated maternal thyroid hormones at no apparent oxidative cost during early postnatal
period. Journal of Avian Biology. https://doi.org/10.1111/jav.01919
Hsu, B.Y., Sarraude, T., Cossin-Sevrin, N., Crombecque, M., Stier, A., Ruuskanen, S., 2020. Testing for
context-dependent effects of prenatal thyroid hormones on offspring survival and
physiology: an experimental temperature manipulation. Scientific Reports.
https://doi.org/10.1038/s41598-020-71511-y

Hsu, B.Y., Verhagen, I., Gienapp, P., Darras, V.M., Visser, M.E., Ruuskanen, S., 2019b. Between- and
Within-Individual Variation of Maternal Thyroid Hormone Deposition in Wild Great Tits
(Parus major). American Naturalist. https://doi.org/10.1086/704738
Jetz, W., Freckleton, R.P., McKechnie, A.E., 2008. Environment, Migratory Tendency, Phylogeny and
Basal Metabolic Rate in Birds. Plos One. https://doi.org/10.1371/journal.pone.0003261
Kärkkäinen, T., Briga, M., Laaksonen, T., Stier, A., n.d. Within-individual repeatability in telomere
length: A meta-analysis in nonmammalian vertebrates. Molecular Ecology n/a.
https://doi.org/10.1111/mec.16155
Kuznetsova, A., Brockhoff, P.B., Christensen, R.H.B., 2017. lmerTest Package: Tests in Linear Mixed
Effects Models. Journal of Statistical Software 82, 1–26.
https://doi.org/10.18637/jss.v082.i13
Larsen, S., Nielsen, J., Hansen, C.N., Nielsen, L.B., Wibrand, F., Stride, N., Schroder, H.D., Boushel, R.,
Helge, J.W., Dela, F., Hey-Mogensen, M., 2012. Biomarkers of mitochondrial content in
skeletal muscle of healthy young human subjects. J Physiol 590, 3349–3360.
https://doi.org/10.1113/jphysiol.2012.230185
Length, R.V. (2021). emmeans: Estimated Marginal Means, aka Least-Squares Means. R package
version 1.6.2-1. https://CRAN.R-project.org/package=emmeans
Liu, J.S., Chen, Y.Q., Li, M., 2006. Thyroid hormones increase liver and muscle thermogenic capacity in
the little buntings (Emberiza pusilla). Journal of Thermal Biology.
https://doi.org/10.1016/j.jtherbio.2006.01.002
López-Otín, C., Blasco, M.A., Partridge, L., Serrano, M., Kroemer, G., 2013. The Hallmarks of Aging.
Cell 153, 1194–1217. https://doi.org/10.1016/j.cell.2013.05.039
Marchetto, N.M., Glynn, R.A., Ferry, M.L., Ostojic, M., Wolff, S.M., Yao, R., Haussmann, M.F., 2016.
Prenatal stress and newborn telomere length. American Journal of Obstetrics and
Gynecology 215, 94.e1-94.e8. https://doi.org/10.1016/j.ajog.2016.01.177
McNabb, F.M.A., 2007. The hypothalamic-pituitary-thyroid (HPT) axis in birds and its role in bird
development and reproduction. Critical Reviews in Toxicology.
https://doi.org/10.1080/10408440601123552
Medici, M., Timmermans, S., Visser, W., Keizer-Schrama, S., Jaddoe, V.W.W., Hofman, A., Hooijkaas,
H., de Rijke, Y.B., Tiemeier, H., Bongers-Schokking, J.J., Visser, T.J., Peeters, R.P., Steegers,
E.A.P., 2013. Maternal Thyroid Hormone Parameters during Early Pregnancy and Birth
Weight: The Generation R Study. Journal of Clinical Endocrinology & Metabolism.
https://doi.org/10.1210/jc.2012-2420
Metcalfe, N.B., Monaghan, P., 2001. Compensation for a bad start: grow now, pay later? Trends in
Ecology & Evolution.
Møllehave, L.T., Skaaby, T., Linneberg, A., Knudsen, N., Jørgensen, T., Thuesen, B.H., 2020. The
association of thyroid stimulation hormone levels with incident ischemic heart disease,
incident stroke, and all-cause mortality. Endocrine 68, 358–367.
https://doi.org/10.1007/s12020-020-02216-5
Monaghan, P., Ozanne, S.E., 2018. Somatic growth and telomere dynamics in vertebrates:
relationships, mechanisms and consequences. Philos Trans R Soc Lond B Biol Sci 373,
20160446. https://doi.org/10.1098/rstb.2016.0446
Mullur, R., Liu, Y.-Y., Brent, G.A., 2014. Thyroid Hormone Regulation of Metabolism. Physiological
Reviews 94, 355–382. https://doi.org/10.1152/physrev.00030.2013
Muñoz-Lorente, M.A., Cano-Martin, A.C., Blasco, M.A., 2019. Mice with hyper-long telomeres show
less metabolic aging and longer lifespans. Nat Commun 10, 4723.
https://doi.org/10.1038/s41467-019-12664-x
Muriel, J., Salmon, P., Nunez-Buiza, A., de Salas, F., Perez-Rodriguez, L., Puerta, M., Gil, D., 2015.
Context-dependent effects of yolk androgens on nestling growth and immune function in a
multibrooded passerine. Journal of Evolutionary Biology. https://doi.org/10.1111/jeb.12668

Noguera, J.C., da Silva, A., Velando, A., n.d. Egg corticosterone can stimulate telomerase activity and
promote longer telomeres during embryo development. Molecular Ecology n/a.
https://doi.org/10.1111/mec.15694
Parolini, M., Possenti, C.D., Caprioli, M., Rubolini, D., Romano, A., Saino, N., 2019. Egg Testosterone
Differentially Affects Telomere Length in Somatic Tissues of Yellow-Legged Gull Embryos.
Physiological and Biochemical Zoology. https://doi.org/10.1086/705037
Podmokla, E., Drobniak, S.M., Rutkowska, J., 2018. Chicken or egg? Outcomes of experimental
manipulations of maternally transmitted hormones depend on administration method - a
meta-analysis. Biological Reviews. https://doi.org/10.1111/brv.12406
Radersma, R., Komdeur, J., Tinbergen, J.M., 2015. Early morning fledging improves recruitment in
Great Tits Parus major. Ibis 157, 351–355. https://doi.org/10.1111/ibi.12230
Reichert, S., Stier, A., 2017. Does oxidative stress shorten telomeres in vivo? A review. Biology
Letters. https://doi.org/10.1098/rsbl.2017.0463
Ruuskanen, S., Darras, V.M., Visser, M.E., Groothuis, T.G.G., 2016a. Effects of experimentally
manipulated yolk thyroid hormone levels on offspring development in a wild bird species.
Hormones and Behavior. https://doi.org/10.1016/j.yhbeh.2016.03.006
Ruuskanen, S., Espin, S., Sanchez-Virosta, P., Sarraude, T., Hsu, B.Y., Pajunen, P., Costa, R.A., Eens, M.,
Hargitai, R., Torok, J., Eeva, T., 2019. Transgenerational endocrine disruption: Does elemental
pollution affect egg or nestling thyroid hormone levels in a wild songbird? Environmental
Pollution. https://doi.org/10.1016/j.envpol.2019.01.088
Ruuskanen, S., Groothuis, T.G.G., Schaper, S.V., Darras, V.M., de Vries, B., Visser, M.E., 2016b.
Temperature-induced variation in yolk androgen and thyroid hormone levels in avian eggs.
General and Comparative Endocrinology. https://doi.org/10.1016/j.ygcen.2016.05.026
Ruuskanen, S., Hsu, B.Y., 2018. Maternal Thyroid Hormones: An Unexplored Mechanism Underlying
Maternal Effects in an Ecological Framework. Physiological and Biochemical Zoology.
https://doi.org/10.1086/697380
Ruuskanen, S., Hsu, B.-Y., Heinonen, A., Darras, V., Vainio, M., Sarraude, T., Rokka, A., 2018. A new
method for measuring thyroid hormones using nano-LC-MS/MS. Journal of Chromatography
B.
Salmón, P., Nilsson, J.F., Nord, A., Bensch, S., Isaksson, C., 2016. Urban environment shortens
telomere length in nestling great tits, Parus major. Biol Lett 12, 20160155.
https://doi.org/10.1098/rsbl.2016.0155
Salmón, P., Nilsson, J.F., Watson, H., Bensch, S., Isaksson, C., 2017. Selective disappearance of great
tits with short telomeres in urban areas. Proc Biol Sci 284, 20171349.
https://doi.org/10.1098/rspb.2017.1349
Sarraude, T., Hsu, B.-Y., Groothuis, T., Ruuskanen, S., 2020a. Manipulation of prenatal thyroid
hormones does not influence growth or physiology in nestling pied flycatchers. Physiological
and Biochemical Zoology.
Sarraude, T., Hsu, B.Y., Groothuis, T., Ruuskanen, S., 2020b. Testing the short-and long-term effects
of elevated prenatal exposure to different forms of thyroid hormones. Peerj.
https://doi.org/10.7717/peerj.10175
Schielzeth, H., 2010. Simple means to improve the interpretability of regression coefficients.
Methods in Ecology and Evolution 1, 103–113. https://doi.org/10.1111/j.2041-
210X.2010.00012.x
Schielzeth, H., Dingemanse, N.J., Nakagawa, S., Westneat, D.F., Allegue, H., Teplitsky, C., Réale, D.,
Dochtermann, N.A., Garamszegi, L.Z., Araya-Ajoy, Y.G., 2020. Robustness of linear mixed-
effects models to violations of distributional assumptions. Methods in Ecology and Evolution
11, 1141–1152. https://doi.org/10.1111/2041-210X.13434
Smith, S.M., Nager, R.G., Costantini, D., 2016. Meta-analysis indicates that oxidative stress is both a
constraint on and a cost of growth. Ecol Evol 6, 2833–2842.
https://doi.org/10.1002/ece3.2080

Stier, A., Bize, P., Hsu, B.Y., Ruuskanen, S., 2019. Plastic but repeatable: rapid adjustments of
mitochondrial function and density during reproduction in a wild bird species. Biology
Letters. https://doi.org/10.1098/rsbl.2019.0536
Stier, A., Delestrade, A., Bize, P., Zahn, S., Criscuolo, F., Massemin, S., 2016. Investigating how
telomere dynamics, growth and life history covary along an elevation gradient in two
passerine species. Journal of Avian Biology 47, 134–140. https://doi.org/10.1111/jav.00714
Stier, A., Hsu, B.Y., Marciau, C., Doligez, B., Gustafsson, L., Bize, P., Ruuskanen, S., 2020. Born to be
young? Prenatal thyroid hormones increase early-life telomere length in wild collared
flycatchers. Biology Letters. https://doi.org/10.1098/rsbl.2020.0364
Stier, A., Massemin, S., Criscuolo, F., 2014. Chronic mitochondrial uncoupling treatment prevents
acute cold-induced oxidative stress in birds. Journal of Comparative Physiology B-Biochemical
Systems and Environmental Physiology. https://doi.org/10.1007/s00360-014-0856-6
Stier, A., Massemin, S., Zahn, S., Tissier, M.L., Criscuolo, F., 2015. Starting with a handicap: effects of
asynchronous hatching on growth rate, oxidative stress and telomere dynamics in free-living
great tits. Oecologia. https://doi.org/10.1007/s00442-015-3429-9
Stier, A., Metcalfe, N.B., Monaghan, P., 2020. Pace and stability of embryonic development affect
telomere dynamics: an experimental study in a precocial bird model. Proc Biol Sci 287,
20201378. https://doi.org/10.1098/rspb.2020.1378
Tissier, M.L., Williams, T.D., Criscuolo, F., 2014. Maternal Effects Underlie Ageing Costs of Growth in
the Zebra Finch (Taeniopygia guttata). PLoS One 9, e97705.
https://doi.org/10.1371/journal.pone.0097705
Tschirren, B., Saladin, V., Fitze, P.S., Schwabl, H., Richner, H., 2005. Maternal yolk testosterone does
not modulate parasite susceptibility or immune function in great tit nestlings. Journal of
Animal Ecology 74: 675-682.

Verhulst, S., 2020. Improving comparability between qPCR-based telomere studies. Molecular
Ecology Resources 20, 11–13. https://doi.org/10.1111/1755-0998.13114
Vrijkotte, T.G.M., Hrudey, E.J., Twickler, M.B., 2017. Early Maternal Thyroid Function During
Gestation Is Associated With Fetal Growth, Particularly in Male Newborns. The Journal of
Clinical Endocrinology & Metabolism 102, 1059–1066. https://doi.org/10.1210/jc.2016-3452
Weber, K., Brück, P., Mikes, Z., Küpper, J-H., Klingenspor, M., Wiesner, R.J.. 2002. Glucocorticoid
Hormone Stimulates Mitochondrial Biogenesis Specifically in Skeletal Muscle. *Endocrinology*,
143, 177–184
Welcker, J., Chastel, O., Gabrielsen, G.W., Guillaumin, J., Kitaysky, A.S., Speakman, J.R., Tremblay, Y.,
Bech, C., 2013. Thyroid Hormones Correlate with Basal Metabolic Rate but Not Field
Metabolic Rate in a Wild Bird Species. Plos One.
https://doi.org/10.1371/journal.pone.0056229
Wilbourn, R.V., Moatt, J.P., Froy, H., Walling, C.A., Nussey, D.H., Boonekamp, J.J., 2018. The
relationship between telomere length and mortality risk in non-model vertebrate systems: a
meta-analysis. Philosophical Transactions of the Royal Society B-Biological Sciences.
https://doi.org/10.1098/rstb.2016.0447
Zhang, C., Yang, X., Zhang, Y., Guo, F., Yang, S., Peeters, R.P., Korevaar, T.I.M., Fan, J., Huang, H.-F.,
2019. Association Between Maternal Thyroid Hormones and Birth Weight at Early and Late
Pregnancy. The Journal of Clinical Endocrinology & Metabolism 104, 5853–5863.
https://doi.org/10.1210/jc.2019-00390
778

779
780

Supplementary material to

Prenatal thyroid hormones accelerate postnatal growth and telomere shortening in wild great tits

Bin-Yan Hsu[1], Nina Cossin-Sevrin[1,2], Antoine Stier[1,3*], Suvi Ruuskanen[1,4*$]
[1] Department of Biology, University of Turku, Finland
[2] Université de Strasbourg, Centre National de la Recherche Scientifique, Institut Pluridisciplinaire Hubert Curien, UMR 7178, 67087 Strasbourg, France
[3] Univ Lyon, Université Claude Bernard Lyon 1, CNRS, ENTPE, UMR 5023 LEHNA, F-69622, Villeurbanne, France
[4] Department of Biological and Environmental Science, University of Jyväskylä

joint senior authors
[$]*Corresponding author:* suvi.k.ruuskanen@jyu.fi

Table S1. Linear mixed models (GLMM) on the effects of prenatal thyroid hormone elevation on nestling body mass and tarsus length at day 2, day 7, and day 14 post-hatching. The denominator degree of freedom (ddf) were approximated by the Kenward-Roger method to calculate p values.

(a) Body mass at day 2 (n = 218 nestlings)

Random effects:		Variance	Std. Dev.			
Nest of origin (n=41)	Intercept	0.481	0.694			
Residual		0.625	0.791			
Fixed factors	**Estimate**	**SE**	**t**	**df**	**F**	**p**
Intercept	3.849	0.123	31.426			
Hormone (TH)	-0.402	0.245	-1.639	1, 38.69	2.686	0.109

Sample size: n = 99 control and 119 TH-supplemented
The total sample size at day 2 (n =218 nestlings from 41 nests) was smaller than that at day 7 because we missed the day 2 for one nest (n = 7 nestlings)

(b) Body mass at day 7 (n = 221 nestlings)

Random effects:		Variance	Std. Dev.			
Nest of origin (n=41)	Intercept	0.944	0.972			
Nest of rearing (n=41)	Intercept	<0.001	<0.001			
Residual		1.482	1.217			
Fixed factors	**Estimate**	**SE**	**t**	**df**	**F**	**p**
Intercept	12.495	0.175	71.459			
Hormone (TH)	-0.367	0.353	-1.040	1, 38.93	1.081	0.305
Cross-foster (yes)	-0.347	0.179	-1.945	1, 179.22	3.782	0.053
Brood size	-0.141	0.072	-1.956	1, 25.71	3.828	0.061
Hormone × Cross-foster	0.668	0.368	1.813	1, 28.49	3.286	0.080

(c) Tarsus length at day 7 (n = 221 nestlings)

Random effects:		Variance	Std. Dev.			
Nest of origin (n=41)	Intercept	0.4916	0.700			
Nest of rearing (n=41)	Intercept	0.000[§]	0.000[§]			
Residual		0.982	0.991			
Fixed factors	**Estimate**	**SE**	**t**	**df**	**F**	**p**
Intercept	18.940	0.205	92.592			
Hormone (TH)	-0.275	0.322	-0.855	1, 42.78	0.731	0.397
Cross-foster (yes)	-0.330	0.146	-2.268	1, 178.84	5.145	0.025
Proportion of TH nestlings	-0.297	0.482	-0.617	1, 54.90	0.381	0.540
Measurer 2	0.211	0.199	1.059	2, 204.00	3.357	0.037
Measurer 3	0.523	0.204	2.561			
Hormone × proportion of TH nestlings	-1.711	0.972	-1.762	1, 58.65	3.103	0.083

Sample size: n = 104 control and 117 TH-supplemented
[§] Singular fit (i.e. insufficient data to reliably estimate the variance)

Table S1 (cont.)

(d) Body mass at day 14 (n = 99 nestlings)

Random effects:		Variance	Std. Dev.			
Nest of origin (n=34)	Intercept	0.310	0.557			
Nest of rearing (n=19)	Intercept	0.228	0.477			
Residual		0.801	0.895			
Fixed factors	**Estimate**	**SE**	**t**	**df**	**F**	**p**
Intercept	18.603	0.176	105.693			
Hormone (TH)	0.099	0.303	0.747	1, 22.45	0.130	0.722

(e) Tarsus length at day 14 (n = 99 nestlings)

Random effects:		Variance	Std. Dev.			
Nest of origin (n=34)	Intercept	0.132	0.363			
Nest of rearing (n=19)	Intercept	$0.000^{§}$	$0.000^{§}$			
Residual		0.283	0.532			
Fixed factors	**Estimate**	**SE**	**t**	**df**	**F**	**p**
Intercept	23.199	0.238	97.313			
Hormone (TH)	0.157	0.175	0.899	1, 27.77	0.808	0.376
Measurer 2	-0.488	0.265	-1.845	2, 61.90	4.014	0.023
Measurer 3	-0.066	0.256	-0.258			

Sample sizes: n = 50 control and 49 TH-supplemented

§ Singular fit (i.e. insufficient data to reliably estimate the variance)

Table S2. Linear mixed models of blood oxidative stress biomarkers in response to prenatal TH-supplementation.

(a) total GSH µmol/mg protein (n = 58 nestlings)

Random effects:		Variance	Std. Dev.			
Nest of origin (n=29)	Intercept	0.031	0.177			
Nest of rearing (n=18)	Intercept	0.050	0.223			
Residual		0.044	0.210			
Fixed factors	**Estimate**	**SE**	**t**	**df**	**F**	**p**
Intercept	-1.492	0.114	-13.148			
Hormone (TH)	0.014	0.105	0.128	1, 15.46	0.016	0.900
GSH assay batch	0.221	0.145	1.526	1, 16.57	2.328	0.146

(b) GSH:GSSG ratio (n = 58 nestlings)

Random effects:		Variance	Std. Dev.			
Nest of origin (n=29)	Intercept	0.210	0.458			
Nest of rearing (n=18)	Intercept	0.031	0.175			
Residual		0.134	0.367			
Fixed factors	**Estimate**	**SE**	**t**	**df**	**F**	**p**
Intercept	-3.106	0.181	-17.207			
Hormone (TH)	-0.171	0.243	-0.703	1, 16.75	0.456	0.509
Body mass	0.106	0.059	1.794	1, 41.88	2.986	0.091
Proportion of TH-nestlings	-0.173	0.462	-0.375	1, 21.30	0.124	0.728
GSH assay batch	0.133	0.237	0.561	1, 16.87	0.307	0.587
Hormone × Proportion of TH-nestlings	1.787	0.926	1.930	1, 19.81	3.553	0.074

(c) MDA nmol/µg protein (n = 59 nestlings)

Random effects:		Variance	Std. Dev.			
Nest of origin (n=30)	Intercept	0.004	0.059			
Nest of rearing (n=18)	Intercept	0.004	0.060			
TBARS batch (n = 6)	Intercept	0.100	0.317			
Residual		0.039	0.198			
Fixed factors	**Estimate**	**SE**	**t**	**df**	**F**	**p**
Intercept	-3.286	0.135	-24.340			
Hormone (TH)	0.036	0.067	0.536	1, 16.86	0.287	0.599

Sample sizes:

total GSH and GSH:GSSG ratio, n = 26 control and 32 TH-supplemented

MDA, n = 26 control and 33 TH-supplemented

Table S3. Linear mixed models of blood plasma thyroid hormones growth in response to prenatal TH-supplementation.

(a) T3 pmol/ml (n = 24 nestlings)						
Random effects:		**Variance**	**Std. Dev.**			
Nest of origin (n=18)	Intercept	0.054	0.232			
Nest of rearing (n=11)	Intercept	0.127	0.357			
Residual		0.132	0.363			
Fixed factors	**Estimate**	**SE**	**t**	**df**	**F**	**p**
Intercept	1.218	0.147	8.273			
Hormone (TH)	-0.026	0.202	-0.128	1, 7.37	0.016	0.902
(b) T4 pmol/ml (n = 24 nestlings)						
Random effects:		**Variance**	**Std. Dev.**			
Nest of origin (n=18)	Intercept	0.738	0.859			
Nest of rearing (n=11)	Intercept	2.934	1.713			
Residual		5.721	2.392			
Fixed factors	**Estimate**	**SE**	**t**	**df**	**F**	**p**
Intercept	7.452	0.758	9.837			
Hormone (TH)	-0.041	1.113	-0.037	1, 6.57	0.001	0.973
Sex	2.101	1.119	1.877	1, 14.99	2.532	0.132

Sample sizes: n = 11 control and 13 TH-supplemented