**Life-stage and sex influence *Philornis* ectoparasitism in a Neotropical woodpecker (*Melanerpes***

***striatus*) with essential male parental care**

Running title: *Philornis* on adult *Melanerpes striatus*

Joshua B. LaPergola [1,2]

[1] Department of Neurobiology and Behavior, Cornell University, Ithaca, NY; Bird Population Studies

program, Cornell Lab of Ornithology, Ithaca, NY, USA

[2] Department of Ecology and Evolutionary Biology, Princeton University, Princeton, NJ, USA

Corresponding author: jl101@princeton.edu; jlapblca@gmail.com

https://orcid.org/0000-0002-2575-0751

**Conflict of interest.** The author declares that he has no conflicts of interest.

**Ethics approval.** All research activities described here were approved by the Dominican Republic's

Ministerio de Medio Ambiente y Recursos Naturales and conducted in accordance with IACUC protocol

2008-0185 at Cornell University.

**Availability of data and material.** The data that support the findings of this study will be openly

available in the repository Dryad upon acceptance of the manuscript.

**Acknowledgments**

I gratefully acknowledge the following individuals for providing invaluable assistance in the field: M.

Angelucci, H. Boyle, C. Cerrilla, W. Coleman, A. Diaz, L. Emerson, N. Gilbert, A. Janik, K. Kauffman,

T. Lacerda, A. Lascher-Posner, M. Larrieu, K. Larsen, C. Mathers-Winn, K. Nelsen, A. Occhialini, S.

Schubert, H. Stapleton, M. Walters, A. Waterman-Snow, P. Werner, A. Wichtendahl. I thank W. D.

Koenig, J. L. Dickinson, P. W. Sherman, M. S. Webster, H. K. Reeve, and I. J. Lovette for providing

constructive criticism at various points in the development of the main project that ultimately led to this

separate analysis. B. L. Bennett, T. C. Hendrix, S. B. Hex, C. Riehl, A. G. Savagian, and M. G. Smith

provided much appreciated constructive feedback and discussions that greatly improved the manuscript.

The following sources provided funding support: American Ornithological Society Wetmore Award,

Cornell Lab of Ornithology Athena Fund, Department of Neurobiology Animal Behavior Research Grant,

Society for the Study of Evolution Rosemary Grant Award, and Sigma Xi Grant in Aid of Research. The

following fellowships supported me during fieldwork: Charles Walcott Graduate Fellowship, Linda and

Samuel Kramer Graduate Student Fellowship, Eleanore Stuart Graduate Fellowship, Andrew '78 and

Margaret Paul Graduate Fellowship, Kramer Graduate Fellowship, Halberstadt Graduate Fellowship,

Anne Marie Brown Summer Graduate Fellowship, and Lab of Ornithology Summer Graduate Fellowship.

**Life-stage and sex influence *Philornis* ectoparasitism in a Neotropical woodpecker (*Melanerpes striatus*) with essential male parental care**

**Abstract**

The nestlings of many Neotropical bird species suffer from *Philornis* (Diptera: Muscidae) ectoparasitism. While nestlings are typically considered the intended targets, recent work indicates that *Philornis* infest adult birds more frequently than previously appreciated, yet few studies have concurrently surveyed nestlings and adults for *Philornis* in the same population. Over six field seasons (2012–2017), I documented the presence of current or recent subcutaneous *Philornis* infestations on adult and nestling Hispaniolan Woodpeckers (*Melanerpes striatus*) from the same population. I tested the following three non-mutually exclusive hypotheses regarding occurrence of *Philornis* on adult birds: (1) nestlings are more vulnerable to *Philornis* parasitism than adults, (2) nesting is associated with *Philornis* parasitism in adults, and (3) *Philornis* parasitism is associated with incubation and brooding investment. While nestling and adult woodpeckers exhibited similar prevalence of parasitism, parasitized nestlings hosted on average 3.5 times more *Philornis* wounds (larvae plus scars) than parasitized adults. Nesting *per se* was not significantly associated with parasitism among adults, as breeding and non-breeding adults showed similar prevalence and intensity. However, adult males, which perform overnight incubation and brooding, were significantly more likely to be parasitized than adult females. This last result supports the hypothesis that incubation and brooding investment increase the risk of *Philornis* parasitism for adults, but this conclusion is complicated by the lack of an association between parasitism and nesting status. Together, these results raise questions about the degree of host life-stage specialization and whether adult parasitism is incidental or part of an alternative parasitic strategy for *Philornis*.

**Resumen.** Los pichones de muchas especies de aves Neotropicales sufren de ectoparasitismo por *Philornis* (Diptera: Muscidae). Mientras que los pichones se consideran típicamente los objetivos previstos, trabajos recientes indican que *Philornis* infestan aves adultas con más frecuencia de lo que se

pensaba anteriormente, sin embargo, pocos estudios han examinado simultáneamente pichones y adultos

para *Philornis* en la misma población. Durante seis temporadas de campo (2012–2017), documenté la

presencia de infestaciones subcutáneas recientes o actuales de *Philornis* en adultos y pichones de la

misma población del pájaro carpintero *Melanerpes striatus*. Probé las siguientes tres hipótesis no

mutuamente excluyentes con respecto a la aparición de *Philornis* en aves adultas: (1) los pichones son

más vulnerables al parasitismo de *Philornis* que los adultos, (2) la anidación está asociada con el

parasitismo de *Philornis* en adultos, y (3) el parasitismo de *Philornis* es asociado con la incubación y la

inversión de crianza. Mientras que los pichones y los adultos exhibieron una prevalencia similar de

parasitismo, los pichones parasitados hospedaron en promedio 3.5 veces más heridas de *Philornis* (larvas

más cicatrices) que los adultos parasitados. La nidificación per se no se asoció significativamente con el

parasitismo entre los adultos, ya que los adultos reproductores y no reproductores mostraron una

prevalencia e intensidad similares. Sin embargo, los machos adultos, que realizan incubación y

empollando durante la noche, tenían una probabilidad significativamente mayor de ser parasitados que las

hembras adultas. Este último resultado apoya la hipótesis de que la inversión de incubación y de

empollando aumentan el riesgo de parasitismo por *Philornis* en adultos, pero esta conclusión se complica

por la falta de una asociación entre el parasitismo y el estado de anidación. Juntos, estos resultados

plantean preguntas sobre el grado de especialización de la etapa de vida del hospedador y si el parasitismo

de adultos es incidental o parte de una estrategia parasitaria alternativa para *Philornis*.

**Keywords**

Botflies, Adult birds, Myiasis, Caribbean, Dominican Republic, Picidae

**INTRODUCTION**

Nestlings of many bird species suffer from myiasis, "the infestation of healthy or necrotic tissue…by dipteran larvae" (Little 2009 p. 546:546), and in the Neotropics, *Philornis* (Diptera: Muscidae) botflies are the primary cause of healthy tissue myiasis (Teixeira 1999, Dudaniec & Kleindorfer 2006). The larvae of at least 23 *Philornis* species are subcutaneous, blood-feeding parasites (Common *et al.* 2019). Botfly effects on nestlings can be severe (reviewed in Dudaniec & Kleindorfer 2006), in some cases reducing survival (Delanoy & Cruz 1991, Rabuffetti & Reboreda 2007, Hayes *et al.* 2019). Indeed, native and introduced *Philornis* have been implicated in the decline of several island endemic birds, most notably in the Galápagos where introduced *P. downsi* have impacted many endemics (Fessl *et al.* 2006, Kleindorfer & Dudaniec 2016, Leuba *et al.* 2020). Yet the extent of *Philornis* infestation's ecological impacts remains poorly known, especially in these botflies' native ranges. For example, questions remain over the degree to which each species is a host specialist or generalist (Löwenberg-Neto 2008), what natural enemies impact *Philornis* (Knutie *et al.* 2017), and how anthropogenic activities like landscape change influence *Philornis*-host dynamics (Hayes *et al.* 2019). Answering these questions will be important for not only advancing basic ornithology but also for understanding whether to account for and how to control *Philornis* in conservation and management efforts.

Another aspect of *Philornis* parasitism that requires deeper exploration is the degree of host life stage specialization. The prevailing wisdom has been that *Philornis* target altricial and semi-altricial nestlings while the occasional observations of larvae on adult birds represent opportunistic infestation (Teixeira 1999). Some researchers have even posited that *Philornis* cannot successfully pupate once host birds have fledged (Arendt 1985a). Understanding the degree to which *Philornis* parasitizes nestlings and adults has important ramifications for bird populations since nestling parasitism directly impacts reproductive success while adult parasitism could impact survival and reproductive success. In a recent review of published records and analysis of new data from adult capture records from three Caribbean islands, Quiroga et al. (2020) reported adult parasitism for 15 bird species representing 12 families and four orders. While these results indicate that adult parasitism by *Philornis* might be more than

opportunistic, much remains unknown, and more precise estimates of adult infestation prevalence are

needed to clarify this relationship.

My objective here is to expand on the findings of Quiroga et al. (2020) by utilizing a species

well-suited for investigating *Philornis* parasitism: the Hispaniolan Woodpecker (*Melanerpes striatus*).

This woodpecker is one of the most abundant birds on Hispaniola, common from sea-level to 2,400 m asl

in a wide range of habitats (Latta *et al.* 2006), providing ample sampling opportunities. Additionally, the

first *Philornis* species (*P. pici*, reported as *Aricia pici*) was described from a subcutaneous larva collected

from an adult Hispaniolan Woodpecker (Macquart 1853). Despite the Hispaniolan Woodpecker's high

abundance, Quiroga et al. (2020) reported only two new records of *Philornis* infestation on adults: one

each from the Cordillera Central (prevalence = 20%, $N = 5$ individuals; H.M. Garrod, pers. comm.) and

Punta Cana (prevalence = 7%, $N = 14$ individuals; L. Soares and S.C. Latta, pers. comm.). Furthermore,

the parasite negatively impacts the reproductive success of at least one other Hispaniolan endemic, the

critically endangered Ridgway's Hawk (*Buteo ridgwayi*; Hayes *et al.* 2019). Yet the woodpecker's

continued abundance in spite of *Philornis* and anthropogenic pressures (Mitchell & Bruggers 1985)

suggests it could be an excellent model system to advance *Philornis* biology. To that end, I test three

hypotheses (Table 1) regarding *Philornis* infestation prevalence and intensity on adult birds.

First, I test two predictions of the hypothesis (H1) that nestlings are more vulnerable than adults

to *Philornis* parasitism (Teixeira 1999). This hypothesis predicts that (P1.1) *Philornis* prevalence (the

proportion of birds infested) should be higher for nestlings than for adults. Assuming nestlings are easier

targets for infestation, this hypothesis also predicts (P1.2) that nestlings should have higher intensity

(number of larvae per individual) *Philornis* infestations compared with adult birds. Due to their mobility,

adult woodpeckers should provide not only fewer opportunities for larval deposition by adult flies across

adults, but also fewer opportunities for repeat deposition on individual adults.

Second, I test two predictions of the hypothesis (H2) that nesting behavior itself is associated with

*Philornis* parasitism of adults. If *Philornis* is more prevalent and intense on nestlings than adults (Arendt

1985a), parasitism of adults might be an opportunistic direct result of nesting activity. This hypothesis

**Table 1.** Summary of hypotheses and predictions regarding *Philornis* infestation status.

| Hypothesis | Prediction |
|---|---|
| H1: Nestlings are more vulnerable than adults to *Philornis* parasitism. | P1.1: *Philornis* prevalence is higher for nestlings than for adults. P1.2: Infested nestlings host more *Philornis* larvae than adults. |
| H2: Nesting is associated with *Philornis* parasitism of adults. | P2.1: *Philornis* prevalence is higher in nesting birds than birds not nesting. P2.2: Nesting birds host more *Philornis* larvae than birds not nesting. |
| H3a: *Philornis* parasitism is associated with diurnal incubation and brooding investment. | P3.1a: *Philornis* prevalence is equal for nesting males and females, who share daytime incubation and brooding. P3.2a: Infested males and females host similar numbers of *Philornis* larvae. |
| H3b: *Philornis* parasitism is associated with total incubation and brooding investment. | P3.1b: *Philornis* prevalence is higher for males, the sex that conducts nocturnal incubation and brooding. P3.2b: Infested males host more *Philornis* larvae. |

thus predicts that *Philornis* (P2.1) prevalence and (P2.2) intensity should be higher for nesting birds than

birds not nesting.

Lastly, I test four predictions of the hypothesis (H3a,b) that adult *Philornis* parasitism is

associated with incubation and brooding investment (Teixeira 1999). While nesting itself might increase

exposure to *Philornis*, intersexual differences in breeding behavior might result in females and males

experiencing different levels of parasitism. Both female and male Hispaniolan Woodpeckers develop

brood patches and share approximately equivalent diurnal incubation and brooding (unpub. data). If

incubation and brooding behavior increase exposure (H3a), *Philornis* (P3.1a) prevalence and (P3.2a)

intensity should be similar in female and male Hispaniolan Woodpeckers. Like most woodpecker species

(Winkler *et al.* 1995), though, male Hispaniolan Woodpeckers perform all overnight incubation and

brooding (pers. obs.), a form of essential parental care. If overnight incubation and brooding increase

exposure (H3b), *Philornis* (P3.1b) prevalence and (P3.2b) intensity should be higher for nesting males

than nesting females.

Tests of these hypotheses and predictions (Table 1), which require data from both nestlings and

adults from the same population, have only been reported for the Caribbean endemic Pearly-eyed

Thrasher (*Margarops fuscatus*). Both Pearly-eyed Thrashers and Hispaniolan Woodpeckers nest in

cavities, a life-history trait that could impact parasitism exposure (Nilsson 1986), so one might predict

similar patterns of *Philornis* prevalence and intensity in both species. In support of H1, nestling Pearly-

eyed Thrashers exhibited a far higher prevalence (96%) and intensity (mean = 37 larvae/nestling) of *P.*

*deceptivus* compared with adult prevalence (31%) and intensity (mean = 3.1 larvae/adult) on Puerto Rico

(Arendt 1985a). To the best of my knowledge, H2 has not been directly tested in Pearly-eyed Thrashers

and has only indirect support from immunological data in the Galápagos endemic medium ground finch

(*Geospiza fortis*), which showed higher *Philornis*-specific antibody levels during nesting than pre-nesting

(Huber *et al.* 2010). Pearly-eyed Thrasher data support H3a since *Philornis* prevalence among nesting

females, which perform all incubation and brooding, was ~3.5 times higher than for nesting males

(Arendt 1985a). Indirect evidence supporting H3a was also found in the medium ground finch: nesting

females, who brood nestlings, had higher *Philornis*-specific antibody levels than nesting males (Huber *et*

*al.* 2010). However, no studies have investigated *Philornis* in a species where males perform essential

incubation and brooding.

**METHODS**

*Field methods***.** I studied Hispaniolan Woodpeckers in the community of Piedra Blanca, 3 km east of

Jarabacoa (19.1193ºN, 70.5819ºW; 550–700 m asl), La Vega, Dominican Republic, between April 2012

and July 2017. The site (~84 ha) comprised several private properties on a landscape of pine (*Pinus*

*occidentalis* and *P. caribaea*) and broadleaf wet forest fragments immersed in a matrix of cattle pastures

with isolated or clustered royal palms (*Roystonea hispaniolana*), small fragments of secondary vegetation,

and "living tree" (predominantly *Gliricidia sepium*) fences. This region experiences a mild, dry winter

(January - March), followed by a short wet spring season (April - May), a long, dry summer season (June

- September), and a short, wet fall season (October - December) coinciding with the latter half of the

Atlantic hurricane season (Climate-data.org 2021). For the remainder of the Methods, I use "we" in lieu

of "I" to describe most activities because they involved a team of tireless volunteer field assistants.

We evaluated *Philornis* infestation status on nestling and adult woodpeckers at trees monitored

for nesting activity, which we selected based on the presence of cavities and nesting activity (e.g., cavity

excavation, adults entering/exiting cavities, etc.). To determine nesting activity, we inspected cavities

using a penlight and small inspection mirror (1-2" diameter) while climbing or with a wireless camera

attached to a 15.2 m telescoping pole that broadcasted images to a portable digital television (Huebner &

Hurteau 2007, Waldstein 2012). Once we detected a nesting attempt (i.e., ≥1 eggs), we typically checked

the clutch every 3-5 days and, when possible, daily if we did not know the clutch completion date.

Incubation typically lasted 11 days (range = 9–14 days). The nestling sampling protocol differed slightly

in timing across years, but in general, sampling involved collecting morphometric measurements and

inspecting the entire body surface for the presence/absence of *Philornis*, including counting the number of

active wounds and scars (Fig. 1). We considered a wound active if it contained ≥1 subcutaneous *Philornis*

larvae, and, in cases where ≥2 larvae inhabited the same wound (see Fig. 1a for example of two sets of

posterior spiracles of larvae visible in a single wound), we recorded the total number of detectable larvae.

*Philornis* scars resembled active wounds in appearance, except that scars tended to look less swollen (Fig.

1b), lacked detectable larvae, and retained an opening where a larva had resided. For all years when we

did not know the nest's hatch date (e.g., nest was found with nestlings), we sampled and banded nestlings

as soon as they were large enough to carry four bands—two color bands on one leg and one color band

and one metal band on the other leg. For nests with known hatch dates from 2013-2015, we sampled and

fully banded nestlings when they were ~14 days old and resampled at ~21 days old. For nests with known

hatch dates in 2016 and 2017, we sampled and metal banded nestlings at ~7 days old, resampled at ~14

days old, and resampled and added three color bands at ~21 days old.

**Figure 1.** Example of active wounds and scars associated with *Philornis* parasitism in Hispaniolan Woodpeckers. (A) Active wounds containing three *Philornis* larvae (indicated by white arrows) on a nestling woodpecker's leg, where the posterior spiracles of larvae are visible. Two larvae inhabit one wound while one larva inhabits an adjacent wound. (B) *Philornis* scar on an adult male woodpecker's face.

196   For adult sampling, we captured birds via two approaches: (1) ambushing adults in nest cavities

197 and (2) an elevated, dual-tower mist-net system (LaPergola & Kenyon in prep.). Ambushing involved

198 setting up ambush traps as in Stanback and Koenig (1994) to allow pre-dawn capture of roosting birds.

199 See Garrod and LaPergola (2018) for more details on implementation. To reduce nest abandonment, we

200 used the ambush method ≥7 days before egg-laying or ≥22 days post-hatch. The mist-net tower system

201 involved erecting two 15.2 m tower poles supported with guy lines (ropes) and, using pulleys and ropes,

202 raising two stacked 12 m mist-nets in front of nesting trees. This method reduced disturbance at nests,

203 enabled capture of woodpeckers using trees too unstable to climb, increased sampling efficacy before

204 nesting, and increased sampling of non-nesting birds. As with nestlings, each adult received a unique

205 four-band combination and was inspected for the absence or presence of *Philornis*. When present, we

counted the number of *Philornis* active wounds and scars. We also recorded sex of adults based on crown

color, which is black for females and red for males.

We determined the nesting and breeder status of captured birds by monitoring nesting attempts

and identifying attendant birds via focal nest watches. Nest watches involved 2 or 3 hour sessions in

which an observer sat 15-20 m from a nest tree in a burlap blind, trained a 15X or 20X spotting scope on

a focal cavity entrance for a nest, and recorded the identities (i.e., band combinations) and behaviors of

woodpeckers that visited the nest. For testing the hypothesis that *Philornis* parasitism in adults is

associated with nesting, I coded adults as belonging to one of two categories: nesting or not nesting. I

counted an adult as nesting if it met two criteria: (1) we observed the bird incubating at or provisioning ≥1

nest within the year of capture, and (2) we captured the bird after the earliest possible clutch initiation

date for its earliest possible nesting attempt within the year of capture. I counted banded birds as not

nesting if they met one of the following criteria: (1) we captured the bird early in the field season before

most nests were initiated (between January and before early April), or (2) the bird was not associated with

a nesting attempt prior to the date of capture in the same calendar year.

Although we did not attempt to identify larvae to species, *Philornis pici* is the only *Philornis*

species currently known to infest birds on Hispaniola, and, as mentioned earlier, was first described from

the Hispaniolan Woodpecker (Macquart 1853). Elsewhere in the Dominican Republic, researchers have

confirmed this species to parasitize Ridgway's Hawk (Hayes *et al.* 2019, Quiroga *et al.* 2020). However,

*P. porteri* has also been identified parasitizing Ridgway's Hawk (M. A. Quiroga, pers. comm.). The

distribution of *P. porteri* on Hispaniola is currently unknown, but it is possible that the *Philornis* detected

in the present study could be *P. pici*, *P. porteri*, or both.

***Statistical analyses.*** See Table 1 for a summary of the hypotheses and their predictions. For testing the

hypothesis that nestlings are more vulnerable to *Philornis* parasitism than adults (H1), I tested the two

predictions with separate generalized linear mixed-effects models (GLMMs). For the prediction that the

probability of being parasitized is higher for nestlings than for adult birds (P1.1), I used a GLMM with a

binomial fit to test for an association between infestation status and age coded as a categorical fixed effect

(adult vs. nestling). Infestation status was treated as a binary response (0 = no evidence of *Philornis*, 1 =

presence of ≥1 *Philornis* larvae, scars associated with infestation, or both) in this model. For the

prediction that nestlings host greater numbers of *Philornis* larvae than adults (P1.2), I used a GLMM with

a Poisson distribution to test for an association between the total number of *Philornis* wounds (summing

the numbers of scars and active wounds, or total number of larvae) and age. Because many birds were

never observed with infestations, including all sampled individuals would lead to zero-inflation for the

total number of *Philornis* wounds; consequently, I used a manual hurdle model approach, including only

infested birds in this model.

To test the predictions of the hypotheses that (H2) nesting and (H3a, H3b) incubation and

brooding investment are associated with *Philornis* parasitism in adults, I used four GLMMs to test for

associations of adult infestation status with nesting status, sex, and the interaction of nesting status and

sex. I coded both nesting status (nesting vs. not nesting) and sex (female vs. male) as categorical fixed

effects for all four models. To test predictions regarding prevalence (P2.1, P3.1a, and P3.2a), I used

GLMMs with a binomial fit: infestation status was treated as a binary response as with the analysis

comparing nestlings and adults. For the first GLMM, I included all adults of known nesting status,

retaining birds known not to have bred within the year of capture. For the second GLMM, I retained only

birds known to have bred within the year of capture to exclude any effects of unanticipated differences

between breeders and non-breeders. For testing the predictions regarding intensity as they relate to

nesting status and sex (P2.2, P3.2a, and P3.2b), I used a GLMM with a Poisson distribution to test for an

association of the total number of *Philornis* wounds (summing the numbers of scars and active wounds,

or total number of larvae) with nesting status, sex, and the interaction effect of nesting status and sex. As

with the nestling-adult comparison, many observations involved no infestation and would lead to zero-

inflation for the total number of *Philornis* wounds; consequently, I used a manual hurdle model approach,

including only infested adults in this model. Additionally, I only ran this model with the dataset that

included all adults of known nesting status, including birds known not to have bred within the year of

capture.

259             For all models, I also included capture date as a continuous fixed effect based on the following.

The capture date range was fairly large (range = 168 d, 28-Feb – 4-Aug), which included the end of the

winter dry season, the short wet spring season, and the long dry summer season. Furthermore, previous

studies have documented a positive association between the probability of adults and nests having

*Philornis* and the timing of breeding (Arendt 1985a, Rabuffetti & Reboreda 2007). For all analyses, I

scaled capture date in day of year format via Z-transformation by subtracting the mean capture date and

dividing by the standard deviation.

266             For all models except for those testing predictions regarding intensity only in adults (P2.2, P3.2a,

and P3.2b), I included the following as random effects: the tree where a bird was captured or, for known

breeders, where it bred in the year of capture (Tree ID); year of capture; and individual ID. I included

Tree ID as a random effect because the Hispaniolan Woodpecker is unique among the Picidae, being one

of only three known woodpecker species to exhibit facultative colonial nesting. Within the same

population, Hispaniolan Woodpeckers pairs can nest singly or in clusters, with two or more pairs nesting

concurrently in separate cavities on the same tree (Short 1974, Winkler *et al.* 1995, LaPergola 2018).

Additionally, I wanted to account for the non-independence of nestlings from the same brood and thus the

same parents, but using a nest ID random effect would have precluded using adults without nests. Using

Tree ID as a random effect is thus a more conservative approach to account for non-independence,

especially for nestlings. I included year as a random effect in all analyses because I was not confident that

interannual variation was sampled adequately enough to interpret the fixed effects of year (Bennington &

Thayne 1994). Lastly, I included individual ID because some individuals were captured multiple times.

For testing predictions regarding intensity only in adults (P2.2, P3.2a, and P3.2b), I included only year as

a random effect because including Tree ID and individual ID led to failed model convergence.

281             I conducted all statistical analyses in RStudio v. 1.1.463 using R v. 3.6.3 (R Core Team 2020).

For fitting GLMMs, I used the *glmer* function in the *lme4* package (Bates *et al.* 2015). All means are

reported ± the standard error of the mean, and all confidence intervals for count data are 95% and were

calculated via the Wald Method.

**RESULTS**

**Summary of *Philornis* parasitism prevalence**

Over six years, I obtained 218 adult records representing 184 unique individuals (83 females and 101

males), which included 26 individuals (eight females and 18 males) recaptured once and four individuals

(one female and three males) recaptured twice. Of all adult records, 40 (18%; CI = 14–24%) included

individuals with evidence of *Philornis* parasitism. Of all individuals ($N$ = 184), 36 (20%; CI = 14–26%)

had evidence of *Philornis* parasitism, which included 24 (67%; CI = 50–80%; $N$ = 36 individuals) with

*Philornis* scars, nine (25%; CI = 14–41%) with active wounds, and three (8%; CI = 2–23%) with both

scars and active wounds. Of all adults with more than one capture ($N$ = 26), 11 individuals exhibited

changed infestation status (Table S1). These records included four individuals recaptured within the same

year, of which two had active wounds on the second capture but no wounds on the first, one had old

wounds on the first capture but not the second, and one individual had old wounds on the first capture but

no visible wounds on the second capture 82 days later.

Across six years, I collected 554 nestling records representing 381 individuals from 127 nesting

attempts. These figures amounted to a mean of 4.4 ± 2.4 records per nesting attempt (range = 1–10

records per nesting attempt) and a mean of 3.0 ± 1.0 nestlings per nesting attempt (range = 1–5 nestlings

per nesting attempt). Of all nestling records, 123 (22%; CI = 19–26%) showed evidence of *Philornis*

parasitism, and of all nestlings observed, 107 (28%; CI = 24–33%) exhibited evidence of *Philornis*

parasitism on at least one sampling event. Of the nestling individuals with evidence of *Philornis*, most

(73%; CI = 64–80%) involved active wounds (45 observations with only active wounds and 33

observations with both active and old wounds), while fewer observations involved only old wounds

(19%; CI = 12–27%; for 8% of nestling observations, the wound status was not recorded). Infested

nestlings came from 43 (34%; CI = 26–43%) of all monitored nesting attempts.

**Table 2.** Results of two generalized linear mixed-effects models testing for an association of *Philornis* parasitism with age in Hispaniolan Woodpeckers. Model (a) included the binary response of *Philornis* parasitism (yes/no) and fixed effects of age (adult vs. nestling), scaled date of capture (DOY scaled), and their interaction, and was fit with a binomial distribution. Model (b) tested for an association of total number of *Philornis* wounds on infested birds only with age, DOY scaled, and their interaction, and was fit with a Poisson distribution. Random effects for both models were individual identity (a: $N = 559$ individuals; b: $N = 143$ individuals), year of capture ($N = 6$ years in both a and b), and tree ID where captured or bred (a: $N = 41$ trees; b: $N = 25$ trees).

| Model and factors | Estimate ± S.E. | z-value | *P* |
|---|---|---|---|
| a. *Philornis* parasitism (yes/no) | | | |
| Intercept | -3.092 ± 0.971 | -3.184 | 0.00145 ** |
| Age | -0.689 ± 0.365 | -1.888 | 0.059 |
| DOY Scaled | 0.170 ± 0.205 | 0.829 | 0.407 |
| *Age * DOY Scaled* | *2.383 ± 0.438* | *5.443* | *5.25e-08 *** |
| b. Total *Philornis* wounds | | | |
| Intercept | 0.529 ± 0.231 | 2.288 | 0.022 * |
| *Age* | *0.961 ± 0.207* | *4.638* | *3.52e-06 *** |
| DOY scaled | -0.076 ± 0.116 | -0.656 | 0.512 |
| Age * DOY scaled | 0.158 ± 0.166 | 0.953 | 0.341 |

**H1: Comparison of adults and nestlings**

Using the full set of adult and nestling capture records, age and scaled day of year captured alone were not significant predictors of the presence/absence of *Philornis* parasitism (Table 2a). However, there was a significant interaction effect of age and scaled day of year captured, such that the probability of exhibiting *Philornis* parasitism increased with the scaled day of year for nestlings but not for adults (Fig. 2A). In contrast to presence/absence, age alone was significantly associated with the total number of *Philornis* wounds (active infestations plus scars) (Table 2b). Infested nestlings had an average of 7.1 ± 0.5

317      *Philornis* wounds (range = 1–39 *Philornis* wounds; *N* = 123 nestling records) while infested adults had an

318      average of only 2.0 ± 0.2 wounds (range = 1–5 *Philornis* wounds; *N* = 40 adult records; Fig. 2b).

**Figure 2.** Probability and intensity of *Philornis* parasitism on adult and nestling Hispaniolan

Woodpeckers. (A) Probability of *Philornis* parasitism plotted as raw data (adults = blue circles, nestlings

= gray triangles) and model predictions from a generalized linear mixed model testing for an association

with age, day of year captured, and their interaction. The blue solid line and black dashed line represent

model predictions for adults and nestlings, respectively. (B) Raincloud plot comparing adults and

nestlings for the total number of *Philornis* wounds observed on infested individuals (i.e., only non-zero

values for the total number of *Philornis* wounds). Sample sizes indicate the number of observations.

320      **H2 and H3: Nesting status and sex**

321      When restricting the analyses to adults of known nesting status, there was no significant association

322      between *Philornis* infestation and whether an adult was currently nesting (Table 3). This result was true

323      for both the analysis including all adults of known nesting status, i.e., retaining birds known not to have

324      bred within the capture year (Table 3a), and for the analysis restricted to only birds that nested within the

**Figure 3.** Proportion of female and male adult Hispaniolan Woodpeckers with ≥1 *Philornis* parasite. Error bars represent 95% confidence intervals. Sample sizes indicate the number of observations.

capture year (Table 3b). Additionally, the scaled day of year was not significantly associated with

infestation status. There was no significant interaction between sex and nesting status, but adult sex was

significantly associated with infestation in both analyses. For all adults of known nesting status, 8% of

females (CI = 4–16%; $N$ = 84 observations) and 27% of males (CI = 20–36%; $N$ = 111 observations)

showed signs of current or past infestation (Figure 3). These proportions remained unchanged for the

subset that included only birds that nested within the capture year (females: CI = 3–18%, $N$ = 61

observations; males: CI = 18–38%, $N$ = 75 observations).

Infested female and male adults had similar numbers of *Philornis* wounds. Infested females had a

mean of 2.0 ± 0.7 wounds (range = 1–5; $N$ = 7 observations), and infested males had a mean of 2.0 ± 0.2

(range = 1–5; $N$ = 29 observations). None of the fixed effects were significant in the model (Table 4).

**Table 3.** Results of two generalized linear mixed-effects models testing for an association of the binary response of *Philornis* parasitism (yes/no) of adult Hispaniolan Woodpeckers with sex, nesting status (actively nesting or not nesting at time of capture), the interaction of sex and nesting status, and the scaled day of the year captured (DOY scaled). Random effects were individual identity (a: $N = 163$ individuals; b: $N = 136$ individuals), tree ID where an individual was captured or bred (a: $N = 30$ trees; b: $N = 29$ trees), and year of capture ($N = 6$ years in both a and b). Model (a) included the full set of individuals with known nesting status within a year, including individuals that never bred (non-breeders) within the capture year ($N = 193$ observations). Model (b) included only individuals that were known to have bred within the capture year ($N = 136$ observations).

| Model and factors | Estimate ± S.E. | z-value | *P* |
|---|---|---|---|
| a. *Philornis* parasitism (yes/no) on breeders and non-breeders | | | |
| Intercept | -3.273 ± 0.973 | -3.362 | 0.0008 |
| *Sex* | *1.758 ± 0.715* | *2.459* | *0.014\** |
| Nesting Status | 0.585 ± 0.959 | 0.610 | 0.542 |
| Sex * Nesting Status | -0.145 ± 0.956 | -0.151 | 0.880 |
| DOY Scaled | 0.150 ± 0.310 | 0.484 | 0.628 |
| b. *Philornis* parasitism (yes/no) on breeders only | | | |
| Intercept | -3.455 ± 1.006 | -3.436 | 0.0006 |
| *Sex* | *1.888 ± 0.719* | *2.624* | *0.009\*\** |
| Nesting Status | 1.304 ± 1.204 | 1.083 | 0.279 |
| Sex * Nesting Status | -0.337 ± 1.203 | -0.280 | 0.779 |
| DOY Scaled | 0.406 ± 0.373 | 1.089 | 0.276 |

The 12 adults with active infestations were mostly (58%) known breeders (Table S2). Two of the three females with active infestations also had infested nestlings at the time of capture. Of the nine males with active infestations, four had infested nestlings at the time of capture and one male had fledged two young (one infested and one not) one month prior to capture. The remaining individuals of uncertain breeding status (one female and four males) were all caught within the known breeding season at the site (Figure S1); the earliest capture was on 8 March and the latest capture 20 July.

**Table 4.** Results of a generalized linear mixed-effect model with a Poisson distribution testing for an association between the total number of *Philornis* wounds on adult Hispaniolan Woodpeckers and sex, nesting status (actively nesting or not nesting at time of capture), the interaction of sex and nesting status, and scaled day of year captured (DOY scaled), including year as a random effect ($N = 4$ years). This analysis used birds of known nesting status ($N = 36$ observations), including birds that bred and those that did not within the year of capture.

| Factors | Estimate ± S.E. | z-value | *P* |
|---|---|---|---|
| Intercept | 0.022 ± 0.675 | 0.032 | 0.974 |
| Sex | 0.659 ± 0.660 | 0.998 | 0.318 |
| Nesting Status | 0.893 ± 0.834 | 1.070 | 0.284 |
| Sex * Nesting Status | -0.065 ± 0.242 | -0.267 | 0.789 |
| DOY Scaled | -0.834 ± 0.725 | -1.150 | 0.250 |

**DISCUSSION**

Most previous work on *Philornis* myiasis has understandably focused on these parasites' impacts on

nestling birds (Arendt 1985b, Dudaniec & Kleindorfer 2006, Hayes *et al.* 2019) since this life stage has

long been considered the primary target of parasitism (Teixeira 1999). The present study is one of very

few that has concurrently documented *Philornis* prevalence and intensity on nestlings and adults from the

same population (see also Arendt 1985a). Intriguingly, adult Hispaniolan Woodpeckers were just as likely

to exhibit evidence of *Philornis* as nestlings, although nestlings experienced greater intensity of *Philornis*.

For adults, while nesting status itself was not a significant predictor of being infested, males, which invest

more in overall incubation and brooding, were significantly more likely to have *Philornis* infestations

than females.

**Nestlings vs. adults**

The present study's results falsify the first prediction of the hypothesis that nestlings are more vulnerable

to *Philornis* parasitism but support the second prediction that nestlings experience more intense

infestations. While adult and nestling Hispaniolan Woodpeckers did not differ in the probability of being

parasitized (Table 2a), the probability of being parasitized for nestlings did increase with passage of the

breeding season yet remained more or less static for adults across the breeding season (Fig. 2a).

Furthermore, when infested, nestlings bore greater numbers of *Philornis* larvae than did adult birds (Table

2b; Fig. 2b.). This difference in intensity is likely due to the increased accessibility of nestlings to

*Philornis* flies in contrast to lower accessibility of adults. Of the two results and corresponding

predictions, the contrast in *Philornis* intensity supports the hypothesis that nestlings are indeed more

vulnerable to parasitism, but the similarity in prevalence suggests a complementary hypothesis that adult

*Philornis* are equally likely to find nestling and adult Hispaniolan Woodpeckers. One mechanistic

explanation for lower parasite intensity on adults is that their well-developed plumage reduces

accessibility by presenting a physical barrier to burrowing larvae (Oniki 1983). This might be especially

relevant for Hispaniolan Woodpeckers as they hatch naked and remain so until 7-8 d post-hatch when

their pin feathers typically begin erupting, and while pin break begins around 15 d post-hatch, these

feathers fail to cover most of the body other than the feather tracts until about 21 d post-hatch (unpub.

data). Thus the young have little to no physical barrier against *Philornis* other than a brooding adult for

roughly their first three weeks. Hispaniolan Woodpecker nestlings also remain in the nest 29–38 d post-

hatch (unpub. data), providing additional exposure time, albeit with an increasing amount of feather

coverage. A non-mutually exclusive alternative hypothesis is that the greater mobility of adults reduces

their accessibility to *Philornis* (Teixeira 1999). When not actively attending a nest, adults are literally

moving targets for flies, covering areas of 1.7–4.2 km$^2$ while foraging (Mitchell & Bruggers 1985),

whereas nestlings remain relatively stationary, confined to the same nest cavity until fledging. Whether

either or both mechanisms can explain differences in *Philornis* infestation intensity for Hispaniolan

Woodpeckers remains to be examined.

The patterns of *Philornis* prevalence and intensity reported here for Hispaniolan Woodpeckers

contrast somewhat with those from the only other study comparing nestlings and adults in the same

population. Whereas *Philornis* prevalence among nestling Pearly-eyed Thrashers was much higher than

for adults (96% vs. 31%, respectively; Arendt 1985a), prevalence was only non-significantly higher for

nestling woodpeckers than for adults (28% vs. 20%). This contrast might be explained by two inter-

related factors that differ between the Pearly-eyed Thrasher study and the present study: habitat and

climate. The thrasher study took place in a tropical rainforest, with annual rainfall averaging 4,460 mm

(Arendt 1985a), while the present study occurred in more open, drier habitat, with annual rainfall of 1,723

mm (Climate-data.org 2021). Rainfall is a significant predictor of *Philornis* infestation, showing a

positive correlation with intensity (Antoniazzi *et al.* 2011, Manzoli *et al.* 2013), and moisture and

humidity predict the geographic distribution of at least one *Philornis* species (Cuervo *et al.* 2021). The

greater canopy cover and humidity of the rainforest might have promoted larger populations of adult

*Philornis* than those in the drier habitat of the Dominican Republic, and these hypothetical larger fly

populations might have more fully exploited the vulnerable nestling thrashers while adults could

effectively avoid or prevent parasitism. Alternatively, the drier habitat on the Dominican Republic might

have reduced access for adult *Philornis* because they would have needed to cross open (i.e., no canopy

cover) habitat to reach woodpecker nests. In other words, the Hispaniolan Woodpecker's habitat structure

provides a barrier for adult *Philornis* so they are prevented from fully exploiting the vulnerable nestling

woodpeckers. The pattern of intensity differences was similar, though: both nestling thrashers and

woodpeckers had greater intensity of *Philornis* than adult birds, and this aligns with the hypothesis that

nestlings in both species are more vulnerable to *Philornis* parasitism.

One limitation of the present study was that the precise timing of active infestation for adults was

often unknown, especially relative to timing of nesting. This issue arose because most evidence of

*Philornis* on adult Hispaniolan Woodpeckers was in the form of scars rather than active wounds

containing larvae. In contrast, most nestling observations involved active wounds. This difference is due

in part to the sampling effort relative to the age of target birds. Nestling Hispaniolan Woodpeckers were

sampled at known ages and within 25 days of hatching so the period of exposure was limited, increasing

the probability of detecting subcutaneous *Philornis* larvae, which can remain attached for 5–8 days

(Arendt 1985a, Young 1993). The exposure period for adult birds bearing scars, however, was

presumably all the days they lived prior to the date of capture, decreasing the probability that we would

detect their wounds when they contained larvae. This limitation is important for two reasons regarding

timing. First, while it might be most parsimonious to assume all adult Hispaniolan Woodpeckers with

*Philornis* scars were infested as adults, we do not know the maximum number of days *Philornis* scars

persist after larval detachment from woodpeckers. This uncertainty means that some adults bearing scars

might have been infested as nestlings. Quiroga et al. (2020 p. 2:2) posited that all adults in their sample

were likely parasitized as adults because "scars usually heal ca. one week after larvae detach from the

host…". However, *Philornis* wounding as a nestling can cause beak and naris deformation that lasts into

adulthood in Galápagos finches (Galligan & Kleindorfer 2009, Kleindorfer *et al.* 2019), so considerable

variation in scar duration might exist among and within species. Second, uncertainty of adult exposure

reduces the accuracy of nestling-adult seasonality comparisons.

The difference between adults and nestling Hispaniolan Woodpeckers with respect to the

seasonality of prevalence begs further consideration. The lack of an effect of day of capture on prevalence

in adults might be related to the abovementioned limitation: i.e., sampling date relative to the day(s) of

active infestation. Because the majority of nestling observations involved active wounds while most adult

records only involved scars, the day of capture for adults was a less reliable indicator of the timing of

infestation for them. In other words, it could be that adults showed the same type of seasonality in

infestation as nestlings, with the probability of being infested increasing as the season progressed, but the

sampling effort precluded detecting such a pattern. If the difference in seasonality between adults and

nestlings was a real pattern, though, Hispaniolan Woodpeckers would differ from Pearly-eyed Thrashers,

in which adults showed increasing prevalence of *Philornis* as the season progressed but prevalence among

nestlings was high throughout the nesting season (Arendt 1985a).

**Nesting status, sex, and brooding/incubation investment**

The nesting status of Hispaniolan Woodpecker adults was not significantly associated with prevalence nor

intensity of *Philornis* parasitism (Tables 3 and 4), refuting the two predictions of the hypothesis that such

parasitism is associated with nesting. To the best of my knowledge, these results represent the first direct

test of this hypothesis. The lack of an effect of nesting status in the present study could be an artifact of

the sampling period, which mostly comprised the nesting season. However, the inclusion of adults known

to not be actively nesting at the time of capture should lessen the impact of such an artifact. Another

possible limitation was the uncertainty around the time when an adult was first infested because it makes

it harder to discern the amount of overlap between infestation and nesting. It will be crucial to more

precisely define the window of infestation for sampled adults to accurately compare prevalence and

intensity among nesting and non-nesting birds in future studies. One could achieve increased accuracy

here by sampling more birds in the non-breeding season and capturing more adults when they have chicks

of known age. However, it is worth considering whether the observed pattern is not an artifact, i.e.,

nesting and non-nesting Hispaniolan Woodpeckers are equally likely to be parasitized. If *Philornis*

typically finds hosts by searching for nest-related cues (e.g., olfactory), adult Hispaniolan Woodpeckers

might be parasitized outside the context of actively breeding if they spend time in nest cavities for other

activities. For example, Hispaniolan Woodpeckers roost in previously used nest cavities (personal

observation). If the cues adult *Philornis* use to find nestlings remain detectable, opportunistic parasitism

of adult woodpeckers could occur. Such a scenario might apply in the non-breeding season or even within

the breeding season prior to active nesting. One could test this idea experimentally by setting un-baited

traps for adult *Philornis* in old or recently used cavities. Another possible reason that nesting status might

be less relevant for Hispaniolan Woodpeckers concerns their habit of colonial nesting. For example,

adults lacking active nests might still be subjected to parasitism when one or more other colony members

are nesting and thus attracting adult *Philornis*. This hypothesis and the impacts of colonial nesting on

*Philornis* parasitism more broadly warrant further study since group-living can either increase (Brown &

Brown 1986) or decrease (Mooring & Hart 1992) the risk of parasitism.

The combined results of adult Hispaniolan Woodpeckers being parasitized regardless of nesting

status and nestlings and adults exhibiting similar prevalence suggest that parasitism of adult woodpeckers

might be part of a mixed strategy by *Philornis* in which they target adult birds. As suggested by Quiroga

et al. (2020), such a strategy might allow flies to reproduce when nestlings are unavailable or in short

supply. In the present study, the Hispaniolan Woodpecker population had a defined breeding season,

beginning in early March, peaking in May, and tapering off in August (LaPergola 2018) so nestling

woodpeckers are unavailable for approximately half the year and only abundant for roughly three months.

Some other local species that might host *Philornis* (e.g., *Crotophaga ani*, *Columbina passerina*, *Zenaida*

*aurita*, *Z. asiatica*, and *Coereba flaveola*) have been suggested to breed year-round (Latta *et al.* 2006), but

the extent to which they do so in addition to whether they are parasitized at the study site remains

unknown. Additionally, capture records of Hispaniolan Woodpeckers from the site are unavailable from

most of the non-breeding season, especially September through December. Fully testing this hypothesis

that *Philornis* target adult birds when nestlings are unavailable or scarce requires year-round monitoring

for infestation of both adults and nestlings and data on the availability and abundance of host nestlings.

Despite the non-significant effect of nesting status, there was some support for the hypothesis that

*Philornis* parasitism is associated with incubation and brooding investment in Hispaniolan Woodpeckers.

Although the sexes did not differ in intensity of infestation, males were 3.4 times more likely than

females to host *Philornis*. This result mirrors the pattern observed in Pearly-eyed Thrashers, where

females, who are the sole incubators/brooders, were 3.5 times more likely to host *Philornis* than males

(Arendt 1985a). Since females perform only diurnal incubation/brooding while males perform diurnal *and*

nocturnal incubation/brooding, males might experience increased *Philornis* exposure at night.

Unfortunately, almost nothing is known about the temporal activity patterns of *Philornis* on Hispaniola

nor for most other *Philornis* with subcutaneous larvae. We still do not know if *Philornis* on Hispaniola

are oviparous or larviparous. Recently, *P. torquans*, a species with subcutaneous larvae, was shown to be

oviparous in a lab study, though it remains unknown if they oviposit directly on hosts or on nesting

material in nature (Saravia-Pietropaolo *et al.* 2018). In the Galápagos, adult *P. downsi* enter host nests to

oviposit when the parent birds are absent during the day when nestlings are young and at night when

nestlings are older (O'Connor *et al.* 2010), but *P. downsi* larvae are free-living and hematophagous and

eggs are oviposited in the nest. An important assumption of the hypothesis that nocturnal incubation

increases exposure thus needs testing. Additionally, the lack of an interaction effect of nesting status and

sex suggests alternative hypotheses warrant testing.

Two major sets of alternative explanations for higher prevalence of *Philornis* among adult male

Hispaniolan Woodpeckers are sexual dimorphisms in behavior and morphology (Zuk & McKean 1996).

One behavioral difference could be that males experience greater exposure by spending more time in a

particular site or habitat (e.g., Tinsley 1989). For example, male Hispaniolan Woodpeckers might spend

more time than females in cavities overall, even when not tending a nest with eggs or young. This could

be the case if males played a larger role in defending cavities from competitors and were thus more likely

to encounter *Philornis* searching for nestlings. Another behavioral difference might be that males simply

invest less in anti-parasitic behaviors like preening and grooming such that they are less likely than

females to remove *Philornis*. At present, it is unknown whether male and female Hispaniolan

Woodpeckers differ in preening behavior. In at least some bird species, though, males tend to spend more

time grooming rather than less (Cotgreave & Clayton 1994, Oswald *et al.* 2019). With respect to

morphology, male Hispaniolan Woodpeckers have bills that are on average 25% longer than those of

females (unpub. data; see also Selander 1966), and it could be that their longer bills reduce their

effectiveness at removing *Philornis* eggs or larvae. Male Hispaniolan Woodpeckers are also larger in

other dimensions of size, including weight, and it could be that their larger size increases the probability

that they miss a parasite during preening. Both the alternative behavioral and morphological explanations

clearly warrant future study.

**Future considerations and implications for *Philornis* biology**

Inter-population comparisons of *Philornis* parasitism in Hispaniolan Woodpeckers could be a fruitful

course of future research since this woodpecker occupies a range of habitats and elevations. The *Philornis*

prevalence on adult woodpeckers documented in the present study was the same as that reported for the

species at nearby Rancho Baiguate (H. M. Garrod, pers. comm.; Quiroga *et al.* 2020) but higher than that

reported from coastal, low elevation Punta Cana (7%; L. Soares and S. C. Latta, pers. comm.; Quiroga *et*

*al.* 2020). Whether these differences correspond to *Philornis* population sizes differing according to

habitat or climatic conditions could be explored with the Hispaniolan Woodpecker.

Though the difference in nestling and adult *Philornis* infestation intensity is suggestive of a

preference for nestlings by botflies, more work is needed to robustly test this hypothesis. In studies of

choice and decision-making, confirming the presence of a positive association provides support for a

preference hypothesis, but a more discriminating test involves an experimental choice assay (Dougherty

2020). Similar approaches reveal host preferences in insects (e.g., Linn *et al.* 2003). In this case,

presenting adult *Philornis* with a choice test between depositing eggs on a nestling or adult bird would be

most revealing. Additionally, it is often assumed that parasitizing nestlings yields a higher fitness payoff,

but this hypothesis, as far as I know, remains untested.

Given the observed prevalence and intensity of *Philornis* on both nestlings and adults, the

Hispaniolan Woodpecker would make an excellent model system to study this parasite's biology. For

example, it would be revealing to conduct *Philornis* exclusion experiments to better understand how non-

nesting use of cavities impacts parasitism outside the breeding season or even for non-breeders during the

nesting season. The woodpecker's abundance would also facilitate testing alternative *Philornis*

management programs before using them with species of conservation concern.

**References**

**Antoniazzi, L.R., Manzoli, D.E., Rohrmann, D., Saravia, M.J., Silvestri, L. & Beldomenico, P.M.**

2011. Climate variability affects the impact of parasitic flies on Argentinean forest birds. *J. Zool.*

**283**: 126–134.

**Arendt, W.J.** 1985a. *Philornis* ectoparasitism of Pearly-eyed Thrashers. II. Effects on adults and

reproduction. *Auk* **102**: 281–292.

**Arendt, W.J.** 1985b. *Philornis* ectoparasitism of Pearly-eyed Thrashers. I. Impact on growth and

development of nestlings. *Auk* **102**: 270–280.

**Bates, D., Mächler, M., Bolker, B. & Walker, S.** 2015. Fitting Linear Mixed-Effects Models using

lme4. *J. Stat. Softw.* **67**: 1–48.

**Bennington, C.C. & Thayne, W. V**. 1994. Use and misuse of mixed model analysis of variance in

ecological studies. *Ecology* **75**: 717–722.

**Brown, C.R. & Brown, M.B.** 1986. Ectoparasitism as a cost of coloniality in Cliff Swallows (*Hirundo*

*pyrrhonota*). *Ecology* **67**: 1206–1218.

**Climate-data.org**. 2021. Climate Jarabacoa (Dominican Republic).

**Common, L.K., Dudaniec, R.Y., Colombelli-Négrel, D. & Kleindorfer, S.** 2019. Taxonomic shifts in

*Philornis* larval behaviour and rapid changes in *Philornis downsi* Dodge & Aitken (Diptera:

Muscidae): an invasive avian parasite on the Galápagos Islands. In: *Life Cycle and Development of*

*Diptera* (M. Sarwar, ed), pp. 1–22. IntechOpen, London, UK.

**Cotgreave, P. & Clayton, D.H.** 1994. Comparative analysis of time spent grooming by birds in relation

to parasite load. *Behaviour* **131**: 171–187.

**Cuervo, P.F., Percara, A., Monje, L., Beldomenico, P.M. & Quiroga, M.A.** 2021. Environmental

variables determining the distribution of an avian parasite: the case of the Philornis torquans

complex in South America. *Med. Vet. Entomol.* **35**: 284–292.

**Delanoy, C.A. & Cruz, A.** 1991. *Philornis* parasitism and nestling survival of the Puerto Rican Sharp-

Shinned Hawk. In: *Bird-parasite interactions* (J. E. Loye & M. Zuk, eds), pp. 93–103. Oxford

University Press, Oxford, UK.

**Dougherty, L.R.** 2020. Designing mate choice experiments. *Biol. Rev.* **95**: 759–781.

**Dudaniec, R.Y. & Kleindorfer, S.** 2006. Effects of the parasitic flies of the genus *Philornis* (Diptera:

Muscidae) on birds. *Emu* **106**: 13–20.

**Fessl, B., Sinclair, B.J. & Kleindorfer, S.** 2006. The life-cycle of *Philornis downsi* (Diptera: Muscidae)

parasitizing Darwin's finches and its impacts on nestling survival. *Parasitology* **133**: 739–747.

**Galligan, T.H. & Kleindorfer, S.** 2009. Naris and beak malformation caused by the parasitic fly,

Philornis downsi (Diptera: Muscidae), in Darwin's small ground finch, Geospiza fuliginosa

(Passeriformes: Emberizidae). *Biol. J. Linn. Soc.* **98**: 577–585.

**Garrod, H.M. & LaPergola, J.B.** 2018. Overlap in molt and breeding phenology in the Hispaniolan

Woodpecker (*Melanerpes striatus*). *Ornitol. Neotrop.* **Special Is**: 29–36.

**Hayes, C.D., Hayes, T.I., McClure, C.J.W., Quiroga, M., Thorstrom, R.K. & Anderson, D.L.** 2019.

Native parasitic nest fly impacts reproductive success of an island-endemic host. *Anim. Conserv.* **22**:

157–164.

**Huber, S.K., Owen, J.P., Koop, J.A.H., King, M.O., Grant, P.R., Grant, B.R. & Clayton, D.H.** 2010.

Ecoimmunity in Darwin's finches: Invasive parasites trigger acquired immunity in the medium

ground finch (Geospiza fortis). *PLoS One* **5**.

**Huebner, D.P. & Hurteau, S.R.** 2007. An economical wireless cavity-nest viewer. *J. F. Ornithol.* **78**:

87–92.

**Kleindorfer, S., Custance, G., Peters, K.J. & Sulloway, F.J.** 2019. Introduced parasite changes host

phenotype, mating signal and hybridization risk: *Philornis downsi* effects on Darwin's finch song.

*Proc. R. Soc. B Biol. Sci.* **286**: 20190461.

**Kleindorfer, S. & Dudaniec, R.Y.** 2016. Host-parasite ecology, behavior and genetics: a review of the

introduced fly parasite *Philornis downsi* and its Darwin's finch hosts. *BMC Zool.* **1**: 1.

**Knutie, S.A., Herman, J.M., Owen, J.P. & Clayton, D.H.** 2017. Tri-trophic ecology of native parasitic

nest flies of birds in Tobago. *Ecosphere* **8**.

**LaPergola, J.B.** 2018. Behavioral ecology of coloniality in the Hispaniolan Woodpecker. Cornell

University.

**Latta, S., Rimmer, C., Allan, K., Wiley, J., Raffaele, H., McFarland, K. & Fernandez, E.** 2006. *Birds*

*of the Dominican Republic and Haiti*. Princeton University Press, Princeton.

**Leuba, C., Tebbich, S., Nemeth, E., Anchundia, D., Heyer, E., Mosquera, D.A., Richner, H., Rojas**

**Allieri, M.L., Sevilla, C. & Fessl, B.** 2020. Effect of an introduced parasite in natural and

anthropogenic habitats on the breeding success of the endemic little vermilion flycatcher

*Pyrocephalus nanus* in the Galápagos. *J. Avian Biol.* **51**: 1–13.

**Linn, C., Feder, J.L., Nojima, S., Dambroski, H.R., Berlocher, S.H. & Roelofs, W.** 2003. Fruit odor

discrimination and sympatric host race formation in *Rhagoletis*. *Proc. Natl. Acad. Sci. U. S. A.* **100**: 11490–11493.

**Little, S.E.** 2009. Myiasis in Wild Birds. In: *Parasitic Diseases of Wild Birds* (C. T. Atkinson, N. J. Thomas, & D. B. Hunter, eds), pp. 546–556. Wiley-Blackwell, Oxford, UK.

**Löwenberg-Neto, P.** 2008. The structure of the parasite–host interactions between *Philornis* (Diptera: Muscidae) and neotropical birds. *J. Trop. Ecol.* **24**: 575–580.

**Macquart, J.** 1853. Notice sur une nouvelle espéce d'aricie, diptère de la tribu des anthomyzides. *Ann. la Société Entomol. Fr.* **Series 3,**: 567–661.

**Manzoli, D.E., Antoniazzi, L.R., Saravia, M.J., Silvestri, L., Rorhmann, D. & Beldomenico, P.M.** 2013. Multi-level determinants of parasitic fly infection in forest passerines. *PLoS One* **8**: e67104.

**Mitchell, B. & Bruggers, R.L.** 1985. Aspects of woodpecker damage to cacao in the Dominican Republic. *Trop. Pest Manag.* **31**: 148–152.

**Mooring, M.S. & Hart, B.L.** 1992. Animal grouping for protection from parasites: selfish herd and encounter-dilution effects. *Behaviour* **123**: 173–193.

**Nilsson, S.G.** 1986. Evolution of hole-nesting in birds: on balancing selection pressures. *Auk* **103**: 432–435.

**O'Connor, J.A., Robertson, J. & Kleindorfer, S.** 2010. Video analysis of host-parasite interactions in nests of Darwins finches. *Oryx* **44**: 588–594.

**Oniki, Y.** 1983. Notes on fly (Musicdae) parasitism of nestlings of South American birds. *Le Gerfaut* **73**: 281–286.

**Oswald, K.N., Smit, B., Lee, A.T.K. & Cunningham, S.J.** 2019. Behaviour of an alpine range-restricted species is described by interactions between microsite use and temperature. *Anim. Behav.* **157**: 177–187.

**Quiroga, M.A., Hayes, T.I., Hayes, C.D., Garrod, H., Soares, L., Knutie, S.A., Latta, S.C. & Anderson, D.L.** 2020. More than just nestlings: incidence of subcutaneous Philornis (Diptera: Muscidae) nest flies in adult birds. *Parasitol. Res.* **119**: 2337–2342.

**R_Core_Team**. 2020. R: A language and environment for statistical computing.

**Rabuffetti, F.L. & Reboreda, J.C.** 2007. Early infestation by bot flies (*Philornis seguyi*) decreases chick

survival and nesting success in chalk-browed mockingbirds (*Mimus saturninus*). *Auk* **124**: 898–906.

**Saravia-Pietropaolo, M.J., Arce, S.I., Manzoli, D.E., Quiroga, M. & Beldomenico, P.M.** 2018.

Aspects of the life cycle of the avian parasite *Philornis torquans* (Diptera: Muscidae) under

laboratory rearing conditions. *Can. Entomol.* **150**: 317–325.

**Selander, R.K.** 1966. Sexual dimorphism and differential niche utilization in birds. *Condor* **68**: 113–151.

**Short, L.L.** 1974. Habits of three endemic West Indian woodpeckers (Aves, Picidae). *Am. Museum Novit.*

**2549**: 1–44.

**Stanback, M.T. & Koenig, W.D.** 1994. Techniques for capturing birds inside natural cavities. *J. F.*

*Ornithol.* **65**: 70–75.

**Teixeira, D.M.** 1999. Myiasis caused by obligatory parasites. Ib. General observations on the biology of

the species of the genus Philornis Meinert, 1890 (Diptera, Muscidae). In: *Myiasis in man and*

*animals in the Neotropical region. Bibliographic database* (J. H. Guimarães & N. Papavero, eds),

pp. 71–96. Editora Pleiade, Sao Paulo, Brazil.

**Tinsley, R.C.** 1989. The effects of host sex on transmission success. *Parasitol. Today* **5**: 190–195.

**Waldstein, A.L.** 2012. An inexpensive camera system for monitoring cavity nests. *J. F. Ornithol.* **83**:

302–305.

**Winkler, H., Christie, D.A. & Nurney, D.** 1995. *Woodpeckers: an identification guide to the*

*woodpeckers of the world*. Houghton Mifflin, Boston, Massachusetts.

**Young, B.E.** 1993. Effects of the parasitic botfly *Philornis carinatus* on nestling House Wrens,

*Troglodytes aedon*, in Costa Rica. *Oecologia* **93**: 256–262.

**Zuk, M. & McKean, K.A.** 1996. Sex differences in parasite infections: patterns and processes. *Int. J.*

*Parasitol.* **26**: 1009–1024.

**SUPPORTING INFORMATION**

**Figure S1.** *Hispaniolan Woodpecker hatch date histogram with earliest and latest adult capture dates.*

**Table S1.** *Individual adult Hispaniolan Woodpeckers recapture records.*

**Table S2.** *Individual adult Hispaniolan Woodpeckers with active* Philornis *infestations.*