SARS-CoV-2 Omicron neutralization by therapeutic antibodies, convalescent sera, and post-mRNA vaccine booster

One Sentence Summary: Third dose of Pfizer/BioNTech COVID-19 vaccine significantly boosts neutralizing antibodies to the Omicron variant compared to a second dose, while neutralization of Omicron by convalescent sera, two-dose vaccine-elicited sera, or therapeutic antibodies is variable and often low.

Authors:

Sabrina Lusvarghi[1], Simon D. Pollett[2,3], Sabari Nath Neerukonda[1], Wei Wang[1], Richard Wang[1], Russell Vassell[1], Nusrat J. Epsi[2,3], Anthony C Fries[4], Brian K Agan[2,3], David A. Lindholm[5,6], Christopher J. Colombo[6,7], Rupal Mody[8], Evan C. Ewers[9], Tahaniyat Lalani[2,3,10], Anuradha Ganesan[2,3,11], Emilie Goguet[3,12], Monique Hollis-Perry[14], Si'Ana A. Coggins[3,12], Mark P. Simons[2], Leah C. Katzelnick[13], Gregory Wang[14,15], David R. Tribble[2], Lisa Bentley[16], Ann E. Eakin[17], Christopher C. Broder[12], Karl J. Erlandson[18], Eric D. Laing[12], Timothy H. Burgess[2], Edward Mitre[12], Carol D. Weiss[1†]

Affiliations:

[1] Division of Viral Products, Office of Vaccine Research and Review, Center for Biologics Evaluation and Research, U.S. Food and Drug Administration, Silver Spring, Maryland, USA

[2] Infectious Diseases Clinical Research Program, Department of Preventive Medicine and Biostatistics, Uniformed Services University of the Health Sciences, Bethesda, MD, USA

[3] Henry M. Jackson Foundation for the Advancement of Military Medicine, Inc., Bethesda, MD, USA

[4] U.S. Air Force School of Aerospace Medicine, Wright-Patterson, OH, USA

[5] Brooke Army Medical Center, Joint Base San Antonio-Fort Sam Houston, TX, USA

[6] Department of Medicine, Uniformed Services University of the Health Sciences, Bethesda, MD, USA

[7] Madigan Army Medical Center, Joint Base Lewis McChord, WA, USA

[8] William Beaumont Army Medical Center, El Paso, TX, USA

[9] Fort Belvoir Community Hospital, Fort Belvoir, VA, USA

[10] Naval Medical Center Portsmouth, Portsmouth, VA, USA

[11] Walter Reed National Military Medical Center, Bethesda, MD, USA

[12] Department of Microbiology and Immunology, Uniformed Services University of the Health Sciences,

Bethesda, MD, USA

[13] Viral Epidemiology and Immunity Unit, Laboratory of Infectious Diseases, National Institute of Allergy

and Infectious Diseases, National Institutes of Health, Bethesda, MD, USA

[14] Clinical Trials Center, Infectious Diseases Directorate, Naval Medical Research Center, Silver Spring,

MD, USA

[15] General Dynamics Information Technology, Falls Church, VA, USA

[16] Office of the Assistance Secretary for Preparedness and Response, U.S. Department of Human

Health and Services, Washington D.C., USA.

[17] Division of Microbiology and Infectious Diseases, National Institute of Allergy and Infectious

Disease, National Institutes of Health, Rockville, Maryland, USA.

[18] Influenza and Emerging Infectious Diseases Division, Biomedical Advanced Research and

Development Authority, U.S. Department of Health and Human Services, Washington, D.C.,

USA.

Abstract:

The rapid spread of the highly contagious Omicron variant of SARS-CoV-2 along with its high number of mutations in the spike gene has raised alarm about the effectiveness of current medical countermeasures. To address this concern, we measured neutralizing antibodies against Omicron in three important settings: (1) post-vaccination sera after two and three immunizations with the Pfizer/BNT162b2 vaccine, (2) convalescent sera from unvaccinated individuals infected by different variants, and (3) clinical-stage therapeutic antibodies. Using a pseudovirus neutralization assay, we found that titers against Omicron were low or undetectable after two immunizations and in most convalescent sera. A booster vaccination significantly increased titers against Omicron to levels comparable to those seen against the ancestral (D614G) variant after two immunizations. Neither age nor sex were associated with differences in post-vaccination antibody responses. Only three of 24 therapeutic antibodies tested retained their full potency against Omicron and high-level resistance was seen against fifteen. These findings underscore the potential benefit of booster mRNA vaccines for protection against Omicron and the need for additional therapeutic antibodies that are more robust to highly mutated variants.

Main Text:

INTRODUCTION

In November 2021 a new SARS-CoV-2 variant, named Omicron (Pango lineage B.1.1.529), was identified as a variant of concern (VOC). Its rapid spread in Africa and unusually high number of mutations, especially in the spike gene, has triggered intense international efforts to track the variant's spread and evaluate its effects on the potency of therapeutics and vaccines. The predominant strain of Omicron has mutations in the spike gene encoding 15 amino acid

changes in the receptor binding domain (RBD) of the spike surface protein (G339D, S371L,

S373P, S375F, K417N, N440K, G446S, S477N, T478K, E484A, Q493R, G496S, Q498R,

N501Y, and Y505H). The RBD mediates virus attachment to the ACE2 receptor on target cells

and is the principal target of neutralizing antibodies that contribute to protection against SARS-

CoV-2. Many of these RBD changes have been previously reported to reduce the effectiveness

of several therapeutic neutralizing antibodies (reviewed in Corti et al(*1*)). A recent study reports

that the full complement of RBD substitutions in the Omicron spike compromises the potency of

over 85% of 247 anti-RBD monoclonal antibodies (mAbs) tested(*2*). Preliminary reports indicate

substantial immune evasion to two-dose vaccine-elicited sera(*3-7*), booster-elicited sera(*8-16*),

genotype-varying convalescent sera(*3, 5, 6*), and several mAbs(*2, 6*). However, study

populations and methods vary widely among the studies to date, and many lack critical details

about host characteristics. Moreover, studies have not examined how host demography predicts

these neutralizing humoral responses, and examination of how infection by a broader diversity of

SARS-CoV-2 Delta and non-Delta genotypes is important for further insights into how genetic

diversity may correlate with cross-neutralizing antibody responses.

Here we used a pseudovirus neutralization assay(*16*) to measure antibody neutralization of

SARS-CoV-2 Omicron in three important contexts: (1) antibodies induced after two or three

doses of the Pfizer-BioNTech Covid-19 (Pfizer/BNT162b2 mRNA) vaccine, (2) antibodies

induced from infection by different SARS-CoV-2 variants and (3) therapeutic antibodies under

emergency use authorization (EUA) or in later stages of clinical development. We compared the

magnitude of neutralization escape by Omicron to D614G (referred to as wild type, WT) and

Delta SARS-CoV-2 variants to help inform public health decisions and offer further data toward

correlate of protection research.

RESULTS

Three immunizations of the Pfizer/BNT162b2 mRNA COVID-19 vaccine significantly boosts neutralizing antibodies to the Omicron variant compared to two-vaccinations.

The emergence of the Omicron variant coincided with recommendations for booster immunizations, particularly for at risk populations. We studied the neutralization titers of 39 generally healthy, adult healthcare workers participating in the Prospective Assessment of SARS-CoV-2 Seroconversion study (PASS study, Table 1)(*17*) who received the full primary series (1st and 2nd) and booster (3rd) immunizations with the Pfizer/BNT162b2 vaccine. We chose to study sera at peak responses after the full primary series vaccination rather than after 6 months because 6-months titers are often very low(*10, 18*).

Neutralization activity against Omicron above the threshold of our assay (1:40 dilution) was only observed in 12.8% (5/39) of serum samples obtained at a mean of 30 ± 11 days after the 2nd Pfizer/BNT162b2 vaccination (Figure 1A). After the 2nd vaccination neutralization titers against Omicron (geometric mean titer, GMT 22) were 25.5-fold lower than neutralization titers against WT (GMT 562). By contrast, neuralization titers against Delta (GMT 292) were 1.9-fold lower than WT. Neutralization titers from the same individuals collected 43 ± 17 days after the 3rd Pfizer/BNT162b2 vaccination were 8.9-fold greater against WT (GMT 5029) compared to titers after the 2nd vaccination. The titers against Omicron after the 3rd vaccination (GMT 700) showed 31.8-fold increase compared to titers after the 2nd vaccination, whereas titers against Delta after the 3rd vaccination (GMT 1673) were only 5.7-fold higher than the titers after the 2nd vaccination. Importantly, all individuals had measurable neutralizing titers against Omicron after the 3rd vaccination, highlighting the potential for increased protection by a booster vaccine.

We also evaluated how sex or age of the individual might affect the titers in sera after

vaccination. We did not observe a trend according to sex or age after the 2^{nd} or 3^{rd} vaccination

(Figure 1B and 1C, respectively). A total of 17/39 individuals had a positive anti-N (SARS-CoV-

2 nucleoprotein) seroconversion during regular scheduled blood sampling between the 2^{nd} and

the 3^{rd} immunization[17]. No symptoms of COVID-19 were reported by the subjects, possibly

indicative of silent infection or exposure to SARS-CoV-2 or other coronaviruses. We did not

find any trends when comparing differences in neutralizing antibody titers between individuals

who had anti-N seroconversion and those who did not (Figure 1D). To assess the breadth of

neutralization responses against Omicron induced by boosting, we compared the change in titers

against Omicron or Delta relative to WT after the 2^{nd} and 3^{rd} vaccination. To account for

variability in the antibody titers from one individual to another, we calculated the ratio between

the neutralization titers after the 3^{rd} and the 2^{nd} vaccination for each variant and plotted this ratio

for Omicron and Delta against WT (Figure 1E). We observed that the ratios of NT_{50} titer from

the 3^{rd} to the 2^{nd} vaccination for Omicron relative to the corresponding ratios for WT were higher

than the ratios for Delta relative to WT, suggesting that the 3^{rd} vaccination broadened responses

to the distantly-related variant.

Neutralization of Omicron by convalescent sera from individuals infected by different

variants shows lower neutralization titers compared to the infecting variant.

To investigate the potency of infection-induced neutralizing antibodies against Omicron, we

compared neutralization titers against WT, Delta, and Omicron in convalescent sera obtained at a

mean of 30.2 ± 9.3 days post-symptom onset from unvaccinated individuals infected with WT,

Alpha, Beta, or Delta variants who were enrolled in the Epidemiology, Immunology, and

Clinical Characteristics of Emerging Infectious Diseases with Pandemic Potential (EPICC)

study(*19*). Genotypes of the infecting variants were sequenced for all cases (Table 2 and

Supplementary Table). These convalescent sera were complemented by a Beta-convalescent

serum from another protocol.

Figure 2 shows that the Delta variant was modestly more resistant to neutralization than WT

by sera from most individuals infected with WT, Alpha, or Beta viruses (1.1 to 1.9-fold), while

neutralization titers were much more reduced against Omicron (2.3 to 70.1- fold). A total of 2/10

and 0/20 individuals infected with WT (B.1 and B.1.2) variants, respectively, had a response

above the threshold against Omicron, whereas 2/5 and 2/2 individuals infected with Alpha

(B.1.1.7) and Beta (B.1.351) variants, respectively, were above threshold. Convalescent sera

from individuals infected with Delta (B.1.617 and AY.14/.25/.44/.47/.62/.74/.119) variants

generally had higher neutralization titers against Delta (2.8 to 13.5-fold) compared to WT,

indicating more focused antibody responses to epitopes in the Delta spike. Convalescent sera

from these Delta-infected individuals showed 22.1 to 74.4-fold lower neutralization titers against

Omicron compared to Delta. However, a total of 6/7 individuals infected with the B.1.617.2

variant and 9/10 individuals infected with an AY variant had titers above threshold against

Omicron.

Boosting reduces apparent antigenic differences between WT and Omicron variants.

We applied antigenic cartography to explore how the convalescent and post-vaccination sera

distinguish the different spike antigens. Antigenic maps were made separately using neutralizing

antibody titers from individuals infected with the different variants or from individuals after the

2nd or 3rd vaccination (Figure 3). Convalescent sera were more heterogeneous compared to the

post-vaccination, with each convalescent serum clustering closer to the infecting variant, as

expected. The 3rd post-vaccination sera were more tightly clustered around WT compared to the

2nd post-vaccination sera. The antigenic distances between Omicron and WT were large for all

three sera sets, but the antigenic distance between Omicron and WT were smaller after the 3rd

vaccination (7.2-fold drop) compared to either convalescent or the 2nd vaccination sera (49.4-fold

drop and 39.4-fold drop, respectively), in agreement with the titer changes in Figure 1A. Small

distance changes between WT and Delta were observed for all three sera sets (2.0 to 3.6-fold

difference).

The potency of most therapeutic antibody products is compromised against Omicron.

As part of the US government COVID-19 response effort to speed development of

therapeutics for COVID-19, we assessed the neutralization of Omicron by 24 therapeutic

antibody products currently under EUA or in late stages of clinical development. This panel

includes 15 single therapeutic neutralizing antibody products (nAbs), six combination nAbs

(cnAbs) and three polyclonal antibody preparations (pAbs). The manufacturers provided these

clinical products for side-by-side comparisons of potency against variants but required blind

coding of these antibody products for publications. Previously, we reported that several single

substitutions in the spike protein of other variants conferred resistance to some of these

products(*20*), but a similar assessment has not been performed on the Omicron spike.

Figure 4A shows the neutralization curves for each product against WT and Omicron. To

quantify the relative drop in nAb potency we calculated ratios between the 50% inhibitory

concentration of Omicron to WT (Figure 4B). These ratios do not account for the absolute

potencies of the nAbs but do allow for a uniform comparison of the changes in potency against

Omicron relative to WT for all the antibody products. We note that the majority of successful

clinical trials were performed when predominant strains had a ratio of near 1 compared to WT.

However, absolute potency using IC_{50} ng/ml as a measure has not been established as a correlate

of protection. We defined ratios between of 5 to 50 as a benchmark for partial or moderate

resistance and ratios of > 50 as a benchmark for more complete resistance relative to WT. While

the clinical relevance of the IC_{50} changes has not been determined for any *in vitro* assay, these

cutoffs were chosen because the therapeutic levels of antibody therapeutics may be high enough

to overcome low levels of resistance. By the fold-change measure, only three of 15 nAbs

retained near full potency against Omicron compared to WT, and only one retained partial

potency. Two cnAbs retained partial potency, while the remaining four cnAbs showed complete

loss of neutralization potency. All three pAbs showed reduced neutralization potency (ratios 13-

17) against Omicron, in agreement with the data from convalescent and vaccinated individuals.

These findings raise concerns that many of the available therapeutic antibody products may not

be effective against Omicron.

DISCUSSION

Neutralizing antibodies are widely accepted as an important component of protection against

SARS-CoV-2 infection and disease (COVID-19), but efforts to assess antibody titers that

correlate with protection are complicated by many factors. These include potential redundancy

and synergism of different components of the humoral, cellular, and innate immune system, and

differences in variant fitness and host genetics, age, and prior immunity. The risk of infection can

also be confounded by human behavior and local public health measures, while measurements of

antibody titers can vary with different laboratory methods. Therefore, differences in study

populations and laboratory methods are important considerations for assessing the impact of

immune evasion by Omicron on medical countermeasures.

Here, using the same lentiviral pseudovirus neutralization platform we measured the change

in potency of 24 clinical-stage therapeutic antibodies against Omicron compared to WT (D614G)

and compared neutralizing antibodies in sera from two well-characterized cohorts of subjects in

prospective clinical studies. Our findings show that most vaccinated individuals have low or

undetectable titers against Omicron after the second Pfizer/BNT162b2 vaccination, similar to

findings reported by others(*3-7*). However, the third vaccination significantly increased

neutralizing titers to levels significantly higher than those elicited by the second vaccination, in

agreement with other preliminary studies(*8-14*). It is notable that the post-2nd vaccination and

post-3rd vaccination sampling times were similar, indicating that this boost does not simply

reflect time since last vaccine. We found no association between sex or age with these

neutralizing immune responses, although it is important to note that the study samples were from

generally healthy adults, and that the post-vaccine-dose sampling time was overall short (43 ± 17

days). It has been shown that infection followed by vaccination results in neutralizing antibody

titers comparable or higher to titers after two vaccinations(*3, 10, 14, 21*). The PASS study

included anti-N antibody testing on all blood samples for detection of silent infections. The

neutralizing antibody titers among the 17 asymptomatic individuals who seroconverted for N

antibodies between the 2nd and 3rd immunization were not higher than those who did not

seroconvert. Other studies have shown high antibody titers in convalescent individuals after the

1st or 2nd vaccination(*10, 11*). We did not see an increase after the 3rd vaccination in individuals

who seroconverted for N-antibodies. This may be due to reduced antigen load from incident

asymptomatic infection or having reached a maximum response after vaccination.

The antigenic cartography analysis suggests that the Omicron variant appears to be

antigenically distant from WT and Delta, but this distance seems to decrease after the 3rd

vaccination. The apparent broadening of responses to Omicron may be due to boosting of titers

to cross-reactive epitopes or improved antibody affinity to shared epitopes, or both. For the

convalescent sera, high titers generally correlated with the highest cross-neutralization to

Omicron. Continued studies of the breadth against multiple variants and duration of responses

after booster vaccinations are urgently needed. In both vaccination and convalescent individuals,

our studies suggest that booster vaccinations, even with the original ancestral vaccine antigen,

could be beneficial in protecting against Omicron, in agreement with the rapidly accumulating

data from many sources(*3, 5, 6*).

Lastly, nine of the fifteen clinical-grade nAbs under EUA or in late stages of clinical

development had no measurable IC_{50} against Omicron compared to WT. Also, while most nAbs

and cnAbs lost measurable potency, all polyclonal antibodies retained measurable, though

reduced with IC_{50}. Careful selection of therapeutic antibodies is needed according to variant

prevalence. As Omicron continues to acquire additional mutations the products that remain

effective could be jeopardized, underscoring the risk associated with this variant and its

derivatives.

In summary, our findings indicate that booster doses of mRNA COVID-19 vaccines may

afford an increase in protection against Omicron by inducing higher levels of neutralizing

antibodies compared to two vaccine doses or the levels of neutralizing antibodies induced by

SARS-CoV-2 infection from different variants. The strengths of this study include representation

of a broad diversity of genotypes, including within-Delta diversity, comparable sampling times

between convalescent and vaccinated subjects, the use of cartography methods, and the

availability of subject demographics to interpret how host characteristics may influence Omicron

humoral immunity. The limitations of our study include the relatively small numbers of study

subjects, restricted timing of sera collection, and the use of a pseudovirus platform as a surrogate

to authentic SARS-CoV-2 viruses. Ultimately neutralization titers need to be tied to clinical

outcomes. The rapid and unpredictable evolution of SARS-CoV-2 requires continued

development and assessments of medical countermeasures.

MATERIALS AND METHODS

Vaccinated participants

Details of the PASS study protocol, including the inclusion/exclusion criteria, have been

published(*17*). Full details are in the Supplemental Material. Briefly, generally healthy

healthcare workers without a history of SARS-CoV-2 infection at screening were enrolled. The

study began in August 2020 and involved monthly research clinic visits to obtain serum for

longitudinal SARS-CoV-2 antibody testing. The subset of participants included in this study

received three doses of Pfizer/BNT162b2 vaccine; none had a PCR-confirmed SARS-CoV-2

infection during follow-up. Participants' serially-collected serum samples were screened for

immunoglobulin G (IgG) reactivity with SARS-CoV-2 spike protein and nucleocapsid protein

(N) using a multiplex microsphere-based immunoassay, as described(*22*).

Unvaccinated infections - study population and general study design

The EPICC study is a cohort study of U.S. Military Health System (MHS) beneficiaries that

includes those with a history of SARS-CoV-2 infection, as described previously(*19*). Full details

are in Supplemental Material. Enrollment occurred at six Military Treatment Facilities (MTFs).

Demographic, comorbidity, COVID illness characteristics, and vaccination status were obtained

from the clinical case report form and/or the MHS Data Repository. Biospecimen collection

included serial serum samples and upper respiratory specimen swabs.

Diagnosis of SARS-CoV-2 infection and genotyping of infections used for convalescent sera

SARS-CoV-2 infection was determined by positive clinical laboratory PCR test performed at the

enrolling clinical site, or a follow-up upper respiratory swab collected as part of the EPICC

study. The specific clinical PCR assay employed at the MTF varied. The follow-up PCR assay

(for EPICC specimens) was the SARS-CoV-2 (2019-nCoV) CDC qPCR Probe Assay research

use only kits (IDT, Coralville, IA)(23); details are in the Supplemental Material. Whole viral

genome sequencing was performed on extracted SARS-CoV-2 RNA from PCR-positive

specimens; details are in the Supplemental Material. The Pango classification tool (https://cov-

lineages.org/) was used for genotype classification (version 3.1.17).

Ethics

The PASS (IDCRP-126) and EPICC (IDCRP-085) studies were approved by the Uniformed

Services University of the Health Sciences Institutional Review Board (IRB) in compliance with

all applicable Federal regulations governing the protection of human participants. All PASS and

EPICC study participants provided informed consent. The convalescent Beta sera, obtained from

a traveler who had moderate-severe Covid-19 in the Republic of South Africa during the peak of

the Beta (B.1.351) wave in January 2021, was obtained with informed consent and covered

under the US Food and Drug Administration IRB approved expedited protocol # 2021-CBER-

045.

SARS-Cov-2 pseudovirus production and neutralization assays

Lentiviral pseudoviruses were generated and used in neutralization assays, as previously

described(24). The Omicron spike expression plasmid was generously provided by Nicole Doria-

Rose (Vaccine Research Center, National Institutes of Health). Neutralization assays were

performed on 293T-ACE2/TMPRSS2 cells stably expressing ACE2 and transmembrane serine

protease 2 (BEI # NR-55293). Twenty-four clinical-stage therapeutic antibody products were

provided by pharmaceutical companies to support the US government COVID-19 response

efforts. The antibody identities are blinded for publications per an agreement with the

companies. Neutralization curves were normalized to virus only controls and fitted using

nonlinear regression curve (GraphPad Prism, La Jolla, CA). The antibody concentration or sera

dilution corresponding to 50% neutralization was defined as NT_{50} for sera or IC_{50} for antibody

products. Data reported are averages from at least two independent experiments, each with intra-

assay duplicates.

Antigenic cartography

ACMACS antigenic cartography software (https://acmacs-web.antigenic-441cartography.org/)

was used to create a geometric interpretation of neutralization titers against the WT, Delta, and

Omicron variants. Each square in the map indicates one antigenic unit, corresponding to two-fold

dilution of the antibody in the neutralization assay. Antigenic distance is measured in any

direction of the map.

Supplementary Materials

Supplementary materials and methods

Table S1: SARS-CoV-2 genotypes for the infecting variants

References and Notes

1. D. Corti, L. A. Purcell, G. Snell, D. Veesler, Tackling COVID-19 with neutralizing monoclonal antibodies.
Cell **184**, 3086-3108 (2021).

2. Y. Cao, J. Wang, F. Jian, T. Xiao, W. Song, A. Yisimayi, W. Huang, Q. Li, P. Wang, R. An, J. Wang, Y.
Wang, X. Niu, S. Yang, H. Liang, H. Sun, T. Li, Y. Yu, Q. Cui, S. Liu, X. Yang, S. Du, Z. Zhang, X. Hao,
F. Shao, R. Jin, X. Wang, J. Xiao, Y. Wang, X. S. Xie, B.1.1.529 escapes the majority of SARS-CoV-2
neutralizing antibodies of diverse epitopes. *bioRxiv*, 2021.2012.2007.470392 (2021).

3. A. Rössler, L. Riepler, D. Bante, D. v. Laer, J. Kimpel, SARS-CoV-2 B.1.1.529 variant (Omicron) evades
neutralization by sera from vaccinated and convalescent individuals. *medRxiv*, 2021.2012.2008.21267491
(2021).

4. L. Lu, B. W.-Y. Mok, L. Chen, J. M.-C. Chan, O. T.-Y. Tsang, B. H.-S. Lam, V. W.-M. Chuang, A. W.-H.
Chu, W.-M. Chan, J. D. Ip, B. P.-C. Chan, R. Zhang, C. C.-Y. Yip, V. C.-C. Cheng, K.-H. Chan, I. F.-N.
Hung, K.-Y. Yuen, H. Chen, K. K.-W. To, Neutralization of SARS-CoV-2 Omicron variant by sera from
BNT162b2 or Coronavac vaccine recipients. *medRxiv*, 2021.2012.2013.21267668 (2021).

5. N. Ikemura, A. Hoshino, Y. Higuchi, S. Taminishi, T. Inaba, S. Matoba, SARS-CoV-2 Omicron variant
escapes neutralization by vaccinated and convalescent sera and therapeutic monoclonal antibodies.
medRxiv, 2021.2012.2013.21267761 (2021).

6. E. Cameroni, C. Saliba, J. E. Bowen, L. E. Rosen, K. Culap, D. Pinto, A. De Marco, S. K. Zepeda, J. di
Iulio, F. Zatta, H. Kaiser, J. Noack, N. Farhat, N. Czudnochowski, C. Havenar-Daughton, K. R. Sprouse, J.
R. Dillen, A. E. Powell, A. Chen, C. Maher, L. Yin, D. Sun, L. Soriaga, C. Gustafsson, H. Chu, N. M.
Franko, J. Logue, N. T. Iqbal, I. Mazzitelli, J. Geffner, R. Grifantini, A. Gori, A. Riva, O. Giannini, A.
Ceschi, P. Ferrari, A. Franzetti-Pellanda, C. Garzoni, C. Hebner, L. Purcell, L. Piccoli, M. S. Pizzuto, A. C.
Walls, A. Telenti, H. W. Virgin, A. Lanzavecchia, D. Veesler, G. Snell, D. Corti, Broadly neutralizing
antibodies overcome SARS-CoV-2 Omicron antigenic shift. *bioRxiv*, 2021.2012.2012.472269 (2021).

7. S. Cele, L. Jackson, K. Khan, D. Khoury, T. Moyo-Gwete, H. Tegally, C. Scheepers, D. Amoako, F.
Karim, M. Bernstein, G. Lustig, D. Archary, M. Smith, Y. Ganga, Z. Jule, K. Reedoy, J. E. San, S.-H. Hwa,
J. Giandhari, J. M. Blackburn, B. I. Gosnell, S. A. Karim, W. Hanekom, NGS-SA, C.-K. Team, A. von
Gottberg, J. Bhiman, R. J. Lessells, M.-Y. S. Moosa, M. Davenport, T. de Oliveira, P. L. Moore, A. Sigal,

SARS-CoV-2 Omicron has extensive but incomplete escape of Pfizer BNT162b2 elicited neutralization

and requires ACE2 for infection. *medRxiv*, 2021.2012.2008.21267417 (2021).

8.   C. I. Kaku, E. R. Champney, J. Normark, C. E. Johnson, C. Ahlm, M. Sakharkar, M. E. Ackerman, M. N.

E. Forsell, L. M. Walker, Broad anti-SARS-CoV-2 antibody immunity induced by heterologous

ChAdOx1/mRNA-1273 prime-boost vaccination. *medRxiv*, 2021.2012.2013.21267598 (2021).

9.   W. F. Garcia-Beltran, K. J. St Denis, A. Hoelzemer, E. C. Lam, A. D. Nitido, M. L. Sheehan, C. Berrios, O.

Ofoman, C. C. Chang, B. M. Hauser, J. Feldman, D. J. Gregory, M. C. Poznansky, A. G. Schmidt, A. J.

Iafrate, V. Naranbhai, A. B. Balazs, mRNA-based COVID-19 vaccine boosters induce neutralizing

immunity against SARS-CoV-2 Omicron variant. *medRxiv*, 2021.2012.2014.21267755 (2021).

10.   H. Gruell, K. Vanshylla, P. Tober-Lau, D. Hillus, P. Schommers, C. Lehmann, F. Kurth, L. E. Sander, F.

Klein, mRNA booster immunization elicits potent neutralizing serum activity against the SARS-CoV-2

Omicron variant. *medRxiv*, 2021.2012.2014.21267769 (2021).

11.   F. Schmidt, F. Muecksch, Y. Weisblum, J. D. Silva, E. Bednarski, A. Cho, Z. Wang, C. Gaebler, M.

Caskey, M. C. Nussenzweig, T. Hatziioannou, P. D. Bieniasz, Plasma neutralization properties of the

SARS-CoV-2 Omicron variant. *medRxiv*, 2021.2012.2012.21267646 (2021).

12.   K. Basile, R. J. Rockett, K. McPhie, M. Fennell, J. Johnson-Mackinnon, J. E. Agius, W. Fong, H. Rahman,

D. Ko, L. Donavan, L. Hueston, C. Lam, A. Arnott, S. C.-A. Chen, S. Maddocks, M. V. O'Sullivan, D. E.

Dwyer, V. Sintchenko, J. Kok, Improved neutralization of the SARS-CoV-2 Omicron variant after Pfizer-

BioNTech BNT162b2 COVID-19 vaccine boosting. *bioRxiv*, 2021.2012.2012.472252 (2021).

13.   I. Nemet, L. Kliker, Y. Lustig, N. S. Zuckerman, O. Erster, C. Cohen, Y. Kreiss, S. Alroy-Preis, G. Regev-

Yochay, E. Mendelson, M. Mandelboim, Third BNT162b2 vaccination neutralization of SARS-CoV-2

Omicron infection. *medRxiv*, 2021.2012.2013.21267670 (2021).

14.   A. Wilhelm, M. Widera, K. Grikscheit, T. Toptan, B. Schenk, C. Pallas, M. Metzler, N. Kohmer, S. Hoehl,

F. A. Helfritz, T. Wolf, U. Goetsch, S. Ciesek, Reduced Neutralization of SARS-CoV-2 Omicron Variant

by Vaccine Sera and monoclonal antibodies. *medRxiv*, 2021.2012.2007.21267432 (2021).

15.   N. Doria-Rose, X. Shen, S. D. Schmidt, S. O'Dell, C. McDanal, W. Feng, J. Tong, A. Eaton, M. Maglinao,

H. Tang, R. L. Atmar, K. E. Lyke, L. Wang, y. Zhang, M. R. Gaudinski, W. P. Black, I. Gordon, M. Guech,

J. E. Ledgerwood, J. N. Misasi, A. Widge, P. C. Roberts, J. Beigel, B. Korber, R. Pajon, J. R. Mascola, D.

C. Montefiori, Booster of mRNA-1273 Vaccine Reduces SARS-CoV-2 Omicron Escape from Neutralizing

Antibodies. *medRxiv*, 2021.2012.2015.21267805 (2021).

16.   J. M. Carreño, H. Alshammary, J. Tcheou, G. Singh, A. Raskin, H. Kawabata, L. Sominsky, J. Clark, D. C.

Adelsberg, D. Bielak, A. S. Gonzalez-Reiche, P. P. S. Group, K. Srivastava, E. M. Sordillo, G. Bajic, H.

van Bakel, V. Simon, F. Krammer, Activity of convalescent and vaccine serum against a B.1.1.529 variant

SARS-CoV-2 isolate. *medRxiv*, 2021.2012.2020.21268134 (2021).

17.   B. M. Jackson-Thompson, E. Goguet, E. D. Laing, C. H. Olsen, S. Pollett, K. M. Hollis-Perry, S. E.

Maiolatesi, L. Illinik, K. F. Ramsey, A. E. Reyes, Y. Alcorta, M. A. Wong, J. Davies, O. Ortega, E.

Parmelee, A. R. Lindrose, M. Moser, E. Graydon, A. G. Letizia, C. A. Duplessis, A. Ganesan, K. P. Pratt,

A. M. Malloy, D. W. Scott, S. K. Anderson, A. L. Snow, C. L. Dalgard, J. H. Powers, 3rd, D. Tribble, T. H.

Burgess, C. C. Broder, E. Mitre, Prospective Assessment of SARS-CoV-2 Seroconversion (PASS) study:

an observational cohort study of SARS-CoV-2 infection and vaccination in healthcare workers. *BMC Infect*

Dis **21**, 544 (2021).

18.   E. D. Laing, C. D. Weiss, E. C. Samuels, S. A. A. Coggins, W. Wang, R. Wang, R. Vassell, S. L. Sterling,

M. A. Tso, T. Conner, E. Goguet, M. Moser, B. M. Jackson-Thompson, L. Illinik, J. Davies, O. Ortega, E.

Parmelee, M. Hollis-Perry, S. E. Maiolatesi, G. Wang, K. F. Ramsey, A. E. Reyes, Y. Alcorta, M. A.

Wong, A. R. Lindrose, C. A. Duplessis, D. R. Tribble, A. M. W. Malloy, T. H. Burgess, S. D. Pollett, C. H.

Olsen, C. C. Broder, E. Mitre, Durability of antibody responses and frequency of clinical and subclinical

SARS-CoV-2 infection six months after BNT162b2 COVID-19 vaccination in healthcare workers.

medRxiv, 2021.2010.2016.21265087 (2021).

19.   S. A. Richard, S. D. Pollett, C. A. Lanteri, E. V. Millar, A. C. Fries, R. C. Maves, G. C. Utz, T. Lalani, A.

Smith, R. M. Mody, A. Ganesan, R. E. Colombo, C. J. Colombo, D. A. Lindholm, C. Madar, S. Chi, N.

Huprikar, D. T. Larson, S. E. Bazan, C. English, E. Parmelee, K. Mende, E. D. Laing, C. C. Broder, P. W.

Blair, J. G. Chenoweth, M. P. Simons, D. R. Tribble, B. K. Agan, T. H. Burgess, E. C.-C. S. Group,

COVID-19 Outcomes Among US Military Health System Beneficiaries Include Complications Across

Multiple Organ Systems and Substantial Functional Impairment. *Open Forum Infect Dis* **8**, ofab556 (2021).

20. S. Lusvarghi, W. Wang, R. Herrup, S. N. Neerukonda, R. Vassell, L. Bentley, A. E. Eakin, K. J. Erlandson,

C. D. Weiss, Key substitutions in the spike protein of SARS-CoV-2 variants can predict resistance to

monoclonal antibodies, but other substitutions can modify the effects. *J Virol*, JVI0111021 (2021).

21. M. Schubert, F. Bertoglio, S. Steinke, P. A. Heine, M. A. Ynga-Durand, F. Zuo, L. Du, J. Korn, M.

Milošević, E. V. Wenzel, H. Maass, F. Krstanović, S. Polten, M. Pribanić-Matešić, I. Brizić, A. Piralla, F.

Baldanti, L. Hammarström, S. Dübel, A. Šustić, H. Marcotte, M. Strengert, A. Protić, Q. Pan

Hammarström, L. Čičin-Šain, M. Hust, Human serum from SARS-CoV-2 vaccinated and COVID-19

patients shows reduced binding to the RBD of SARS-CoV-2 Omicron variant in comparison to the original

Wuhan strain and the Beta and Delta variants. *medRxiv*, 2021.2012.2010.21267523 (2021).

22. E. D. Laing, S. L. Sterling, S. A. Richard, N. J. Epsi, S. Coggins, E. C. Samuels, S. Phogat, L. Yan, N.

Moreno, C. L. Coles, M. Drew, J. Mehalko, S. Merritt, K. Mende, V. Munster, E. de Wit, K. K. Chung, E.

V. Millar, D. R. Tribble, M. P. Simons, S. D. Pollett, D. Esposito, C. Lanteri, G. T. Clifton, E. Mitre, T. H.

Burgess, C. C. Broder, Antigen-based multiplex strategies to discriminate SARS-CoV-2 natural and

vaccine induced immunity from seasonal human coronavirus humoral responses. *medRxiv*, (2021).

23. S. Jangra, C. Ye, R. Rathnasinghe, D. Stadlbauer, g. Personalized Virology Initiative study, F. Krammer,

V. Simon, L. Martinez-Sobrido, A. Garcia-Sastre, M. Schotsaert, SARS-CoV-2 spike E484K mutation

reduces antibody neutralisation. *Lancet Microbe*, (2021).

24. S. N. Neerukonda, R. Vassell, S. Lusvarghi, R. Wang, F. Echegaray, L. Bentley, A. E. Eakin, K. J.

Erlandson, L. C. Katzelnick, C. D. Weiss, W. Wang, SARS-COV-2 Delta variant displays moderate

resistance to neutralizing antibodies and spike protein properties of higher soluble ACE2 sensitivity,

enhanced cleavage and fusogenic activity. *bioRxiv*, (2021).

Acknowledgments:

The authors wish to also acknowledge all who have contributed to the EPICC (IDCRP-085)

COVID-19 study.

Brooke Army Medical Center, Fort Sam Houston, TX: Col J. Cowden; LTC M. Darling; S.

DeLeon; Maj D. Lindholm; LTC A. Markelz; K. Mende; S. Merritt; T. Merritt; LTC N. Turner;

CPT T. Wellington

Carl R. Darnall Army Medical Center, Fort Hood, TX: LTC S. Bazan; P.K Love

Fort Belvoir Community Hospital, Fort Belvoir, VA: N. Dimascio-Johnson; MAJ E. Ewers;

LCDR K. Gallagher; LCDR D. Larson; A. Rutt

Henry M. Jackson Foundation, Inc., Bethesda, MD: P. Blair; J. Chenoweth; D. Clark

Madigan Army Medical Center, Joint Base Lewis McChord, WA: S. Chambers; LTC C.

Colombo; R. Colombo; CAPT C. Conlon; CAPT K. Everson; COL P. Faestel; COL T. Ferguson;

MAJ L. Gordon; LTC S. Grogan; CAPT S. Lis; COL C. Mount; LTC D. Musfeldt; CPT D.

Odineal; LTC M. Perreault; W. Robb-McGrath; MAJ R. Sainato; C. Schofield; COL C. Skinner;

M. Stein; MAJ M. Switzer; MAJ M. Timlin; MAJ S. Wood

Naval Medical Center Portsmouth, Portsmouth, VA: S. Banks; R. Carpenter; L. Kim; CAPT K.

Kronmann; T. Lalani; LCDR T. Lee; LCDR A. Smith; R. Smith; R. Tant; T. Warkentien

Naval Medical Center San Diego, San Diego, CA: CDR C. Berjohn; S. Cammarata; N. Kirkland;

D. Libraty; CAPT (Ret) R. Maves; CAPT (Ret) G. Utz

Tripler Army Medical Center, Honolulu, HI: S. Chi; LTC R. Flanagan; MAJ M. Jones; C. Lucas;

LTC (Ret) C. Madar; K. Miyasato; C. Uyehara

Uniformed Services University of the Health Sciences, Bethesda, MD: B. Agan; L. Andronescu;

A. Austin; C. Broder; CAPT T. Burgess; C. Byrne; COL K Chung; J. Davies; C. English; N.

Epsi; C. Fox; M. Fritschlanski; M. Grother; A. Hadley; COL P. Hickey; E. Laing; LTC C.

Lanteri; LTC J. Livezey; A. Malloy; R. Mohammed; C. Morales; P. Nwachukwu; C. Olsen; E.

Parmelee; S. Pollett; S. Richard; J. Rozman; J. Rusiecki; E. Samuels; M. Sanchez; A. Scher;

CDR M. Simons; A. Snow; K. Telu; D. Tribble; L. Ulomi

United States Air Force School of Medicine, Dayton, OH: SSgt T. Chao; R. Chapleau; M.

Christian; A. Fries; C. Harrington; S. Huntsberger; K. Lanter; E. Macias; J. Meyer; S. Purves; K.

Reynolds; J. Rodriguez; C. Starr

Womack Army Medical Center, Fort Bragg, NC: B. Barton; LTC D. Hostler; LTC J. Hostler;

MAJ K. Lago; C. Maldonado; J. Mehrer

United States Army Medical Research Institute of Infectious Diseases, Frederick, MD: MAJ J.

Kugelman

William Beaumont Army Medical Center, El Paso, TX: MAJ T. Hunter; J. Mejia; R. Mody; R.

Resendez; P. Sandoval; M. Wayman

Walter Reed National Military Medical Center, Bethesda, MD: I. Barahona; A. Baya; A.

Ganesan; MAJ N. Huprikar; B. Johnson

Walter Reed Army Institute of Research, Silver Spring, MD: S. Peel

We also thank Douglas Pratt and Hector Izurieta (US Food and Drug Administration) for critical

review of the manuscript.

Funding Statement:

This work was supported by awards from the Defense Health Program (HU00012020067,

HU00012120094, HU00012120104) and the National Institute of Allergy and Infectious Disease

(HU00011920111). The PASS and EPICC protocols were executed by the Infectious Disease

Clinical Research Program (IDCRP), a Department of Defense (DoD) program executed by the

Uniformed Services University of the Health Sciences (USUHS) through a cooperative

agreement by the Henry M. Jackson Foundation for the Advancement of Military Medicine, Inc.

(HJF). This project has also been funded in part by the National Institute of Allergy and

Infectious Diseases at the National Institutes of Health (NIAID/NIH), under an interagency

agreement with USUHS (Y1-AI-5072), and by the Intramural Research Program of NIAID.

Additional support was provided by institutional research funds from the US Food and Drug

Administration (FDA) and NIAID/NIH through an interagency agreement (AAI21013-001-

00000) with Center for Biologics Evaluation and Research, FDA, as part of the US Government

COVID-19 response efforts.

Author contributions:

Conceptualization: SL, SDP, RV, EG, CCB, LB, KJE, EDL, THB, EM, CDW

Methodology: SL, SDP, SNN, WW, RV, ACF, EG, SAC, CCB, LB, KJE, EDL, LCK, EM,

CDW

Investigation: SL, SDP, SNN, WW, RW, RV, NJE, ACF, DAL, CJC, RM, AG, EG, SAC, GW,

LB, KJE, ECE, TL, CDW

Visualization: SL, SDP, WW, RV, LCK, CDW

Funding acquisition: SDP, AEE, DRT, THB, EM, CDW

Project administration: SDP, RV, BKA, MH-P, MPS, DRT, AEE, KJE, EDL, THB, EM, CDW

Supervision: SDP, CCB, EDL, EM, THB, CDW

Writing – original draft: SL, SDP, CDW

Writing – review & editing: SL, SDP, SNN, WW, RW, RV, NJE, ACF, BKA, DAL, CJC, RM,

ECE, TL, AG, EG, MH-P, SAC, MPS, LCK, GW, CCB, DRT, LB, AEE, KJE, EDL, THB, EM,

CDW

Competing interests:

Potential conflicts of interest. S. D. P., T. H. B, and D.R.T. report that the Uniformed Services

University (USU) Infectious Diseases Clinical Research Program (IDCRP), a US Department of

Defense institution, and the Henry M. Jackson Foundation (HJF) were funded under a

Cooperative Research and Development Agreement to conduct an unrelated phase III COVID-19

monoclonal antibody immunoprophylaxis trial sponsored by AstraZeneca. The HJF, in support

of the USU IDCRP, was funded by the Department of Defense Joint Program Executive Office

for Chemical, Biological, Radiological, and Nuclear Defense to augment the conduct of an

unrelated phase III vaccine trial sponsored by AstraZeneca. Both of these trials were part of the

US Government COVID-19 response.

Government. The investigators have adhered to the policies for protection of human subjects as

prescribed in 45 CFR 46.

Group authors EPICC and PASS (MEDLINE indexed):

Brooke Army Medical Center, Fort Sam Houston, TX: Katrin Mende

Clinical Trials Center, Infectious Diseases Directorate, Naval Medical Research Center, Silver

Spring, MD, USA: Christopher Duplessis, Kathleen F. Ramsey, Anatalio E. Reyes, Yolanda

Alcorta, Mimi A. Wong, Santina Maiolatesi, Keishla Morales Padilla

Fort Belvoir Community Hospital, Fort Belvoir, VA: Derek Larson

Madigan Army Medical Center, Joint Base Lewis McChord, WA: Rhonda E. Colombo

Naval Medical Center Portsmouth, Portsmouth, VA: Alfred Smith

Uniformed Services University of the Health Sciences, Bethesda, MD: Stephanie Richard, Julia

Rozman, Alyssa R. Lindrose, Matthew Moser, Belinda Jackson-Thompson, Margaret Sanchez-

Edwards, Luca Illinik, Julian Davies, Orlando Ortega, Edward Parmelee

United States Army Medical Research Institute of Infectious Diseases, Frederick, MD: MAJ J.

Kugelman

Walter Reed National Military Medical Center, Bethesda, MD: Nikhil Huprikar

Fig. 1. Sensitivity of the Omicron variant to neutralization by Pfizer/BNT162b2 vaccinee

sera. (A) Neutralization assays were performed using lentiviral pseudoviruses bearing SARS-CoV-2 WT (B.1 lineage, D614G), Delta (Δ, B.1.617.2 lineage, T19R, G142D, E156-, F157-, R158G, L452R, T478K, D614G, P681R and D950N) and Omicron (O, B.1.1.529 lineage, A67V, del69-70, T95I, del142-144, Y145D, del211, L212I, ins214EPE, G339D, S371L, S373P, S375F, K417N, N440K, G446S, S477N, T478K, E484A, Q493R, G496S, Q498R, N501Y, Y505H, T547K, D614G, H655Y, N679K, P681H, N764K, D796Y, N856K, Q954H, N969K and L981F) with sera from 39 healthcare workers after immunization with two and three doses of Pfizer/BNT162b2 vaccine. Sera was obtained at a mean of 30 ± 11 days after immunization with two doses and at a mean of 43 ± 17 days after booster vaccination. Each serum was run in duplicate in two independent experiments against each pseudovirus to determine the 50% neutralization titer (NT_{50}). The geometric means titers (GMT), the number of NT_{50} above threshold (1:40) and the fold change are indicated. Titers below 1:40 were set at 20 to calculate GMTs. Arrows indicate increase or decrease relative to WT. Connecting lines indicate serum from the same individual. The demographic information for this sera cohort is provided in Table 1. (B) NT_{50} by sex after 2nd or 3rd vaccination. (C) NT_{50} by age after 2nd and 3rd vaccination. (D) 2nd or 3rd vaccination NT_{50} according to anti-N (nucleocapsid protein) seroconversion between 2nd and 3rd vaccination. (E) The ratios between the neutralization titers after the 3rd and the 2nd immunization for Omicron and Delta were plotted against the corresponding ratios for WT. For panels B-E, black squares correspond to WT, blue triangles correspond to Delta and red circles correspond to Omicron.

Fig. 2. Sensitivity of the Omicron variant to neutralization by convalescent sera

Neutralization assays were performed using convalescent sera from persons infected with

genotyped variants from B.1, B.1.2, B.1.1.7, B.1.351, B.1.617.2, AY.14, AY.25, AY.44, AY.47,

AY.62, AY.74 or AY.119 lineages (Table 2 and Supplementary Table). Both B.1 and B.1.2 have

no mutations in the receptor binding domain and were therefore considered WT, whereas some

of the AY mutants have additional mutations in the RBD relative to B.1.617.2. Each serum was

run in duplicate against WT, Delta, and Omicron to determine the NT_{50}. The geometric means

(GMT), the number of NT_{50} above threshold (1:40) and the fold change are indicated. Titers that

did not inhibit at the lowest dilution tested (1:40) were assigned a titer of 20 for GMT

calculations. Arrows indicate decrease relative to the infecting variant. Connecting lines indicate

serum from the same individual. Data shown represent two independent experiments each with

an intra-assay duplicate. Squares correspond to WT, triangles correspond to Delta, and circles

correspond to Omicron.

Fig. 3. Antigenic cartography of convalescent and vaccinee sera against WT, Delta and

Omicron. Antigenic maps were separately generated from convalescent (left panel), 2nd

vaccination (middle panel) or 3rd vaccination (right panel) sera. Convalescent sera are shown in

diamonds as follows: B.1 (dark green), B.1.2 (light green), B.1.1.7 (purple), B.1.351 (brown),

AY variants (light blue), and B.1.617.2 (dark blue). Gray diamonds correspond to post-

vaccination sera. Each grid square corresponds to 2-fold dilution in the neutralization assay.

Black squares correspond to WT variant. Blue triangles correspond to Delta variant. Red circle

corresponds to Omicron variant.

Fig. 4. Neutralization of Omicron by therapeutic antibodies. (A) Neutralization curves for

each one of the 24 therapeutic antibody products against WT (black) and Omicron (red). (B) Bar

graph showing the ratio between the IC_{50} of Omicron and WT for all the antibody products. The

sensitivity of the Omicron variant against 15 monoclonal antibodies (nAbs), 6 combination nAbs

products (cnAbs), and 3 polyclonal antibodies (pAbs). Red indicates IC_{50} resistance ratios >50,

yellow indicates moderate resistance with IC_{50} ratios between 5-50, and green indicates

sensitivity comparable to WT with IC_{50} ratios <5. Antibodies for which complete neutralization

was not achieved at the highest concentration tested are denoted by *. Data shown represent two

independent experiments each with an intra-assay duplicate.

616 **Table 1. Demographic data for participants receiving Pfizer/BNT162b2 initial vaccine**

617 **series and booster**

	N (%)
Sex	
Female	25 (64.1)
Male	14 (35.9)
Race	
White	26 (66.6%)
Asian	8 (20.5%)
Black	4 (10.3%)
Multiracial	1 (2.6%)
Occupation	
Nurse	11 (28.2%)
Physician	11 (28.2%)
Physical/Occupational/Recreational Therapist	9 (23.1%)
Medical Technician	3 (7.7%)
Lab Personnel	3 (7.7%)
Social Worker	1 (2.6%)
Psychologist	1 (2.6%)
Anti-N seroconversion after vaccination and before boost	
Positive	17 (43.6%)
Negative	22 (56.3%)

Age	
Mean age ± SD (range)	45 ± 11 (26 - 69)
Time between second vaccine and sample collection	
Mean days ± SD (range)	30 ± 11 (28 - 34)
Time between second vaccine and booster dose	
Mean days ± SD (range)	267 ± 14 (218-310)
Time between booster dose and sample collection	
Mean days +/- SD (range)	43 ± 17 (7 - 93)

Table 2: Characteristics of unvaccinated infections providing convalescent sera

	N = 39
Gender	
Female	14 (35.9%)
Male	25 (64.1%)
Race	
White	29 (74.4%)
Asian	1 (2.6%)
Black	6 (15.4%)
Multiracial	3 (7.7%)
Age	
Mean age ± SD (range)	41.1 ± 20 (1.4 - 73.2)
Charlson comorbidity index	
0	20 (51.3%)
1-2	10 (25.6%)
3-4	5 (12.8%)
>5	4 (10.3%)
Time between infection symptom onset and sample collection	
Mean days ± SD (range)	30.2 ± 9.3 (14.0 - 51.0)
Severity of initial infection	
Outpatient	23 (59.0%)
Hospitalized	16 (41.0%)
Infecting genotype[*]	

AY.119	1 (2.6%)
AY.14	2 (5.1%)
AY.25	3 (7.7%)
AY.44	1 (2.6%)
AY.47	1 (2.6%)
AY.62	1 (2.6%)
AY.74	1 (2.6%)
B.1	10 (25.6%)
B.1.1.7	5 (12.8%)
B.1.2	6 (15.4%)
B.1.351	1 (2.6%)
B.1.617.2	7 (17.9%)

620 [*] Genotypes assigned based on Pango 3.1.17 (2021-12-06). The genotype of the infecting variant

621 was determined in all cases except for one, a traveler who had moderate-severe Covid-19

622 (outpatient) in the Republic of South Africa during the peak of the Beta (B.1.351) wave in

623 January 2021 (FDA IRB Study # 2021-CBER-045).