# SARS-CoV-2 Omicron neutralization by therapeutic antibodies, convalescent sera, and post-mRNA vaccine booster

**Authors:**
Sabrina Lusvarghi[1], Simon D. Pollett[2,3], Sabari Nath Neerukonda[1], Wei Wang[1], Richard Wang[1], Russell Vassell[1], Nusrat J. Epsi[2,3], Anthony C Fries[4], Brian K Agan[2,3], David A. Lindholm[5,6], Christopher J. Colombo[6,7], Rupal Mody[8], Evan C. Ewers[9], Tahaniyat Lalani[2,3,10], Anuradha Ganesan[2,3,11], Emilie Goguet[3,12], Monique Hollis-Perry[14], Si'Ana A. Coggins[3,12], Mark P. Simons[2], Leah C. Katzelnick[13], Gregory Wang[14,15], Christopher C. Broder[12], David R. Tribble[2], Lisa Bentley[16], Ann E. Eakin[17], Karl J. Erlandson[18], Eric D. Laing[12], Timothy H. Burgess[2], Edward Mitre[12], Carol D. Weiss[1†]

**Affiliations:**
[1] Division of Viral Products, Office of Vaccine Research and Review, Center for Biologics Evaluation and Research, U.S. Food and Drug Administration, Silver Spring, Maryland, USA.
[2] Infectious Diseases Clinical Research Program, Department of Preventive Medicine and Biostatistics, Uniformed Services University of the Health Sciences, Bethesda, MD, USA
[3] Henry M. Jackson Foundation for the Advancement of Military Medicine, Inc., Bethesda, MD, USA
[4] U.S. Air Force School of Aerospace Medicine, Wright-Patterson, OH, USA
[5] Brooke Army Medical Center, Joint Base San Antonio-Fort Sam Houston, TX, USA
[6] Department of Medicine, Uniformed Services University of the Health Sciences, Bethesda, MD, USA
[7] Madigan Army Medical Center, Joint Base Lewis McChord, WA, USA
[8] William Beaumont Army Medical Center, El Paso, TX, USA
[9] Fort Belvoir Community Hospital, Fort Belvoir, VA, USA
[10] Naval Medical Center Portsmouth, Portsmouth, VA, USA
[11] Walter Reed National Military Medical Center, Bethesda, MD, USA
[12] Department of Microbiology and Immunology, Uniformed Services University of the Health Sciences, Bethesda, MD, USA
[13] Viral Epidemiology and Immunity Unit, Laboratory of Infectious Diseases, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, MD, USA
[14] Clinical Trials Center, Infectious Diseases Directorate, Naval Medical Research Center, Silver Spring, MD, USA
[15] General Dynamics Information Technology, Falls Church, VA, USA
[16] Office of the Assistance Secretary for Preparedness and Response, U.S. Department of Human Health and Services, Washington D.C., USA.
[17] Division of Microbiology and Infectious Diseases, National Institute of Allergy and Infectious Disease, National Institutes of Health, Rockville, Maryland, USA.
[18] Influenza and Emerging Infectious Diseases Division, Biomedical Advanced Research and Development Authority, U.S. Department of Health and Human Services, Washington, D.C., USA.

**Abstract:**

The rapid spread of the highly contagious Omicron variant of SARS-CoV-2 along with its high number of mutations in the spike gene has raised alarm about the effectiveness of current medical countermeasures. To address this concern, we measured neutralizing antibodies against Omicron in three important settings: (1) post-vaccination sera after two and three immunizations with the Pfizer/BNT162b2 vaccine, (2) convalescent sera from unvaccinated individuals infected by different variants, and (3) clinical-stage therapeutic antibodies. Using a pseudovirus neutralization assay, we found that titers against Omicron were low or undetectable after two immunizations and in most convalescent sera from individuals infected by different variants. A booster vaccination significantly increased titers against Omicron to levels comparable to those seen against the ancestral (D614G) variant after two immunizations. Neither age nor sex were associated with differences in post-vaccination antibody responses. Only three of 24 therapeutic antibodies tested retained their full potency against Omicron and high-level resistance was seen against fifteen. These findings underscore the potential benefit of booster mRNA vaccines for protection against Omicron and the need for additional therapeutic antibodies that are more robust to highly mutated variants.

## INTRODUCTION

In November 2021 a new SARS-CoV-2 variant, named Omicron (Pango lineage B.1.1.529), was identified as a variant of concern (VOC). Its rapid spread in Africa and unusually high number of mutations, especially in the spike gene, has triggered intense international efforts to track the variant's spread and evaluate its effects on the potency of therapeutics and vaccines. The predominant strain of Omicron has mutations in the spike gene encoding 15 amino acid changes in the receptor binding domain (RBD) of the spike surface protein (G339D, S371L, S373P, S375F, K417N, N440K, G446S, S477N, T478K, E484A, Q493R, G496S, Q498R, N501Y, and Y505H). The RBD mediates virus attachment to the ACE2 receptor on target cells and is the principal target of neutralizing antibodies that contribute to protection against SARS-CoV-2. Many of these RBD changes have been previously reported to reduce the effectiveness of several therapeutic neutralizing antibodies (reviewed in Corti et al(*1*)). A recent study reports that the full complement of RBD substitutions in the Omicron spike compromises the potency of over 85% of 247 anti-RBD monoclonal antibodies (mAbs) tested(*2*). Preliminary reports indicate substantial immune evasion to two-dose vaccine-elicited sera(*3-7*), booster-elicited sera(*8-15*), genotype-varying convalescent sera(*3, 5, 6*), and several mAbs(*2, 6*). However, study populations and methods vary widely among the studies to date, and many lack critical details about host characteristics. Moreover, studies have not examined how host demography predicts these neutralizing humoral responses, and examination of how infection by a broader diversity of SARS-CoV-2 Delta and non-Delta genotypes is important for further insights into how genetic diversity may correlate with cross-neutralizing antibody responses.

Here we used a pseudovirus neutralization assay(*16*) to measure antibody neutralization of SARS-CoV-2 Omicron in three important contexts: (1) antibodies induced after two or three doses of the Pfizer-BioNTech Covid-19 (Pfizer/BNT162b2 mRNA) vaccine, (2) antibodies

induced from infection by different SARS-CoV-2 variants and (3) therapeutic antibodies under emergency use authorization (EUA) or in later stages of clinical development. We compared the magnitude of neutralization escape by Omicron to D614G (referred to as wild type, WT) and Delta SARS-CoV-2 variants to help inform public health decisions and offer further data toward correlate of protection research.

## RESULTS

**Three immunizations of the Pfizer/BNT162b2 mRNA COVID-19 vaccine significantly boosts neutralizing antibodies to the Omicron variant compared to two-vaccinations.**
The emergence of the Omicron variant coincided with recommendations for booster immunizations, particularly for at risk populations. We studied the neutralization titers of 39 generally healthy, adult healthcare workers participating in the Prospective Assessment of SARS-CoV-2 Seroconversion study (PASS study, Table 1)(17) who received the full primary series (1st and 2nd) and booster (3rd) immunizations with the Pfizer/BNT162b2 vaccine. We chose to study sera at peak responses after the full primary series vaccination rather than after 6 months because 6-months titers are often very low(10, 18). Neutralization activity against Omicron above the threshold of our assay (1:40 dilution) was only observed in 12.8% (5/39) of serum samples obtained at a mean of $30 \pm 11$ days after the 2nd Pfizer/BNT162b2 vaccination (Figure 1A). After the 2nd vaccination neutralization titers against Omicron (geometric mean titer, GMT 22) were 25.5-fold lower than neutralization titers against WT (GMT 562). By contrast, neuralization titers against Delta (GMT 292) were 1.9-fold lower than WT. Neutralization titers from the same individuals collected $43 \pm 17$ days after the 3rd Pfizer/BNT162b2 vaccination were 8.9-fold greater against WT (GMT 5029) compared to titers after the 2nd vaccination. The titers against Omicron after the 3rd vaccination (GMT 700) showed 31.8-fold increase compared to titers after the 2nd vaccination, whereas titers against Delta after the 3rd vaccination (GMT 1673) were only 5.7-fold higher than the titers after the 2nd vaccination. Importantly, all individuals had measurable neutralizing titers against Omicron after the 3rd vaccination, highlighting the potential for increased protection by a booster vaccine.

We also evaluated how sex or age of the individual might affect the titers in sera after vaccination. We did not observe a trend according to sex or age after the 2nd or 3rd vaccination (Figure 1B and 1C, respectively). A total of 17/39 individuals had a positive anti-N (SARS-CoV-2 nucleoprotein) seroconversion during regular scheduled blood sampling between the 2nd and the 3rd immunization(17). No symptoms of COVID-19 were reported by the subjects, possibly indicative of silent infection or exposure to SARS-CoV-2 or other coronaviruses. We did not find any trends when comparing differences in neutralizing antibody titers between individuals who had anti-N seroconversion and those who did not (Figure 1D). To assess the breadth of neutralization responses against Omicron induced by boosting, we compared the change in titers against Omicron or Delta relative to WT after the 2nd and 3rd vaccination. To account for variability in the antibody titers from one individual to another, we calculated the ratio between the neutralization titers after the 3rd and the 2nd vaccination for each variant and plotted this ratio for Omicron and Delta against WT (Figure 1E). We observed that the ratios of $NT_{50}$ titer from the 3rd to the 2nd vaccination for Omicron relative to the corresponding ratios for WT were higher than the ratios for Delta relative to WT, suggesting that the 3rd vaccination broadened responses to the distantly-related variant.

**Neutralization of Omicron by convalescent sera from individuals infected by different variants shows lower neutralization titers compared to the infecting variant.**

To investigate the potency of infection-induced neutralizing antibodies against Omicron, we compared neutralization titers against WT, Delta, and Omicron in convalescent sera obtained at a mean of $30.2 \pm 9.3$ days post-symptom onset from unvaccinated individuals infected with WT, Alpha, Beta, or Delta variants who were enrolled in the Epidemiology, Immunology, and Clinical Characteristics of Emerging Infectious Diseases with Pandemic Potential (EPICC) study(19). Genotypes of the infecting variants were sequenced for all cases (Table 2 and Supplementary Table). These convalescent sera were complemented by a reference Beta-convalescent serum.

Figure 2 shows that the Delta variant was modestly more resistant to neutralization than WT by sera from most individuals infected with WT, Alpha, or Beta viruses (1.1 to 1.9-fold), while neutralization titers were much more reduced against Omicron (2.3 to 70.1- fold). A total of 2/10 and 0/20 individuals infected with WT (B.1 and B.1.2) variants, respectively, had a response above the threshold against Omicron, whereas 2/5 and 2/2 individuals infected with Alpha (B.1.1.7) and Beta (B.1.351) variants, respectively, were above threshold. Convalescent sera from individuals infected with Delta (B.1.617 and AY.14/.25/.44/.47/.62/.74/.119) variants generally had higher neutralization titers against Delta (2.8 to 13.5-fold) compared to WT, indicating more focused antibody responses to epitopes in the Delta spike. Convalescent sera from these Delta-infected individuals showed 22.1 to 74.4-fold lower neutralization titers against Omicron compared to Delta. However, a total of 6/7 individuals infected with the B.1.617.2 variant and 9/10 individuals infected with an AY variant had titers above threshold against Omicron.

**Boosting reduces apparent antigenic differences between WT and Omicron variants.**

We applied antigenic cartography to explore how the convalescent and post-vaccination sera distinguish the different spike antigens. Antigenic maps were made separately using neutralizing antibody titers from individuals infected with the different variants or from individuals after the $2^{nd}$ or $3^{rd}$ vaccination (Figure 3). Convalescent sera were more heterogeneous compared to the post-vaccination, with each convalescent serum clustering closer to the infecting variant, as expected. The $3^{rd}$ post-vaccination sera were more tightly clustered around WT compared to the $2^{nd}$ post-vaccination sera. The antigenic distances between Omicron and WT were large for all three sera sets, but the antigenic distance between Omicron and WT were smaller after the $3^{rd}$ vaccination (7.2-fold drop) compared to either convalescent or the $2^{nd}$ vaccination sera (49.4-fold drop and 39.4-fold drop, respectively), in agreement with the titer changes in Figure 1A. Small distance changes between WT and Delta were observed for all three sera sets (2.0 to 3.6-fold difference).

**The potency of most therapeutic antibody products is compromised against Omicron.**

As part of the US government COVID-19 response effort to speed development of therapeutics for COVID-19, we assessed the neutralization of Omicron by 24 therapeutic antibody products currently under EUA or in late stages of clinical development. This panel includes 15 single therapeutic neutralizing antibody products (nAbs), six combination nAbs (cnAbs) and three polyclonal antibody preparations (pAbs). The manufacturers provided these clinical products for

side-by-side comparisons of potency against variants but required blind coding of these antibody products for publications. Previously, we reported that several single substitutions in the spike protein of other variants conferred resistance to some of these products(20), but a similar assessment has not been performed on the Omicron spike.

Figure 4A shows the neutralization curves for each product against WT and Omicron. To quantify the relative drop in nAb potency we calculated ratios between the 50% inhibitory concentration of Omicron to WT (Figure 4B). These ratios do not account for the absolute potencies of the nAbs but do allow for a uniform comparison of the changes in potency against Omicron relative to WT for all the antibody products. We note that the majority of successful clinical trials were performed when predominant strains had a ratio of near 1 compared to WT. However, absolute potency using $IC_{50}$ ng/ml as a measure has not been established as a correlate of protection. We defined ratios between of 5 to 50 as a benchmark for partial or moderate resistance and ratios of $> 50$ as a benchmark for more complete resistance relative to WT. While the clinical relevance of the $IC_{50}$ changes has not been determined for any *in vitro* assay, these cutoffs were chosen because the therapeutic levels of antibody therapeutics may be high enough to overcome low levels of resistance. By the fold-change measure, only three of 15 nAbs retained near full potency against Omicron compared to WT, and only one retained partial potency. Two cnAbs retained partial potency, while the remaining four cnAbs showed complete loss of neutralization potency. All three pAbs showed reduced neutralization potency (ratios 13-17) against Omicron, in agreement with the data from convalescent and vaccinated individuals. These findings raise concerns that many of the available therapeutic antibody products may not be effective against Omicron.

## DISCUSSION

Neutralizing antibodies are widely accepted as an important component of protection against SARS-CoV-2 infection and disease (COVID-19), but efforts to assess antibody titers that correlate with protection are complicated by many factors. These include potential redundancy and synergism of different components of the humoral, cellular, and innate immune system, and differences in variant fitness and host genetics, age, and prior immunity. The risk of infection can also be confounded by human behavior and local public health measures, while measurements of antibody titers can vary with different laboratory methods. Therefore, differences in study populations and laboratory methods are important considerations for assessing the impact of immune evasion by Omicron on medical countermeasures.

Here, using the same lentiviral pseudovirus neutralization platform we measured the change in potency of 24 clinical-stage therapeutic antibodies against Omicron compared to WT (D614G) and compared neutralizing antibodies in sera from two well-characterized cohorts of subjects in prospective clinical studies. Our findings show that most vaccinated individuals have low or undetectable titers against Omicron after the second Pfizer/BNT162b2 vaccination, similar to findings reported by others(3-7). However, the third vaccination significantly increased neutralizing titers to levels significantly higher than those elicited by the second vaccination, in agreement with other preliminary studies(8-14). It is notable that the post-2[nd] vaccination and post-3[rd] vaccination sampling times were similar, indicating that this boost does not simply

reflect time since last vaccine. We found no association between sex or age with these neutralizing immune responses, although it is important to note that the study samples were from generally healthy adults, and that the post-vaccine-dose sampling time was overall short ($43 \pm 17$ days). It has been shown that infection followed by vaccination results in neutralizing antibody titers comparable or higher to titers after two vaccinations(*3, 10, 14, 21*). The PASS study included anti-N antibody testing on all blood samples for detection of silent infections. The neutralizing antibody titers among the 17 asymptomatic individuals who seroconverted for N antibodies between the 2[nd] and 3[rd] immunization were not higher than those who did not seroconvert. Other studies have shown high antibody titers in convalescent individuals after the 1[st] or 2[nd] vaccination(*10, 11*). We do not see an increase after the 3[rd] vaccination in individuals who seroconverted for N-antibodies. This may be due to reduced antigen load from incident asymptomatic infection or having reached a maximum response after vaccination.

The antigenic cartography analysis suggests that the Omicron variant appears to be antigenically distant from WT and Delta, but this distance seems to decrease after the 3[rd] vaccination. The apparent broadening of responses to Omicron may be due to boosting of titers to cross-reactive epitopes or improved antibody affinity to shared epitopes, or both. For the convalescent sera, high titers generally correlated with the highest cross-neutralization to Omicron. Continued studies of the breadth against multiple variants and duration of responses after booster vaccinations are urgently needed. In both vaccination and convalescent individuals, our studies suggest that booster vaccinations, even with the original ancestral vaccine antigen, could be beneficial in protecting against Omicron, in agreement with the rapidly accumulating data from many sources(*3, 5, 6*).

Lastly, nine of the fifteen clinical-grade nAbs under EUA or in late stages of clinical development had no measurable $IC_{50}$ against Omicron compared to WT. Also, while most nAbs and cnAbs lost measurable potency, all polyclonal antibodies retained measurable, though reduced with $IC_{50}$. Careful selection of therapeutic antibodies is needed according to variant prevalence. As Omicron continues to acquire additional mutations the products that remain effective could be jeopardized, underscoring the risk associated with this variant and its derivatives.

In summary, our findings indicate that booster doses of mRNA COVID-19 vaccines may afford an increase in protection against Omicron by inducing higher levels of neutralizing antibodies compared to two vaccine doses or the levels of neutralizing antibodies induced by SARS-CoV-2 infection from different variants. The strengths of this study include representation of a broad diversity of genotypes, including within-Delta diversity, comparable sampling times between convalescent and vaccinated subjects, the use of cartography methods, and the availability of subject demographics to interpret how host characteristics may influence Omicron humoral immunity.  The limitations of our study include the relatively small numbers of study subjects, restricted timing of sera collection, and the use of a pseudovirus platform as a surrogate to authentic SARS-CoV-2 viruses. Ultimately neutralization titers need to be tied to clinical outcomes. The rapid and unpredictable evolution of SARS-CoV-2 requires continued development and assessments of medical countermeasures.

**MATERIALS AND METHODS**

**Vaccinated participants**
Details of the PASS study protocol, including details of the inclusion/exclusion criteria, have been published(17). Inclusion criteria included being generally healthy, ≥ 18 years old, and employed at the WRNMMC. Exclusion criteria included history of COVID-19, IgG seropositivity for SARS-CoV-2, and being severely immunocompromised at time of screening. The study was initiated in August 2020, with rolling enrollment and monthly research clinic visits to obtain serum for longitudinal SARS-CoV-2 antibody testing. The subset of participants included in this study received two doses of Pfizer/BNT162b2 vaccine by January 26, 2021 and had no serological or PCR evidence of SARS-CoV-2 infection prior to two doses of vaccine. These subjects received a $3^{rd}$ dose of Pfizer/BNTech162b2 vaccine by Nov 18, 2021. No subject included in this analysis had a clinically apparent PCR-confirmed SARS-CoV-2 infection during follow-up. Participants' serum samples were collected monthly through September of 2021, and then quarterly, diluted 1:400 and 1:8000, and screened for immunoglobulin G (IgG) reactivity with SARS-CoV-2 spike protein and nucleocapsid protein (N), and four human coronavirus (HCoV) spike proteins using a multiplex microsphere-based immunoassay, as previously described (. For this study sample, we selected individuals with sera available at 3 to 6 weeks after symptom onset, with complete or near complete and high coverage spike gene sequence availability, with no prior vaccination, and with a diverse set of genotypes including Delta and non-Delta variants.

**Unvaccinated infections - study population and general study design**
The Epidemiology, Immunology, and Clinical Characteristics of Emerging Infectious Diseases with Pandemic Potential (EPICC) study is a cohort study of U.S. Military Health System (MHS) beneficiaries which includes those with a history of SARS-CoV-2 infection(19). Eligibility criteria for enrollment included those presenting to clinical care with COVID-19-like illness and for SARS-CoV-2 polymerase chain reaction (PCR) testing. The EPICC study has been enrolling since March 2020. For this sample set, EPICC enrollment occurred at six Military Treatment Facilities (MTFs): Brooke Army Medical Center, Fort Belvoir Community Hospital, Madigan Army Medical Center, Naval Medical Center Portsmouth, Walter Reed National Military Medical Center, and the William Beaumont Army Medical Center.

Study procedures for these subjects with SARS-CoV-2 infection included collection of demographic data and completion of a clinical case report form (CRF) to characterize the acute COVID-19 illness. Biospecimen collection included serial serum samples for immune response analysis, and upper respiratory specimen swabs for virological analysis. For all enrolled subjects, we also abstracted MHS-wide healthcare encounter data from the Military Health System Data Repository (MDR) to determine comorbidities. Vaccination status was ascertained by the MDR record in addition to the CRF and questionnaire self-report.

**Diagnosis of SARS-CoV-2 infection and genotyping of infections used for convalescent sera**
SARS-CoV-2 infection was determined by positive PCR clinical laboratory test performed at the enrolling clinical site, or a follow-up upper respiratory swab collected as part of the EPICC study. The specific PCR assay employed at the MTF varied. The follow-up PCR assay (used for EPICC specimens) was the SARS-CoV-2 (2019-nCoV) CDC qPCR Probe Assay research use

only kits (IDT, Coralville, IA)(*22*). This assay targets two regions of the SARS-CoV-2 nucleocapsid (N) gene (N1 and N2), with an additional control target to detect the human RNase P gene (RP). We considered a positive SARS-CoV-2 infection as positive based on a cycle threshold value of less than 40 for both N1/N2 gene targets.

Whole viral genome sequencing was performed on extracted SARS-CoV-2 RNA from PCR positive specimens.  A 1200bp amplicon tiling strategy was used (dx.doi.org/10.17504/protocols.io.btsrnnd6). Amplified product was prepared for sequencing using NexteraXT library kits (Illumina Inc., San Diego, CA).  Libraries were run on the Illumina NextSeq 550 sequencing platform. BBMap v. 38.86 and iVar v. 1.2.2 tools were used for genome assembly. The Pango classification tool ([https://cov-lineages.org/](https://cov-lineages.org/))  was used for genotype classification (version 3.1.17).

The infecting genotype for one subject (Conv-19, Supplementary Table) was determined from a viral sequence derived from an alternative sequencing platform (Illumina MiSeq). Briefly, cDNA synthesis was performed with the Superscript IV first-strand synthesis system (Life Technologies/Invitrogen, Carlsbad, CA). Multiplex PCR was performed with the ARTIC v3 primer set, designed to amplify overlapping regions of the Sars-CoV-2 reference genome (MN908947.3). Primer and genomic alignment position information is available here: https://github.com/artic-network/artic-ncov2019/tree/master/primer_schemes/nCoV-2019/V1. PCR products were purified with the MinElute PCR purification kit (QIAgen, Valencia, CA). Libraries were prepared with the SMARTer PrepX DNA Library Kit (Takara Bio, Mountain View, CA), using the Apollo library prep system (Takara Bio, Mountain View, CA). The libraries were evaluated for quality using the Agilent 2200 TapeStation (Agilent, Santa Clara, CA). After quantification by real-time PCR with the KAPA SYBR FAST qPCR Kit (Roche, Pleasanton, CA), libraries were diluted to 10 nM.

**Ethics**
The PASS (IDCRP-126) and EPICC (IDCRP-085) studies were approved by the Uniformed Services University of the Health Sciences Institutional Review Board in compliance with all applicable Federal regulations governing the protection of human participants. All PASS and EPICC study participants provided informed consent. The sample for the reference convalescent Beta sera obtained from a traveler who had moderate-severe Covid-19 in the Republic of South Africa during the peak of the Beta (B.1.351) wave in January 2021 was obtained with informed consent and covered under the US Food and Drug Administration Institutional Review Board approved expedited protocol # 2021-CBER-045.

**Plasmids, cell lines and therapeutic antibodies**
The gene to express full-length wild-type (D614G-WT) SARS-CoV-2 spike or its variants was codon optimized and cloned into pcDNA3.1 (+) or pVRC7480 plasmid (GeneScript, Piscataway, NJ). The Omicron spike expression plasmid was generously provided by Nicole Doria-Rose (Vaccine Research Center (VRC), National Institutes of Health (NIH), Bethesda, MD) The HIV gag/pol (pCMVΔR8.2) and the Luciferase reporter (pHR'CMV-Luc) plasmids were obtained from the VRC/NIH, Bethesda, MD). 293T-ACE2/TMPRSS2 cell stably expressing ACE2 and transmembrane serine protease 2 (TMRSS2) were established and maintained as previously described (BEI # NR-55293)(*23*).A total of 24 therapeutic antibody products in various stages of

development were provided by pharmaceutical companies to support the US-government COVID-19 therapeutics response. The identity of these antibodies has been blinded as part of the agreement with the companies.

**SARS-Cov-2 pseudovirus production and neutralization**
HIV-based lentiviral pseudoviruses were produced as previously described(*23*). Briefly, three plasmids were co-transfected, the spike encoding plasmid (4µg for the pcDNA3.1 (+) or 0.5 µg for pVRC7480), the pCMVΔR8.2 plasmid (HIV gag/pol, 5µg) and the pHR'CMV-Luc plasmid (luciferase reporter, 5µg) in 293T cells in 10% FBS/DMEM. The pseudoviruses were expressed for 48h at 37°C after which supernatants were collected and filtered through 0.45 µM low protein binding filters and aliquoted at -80°C until further use.

The neutralization assays were preformed using 293T ACE2/TMRPSS2 cells as previously described(*23*). Briefly, pseudoviruses with titers $0.8-1.5^6$ relative luminescence units (RLU)/mL were incubated with 4x serial dilutions of antibodies or 3x serial dilutions of sera for 2h at 37°C. The mixtures (100 µL) were used to inoculate plates with 293T ACE2/TMPRSS2 cells plated the day before at $3.0 \times 10^4$ cells/well. The virus infectivity was determined 48 h later by quantifying the luciferase activity. Neutralization curves were normalized to a virus only control. Neutralization curves were fitted using nonlinear regression curve (GraphPad Prism, La Jolla, CA). The antibody concentration or sera dilution corresponding to 50% neutralization was defined as $NT_{50}$ for sera or $IC_{50}$ for antibody products. All experiments were done at least in duplicates.

**Antigenic cartography**
ACMACS antigenic cartography software (https://acmacs-web.antigenic-441cartography.org/) was used to create a geometric interpretation of neutralization titers against the WT, Delta and Omicron variants. The map is presented on a grid in which each square indicates one antigenic unit, corresponding to two-fold dilution of the antibody in the neutralization assay. Antigenic distance is measured in any direction of the map.

**Supplementary Materials**

**Supplementary Table. SARS-CoV-2 genotypes for the infecting variants**

**References and Notes**

1.    D. Corti, L. A. Purcell, G. Snell, D. Veesler, Tackling COVID-19 with neutralizing monoclonal antibodies. *Cell* **184**, 3086-3108 (2021).
2.    Y. Cao, J. Wang, F. Jian, T. Xiao, W. Song, A. Yisimayi, W. Huang, Q. Li, P. Wang, R. An, J. Wang, Y. Wang, X. Niu, S. Yang, H. Liang, H. Sun, T. Li, Y. Yu, Q. Cui, S. Liu, X. Yang, S. Du, Z. Zhang, X. Hao, F. Shao, R. Jin, X. Wang, J. Xiao, Y. Wang, X. S. Xie, B.1.1.529 escapes the majority of SARS-CoV-2 neutralizing antibodies of diverse epitopes. *bioRxiv*, 2021.2012.2007.470392 (2021).
3.    A. Rössler, L. Riepler, D. Bante, D. v. Laer, J. Kimpel, SARS-CoV-2 B.1.1.529 variant (Omicron) evades neutralization by sera from vaccinated and convalescent individuals. *medRxiv*, 2021.2012.2008.21267491 (2021).

4. L. Lu, B. W.-Y. Mok, L. Chen, J. M.-C. Chan, O. T.-Y. Tsang, B. H.-S. Lam, V. W.-M. Chuang, A. W.-H. Chu, W.-M. Chan, J. D. Ip, B. P.-C. Chan, R. Zhang, C. C.-Y. Yip, V. C.-C. Cheng, K.-H. Chan, I. F.-N. Hung, K.-Y. Yuen, H. Chen, K. K.-W. To, Neutralization of SARS-CoV-2 Omicron variant by sera from BNT162b2 or Coronavac vaccine recipients. *medRxiv*, 2021.2012.2013.21267668 (2021).

5. N. Ikemura, A. Hoshino, Y. Higuchi, S. Taminishi, T. Inaba, S. Matoba, SARS-CoV-2 Omicron variant escapes neutralization by vaccinated and convalescent sera and therapeutic monoclonal antibodies. *medRxiv*, 2021.2012.2013.21267761 (2021).

6. E. Cameroni, C. Saliba, J. E. Bowen, L. E. Rosen, K. Culap, D. Pinto, A. De Marco, S. K. Zepeda, J. di Iulio, F. Zatta, H. Kaiser, J. Noack, N. Farhat, N. Czudnochowski, C. Havenar-Daughton, K. R. Sprouse, J. R. Dillen, A. E. Powell, A. Chen, C. Maher, L. Yin, D. Sun, L. Soriaga, C. Gustafsson, H. Chu, N. M. Franko, J. Logue, N. T. Iqbal, I. Mazzitelli, J. Geffner, R. Grifantini, A. Gori, A. Riva, O. Giannini, A. Ceschi, P. Ferrari, A. Franzetti-Pellanda, C. Garzoni, C. Hebner, L. Purcell, L. Piccoli, M. S. Pizzuto, A. C. Walls, A. Telenti, H. W. Virgin, A. Lanzavecchia, D. Veesler, G. Snell, D. Corti, Broadly neutralizing antibodies overcome SARS-CoV-2 Omicron antigenic shift. *bioRxiv*, 2021.2012.2012.472269 (2021).

7. S. Cele, L. Jackson, K. Khan, D. Khoury, T. Moyo-Gwete, H. Tegally, C. Scheepers, D. Amoako, F. Karim, M. Bernstein, G. Lustig, D. Archary, M. Smith, Y. Ganga, Z. Jule, K. Reedoy, J. E. San, S.-H. Hwa, J. Giandhari, J. M. Blackburn, B. I. Gosnell, S. A. Karim, W. Hanekom, NGS-SA, C.-K. Team, A. von Gottberg, J. Bhiman, R. J. Lessells, M.-Y. S. Moosa, M. Davenport, T. de Oliveira, P. L. Moore, A. Sigal, SARS-CoV-2 Omicron has extensive but incomplete escape of Pfizer BNT162b2 elicited neutralization and requires ACE2 for infection. *medRxiv*, 2021.2012.2008.21267417 (2021).

8. C. I. Kaku, E. R. Champney, J. Normark, C. E. Johnson, C. Ahlm, M. Sakharkar, M. E. Ackerman, M. N. E. Forsell, L. M. Walker, Broad anti-SARS-CoV-2 antibody immunity induced by heterologous ChAdOx1/mRNA-1273 prime-boost vaccination. *medRxiv*, 2021.2012.2013.21267598 (2021).

9. W. F. Garcia-Beltran, K. J. St Denis, A. Hoelzemer, E. C. Lam, A. D. Nitido, M. L. Sheehan, C. Berrios, O. Ofoman, C. C. Chang, B. M. Hauser, J. Feldman, D. J. Gregory, M. C. Poznansky, A. G. Schmidt, A. J. Iafrate, V. Naranbhai, A. B. Balazs, mRNA-based COVID-19 vaccine boosters induce neutralizing immunity against SARS-CoV-2 Omicron variant. *medRxiv*, 2021.2012.2014.21267755 (2021).

10. H. Gruell, K. Vanshylla, P. Tober-Lau, D. Hillus, P. Schommers, C. Lehmann, F. Kurth, L. E. Sander, F. Klein, mRNA booster immunization elicits potent neutralizing serum activity against the SARS-CoV-2 Omicron variant. *medRxiv*, 2021.2012.2014.21267769 (2021).

11. F. Schmidt, F. Muecksch, Y. Weisblum, J. D. Silva, E. Bednarski, A. Cho, Z. Wang, C. Gaebler, M. Caskey, M. C. Nussenzweig, T. Hatziioannou, P. D. Bieniasz, Plasma neutralization properties of the SARS-CoV-2 Omicron variant. *medRxiv*, 2021.2012.2012.21267646 (2021).

12. K. Basile, R. J. Rockett, K. McPhie, M. Fennell, J. Johnson-Mackinnon, J. E. Agius, W. Fong, H. Rahman, D. Ko, L. Donavan, L. Hueston, C. Lam, A. Arnott, S. C.-A. Chen, S. Maddocks, M. V. O'Sullivan, D. E. Dwyer, V. Sintchenko, J. Kok, Improved neutralization of the SARS-CoV-2 Omicron variant after Pfizer-BioNTech BNT162b2 COVID-19 vaccine boosting. *bioRxiv*, 2021.2012.2012.472252 (2021).

13. I. Nemet, L. Kliker, Y. Lustig, N. S. Zuckerman, O. Erster, C. Cohen, Y. Kreiss, S. Alroy-Preis, G. Regev-Yochay, E. Mendelson, M. Mandelboim, Third BNT162b2 vaccination neutralization of SARS-CoV-2 Omicron infection. *medRxiv*, 2021.2012.2013.21267670 (2021).

14. A. Wilhelm, M. Widera, K. Grikscheit, T. Toptan, B. Schenk, C. Pallas, M. Metzler, N. Kohmer, S. Hoehl, F. A. Helfritz, T. Wolf, U. Goetsch, S. Ciesek, Reduced Neutralization of SARS-CoV-2 Omicron Variant by Vaccine Sera and monoclonal antibodies. *medRxiv*, 2021.2012.2007.21267432 (2021).

15. N. Doria-Rose, X. Shen, S. D. Schmidt, S. O'Dell, C. McDanal, W. Feng, J. Tong, A. Eaton, M. Maglinao, H. Tang, R. L. Atmar, K. E. Lyke, L. Wang, y. Zhang, M. R. Gaudinski, W. P. Black, I. Gordon, M. Guech, J. E. Ledgerwood, J. N. Misasi, A. Widge, P. C. Roberts, J. Beigel, B. Korber, R. Pajon, J. R. Mascola, D. C. Montefiori, Booster of mRNA-1273 Vaccine Reduces SARS-CoV-2 Omicron Escape from Neutralizing Antibodies. *medRxiv*, 2021.2012.2015.21267805 (2021).

16. S. N. Neerukonda, R. Vassell, R. Herrup, S. Liu, T. Wang, K. Takeda, Y. Yang, T.-L. Lin, W. Wang, C. D. Weiss, Establishment of a well-characterized SARS-CoV-2 lentiviral pseudovirus neutralization assay using 293T cells with stable expression of ACE2 and TMPRSS2. *PLOS ONE* **16**, e0248348 (2021).

17. B. M. Jackson-Thompson, E. Goguet, E. D. Laing, C. H. Olsen, S. Pollett, K. M. Hollis-Perry, S. E. Maiolatesi, L. Illinik, K. F. Ramsey, A. E. Reyes, Y. Alcorta, M. A. Wong, J. Davies, O. Ortega, E. Parmelee, A. R. Lindrose, M. Moser, E. Graydon, A. G. Letizia, C. A. Duplessis, A. Ganesan, K. P. Pratt, A. M. Malloy, D. W. Scott, S. K. Anderson, A. L. Snow, C. L. Dalgard, J. H. Powers, 3rd, D. Tribble, T. H. Burgess, C. C. Broder, E. Mitre, Prospective Assessment of SARS-CoV-2 Seroconversion (PASS) study:

an observational cohort study of SARS-CoV-2 infection and vaccination in healthcare workers. *BMC Infect Dis* **21**, 544 (2021).

18.    E. D. Laing, C. D. Weiss, E. C. Samuels, S. A. A. Coggins, W. Wang, R. Wang, R. Vassell, S. L. Sterling, M. A. Tso, T. Conner, E. Goguet, M. Moser, B. M. Jackson-Thompson, L. Illinik, J. Davies, O. Ortega, E. Parmelee, M. Hollis-Perry, S. E. Maiolatesi, G. Wang, K. F. Ramsey, A. E. Reyes, Y. Alcorta, M. A. Wong, A. R. Lindrose, C. A. Duplessis, D. R. Tribble, A. M. W. Malloy, T. H. Burgess, S. D. Pollett, C. H. Olsen, C. C. Broder, E. Mitre, Durability of antibody responses and frequency of clinical and subclinical SARS-CoV-2 infection six months after BNT162b2 COVID-19 vaccination in healthcare workers. *medRxiv*, 2021.2010.2016.21265087 (2021).

19.    S. A. Richard, S. D. Pollett, C. A. Lanteri, E. V. Millar, A. C. Fries, R. C. Maves, G. C. Utz, T. Lalani, A. Smith, R. M. Mody, A. Ganesan, R. E. Colombo, C. J. Colombo, D. A. Lindholm, C. Madar, S. Chi, N. Huprikar, D. T. Larson, S. E. Bazan, C. English, E. Parmelee, K. Mende, E. D. Laing, C. C. Broder, P. W. Blair, J. G. Chenoweth, M. P. Simons, D. R. Tribble, B. K. Agan, T. H. Burgess, E. C.-C. S. Group, COVID-19 Outcomes Among US Military Health System Beneficiaries Include Complications Across Multiple Organ Systems and Substantial Functional Impairment. *Open Forum Infect Dis* **8**, ofab556 (2021).

20.    S. Lusvarghi, W. Wang, R. Herrup, S. N. Neerukonda, R. Vassell, L. Bentley, A. E. Eakin, K. J. Erlandson, C. D. Weiss, Key substitutions in the spike protein of SARS-CoV-2 variants can predict resistance to monoclonal antibodies, but other substitutions can modify the effects. *J Virol*, JVI0111021 (2021).

21.    M. Schubert, F. Bertoglio, S. Steinke, P. A. Heine, M. A. Ynga-Durand, F. Zuo, L. Du, J. Korn, M. Milošević, E. V. Wenzel, H. Maass, F. Krstanović, S. Polten, M. Pribanić-Matešić, I. Brizić, A. Piralla, F. Baldanti, L. Hammarström, S. Dübel, A. Šustić, H. Marcotte, M. Strengert, A. Protić, Q. Pan Hammarström, L. Čičin-Šain, M. Hust, Human serum from SARS-CoV-2 vaccinated and COVID-19 patients shows reduced binding to the RBD of SARS-CoV-2 Omicron variant in comparison to the original Wuhan strain and the Beta and Delta variants. *medRxiv*, 2021.2012.2010.21267523 (2021).

22.    S. Jangra, C. Ye, R. Rathnasinghe, D. Stadlbauer, g. Personalized Virology Initiative study, F. Krammer, V. Simon, L. Martinez-Sobrido, A. Garcia-Sastre, M. Schotsaert, SARS-CoV-2 spike E484K mutation reduces antibody neutralisation. *Lancet Microbe*, (2021).

23.    S. N. Neerukonda, R. Vassell, R. Herrup, S. Liu, T. Wang, K. Takeda, Y. Yang, T. L. Lin, W. Wang, C. D. Weiss, Establishment of a well-characterized SARS-CoV-2 lentiviral pseudovirus neutralization assay using 293T cells with stable expression of ACE2 and TMPRSS2. *PLoS One* **16**, e0248348 (2021).

## Acknowledgments:

The authors wish to also acknowledge all who have contributed to the EPICC (IDCRP-085) COVID-19 study

*Brooke Army Medical Center, Fort Sam Houston, TX*: Col J. Cowden; LTC M. Darling; S. DeLeon; Maj D. Lindholm; LTC A. Markelz; K. Mende; S. Merritt; T. Merritt; LTC N. Turner; CPT T. Wellington

*Carl R. Darnall Army Medical Center, Fort Hood, TX*:  LTC S. Bazan; P.K Love

*Fort Belvoir Community Hospital, Fort Belvoir, VA:* N. Dimascio-Johnson; MAJ E. Ewers; LCDR K. Gallagher; LCDR D. Larson; A. Rutt

*Henry M. Jackson Foundation, Inc., Bethesda, MD:* P. Blair; J. Chenoweth; D. Clark

*Madigan Army Medical Center, Joint Base Lewis McChord, WA*: S. Chambers; LTC C. Colombo; R. Colombo; CAPT C. Conlon; CAPT K. Everson; COL P. Faestel; COL T. Ferguson; MAJ L. Gordon; LTC S. Grogan; CAPT S. Lis; COL C. Mount; LTC D. Musfeldt; CPT D.

Odineal; LTC M. Perreault; W. Robb-McGrath; MAJ R. Sainato; C. Schofield; COL C. Skinner; M. Stein; MAJ M. Switzer; MAJ M. Timlin; MAJ S. Wood

*Naval Medical Center Portsmouth, Portsmouth, VA*: S. Banks; R. Carpenter; L. Kim; CAPT K. Kronmann; T. Lalani; LCDR T. Lee; LCDR A. Smith; R. Smith; R. Tant; T. Warkentien

*Naval Medical Center San Diego, San Diego, CA:* CDR C. Berjohn; S. Cammarata; N. Kirkland; D. Libraty; CAPT (Ret) R. Maves; CAPT (Ret) G. Utz

*Tripler Army Medical Center, Honolulu, HI*: S. Chi; LTC R. Flanagan; MAJ M. Jones; C. Lucas; LTC (Ret) C. Madar; K. Miyasato; C. Uyehara

*Uniformed Services University of the Health Sciences, Bethesda, MD*: B. Agan; L. Andronescu; A. Austin; C. Broder; CAPT T. Burgess; C. Byrne; COL K Chung; J. Davies; C. English; N. Epsi; C. Fox; M. Fritschlanski; M. Grother; A. Hadley; COL P. Hickey; E. Laing; LTC C. Lanteri; LTC J. Livezey; A. Malloy; R. Mohammed; C. Morales; P. Nwachukwu; C. Olsen; E. Parmelee; S. Pollett; S. Richard; J. Rozman; J. Rusiecki; E. Samuels; M. Sanchez; A. Scher; CDR M. Simons; A. Snow; K. Telu; D. Tribble; L. Ulomi

*United States Air Force School of Medicine, Dayton, OH*: SSgt T. Chao; R. Chapleau; M. Christian; A. Fries; C. Harrington; S. Huntsberger; K. Lanter; E. Macias; J. Meyer; S. Purves; K. Reynolds; J. Rodriguez; C. Starr

Womack Army Medical Center, Fort Bragg, NC: B. Barton; LTC D. Hostler; LTC J. Hostler; MAJ K. Lago; C. Maldonado; J. Mehrer

*United States Army Medical Research Institute of Infectious Diseases, Frederick, MD*: MAJ J. Kugelman

*William Beaumont Army Medical Center, El Paso, TX*:  MAJ T. Hunter; J. Mejia; R. Mody; R. Resendez; P. Sandoval; M. Wayman

*Walter Reed National Military Medical Center, Bethesda, MD:* I. Barahona; A. Baya; A. Ganesan; MAJ N. Huprikar; B. Johnson

*Walter Reed Army Institute of Research, Silver Spring, MD*: S. Peel

We also thank Douglas Pratt and Hector Izurieta (US Food and Drug Administration) for critical review of the manuscript.

**Funding:**

**Funding Statement:**
This work was supported by awards from the Defense Health Program (HU00012020067, HU00012120094, HU00012120104) and the National Institute of Allergy and Infectious Disease

(HU00011920111). The PASS and EPICC protocols were executed by the Infectious Disease Clinical Research Program (IDCRP), a Department of Defense (DoD) program executed by the Uniformed Services University of the Health Sciences (USUHS) through a cooperative agreement by the Henry M. Jackson Foundation for the Advancement of Military Medicine, Inc. (HJF). This project has also been funded in part by the National Institute of Allergy and Infectious Diseases at the National Institutes of Health (NIAID/NIH), under an interagency agreement with USUHS (Y1-AI-5072), and by the Intramural Research Program of NIAID.. Additional support was provided by institutional research funds from the US Food and Drug Administration (FDA) and NIAID/NIH through an interagency agreement (AAI21013-001-00000) with Center for Biologics Evaluation and Research, FDA, as part of the US Government COVID-19 response efforts.

**Author contributions:**
Conceptualization: SL, SDP, RV, EG, CCB, LB, KJE, EDL, THB, EM, CDW
Methodology: SL, SDP, SNN, WW, RV, ACF, EG, SAC, CCB, LB, KJE, EDL, LCK, EM, CDW
Investigation: SL, SDP, SNN, WW, RW, RV, NJE, ACF, DAL, CJC, RM, AG, EG, SAC, GW, LB, KJE, ECE, TL, CDW
Visualization: SL, SDP, WW, RV, LCK, CDW
Funding acquisition: SDP, AEE, DRT, THB, EM, CDW
Project administration: SDP, RV, BKA, MH-P, MPS, DRT, AEE, KJE, EDL, THB, EM, CDW
Supervision: SDP, CCB, EDL, EM, THB, CDW
Writing – original draft: SL, SDP, CDW
Writing – review & editing: SL, SDP, SNN, WW, RW, RV, NJE, ACF, BKA, DAL, CJC, RM, ECE, TL, AG, EG, MH-P, SAC, MPS, LCK, GW, CCB, DRT, LB, AEE, KJE, EDL, THB, EM, CDW

**Competing interests:**
**Potential conflicts of interest.** S. D. P., T. H. B, and D.R.T. report that the Uniformed Services University (USU) Infectious Diseases Clinical Research Program (IDCRP), a US Department of Defense institution, and the Henry M. Jackson Foundation (HJF) were funded under a Cooperative Research and Development Agreement to conduct an unrelated phase III COVID-19 monoclonal antibody immunoprophylaxis trial sponsored by AstraZeneca. The HJF, in support of the USU IDCRP, was funded by the Department of Defense Joint Program Executive Office for Chemical, Biological, Radiological, and Nuclear Defense to augment the conduct of an unrelated phase III vaccine trial sponsored by AstraZeneca. Both of these trials were part of the US Government COVID-19 response.

**Disclaimer:** E. Laing, T. Burgess, D. Tribble, A Fries, C. Broder, E. Mitre, M. Hollis-Perry, S. Lusvarghi, S. Pollett, SN. Neerukonda, W. Wang, R. Wang, R. Vassell, LC. Katzelnick, L Bentley, AE. Eakin, KJ. Erlandson and CD. Weiss are service members or employees of the U.S. Government. This work was prepared as part of their official duties. Title 17 U.S.C. §105 provides that 'Copyright protection under this title is not available for any work of the United States Government.' Title 17 U.S.C. §101 defines a U.S. Government work as a work prepared

**Group authors EPICC and PASS (MEDLINE indexed):**

Brooke Army Medical Center, Fort Sam Houston, TX: Katrin Mende

Clinical Trials Center, Infectious Diseases Directorate, Naval Medical Research Center, Silver Spring, MD, USA: Christopher Duplessis, Kathleen F. Ramsey, Anatalio E. Reyes, Yolanda Alcorta, Mimi A. Wong, Santina Maiolatesi, Keishla Morales Padilla

Fort Belvoir Community Hospital, Fort Belvoir, VA: Derek Larson

Madigan Army Medical Center, Joint Base Lewis McChord, WA: Rhonda E. Colombo

Naval Medical Center Portsmouth, Portsmouth, VA: Alfred Smith

Uniformed Services University of the Health Sciences, Bethesda, MD: Stephanie Richard, Julia Rozman, Alyssa R. Lindrose, Matthew Moser, Belinda Jackson-Thompson, Margaret Sanchez-Edwards, Luca Illinik, Julian Davies, Orlando Ortega, Edward Parmelee

United States Army Medical Research Institute of Infectious Diseases, Frederick, MD: MAJ J. Kugelman

Walter Reed National Military Medical Center, Bethesda, MD: Nikhil Huprikar

**Fig. 1. Sensitivity of the Omicron variant to neutralization by Pfizer/BNT162b2 vaccinee sera.** (A) Neutralization assays were performed using lentiviral pseudoviruses bearing SARS-CoV-2 WT (B.1 lineage, D614G), Delta (Δ, B.1.617.2 lineage, T19R, G142D, E156-, F157-, R158G, L452R, T478K, D614G, P681R and D950N) and Omicron (O, B.1.1.529 lineage, A67V, del69-70, T95I, del142-144, Y145D, del211, L212I, ins214EPE, G339D, S371L, S373P, S375F,

K417N, N440K, G446S, S477N, T478K, E484A, Q493R, G496S, Q498R, N501Y, Y505H, T547K, D614G, H655Y, N679K, P681H, N764K, D796Y, N856K, Q954H, N969K and L981F) with sera from 39 healthcare workers after immunization with two and three doses of Pfizer/BNT162b2 vaccine. Sera was obtained at a mean of $30 \pm 11$ days after immunization with two doses and at a mean of $43 \pm 17$ days after booster vaccination. Each serum was run in duplicate in two independent experiments against each pseudovirus to determine the 50% neutralization titer ($NT_{50}$). The geometric means titers (GMT), the number of $NT_{50}$ above threshold (1:40) and the fold change are indicated. Titers below 1:40 were set at 20 to calculate GMTs. Arrows indicate increase or decrease relative to WT. Connecting lines indicate serum from the same individual. The demographic information for this sera cohort is provided in Table 1. (B) $NT_{50}$ by sex after $2^{nd}$ or $3^{rd}$ vaccination. (C) $NT_{50}$ by age after $2^{nd}$ and $3^{rd}$ vaccination. (D) $2^{nd}$ or $3^{rd}$ vaccination $NT_{50}$ according to anti-N (nucleocapsid protein) seroconversion between $2^{nd}$ and $3^{rd}$ vaccination. (E) The ratios between the neutralization titers after the $3^{rd}$ and the $2^{nd}$ immunization for Omicron and Delta were plotted against the corresponding ratios for WT. For panels B-E, black squares correspond to WT, blue triangles correspond to Delta and red circles correspond to Omicron.

**Fig. 2. Sensitivity of the Omicron variant to neutralization by convalescent sera**

Neutralization assays were performed using convalescent sera from persons infected with genotyped variants from B.1, B.1.2, B.1.1.7, B.1.351, B.1.617.2, AY.14, AY.25, AY.44, AY.47, AY.62, AY.74 or AY.119 lineages (Table 2 and Supplementary Table). Both B.1 and B.1.2 have no mutations in the receptor binding domain and were therefore considered WT, whereas the AY mutants have additional mutations in the RBD. Each serum was run in duplicate against WT, Delta, and Omicron to determine the $NT_{50}$. The geometric means (GMT), the number of $NT_{50}$ above threshold (1:40) and the fold change are indicated. Titers that did not inhibit at the lowest dilution tested (1:40) were assigned a titer of 20 for GMT calculations. Arrows indicate decrease relative to the infecting variant. Connecting lines indicate serum from the same individual. Data shown represent two independent experiments each with an intra-assay duplicate. Squares correspond to WT, triangles correspond to Delta, and circles correspond to Omicron.

**Fig. 3. Antigenic cartography of convalescent and vaccinee sera against WT, Delta and Omicron.** Antigenic maps were separately generated from convalescent (left panel), 2nd vaccination (middle panel) or 3rd vaccination (right panel) sera. Convalescent sera are shown in diamonds as follows: B.1 (dark green), B.1.2 (light green), B.1.1.7 (purple), B.1.351 (brown), AY variants (light blue), and B.1.617.2 (dark blue). Gray diamonds correspond to post-vaccination sera. Each grid square corresponds to 2-fold dilution in the neutralization assay. Black squares correspond to WT variant. Blue triangles correspond to Delta variant. Red circle corresponds to Omicron variant.

**Fig. 4. Neutralization of Omicron by therapeutic antibodies.** (A) Neutralization curves for each one of the 24 therapeutic antibody products against WT (black) and Omicron (red). (B) Bar graph showing the ratio between the $IC_{50}$ of Omicron and WT for all the antibody products. The sensitivity of the Omicron variant against 15 monoclonal antibodies (nAbs), 6 combination nAbs products (cnAbs), and 3 polyclonal antibodies (pAbs). Red indicates $IC_{50}$ resistance ratios >50, yellow indicates moderate resistance with $IC_{50}$ ratios between 5-50, and green indicates sensitivity comparable to WT with $IC_{50}$ ratios <5. Antibodies for which complete neutralization was not achieved at the highest concentration tested are denoted by *. Data shown represent two independent experiments each with an intra-assay duplicate.

**Table 1. Demographic data for participants receiving Pfizer/BNT162b2 initial vaccine series and booster**

| | N (%) |
|---|---|
| **Sex** | |
| Female | 25 (64.1) |
| Male | 14 (35.9) |
| **Race** | |
| White | 26 (66.6%) |
| Asian | 8 (20.5%) |
| Black | 4 (10.3%) |
| Multiracial | 1 (2.6%) |
| **Occupation** | |
| Nurse | 11 (28.2%) |
| Physician | 11 (28.2%) |
| Physical/Occupational/Recreational Therapist | 9 (23.1%) |
| Medical Technician | 3 (7.7%) |
| Lab Personnel | 3 (7.7%) |
| Social Worker | 1 (2.6%) |
| Psychologist | 1 (2.6%) |
| **Anti-N seroconversion after vaccination and before boost** | |
| Positive | 17 (43.6%) |
| Negative | 22 (56.3%) |
| **Age** | |
| Mean age ± SD (range) | 45 ± 11 (26 - 69) |
| **Time between second vaccine and sample collection** | |
| Mean days ± SD (range) | 30 ± 11 (28 - 34) |
| **Time between second vaccine and booster dose** | |
| Mean days ± SD (range) | 267 ± 14 (218-310) |
| **Time between booster dose and sample collection** | |
| Mean days +/- SD (range) | 43 ± 17 (7 - 93) |

**Table 2: Characteristics of unvaccinated infections providing convalescent sera**

| | N = 39 |
|---|---|
| **Gender** | |
| Female | 14 (35.9%) |
| Male | 25 (64.1%) |
| **Race** | |
| White | 29 (74.4%) |
| Asian | 1 (2.6%) |
| Black | 6 (15.4%) |
| Multiracial | 3 (7.7%) |
| **Age** | |
| Mean age ± SD (range) | 41.1 ± 20 (1.4 - 73.2) |
| **Charlson comorbidity index** | |
| 0 | 20 (51.3%) |
| 1-2 | 10 (25.6%) |
| 3-4 | 5 (12.8%) |
| >5 | 4 (10.3%) |
| **Time between infection symptom onset and sample collection** | |
| Mean days ± SD (range) | 30.2 ± 9.3 (14.0 - 51.0) |
| **Severity of initial infection** | |
| Outpatient | 23 (59.0%) |
| Hospitalized | 16 (41.0%) |
| **Infecting genotype**[*] | |
| AY.119 | 1 (2.6%) |
| AY.14 | 2 (5.1%) |
| AY.25 | 3 (7.7%) |
| AY.44 | 1 (2.6%) |
| AY.47 | 1 (2.6%) |
| AY.62 | 1 (2.6%) |
| AY.74 | 1 (2.6%) |
| B.1 | 10 (25.6%) |
| B.1.1.7 | 5 (12.8%) |
| B.1.2 | 6 (15.4%) |
| B.1.351 | 1 (2.6%) |
| B.1.617.2 | 7 (17.9%) |

[*] Genotypes assigned based on Pango 3.1.17 (2021-12-06). The genotype of the infecting variant was determined in all cases except for one, a traveler who had moderate-severe Covid-19 (outpatient) in the Republic of South Africa during the peak of the Beta (B.1.351) wave in January 2021 (FDA IRB Study # 2021-CBER-045).