Distinct patterns of genetic variation at low-recombining genomic regions represent haplotype structure

Jun Ishigohoka[1,*]   Karen Bascón-Cardozo[1]   Andrea Bours[1]   Janina Fuß[2]
Arang Rhie[3]   Jacquelyn Mountcastle[4]   Bettina Haase[4]   William Chow[5]
Joanna Collins[5]   Kerstin Howe[5]   Marcela Uliano-Silva[5]   Olivier Fedrigo[4]
Erich D. Jarvis[4,6,7]   Javier Pérez-Tris[8]   Juan Carlos Illera[9]

Miriam Liedvogel[1,10,*]

[1]Max Planck Institute for Evolutionary Biology, Plön, Germany
[2]Institute of Clinical Molecular Biology (IKMB), Kiel University, Kiel, Germany
[3]Genome Informatics Section, Computational and Statistical Genomics Branch, National Human Genome Research Institute, National Institutes of Health, Bethesda, MD, USA
[4]The Vertebrate Genome Lab, Rockefeller University, New York, NY, USA
[5]Wellcome Sanger Institute, Cambridge, UK
[6]Laboratory of Neurogenetics of Language, Rockefeller University, New York, NY, USA
[7]The Howards Hughes Medical Institute, Chevy Chase, MD, USA
[8]Department of Biodiversity, Ecology and Evolution, Complutense University of Madrid, Madrid, Spain
[9]Biodiversity Research Institute (CSIC-Oviedo University-Principality of Asturias), Oviedo University, Mieres, Spain
[10]Institute of Avian Research, Wilhelmshaven, Germany

[*] Correspondence: Jun Ishigohoka <ishigohoka@evolbio.mpg.de>, Miriam Liedvogel <liedvogel@evolbio.mpg.de>

Abstract

Genetic variation of the entire genome represents population structure, yet individual loci can show distinct patterns. Such deviations identified through genome scans have often been attributed to effects of selective factors instead of randomness, assuming that the genomic intervals are long enough to average out randomness in underlying genealogies. However, an alternative explanation to distinct patterns has not been fully addressed: too few genealogies to average out the effect of randomness. Specifically, distinct patterns of genetic variation may be due to reduced local recombination rate, since the number of genealogies in a genomic interval corresponds to the number of ancestral recombination events. Here, we associate distinct patterns of local genetic variation with reduced recombination rate in a songbird, the Eurasian blackcap, using genome sequences and recombination maps. We find that distinct patterns of local genetic variation represent haplotype structure at low-recombining regions present either in all populations or only in a few populations. At the former species-wide low-recombining regions, genetic variation depicts conspicuous haplotypes segregating in multiple populations. On the contrary, at the latter population-specific low-recombining regions, genetic variation primarily represents cryptic haplotype structure among individuals of the low-recombining populations. With simulations, we confirm that reduction in recombination rate alone can cause distinct patterns of genetic variation mirroring our empirical data. Our results highlight that distinct patterns of genetic variation can emerge through evolution of reduced local recombination rate. Recombination landscape as an evolvable trait therefore plays an important role determining the heterogeneous distribution of genetic variation along the genome.

Introduction

Patterns of genetic variation in the genome represent ancestries of sequences and are influenced by population history. While genome-wide genetic variation represents population structure (McVean, 2009; Patterson et al., 2006), randomness in genealogies also contributes to fluctuation of local genetic variation along recombining chromosomes. Specifically, genealogies can differ between loci even under the same population history (Dutheil et al., 2009; Martin & Van Belleghem, 2017; McVean & Cardin, 2005; Pamilo & Nei, 1988; Wakeley, 2008, 2020; Wiuf & Hein, 1999). This is because realisation of a genealogy under a given population history is a probabilistic process: an ancestral haplotype for a set of individuals at one locus is not necessarily a common ancestor of the same set of individuals at another locus (Shipilina et al., 2023). Patterns of local genetic variation along the genome tend to conform with the population structure with random fluctuation (Fig. 1).

Figure 1: Distinct patterns of genetic variation can be due to reduced recombination rate. Population history (**A**) affects the distribution of possible genealogies (**B**) from which local genealogies are drawn (**C**). The number of genealogies in a genomic interval with a fixed physical length depends on the local recombination rate (**C**). Mutations occurring on the genealogies (not shown) determine the patterns of realised genetic variation (**D**). (1) In freely recombining neutral regions, mutations represent many genealogies and hence the pattern of genetic variation converges to the population structure. (2, 3) In low-recombining neutral regions, mutations represent few genealogies covering the region leading to patterns of genetic variation distinct from the population structure. (3) Due to randomness in sampling of genealogies, some of such distinct patterns can be similar to patterns expected at targets of selective factors (c.f. 4). (4) At targets of selection, distribution of possible genealogies is different from that at neutral regions, which is depicted as a different set of possible genealogies in **B** and the dotted arrow.

³⁵   Inference of population structure as well as other genome-wide analyses based on genetic

³⁶ variation take advantage of a sufficient number of variable sites (e.g. single nucleotide poly-

³⁷ morphisms (SNPs)) to eliminate the effect of randomness. One of the most common methods

³⁸ to summarise population structure based on this approach is principal component analysis

³⁹ (PCA) applied on a whole-genome genotype table (McVean, 2009; Price et al., 2006). In a

⁴⁰ whole-genome PCA, variation among individuals based on variable sites of the entire genome

⁴¹ are projected onto a few major axes, and the distances among individuals on these reduced

dimensions represent genetic differences. Summarising population structure and other related measures using the entire genome has been proven to be an effective approach to eliminate random fluctuation of genealogies along the genome (Bhatia et al., 2013; Cao et al., 2020; Fedorova et al., 2013; Peter, 2022; Shao et al., 2023).

However, some fundamental biological questions concern selective factors that systematically bias the shape of genealogies at a genomic local scale, shifting the expected patterns of genetic variation from the population structure. For example, patterns of local genetic variation are distinct under selection against gene flow (Fig. 1C4), positive selection and adaptive introgression because they affect coalescence rate, topology, and branch lengths of the underlying genealogies (Hejase et al., 2020; Setter et al., 2020; Speidel et al., 2019; Wolf & Ellegren, 2017). Empirically, genome scans of population genetic summary statistics have been used to identify regions with shifted patterns of genetic variation (Irwin et al., 2018; Martin et al., 2015).

Distinct patterns of local genetic variation identified with genome scans are often attributed to the effects of selective factors instead of randomness (Burri, 2017; Mérot et al., 2021) based on the assumption that the genomic intervals are large enough to eliminate random fluctuation of genealogies (Li & Ralph, 2019). However, an alternative non-selective explanation is equally conceivable and yet often overlooked: the genomic region may contain an insufficient number of genealogies to eliminate the effect of randomness. Specifically, low-recombining regions may harbour too few genealogies to eliminate the effect of random fluctuation (Lotterhos, 2019).

We address the effect of reduced recombination rate on local genetic variation using a songbird species, Eurasian blackcap (*Sylvia atricapilla*, hereafter "blackcap"), which is characterised by variability in seasonal migration across its distribution range (Berthold, 1988, 1991; Delmore et al., 2020a; Helbig, 1991). Populations with diverged migratory phenotypes split as recently as ~30,000 years ago, likely corresponding to the last glacial period and now exhibit population structure (Fig. 2A-C, Sup. Fig. 1) (Delmore et al., 2020b). Due to their recent split and relatively large effective population size, genetic differentiation is very low among blackcap populations (Delmore et al., 2020b). The presence of population

structure albeit with the low levels of differentiation makes the blackcap a perfect system to investigate local deviations of genetic variation: even the slightest effects of factors that change local genetic variation are likely detectable because such effects are not obscured by population structure. In addition, fine-scale recombination maps for multiple populations are available for this species (Bascón-Cardozo et al., 2022a), facilitating investigation of the relationship between changes in the recombination landscape and locally distinct patterns of genetic variation.

By leveraging a large-scale genomic re-sequencing dataset, we first systematically explore distinct patterns of local genetic variation along the blackcap genome, and compare these with genomic regions exhibiting reduced recombination rate. We further investigate the patterns of genetic variation in outlier regions with distinct patterns of genetic variation and associate them with the prevalence of recombination suppression across populations. We also conduct simulations to analyse how reduced local recombination rate in the entire species and in a subpopulation affects patterns of genetic variation through time. Finally, we propose a model of local genetic variation representing haplotype structure corresponding to evolutionary changes in local recombination rate.

Results

Deviation of genetic variation coincides with low-recombining regions

To investigate the genome-wide distribution of genetic variation, we mapped short reads of the whole-genomes of 179 blackcaps including 69 newly sequenced individuals (Sup. Table 1) on a *de novo*-assembled reference genome generated through the Vertebrate Genomes Project (VGP, Rhie et al., 2021), and called SNPs (Materials and Methods). To characterise genome-wide genetic variation, we performed PCA using SNPs in all autosomes, revealing population structure. While PC1 and PC2 represented differentiation of island populations (Fig. 2B), PC3 represented structure within continental populations with different migratory phenotypes (Fig. 2C). To identify genomic regions with patterns of genetic variation distinct from the population structure, we performed local PCA using `lostruct` (Li & Ralph, 2019).

Specifically, PCA was performed separately in sliding genomic windows using SNPs, and windows with distinct patterns were identified by evaluating dissimilarity among windows with multidimensionality scaling (MDS). This approach allowed systematic and unbiased exploration unaffected by our definition of populations. By applying a threshold of the MDS values, we identified 32 genomic regions with distinct patterns of genetic variation (hereafter "outlier regions", Fig. 2D, Sup. Table 3, Sup. Fig. 2). Comparing the genomic distribution of these outlier regions to population-level recombination maps, we found that low-recombining regions were significantly enriched in the outlier regions (permutation tests, p-value < 0.001 (Sup. Fig. 8)). The outlier regions coincided with regions in which recombination rate was reduced either in all tested populations ("species-wide") or in certain populations ("population-specific") (Fig. 2E, F, Sup. Fig. 7).

Figure 2: Local PCA outliers coincide with species-wide and population-specific low-recombining regions A. Geographic location of blackcap populations included in this study. Each point on the map represents a sampling location where multiple individuals were sampled. Populations were defined based on the geographic location, migratory phenotype, and genomic-wide population structure. **B, C.** Genome-wide PCA illustrating population structure. **D.** Distribution of outlier regions based on local PCA using `lostruct`. **E, F** Inferred recombination rates along two exemplified chromosomes (chromosomes 1 and 14) in two blackcap populations (cont_medlong and Azores). In **D-F**, purple and green shades respectively indicate positions of outliers that coincide with species-wide and population-specific low-recombining regions. cont_medlong: medium and long distance migrant population breeding on the continent; cont_short: short distance migrant population breeding on the continent; cont_res: resident (non-migrant) population breeding on the continent. All island populations (Canary, Madeira, Azores, Cape Verde, Mallorca, and Crete) are resident.

To further investigate the outlier regions, we separately performed PCA using SNPs in each region, revealing diverse patterns of distinct genetic variation (Fig. 3A-C top). First, species-wide low-recombining regions showed different levels of clustering of individuals in PCA. Specifically, the PCA projections consisted of either three distinct clusters (Fig. 3A top, Sup. Fig. 6), six loose clusters (Fig. 3B top, Sup. Fig. 6), or mixture of all individuals without apparent clustering (Sup. Fig. 6), suggesting that they represent haplotype structure with different numbers of low-recombining alleles. These clusters did not clearly separate populations, indicating a greater contribution of haplotype structure than the population structure. Five of these (e.g. Fig. 3A top, Sup. Figs. 6, 9) had the clearest clustering patterns with three groups of individuals in PCA , which is expected for a haplotype block with two distinct alleles (Huang et al., 2020; Ma & Amos, 2012; Todesco et al., 2020). Two of these regions showed LD patterns consistent with segregating inversions (Fig. 3A bottom, Sup. Fig. 10), and the other three showed patterns of non-inversion haplotype blocks (Sup. Fig. 10), indicating that recombination suppression with different mechanisms resulted in similar patterns of genetic variation due to presence of two distinct segregating haplotypes.

Figure 3: Patterns of genetic variation and linkage disequilibrium at local PCA outliers
Top: PCA at exemplified outlier regions visualising the patterns of local genetic variation. Data points represent blackcap individuals colour-coded by population as depicted in Fig. 2. **Bottom**: LD calculated for all individuals (top-left diagonal) and for subset individuals (bottom-right diagonal). **A.** A putative inversion. Three clusters correspond to combination of two non-recombining alleles possessed by individuals, depicted as AA, AB, and BB. LD calculated using AA individuals is not elevated, in line with heterozygote-specific recombination suppression at an inversion locus (Sup. Fig. 12). **B.** A species-wide low-recombining region with six loose clusters of individuals. LD calculated using subset individuals was elevated, suggesting genotype-non-specific recombination suppression. **C.** A population-specific low-recombining region. The variance in genetic distances between individuals of the low-recombining populations (Azores (blue) and Cape Verde (light blue)) is greater than between other pairs of individuals (top). LD calculated using individuals of the low-recombining populations is elevated (bottom).

Second, population-specific low-recombining regions exhibited distinct patterns of genetic variation consistently across the outlier regions. While individuals from the low-recombining populations were spread in PCA projections, individuals of other populations were more densely clustered (Fig. 3C top). This pattern indicates that the variance in genetic distances between a pair of individuals of the low-recombining populations is greater than between individuals of normally recombining populations. LD was elevated only in the low-recombining populations (Fig. 3C bottom), supporting population-specific reduction in recombination rate.

Reduced recombination rate generates distinct patterns of genetic variation

To address whether species-wide and population-specific reduction in recombination rate generates the distinct patterns of genetic variation that we observed above, we performed simulations.

First, to investigate the effects of species-wide reduction in local recombination rate, we simulated one ancestral population of 1,000 diploids with a low-recombining genomic region that splits into three subpopulations (pop1, pop2, pop3. Fig. 4A). We sampled individuals over time after the populations split and conducted PCA both in the low-recombining and normally recombining genomic regions. PCA patterns at low-recombining regions (Fig. 4B, C, Sup. Fig. 17) were distinct from normally recombining regions (Fig. 4D). The low-recombining regions exhibited three, six, or more clusters of individuals resembling our empirical results. The clusters of individuals represented genotypes consisting of different combinations of ancestral haplotypes (Sup. Fig. 18). The distinct patterns representing haplotype structure persisted until population structure emerged along the PC axes (Fig. 4B, C). Accordingly, the percentages of variation explained by PC1 and PC2 were higher at low-recombining regions than in normally recombining region until this transition (Fig. 4C). Distinct patterns in the low-recombining regions persisted over longer times than it took for population structure in normally recombining region to emerge (Fig. 4D). These results suggest that distinct patterns of genetic variation in species-wide low-recombining regions represent transient haplotype structure where transition to the population structure is slower than in normally recombining regions.

Figure 4: Simulation of a species-wide low-recombining region. A. Simulated demography scenario. Our simulated genome contained two chromosomes, one with a low-recombining region and the other without. **B, C.** PCA showing patterns of genetic variation at the species-wide low-recombining region at three time points in three exemplified simulation replicates. **D.** PCA showing patterns of genetic variation at a normally recombining chromosome at three time points in the same replicates as **B**.

Second, to investigate the effects of population-specific reduction in local recombination rate, we performed simulations under two scenarios. In both scenarios, three populations (pop1, pop2, and pop3) and their ancestral population had 1,000 diploid individuals, and pop1 evolved a reduced local recombination rate. The difference between the two scenarios was the timing of introduction of reduced recombination rate. In the first scenario (Sup. Fig. 19), recombination suppression was introduced at the same time as the three populations split, while in the second scenario (Fig. 5A) recombination suppression was introduced once the three populations differentiated. We conducted PCA in genomic regions with and without population-specific recombination suppression using individuals sampled over time. In both scenarios, the genomic region with population-specific recombination suppression transiently showed distinct patterns of genetic variation (Fig. 5B, Sup. Fig. 19B) resembling the empirical results, while regions without population-specific suppression showed population structure (Fig. 5C). Haplotype structure was not as conspicuous in species-wide low-recombining regions (Sup. Fig. 20B, F, c.f. Sup. Fig. 18) due to standing genetic variation. Mutations originating in the non-recombining population were enriched in the set of mutations that have the greatest contribution to the distinct pattern of PCA (Sup. Fig. 20C, G. χ^2 tests, p-value < 0.001 for both models). These mutations were significantly associated with each other in the underlying genealogy sharing common branches compared to other mutations originating in the same population (Sup. Fig. 20D, H. Materials and Methods, Kolmogorov-Smirnov tests, p-value < 0.005 for both models), indicating that the distinct pattern of genetic variation represents sets of mutations that occurred in ancestral haplotypes. Associations between these population-specific mutations on ancestral haplotypes would have eventually decayed by recombination events, but in the low-recombining population the association was maintained due to suppressed recombination, resulting in the cryptic haplotype structure.

Figure 5: Simulation of a population-specific low-recombining region. A. Simulated scenario. Simulated genome contained two chromosomes, one with a population-specific low-recombining region and the other without. **B, C.** PCA showing patterns of genetic variation at the population-specific low-recombining region (**B**) and the normally recombining chromosome (**C**) at three time points in one exemplified simulation replicate.

Effect of selection on patterns of genetic variation

Selection is known to cause distinct patterns of genetic variation (Nielsen, 2005). To test whether the outlier regions based on `lostruct` identified in the blackcap genome are also targets of selection, we measured nucleotide diversity (π) and Tajima's D in each population, as well as ratio between non-synonymous and synonymous substitutions (d_N/d_S) for annotated

genes. Many species-wide low-recombining regions showed reduced nucleotide diversity (Sup. Fig. 22) and Tajima's D (Sup. Fig. 21), suggesting that they are under either positive or purifying selection. Most genes within outlier regions had d_N/d_S below 0 (Sup. Fig. 23) with a few genes with positive d_N/d_S, indicating that most genes are under purifying selection and a few others are under positive selection. Furthermore, sequence analysis indicated that some but not all species-wide low-recombining outlier regions coincide with putative pericentromeric regions with enrichment of long tandem repeats (Sup. Figs. 26, 27). These results indicate that the outlier regions may experience effects of selection in addition to reduced recombination rates.

We asked whether the distinct patterns of local genetic variation at the outlier regions observed in blackcaps represent the effect of selection or reduced recombination rates. To this end, we simulated positive and purifying selection in with and without reduction in recombination rate (Materials and Methods), and investigated local genetic variation over time by PCA. Overall, representation of haplotype structure in local PCA occurred primarily when recombination rate was reduced at the focal region (Sup. Figs. 24, 25). Decreased genetic diversity due to selection was represented as small dispersal of individuals around a cluster, while variation between non-selected haplotypes, if present, was represented in the primary PC axes. Separation of populations in local PCA at the low-recombining region occurred faster under a higher level of background selection (Sup. Fig. 24). These results indicate that the distinct patterns of genetic variation represented in local PCA primarily reflect haplotype structure due to reduced recombination rate, on which the effect of selection can be overlaid.

Discussion

A number of empirical population genomics studies have identified ecologically and evolutionarily important genomic regions by locating outlier regions with distinct patterns of genetic variation. Genomic windows in such studies should be ideally both large enough to eliminate the effect of random fluctuation in local genetic variation and small enough to capture the localised signatures of selection. Our results illustrate that distinct patterns of genetic variation in outlier regions based on sliding window approaches can represent haplotype structure reflecting

reduced local recombination rates instead of selection. The exact patterns vary depending on the number of haplotypes and prevalence of recombination suppression across populations. Distinct clusters of individuals based on local genetic variation at species-wide low-recombining regions represent combinations of distinct haplotypes between which shuffling of variation is hindered. Population-specific recombination suppression creates unequal variance in genetic distances between individuals of low-recombining and normally recombining populations.

Distinct patterns of genetic variation at low-recombining regions: Genealogical interpretations

We discuss our findings from the perspective of underlying genealogies. We first define three terms: (1) genealogical noise, (2) genealogical bias, and (3) mutational noise. (1) By "genealogical noise" we refer to the fact that gene genealogies vary along the genome following a null distribution given a population history (Dutheil et al., 2009; Martin & Van Belleghem, 2017; McVean & Cardin, 2005; Wakeley, 2008, 2020; Wiuf & Hein, 1999). (2) By "genealogical bias" we refer to the fact that selective processes can systematically shift the distribution of local genealogies away from the null distribution. For example, genealogies under positive selection, selection against gene flow, adaptive introgression, and balancing selection are biased due to bursts of coalescence, faster lineage sorting, and introduction and maintenance of long branches (Barton & Etheridge, 2004; Guerrero et al., 2012; Hejase et al., 2020; Martin et al., 2019; Setter et al., 2020; Speidel et al., 2019; Taylor, 2013). On top of these, (3) randomness in the process of mutation causes additional noise in realised genetic variation (Ralph et al., 2020), which we call "mutational noise". For example, the first and the second halves of a chromosomal interval with a single genealogy can still have slightly different patterns of genetic variation because they represent independent sets of mutations. Here we show that distinct patterns of local genetic variation at low-recombining regions can be explained primarily by haplotype structure due to non-selective genealogical noise instead of selective genealogical bias.

We showed that some distinct patterns of genetic variation are associated with species-wide low-recombining regions. This is in line with previous studies reporting negative correlation

between recombination rate and genetic differentiation (Burri et al., 2015). To investigate this relationship in more detail, we performed simulations and demonstrated that haplotype structure underlies the distinct patterns and that it persists only transiently until the effect of the population structure emerges. This transiency reflects a shift from local genetic variation primarily representing haplotype structure (Lotterhos, 2019; Ma & Amos, 2012) to that representing population structure, which can be interpreted based on the underlying genealogies. Low-recombining regions have few underlying genealogies and haplotype structure at such regions tend to reflect their basal branches because basal branches tend to be longer than peripheral branches. At a time point soon after a population split event, peripheral branches covering more recent times than the population split harbour fewer mutations than basal branches. Therefore, the realised pattern of genetic variation at this stage has the greatest contributions by mutations on the long basal branches undifferentiated among populations (i.e. consisting standing genetic variation), representing a few ancestral haplotypes that descend the current sample. As time passes after the population split, the proportion of mutations that have occurred after the population split increases while some ancestral haplotypes can be lost by chance (i.e. drift), increasing the contribution of population structure on genetic variation. This type of distinct patterns of genetic variation arises predominantly in low-recombining regions but less so in normally recombining regions. This is because haplotype structure representing a few ancestral lineages would become less prominent with recombination as different segments of a current haplotype can follow distinct ancestries and thus the genealogical noise is effectively averaged out.

Some low-recombining regions may have genealogies with much shorter basal branches than other low-recombining regions because the variance in the basal branch length is greater than peripheral branches (Wakeley, 2008). The over-representation of a few ancestral haplotypes in genetic variation requires long basal branches in the underlying genealogies, and thus low-recombining regions with relatively short basal branches cannot accommodate sufficient mutations to represent distinct ancestral haplotypes. This decreases the relative contribution of genealogical noise compared to mutational noise (Supplementary Notes 1.1). Distinct patterns of genetic variation with varying levels of clustering of individuals in PCA in our empirical

²⁶⁶ results may correspond to different ratios between genealogical and mutational noise due to
²⁶⁷ large variance in the basal branch lengths of underlying genealogies. Specifically, some outlier
²⁶⁸ regions with mixture of individuals from multiple populations without distinct clusters and
²⁶⁹ population subdivision in PCA may have underlying genealogies with short basal branches
²⁷⁰ leading to greater contributions of mutational noise on the realised genetic variation.

²⁷¹   We both empirically and with simulations showed that population-specific low-recombining
²⁷² regions exhibit distinct patterns of genetic variation in which individuals of low-recombining
²⁷³ and normally recombining populations have different variance in genetic distances. This
²⁷⁴ unequal variance in low-recombining and normally recombining populations can be interpreted
²⁷⁵ based on the underlying genealogies (Sup. Fig. 28). We consider the ancestry of current
²⁷⁶ samples of low-recombining and normally recombining populations and split the ancestry at
²⁷⁷ the time T when the population-specific recombination suppression initiated (Sup. Fig. 28A).
²⁷⁸ At time T, there were n ancestral haplotypes that descend all current samples. At times older
²⁷⁹ than T, the ancestors of these n haplotypes freely recombine, making the genetic distances
²⁸⁰ among the ancestral haplotypes close to equidistant (Sup. Fig. 28B). After the initiation of the
²⁸¹ population-specific reduction in recombination rate, the ancestry of one current sequence of
²⁸² the low-recombining population can be traced back to either one of the n ancestral haplotypes
²⁸³ present at the time T (Sup. Fig. 28A). On the contrary, the ancestry of one current sequence
²⁸⁴ of the normally recombining population can be traced back to multiple ancestral haplotypes
²⁸⁵ of the n sequences because of the presence of recombination (Sup. Fig. 28A). From the
²⁸⁶ perspective of mutations, in the low-recombining population, mutations that arose on the same
²⁸⁷ haplotype tend to be linked until the present time because of the suppressed recombination.
²⁸⁸ On the other hand, in the normally recombining population, mutations that arose on the same
²⁸⁹ ancestral haplotype less likely stay linked until the present time because recombination can
²⁹⁰ dissociate them. Because shuffling of haplotypes reduces the variance of genetic distances
²⁹¹ among sequences, population-specific reduction in recombination rates leads to greater variance
²⁹² in low-recombining population than in normally recombining population as observed in our
²⁹³ empirical results and simulations. In short, because of the different recombination rates
²⁹⁴ between the populations, genealogical noise is more efficiently eliminated in the normally

recombining population than in the low-recombining population.

We also demonstrated with simulations that distinct patterns of genetic variation at population-specific low-recombining regions represent cryptic haplotype structure within the low-recombining population. The haplotype structure is only cryptic and less apparent than in species-wide low-recombining regions because other standing mutations coexist on the same haplotype, which are older than the initiation of the population-specific recombination suppression. The elevated PC loadings at linked mutations originating in the low-recombining population could be informative to study evolutionary change in local recombination rate: the ages of such mutations mapped on inferred genealogies might be useful to estimate the timing at which the population-specific recombination suppression initiated.

In our empirical analyses in blackcaps, we detected the effect of population-specific reduction of recombination rate in Azores and Cape Verde island populations. It remains unclear why reduced recombination rate in certain populations but not others is reflected as distinct patterns of genetic variation by `lostruct`. The recent split of Azores and Cape Verde populations from other populations, accompanied by reduction in population size and the level of isolation (Delmore et al., 2020b) may have contributed to more efficient spread of reduced recombination rate.

Recombination landscape as a driver of evolution of local genetic variation

Species-wide and population-specific recombination suppression underlying distinct patterns of local genetic variation are probably not independent: reduction in recombination rates that initiates formation of haplotype blocks likely originates from one population and may spread to multiple populations. For example, local recombination rate may be initially reduced in one population in which a segregating inversion originates before it may spread in multiple populations by gene flow (Faria et al., 2019). In line with this view of recombination map as an evolvable trait diverging across populations according to subdivision, recent studies find that divergence in local recombination rate among populations is correlated with genetic divergence (Bascón-Cardozo et al., 2022a; Spence & Song, 2019). Future work on the effects of transition from population-specific to species-wide suppression of recombination will fill the

gap between the two states.

Besides spread of recombination suppression across populations, there are other paths along which patterns of local genetic variation may change over time. First, change in frequency of one haplotypic variant by drift or gene flow and selection and accumulation of novel mutations may shift the distinct pattern of genetic variation (Rubin et al., 2022). Second, an increase in recombination rate in the region may resolve the distinct pattern of genetic variation and result in emergence of the population structure, because recombination breaks down discrete haplotypes and generates mixed types whereby reducing the variance of genetic variation (Hudson, 1983). These two types of shifts in distinct patterns of genetic variation are not mutually exclusive. For example, fixation of an inversion results in elevated recombination rate (Smukowski Heil et al., 2015; Stevison et al., 2011) because there are no longer non-recombining heterozygotes in the population. Due to resumed recombination, patterns of local genetic variation in such regions are expected to reflect population structure eventually. The question of how long it takes for an outlier region with distinct patterns of genetic variation to disappear after these events should be focally studied in the future.

In Fig. 6A, we illustrate a model for the evolution of local genetic variation that changes according primarily to the evolution of local recombination rates. Local genetic variation can become distinct from the population structure first by representing emerging haplotype structure associated with population-specific recombination suppression or other types of haplotype blocks (e.g. inversions) in one population. If this recombination suppression spreads throughout all populations, then local genetic variation will start to reflect species-wide haplotype structure. Once the relative contribution of haplotype structure on local genetic variation is reduced by differentiation or disappears by elevated recombination rates, then genetic variation returns to reflect the population structure and consequently the outlier region disappears. The effect of selection on local genetic variation may be overlaid on top (Supplementary Notes 1.2).

Figure 6: Evolutionary changes in local recombination rate influence evolution of local genetic variation. A. Local genetic variation is shown in hypothetical PCA plots. Their underlying genealogies are shown in simplified ancestral recombination graphs (ARGs), on which black dots represent ancestral recombination events contributing to the sampled sequences. Points in PCA depict diploid individuals, while those on the ARGs represent haploid sequences. Two colours of these points (blue and orange) indicate two populations. (1) Local genetic variation concordant to population structure. Genetic variation shows separation of individuals from two populations. ARG shows that recombination is suppressed in neither population. (2) Population-specific recombination suppression in the blue population. ARG shows that recombination is suppressed in the blue population. (3) Species-wide recombination suppression. Top: A case in which there are few mutations representing the basal splits of the underlying genealogy at species-wide low-recombining region. Middle: A case in which there are two haplotypic variants at the species-wide low-recombining region. If this is due to presence of an inversion (right ARG), recombination is suppressed between but not within the two clades representing two alleles. Bottom: A case in which there are three haplotypic variants at the species-wide low-recombining region. **B** Evolution of recombination map influences difference in genomic distributions of distinct patterns of genetic variation between species/populations.

Implications

Finally, we discuss technical and biological implications of our study. The technical implication concerns interpretation of genome scans based on local genetic variation. A number of methods based on local genetic variation have been used to detect loci involved in different kinds of selective processes. For example, F_{ST} (differentiation), d_{XY} (divergence), and other population parameters are inferred to detect genomic islands of speciation (Delmore et al., 2018; Hejase et al., 2020; Huang et al., 2020; Malinsky et al., 2015). Reduced diversity (π) is a signature of selection (Delmore et al., 2018; Irwin et al., 2018; Pracana et al., 2017), and by combining it with variation among populations, loci associated with population-specific selection can be also inferred (Yi et al., 2010). Targets of adaptive introgression have been identified by applying statistics based on ABBA-BABA test, which is related to genetic variation (Peter, 2016, 2022), in sliding windows (Kronforst et al., 2013; Martin et al., 2015; Patterson et al., 2012; Reich et al., 2009). However, there are confounding factors that affect inference of these statistics. For example, it has been shown that low diversity can cause elevation in some of these statistics (Cruickshank & Hahn, 2014; Noor & Bennett, 2009). In addition to reduced diversity, this study and others (Lotterhos, 2019; Renaut et al., 2013) show that reduced recombination rate also causes distinct patterns of genetic variation which can lead to erroneous identification of regions under influence of selective factors. Examining recombination rates at identified regions and comparing them to other regions are necessary to avoid this. Furthermore, corroborating methods based on different aspects of distinct patterns of variation, such as site frequency spectrum (DeGiorgio et al., 2016; Fay & Wu, 2000; Tajima, 1989), LD (Sabeti et al., 2002, 2007; Voight et al., 2006), inferred genealogies (Hejase et al., 2020; Speidel et al., 2019; Stern et al., 2019), local landscape of variation (Setter et al., 2020), and sites of mutations in genes (Nei & Gojobori, 1986), as well as approaches with explicit simulation based on inferred demography (Hager et al., 2022), may be informative.

The biological implication is about evolution of recombination rates and genetic variation along the genome. Based on our findings of a link between these, we predict that organisms with more conserved recombination landscape along the genome may have more conserved genomic landscapes of distinct patterns of genetic variation (Fig. 6B). In other words, the

more conserved recombination maps are, the more correlated genomic distribution of distinct genetic variation may be between species. In vertebrates including placental mammals (with some exceptions), recombination landscape along the genome evolves fast due to continuous turnovers of alleles of PRDM9 (the gene coding a protein that determines recombination hot spots) and its target DNA sequences (Baudat et al., 2010; Myers et al., 2008). For instance, in mammals that possess functional PRDM9, the genomic landscape of recombination rates is distinct between and even within species (Kong et al., 2010; Spence & Song, 2019; Stevison et al., 2016). Importantly, PRDM9 has been pseudogenised (Birtle & Ponting, 2006) or lost (Baker et al., 2017) independently in multiple vertebrate lineages. This shifted the determinants of recombination map from the PRDM9 allele and its target to genomic features such as CpG islands and transcription start sites, stabilising the recombination landscape (Auton et al., 2013; Baker et al., 2017; Singhal et al., 2015). Our results shown in birds, a group lacking PRDM9 (Birtle & Ponting, 2006; Singhal et al., 2015), raises a question whether the evolution of local recombination rates may play an even more important role in shaping local genetic variation in organisms with functional PRDM9. Comparative studies using taxa with and without functional PRDM9 will address this and may link the evolution of genomic landscape of distinct patterns of genetic variation and (in)stability of recombination maps.

Materials and Methods

Empirical analyses

de novo genome assembly

A chromosome-level blackcap reference genome was *de novo* assembled within the Vertebrate Genomes Project (VGP), following pipeline version 1.5 (Rhie et al., 2021). In brief, blood of a female blackcap from the resident Tarifa population in Spain was collected in 100% ethanol on ice and stored at -80 °C (NCBI BioSample accession SAMN12369542). The ethanol supernatant was removed and the blood pellet was resuspended in Bionano Cell Buffer in a 1:2 dilution. Ultra-long high molecular weight (HMW) DNA was isolated using Bionano agarose plug method (Bionano Frozen Whole Nucleated Blood Stored in Ethanol – DNA

Isolation Guidelines (document number 30033)) using the Bionano Prep Blood and Cell Culture DNA Isolation Kit. Four DNA extractions were performed yielding a total of 13.5 µg HMW DNA. About 6 µg of DNA was sheared using a 26G blunt end needle (PacBio protocol PN 101-181-000 Version 05) to ~40 kb fragments. A large-insert PacBio library was prepared using the Pacific Biosciences Express Template Prep Kit v1.0 following the manufacturer protocol. The library was then size selected (>15 kb) using the Sage Science BluePippin Size-Selection System. The library was then sequenced on 8 PacBio 1M v3 smrtcells on the Sequel instrument with the sequencing kit 3.0 and 10 hours movie with 2 hours pre-extension time, yielding 77.51 Gb of data (~66.29X coverage) with N50 read length averaging around 22,927 bp. We used the unfragmented HMW DNA to generate a linked-reads library on the 10X Genomics Chromium (Genome Library Kit & Gel Bead Kit v2 , Genome Chip Kit v2 , i7 Multiplex Kit PN-120262). We sequenced this 10X library on an Illumina Novaseq S4 150 bp PE lane to ~60X coverage. Unfragmented HMW DNA was also used for Bionano Genomics optical mapping. Briefly, DNA was labeled using the Bionano Prep Direct Label and Stain (DLS) Protocol (30206E) and run on one Saphyr instrument chip flowcell. 136.31 Gb of data was generated (N50 = 301.9kb with a label density = 16.91 labels/100kb). Optical maps were assembled using Bionano Access (N50 = 27.48 Mb and total length = 1.41 Gb). Hi-C libraries were generated by Arima Genomics and Dovetail Genomics and sequenced on HiSeq X at ~60X coverage following the manufacturer's protocols. Proximally ligated DNA was produced using the Arima-HiC kit v1 , sheared and size selected (200 – 600 bp) with SRI beads, and fragments containing proximity-ligated DNA were enriched using streptavidin beads. A final Illumina library was prepared using the KAPA Hyper Prep kit following the manufacturer guidelines. FALCON v1.9.0 and FALCON unzip v1.0.6 were used to generate haplotype phased contigs, and purge_haplotigs v1.0.3 was used to further sort out haplotypes (Guan et al., 2020). The phased contigs were first scaffolded with 10X Genomics linked reads using scaff10X 4.1.0 software, followed with Bionano Genomics optical maps using Bionano Solve single enzyme DLS 3.2.1, and Arima Genomics in-vitro cross-linked Hi-C maps using Salsa Hi-C 2.2 software (Ghurye et al., 2019). Base call errors were polished with both PacBio long reads and Arrow short reads to achieve above Q40 accuracy (no more than 1 error every 10,000 bp). Manual curation was conducted using gEVAL software by the Sanger Institute Curation team (Howe

et al., 2021). Curation identified 33 autosomes and Z and W chromosomes (plus 1 unlocalised W). Autosomes were named in decreasing order of size, and autosomes 1 through 30 and sex chromosomes had counterparts in the commonly used VGP reference zebra finch assembly (Sup. Table 2). The total length of the primary haplotype assembly was 1,066,786,587 bp, with 99.14% assigned to chromosomes. The final 1.1 Gb assembly consisted of 601 contigs in 189 scaffolds, with a contig N50 of 7.4 Mb, and scaffold N50 of 73 Mb, indicating a high-quality assembly that fulfills the VGP standard metrics.

Whole-genome resequencing

We resequenced 69 blackcap samples from various populations across the species distribution range (Sup. Table 1) to complement an existing dataset of 110 blackcaps, 5 garden warblers, and 3 African hill babblers that had been sequenced previously (Delmore et al., 2020b). Blood samples from the additional 69 blackcaps were collected from the brachial vein and stored in 100% ethanol. High molecular weight genomic DNA was extracted with a standard salt extraction protocol or through the Nanobind CBB Big DNA Kit Beta following the manufacturer's instructions. Libraries for short insert fragments between 300 and 500 bp were prepared and were then sequenced for short paired-end reads on either Illumina NextSeq 500, HiSeq 4000 or NovaSeq 5000 (Sup. Table 1).

We performed quality control of the reads with FastQC version 0.11.8 (https://www.bioinformatics.babraham.ac.uk/projects/fastqc/). Reads from all samples were mapped against the blackcap reference genome following an adjusted pipeline of `Genome Analysis Toolkit` (`GATK` version 4.1.7.0, McKenna et al. (2010)) and `Picard` version 2.21.9 (http://broadinstitute.github.io/picard/). After resetting the base quality of adapter bases in the sequenced reads to 2 with `Picard MarkIlluminaAdapters`, paired-end reads were mapped to the reference using `BWA mem` (Li, 2013). To ensure that both unmapped mates and secondary/supplementary reads were marked for duplicates, we ran `Picard MarkDuplicates` for sorted reads with the default pixel distance of 100 for reads from Illumina NextSeq 500 or with a pixel distance of 2,500 for reads from HiSeq 4000 and NovaSeq 5000. Due to low coverage, 10 samples (Sup. Table 1) were sequenced multiple times. Alignment files for these

samples (in BAM format) were merged with `Picard MergeSamFiles`. Per-sample quality control of BAM files were performed using `QualiMap` version 2.2.1 (Okonechnikov et al., 2016), `Picard CollectMultipleMetrics`, `CollectRawWgsMetrics` and `CollectWgsMetrics`; and `MultiQC` version 1.8 (Ewels et al., 2016). We called bases at all positions per sample using `GATK HaplotypeCaller`. We combined gVCF files of 189 individuals into ten evenly sized subsets (to allow parallelisation of the following variant calling step) with `GATK CombineGVCFs`. We genotyped SNPs and INDELs using `GATK GenotypeGVCFs`, and the 10 subsets were concatenated using `Picard GatherVcfs` into one VCF file covering the entire genome. From the VCF file, SNPs were selected (i.e. indels were excluded) using `GATK SelectVariants`, after which we filtered SNPs with the following criteria: `QD < 2.5; FS > 45.0; SOR > 3.0; MG < 40; MQRankSum < -12.5; ReadPosRankSum < -8.0`. We removed garden warblers and African hill babblers from the multi-species VCF and kept only biallelic sites. We estimated blackcap haplotypes using `SHAPEIT2` (r837) (Delaneau et al., 2013) with the blackcap recombination map (Bascón-Cardozo et al., 2022a), yielding 142,083,056 SNPs.

Local PCA

To identify genomic regions with distinct patterns of genetic variation in blackcaps, we performed local PCA in sliding genomic windows of 1,000 SNPs and summarised dissimilarity of windows by multidimensional scaling using `lostrct` (Li & Ralph, 2019) in R version 3.5.3. First, we prepared a genotype and a haplotype table for each chromosome in which rows and columns represented positions and individuals from the phased VCF file using `BCFtools`. Specifically, genotypes were encoded 0, 1, and 2 for the reference allele homozygotes, heterozygotes, and non-reference allele homozygotes in the genotype table, and 0 and 2 for the reference and the non-reference allele in the haplotype table (encoding 0 and 1 instead of 0 and 2 in haplotype-based analysis gives the same results). Chromosomes shorter than 10 Mb were concatenated to avoid misidentification of short chromosomal background as an outlier region. Distance matrices of windows were computed based on the coordinates (PC1 and PC2) of samples (individuals for genotype-based local PCA, and haplotype for haplotype-based local PCA) within R using `lostruct`. Multidimensional scaling (MDS) was performed to summarise similarities of local genetic variation patterns among windows into 20 axes (MDS1

through MDS20).

Using the `lostruct` output, we identified chromosomal intervals with distinct patterns of genetic variation. In each chromosome, windows with MDS value apart from the mode of the distribution by greater than 0.3 for any one of the 20 axes were defined as outlier windows. This threshold was determined by visualising the distribution of MDS values in each chromosome (Sup. Fig. 3). For each MDS axis, we defined genomic intervals with at least 10 outlier windows as "outlier regions" with distinct patterns of genetic variation. Overlapping intervals across different MDS axes as well as intervals identified based on genotypes and haplotypes were merged using `BEDtools`. To verify that the outliers show pattern of genetic variation distinct from the whole-genome PCA, we performed PCA using all SNPs within each outlier region using `PLINK` (Purcell et al., 2007). Genomic regions showing similar pattern to the whole genome PCA were identified with visual inspection and discarded from the outliers.

To assess consistency between the pipelines using genotypes and haplotypes, we compared MDS results of genotype- and haplotype-based `lostruct`. We calculated Euclidean distance of windows from the centre of the 20 dimensional space to enable comparison of the same window in genotype- and haplotype-based MDS. We measured this distance instead of comparing the coordinates directly to account for possible rotations of MDS patterns between genotype- and haplotype-based `lostruct`. Because dissimilarity of windows in terms of the pattern of genetic variation was computed per chromosome, we calculated correlation of the above distance between genotype- and haplotype-based methods per chromosome. The comparison of genotype-based and haplotype-based `lostruct` is in Sup. Fig. 4.

To assess whether `lostruct` can identify outliers irrespective of presence/absence of other outliers on the same chromosome as well as the chromosome length, we ran `lostruct` treating either one part of a blackcap chromosome ("split chromosomes") or multiple blackcap chromosomes as a single chromosome ("joined chromosome"). If `lostruct` is robust to the chromosomal background, it is expected that the same regions should be detected as outliers with distinct patterns of genetic variation in both split and joined chromosomes compared to per-chromosome results. We prepared four split chromosomes by splitting chromosomes 1 and 2 at the middle, and one joined chromosome by concatenating chromosomes 20, 21,

and 28. We performed `lostruct` analysis based both on genotype and haplotype and merged the identified regions. The comparison of `lostruct` between using single chromosomes and split/joined chromosomes is in Sup. Fig. 5.

LD and recombination landscape

To calculate LD around outlier regions, we first extracted SNPs within and 30% length outside each outlier. We then thinned SNPs so that all neighbouring SNP positions were at least 10 kb away from each other. Linkage disequilibrium (LD) between all pairs of thinned SNPs was calculated with `VCFtools` with the `--geno-r2`.

We inferred recombination landscape along blackcap chromosomes using `Pyrho` (Spence & Song, 2019). `Pyrho` infers demography-aware recombination rates with a composite-likelihood approach from SNPs data of unrelated samples making use of likelihood lookup tables generated by simulations based on demography and sample size of each population. In all inferences, we used demography of focal populations inferred in Delmore et al. (2020b). Before the recombination inference, focal samples were filtered and singletons were removed. We ran `Pyrho` with mutation rate of 4.6×10^{-9} per site per generation (Smeds et al., 2016), block penalty of 20, and window size of 50 kb to infer population-level recombination landscape in Azores, Cape Verde, continental resident, and medium-long distance migrants (represented by medium distance south-west migrants). We computed mean recombination rate in 10 kb sliding windows for each population.

To test association between local PCA outlier regions and low-recombining regions, we performed permutation tests. For this analysis we defined low-recombining regions as genomic intervals with recombination rate below 1 percentile for the entire genome. We counted the number of overlaps in the observed data using `BEDtools`. For resampling, we shuffled intervals corresponding to local PCA outlier regions within the genome 1,000 times and counted the number of intervals overlapping the low-recombining regions for each population using `BEDtools`.

To characterise genotype-specific LD and recombination landscape at the five outlier regions with three clusters of individuals in PCA, we applied `vcftools --geno-r2` and `Pyrho`

 (Spence & Song, 2019) to our empirical data using each genotype (AA, AB, and BB in Sup.

550 Fig. 9) separately.

Inversion breakpoints

552 Three clusters of individuals observed in PCA with genotype-specific LD at two outlier regions
553 on chromosomes 12 and 30 were indicative of polymorphic inversion (Ma & Amos, 2012; Ruiz-
554 Arenas et al., 2019). To further characterise whether they represent polymorphic inversions,
555 we intended to locate breakpoints by two independent approaches.

Soft-clip reads

We attempted to identify positions where presence of soft-clipping of mapped reads is
associated with PCA-based genotype of the putative inversions. First, we extracted focal
regions around boundaries of the outliers (Sup. Table 4) from read mapping file of all
individuals using `SAMtools` (Danecek et al., 2021). Next, we identified soft clip reads in
each extracted region using `samextractclip` (Lindenbaum, 2015), and obtained reference
position corresponding to the position of soft clipping in mapped reads using a custom script.
At all extracted soft-clip positions, we counted the number of reads that switch to soft-clip
("soft-clip depth"), as well as the depth of mapped reads, using `SAMtools`. At each of all
positions with at least one read supporting soft-clip switch, we calculated proportion of reads
with soft-clip switch relative to all mapped reads (depth of the position) for each individual
("soft-clip proportion"). This resulted in "position-by-individual" matrix whose entry depicts
the proportion of soft-clip in all reads mapped at the focal position for the focal individual.
Using this matrix, we fit a linear model (soft-clip proportion $\sim$ PCA $-$ based genotype) in `R` at
each position treating genotypes AA, AB, and BB as 0, 1, and 2. Based on the significance of
genotype and R^2 of the linear models, we generated a list of 14 positions at which soft-clip
proportion was significantly associated with genotype of the putative inversions. We visualised
the distribution of the soft-clip proportion at these positions (Sup. Fig. 13) and selected six
positions for which the soft-clip proportion of BB was high enough and that of AB was around a
half of BB based on the assumption that soft clip reads covering an inversion breakpoint should
originate from haplotype B and non-soft clip reads should originate from haplotype A (Sup.

Table 5). To investigate whether some of these six positions represent inversion breakpoints, we asked whether the soft-clipped segments of the reads have homologous sequences at the other end of the outlier regions. We extracted soft-clipped segments of reads mapped at the focal six positions in AB and BB individuals using a custom script, and re-mapped these segments (instead of the entire reads) to the blackcap reference using `BWA mem`. We computed the depth of mapped segments in each position using `SAMtools` (Sup. Table 5).

10x linked read

We used an independent set of blackcap individuals (hereafter "10x individuals") whose genomes were sequenced with the 10x linked-read technology (Delmore et al., 2023, NCBI BioProject PRJEB65115). We genotyped the 10x individuals at the two putative inversion loci (i.e. AA, AB, or BB) based on genotypes at diagnostic SNP positions. We started by determining diagnostic SNP positions using our Illumina short read-based resequence data. Because usable diagnostic SNP positions should have genotypes perfectly associated with PCA-based genotype, we focused on positions at which F_{ST} was 1 between AA and BB, and all AB were heterozygous, using `VCFtools` and `BCFtools`. We also recorded mapping between an allele at the diagnostic positions and a genotype of the putative inversion ("A- and B-diagnostic alleles", e.g. G for haplotype A, T for haplotype B).

We then counted the number of sites with A- and B-diagnostic allele in each of 10x samples. To convert coordinates of 10x assemblies to the reference coordinate, we mapped the 10x pseudo-haplotyped assemblies to the blackcap reference using `minimap2` (Li, 2018). To determine the putative inversion genotype in the 10x individuals, we counted the number of positions with A-diagnostic and B-diagnostic alleles for each 10x pseudo-haplotype, and calculated the proportion of sites with A-diagnostic and B-diagnostic sites. In principle, an AA and a BB individual respectively are expected to have proportion of 100% and 0% of A-diagnostic sites in both of two pseudo-haplotypes, while an AB individual is expected to have 100% of A-diagnostic sites in one pseudo-haplotype and 0% for the other. For genotyping, we set the following three thresholds.

1. Missingness at the diagnostic positions is less than 10%, after removing positions with

⁶⁰⁵ non-unique `minimap2` mapping (i.e. at least 90% of all diagnostic positions should have

⁶⁰⁶ depth of 1x).

⁶⁰⁷ 2. More than 90% of all diagnostic sites should agree per pseudo-haplotype.

⁶⁰⁸ 3. The second criterion should be fulfilled for both pseudo-haplotypes of an individual.

⁶⁰⁹ We identified two BB individuals for each of the putative inversions on chromosomes 12

⁶¹⁰ and 30. There were no AB individuals passing the above threshold, indicating 10x pseudo-

⁶¹¹ haplotyping is not accurate in separating two diverged non-recombining alleles at a long range

⁶¹² in an individual that has both. To identify breakpoints, we aligned the pseudo-haplotype

⁶¹³ assemblies of these BB individuals as well as one AA individual for each putative inversion to

⁶¹⁴ the blackcap reference using `Nucmer4` (Marçais et al., 2018), and generated dot plots (Sup.

⁶¹⁵ Fig. 14).

Sequence analysis at breakpoint of putative inversion on chromosome 12

⁶¹⁷ 10x contigs of pseudo-haplotype B aligned next to the putative breakpoint position of

⁶¹⁸ blackcap reference chromosome 12 had an un-aligned flanking sequence. To characterise the

⁶¹⁹ DNA sequence of these flanking segments, we extracted the flanking sequences using `SAMtools`,

⁶²⁰ aligned the sequences to themselves using `minimap2`, and generated self-dot plots (Sup. Fig.

⁶²¹ 15), revealing presence of tandem repeats. To identify unit of tandem repeats within the

⁶²² flanking sequences, we ran `TandemRepeatsFinder` against these extracted sequences, resulting

⁶²³ in four consensus unit sequences of 144 bp based on two contigs from two individuals. To

⁶²⁴ confirm that the four consensus sequences represent the same tandem repeat (because the unit

⁶²⁵ of identical tandem repeat can have different phases), we ran `BLASTn` (version 2.10.1, Altschul

⁶²⁶ et al., 1990) with each consensus as query against dimers of the consensus. To investigate

⁶²⁷ whether the tandem repeat found at the putative breakpoint of chromosome 12 in haplotype

⁶²⁸ B is present in chromosome 12 and other chromosomes of the reference and corresponding

⁶²⁹ position of haplotype A, we ran `BLASTn` with the 144 bp consensus of the tandem repeat unit as

⁶³⁰ the query against blackcap reference and a contig of an AA individual that spans the breakpoint

⁶³¹ position, and counted how many copies were found in each reference chromosome/scaffold and

⁶³² the 10x contig (Sup. Fig. 16).

Selection in blackcaps

To test for selection in different outlier regions and to compare them with the genome-wide base line, we computed nucleotide diversity (π) and Tajima's D in 10 kb sliding windows per population using `PopGenome` (Pfeifer et al., 2014) and `VCFtools` (Danecek et al., 2011) respectively. The effects of the outlier regions on these statistics were tested using a linear mixed effects model (`nlme::lme` (Pinheiro et al., 2021)) and a generalised linear mixed effects model with a Gamma distribution (`lme4::glmer` (Bates et al., 2015)). To test for selection in genes d_N/d_S were computed following the counting method by Nei & Gojobori (1986). Gene annotation of the blackcap was obtained from Bascón-Cardozo et al. (2022b).

Tandem repeats within and outside outlier regions

To characterise correlation between outlier regions with distinct patterns of genetic variation and tandem repeats, we identified tandem repeats in the reference genome and compared the distribution of the tandem repeats with genomic regions with distinct patterns of genetic variation. First, `TandemRepeatsFinder` (Benson, 1999) was run on the blackcap reference genome with the parameter set recommended on the documentation (`trf </path/to/fasta> 2 7 7 80 10 50 500 -f -d -m -h`). The output was formatted and summarised for visualisation using custom scripts. Briefly, distribution of tandem repeats with a different unit size along the genome was summarised in 100 kb sliding windows in blocks of repeat unit sizes of 10 bp step (Sup. Fig. 26). Tandem repeats with the six longest repeat unit size were extracted per chromosome, and copy number for each tandem repeat was counted (Sup. Fig. 27).

Next, we tested whether the number of tandem repeats with long repeat unit were enriched in outlier regions at species-wide and population-specific low-recombining regions. We extracted tandem repeats with repeat unit size greater than or equal to 150 bp, and counted the number of tandem repeats (instead of total copy number) within and outside outlier regions. We performed Fisher's exact tests to test independence between the number of long tandem repeats and the mode of recombination suppression (species-wide/population-specific) (Sup. Table 7) using `fisher.test` function in R.

Simulation

Validation of LD-based inference of recombination landscape using non-randomly selected sample

Effects of recombination suppression model on recombination rate inference at an inversion

Three clusters of individuals observed in PCA at five outlier regions indicate presence of distinct haplotypes. Polymorphic inversions are known to show this pattern due to suppression of recombination between the normal and inverted alleles (Wellenreuther & Bernatchez, 2018). To test whether some of the five outlier regions represent polymorphic inversions, we intended to infer recombination rates using AA, AB, and BB individuals separately based on linkage disequilibrium (LD) patterns. Before addressing this in blackcaps empirically, we assessed how different types of recombination suppression at a haplotype block affect inference of recombination landscape using a set of individuals with a certain combination of haplotypes. To investigate the effect of a genotype-specific suppression of recombination on LD-based inference of recombination rate, we simulated different modes of recombination suppression using `SLiM` version 3.5 (Haller & Messer, 2019) under six scenarios listed in Sup. Table 6. Specifically, we performed 1,000 replicates of forward-time simulations of two 500 kb-long chromosomes with neutral mutation rate of 1×10^{-7} [per site per generation] and recombination rate of 1×10^{-6} [per site per generation] in a population of 1,000 diploid individuals under the Wright-Fisher model (We downscaled the population size and upscaled mutation rate to minimise the time and computational resource for simulation). We introduced a mutation (inversion marker) on one chromosome at 100 kb position at the 50th generation. We modelled an inversion by suppressing recombination in an interval from 100 kb to 400 kb position if the inversion marker site was heterozygous. We defined additional suppression according to different scenarios (models 1-6). To allow for the inversion to remain in the population, we applied negative frequency-dependent selection (fitness of inversion is $1 - (p_{inv} - 0.2)$ for models 1-3 and $1 - (p_{inv} - 0.8)$ for models 4-6 where p_{inv} is the frequency of the inversion allele). 1,000 generations after the inversion event, we recorded the mutations in all samples, making a VCF file including all individuals. Although 1,000 generations is relatively short

given the population size of 1,000, the haplotype structure at the inversion locus was stable in test runs of model-1 (inversion frequency of 0.2 without additional recombination suppression). Based on the genotype at the marker, we randomly sampled 10 individuals for each inversion genotype. `Pyrho` was run to estimate recombination rates using the sampled 10 individuals, with the block penalty 50 and window size 50. The inferred recombination landscape is in Sup. Fig. 11.

Species-wide reduction of recombination rate

To investigate how species-wide low-recombining regions affect patterns of local genetic variation depicted in local PCA, we performed forward simulation with `SLiM` version 4.0.1 (Haller & Messer, 2022). We simulated 100 replicates of two 500 kb-long chromosomes with neutral mutation rate of 1×10^{-7} [per site per generation] and recombination rate of 1×10^{-6} [per site per generation] except for an interval from 100 to 400 [kb] of the first chromosome where recombination rate was set to 1×10^{-9}, which is 1/1000 of the normally recombining chromosome. First, we ran a burn-in of 4,000 generations for an ancestral population of 1,000 diploids. After the burn-in, we made three populations of 1,000 diploids (pop1, pop2, and pop3) split from the ancestral population. We sampled 50 individuals per population every 20 generations over 1,000 generations after the population split and recorded SNPs in VCF. For each time point of each of 100 simulation replicates, we performed PCA with `PLINK`, using SNPs either within 100 to 400 [kb] of the first chromosome (pop1-specific suppression) or the normally recombining chromosome.

We investigated how reduced recombination rate affects representation of population subdivision in local PCA. To evaluate whether the individuals from different populations were distributed differently in local PCA at the low-recombining region, we performed Fasano-Franceschini test (Fasano & Franceschini, 1987), which is a multi-dimensional extension of Kolmogorov-Smirnov test, in three pairs of populations (pop1-pop2, pop1-pop3, pop2-pop3). We counted the number of significant pairs of populations (0, 1, 2, or 3) for each time point of each replicate. We compared between the low-recombining and normally recombining regions the number of pairs of populations with distinct distribution in PCA (Sup. Fig. 24).

Population-specific reduction of recombination rate

To investigate how evolution of low-recombining regions in population(s) affect patterns of local genetic variation depicted in local PCA, we performed forward simulation with `SLiM` version 4.0.1. We simulated two 500kb-long chromosomes with neutral mutation rate and recombination rate of 1×10^{-7} [per site per generation] and 1×10^{-6} [per site per generation]. First, we ran a burn-in of 4,000 generations for an ancestral population of 1,000 diploids. After the burn-in, we made three populations of 1,000 diploids (pop1, pop2, and pop3) split from the ancestral population, after which gene flow between all pairs of populations were set to 0.0025. We introduced recombination suppression in pop1 from 100 to 400 [kb] of the first chromosome in two scenarios. In the first scenario, recombination suppression was introduced at the same time of the split. In the second scenario, recombination suppression was introduced 4,000 generations after the population split event, allowing the three populations to differentiate before population-specific recombination suppression was introduced in pop1. We sampled 50 individuals per population every 20 generations over 1,000 generations after the introduction of the population-specific suppression of recombination and recorded SNPs in VCF. For each time point of each of 1,000 simulation replicates, we performed PCA with `PLINK`, using SNPs either within 100 to 400 [kb] of the first chromosome (pop1-specific suppression) or the normally recombining chromosome.

To characterise factors represented in the primary axes of distinct local PCA at population-specific low-recombining regions, we performed one replicate of `SLiM` simulation with the same scenarios of models 1 and 2 recording the full ancestry and mutations in tree sequence, with an increased duration of burn-in (40,000 generations) to make sure that all lineages at sampling time coalesce. We loaded the tree sequence with mutations in `tskit` (Kelleher et al., 2018) and sampled 50 diploids per population, and saved SNPs in VCF. Using the VCF files for each time point for each model, we performed PCA using `PLINK` at the population-specific low-recombining region, and determined one time point per model showing typical spread of individuals from the low-recombining population in PCA (Sup. Fig. 20A, E). For these PCAs we identified 5% SNPs with the highest loadings to the first two PC axes. We analysed these mutations on the underlying genealogies using `tskit`. Specifically, we investigated

 whether mutations originating from the low-recombining population were enriched in the
 high-loading mutations (Sup. Fig. 20C, G) with a χ^2 test. We also assessed whether multiple
 mutations originating in the low-recombining population occurring on the same genealogical
 branches (i.e. mutations on the same ancestral haplotypes) were enriched in the high-loading
 mutations (Sup. Fig. 20D, H). For this, we compared the number of mutations sharing
 the same genealogical branches among the high-loading mutations originating from the low-
 recombining population and the same number of randomly-selected mutations originating from
 the low-recombining population by a Kolmogorov-Smirnov test.

Effects of linked selection on local PCA

Background selection

To investigate the linked effect of purifying selection at low-recombining regions (back-
ground selection) on patterns of local genetic variation represented in local PCA, we performed
forward simulation with `SLiM` version 4.0.1. We simulated a species-wide low-recombining
region in three populations as described above, except we changed the distribution of fitness
effect of mutations with three different ratios between neutral ("n", $s = 0$) and deleterious
("d", $s = -0.05$ and $h = 0.5$) mutations of $n/(n + d) = 0, 0.25, 0.5, 0.75$. To evaluate whether
individuals from different populations were distributed differently in the local PCA at the
low-recombining region, we performed Fasano-Franceschini test between three pairs of popu-
lations (pop1-pop2, pop1-pop3, pop2-pop3). We counted the number of significant pairs of
populations (0, 1, 2, or 3) for each sampled time point of each replicate (out of 100) for each
DFE (Sup. Fig. 24).

Positive selection

To investigate the linked effect of positive selection at low-recombining regions on patterns
of local genetic variation represented in local PCA, we performed forward simulation with `SLiM`
version 4.0.1 under four scenarios: population-specific sweep and sweep before populations
split, with and without reduced local recombination rate. We simulated 10 replicates of one
500 kb-long chromosome with neutral mutation rate of 1×10^{-7} [per site per generation]

and recombination rate of 1×10^{-6} [per site per generation]. In scenarios with reduced recombination rate, we introduced a reduced recombination rate within an interval from 100 to 400 [kb] of the chromosome where recombination rate was set to 1×10^{-9}, which is 1/1000 of the normally recombining regions. For all scenarios, we ran a burn-in of 4,000 generations for an ancestral population of 1,000 diploids. In the scenarios with population-specific sweep, we made three populations of 1,000 diploids (pop1, pop2, and pop3) split from the ancestral population at the 4000-th generation. We introduced a strongly beneficial mutation ($s = 1$ and $h = 0.5$) in the middle of a chromosome of one randomly selected sample of the first population at the 100-th generation after the populations split. In the scenarios with sweep before split, we introduced a strongly beneficial mutation ($s = 1$ and $h = 0.5$) in the middle of the chromosome of one randomly selected sample of the ancestral population, and made the three populations of 1,000 diploids split at the 100-th generation after the introduction of the beneficial mutation. We sampled 100 diploid individuals per population every 20 generations since the introduction of the beneficial mutation (scenarios of population-specific sweep) or the split (scenarios of ancestral sweep) and recorded the SNPs in VCF format. We performed PCA using `PLINK`.

Acknowledgment

This work was supported by the Max Planck Society (Max Planck Research Group grant MFFALIMN0001 to ML), the DFG (project Z02 and Nav05 within SFB 1372 – Magnetoreception and Navigation in Vertebrates to ML), and DFG Research Infrastructure NGS_CC (project 407495230) as part of the Next Generation Sequencing Competence Network (project 423957469). AR was supported by the Intramural Research Program of the NHGRI, NIH (1ZIAHG200398). We thank Britta Meyer, Tianhao Zhao, Hanna Koch, Conny Burghardt, and Sven Künzel for DNA extraction, library preparation, and/or sequencing. We are grateful to Julien Dutheil, Diethard Tautz, Linda Odenthal-Hesse, Tobias Kaiser, Carolina Peralta, and Matthias Weissensteiner for constructive discussion. We are grateful to Thord Fransson, Christos Barboutis, Zura Javakhishvili, Martim Melo, Álvaro Ramírez, and Helena Batalha for providing us with samples. Permits were provided to JCI for samples collected on Cape

Verde (Ministerio do Ambiente - Habitacao e Ordenamento do Territorio, 18/CITES/DNA, 17 Dec 2015), Canary Islands (Ref.: 2012/0710), Madeira (Ref.: 02/2016), and the Azores (Instituto da Conservacao da Natureza e da Biodiversidade, 171/2008, 31 Mar 2009); to JP-T for samples collected on Mallorca (CAP 64/2009); to Thord Fransson for samples collected on Crete (6Υ0Ξ4653Π8-ΥΓ5 issued by the Hellenic Ministry of Environment and Energy), and to Zura Javakhishvili for samples collected in Georgia (889-0-2-202303291450 by the Ministry of Environment and Agriculture of Georgia).

Data availability

The primary and alternate haplotype assemblies of the blackcap reference genome can be found under NCBI BioProject PRJNA558064 (accenssion GCA_009819655.1) and PRJNA558065 (accession GCA_009819715.1). Raw Illumina reads for the resequencing data can be accessed under NCBI BioProject PRJEB66075 (SRA accession ERP151147). Processed data and scripts for analysis and simulation are found in Zenodo (https://doi.org/10.5281/zenodo.8358874).

Conflict of interest

The authors declare no conflict of interest.

Author contributions

JI and ML designed the study. Reference genome was generated by JF, AR, JM, BH, WC, JC, KH, MU, OF, and EDJ. JP-T and JCI collected samples for resequencing. AB performed read mapping, variant calling, and data filtration. KB-C inferred recombination maps. JI conducted haplotype inference, population genomics analyses, simulations, sequence analyses, statistical modelling, and data visualisation. JI and ML wrote the manuscript with inputs from other authors.

References

Altschul, S. F., Gish, W., Miller, W., Myers, E. W., & Lipman, D. J. (1990). Basic local alignment search tool. *Journal of Molecular Biology*, *215*(3), 403–410. https://doi.org/10.1016/S0022-2836(05)80360-2

Auton, A., Li, Y. R., Kidd, J., Oliveira, K., Nadel, J., Holloway, J. K., Hayward, J. J., Cohen, P. E., Greally, J. M., Wang, J., Bustamante, C. D., & Boyko, A. R. (2013). Genetic Recombination Is Targeted towards Gene Promoter Regions in Dogs. *PLOS Genetics*, *9*(12), e1003984. https://doi.org/10.1371/journal.pgen.1003984

Baker, Z., Schumer, M., Haba, Y., Bashkirova, L., Holland, C., Rosenthal, G. G., & Przeworski, M. (2017). Repeated losses of PRDM9-directed recombination despite the conservation of PRDM9 across vertebrates. *eLife*, *6*, e24133. https://doi.org/10.7554/eLife.24133

Barton, N. H., & Etheridge, A. M. (2004). The Effect of Selection on Genealogies. *Genetics*, *166*(2), 1115–1131. https://doi.org/10.1093/genetics/166.2.1115

Bascón-Cardozo, K., Bours, A., Ishigohoka, J., Odenthal-Hesse, L., & Liedvogel, M. (2022a). *Historical recombination maps diverge between Eurasian blackcap populations with distinct migratory strategies* [Preprint]. Authorea. https://doi.org/10.22541/au.166790167.72861799/v1

Bascón-Cardozo, K., Bours, A., Manthey, G., Pruisscher, P., Durieux, G., Dutheil, J., Odenthal-Hesse, L., & Liedvogel, M. (2022b). *Fine-scale map reveals highly variable recombination rates associated with genomic features in the European blackcap* [Preprint]. Authorea. https://doi.org/10.22541/au.165423614.49331155/v1

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software*, *67*(1), 1–48. https://doi.org/10.18637/jss.v067.i01

Baudat, F., Buard, J., Grey, C., Fledel-Alon, A., Ober, C., Przeworski, M., Coop, G., & Massy, B. de. (2010). PRDM9 Is a Major Determinant of Meiotic Recombination Hotspots in Humans and Mice. *Science*, *327*(5967), 836–840. https://doi.org/10.1126/science.1183439

Benson, G. (1999). Tandem repeats finder: A program to analyze DNA sequences. *Nucleic Acids Research*, *27*(2), 573–580. https://doi.org/10.1093/nar/27.2.573

Berthold, P. (1988). Evolutionary aspects of migratory behavior in European warblers. *Journal of Evolutionary Biology*, *1*(3), 195–209. https://doi.org/https://doi.org/10.1046/j.1420-9101.1998.1030195.x

Berthold, P. (1991). Genetic control of migratory behaviour in birds. *Trends in Ecology and Evolution*, *6*(8), 254–257. https://doi.org/10.1016/0169-5347(91)90072-6

Bhatia, G., Patterson, N., Sankararaman, S., & Price, A. L. (2013). Estimating and interpreting FST: The impact of rare variants. *Genome Research*, *23*(9), 1514–1521. https://doi.org/10.1101/gr.154831.113

Birtle, Z., & Ponting, C. P. (2006). Meisetz and the birth of the KRAB motif. *Bioinformatics*,

22(23), 2841–2845. https://doi.org/10.1093/bioinformatics/btl498

Burri, R. (2017). Interpreting differentiation landscapes in the light of long-term linked selection.
Evolution Letters, *1*(3), 118–131. https://doi.org/https://doi.org/10.1002/evl3.14

Burri, R., Nater, A., Kawakami, T., Mugal, C. F., Olason, P. I., Smeds, L., Suh, A., Dutoit,
L., Bureš, S., Garamszegi, L. Z., Hogner, S., Moreno, J., Qvarnström, A., Ružić, M.,
Sæther, S.-A., Sætre, G.-P., Török, J., & Ellegren, H. (2015). Linked selection and
recombination rate variation drive the evolution of the genomic landscape of differentiation
across the speciation continuum of Ficedula flycatchers. *Genome Research*, *25*(11), 1656–
1665. https://doi.org/10.1101/gr.196485.115

Cao, Y., Li, L., Xu, M., Feng, Z., Sun, X., Lu, J., Xu, Y., Du, P., Wang, T., Hu, R., Ye,
Z., Shi, L., Tang, X., Yan, L., Gao, Z., Chen, G., Zhang, Y., Chen, L., Ning, G., et al.
(2020). The ChinaMAP analytics of deep whole genome sequences in 10,588 individuals.
Cell Research, *30*(9), 717–731. https://doi.org/10.1038/s41422-020-0322-9

Cruickshank, T. E., & Hahn, M. W. (2014). Reanalysis suggests that genomic islands of
speciation are due to reduced diversity, not reduced gene flow. *Molecular Ecology*, *23*(13),
3133–3157. https://doi.org/https://doi.org/10.1111/mec.12796

Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A., Handsaker,
R. E., Lunter, G., Marth, G. T., Sherry, S. T., McVean, G., Durbin, R., & 1000 Genomes
Project Analysis Group. (2011). The variant call format and VCFtools. *Bioinformatics*,
27(15), 2156–2158. https://doi.org/10.1093/bioinformatics/btr330

Danecek, P., Bonfield, J. K., Liddle, J., Marshall, J., Ohan, V., Pollard, M. O., Whitwham,
A., Keane, T., McCarthy, S. A., Davies, R. M., & Li, H. (2021). Twelve years of SAMtools
and BCFtools. *GigaScience*, *10*(giab008). https://doi.org/10.1093/gigascience/giab008

DeGiorgio, M., Huber, C. D., Hubisz, M. J., Hellmann, I., & Nielsen, R. (2016). SweepFinder2:
Increased sensitivity, robustness and flexibility. *Bioinformatics*, *32*(12), 1895–1897. https:
//doi.org/10.1093/bioinformatics/btw051

Delaneau, O., Zagury, J.-F., & Marchini, J. (2013). Improved whole-chromosome phasing for
disease and population genetic studies. *Nature Methods*, *10*(1), 5–6. https://doi.org/10.
1038/nmeth.2307

Delmore, K., Doren, B. M. V., Ullrich, K., Curk, T., Jeugd, H. P. van der, & Liedvogel, M.
(2023). *Structural genomic variation and migratory behavior in wild songbirds* [Preprint].
bioRxiv. https://doi.org/10.1101/2023.04.24.538030

Delmore, K. E., Lugo Ramos, J. S., Van Doren, B. M., Lundberg, M., Bensch, S., Irwin, D. E.,
& Liedvogel, M. (2018). Comparative analysis examining patterns of genomic differentiation
across multiple episodes of population divergence in birds. *Evolution Letters*, *2*(2), 76–87.
https://doi.org/10.1002/evl3.46

Delmore, K. E., Van Doren, B. M., Conway, G. J., Curk, T., Garrido-Garduño, T., Germain,
R. R., Hasselmann, T., Hiemer, D., Jeugd, H. P. van der, Justen, H., Lugo Ramos, J. S.,
Maggini, I., Meyer, B. S., Phillips, R. J., Remisiewicz, M., Roberts, G. C. M., Sheldon,
B. C., Vogl, W., & Liedvogel, M. (2020a). Individual variability and versatility in an
eco-evolutionary model of avian migration. *Proceedings of the Royal Society B: Biological*

Sciences, *287*(1938), 20201339. https://doi.org/10.1098/rspb.2020.1339

Delmore, K., Illera, J. C., Pérez-Tris, J., Segelbacher, G., Ramos, J. S., Durieux, G., Ishigohoka, J., & Liedvogel, M. (2020b). The evolutionary history and genomics of european blackcap migration. *eLife*, *9*, e54462. https://doi.org/10.7554/eLife.54462

Dutheil, J. Y., Ganapathy, G., Hobolth, A., Mailund, T., Uyenoyama, M. K., & Schierup, M. H. (2009). Ancestral Population Genomics: The Coalescent Hidden Markov Model Approach. *Genetics*, *183*(1), 259–274. https://doi.org/10.1534/genetics.109.103010

Ewels, P., Magnusson, M., Lundin, S., & Käller, M. (2016). MultiQC: Summarize analysis results for multiple tools and samples in a single report. *Bioinformatics*, *32*(19), 3047–3048. https://doi.org/10.1093/bioinformatics/btw354

Faria, R., Johannesson, K., Butlin, R. K., & Westram, A. M. (2019). Evolving Inversions. *Trends in Ecology & Evolution*, *34*(3), 239–248. https://doi.org/10.1016/j.tree.2018.12.005

Fasano, G., & Franceschini, A. (1987). A multidimensional version of the Kolmogorov–Smirnov test. *Monthly Notices of the Royal Astronomical Society*, *225*(1), 155–170. https://doi.org/10.1093/mnras/225.1.155

Fay, J. C., & Wu, C.-I. (2000). Hitchhiking Under Positive Darwinian Selection. *Genetics*, *155*(3), 1405–1413. https://doi.org/10.1093/genetics/155.3.1405

Fedorova, S. A., Reidla, M., Metspalu, E., Metspalu, M., Rootsi, S., Tambets, K., Trofimova, N., Zhadanov, S. I., Kashani, B. H., Olivieri, A., Voevoda, M. I., Osipova, L. P., Platonov, F. A., Tomsky, M. I., Khusnutdinova, E. K., Torroni, A., & Villems, R. (2013). Autosomal and uniparental portraits of the native populations of Sakha (Yakutia): Implications for the peopling of Northeast Eurasia. *BMC Evolutionary Biology*, *13*(1), 127. https://doi.org/10.1186/1471-2148-13-127

Ghurye, J., Rhie, A., Walenz, B. P., Schmitt, A., Selvaraj, S., Pop, M., Phillippy, A. M., & Koren, S. (2019). Integrating Hi-C links with assembly graphs for chromosome-scale assembly. *PLOS Computational Biology*, *15*(8), e1007273. https://doi.org/10.1371/journal.pcbi.1007273

Guan, D., McCarthy, S. A., Wood, J., Howe, K., Wang, Y., & Durbin, R. (2020). Identifying and removing haplotypic duplication in primary genome assemblies. *Bioinformatics*, *36*(9), 2896–2898. https://doi.org/10.1093/bioinformatics/btaa025

Guerrero, R. F., Rousset, F., & Kirkpatrick, M. (2012). Coalescent patterns for chromosomal inversions in divergent populations. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *367*(1587), 430–438. https://doi.org/10.1098/rstb.2011.0246

Hager, E. R., Harringmeyer, O. S., Wooldridge, T. B., Theingi, S., Gable, J. T., McFadden, S., Neugeboren, B., Turner, K. M., Jensen, J. D., & Hoekstra, H. E. (2022). A chromosomal inversion contributes to divergence in multiple traits between deer mouse ecotypes. *Science*, *377*(6604), 399–405. https://doi.org/10.1126/science.abg0718

Haller, B. C., & Messer, P. W. (2019). SLiM 3: Forward Genetic Simulations Beyond the Wright–Fisher Model. *Molecular Biology and Evolution*, *36*(3), 632–637. https://doi.org/10.1093/molbev/msy228

Haller, B. C., & Messer, P. W. (2022). SLiM 4: Multispecies Eco-Evolutionary Modeling. *The American Naturalist*, E000–E000. https://doi.org/10.1086/723601

Hejase, H. A., Salman-Minkov, A., Campagna, L., Hubisz, M. J., Lovette, I. J., Gronau, I., & Siepel, A. (2020). Genomic islands of differentiation in a rapid avian radiation have been driven by recent selective sweeps. *Proceedings of the National Academy of Sciences*, *117*(48), 30554–30565. https://doi.org/10.1073/pnas.2015987117

Helbig, A. J. (1991). Inheritance of migratory direction in a bird species: A cross-breeding experiment with SE- and SW-migrating blackcaps (*Sylvia atricapilla*). *Behavioral Ecology and Sociobiology*, *28*(1), 9–12. https://doi.org/10.1007/BF00172133

Howe, K., Chow, W., Collins, J., Pelan, S., Pointon, D.-L., Sims, Y., Torrance, J., Tracey, A., & Wood, J. (2021). Significantly improving the quality of genome assemblies through curation. *GigaScience*, *10*(1), giaa153. https://doi.org/10.1093/gigascience/giaa153

Huang, K., Andrew, R. L., Owens, G. L., Ostevik, K. L., & Rieseberg, L. H. (2020). Multiple chromosomal inversions contribute to adaptive divergence of a dune sunflower ecotype. *Molecular Ecology*, *29*(14), 2535–2549. https://doi.org/https://doi.org/10.1111/mec.15428

Hudson, R. R. (1983). Properties of a neutral allele model with intragenic recombination. *Theoretical Population Biology*, *23*(2), 183–201. https://doi.org/10.1016/0040-5809(83)90013-8

Irwin, D. E., Milá, B., Toews, D. P. L., Brelsford, A., Kenyon, H. L., Porter, A. N., Grossen, C., Delmore, K. E., Alcaide, M., & Irwin, J. H. (2018). A comparison of genomic islands of differentiation across three young avian species pairs. *Molecular Ecology*, *27*(23), 4839–4855. https://doi.org/https://doi.org/10.1111/mec.14858

Kelleher, J., Thornton, K. R., Ashander, J., & Ralph, P. L. (2018). Efficient pedigree recording for fast population genetics simulation. *PLOS Computational Biology*, *14*(11), e1006581. https://doi.org/10.1371/journal.pcbi.1006581

Kong, A., Thorleifsson, G., Gudbjartsson, D. F., Masson, G., Sigurdsson, A., Jonasdottir, A., Walters, G. B., Jonasdottir, A., Gylfason, A., Kristinsson, K. T., Gudjonsson, S. A., Frigge, M. L., Helgason, A., Thorsteinsdottir, U., & Stefansson, K. (2010). Fine-scale recombination rate differences between sexes, populations and individuals. *Nature*, *467*(7319), 1099–1103. https://doi.org/10.1038/nature09525

Kronforst, M. R., Hansen, M. E. B., Crawford, N. G., Gallant, J. R., Zhang, W., Kulathinal, R. J., Kapan, D. D., & Mullen, S. P. (2013). Hybridization Reveals the Evolving Genomic Architecture of Speciation. *Cell Reports*, *5*(3), 666–677. https://doi.org/10.1016/j.celrep.2013.09.042

Li, H. (2013). *Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM* [Preprint]. arXiv. https://doi.org/https://doi.org/10.48550/arXiv.1303.3997

Li, H. (2018). Minimap2: Pairwise alignment for nucleotide sequences. *Bioinformatics*, *34*(18), 3094–3100. https://doi.org/10.1093/bioinformatics/bty191

Li, H., & Ralph, P. (2019). Local PCA Shows How the Effect of Population Structure Differs Along the Genome. *Genetics*, *211*(1), 289–304. https://doi.org/10.1534/genetics.118.301747

Lindenbaum, P. (2015). *JVarkit: Java-based utilities for Bioinformatics*. https://doi.org/10.6084/m9.figshare.1425030.v1

Lotterhos, K. E. (2019). The Effect of Neutral Recombination Variation on Genome Scans for Selection. *G3 GenesGenomesGenetics*, *9*(6), 1851–1867. https://doi.org/10.1534/g3.119.400088

Ma, J., & Amos, C. I. (2012). Investigation of Inversion Polymorphisms in the Human Genome Using Principal Components Analysis. *PLOS ONE*, *7*(7), e40224. https://doi.org/10.1371/journal.pone.0040224

Malinsky, M., Challis, R. J., Tyers, A. M., Schiffels, S., Terai, Y., Ngatunga, B. P., Miska, E. A., Durbin, R., Genner, M. J., & Turner, G. F. (2015). Genomic islands of speciation separate cichlid ecomorphs in an East African crater lake. *Science*, *350*(6267), 1493–1498. https://doi.org/10.1126/science.aac9927

Marçais, G., Delcher, A. L., Phillippy, A. M., Coston, R., Salzberg, S. L., & Zimin, A. (2018). MUMmer4: A fast and versatile genome alignment system. *PLOS Computational Biology*, *14*(1), e1005944. https://doi.org/10.1371/journal.pcbi.1005944

Martin, S. H., Davey, J. W., & Jiggins, C. D. (2015). Evaluating the Use of ABBA–BABA Statistics to Locate Introgressed Loci. *Molecular Biology and Evolution*, *32*(1), 244–257. https://doi.org/10.1093/molbev/msu269

Martin, S. H., Davey, J. W., Salazar, C., & Jiggins, C. D. (2019). Recombination rate variation shapes barriers to introgression across butterfly genomes. *PLoS Biology*, *17*(2), 1–28. https://doi.org/10.1371/journal.pbio.2006288

Martin, S. H., & Van Belleghem, S. M. (2017). Exploring Evolutionary Relationships Across the Genome Using Topology Weighting. *Genetics*, *206*(1), 429–438. https://doi.org/10.1534/genetics.116.194720

McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A., Garimella, K., Altshuler, D., Gabriel, S., Daly, M., & DePristo, M. A. (2010). The Genome Analysis Toolkit: A MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Research*, *20*(9), 1297–1303. https://doi.org/10.1101/gr.107524.110

McVean, G. (2009). A Genealogical Interpretation of Principal Components Analysis. *PLOS Genetics*, *5*(10), e1000686. https://doi.org/10.1371/journal.pgen.1000686

McVean, G. A., & Cardin, N. J. (2005). Approximating the coalescent with recombination. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *360*(1459), 1387–1393. https://doi.org/10.1098/rstb.2005.1673

Mérot, C., Berdan, E. L., Cayuela, H., Djambazian, H., Ferchaud, A.-L., Laporte, M., Normandeau, E., Ragoussis, J., Wellenreuther, M., & Bernatchez, L. (2021). Locally Adaptive Inversions Modulate Genetic Variation at Different Geographic Scales in a Seaweed Fly. *Molecular Biology and Evolution*, *38*(9), 3953–3971. https://doi.org/10.1093/molbev/msab143

Myers, S., Freeman, C., Auton, A., Donnelly, P., & McVean, G. (2008). A common sequence motif associated with recombination hot spots and genome instability in humans. *Nature Genetics*, *40*(9), 1124–1129. https://doi.org/10.1038/ng.213

Nei, M., & Gojobori, T. (1986). Simple methods for estimating the numbers of synonymous and nonsynonymous nucleotide substitutions. *Molecular Biology and Evolution*, *3*(5), 418–426. https://doi.org/10.1093/oxfordjournals.molbev.a040410

Nielsen, R. (2005). Molecular Signatures of Natural Selection. *Annual Review of Genetics*, *39*(1), 197–218. https://doi.org/10.1146/annurev.genet.39.073003.112420

Noor, M., & Bennett, S. (2009). Islands of speciation or mirages in the desert? Examining the role of restricted recombination in maintaining species. *Heredity*, *103*, 439–444.

Okonechnikov, K., Conesa, A., & García-Alcalde, F. (2016). Qualimap 2: Advanced multi-sample quality control for high-throughput sequencing data. *Bioinformatics*, *32*(2), 292–294. https://doi.org/10.1093/bioinformatics/btv566

Pamilo, P., & Nei, M. (1988). Relationships between gene trees and species trees. *Molecular Biology and Evolution*, *5*(5), 568–583. https://doi.org/10.1093/oxfordjournals.molbev.a040517

Patterson, N., Moorjani, P., Luo, Y., Mallick, S., Rohland, N., Zhan, Y., Genschoreck, T., Webster, T., & Reich, D. (2012). Ancient Admixture in Human History. *Genetics*, *192*(3), 1065–1093. https://doi.org/10.1534/genetics.112.145037

Patterson, N., Price, A. L., & Reich, D. (2006). Population Structure and Eigenanalysis. *PLOS Genetics*, *2*(12), e190. https://doi.org/10.1371/journal.pgen.0020190

Peter, B. M. (2016). Admixture, Population Structure, and F-Statistics. *Genetics*, *202*(4), 1485–1501. https://doi.org/10.1534/genetics.115.183913

Peter, B. M. (2022). A geometric relationship of F2, F3 and F4-statistics with principal component analysis. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *377*(1852), 20200413. https://doi.org/10.1098/rstb.2020.0413

Pfeifer, B., Wittelsbuerger, U., Ramos-Onsins, S. E., & Lercher, M. J. (2014). PopGenome: An Efficient Swiss Army Knife for Population Genomic Analyses in R. *Molecular Biology and Evolution*, *31*, 1929–1936. https://doi.org/10.1093/molbev/msu136

Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., & R Core Team. (2021). *Nlme: Linear and Nonlinear Mixed Effects Models*. https://CRAN.R-project.org/package=nlme

Pracana, R., Priyam, A., Levantis, I., Nichols, R. A., & Wurm, Y. (2017). The fire ant social chromosome supergene variant Sb shows low diversity but high divergence from SB. *Molecular Ecology*, *26*(11), 2864–2879. https://doi.org/10.1111/mec.14054

Price, A. L., Patterson, N. J., Plenge, R. M., Weinblatt, M. E., Shadick, N. A., & Reich, D. (2006). Principal components analysis corrects for stratification in genome-wide association studies. *Nature Genetics*, *38*(8), 904–909. https://doi.org/10.1038/ng1847

Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M. A. R., Bender, D., Maller, J., Sklar, P., Bakker, P. I. W. de, Daly, M. J., & Sham, P. C. (2007). PLINK: A Tool Set for Whole-Genome Association and Population-Based Linkage Analyses. *The American Journal of Human Genetics*, *81*(3), 559–575. https://doi.org/10.1086/519795

Ralph, P., Thornton, K., & Kelleher, J. (2020). Efficiently Summarizing Relationships in Large

Samples: A General Duality Between Statistics of Genealogies and Genomes. *Genetics*, *215*(3), 779–797. https://doi.org/10.1534/genetics.120.303253

Reich, D., Thangaraj, K., Patterson, N., Price, A. L., & Singh, L. (2009). Reconstructing Indian population history. *Nature*, *461*(7263), 489–494. https://doi.org/10.1038/nature08365

Renaut, S., Grassa, C. J., Yeaman, S., Moyers, B. T., Lai, Z., Kane, N. C., Bowers, J. E., Burke, J. M., & Rieseberg, L. H. (2013). Genomic islands of divergence are not affected by geography of speciation in sunflowers. *Nature Communications*, *4*(1), 1827. https://doi.org/10.1038/ncomms2833

Rhie, A., McCarthy, S. A., Fedrigo, O., Damas, J., Formenti, G., Koren, S., Uliano-Silva, M., Chow, W., Fungtammasan, A., Kim, J., Lee, C., Ko, B. J., Chaisson, M., Gedman, G. L., Cantin, L. J., Thibaud-Nissen, F., Haggerty, L., Bista, I., Smith, M., et al. (2021). Towards complete and error-free genome assemblies of all vertebrate species. *Nature*, *592*(7856), 737–746. https://doi.org/10.1038/s41586-021-03451-0

Rubin, C.-J., Enbody, E. D., Dobreva, M. P., Abzhanov, A., Davis, B. W., Lamichhaney, S., Pettersson, M., Sendell-Price, A. T., Sprehn, C. G., Valle, C. A., Vasco, K., Wallerman, O., Grant, B. R., Grant, P. R., & Andersson, L. (2022). Rapid adaptive radiation of Darwin's finches depends on ancestral genetic modules. *Science Advances*, *8*(27), eabm5982. https://doi.org/10.1126/sciadv.abm5982

Ruiz-Arenas, C., Cáceres, A., López-Sánchez, M., Tolosana, I., Pérez-Jurado, L., & González, J. R. (2019). scoreInvHap: Inversion genotyping for genome-wide association studies. *PLOS Genetics*, *15*(7), e1008203. https://doi.org/10.1371/journal.pgen.1008203

Sabeti, P. C., Reich, D. E., Higgins, J. M., Levine, H. Z. P., Richter, D. J., Schaffner, S. F., Gabriel, S. B., Platko, J. V., Patterson, N. J., McDonald, G. J., Ackerman, H. C., Campbell, S. J., Altshuler, D., Cooper, R., Kwiatkowski, D., Ward, R., & Lander, E. S. (2002). Detecting recent positive selection in the human genome from haplotype structure. *Nature*, *419*(6909), 832–837. https://doi.org/10.1038/nature01140

Sabeti, P. C., Varilly, P., Fry, B., Lohmueller, J., Hostetter, E., Cotsapas, C., Xie, X., Byrne, E. H., McCarroll, S. A., Gaudet, R., Schaffner, S. F., Lander, E. S., Frazer, K. A., Ballinger, D. G., Cox, D. R., Hinds, D. A., Stuve, L. L., Gibbs, R. A., Belmont, J. W., et al. (2007). Genome-wide detection and characterization of positive selection in human populations. *Nature*, *449*(7164), 913–918. https://doi.org/10.1038/nature06250

Setter, D., Mousset, S., Cheng, X., Nielsen, R., DeGiorgio, M., & Hermisson, J. (2020). VolcanoFinder: Genomic scans for adaptive introgression. *PLOS Genetics*, *16*(6), e1008867. https://doi.org/10.1371/journal.pgen.1008867

Shao, C., Sun, S., Liu, K., Wang, J., Li, S., Liu, Q., Deagle, B. E., Seim, I., Biscontin, A., Wang, Q., Liu, X., Kawaguchi, S., Liu, Y., Jarman, S., Wang, Y., Wang, H.-Y., Huang, G., Hu, J., Feng, B., et al. (2023). The enormous repetitive Antarctic krill genome reveals environmental adaptations and population insights. *Cell*, *186*(6), 1279–1294.e19. https://doi.org/10.1016/j.cell.2023.02.005

Shipilina, D., Pal, A., Stankowski, S., Chan, Y. F., & Barton, N. H. (2023). On the origin and structure of haplotype blocks. *Molecular Ecology*, *32*(6), 1441–1457. https://doi.org/10.1111/mec.16793

Singhal, S., Leffler, E. M., Sannareddy, K., Turner, I., Venn, O., Hooper, D. M., Strand, A. I., Li, Q., Raney, B., Balakrishnan, C. N., Griffith, S. C., McVean, G., & Przeworski, M. (2015). Stable recombination hotspots in birds. *Science*, *350*(6263), 928–932. https://doi.org/10.1126/science.aad0843

Smeds, L., Qvarnström, A., & Ellegren, H. (2016). Direct estimate of the rate of germline mutation in a bird. *Genome Research*, *26*(9), 1211–1218. https://doi.org/10.1101/gr.204669.116

Smukowski Heil, C. S., Ellison, C., Dubin, M., & Noor, M. A. (2015). Recombining without Hotspots: A Comprehensive Evolutionary Portrait of Recombination in Two Closely Related Species of Drosophila. *Genome Biology and Evolution*, *7*(10), 2829–2842. https://doi.org/10.1093/gbe/evv182

Speidel, L., Forest, M., Shi, S., & Myers, S. R. (2019). A method for genome-wide genealogy estimation for thousands of samples. *Nature Genetics*, *51*(9), 1321–1329. https://doi.org/10.1038/s41588-019-0484-x

Spence, J. P., & Song, Y. S. (2019). Inference and analysis of population-specific fine-scale recombination maps across 26 diverse human populations. *Science Advances*, *5*(10), eaaw9206. https://doi.org/10.1126/sciadv.aaw9206

Stern, A. J., Wilton, P. R., & Nielsen, R. (2019). An approximate full-likelihood method for inferring selection and allele frequency trajectories from DNA sequence data. *PLoS Genetics*, *15*(9), 1–32. https://doi.org/10.1371/journal.pgen.1008384

Stevison, L. S., Hoehn, K. B., & Noor, M. A. F. (2011). Effects of Inversions on Within- and Between-Species Recombination and Divergence. *Genome Biology and Evolution*, *3*, 830–841. https://doi.org/10.1093/gbe/evr081

Stevison, L. S., Woerner, A. E., Kidd, J. M., Kelley, J. L., Veeramah, K. R., McManus, K. F., Great Ape Genome Project, Bustamante, C. D., Hammer, M. F., & Wall, J. D. (2016). The Time Scale of Recombination Rate Evolution in Great Apes. *Molecular Biology and Evolution*, *33*(4), 928–945. https://doi.org/10.1093/molbev/msv331

Tajima, F. (1989). Statistical method for testing the neutral mutation hypothesis by DNA polymorphism. *Genetics*, *123*(3), 585–595. https://doi.org/PMC1203831

Taylor, J. E. (2013). The effect of fluctuating selection on the genealogy at a linked site. *Theoretical Population Biology*, *87*, 34–50. https://doi.org/10.1016/j.tpb.2013.03.004

Todesco, M., Owens, G. L., Bercovich, N., Légaré, J.-S., Soudi, S., Burge, D. O., Huang, K., Ostevik, K. L., Drummond, E. B. M., Imerovski, I., Lande, K., Pascual-Robles, M. A., Nanavati, M., Jahani, M., Cheung, W., Staton, S. E., Muños, S., Nielsen, R., Donovan, L. A., et al. (2020). Massive haplotypes underlie ecotypic differentiation in sunflowers. *Nature*, *584*(7822), 602–607. https://doi.org/10.1038/s41586-020-2467-6

Voight, B. F., Kudaravalli, S., Wen, X., & Pritchard, J. K. (2006). A Map of Recent Positive Selection in the Human Genome. *PLOS Biology*, *4*(3), e72. https://doi.org/10.1371/journal.pbio.0040072

Wakeley, J. (2008). *Coalescent Theory: An Introduction* (1st ed.). W. H. Freeman.

Wakeley, J. (2020). Developments in coalescent theory from single loci to chromosomes. *Theoretical Population Biology*, *133*, 56–64. https://doi.org/10.1016/j.tpb.2020.02.002

Wellenreuther, M., & Bernatchez, L. (2018). Eco-Evolutionary Genomics of Chromosomal Inversions. *Trends in Ecology & Evolution*, *33*(6), 427–440. https://doi.org/10.1016/j.tree.2018.04.002

Wiuf, C., & Hein, J. (1999). Recombination as a Point Process along Sequences. *Theoretical Population Biology*, *55*(3), 248–259. https://doi.org/10.1006/tpbi.1998.1403

Wolf, J. B. W., & Ellegren, H. (2017). Making sense of genomic islands of differentiation in light of speciation. *Nature Reviews Genetics*, *18*(2), 87–100. https://doi.org/10.1038/nrg.2016.133

Yi, X., Liang, Y., Huerta-Sanchez, E., Jin, X., Cuo, Z. X. P., Pool, J. E., Xu, X., Jiang, H., Vinckenbosch, N., Korneliussen, T. S., Zheng, H., Liu, T., He, W., Li, K., Luo, R., Nie, X., Wu, H., Zhao, M., Cao, H., et al. (2010). Sequencing of 50 Human Exomes Reveals Adaptation to High Altitude. *Science*, *329*(5987), 75–78. https://doi.org/10.1126/science.1190371