BK channels of five different subunit combinations underlie the *de novo* KCNMA1 G375R channelopathy

Yanyan Geng[1], Ping Li[2,4], Alice Butler[2] Bill Wang[2], Lawrence Salkoff[2,3,5], and Karl L. Magleby[1,5,*]

[1]Department of Physiology and Biophysics, University of Miami Miller School of Medicine,

Miami FL 33136, USA; [2]Department of Neuroscience and [3]Department of Genetics, Washington

University St. Louis, MO 63110, USA. [4]Present address: Department of OBGYN, Washington

University St. Louis, MO 63110, USA. [5]These authors contributed equally.

[*]Corresponding Author: Karl L. Magleby, Department of Physiology and Biophysics, University of

Miami Miller School of Medicine, Miami FL 33136

Email: kmagleby@miami.edu

Abstract

The molecular basis of a severe developmental and neurological disorder associated with a *de*
novo G375R variant of the tetrameric BK channel is unknown. Here we address this question by
recording from single BK channels expressed for a heterozygous G375R mutation. Five different
types of functional BK channels were observed: 3% were WT, 12% were homomeric mutant,
and 85% were three different types of hybrid channels. All channel types except WT showed a
marked gain-of-function in voltage activation and a smaller loss-of-function in single channel
conductance, with both becoming more pronounced as the number of mutant subunits per
tetrameric channel increased. The molecular phenotype suggested codominance for the two
homomeric channels and partial dominance for the hybrid channels. A model in which BK
channels are randomly assembled from mutant and WT subunits, with each subunit
contributing increments of activation and conductance, approximated the molecular phenotype
of the heterozygous G375R mutation.

Introduction

The BK channel (Slo1, KCa1.1) is a large conductance K^+ selective channel that is synergistically activated by Ca^{2+} and voltage[1-16]. BK channels are homotetrameric proteins comprised of four large pore-forming (alpha) subunits >1200 amino acids[15] (Supplementary Fig. 1) encoded by the *KCNMA1* gene. BK channels are widely expressed in many cell types where they modulate smooth muscle contraction[17], transmitter release[18], circadian rhythms[19], repetitive firing[20,21], and cellular excitability[22]. Mutations in the *KCNMA1* gene that encode the (alpha) subunit of BK channels are associated with a wide range of disease, including epilepsy, dyskinesis, autism, multiple congenital abnormalities, developmental delay, intellectual disability, axial hypotonia, ataxia, cerebral and cerebellar atrophy, bone thickening, and tortuosity of arteries[23-28].

Studies of the pathogenic properties of BK channels associated with disease have often been incomplete, focusing on the homomeric mutant channels. Yet, for a mutation heterozygous with the wild-type (WT) allele, mutant and WT subunits have the potential to assemble into five different stoichiometries for tetrameric channels with likely differences in functional properties (Fig. 1)[29-33].

Here we show that this is the case for a *de novo* G375R mutation in the (alpha) subunit of BK channels associated with the Liang-Wang Syndrome[25]. Three unrelated children who carried this mutation had a syndromic neurodevelopmental disorder associated with severe developmental delay and polymalformation syndrome[25]. The G375R mutation, located on the back side of the S6 pore-lining helix of the alpha subunit (Supplementary Fig. 1), replaces the hydrogen side chain of glycine with a large arginine side chain that might be expected to distort the subunit structure and gating[34] as well as add a positive charge that may alter conductance.

To assess the functional effects of a G375R mutation, we recorded currents from whole cells and excised macro patches of membrane after injecting a 1:1 mixture of G375R mutant and WT cRNA encoding for mutant and WT subunits into *Xenopus* oocytes, to mimic a *de novo* mutation heterozygous with respect to the WT allele. When compared to WT, the currents from the 1:1 injection were left shifted more than -120 mV. This large negative shift in activation indicated a pronounced gain-of-function (GOF) mutation at the cellular level, providing a possible explanation for the severity of the disease associated with the heterozygous G375R mutation.

To investigate the molecular basis underlying the cellular response, detailed single-channel recording[35] after the 1:1 injection suggested that five different types of functional BK channels were expressed: 3% were WT, 12% were homomeric mutant, and 85% were three different types of hybrid (heteromeric) channels. All channel types except WT showed a marked GOF in voltage activation and a smaller LOF in single channel conductance, with both becoming more pronounced as the number of mutant subunits per channel increased. Codominance was observed for the two homotetrameric channels, with homomeric WT channels active at the most positive voltage range of channel activity and homomeric mutant channels active at the most negative. Partial dominance was observed for the three types of hybrid channels, which were activated at voltages intermediate between those of WT and homomeric mutant channels. A model in which BK channels were randomly assembled from mutant and WT subunits, with each subunit

74   contributing increments of activation and conductance, could approximate the molecular
75   phenotype of the heterozygous G375R BK mutation. The possibility that a channelopathy patient
76   with a heterozygous BK channel mutation synthesizes five different types of BK channels in
77   their neurons and other cells, four with aberrant properties, presents a daunting challenge for
78   treatment.
79
80

Results
G375R subunits left shift BK channel activation. To examine what effect a *de novo* G375R
mutation of the pore forming subunit of BK channels would have on channel function when co-
expressed with WT subunits, we first compared macroscopic currents flowing through large
numbers of WT BK channels expressed following injection of WT cRNA into *Xenopus* oocytes,
to those currents expressed following injection of a 1:1 mixture of mutant and WT cRNA to
mimic a heterozygous mutation. Injection of a blank without cRNA served as a control. Voltages
were stepped from a holding potential of -80 mV to more negative and more positive voltages to
reveal the voltage dependent activation of the expressed currents (Fig. 2A; Supplementary Fig.
2). Voltage steps to +50 mV were required to appreciably activate WT BK currents generated
from the injection of WT cRNA. In contrast, following a 1:1 injection of G375R mutant and WT
cRNA, the expressed BK channels currents were hyperactive, being already active at -140 mV,
with depolarization further increasing the response (Fig. 2B, Supplementary Fig. 2). The aberrant
BK channels from the 1:1 injection that are responsible for these hyperactive currents could be
either homomeric mutant channels, hybrid channels, or both (Fig. 1).
The negative shift in activation for currents expressed from the 1:1 injection was so
pronounced that large numbers of aberrant BK channels would have been open at voltages near
the resting membrane potential of -60 mV. If aberrant currents expressed similarly in neurons,
the flux of K^+ through opened BK channels would act to drive the membrane potential towards
the equilibrium potential for K^+, opposing depolarization of the cell and facilitating
repolarization, which could interfere with normal neuronal function. Thus, at the level of whole
cell and macro patch recording, the heterozygous G375R mutation gave a pronounced gain-of-
function (GOF) phenotype because much less depolarization was required to activate BK
currents. Some of the disease phenotypes associated with this mutation might be due to such
GOF behavior. In contrast, Liang et al. reported a LOF for G375R because they observed no BK
currents for homomeric mutant channels[25]. Their conclusion of a LOF vs. our conclusion of a
GOF will be considered in detail in a later section.
To quantify the average negative voltage shift in activation for the BK currents following
injection of a 1:1 mixture of G375R and WT cRNA, when compared to WT currents following
injection of only WT cRNA, we recorded currents from macro patches of membrane which were
excised from oocytes after channel expression. This allowed the composition of the solution at
the inner membrane surface to be controlled, which was not the case for the whole cell

recordings in Fig. 2. For a solution at the intracellular membrane surface containing <0.01 µM

Ca^{2+} (~0 Ca^{2+}), WT BK currents did not activate appreciably until the membrane potential

exceeded 100 mV (black circles), whereas BK currents from the 1:1 injection of G375R mutant

and WT cRNAs started to activate at very negative voltages of -80 mV (Fig. 3B). The mean

voltage for half activation (V_h) was 182 ± 5.4 mV (n = 6) for WT BK currents and 61.5 ± 16 mV

(n = 12) for BK currents from the 1:1 injection, giving a mean left shift of -120 mV in voltage

activation for the currents following a 1:1 injection when compared to WT (p <0.0001). This left

shift in voltage activation was even greater in the presence of 300 µM Ca^{2+} (Supplementary Fig.

3). The 1:1 injection also decreased the voltage sensitivity of activation compared to WT, with a

slope of 38.5 ± 1.8 (n = 8) mV per e-fold change compared to 23.2 ± 1.36 mV (n = 6) for WT

currents (p < 0.0001).

The large negative shifts in voltage activation observed for whole cell recordings following a

1:1 injection of mutant and WT cRNA (Fig. 2; Supplementary Fig. 2) are thus also observed for

macro patch recordings under conditions in which the intracellular solution and Ca^{2+} are

controlled (Fig. 3; Supplementary Fig. 3).

Whole cell and macro patch recordings are useful to show the average response of many

hundreds to thousands of BK channels, but they provide limited information about the properties

of the underlying channels, unless all channels are identical, which is unlikely to be the case for a

heterozygous mutation, as shown in Fig. 1. To investigate the channels underlying the negative

shifts in BK currents in Figs. 2 and 3 for a 1:1 injection of mutant and WT cRNA to mimic a

mutation heterozygous with the WT allele, we first consider the possible channel types and

frequencies of expression that might be expected for a heterozygous mutation. We then record

from individual channels expressed following a 1:1 injection of mutant and WT cRNA to see if

the expectations are met.

BK channels of five different potential stoichiometries and six different subunit arrangements

could theoretically be expressed for a mutation heterozygous with WT. WT and mutant

subunits arising from a heterozygous mutation could potentially assemble into tetrameric BK

channels with five different stoichiometries and six different subunit arrangements, each with

potentially different properties (Fig. 1)[29-33]. If it is assumed that there is equal production of

mutant and WT subunits, that subunits assemble randomly to form tetrameric channels, and that

each assembled channel has the same probability of reaching the surface membrane, then the

channels expressed for a heterozygous mutation would consist of 6% WT channels, 25%

comprised of 1 mutant and 3 WT subunits, 38% comprised of 2 mutant and 2 WT subunits, 25%

comprised of 3 mutant and 1 WT subunit, and 6% homomeric mutant channels (Fig. 1). On this

basis, 94%, of the expressed channels could be pathogenic, comprised of 88% hybrid

(heteromeric) channels of mixed subunits and 6% homomeric mutant channels, with only 6%

WT channels (Fig. 1).

Assembled channels. The term assembled channels will be used to refer to those BK channels that are assembled and expressed following injection of a 1:1 mixture of G375R mutant and WT cRNA into *Xenopus* oocytes, to mimic a heterozygous mutation. Assembled channels could theoretically include five types of channels with different stoichiometries and six types of channels if subunit arrangement is considered (Fig. 1). An individual assembled channel could then be any one of the six channel types. It is the combined activity of the six potential types of assembled channels that would determine the BK currents for a heterozygous mutation.

Assembled channels display an abnormally wide range of V_h. To explore if multiple channel types contribute to the negative shift in V_h resulting from a G375R BK subunit mutation heterozygous with WT (Figs. 2, 3, Supplementary Figs. 2, 3), we used the unique ability of the patch clamp technique[31,35-37] to isolate and record from individual assembled channels expressed following a 1:1 injection of mutant and WT cRNA into *Xenopus* oocytes. For each of the 33 individual assembled channels studied, single-channel currents were recorded over a range of voltages (Fig. 4*A*) to obtain plots of open probability (Po) vs. voltage (Fig. 4*B*, red curves). The Po-V plots for the 33 individual assembled channels had a voltage for half activation (V_h) that ranged from -152 mV to +151 mV, for a span of 303 mV (Fig. 4*B*, red Po-V curves). This very wide range in V_h supports the idea that there are multiple types of assembled channels with different properties arising from different subunit composition (Fig. 1).

Assembled channels include WT, homomeric mutant, and hybrid channels. To identify the types of BK channels expressed following a 1:1 injection of mutant and WT cRNA, the Po-V curves of the 33 individual assembled channels in Fig. 4B were compared to Po-V curves obtained from individual WT and homomeric mutant channels. The WT and homomeric mutant channels were obtained by single channel recording after injection of only WT cRNA or only mutant cRNA, respectively. The WT channels had V_h values that spanned a narrow range from +140 to +176 mV (Fig. 4C, black Po-V curves), and the homomeric mutant channels had V_h values that ranged from -272 mV to -38 mV (Fig. 4C, blue Po-V curves). The homomeric mutant channels had a surprisingly wide range of V_h for channels of presumed identical composition of four mutant subunits, but all were in a far more negative voltage range than WT channels. The reason for such a wide voltage range in V_h for the homomeric mutant channels is not known, but perhaps channels with four mutant subunits can assume different conformations with markedly different activation properties, depending on mutant side-chain orientation (Supplementary Fig. 1).

To facilitate the identification of the types of assembled channels, the Po-V curves of the individual 33 assembled channels from Fig. 4B were overlaid on plots of the V_h ranges of the Po-V curves for known WT (gray shading) and homomeric mutant (blue shading) channel controls in Fig. 4D. One of the 33 assembled channels had a Po-V curve that overlapped with the known WT channel controls (Fig. 4D), suggesting that this assembled channel was WT with four WT

subunits. Four of the 33 assembled channels had Po-V curves that overlapped with the known homomeric mutant channel controls (Fig. 4D), suggesting that these four assembled channels were homomeric mutant, comprised of four mutant subunits. The remaining 28 assembled channels had Po-V curves that did not overlap with the known WT or homomeric mutant channel controls (Figs. 4D), but fell in between, suggesting that these 28 assembled channels were all hybrid channels comprised of a mix of mutant and WT subunits, because hybrid channels were the only types of channels remaining (Fig. 1). In this study, assembled channels with V_h values in the ranges of WT, hybrid, and homomeric mutant channels will be referred to as WT, hybrid, and homomeric mutant channels, respectively, with the understanding that these classifications are based on V_h values.

For the sampled group of 33 assembled channels, ~3% (1/33) were WT, ~85% (28/33) were hybrid, and ~12% (4/33) were homomeric mutant (Fig. 4B-D). Fig. 1 predicts ~6% WT, ~88% hybrid, and ~6% homomeric mutant channels. Thus, for the G375R mutation heterozygous with WT, both experimental and theoretical considerations suggest that most (85-88%) of the expressed assembled channels will be hybrid, with much smaller fractions of homomeric mutant and WT channels. All hybrid and homomeric mutant channels displayed negative shifts in activation compared to WT, with the greatest negative shifts for the homomeric mutant channels. Consequently, both theoretical and experimental considerations suggest that 94-97% of the BK channels arising from a G375R mutation heterozygous with WT would display aberrant negative shifts in activation, even though only 50% of the subunits synthesized in a cell would be mutant.

Three functional types of hybrid assembled channels. Assembled channels were found to be comprised of 85% hybrid channels, 3% WT, and 12% homomeric mutant (Fig. 4). The 85% hybrid channels themselves could consist of three types, with one, two, or three mutant subunits per channel replacing an equal number of WT subunits (Fig. 1). To explore this possibility, a histogram of the V_h values of the 33 assembled channels from Fig. 4B was plotted in Fig. 5A as red bars. Histograms of known WT channel controls (black bars) and homomeric mutant channel controls (blue bars) from Fig. 4C are also plotted. The subunit compositions of the homomeric mutant and WT channel controls are known and placed above these two types of channels in Fig. 5A. As expected from Fig. 4D, four of the assembled channels had V_h values that overlapped with those of homomeric mutant channels, suggesting that they were homomeric mutant channels; one assembled channel overlapped with WT, suggesting that it was a WT channel; and the remaining 28 assembled channels had V_h values falling in between those of homomeric mutant and WT channels, identifying them as hybrid channels (Fig. 1).

The V_h values of the 28 hybrid channels fell into three apparent clusters with mean values of about 1.4, 59 and 96 mV (Fig. 5A). Three clusters of V_h for hybrid channels would be expected from Fig. 1 if each mutant subunit replacing a WT subunit added an increment of negative shift in V_h, as suggested by Fig. 4 and in Fig. 6 in a later section. Accordingly, as a working hypothesis, the different subunit combinations for hybrid channels from Fig. 1 have been placed

above the three clusters of hybrid channels in Fig. 5A to obtain a stepwise negative shift in V_h for
each additional mutant subunit replacing a WT subunit. Because only three clusters of hybrid
channels were observed, instead of four, it has been assumed that the hybrid channels with two
mutant and two WT subunits, but of different subunit arrangement, contribute to the same
cluster. Fig. 5A then suggests five different types of functional assembled channels: homomeric
mutant, three types of hybrid channels, and WT channels, as indicated. The percentages of
expression of the five functional types of assembled channels were then calculated from Fig. 5A
and presented in Fig. 5B. Simulation of 1 million groups of 33 assembled channels, assuming
equal production of mutant and WT subunits followed by random assembly into tetrameric
channels, indicated that the experimentally observed percentages in Fig. 5B were not
significantly different from the predictions of Fig. 1 (Supplementary Fig. 4).
Whereas the observation of three apparent clusters of V_h values for the hybrid assembled
channels (Fig. 5) is consistent with theoretical predictions (Fig. 1), peaks can occur by chance
alone in histograms of binned data of limited sample size[38]. Consequently, additional
experiments would be needed to determine whether V_h values for hybrid channels can
consistently be resolved into three peaks, but an observation of such distinct peaks is not required
for support of Fig. 1, as variability in V_h among channels of the same type[39] might be sufficient
to obscure distinct peaks.
248
Linear Incremental model for the contributions of mutant and WT subunits to V_h. WT BK
channels comprised of four WT subunits had a mean V_h of about 160 mV, and homomeric
mutant BK channels comprised of four mutant subunits had a mean V_h of about -80 mV (Figs.
4A, C; Fig. 5A). As all four subunits contribute to the gating of BK channels[4,5,31], these
observations suggest that each WT subunit contributes an increment of positive shift to V_h and
each mutant subunit contributes an increment of negative shift. In support of this hypothesis, a
linear incremental model in which each WT subunit in a channel added +40.5 mV to V_h and each
mutant subunit added -16.7 mV, provided an approximate description of the observed V_h's for
the five types of assembled channels (Fig. 5C dashed line; model in figure legend). The net effect
of replacing a WT subunit with a mutant subunit was a -57.2 mV shift in V_h to account for the
loss of the positive shift contributed by the removed WT subunit and the addition of the negative
shift contributed by the added mutant subunit.
261
Dual action of G375R subunits on V_h and single channel conductance g. In addition to the
negative shift in activation induced by replacing WT subunits with G375R mutant subunits (Figs.
2-5), replacing WT subunits with mutant subunits also decreased single-channel conductance
(Fig. 6A, B). The mean single-channel conductance of WT channels was 312 ± 4 pS. This
decreased to 245 ± 6 pS for hybrid channels, and further decreased to 190 ± 14 pS for
homomeric mutant channels. These decreases were significant (Fig. 6 legend). Hence, single-
channel conductance decreased as mutant subunits replaced WT subunits. The decreased single-

channel conductance may arise from the larger volume and positive charge of the mutant
arginine side chains replacing the single hydrogen atom of the glycine side chains on the four S6
segments lining the conductance pathway of the BK channel (Supplementary Fig. 1). The added
volume of the sidechains could decrease the volume of the inner vestibule and the added positive
charge may act to repel K^+ from the inner cavity. Both actions can reduce single channel
conductance in BK channels[40-42].

To examine the relationship between single-channel conductance and V_h, the single-channel
conductance for each channel was plotted against V_h for the same channel in Fig. 6C for the
indicated channel types. A linear relationship was observed (Fig. 6C; R = 0.86, P < 0.0001).
When taken together, the data in Figs. 6B and 6C are consistent with the idea that replacing a
WT subunit with a mutant subunit adds both an increment of negative voltage shift to V_h and a
step decrease in single-channel conductance. Thus, at the single-channel level, the heterozygous
G375R mutation acts simultaneously as a GOF mutation to shift voltage activation to more
negative voltages (Figs. 4, 5), and as a loss-of-function (LOF) mutation to decrease single-
channel conductance (Fig. 6). At the whole cell and macro patch level, the increase in currents
from the GOF negative shift in activation would dominate the LOF reduction in single-channel
conductance, producing large left shifted currents (Figs. 2, 3; Supplementary Figs. 2 and 3). The
LOF in conductance would act to decrease the consequences of the negative shift in activation.

Linear Incremental model for the contributions of mutant and WT subunits to single-channel
conductance. A linear incremental model in which each WT subunit contributes 76.5 pS to
single channel conductance and each mutant subunit contributes 46.1 pS could approximate the
single channel conductance for the five types of assembled channels (Fig. 6D dashed line; model
in figure legend).

WT, hybrid, and G375R homomeric mutant channels are expressed in the HEK293 cell
expression system. Liang et al.[25] reported that no potassium currents were recorded from
excised macro patches following transfection of HEK293T cells with plasmids coding for
G375R mutant BK subunits. In contrast, we found that functional single homomeric mutant
channels were readily expressed following injection of *Xenopus* oocytes with cRNA coding for
mutant BK G375R subunits (Fig. 4A, C). To examine if these differences in channel expression
were due to differences in expression systems, we transfected HEK293 cells with WT cDNA to
generate WT channels, with mutant G375R cDNA to generate homomeric mutant channels, and
with a 1:1 mixture of WT and mutant cDNA to mimic the generation of channels that would
normally arise from a heterozygous mutation. We found that the transfected HEK293 cells
displayed robust whole-cell BK currents for each of the three different types of transfection (Fig.
7A). The G-V plots for transfection with WT cDNA or 1:1 transfection with WT and mutant
cDNA were very similar to those we obtained using the *Xenopus* oocyte expression system
(compare Fig. 7 with Fig. 3). We also found that whole cell G375R homomeric mutant currents

from HEK293 cells had V_h values within the broad range of V_h values of single-channel

recordings from patches of membrane excised from *Xenopus* oocytes (Fig. 4).

Hence, in both *Xenopus* oocytes and HEK293 expression systems we observed functional

G375R homomeric mutant currents that displayed a marked negative shift in activation

consistent with a GOF mutation. This contrasts with the observations of Liang et al.[25] who did

not observe potassium currents for G375R transfection of HEK293T cells. They reported G375R

as a LOF mutation. The reason for the difference in observations is not known, but we found that

the viability of *Xenopus* oocytes injected with G375R cRNA was reduced (see Methods).

Whole cell and macro-patch currents can obscure mechanism. The narrow error bars for the

macro currents recorded from whole oocytes and excised macro patches (Figs. 2 and 3;

Supplemental Figs. 2 and 3) following injection of WT cRNA or a 1:1 injection of mutant and

WT cRNA, indicated that the mean responses recorded from hundreds to thousands of channels

were repeatable. Narrow error bars for mean responses do not necessarily indicate that the

individual channels underlying the mean response have near identical properties, as this was not

the case for the macro currents following 1:1 injections of mutant and WT cRNA, where

individual assembled channels for 1:1 injections had a very wide range of V_h that spanned over

300 mV (Fig. 4B). In spite of this wide range for individual channels, the mean G-V data

following the 1:1 injections were reasonably well described by a single Boltzmann function (Fig.

3B, blue line through red diamonds), but were better described (red line) by assuming five

underlying channel types (Fig. 1) as described in the legend to Fig. 3.

Discussion

The heterozygous G375R BK channel variant has been associated with a devastating human

phenotype that includes malformation syndrome and severe neurological and developmental

disorders[25]. This variant has only appeared in the human population when heterozygous with

WT, perhaps because a homozygous G375R variant may not permit viability because of the

extreme GOF phenotype we observed for homomeric mutant channels. To gain insight into the

pathogenicity of this variant, currents from whole cells, macro patches, and single channels were

recorded from BK channels expressed following a 1:1 injection of G375R mutant and WT cRNA

to mimic a G375R mutation heterozygous with WT.

Recordings from whole cells and macro patches, both of which contain many hundreds to

thousands of channels, indicated that the voltage required for half activation, V_h, of the current

following a 1:1 injection, was left shifted to more negative potentials by about -120 mV

compared to WT currents (Figs. 2 and 3; Supplementary Figs. 2 and 3), indicating a pronounced

net GOF phenotype on channel activation. The aberrant BK channels underlying the negative

shifts in activation would lead to a much greater fraction of BK channels being open at negative

membrane potentials, including at potentials near the resting potential (Figs. 2, 3 and

Supplementary Figs. 2, 3), which would oppose cellular depolarization, altering cellular function.

To explore the underlying molecular basis for the aberrant current activation associated with

the heterozygous G375R variant, we used high resolution single channel recording to isolate and

characterize the BK channels, termed assembled channels, that are assembled and expressed

following injection of a 1:1 mixture of G375R mutant and WT cRNA. Theoretical considerations

based on equal production and random assembly of mutant and WT subunits suggest there could

be multiple types of assembled channels with different subunit combinations[29,30,32,33] (Fig. 1)

Consistent with this possibility, our analysis suggested that five different types of functional BK

channels were expressed: 3% were WT, 12% were homomeric mutant, and 85% were three

different types of hybrid channels of mixed subunits (Fig. 4, Fig. 5A, B). The frequencies of

expression of these five types of functional assembled channels were not significantly different

(Supplementary Fig. 4) from the theoretical predictions of Fig. 1 based on equal production and

random assembly of subunits. This suggests that the processes involved in subunit production,

assembly, and expression do not distinguish significantly between the mutant and WT alleles or

subunits, so that phenotypic differences arise at the level of individual channels.

The three types of hybrid channels comprising 85% of the expressed assembled channels,

had properties falling between those of mutant and WT channels, that varied with their apparent

subunit composition (Figs. 4-6). Ninety-seven percent of the assembled channels, all except for

the 3% WT, displayed both GOF negative shifts in activation and LOF decreases in single-

channel conductance (Figs. 4-6). Hence, most of the expressed channels for the heterozygous

G375R mutation displayed aberrant properties. The values of V_h and single channel conductance

for each of the five types of functional assembled channels could be predicted with a linear

incremental model in which each mutant and WT subunit in each of the five types of functional

assembled channels acted relatively independently to contribute increments of both V_h and

single-channel conductance to the molecular phenotype of the channel (Figs. 5C, 6D).

A potential mechanism to produce the five functional types of assembled channels for a

heterozygous G375R BK channel mutation, then, is equal production and random assembly of

mutant and WT subunits into channels of five different subunit compositions, where the V_h and

single-channel conductance of each channel type are determined by independent contributions

from each of the four subunits in a channel. Each WT subunit adds increments of 40.5 mV to V_h

and 76.5 pS to single channel conductance, and each mutant subunit adds increments of -16.7

mV to V_h and 46.1 pS to single channel conductance.

To our knowledge, this is the first time that the types of functional channels expressed for a

heterozygous mutation, their specific phenotypes, frequencies of expression, and contributions of

individual subunits have been revealed in such molecular detail (Figs. 4-6). With this molecular

information in hand, it is now possible to classify the heterozygous G375R mutation with regards

to inheritance of genetic disease, and also the individual molecular phenotypes of the five types

of expressed functional channels.

For classification with regard to genetic disease, we suggest that the heterozygous G375R

mutation be labeled as an assembly-mediated dominant-positive mutation, following the

terminology of Backwell and Marsh[33]. Assembly-mediated to indicate that mutant and WT
subunits assemble into multiple types of functional channels for the tetrameric BK channel,
dominant because the inclusion of only a single copy of the mutant *KCNMA1* gene leads to
disease[25], and positive to indicate an aberrant GOF in the activation of BK currents (Figs. 2, 3,
7). The net result of this assembly-mediated dominant-positive mutation in the tetrameric BK
channels is that a disproportionate fraction (94-97%) of the expressed channels display aberrant
activation due to the random assembly of mutant and WT subunits into functional channels
(Figs. 1, 4, 5). A dominant-positive GOF phenotype has been described previously for the
heterozygous G88R mutation of the TASK-4 K^+ channel in the heart, based on a study using
whole cell currents[43].

A dominant-positive effect is an analogy[33] to the well-known dominant-negative effect
observed for some types of protein complexes and channelopathies[33,44-49] Both can cause
disease: dominant positive from a GOF and dominant-negative from complete or partial LOF,
and both have a disproportionate fraction of the channels affected.

Whereas an assembly-mediated dominant-positive classification of the heterozygous G375R
mutation is useful to suggest a basis of genetic disease at the level of whole cell and macro patch
currents through BK channels, it describes only part of the molecular phenotype. Consequently,
to characterize the molecular phenotype we suggest that random assembly of mutant and WT
subunits for the heterozygous G375R mutation produces five functional types of assembled
channels: WT, mutant, and three types of hybrid channels of mixed subunit composition. All
channel types except WT simultaneously display a GOF in activation and a partial LOF in
single-channel conductance.

The genetic dominance of the various channel types expressed for a heterozygous mutation is
mixed and complex. Codominance was observed for the WT and homomeric mutant channels
and partial dominance was observed for the three types of hybrid channels. Support for
codominance is that the mutant and WT channels expressed when mimicking a heterozygous
mutation had the same molecular phenotypes as homomeric mutant and WT channels expressed
after injection of only mutant or WT cRNA, respectively (Figs. 4 and 5). Support for partial
dominance is that the three types of hybrid channels had their own values of V_h and single
channel conductance that were distinct from each other and fell in the range between those of the
homomeric mutant and WT channels (Figs. 5C, 6D). The levels of partial dominance in the
linear incremental model (Figs. 5C and 6D) were determined by the numbers of mutant and WT
subunits per channel acting relatively independently of one another, rather than by mutant
subunits altering the function of WT subunits, as in some classical descriptions of dominant-
negative. It is remarkable that a single base pair substitution in one allele of a pair of alleles that
encode for the alpha subunit of BK channels would result in WT channels plus four aberrant
channel types, each with different molecular phenotypes that simultaneously display dominant
positive and dominant negative effects, with the five types of functional channels displaying
either co-dominance or one of three levels of partial dominance.

How do our observations of a relatively independent interactions of the G375R mutant and
WT pore-forming subunits compare to interactions of these subunits with regulatory subunits?
Wang et al.[50] found that single BK channels comprised of four subunits could be associated with
from 0 to 4 regulatory β2-subunits per channel, with each β2-subunit giving incremental changes
in V_h. Hence both mutated and WT pore forming subunits[36] (Fig. 5) and the non-pore forming
regulatory β2-subunits[50] can give incremental changes in gating, but such incremental changes
per subunit are not necessarily universal, so each type of subunit will need to be checked. A
single BK channel comprised of four subunits can also include up to four regulatory γ1-subunits,
but a single γ1-subunit per channel is sufficient to induce the full gating shift induced by γ1-
subunits[51,52].
Devising therapies for the *de novo* G375R heterozygous mutation will be challenging.
Theoretical and experimental observations (Figs. 1, 4, 5) suggest that most (94 to 97%) of the
channels for a mutation heterozygous with WT would be pathogenic, with only 3 to 6% of the
channels WT. The pathogenic channels would consist of multiple channel types, each with large
differences in activation properties and smaller differences in conductance (Figs. 4 - 6).
Therapies to block or inactivate the pathogenic channels would ideally silence the multiple types
of pathogenic channels while leaving any WT channels intact. Even if such selective blockers
could be devised, it is unlikely that the remaining 3 to 6% of WT channels would be sufficient to
restore normal cellular function. Effective therapies will likely require replacing or silencing the
mutant alleles, or preventing mutant subunits from assembling with themselves and WT subunits
if/when such techniques become practical in humans.
446

Methods

Constructs. Experiments were performed using the human large conductance calcium-activated
potassium (BK) channel (hSlo1) KCNMA1 transcript: GenBank: U23767.1[3] for WT, and a
G375R mutation (VFFIL**GG**GLAMF to VFF IL**R**GLAMF) was constructed to match the G375R
de novo variant described by Liang et al.[25]. G375R is at the same position as G310 in Tao and
MacKinnon[15], which they called the gating hinge residue. The numbering differs because Tao
and MacKinnon[15] used an alternative transcript missing the first 65 amino acids compared to
U23767.1. Both our hSlo1 WT cRNA plasmid and WT mammalian transfection plasmid contain
the same hSlo1 insert and are described in McCobb et. al.[3]. Overlap extension PCR cloning was
used to generate the G375R mutation, which was verified by sequencing. The new construct was
linearized downstream of the end of coding and transcribed with T3 using Invitrogen's T3
mMessage mMachine kit to make cRNA for injection into *Xenopus* oocytes. To mimic the
effects of a heterozygous mutation, G375R mutant and WT cRNA were mixed in a 1:1 ratio by
weight before injection into oocytes. For expression in HEK293 cells, two mammalian
transfection plasmids were created. DNA fragments containing the complete channel cDNAs of
either the WT or G375R mutant and two unique restriction sites flanking it were amplified by
PCR and ligated into mammalian transfection plasmids, which were verified by sequencing.

Expression of BK channels in oocytes for whole cell recording. Defolliculated *Xenopus* oocytes were injected with 0.5-150 ng of cRNA using a Nanoject II (Drummond Scientific) and incubated at 18 °C for 2-5 days before recording. Incubation was in ND96 complete medium consisting of (mM) 96 NaCl, 2 KCl, 1.8 CaCl$_2$, 5 MgCl$_2$, and 5 HEPES adjusted to pH 7.5, supplemented with 2.5 mM sodium pyruvate and 100 µg/ml each penicillin and streptomycin. The two-microelectrode whole cell voltage-clamp recordings from oocytes were obtained in ND96 medium with 1 mM added 4,4′-diisothiocyanatostilbene-2,2′-disulphonic acid disodium salt hydrate (DIDS) to block the endogenous chloride conductance. Currents were obtained with an Oocyte Clamp OC-725C amplifier (Warner Instrument Corp.) Recordings were low-pass filtered at 1 kHz and digitized at 10 kHz. Electrodes were made with borosilicate glass capillaries (World Precision Instruments) pulled with a Sutter Instrument Co. P-87 pipette puller and filled with 3 M KCl.

Macro patch and single-channel recordings from BK channels expressed in *Xenopus* oocytes. Oocytes were injected with 0.1-18 ng of cRNA and incubated at 18°C for 2-5 days in Barth's Solution (mM): 88 NaCl, 1 KCl, 2.4 NaHC0$_3$, 0.33 Ca(NO$_3$)$_2$, 0.41 CaCl$_2$, 0.82 MgSO$_4$, 15 mM HEPES, pH 7.6, plus 12 µM tetracycline. Macro and single channel currents were recorded from inside-out patches of membrane[35] excised from oocytes at room temperature (21°C to 24°C). pClamp 9.0 software (Molecular Devices) was used to drive an Axopatch 200B amplifier to collect the currents. For macro patch current recordings, borosilicate pipettes with 0.5–2 MΩ resistance were used. The macro currents were filtered at 10 KHz and sampled at 100 kHz. A minus P/4 protocol was used to remove capacitive transients and leak currents. For single-channel recordings, borosilicate pipettes with 8-12 MΩ resistance were used. The single-channel currents were filtered at 5 KHz and sampled at 200 KHz. The pipette (external) solution contained (mM): 160 KCl, 2 MgCl$_2$, 5 TES buffer, pH 7.0. The internal membrane surface of the excised patches was perfused by two different solutions. The designated 0 Ca^{2+} solution had a free Ca^{2+} <0.01 µM and contained (mM): 160 KCl, 1 EGTA, 1 HEDTA, and 10 HEPES, pH 7.0. The solution with 300 µM internal Ca^{2+} contained (mM): 160 KCl, 0.3 CaCl$_2$, 10 HEPES, pH 7.0. Procedures to obtain oocytes from *Xenopus* laevis were approved by the University of Miami Animal Care and Use Committee. Macro patch and single channel recordings were analyzed with Clampfit 10.7 software (Molecular Devices) and SigmaPlot 12.

For injection of only G375R cRNA into *Xenopus* oocytes we found that it was difficult to get giga-ohm seals of sufficient quality for macro patch recordings using the lower resistance electrodes required for such recordings. Hence, macro patch currents are not presented for injection of only G375R cRNA into oocytes. However, it was still possible to obtain high quality giga-ohm seals for single-channel recording using the higher resistance pipettes required for such recordings (Fig. 4). We also found that the viability of the oocytes was greatly reduced following injection of only G375R cRNA, with the oocytes often starting to die by the second or third day

after injection, rather than after a week or more, perhaps because a large fraction of the G375R
homomeric mutant channels would be expected to be open at resting membrane potentials (Figs.
4, 7). These viability problems were not observed for injection of 1:1 mixtures of G375R mutant
and WT cRNA, which gave less pronounced negative shifts in activation or for injections of only
WT cRNA (Figs. 3, 4, 7),
The open probability (Po) of each single channel analyzed in detail was determined for a
range of voltages with Clampfit 10.7 using 50% threshold analysis to measure open and closed
interval durations. Po at each voltage was calculated by dividing the total open time by the sum
of the open and closed times. The durations of recordings to estimate Po ranged from ~ 5 s to ~3
min with the time increasing as the Po decreased. For macro patch current recordings, relative
conductance was determined from macroscopic tail current amplitudes using the voltage
protocols indicated in the figure legends. G/G_{max} vs. V (G-V) plots for macro current recordings
and Po vs. V (Po-V) plots for single channels were fitted with a Boltzmann function to estimate
V_h, the voltage required for half maximal activation, and b, a measure of voltage sensitivity,
using
518
519          $$G/G_{max} = 1/(1 + \exp((V_h - V)/b)) \qquad\qquad\qquad (1)$$
520
where G/G_{max} is the ratio of conductance to maximum conductance and b is the slope factor
which gives a measure of voltage sensitivity, where b indicates the change in millivolts required
to increase G/G_{max} (or Po for single channel recording) e-fold at very low G/G_{max} (or Po). Note
that an increase in b indicates a decrease in slope and voltage sensitivity.
Single-channel conductance was determined at 100 mV by setting horizontal cursor lines by
eye to the open and closed current levels of single-channel recordings.
527
528 **HEK293 cell culture.** As described previously[53], Human Embryonic Kidney HEK293 cells
529 (ATCC CRL-1573) (ATCC, Manassas, VA) were cultured in Gibco Dulbecco's modified Eagle's
530 medium (DMEM) with 10% fetal bovine serum, 100 units ml^{-1} penicillin and 100 μg ml^{-1}
531 streptomycin (ThermoFisher Scientific, Waltham, MA) and incubated at 37°C with 5% CO_2.
532 These cells were grown and passaged twice a week in T25 flasks (MidSci, St. Louis, MO).
533
Expression of BK channels in HEK293 cells. HEK293 cells were plated at a density of ~400,000
cells per 40 mm petri dish (TPP catalog #93040) a day before transfection. For transfection,
plasmid DNA containing cDNA encoding for BK WT and/or BK G375R (mutant) subunits was
added to cell layers that were 70-90% confluent using Lipofectamine 2000 transfection reagent
(catalog #11668-027, ThermoFisher Scientific, Waltham, MA) following manufacturer's
instructions. 2 μg cDNA of WT, mutant, or a mix of WT and mutant (1:1 ratio by weight) was
transfected per 40 mm dish. As a marker for transfection, cells were co-transfected with
pmaxGFP a CMV plasmid expressing green fluorescent protein (Amaxa Biosystems), at 0.2 μg

per dish. After transfection, the cells were incubated at 37°C with 5% CO_2 for 2-4 days until
recording. The recordings were then performed at room temperature (22°C). Fluorescent cells
were used to study the function of the ion channels that the HEK293 cells were transfected with;
non-fluorescing cells were used as a non-transfected controls[54].
546
Whole cell recording from HEK293 cells. Whole cell recordings were obtained from HEK293
cells using an Axopatch 200B amplifier (Molecular Devices, San Jose, CA). Recordings were
filtered at 5 kHz with the amplifier internal filter and digitized at 50 kHz using a Digidata 1550B
digitizer (Molecular Devices, San Jose, CA). The recording pipette was filled with (in mM): 140
KMES, 5 EGTA, 10 HEPES, pH 7.4 with KOH. Bath solutions contained (in mM): 135 NaMES,
5 KMES, 1 $MgCl_2$, 10 HEPES, pH 7.4 with NaOH.

Calculating the percentages of the five different stoichiometries of BK channels that can be
expressed when mimicking a heterozygous mutation. As shown in Fig. 1, a mutation
heterozygous with WT can potentially produce five different types of channels, defined by their
stoichiometry. The binomial equation was used to calculate the expected percentages of different
combinations of mutant and WT subunits for tetrameric BK channels[29,30]. For the calculations in
Fig. 1 this approach assumes that WT and mutant alleles produce equal numbers of mutant and
WT subunits that assemble randomly into tetrameric channels of different stoichiometries,
labelled assembled channels in Fig. 1, and that each assembled channel has the same probability
of being expressed, independent of subunit composition, such that
562
563   $$F_i = (4!/((4-i)!\, i!))\, F_{WT}^{(4-i)}\, F_{MUT}^{i} \quad\quad\quad\quad (2)$$
564
565   where F_i is the fraction of channel types with i mutant subunits, F_{WT} is the fraction of WT cRNA
566   injected, and F_{MUT} is the fraction of mutant cRNA injected, where $F_{WT} + F_{MUT} = 1$. For BK
567   channels, i ranges from 0 to 4, giving five channel types based on subunit composition. Setting
568   both F_{WT} and F_{MUT} to 0.5 gives the fractions of the various channel types for a heterozygous
569   mutation.
570
Simulating the discrete probability distributions of the number of BK channels of each
stoichiometry expected for equal production and random assembly of mutant and WT
subunits. The simulated discrete probability distributions were used to determine if the
experimentally observed number of channels of each stoichiometry differed significantly from
the number expected for random assembly of mutant and WT subunits. Discrete probability
distributions for the five stoichiometries of assembled channels in Fig. 1 for groups of 33
channels were simulated as follows. The first step was to generate a group of 33 channels by
assuming random assembly of mutant and WT subunits drawn from equal numbers of mutant
and WT subunits (with replacement) for each channel in the group. To assemble each channel,

four random numbers between 0 and 1 were drawn. Each random number < 0.5 indicated a WT

subunit and each random number ≥ 0.5 indicated a mutant subunit. The number of channels of

each of the five stoichiometries for each group of 33 channels were then binned into separate

histograms for each stoichiometry. This process was then repeated for 10^6 groups of 33 channels.

Each histogram of the numbers of channels of each type for the 10^6 groups was then normalized

to a discrete probability distribution with a summed probability of 1.0, to indicate the probability

of observing the indicated number of channels of that stoichiometry in a group of 33 channels.

These predicted discrete probability distributions are presented in Supplementary Fig. 4 for

comparison to the experimentally observed number indicated by red histogram bars in the same

figure.

Software. The computer program for simulating the discrete probability distributions for the

different types of channels assuming random assembly of mutant and WT subunits,

Binomdouble4-33.bas, is written in QB64 and is available from the corresponding author upon

request. The increments of single channel conductance and V_h added by each G375R mutant and

WT subunit for Figs. 5C and 6D were optimized by minimizing the least squared differences

between the experimental values and those calculated with the equations in the legends of Figs. 5

and 6 by running Solver in Excel.

Statistics. Error bars in the figures and error estimates in the text are SEM. Significance was

determined with the two-tailed *t*-test unless otherwise indicated.

Data Availability. The data for this study are contained in the figures and text of the manuscript

and the Supplementary information.

References

1. Barrett, J.N., Magleby, K.L. & Pallotta, B.S. Properties of single calcium-activated potassium channels in cultured rat muscle. *J. Physiol* **331**, 211-230 (1982).
2. Latorre, R., Vergara, C. & Hidalgo, C. Reconstitution in planar lipid bilayers of a Ca2+-dependent K+ channel from transverse tubule membranes isolated from rabbit skeletal muscle. *Proc. Natl. Acad. Sci. U. S. A* **79**, 805-809 (1982).
3. McCobb, D.P. *et al.* A human calcium-activated potassium channel gene expressed in vascular smooth muscle. *Am. J. Physiol* **269**, H767-H777 (1995).
4. Rothberg, B.S. & Magleby, K.L. Voltage and Ca2+ activation of single large-conductance Ca2+-activated K+ channels described by a two-tiered allosteric gating mechanism. *J. Gen. Physiol* **116**, 75-99 (2000).
5. Horrigan, F.T. & Aldrich, R.W. Coupling between voltage sensor activation, Ca2+ binding and channel opening in large conductance (BK) potassium channels. *J. Gen. Physiol* **120**, 267-305 (2002).
6. Xia, X.M., Zeng, X. & Lingle, C.J. Multiple regulatory sites in large-conductance calcium-activated potassium channels. *Nature* **418**, 880-884 (2002).
7. Salkoff, L., Butler, A., Ferreira, G., Santi, C. & Wei, A. High-conductance potassium channels of the SLO family. *Nat. Rev. Neurosci* **7**, 921-931 (2006).
8. Horrigan, F.T. Perspectives on: conformational coupling in ion channels: conformational coupling in BK potassium channels. *J. Gen. Physiol* **140**, 625-634 (2012).
9. Geng, Y. & Magleby, K.L. Single-channel kinetics of BK (Slo1) channels. *Front Physiol* **5**, 532 (2014).
10. Pantazis, A. & Olcese, R. Biophysics of BK Channel Gating. *Int. Rev. Neurobiol* **128**, 1-49 (2016).
11. Latorre, R. *et al.* Molecular Determinants of BK Channel Functional Diversity and Functioning. *Physiol Rev* **97**, 39-87 (2017).
12. Zhou, Y., Yang, H., Cui, J. & Lingle, C.J. Threading the biophysics of mammalian Slo1 channels onto structures of an invertebrate Slo1 channel. *J. Gen. Physiol* **149**, 985-1007 (2017).
13. Tao, X., Hite, R.K. & MacKinnon, R. Cryo-EM structure of the open high-conductance Ca(2+)-activated K(+) channel. *Nature* **541**, 46-51 (2017).
14. Hite, R.K., Tao, X. & MacKinnon, R. Structural basis for gating the high-conductance Ca(2+)-activated K(+) channel. *Nature* **541**, 52-57 (2017).
15. Tao, X. & MacKinnon, R. Molecular structures of the human Slo1 K(+) channel in complex with beta4. *Elife* **8**(2019).
16. Geng, Y. *et al.* Coupling of Ca(2+) and voltage activation in BK channels through the alphaB helix/voltage sensor interface. *Proc Natl Acad Sci U S A* **117**, 14512-14521 (2020).
17. Brenner, R. *et al.* Vasoregulation by the beta1 subunit of the calcium-activated potassium channel. *Nature* **407**, 870-876 (2000).
18. Robitaille, R., Garcia, M.L., Kaczorowski, G.J. & Charlton, M.P. Functional colocalization of calcium and calcium-gated potassium channels in control of transmitter release. *Neuron* **11**, 645-655 (1993).
19. Harvey, J.R.M., Plante, A.E. & Meredith, A.L. Ion Channels Controlling Circadian Rhythms in Suprachiasmatic Nucleus Excitability. *Physiol Rev* **100**, 1415-1454 (2020).
20. Gu, N., Vervaeke, K. & Storm, J.F. BK potassium channels facilitate high-frequency firing and cause early spike frequency adaptation in rat CA1 hippocampal pyramidal cells. *J Physiol* **580**, 859-882 (2007).
21. Park, S.M. *et al.* BK channel properties correlate with neurobehavioral severity in three KCNMA1-linked channelopathy mouse models. *Elife* **11**(2022).

22. Montgomery, J.R. & Meredith, A.L. Genetic activation of BK currents in vivo generates
bidirectional effects on neuronal excitability. *Proc Natl Acad Sci U S A* **109**, 18997-9002 (2012).
23. Wang, B., Rothberg, B.S. & Brenner, R. Mechanism of increased BK channel activation from a
channel mutation that causes epilepsy. *J. Gen. Physiol* **133**, 283-294 (2009).
24. Yang, J. *et al.* An epilepsy/dyskinesia-associated mutation enhances BK channel activation by
potentiating Ca2+ sensing. *Neuron* **66**, 871-883 (2010).
25. Liang, L. *et al.* De novo loss-of-function KCNMA1 variants are associated with a new multiple
malformation syndrome and a broad spectrum of developmental and neurological phenotypes.
Hum Mol Genet **28**, 2937-2951 (2019).
26. Bailey, C.S., Moldenhauer, H.J., Park, S.M., Keros, S. & Meredith, A.L. KCNMA1-linked
channelopathy. *J Gen Physiol* **151**, 1173-1189 (2019).
27. Miller, J.P., Moldenhauer, H.J., Keros, S. & Meredith, A.L. An emerging spectrum of variants and
clinical features in KCNMA1-linked channelopathy. *Channels (Austin)* **15**, 447-464 (2021).
28. Cui, J. BK Channel Gating Mechanisms: Progresses Toward a Better Understanding of Variants
Linked Neurological Diseases. *Front Physiol* **12**, 762175 (2021).
29. MacKinnon, R. Determination of the subunit stoichiometry of a voltage-activated potassium
channel. *Nature* **350**, 232-235 (1991).
30. Blaine, J.T. & Ribera, A.B. Heteromultimeric potassium channels formed by members of the Kv2
subfamily. *J Neurosci* **18**, 9585-93 (1998).
31. Niu, X. & Magleby, K.L. Stepwise contribution of each subunit to the cooperative activation of BK
channels by Ca2+. *Proc Natl Acad Sci U S A* **99**, 11441-6 (2002).
32. Bergendahl, L.T. *et al.* The role of protein complexes in human genetic disease. *Protein Sci* **28**,
1400-1411 (2019).
33. Backwell, L. & Marsh, J.A. Diverse Molecular Mechanisms Underlying Pathogenic Protein
Mutations: Beyond the Loss-of-Function Paradigm. *Annu Rev Genomics Hum Genet* **23**, 475-498
(2022).
34. Chen, X., Yan, J. & Aldrich, R.W. BK channel opening involves side-chain reorientation of multiple
deep-pore residues. *Proc. Natl. Acad. Sci. U. S. A* **111**, E79-E88 (2014).
35. Hamill, O.P., Marty, A., Neher, E., Sakmann, B. & Sigworth, F.J. Improved patch-clamp techniques
for high-resolution current recording from cells and cell-free membrane patches. *Pflugers Arch*
391, 85-100 (1981).
36. Niu, X. & Magleby, K.L. Stepwise contribution of each subunit to the cooperative activation of BK
channels by Ca2+. *Proc. Natl. Acad. Sci. U. S. A* **99**, 11441-11446 (2002).
37. Gonzalez-Perez, V., Xia, X.M. & Lingle, C.J. Two classes of regulatory subunits coassemble in the
same BK channel and independently regulate gating. *Nat. Commun* **6**, 8341 (2015).
38. Miller, D.C., Weinstock, M.M. & Magleby, K.L. Is the quantum of transmitter release composed of
subunits? *Nature* **274**, 388-390 (1978).
39. McManus, O.B. & Magleby, K.L. Accounting for the Ca(2+)-dependent kinetics of single large-
conductance Ca(2+)-activated K+ channels in rat skeletal muscle. *J. Physiol* **443**, 739-777 (1991).
40. Brelidze, T.I., Niu, X. & Magleby, K.L. A ring of eight conserved negatively charged amino acids
doubles the conductance of BK channels and prevents inward rectification. *Proc. Natl. Acad. Sci. U.*
S. A **100**, 9017-9022 (2003).
41. Nimigean, C.M., Chappie, J.S. & Miller, C. Electrostatic tuning of ion conductance in potassium
channels. *Biochemistry* **42**, 9263-9268 (2003).
42. Geng, Y., Niu, X. & Magleby, K.L. Low resistance, large dimension entrance to the inner cavity of BK
channels determined by changing side-chain volume. *J. Gen. Physiol* **137**, 533-548 (2011).
43. Friedrich, C. *et al.* Gain-of-function mutation in TASK-4 channels and severe cardiac conduction
disorder. *EMBO Mol Med* **6**, 937-51 (2014).

44. Du, X. *et al.* Loss-of-function BK channel mutation causes impaired mitochondria and progressive
cerebellar ataxia. *Proc Natl Acad Sci U S A* **117**, 6023-6034 (2020).
45. Fink, M. *et al.* Dominant negative chimeras provide evidence for homo and heteromultimeric
assembly of inward rectifier K+ channel proteins via their N-terminal end. *FEBS Lett* **378**, 64-68
(1996).
46. Ribera, A.B., Pacioretty, L.M. & Taylor, R.S. Probing molecular identity of native single potassium
channels by overexpression of dominant negative subunits. *Neuropharmacology* **35**, 1007-1016
(1996).
47. Silberberg, S.D., Chang, T.H. & Swartz, K.J. Secondary structure and gating rearrangements of
transmembrane segments in rat P2X4 receptor channels. *J Gen Physiol* **125**, 347-59 (2005).
48. Hichri, E., Selimi, Z. & Kucera, J.P. Modeling the Interactions Between Sodium Channels Provides
Insight Into the Negative Dominance of Certain Channel Mutations. *Front Physiol* **11**, 589386
(2020).
49. Reed, A.P., Bucci, G., Abd-Wahab, F. & Tucker, S.J. Dominant-Negative Effect of a Missense Variant
in the TASK-2 (KCNK5) K+ Channel Associated with Balkan Endemic Nephropathy. *PLoS One* **11**,
e0156456 (2016).
50. Wang, Y.W., Ding, J.P., Xia, X.M. & Lingle, C.J. Consequences of the stoichiometry of Slo1 alpha and
auxiliary beta subunits on functional properties of large-conductance Ca2+-activated K+ channels.
J. Neurosci **22**, 1550-1561 (2002).
51. Gonzalez-Perez, V., Ben, J.M., Xia, X.M. & Lingle, C.J. Regulatory gamma1 subunits defy symmetry
in functional modulation of BK channels. *Proc. Natl. Acad. Sci. U. S. A* **115**, 9923-9928 (2018).
52. Gonzalez-Perez, V., Xia, X.M. & Lingle, C.J. Functional regulation of BK potassium channels by
gamma1 auxiliary subunits. *Proc. Natl. Acad. Sci. U. S. A* **111**, 4868-4873 (2014).
53. Santi, C.M. *et al.* Opposite regulation of Slick and Slack K+ channels by neuromodulators. *J.
Neurosci* **26**, 5059-5068 (2006).
54. Han, Y., Li, P. & Slaughter, M.M. Selective antagonism of rat inhibitory glycine receptor subunits. *J
Physiol* **554**, 649-58 (2004).
55. Ding, J.P., Li, Z.W. & Lingle, C.J. Inactivating BK channels in rat chromaffin cells may arise from
heteromultimeric assembly of distinct inactivation-competent and noninactivating subunits.
Biophys. J **74**, 268-289 (1998).

Acknowledgements This work was supported in part by NIH grant R01-GM114694 to L.S. and
K.L.M. We thank Julia Bulova for technical help during the initial phase of this project.

Author contributions

Y.G., P.L., L.S., and K.L.M. designed research; Y.G., P.L., A.B., B.W., L.S., and K.L.M

performed research; L.S. and K.L.M. wrote the manuscript with input from the other authors.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary information is included.

Correspondence and requests for materials can be addressed to Lawrence Salkoff

(salkoffl@wustl.edu) or Karl L. Magleby (kmagleby@miami.edu).

Figure legends

Fig. 1 Protein subunits encoded by WT and mutant alleles for a heterozygous mutation could potentially assemble into five different stoichiometries of tetrameric BK channels and six different subunit arrangements[29,30,32,36,55]**.** In this diagram, WT and mutant alleles (left) encode for WT and mutant protein subunits (middle), which then assemble into tetrameric assembled channels with five different subunit stoichiometries and six different subunit arrangements (right). The listed theoretical frequencies of expression of the five different stoichiometries of assembled channels were calculated with Eqn. 2 in the Methods, which assumes that equal numbers of mutant and WT subunits randomly assemble into tetrameric channels, each with equal probability of reaching the surface membrane. With random assembly, ~6% of the assembled channels are WT comprised of four WT subunits per channel, ~88% are hybrid (mixed subunit) channels with from 1 to 3 mutant subunits per channel, and ~6% are homomeric mutant channels comprised of four mutant subunits per channel. Note that ~94% of the assembled channels can be pathogenic, as they contain from 1 to 4 mutant subunits per channel. The listed percentages of ~6%, ~38%, and ~94% here and throughout the paper have been rounded from the calculated values of 6.25%, 37.5%, and 93.75%. The 37.5% reflects the sum of 25% for hybrid channels with two adjacent mutant subunits and 12.5% for hybrid channels with two diagonal mutant subunits[32]. For experimentation, assembled channels are those channels expressed following injection of a 1:1 mixture of mutant and WT cRNA into *Xenopus* oocytes, to mimic a mutation in which the mutant allele is heterozygous with the WT allele.

Fig. 2 BK channels expressed in oocytes following injection of a 1:1 mixture of G375R mutant and WT cRNA activate at greatly left shifted negative potentials compared to BK channels expressed following injection of only WT cRNA. (**A**) Whole cell currents recorded from oocytes with the two-electrode voltage clamp for the indicated injections of cRNA. The whole cell currents were generated by holding the potential at -80 mV and then jumping to voltages ranging from -140 mV to +60 mV in 10 mV increments (1:1) or to +100 mV (WT and Blank). (**B**) Plots of relative conductance versus the voltage of the steps following: injection of: a 1:1 mixture of mutant and WT cRNA (red diamonds); injection of WT cRNA alone (black circles); or injection of carrier only (purple triangles). The high conductance at negative potentials would act to drive the membrane potential to about -80 mV, the equilibrium potential of K^+. Channel activation is left shifted in the whole cell recordings in this figure compared to the ~0 Ca^{2+} macro patch recordings in Fig. 3, because the resting free Ca^{2+} in the oocytes of a few

micromolar left shifts BK channel activation. The relative cord conductance was calculated from

the plotted steady-state currents in Supplementary Fig. 2 using the step potential minus the

reversal potential for the voltage driving force. Mean ± SEM, n = 4 in each case.

Fig. 3 Quantifying the mean negative shift in V_h following injection of a 1:1 mixture of G375R

mutant and WT cRNA when compared to injection of only WT cRNA. Data are for essentially 0

Ca^{2+} at the intracellular side of the membrane. (**A**) Currents recorded from inside-out macro

patches of membrane excised from oocytes. For WT channels following injection of only WT

cRNA, holding potential was -80 mV and voltage pulses were from -80 mV to 240 mV with 20

mV steps followed by a step back to -80 mV to elicit tail currents for G/Gmax measurements.

For channels expressed following injection of a 1:1 mixture of G375R mutant and WT cRNA,

holding potential was -160 mV and voltage pulses were from -120 to 240 mV with 20 mV steps

followed by a step back to -120 mV to elicit tail current for measurement. The currents are the

mean response from the many hundreds to thousands of BK channels in each excised macro

patch of membrane. (**B**) G/G_{max} vs. voltage plots following the injection of WT cRNA alone

(black circles, n = 6), or 1:1 injection of mutant and WT cRNA (red diamonds, n = 12). The black

line through the WT data is a single Boltzmann function with V_h = 181.3 mV and slope factor b =

24.04 mV/e-fold change in Po at low Po. The blue line through the 1:1 data (red diamonds) is a

single Boltzmann function with V_h = 59.04 mV and b = 38.06 mV/e-fold change. The red line

through the 1:1 data is the sum of five Boltzmann functions with: 1) V_h = -75.7 mV, b =21.6

mV/e-fold change, fractional area = 0.0547; 2) 1.5, 21.0, 0.236; 3) 54.6, 13.8, 0.316; 4) 96.6,

14.4, 0.244; 5) 155, 14.0, 0.139. These parameters for the five summed Boltzmann functions

that describe the 1:1 data are generally consistent with the V_h values for the apparent clusters

of assembled channels (Fig. 5A) and theoretical percentages in Fig. 1.

Fig. 4 Identifying the types of assembled BK channels expressed following a 1:1 injection of

G375R mutant and WT cRNA. (**A**) Representative single channel recordings from three different

single BK channel molecules: WT channel (top row) following injection of WT cRNA; an

assembled channel following 1:1 injection of G375R mutant and WT cRNA, (second row); and a

G375R homomeric mutant channel following injection of G375R mutant cRNA (third row).

Recordings are shown at three different voltages for each channel type. Depolarization

activates each channel type with a markedly different V_h of 140 mV for the WT channel, 6 mV

for the assembled channel, and -100 mV for the G375R homomeric mutant channel. Arrows

indicate closed channel current levels. (**B**) Plots of Po vs. V for 33 individual assembled channels

following a 1:1 injection of mutant and WT cRNA (red diamonds with Boltzmann fits). (**C**) Plots

of Po vs. V for 12 single G375R homomeric mutant channels following injection of G375R

mutant cRNA (blue squares with Boltzmann fits), and for 9 WT channels following injection of

WT cRNA (black circles and Boltzmann fits). (**D**) Identifying the types of assembled channels.

Blue and gray areas indicate the range of observed V_h values for G375R homomeric mutant

channels and WT channels, respectively, from *C*. One of the assembled channels (rightmost red

Po-*V* curve) overlaps with the WT channels (gray shading), indicating that this assembled

channel is likely a WT channel with four WT subunits. Four of the assembled channels (four

leftmost red Po-*V* curves) overlap with the G375R homomeric mutant channels (blue shading)

indicating that these four assembled channels are likely G375R homomeric mutant channels

assembled from four mutant subunits. The remaining 28 assembled channels would be hybrid

channels with mixed subunits (Fig. 1), as their V_h values do not overlap with those of either

homomeric mutant or WT channels.

Fig. 5 Five different types of functional BK channels are observed following injection of a 1:1

mixture of mutant and WT cRNA into *Xenopus* oocytes. (A) Histograms of V_h values of single

channels from Fig. 4. WT channels expressed after injection of only WT cRNA are indicated by

black bars, and G375R homomeric mutant channels expressed after injection of only G375R

cRNA are indicated by blue bars. Assembled channels expressed after a 1:1 injection of mutant

and WT cRNA are indicated by red bars. The red bar in the cluster of black bars is an assembled

channel comprised of four WT subunits. The four red bars interspaced between the blue bars

are assembled channels comprised of four mutant subunits. The 28 red bars with V_h values in

between the blue bars and black bars are hybrid assembled channels of mixed subunits. The

hybrid channels appear to cluster into three groups with mean V_h values of 1.4, 59 and 96 mV.

Hypothesized subunit compositions of the three apparent clusters of hybrid channels are

indicated, where black indicates WT subunits and blue indicates G375R mutant subunits. (**B**)

The experimentally observed frequencies of the five types of assembled channels are listed for

comparison to the theoretical frequencies in Fig. 1. (**C**) V_h for the five types of assembled

channels is approximated by a linear incremental model where each subunit contributes an

increment of V_h,

$$V_h(N_M) = N_M \ (\text{-16.7 mV}) + (4\text{-}N_M) \ 40.5 \text{ mV},$$

where N_M is the number of G375R mutant subunits per channel, $V_h(N_M)$ is V_h as a function of

N_M, and -16.7 mV and 40.5 mV are the increments of V_h added by each mutant and WT subunit,

respectively. Filled red circles are the experimentally observed mean ± SEM values of V_h for the

five types of assembled channels from part *A*, and the dashed line indicates the predicted

values. Predictions are for 0 intracellular Ca^{2+}. Mean ± SEM. Histograms of single-channel

kinetic parameters have been used previously to investigate the number of β2 regulatory

subunits per BK channel[50].

Fig. 6 Each replacement of a WT subunit with a G375R mutant subunit in a BK channel acts to

both left shift V_h to more negative potentials and decrease single channel conductance. (**A**)

Single-channel currents recorded from single channels at +100 mV after injecting the indicated

cRNA. (**B**) Single-channel conductance at +100 mV decreases as the number of mutant subunits

increase. Mean single-channel conductance for WT channels was 312 ± 4 pS (n = 7), decreasing

to 245 ± 6 pS for hybrid channels (n = 25, p < 0.0001), and further decreasing to 190 ± 14 pS for

G375R homomeric mutant channels (n = 8, p = 0.0003). Mean ± SEM. WT and G375R

homomeric mutant channels were those channels expressed after injecting only WT cRNA or

only G375R mutant cRNA, respectively. Hybrid channels were identified as in Fig. 4D. (**C**) Plot of

single channel conductance at +100 mV vs. V_h for BK channels expressed following injection of

the indicated cRNA. The linear regression line plots single channel conductance g vs. V_h, where:

$g = g(0) + aV$, where $g(0) = 231.9 ± 3.9$ pA and $a = 0.408 ± 0.0375$ pS/mV (p <0.0001 for slope

significantly different from 0. R = 0.86). (**D**) Single channel conductance for the five types of

assembled channels is predicted by a linear incremental model, where each subunit contributes

an increment of conductance,

$$g(N_M) = N_M\ 46.1\ \text{pS} + (4\text{-}N_M)\ 76.5\ \text{pS},$$

where N_M is the number of G375R mutant subunits per channel, $g(N_M)$ is single channel

conductance as a function of N_M, and 46.1 pS and 76.5 pS are the increments of g added by

each mutant and WT subunit, respectively. The homomeric mutant and WT data are from part

B, and the three red hybrid conductance bars plot g from the three groups of hybrid channels in

Fig. 5A. The red dashed line indicates the predicted values of g. Mean ± SEM.

Fig. 7 G375R homomeric mutant channels, assembled channels, and WT BK channels are

expressed in HEK293 cells. (**A**) Whole cell currents recorded from HEK293 cells following

transfection with the indicated types of cDNA. The whole cell currents were generated by

holding at -60 mV, with a pre-step to -100 mV, followed by steps from −200 or −100 to +200 mV

in 10 mV increments followed by a post-step to -120 mV. The dashed line indicates the level of

0 current. Evoked currents were not observed in HEK293 cells that were not transfected. (**B**)

Plots of normalized conductance versus the voltage of the activating steps following

transfections with G375R mutant cDNA (blue squares), a 1:1 mixture of mutant and WT cDNA

(red diamonds) to express assembled channels, or WT cDNA (black circles). Unlike single

channel recordings (Fig. 4), whole cell recordings reflect the average response from many

hundreds to thousands of channels. There were large negative shifts in V_h for G/Gmax curves

determined from G375R homomeric mutant channels and from assembled channels, when

compared to WT channels. The mean voltage for half activation (V_h) was 130.3 ± 4.4 mV (n = 4)

for WT BK currents, 37.5 ± 2.7 mV (n = 5) for BK currents from the 1:1 injection, and -142.5 ± 3.6

mV (n = 3) for the G375R homomeric mutant channels. The 1:1 transfection also decreased the

voltage sensitivity of activation compared to WT, with a slope of 52.0 ± 2.0 mV (n = 5) per e-fold

change for current after a 1:1 transfection, compared to 32.3 ± 2.5 mV (n = 4) for WT currents.

The G375R homomeric mutant channels had a slope of 45.0 ± 4.0 mV. Mean ± SEM.

Figure 1

Figure 2

Figure 3

Figure 4

A

B

C

Figure 5

Figure 6

Figure 7

BK channels of five different subunit combinations underlie the *de novo* KCNMA1 G375R channelopathy

Yanyan Geng, Ping Li, Alice Butler, Bill Wang, Lawrence Salkoff, and Karl L. Magleby

Contents: Supplementary Figs. 1-4

Supplementary Fig. 1 Cryo-EM ribbon structures of the Ca²⁺-free human BK
channel (PDB ID:6V3G)[1] showing the location of the mutant arginine side
chain G375R. (**A**) BK channel structure with the front and back (pore-forming alpha)
subunits removed. The arginine side chain of the G375R mutation is shown in space
filling format on the back side of S6 away from the conduction pathway on the red
colored subunit. Carbon atoms are gray, nitrogen blue and hydrogen not indicated.
(**B**) Structure after rotating 180° so that the location (blue mark) of the WT glycine
side chain of G375 is visible on the back side of S6 on the yellow subunit. (**C**) Six
013 possible orientations of the mutant arginine side chain are shown, generated
by ChimeraX software www.cgl.ucsf.edu/chimerax. The mutation G375D (G310D in
the referenced publication) also produces a left shift in activation[2] as G375R does, but
of half the magnitude. This raises the possibility that the left shift may be related to the
bulk of the substituted side chain rather than the specific charge, which is reversed for
these two mutations.

Supplementary Fig. 2 BK currents in *Xenopus* oocytes following injection of a 1:1 mixture
of G375R mutant and WT cRNA activate at greatly left shifted negative
voltages compared to BK currents following injection of only WT cRNA. I-V plots for
the indicated injections of cRNA and blank. Two-electrode whole cell voltage clamp. These
currents were used to calculate the relative conductance plotted in Fig. 2. Protocol
details are in Fig. 2 legend. The amount of WT cRNA injected was 25 times greater than
for the injection of the 1:1 mixture of G375R mutant and WT cRNA, so that the initial
deviation of WT currents from the baseline could be readily detected.

Supplementary Fig. 3 The aberrant negative shift in V_h for BK currents expressed following a 1:1 injection of G375R mutant and WT cRNA is also observed with 300 μM intracellular Ca^{2+}. (*A*) Currents recorded from inside-out macro patches of membrane with 300 μM Ca^{2+} at the inner membrane surface. For WT channels voltage pulses were from -160 mV to 180 mV with 20 mV steps from a holding potential of -160 mV. Conductance was measured from tail currents after stepping to -160 mV. For channels expressed following injection of a 1:1 mixture of G375R mutant and WT cRNA, voltage pulses were from -200 mV to 180 mV with 20 mV steps from a holding potential of -200 mV. Conductance was measured from tail currents after stepping to -200 mV. The dashed line indicates the level of 0 current. In the presence of 300 μM intracellulear Ca^{2+}, more than half of the channels expressed following the 1:1 injection of G375R mutant and WT cRNA remain open at -200 mV. The macro currents are the average response from many hundreds of BK channels in each macro patch. (*B*) G/Gmax vs. voltage plots following injection of WT cRNA alone (black circles) or a 1:1 injection of mutant and WT cRNA (red diamonds). The mean voltage for half activation V_h for WT currents was -7.2 ± 3.3 mV, shifting to -205 ± 9.6 mV for BK currents expressed from a 1:1 mixture of mutant and WT cRNA, producing a left shift of -198 mV with 300 μM Ca^{2+} at the inner membrane surface. This can be compared to a left shift of -120 mV with 0 Ca^{2+} (Fig. 3). Hence, the aberrant left shift in V_h induced by a 1:1 injection of mutant and WT cRNA occurs in the presence and absence of intracellular Ca^{2+}.

Supplementary Fig. 4 The experimentally observed percentages of the five types of assembled channels expressed following a 1:1 injection of G375R mutant and WT cRNA did not differ significantly from the theoretical percentages in Fig. 1 calculated assuming random assembly of mutant and WT subunits. To examine if the experimentally observed number of assembled channels of each type in Fig. 6 for the 33 assembled channels are consistent with random assembly of subunits, we compared the experimental number of each type to the expectations for random assembly. The expected number were determined by simulating 106 groups of 33 assembled channels and binning the number of assembled channels of each type for each group of 33 into five discrete probability distributions. The five panels in this figure present the five discrete probability distributions of the expected number of assembled channels of each type for a group of 33 channels. The channel type of each distribution is indicated by subunit composition, where black circles are WT subunits and blue circles are mutant

subunits. The topmost discrete probability distribution gives the mean probabilities based on random assembly of subunits (Fig. 1) of observing the indicated number of WT channels in a group of 33 assembled channels: 11.9% of the groups of 33 assembled channels would have 0 WT assembled channels, 26.2% of the groups of 33 assembled channels would have 1 WT assembled channel (red bar), 27.8% of the groups of 33 assembled channels would have 2 WT assembled channels, and so forth. The arrow at 2.06 WT channels indicates the mean number of predicted WT channels per group of 33 assembled channels, determined from the discrete probability distribution. The mean number can also be calculated directly from the percentages in Fig. 1: 6.25% of the assembled channels are WT times 33 assembled channels equals a mean of 2.06 WT assembled channels for each group of 33. Discrete probability distributions for each of the four other types of assembled channels in Fig. 1, are also presented and are interpreted in a similar manner. For example, the discrete probability distribution for channels with 1 mutant subunit and 3 WT subunits indicates that 7.34% of the groups of 33 assembled channels would contain 5 channels of this type, 11.5% of the groups of 33 channels would contain 6 channels of this type, and so forth.

Our experimental observations of the numbers of assembled channels of each type for the experimental group of 33 assembled channels (Fig. 4D and 6A, B) are indicated by the locations of the red histogram bars on the abscissa. The amplitude of the red bars gives the probability of observing the indicated number of channels of that type in the experimental group of 33 assembled channels. For example, the probability, assuming random assembly of subunits, for our observation of 1 WT channel in the experimental group of 33 assembled channels was 0.262, and the probabilities for our observations of 10, 11, 7, and 4 assembled channels for the other indicated types of assembled channels in the experimental group of 33 assembled channels were 0.118, 0.128, 0.147, and 0.0961. All of these probabilities for the observed numbers of assembled channels of each type were > 0.05 and none of the observed numbers of assembled channel types fell within the 0.05 cumulative probability of either the left or right tails of the discrete probability distributions, indicating that the experimental observations for the number of the five types of assembled channels were not significantly different[3] from the theoretical predictions based on an assumption of random subunit assembly. A lack of significant difference does not exclude that there may be underlying differences between experimental and theoretical predictions, which could be revealed by larger sample sizes.

References for the Supplementary Information

1. Tao, X. & MacKinnon, R. Molecular structures of the human Slo1 K$^+$ channel in complex with beta4. Elife 2019 Dec 9;8:e51409. doi: 10.7554/eLife.51409

2. Chen, X., Yan, J. & Aldrich, R. W. BK channel opening involves side-chain reorientation of multiple deep-pore residues. Proc. Nat. Acad. Sci. 111:E79-88 (2014).

3. D. Colquhoun, Lectures on Biostatistics (Clarendon Press, Oxford, 1971).