Transcriptome-wide analysis of microRNA-mRNA correlations in unperturbed tissue transcriptomes identifies microRNA targeting determinants

Juan Manuel Trinidad Barnech[123], Rafael Sebastián Fort[13], Guillermo Trinidad Barnech[4], Beatriz Garat[1], María Ana Duhagon[12*]

1 Laboratorio de Interacciones Moleculares, Facultad de Ciencias, UDELAR, Iguá 4225, 11400 Montevideo, Uruguay

2 Departamento de Genética, Facultad de Medicina, Universidad de la República, Montevideo, C.P. 11800, Uruguay

3 Departamento de Genómica, Instituto de Investigaciones Biológicas Clemente Estable, Av. Italia 3318, Montevideo CP 11600, Uruguay

4 Instituto de Computación, Facultad de Ingeniería, Universidad de la República, Montevideo, CP 11300, Uruguay

* Corresponding author: mduhagon@fcien.edu.uy/mduhagon@fmed.edu.uy (M.A.D.); Tel.: +598-2-525-8618 (ext. 7237), +598-2-924-3414 (ext. 3468)

Abstract

MicroRNAs are small RNAs that regulate gene expression through complementary base pairing with their target mRNAs. Given the small size of the pairing region and the large number of mRNAs that each microRNA can control, identifying biologically relevant targets is difficult. Since current knowledge of target recognition and repression has mainly relied on *in vitro* studies, we sought to determine if the interrogation of gene expression data of unperturbed tissues could yield new insight into these processes. The transcriptome-wide repression at the microRNA-mRNA canonical interaction sites (seed and 3′-supplementary regions, identified by sole base complementarity) was calculated as a normalized Spearman correlation (Z-score) between the abundance of the transcripts in the PRAD-TCGA tissues (RNA-seq and small RNA-seq data of 546 samples). Using the repression values obtained, we confirmed established properties or microRNA targeting efficacy, such as the preference for gene regions (3'UTR > CDS > 5'UTR), the correspondence between repression and seed length (6mer < 7mer < 8mer), and the contribution to the repression exerted by the supplementary pairing at 13-16nt of the microRNA. Our results suggest that the 7mer-m8 seed could be more repressive than the 7mer-A1, while they have similar efficacy when they interact using the 3'-supplementary pairing. Strikingly, the 6mer+suppl sites yielded a normalized Z-score of repression similar to the sole 7mer-m8 or 7mer-A1 seeds, alerting its potential biological relevance. We then used the approach to characterize the 3′-supplementary pairing further using 39 microRNAs that hold repressive 3'-supplementary interactions. The analysis of the bridge between seed and 3'-supplementary pairing site confirmed the optimum +1 offset previously evidenced, but higher offsets appear to have similar repressive strength. In addition, they show a low GC content at positions 13-16 and base preferences that allow a sequence motif identification. Overall, our study demonstrates that transcriptome-wide analysis of microRNA-mRNA correlations in large, matched RNA-seq and small-RNA-seq data can uncover hints of microRNA targeting determinants operating in the *in vivo* unperturbed set. Finally, we made available a bioinformatic tool to analyze microRNA-target mRNA interactions using our approach.

Keywords: microRNA, seed, 3´-supplementary, transcriptome, RNA-seq, small RNA-seq, TCGA, offset, bridge, loop.

Introduction

MicroRNAs are small, single-stranded RNAs that shape the expression of messenger RNAs (mRNAs) and are essential in metazoans. Because of this, microRNAs influence all developmental processes and diseases, including cancer (Bartel, 2018). In microRNA targeting, the seed region (nt 2–7 or 2–8, from the microRNA 5' end) is the primary determinant for targeting efficacy and specificity, which interact with the 3' untranslated region (UTR) and coding sequence (CDS) of target mRNAs (Grimson et al., 2007; Hafner et al., 2010; Helwak et al., 2013; Moore et al., 2015; Schnall-Levin et al., 2010). Since the seed region is only composed of 6–7 contiguous nucleotides, potential target sites are frequent in the transcriptome, but not all are functionally associated with microRNA-mediated repression (Baek et al., 2008; Selbach et al., 2008; Yue et al., 2009). Despite several years of effort to better understand the underlying factors and contexts involved in the efficient microRNAs repression of mammalian mRNAs, it is still challenging to identify functionally relevant interactions (Bartel, 2018).

It has been demonstrated that nucleotides outside of the seed of the microRNA can contribute to target recognition (Broughton et al., 2016; Duan et al., 2022; Grimson et al., 2007; Helwak et al., 2013; Moore et al., 2015; Salomon et al., 2015; Sheu-Gruttadauria et al., 2019; Wahlquist et al., 2014; Wee et al., 2012). The systematic analysis of microRNA likelihood pairing revealed the conservation of microRNA nucleotides 13–16 (Grimson et al., 2007). However, this 3'-supplementary pairing has shown little influence on on-site affinity and efficacy (Grimson et al., 2007; Salomon et al., 2015; Wee et al., 2012), and actually, after purifying selection, only about 5% of the seed-matched regulatory sites appear to involve it (Friedman et al., 2009).

More recent work has resolved the crystallography of human Ago2 with microRNA-122, uncovering microRNA-mRNA interactions through the 2-8 seed and the 3'-supplementary region (nucleotides 13-16) of the microRNA (Sheu-Gruttadauria et al., 2019). Contrary to previous reports, they demonstrate that optimal supplementary interactions can increase target affinity. They also show that the seed and the 3´-supplementary region can be bridged by an unstructured target loop of 1–15 nucleotides in the mRNA. Even when the loop region has base-pairing potential, Ago2 molecules maintain a stable conformation that avoids it (Sheu-Gruttadauria et al., 2019). To explain the contradictory results, Xiao & Macrae suggested that the differences found in the repression level between the global analyzes and those of specific microRNAs might be due to a "dilution" effect (Xiao & Macrae, 2020). Examining targets in batch may have masked the small number of targets undergoing strong supplementary repression among a larger number that are predicted to make supplementary interactions but weakly (Xiao & Macrae, 2020).

Members of a microRNA family have identical 5' sequences but divergent 3' sequence regions, thus providing an ideal source for assessing the contribution of the 3'-supplementary region to specificity and function (Broughton et al., 2016; Moore et al., 2015; Xiao & Macrae, 2020). Indeed, single-cell reporters on human cell lines have demonstrated that some 3'-supplementary pairing can provide target specificity for individual members of a microRNA family (Moore et al., 2015). Many reports illustrate the potential for microRNA 3'-pairing to

enable high-affinity target binding and suggest that the benefit of this pairing varies depending on the microRNA sequence and the 3′-pairing architecture. Nevertheless, understanding what features determine the use of 3′-pairing would require large measurements of multiple microRNA sequences (McGeary et al., 2022).

In the present study, we evaluate the contribution of the microRNA canonical seed and the 3'-supplementary region to target mRNA repression by analyzing all the evolutionary conserved microRNAs (Agarwal et al., 2015) and all mRNA transcripts expressed in prostate tissue samples of the PRAD-TCGA cohort. MicroRNA-mRNA interaction strength was evaluated by the Spearman correlation coefficient of the expression of both partners obtained from the RNA-seq transcriptomes of the PRAD-TCGA. Overall, our investigation suggests that the comparative analysis of RNA-seq transcriptomes of mRNAs and small RNAs of a large number of unperturbed tissues can analyze microRNA-mRNA interactions allowing the confirmation of previous rules and the discovery of potential novel variables of the interaction.

Methods and Data Sets

MicroRNAs and mRNAs transcriptomic data

RNAseq and small RNA-seq data of the TCGA-PRAD (normal and tumor conditions) were obtained from Firebrowser (firebrowser.org).

The microRNA conservation scores were obtained from TargetScan 7.2 database (Agarwal et al., 2015), and the 221 microRNAs with the highest "Family Conservation score" (score 2 defined by TargetScan), hereafter denominated "conserved microRNAs", were selected for this study (Supplementary Table 1).

Only microRNAs and mRNAs detected in at least 80% of the samples were included in the analysis (Supplementary Figure 1, Supplementary Table 1, see complete pipeline analysis in Figure 1). After filtering for transcript expression, 143 out of 221 microRNAs and 16786 out of 20531 large RNAs (mostly mRNA) were obtained.

Canonical seeds and 3'-supplementary microRNA-mRNA interactions identification

Mature human microRNA sequences were obtained from TargetScan 7.2 database (Agarwal et al., 2015). Human mRNA sequences of all protein-coding genes were obtained using BioMart (Ensembl GRCh38.13).

An *in-house* script written in Python programming language was developed to identify all the sequences complementary to all the canonical microRNA seeds (as described by TargetScan) of the 143 microRNAs (script available on GitHub). The TargetScan nomenclature was used to name the microRNA-mRNA interaction types, which comprise 6mer, 7mer-A1, 7mer-m8, and 8mer microRNA seeds (Supplementary Table 2 and Supplementary Figure 2). For this study, the 3'-supplementary region of an mRNA-microRNA interaction refers to four microRNA nucleotides at positions 13-16 that are complementary to mRNA nucleotides in the vicinity (Grimson et al., 2007; Sheu-Gruttadauria et al., 2019) (Figure 2A). A variable-length bridge of unpaired nucleotides can separate these two sequence modules. We investigated mRNA loops of a maximum of 15 nucleotides length based on previous functional studies (Sheu-Gruttadauria et al., 2019). The difference in the length of the microRNA and the mRNA unpaired bridges is referred to as the "offset" of the interaction involving supplementary sites.

We identified the eight types of putative microRNA-mRNA interactions described using Python regular expressions, comprising 6mer, 7mer-A1, 7mer-m8, 8mer, 6mer+suppl 7mer-A1+suppl, 7mer-m8+suppl, 8mer+suppl (Figure 2A). We chose the transcript variant with the highest number of putative microRNA interactions for each gene. In addition, per each microRNA, we selected the mRNAs that presented only one predicted microRNA-mRNA interaction, pondering the CDS, 3'UTR and 5'UTR independently. As a result, the computed mRNAs only have a single microRNA site at the region considered for every given microRNA analyzed, thus avoiding the co-occurrence of the sites. Of the 16,786 mRNAs and 143 microRNAs analyzed, 1,116,787 unique mRNA sites were identified for 193,203, 443,323 and 480,261 at 5'UTR, CDS and 3'UTR respectively (Supplementary Table 2, Supplementary Figure 2).

MicroRNA-mRNA pairs correlation analysis

For the selected list of microRNA and mRNAs, the normalized Spearman correlation coefficient (Z-score centered on microRNA) for the abundance of all putative microRNA-mRNA interacting pairs were calculated with the Python's Pandas library (McKinney Wes, 2010) using TCGA RNA-seq (mRNAs) and small RNA-seq (microRNAs) normalized data (546 tissue samples) obtained from FireBrowse (firebrowse.org).

Statistical analysis

Unless specified, statistical differences were assessed by Two-tailed T-test, and the p-values were adjusted by Bonferroni correction, using SciPy library version 1.6.2. Fisher exact test was performed using SciPy library version 1.6.2. In all cases, adjusted p-values < 0.05 were considered significant. All the plots were generated using Seaborn library 0.11.2.

Executable files

The executable files to search for microRNA/mRNA interaction sites were developed using PySimpleGUI 4.55.1 (graphical user interface) and PyInstaller 4.10 and are available on GitHub.

Results

Identification of microRNA-mRNA interaction sites in the PRAD-TCGA transcriptome based on nucleotide base complementarity.

Given the tissue specificity of microRNA expression, we performed this study within a single tissue type. We chose the PRAD cohort because it is large and extensively studied for microRNAs (Lin et al., 2021; Wei et al., 2020; Yang et al., 2019; Ye et al., 2018); in addition, PRAD is a very heterogeneous cancer thus biases result in of cancer subtypes are unlikely (Abeshouse et al., 2015). We investigated the expression of the conserved human microRNAs (221) and all the human protein-coding transcripts annotated in Ensembl (20,440) in the 546 samples of the PRAD-TCGA cohort (20,531) (see pipeline analysis in Figure 1). To increase the signal/noise ratio, we filtered out low expressed transcripts, arbitrarily defined as those undetected in ≥ 20% of the samples, narrowing the list of microRNAs and mRNA to 143 and

16,786, respectively (Supplementary Table 1 and Supplementary Figure 1). Aiming to compare the repressive power of different microRNA pairing sites, we studied 6mer, 7mer-A1, 7mer-m8, and 8mer microRNA seeds with or without the 3'-supplementary pairing regions, defined as the four nucleotides at positions 13-16 of the microRNA (hereafter denominated 6mer+suppl, 7mer-A1+suppl, 7mer-m8+suppl, and 8mer+suppl) (Figure 2A). The choice of the supplementary pairing site was based on the literature since mounting evidence of its repressive contribution has been published (references in Bartel 2018, Sheu-Gruttadaria et al. 2019). We determined all the mRNA sites with perfect complementarity to these eight canonical microRNA pairing sites, presuming they represent putative microRNA binding sites. We performed this analysis independently for each of the three regions of the mRNAs (5'UTR, CDS, and 3'UTR). The examination of seed sequence complementarity between the 143 microRNA and the 16,786 mRNA identified 3,065,731 putative microRNA interaction sites on the mRNAs (285,237 at 5'UTR, 1,226,568 at CDS, 1,553,926 at 3'UTR). Since each mRNA can bear more than one putative microRNA site for a given microRNA, in subsequent calculations, we only included the interactions involving microRNAs with a unique site in the mRNAs. Therefore, if an mRNA has more than one site for a given microRNA, these interactions were excluded from the analysis. Yet, if the same mRNA had single sites for other microRNAs, these interactions were compiled. We performed this calculation for each of the three mRNA regions independently. We iterated this analysis for each of the 143 selected microRNAs, obtaining a total of 1,116,787 sites (193,203 at 5'UTR, 443,323 at CDS, 480,261 at 3'UTR), corresponding to an average microRNA interaction number of 1,351±621 at the 5'UTR, 3,100±748 CDS, and 3,358±602 at the 3'UTR (Supplementary Table 2). As expected, for the eight interaction types in the three regions studied, the number of predicted interactions decreases with the seed's size and the incorporation of the 3'-supplementary regions (Supplementary Figure 2).

Figure 1. Data analysis pipeline

Inference of microRNA activity from microRNA-mRNA expression correlation in the PRAD-TCGA transcriptomes.

The availability of matched RNAseq and small-RNAseq and transcriptomes of hundreds of tissues due to global cancer initiatives allows the high-throughput transcriptome-wide assessment of microRNA-mRNA correlations. Our approach evaluates the strength of microRNA-mRNA interaction by the Spearman correlation coefficient of the expression of both molecules calculated from the RNAseq data of the 546 patient tissues (normal and tumor conditions) of the PRAD-TCGA cohort (Figure 1). Since the microRNAs can exert different effects on the expression of their target genes, we included both positive and negative correlations between the mRNA-microRNA pairs. Additionally, in the absence of consensus about the threshold value of significant biological correlation expected in tissues for direct microRNA repression, we analyzed all the correlations regardless of their magnitude.

Validation of microRNA repressive mechanisms using microRNA activity inferred from microRNA-mRNA expression correlations in PRAD-TCGA.

We initially sought to evaluate if the sole correlation of microRNA-mRNA pairs abundance in the tissues validates established microRNA repression rules. The computation of a Z-score of microRNA-mRNA correlation for each type of seed showed that repression levels increase with seed's length, i.e., the number of nucleotide bases involved in the interaction (6mer < 7mer-m8 < 7mer-A1 < 8mer) (Figure 2B and Supplementary Figure 3) (Table 1 and Supplementary Table 3), thus confirming a previously established rule of microRNA targeting (Friedman et al., 2009; Grimson et al., 2007; Hafner et al., 2010). These Z-score differences are statistically more significant for intra 3'UTR (average p-value = E-25), followed by intra CDS (average p-value = E-16) seed comparisons. Meanwhile, only 6mer/7mer-A1 and 7mer-A1/7mer-m8 differences are marginally significant at the 5'UTR (p-value 1.14 E-02 and 2.93 E-03, respectively). As expected, the strength of the mRNA target repression inferred from the correlations increases with the length of the microRNA/mRNA pairing region (Figures 2B and 2C), although the significance of the differences is influenced by the number of interactions in each dataset. Thus, the hierarchy of site efficacy is as follows: 8mer > 7mer > 6mer (Grimson et al., 2007; McGeary et al., 2019; Nielsen et al., 2007) (Figure 2B). In contrast to the previous findings (Friedman et al., 2009; Grimson et al., 2007), the Z-score indicates that the 7mer-A1 seed interaction is significantly more repressive than the 7mer-m8 in the three regions of the mRNA studied and also in comparison to the 6mer seed (Figure 2B). In contrast, the 7mer-m8 shows no difference from the 6mer seed in the UTRs (Figure 2B).

Figure 2. Repression of mRNA-target interactions with different pairing sites. The average Z-score correlations between the 143 conserved microRNAs selected and their predicted target mRNAs are represented for microRNA/mRNA interactions at the CDS, 5'UTR, and 3'UTR. Vertical bars represent the confidence interval (95%). Letter A in the sequences corresponds to Adenine, and Letter B corresponds to the other three nucleotides without Adenine.

Table 1

Group1	Group2	p-value		
		5′UTR	CDS	3′UTR
6mer	7mer-m8	n.s.	6.09E-03	n.s.
6mer	7mer-A1	1.14E-02	2.36E-42	4.82E-45
6mer	8mer	n.s.	1.74E-25	9.23E-35
7mer-m8	8mer	n.s.	2.24E-13	4.41E-23
7mer-A1	7mer-m8	2.93E-03	3.94E-14	9.48E-21
7mer-A1	8mer	n.s.	n.s.	1.24E-02
6mer	6mer suppl	n.s.	2.84E-08	9.90E-07
6mer	7mer-m8 suppl	n.s.	n.s.	3.66E-08
6mer	7mer-A1 suppl	n.s.	1.71E-04	2.48E-09
6mer	8mer suppl	n.s.	1.23E-02	3.72E-03
7mer-m8	7mer-m8 suppl	n.s.	n.s.	1.74E-06
7mer-m8	8mer suppl	n.s.	n.s.	1.52E-02
6mer suppl	7mer-m8	n.s.	2.84E-03	6.92E-04
6mer suppl	8mer	n.s.	n.s.	1.74E-03
7mer-A1 suppl	7mer-m8	n.s.	2.20E-02	2.43E-07

Table 2

Group1	Group2	p-value			
		6mer	7mer-A1	7mer-m8	8mer
3UTR	5UTR	5.98E-09	2.69E-14	4.34E-07	7.30E-12
3UTR	CDS	2.62E-08	1.37E-02	n.s.	n.s.
5UTR	CDS	n.s.	6.25E-08	6.82E-05	2.06E-07

Group1	Group2	p-value			
		6mer suppl	7mer-A1 suppl	7mer-m8 suppl	8mer suppl
3UTR	5UTR	2.81E-06	3.62E-02	1.14E-05	1.29E-03
3UTR	CDS	n.s.	n.s.	5.11E-03	n.s.
5UTR	CDS	7.30E-05	n.s.	n.s.	4.49E-03

Figure 2C presents an identical analysis carried out for the interaction sites involving seeds with base pairing possibilities at the 3'-supplementary regions. Again, most of the comparisons show more repression for longer seeds, which is more notoriously at the CDS and 3'UTR; the comparatively lower significance of the observed differences could be influenced by the smaller number of interactions considered in comparison to the sole seeds analysis (Figure 2B). The comparison of sole seeds and seeds plus 3'-supplementary regions shows that the supplementary interaction mainly contributes to repression, particularly at the 3'UTR and the CDS (Figure 2B and 2C and Supplementary Figure 3), although only the 6mer (CDS and 3'UTR) and 7mer-m8 (3'UTR) differences are statistically significant (p-value < 0.05) (Table 1). This finding from tissue data supports the contribution of the supplementary 13-16 nt microRNA-mRNA interaction to the microRNA repression that has been proposed using *in*

vitro approaches (Grimson et al., 2007; Helwak et al., 2013; Moore et al., 2015; Sheu-Gruttadauria et al., 2019; Xiao & Macrae, 2020). Additional validation of established rules of microRNA targeting mechanism comes from the observation that the relative magnitude of the repression of the microRNA/mRNA interactions is 3'UTR > CDS > 5'UTR (Figure 2B-C) (Grimson et al., 2007; Hafner et al., 2010; Helwak et al., 2013), which is supported by the p-values of the comparison among the sites in the three different regions (Table 2). This finding provides another proof of principle for using microRNA/mRNA correlations in unperturbed tissues to study *bona fide* microRNA repression characteristics.

Insight into the characteristics of microRNA-mRNA interaction involving beneficial supplementary pairing regions withdrawn from microRNA/mRNA correlations.

Our previous results drew our attention to the relevance of the 3'-supplementary region interaction for all types of seed; thus, we sought to investigate if the analysis of the microRNA activity inferred from the tissue transcriptomes could uncover features of the microRNA/mRNA pairing involving supplementary regions. For the following analyzes, we focus on the seed plus 3'-supplementary sites of 6mer+suppl (at 3'UTR and CDS) and 7mer+suppl. (at 3'UTR), since they showed a statistically significant increase in target repression compared to sole seeds (Table 1); therein, we selected the microRNAs that exert more repression on targets bearing 3'-supplementary interactions than on targets without supplementary pairing possibilities, as determined by a T-test of Z-score differences (p-value ≤ 0.05) (Supplementary Table 4). Overall, we found 39 repressive microRNAs in one or more of the three gene regions studied and 18 microRNAs with the opposite behavior (i.e., positively correlated with their target mRNAs). If each dataset is analyzed separately, the microRNAs that follow or oppose the repressive behavior are comparatively 19 vs. 11 (6mer 3'UTR, p-value = 0.2 binomial test of significance), 16 vs. 4 (7mer-m8 3'UTR, p-value = 0.01 binomial test of significance), and 21 vs. 8 (6mer CDS, p-value = 0.02 binomial test of significance). The 39 repressive microRNAs have no significant difference in expression (Supplementary Figure 4) or number of mRNA targets (Supplementary Figure 5) than the total 143 microRNAs analyzed (p-value ≤ 0.05).

High-throughput approaches showed that nucleotides within positions 8-13 of the microRNA are not paired in the AGO-microRNA-target complex (Hafner et al., 2010). Consistently, it was later shown that the seed and the 3'-supplementary regions could be bridged by an unstructured target loop of 1-15 nucleotides when complexed with AGO2, even when central complementary bases are available for pairing (Sheu-Gruttadauria et al., 2019). Moreover, 8mer+suppl interactions with bridges up to 10 nucleotides long in the target mRNA were more repressive than 8mer sole seeds. The same study proposes that interactions involving 3'-supplementary regions with high GC content can be established using bridges of up to 15 nt (Sheu-Gruttadauria et al., 2019). Since the bridge between the seed and the 3'-supplementary region of the microRNA is five nucleotides for the 6mer+suppl (8-12 nt) and four nucleotides for 7mer-m8+suppl (9-12 nt), the same length at the mRNA bridge means no loop formation in any of the two molecules (zero offset) (Figure 3A). Longer and shorter mRNA bridges imply a loop formation in the mRNA (positive offset values) or microRNA (negative offset values). Seeking to assess if the Z-score of microRNA repression withdrawn from the tissue PRAD-TCGA data was influenced by the offset of the microRNA/mRNA interaction, we analyzed the repression exerted by the 39 repressive microRNAs using the different offsets defined for their predicted interaction with their mRNA targets. As a control, we performed the same analysis for all the interactions involving 3'-supplementary region pairing of the 143 microRNAs. As

expected, their Z-scores are higher than those of the 39 repressive microRNAs, regardless of the seed type (6mer or 7merm8), gene location (CDS or 3´UTR), or the offset of the interaction (Figure 3B). Meanwhile, the supplementary interactions of the 39 repressive microRNAs show heterogeneous repression depending on the offset (Figure 3B, right panels). For the 6mer interactions occurring at the 3'UTR, a gradual gain of repression is observed with shorter microRNA loops (negative offsets), reaching their maximum repression at +0 offset. On the contrary, the length of the mRNA loops (offsets +1 to +10) is not directly proportional to the repressiveness of the 6mer interaction at the 3´UTR, yet the +1 offset is one of the most repressive lengths observed. A similar pattern of repression is depicted for the two other datasets (6mer CDS and 7mer-m8 3'UTR), though the differences are smoother, probably due to the smaller number of interactions composing them. To add up the information from the three datasets, we scaled their Z-scores using a minimum-maximum scaling (to normalize for their different Z-score ranges), and then we combined the three values. This integrated analysis indicates that the length of the microRNA loop is inversely proportional to the repression, while a one nucleotide loop in the mRNA (+1 offset) provides maximum repression (Supplementary Figure 6). Consistently, the optimum repression has been shown at +1 (McGeary et al., 2022; Sheu-Gruttadauria et al., 2019). Nevertheless, our results suggest that other offsets may produce similar repression.

Figure 3. Contribution of the mRNA bridge length to the microRNA repression. A) Graphic representation. The bridge region is defined as the unpaired nucleotides from the end of the seed to the beginning 3'-supplementary region. Since microRNA/mRNA supplementary interaction occurs at positions 13-16 of the microRNA, 6mer and 7mer-m8 seeds have 5 and 4 nt microRNA bridges, respectively; thus, these mRNA bridge lengths mean no loop formation for the respective seeds and are denominated "zero offset". Positive offset values

imply loop formation in the mRNA (positive offset values) and negative offset values in the microRNA (negative offset values). **B)** Analysis of bridge length effect on mRNA target repression (Z-transformed score) using the TCGA transcriptomes and the 143 total microRNAs included in this study as a control (left panels) and the 39 microRNAs with repressive supplementary interactions (right panels). The region of the genes and the type of seeds analyzed are indicated. The -log10(p- values) of the differences between each seed+suppl length and the respective sole seed interactions are indicated by the color intensity bar. Letter S stands for "seed" and indicates the Z-score of sole seed interactions.

Since base content bias has been previously proposed at the supplementary microRNA interaction sites (Sheu-Gruttadauria et al., 2019), we analyzed the GC content of the microRNAs with repressive supplementary pairing in comparison with the complete set of microRNAs. We observed a notorious decrease of the GC content in the stretch between positions 13-16 of the microRNAs with a repressive supplement for the three types of interactions analyzed (Figure 4A). In addition, positions 5 and 19 of 6mer 3'UTR sites have a GC content deviated from the total. The base composition analysis per position shows a preference for base A at positions 13-15 of the 3'-supplementary pairing region of the microRNA and for base T at position 16 (Figure 4B-D). The logo represented in Figure 4B resulted from the comparison of the full-length 39 microRNA mature sequences with the 143 microRNAs using the Discriminative Mode of the Motif Discovery of The MEME Suite (Bailey et al., 2015); it is supported by 16 microRNAs (E-value =4.7e-15). Although the enrichment of base T and A observed at positions 1 and 2 of all the microRNA in Figure 4C has been previously described in mammals and other species (Friedman et al., 2009; Wang, 2013), the logo enriched in the 39 selected microRNAs suggests an overrepresentation of these bases (Figure 4B).

Overall, microRNAs that establish repressive supplementary interactions with their targets seem to have low GC content at positions 13-16. Although the 6mer analysis suggests additional compositional skews, more interactions should be needed to confirm these hypotheses.

Figure 4. Nucleotide composition of the microRNAs with repressive supplementary interactions. A) Distribution of GC content along the positions of the microRNAs with the three repressive supplementary interactions indicated (colored) versus the whole set of microRNAs analyzed (black). Fisher exact test was performed to compare each of the represented sites with the total 143 microRNAs (● < 0.1, * < 0.05, *** < 0.001). **B)** MicroRNA site identified by the MEME suite (Discriminative Mode), using the 39 versus all the 143 microRNAs. The motif is represented by 16 of the 39 microRNAs (E-value = 4.7e-15). **C-F)** Base composition per microRNA nucleotide position. Seed types are represented by colors as indicated in A (B: blue/6mer 3'UTR, C: orange/7mer-m8 3'UTR, and D: green/6mer CDS, E: all microRNAs). Horizontal lines represent the ¼ frequency expected for a random nucleotide distribution.

An executable tool to identify the microRNA-mRNA seed+suppl interactions

Finally, we developed two Windows compiled executable scripts to search for seed+suppl interaction in human genes solely based on mRNA/microRNA base-pairing potential. One tool searches all the mRNA targets for a given microRNA, and the other finds all microRNAs that interact with a given mRNA transcript. The executable files identify 6-8mer seeds and supplementary interactions intervened by 1-15 nucleotides between the seed and the supplementary pairing regions. The search discriminates against CDS, 3' and 5' UTR, and all annotated transcript variants. The software, manual, and files are available on GitHub.

Discussion

The interaction between microRNAs and their target mRNAs can be inferred from the regulation of the target mRNA and protein levels provoked by the enforced modulation of the microRNA abundance in experimental settings (Baek et al., 2008; Grimson et al., 2007; Hafner et al., 2010; Helwak et al., 2013; Selbach et al., 2008). Despite the contribution to discovering target recognition features, these experimental approaches have the disadvantage of being limited to a few microRNAs assessed *in vitro*. Further evidence of the microRNA regulation *in vivo* has been withdrawn from the correlation between microRNA/mRNA pairs abundance in unperturbed cells or tissues, which is usually used as a criterion to support the microRNA regulation on single mRNA targets (Elton & Yalowich, 2015). Since Cancer Genome projects involving multi-omics methods have provided matched RNAseq and small-RNAseq of hundreds of tissues, they allow the high-throughput transcriptome-wide assessment of microRNA-mRNA expression correlations.

Here we sought to investigate if the sole analysis of microRNA-mRNA correlations in large datasets can validate established microRNA-mRNA pairing rules, which may indicate its putative power to uncover novel findings. We thus performed a transcriptome-wide approach to measure the correlation between all the mRNA transcripts and conserved microRNAs expressed in the PRAD-TCGA cohort. Our strategy is solely based on perfect Watson-Crick base complementarity at the 2-7 or 2-8 positions of the microRNA seed region to assign candidate microRNA interaction sites on the target mRNAs. This criterion reduces the various biases of the microRNA/mRNA target prediction algorithms (target site conservation, the free energy of the interaction, site accessibility, target-site abundance, local AU content, GU wobble pairing in the seed, position in the transcript, length of the transcript regions, machine learning features) (Peterson et al., 2014). In support of this notion, it has been stated that a simple text search yields predictions more reliable than some common algorithms (Bartel, 2018).

Furthermore, using this single criterion allows the analysis of a vast number of interactions, thus increasing the statistical significance of the findings. Indeed, although general rules of microRNA interactions represent major repression mechanisms, many interactions are expected not to comply with them. Another advantage of our approach is using the endogenous level of the microRNA-mRNA partners since concerns have been raised about artifactual observations of the perturbation-based approaches (Xiao & Macrae, 2020). Meanwhile, a limitation of the study is the exclusion of seed-site interactions with mismatches. In addition, the lack of inclusion of standard target prediction rules may decrease the biological meaning of the findings. Nevertheless, the choice to analyze conserved microRNAs favors the compliance to mechanistically conserved microRNA-target interaction properties.

Our initial analysis allowed the confirmation of well-established rules or microRNA target regulation, such as the dependence of repression on the gene region (3'UTR > CDS > 5'UTR), the seed length (6mer < 7mer < 8mer), and the contribution of the supplementary pairing at 13-16 of the microRNA (Agarwal et al., 2015; Bartel, 2009, 2018; Grimson et al., 2007). It is worth noting that, in terms of inferred repression, the 6mer+suppl interaction is not significantly different from the 7mer-A1, which raises the awareness of the possibly underestimated relevance of the 6mer interaction *in vivo*. Although the 6mer site *per se* is known to be less specific for repression than longer seeds and is classified as a "marginal site" (Agarwal et al.,

2015; Grimson et al., 2007), the addition of a supplementary four nucleotide base pairing (microRNA position 13-16) may confer a more significant Argonaut mediated repressive activity. Additionally, contrary to previous reports (Grimson et al., 2007; Nielsen et al., 2007), our data suggest that the 7mer-A1 seed is significantly more repressive than the 7mer-m8 in the three transcript regions analyzed. Further investigation is needed to understand if this unexpected finding is due to the nature of the models studied (*in vitro* vs. *in vivo*), the bioinformatic pipelines used, or another source variation.

The functionality of the 3'-supplementary pairing in microRNA-mRNA interactions was described several years ago (Grimson et al., 2007), and further insight into its molecular basis was recently achieved (Bibel et al., 2022; McGeary et al., 2022; Sheu-Gruttadauria et al., 2019; Xiao & Macrae, 2020). Unlike the first studies that used a 2-6 nt bridge length, recent publications showed that up to 15 nt long loops on the target mRNA could bridge the seed and the 3'-supplementary region (Sheu-Gruttadauria et al., 2019). The different views of the relative contribution of the supplementary pairing to the target repression have been ascribed to the different methodological strategies of the analyses (Xiao & Macrae, 2020). In this context, we thought our approach would be useful to contribute to the understanding of the supplementary 13-16 nt interaction because it includes a large number of interactions, potentially allowing the identification of features of the supplementary interaction, such as the bridge length, base preferences, or repression strength.

We found that supplementary nucleotide pairing potentiates the repressor activity of 6mer and 7mer-m8 seeds in at least 39 conserved microRNAs. These interactions appear functional in the 3'UTR and coding sequence of mRNAs targets. The analysis of bridge length contribution to repression using our pipeline results consistent with those reported in structural and high throughput *in vitro* studies, where the optimal length was found near or at the +1 offset (McGeary et al., 2022; Sheu-Gruttadauria et al., 2019). Moreover, the pattern of repression for offsets larger than +1 (until +10-11 was here analyzed) suggests that the 3´-supplementary pairing can be equally repressive at higher offsets. However, the heterogeneity of the repression per offset would be compatible with heterogeneous 3' microRNA architecture (McGeary et al., 2022). Interestingly, the two binding modes recently reported for compensatory interactions in McGeary et al. (offset 0-1 and +3-4) are supported by our findings of a second repressive offset at +3-4 offset. In addition, we found strong repression of a +7 loop that has not been previously reported. It is likely that more data, *i.e.*, more transcriptomes and additional tissues, would be needed to increase the power of the analysis. We also identified sequence patterns enriched in the 39 repressive supplementary microRNAs, including a low GC content with a higher A proportion at the 3'-supplementary region and a preferred sequence motif, whose relevance must be validated by additional methods. Previously, a low GC content in this region was associated with a lower affinity of the interaction evaluated *in vitro* using a specific microRNA-RNA target pair (McGeary et al., 2022; Sheu-Gruttadauria et al., 2019; Xiao & Macrae, 2020); this discrepancy may be due to the number of microRNAs, the context of the interaction (*in vitro*/*in vivo*) or to still unknown factors.

Our study represents a novel high-throughput approach to investigate the microRNA-mRNA target interaction by withdrawing patterns from transcriptome-wide expression correlations in large unperturbed tissue datasets. The findings confirm existing knowledge, which supports the biological meaning of the approach, so we used it to get a deeper understanding of the 3'-supplementary interaction sites. Due to the relevance of 6mer+suppl and the length of the bridge between the seeds and supplementary sites, we built two Windows compiled

executable scripts that predict the microRNA-mRNA interactions defined in our current pipeline, which are available on [GitHub](#).

Author Contributions: Conceptualization, M.A.D., R.S.F.; methodology, R.S.F, J.M.T.; software, R.S.F, J.M.T., G.T.; formal analysis, M.A.D., R.S.F, J.M.T.; data curation, R.S.F, J.M.T.; writing—original draft preparation, J.M.T. and M.A.D.; writing—review and editing M.A.D., R.S.F., J.M.T.; project administration, M.A.D. and B.G.; funding acquisition, M.A.D. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by the Comisión Sectorial de Investigación Científica (CSIC-UDELAR, Uruguay) (Research Grants M.A.D. I+D 2016 #487 and I+D 2020 # 566) and the Programa para el Desarrollo de las Ciencias Básicas (PEDECI-BA-MEC, Uruguay, Annual Aliquots and Equipment Aid Programs of). Postgraduate Students fellowships for J.M.T. and R.F. were provided by the University of the Republic (UDELAR), the Comision Academica de Posgrado (CAP-CSIC-UDELAR) and the Agencia Nacional de Investigación e Innovación (ANII).

References

Abeshouse, A., Ahn, J., Akbani, R., Ally, A., Amin, S., Andry, C. D., Annala, M., Aprikian, A., Armenia, J., Arora, A., Auman, J. T., Balasundaram, M., Balu, S., Barbieri, C. E., Bauer, T., Benz, C. C., Bergeron, A., Beroukhim, R., Berrios, M., … Zmuda, E. (2015). The Molecular Taxonomy of Primary Prostate Cancer. *Cell*, *163*(4), 1011–1025. https://doi.org/10.1016/j.cell.2015.10.025

Agarwal, V., Bell, G. W., Nam, J. W., & Bartel, D. P. (2015). Predicting effective microRNA target sites in mammalian mRNAs. *ELife*, *4*(AUGUST2015), 1–38. https://doi.org/10.7554/eLife.05005

Baek, D., Villén, J., Shin, C., Camargo, F. D., Gygi, S. P., & Bartel, D. P. (2008). The impact of microRNAs on protein output. *Nature*, *455*(7209), 64–71. https://doi.org/10.1038/nature07242

Bailey, T. L., Johnson, J., Grant, C. E., & Noble, W. S. (2015). The MEME Suite. *Nucleic Acids Research*, *43*(W1), W39–W49. https://doi.org/10.1093/NAR/GKV416

Bartel, D. P. (2009). MicroRNAs: Target Recognition and Regulatory Functions. *Cell*, *136*(2), 215–233. https://doi.org/10.1016/j.cell.2009.01.002

Bartel, D. P. (2018). *Metazoan MicroRNAs*. https://doi.org/10.1016/j.cell.2018.03.006

Bibel, B., Elkayam, E., Silletti, S., Komives, E. A., Joshua-Tor, L., & Leemor, J.-T. (2022). Target binding triggers hierarchical phosphorylation of human Argonaute-2 to promote target release. *Biorxiv*. https://doi.org/10.1101/2022.01.06.475261

Broughton, J. P., Lovci, M. T., Huang, J. L., Yeo, G. W., & Pasquinelli, A. E. (2016). Pairing beyond the Seed Supports MicroRNA Targeting Specificity. *Molecular Cell*, *64*(2), 320–333. https://doi.org/10.1016/j.molcel.2016.09.004

Duan, Y., Veksler-Lublinsky, I., & Correspondence, V. A. (2022). Critical contribution of 3' non-seed base pairing to the in vivo function of the evolutionarily conserved let-7a microRNA. *CellReports*, *39*, 110745. https://doi.org/10.1016/j.celrep.2022.110745

Elton, T. S., & Yalowich, J. C. (2015). Experimental procedures to identify and validate specific mRNA targets of miRNAs. *EXCLI Journal*, *14*, 758. https://doi.org/10.17179/EXCLI2015-319

Friedman, R. C., Farh, K. K. H., Burge, C. B., & Bartel, D. P. (2009). Most mammalian mRNAs are conserved targets of microRNAs. *Genome Research*, *19*(1), 92–105. https://doi.org/10.1101/gr.082701.108

Grimson, A., Farh, K. K. H., Johnston, W. K., Garrett-Engele, P., Lim, L. P., & Bartel, D. P. (2007). MicroRNA Targeting Specificity in Mammals: Determinants beyond Seed Pairing. *Molecular Cell*, *27*(1), 91–105. https://doi.org/10.1016/j.molcel.2007.06.017

Hafner, M., Landthaler, M., Burger, L., Khorshid, M., Hausser, J., Berninger, P., Rothballer, A., Ascano, M., Jungkamp, A. C., Munschauer, M., Ulrich, A., Wardle, G. S., Dewell, S., Zavolan, M., & Tuschl, T. (2010). Transcriptome-wide Identification of RNA-Binding Protein and MicroRNA Target Sites by PAR-CLIP. *Cell*, *141*(1), 129–141. https://doi.org/10.1016/j.cell.2010.03.009

Helwak, A., Kudla, G., Dudnakova, T., & Tollervey, D. (2013). Mapping the human miRNA interactome by CLASH reveals frequent noncanonical binding. *Cell*, *153*(3), 654–665. https://doi.org/10.1016/j.cell.2013.03.043

Lin, B. B., Lei, H. Q., Xiong, H. Y., Fu, X., Shi, F., Yang, X. W., Yang, Y. F., Liao, G. L., Feng, Y. P., Jiang, D. G., & Pang, J. (2021). MicroRNA-regulated transcriptome analysis identifies four major subtypes with prognostic and therapeutic implications in prostate cancer. *Computational and Structural Biotechnology Journal*, *19*, 4941–4953. https://doi.org/10.1016/J.CSBJ.2021.08.046

McGeary, S. E., Bisaria, N., Pham, T. M., Wang, P. Y., & Bartel, D. P. (2022). MicroRNA 3′-compensatory pairing occurs through two binding modes, with affinity shaped by nucleotide identity and position. *ELife*, *11*, 1–37. https://doi.org/10.7554/ELIFE.69803

McGeary, S. E., Lin, K. S., Shi, C. Y., Pham, T. M., Bisaria, N., Kelley, G. M., & Bartel, D. P. (2019). The biochemical basis of microRNA targeting efficacy. *Science*, *366*(6472), 1–20. https://doi.org/10.1126/science.aav1741

McKinney Wes. (2010). Data Structures for Statistical Computing in Python. *Proceedings of the 9th Python in Science Conference (SCIPY 2010)*, *41*(3), 51–56.

Moore, M. J., Scheel, T. K. H., Luna, J. M., Park, C. Y., Fak, J. J., Nishiuchi, E., Rice, C. M., & Darnell, R. B. (2015). miRNA–target chimeras reveal miRNA 3′-end pairing as a major determinant of Argonaute target specificity. *Nature Communications*, *6*(May), 8864. https://doi.org/10.1038/ncomms9864

Nielsen, C. B., Shomron, N., Sandberg, R., Hornstein, E., Kitzman, J., & Burge, C. B. (2007). Determinants of targeting by endogenous and exogenous microRNAs and siRNAs. *RNA (New York, N.Y.)*, *13*(11), 1894–1910. https://doi.org/10.1261/RNA.768207

Peterson, S. M., Thompson, J. A., Ufkin, M. L., Sathyanarayana, P., Liaw, L., & Congdon, C. B. (2014). Common features of microRNA target prediction tools. *Frontiers in Genetics*, *5*(FEB), 23. https://doi.org/10.3389/FGENE.2014.00023/BIBTEX

Salomon, W. E., Jolly, S. M., Moore, M. J., Zamore, P. D., & Serebrov, V. (2015). Single-Molecule Imaging Reveals that Argonaute Reshapes the Binding Properties of Its Nucleic Acid Guides. *Cell*, *162*(1), 84–95. https://doi.org/10.1016/j.cell.2015.06.029

Schnall-Levin, M., Zhao, Y., Perrimon, N., & Berger, B. (2010). Conserved microRNA targeting in Drosophila is as widespread in coding regions as in 3′UTRs. *Proceedings of the National Academy of Sciences of the United States of America*, *107*(36), 15751–15756. https://doi.org/10.1073/pnas.1006172107

Selbach, M., Schwanhäusser, B., Thierfelder, N., Fang, Z., Khanin, R., & Rajewsky, N. (2008). Widespread changes in protein synthesis induced by microRNAs. *Nature*, *455*(7209), 58–63. https://doi.org/10.1038/nature07228

Sheu-Gruttadauria, J., Xiao, Y., Gebert, L. F., & MacRae, I. J. (2019). Beyond the seed: structural basis for supplementary microRNA targeting by human Argonaute2. *The EMBO Journal*, e101153. https://doi.org/10.15252/embj.2018101153

Wahlquist, C., Jeong, D., Rojas-Muñoz, A., Kho, C., Lee, A., Mitsuyama, S., Van Mil, A., Jin Park, W., Sluijter, J. P. G., Doevendans, P. A. F., Hajjar, R. J., & Mercola, M. (2014). Inhibition of miR-25 improves cardiac contractility in the failing heart. *Nature*, *508*(7497), 531–535. https://doi.org/10.1038/nature13073

Wang, B. (2013). Base composition characteristics of mammalian mirnas. *Journal of Nucleic Acids*, *2013*. https://doi.org/10.1155/2013/951570

Wee, L. M., Flores-Jasso, C. F., Salomon, W. E., & Zamore, P. D. (2012). Argonaute divides Its RNA guide into domains with distinct functions and RNA-binding properties. *Cell*, *151*(5), 1055–1067. https://doi.org/10.1016/j.cell.2012.10.036

Wei, J., Yin, Y., Deng, Q., Zhou, J., Wang, Y., Yin, G., Yang, J., & Tang, Y. (2020). Integrative Analysis of MicroRNA and Gene Interactions for Revealing Candidate Signatures in

Prostate Cancer. *Frontiers in Genetics*, *11*, 176. https://doi.org/10.3389/FGENE.2020.00176/BIBTEX

Xiao, Y., & Macrae, I. J. (2020). Robust differential microRNA targeting driven by supplementary interactions in vitro. *Rna*, *26*(2), 162–174. https://doi.org/10.1261/rna.072264.119

Yang, Y., Jia, B., Zhao, X., Wang, Y., & Ye, W. (2019). miR-93-5p may be an important oncogene in prostate cancer by bioinformatics analysis. *Journal of Cellular Biochemistry*, *120*(6), 10463–10483. https://doi.org/10.1002/jcb.28332

Ye, Y., Li, S. L., & Wang, S. Y. (2018). Construction and analysis of mRNA, miRNA, lncRNA, and TF regulatory networks reveal the key genes associated with prostate cancer. *PLOS ONE*, *13*(8), e0198055. https://doi.org/10.1371/JOURNAL.PONE.0198055

Yue, D., Liu, H., & Huang, Y. (2009). Survey of Computational Algorithms for MicroRNA Target Prediction. *Current Genomics*, *10*(7), 478–492. https://doi.org/10.2174/138920209789208219