The antique genetic plight of the Mediterranean monk seal (*Monachus monachus*)

Jordi SALMONA[1*], Julia DAYON[1,2], Emilie LECOMPTE[1], Alexandros A. KARAMANLIDIS[3], Alex AGUILAR[4], Pablo FERNANDEZ DE LARRINOA[5], Rosa PIRES[6], Giulia MO[7], Aliki PANOU[8], Sabrina AGNESI[7], Asunción BORRELL[4], Erdem DANYER[9,10], Bayram ÖZTÜRK[9,11], Arda M. TONAY[9,11], Anastasios K. ANESTIS[8], Luis M. GONZÁLEZ[12], Panagiotis DENDRINOS[3], Philippe GAUBERT[1*]

1. Laboratoire Évolution & Diversité Biologique, IRD-CNRS-UPS, Université Paul Sabatier, 118 route de Narbonne, 31062 Toulouse, France
2. CEFE, Univ Montpellier, CNRS, EPHE-PSL University, IRD, Montpellier, France
3. MOm/Hellenic Society for the Study and Protection of the Monk seal, Solomou Str. 18, 10682 Athens, Greece
4. Universitat de Barcelona, IRBio and Department of Evolutionary Biology, Ecology and Environmental Sciences, Faculty of Biology, Diagonal 643, 08028 Barcelona, Spain
5. Monk Seal Conservation Program, Fundación CBD-Habitat, Gustavo Fernández Balbuena 2, 28002 Madrid, Spain
6. Instituto das Florestas e Conservação da Natureza IP-RAM, Jardim Botânico da Madeira, Caminho do Meio, Bom Sucesso, 9064-512, Funchal, Madeira, Portugal
7. Istituto Superiore per la Protezione e la Ricerca Ambientale (ISPRA), Via Vitaliano Brancati 48, 00144, Rome, Italy
8. Archipelagos - environment and development, Lourdata, 281 00 Kefalonia, Greece
9. Turkish Marine Research Foundation (TUDAV), PO Box 10, Beykoz, Istanbul, Turkey
10. Veterinary Control Central Research Institute, Ankara, Turkey
11. Istanbul University, Faculty of Aquatic Sciences, Ordu Cad. No: 8 Laleli, Istanbul, Turkey
12. Subdirección General de Biodiversidad Terrestre y Marina, Ministerio para la Transición Ecológica y el Reto Demográfico, Pza. San Juan de la Cruz, 10. E-28071. Madrid, Spain

*Corresponding authors:
Jordi SALMONA: jordi.salmona@gmail.com
Philippe GAUBERT: philippe.gaubert@univ-tlse3.fr

Abstract

Disentangling the impact of Late Quaternary climate change from human activities can have crucial implications on the conservation of endangered species. We investigated the population genetics and demography of the Mediterranean monk seal (*Monachus monachus*), one of the world's most endangered marine mammals, through an unprecedented dataset encompassing historical (extinct) and extant populations from the eastern North Atlantic to the entire Mediterranean Basin. We show that Cabo Blanco (Western Sahara/Mauritania), Madeira, Western Mediterranean (historical range), and Eastern Mediterranean regions segregate into four populations. This structure is likely the consequence of recent drift, combined with long-term isolation by distance (R^2 = 0.7), resulting from prevailing short-distance (< 500 km) and infrequent long-distance dispersal (< 1,500 km). All populations (Madeira especially), show high levels of inbreeding and low levels of genetic diversity, seemingly declining since historical time, but surprisingly not being impacted by the 1997 massive die-off in Cabo Blanco. Approximate Bayesian Computation analyses support scenarios combining local extinctions and a major effective population size decline in all populations during Antiquity. Our results suggest that the early densification of human populations around the Mediterranean Basin coupled with the development of seafaring techniques were the main drivers of the decline of Mediterranean monk seals.

Keywords

Mediterranean monk seal, Population decline, Demographic history, Marine mammals, Genetic diversity, Isolation by distance.

Background

Distinguishing between the respective impact of recent climate- and human-driven changes on the biosphere has proven challenging [2] because Late Quaternary extinctions were caused by the superimposed effects of climate change and anthropization from the Last Glacial Maximum (LGM) onwards (ca. 25–10 ka; [1]). Differentiating climate- and human-driven impacts is similarly relevant to currently endangered species. Indeed, informed conservation planning relies on a systemic approach including knowledge of species' demographic history [2], which in turn can be used to predict species' ability to adapt to future climate changes [3,4].

Pinnipeds are marine mammals that rely on coastal haul-out areas during their annual life cycle. As such, they have been affected by LGM climate changes and early human activities [5,6], although the impact of the latter (through targeted hunting) occurred after the end of the LGM in this case (but see [7]). Despite the fact that the Mediterranean monk seal (MMS; *Monachus monachus*) is arguably the world's most endangered pinniped [8], the factors responsible for its critical conservation status are not well understood. The MMS once ranged across the entire Temperate Northern Atlantic province [9], from the Black Sea and the Mediterranean Basin into North Atlantic eastern waters encompassing the coasts of western Africa, the Macaronesian islands, and the northern Iberian Peninsula [10–12]. Nowadays, the species is fragmented into three isolated areas, distributed in the eastern Mediterranean (ca. 187-240 mature individuals [13]) and the eastern North Atlantic, in Cabo Blanco (Western Sahara/Mauritania; ca. 350 individuals) and the archipelago of Madeira (ca. 20 individuals; [14]) (Fig. 1).

Figure 1. *Monachus monachus* nuclear genetic structure.
Individual posterior cluster membership coefficients for *K* = 4 represented using vertical barplot (**A**) and geolocated pie-charts (**B**). Representation of the samples from the six main sampling areas on the first two axes of principal

component analysis (**C**) showing a strong West-East cline of differentiation on the first axis. In **A**, the samples are ordered by regions, sorted from West to East, separated by vertical black lines, and stars denote historical samples. In **B**, black circles denote locations that include historical samples and the cyan area represents the putative ancient MMS distribution. In **C**, full and empty symbols represent historical and modern samples, respectively. W-Sahara: Cabo Blanco (Western Sahara/Mauritania), W-Med: western Mediterranean Sea, C-Med: central Mediterranean Sea, E-Med: eastern Mediterranean Sea.

A combination of extrinsic (human activities) and intrinsic (e.g., low genomic diversity, high susceptibility to diseases) factors have been proposed to explain the local extinctions affecting the species [15,16], and to a larger extent, the extinction proneness observed in the entire Monachinae subfamily (which also includes the endangered Hawaiian monk seal *Neomonachus schauinslandii* and the extinct Caribbean monk seal *N. tropicalis* [17]). As an iconic representative of the Mediterranean Basin biodiversity hotspot [18], the MMS could have suffered from targeted hunting since the advent of the early Mediterranean sailors (Bronze Age, ~3,300 to 1,200 BCE; [19–21]). From the end of the Middle Age (15th century), the massive exploitation of the species−notably in the Atlantic Ocean−became documented through the logbooks of European marine explorers [11,22]. Over the last centuries, MMS populations have been further impacted by the expansion of the fishing industry, deliberate killing by fishermen, marine pollution, and human coastal encroachment, leading to local extinctions in most of the species' range, notably in the western Mediterranean and the Black Sea [12,23–25].

The decline of the MMS is documented from historical mass killing [10,19,26], local extinctions [27], massive die-off [28], and genetic inferences [29]. However, the respective impacts of Late Quaternary climatic fluctuations and anthropogenic pressures on the demographic history of the species remain poorly understood. Yet, the LGM induced significant changes in the Mediterranean Basin, such as lowering of sea level (down to -120 m; [30]) and sea surface temperatures, and local variations in salinity [31,32]. Moreover, the impacts of human activities on MMS populations may have applied at different periods across the species' range, with early exploitation since the Middle Paleolithic [20,33], followed by a gradual dispersal of seafaring civilizations from the eastern Mediterranean Basin (where coastal encroachment and marine resources' exploitation have been documented since ca. 10,000 yrs) towards the western Mediterranean Basin and North Atlantic Ocean [21,30,33,34].

Previous genetic studies have revealed the genetic isolation of the three extant populations of MMS and their low levels of genetic diversity (since at least the mid-19th century), and the local extinction of mitochondrial DNA (mtDNA) haplotypes in the western Mediterranean Basin [35–38]. Genetic data additionally disclosed a demographic bottleneck in the Cabo Blanco [29,39], and a signature of past metapopulation dynamics across the species' range [37,40]. In this study, we investigate the genetic patterns and demographic history of the MMS through an unprecedented dataset, including historical individuals collected across the majority of the species' range and the three extant populations. We use microsatellite genotyping and mtDNA sequencing of 375 MMS to assess the historical population structure and dynamics of the MMS across the Mediterranean Basin and Northern Atlantic Ocean. Specifically, we delineate past and present population structure and diversity and assess whether such patterns are consistent with scenarios of local extinctions together with genetic drift and inbreeding in extant populations. We also investigate the different historical drivers (LGM *vs.* human activities) potentially affecting the MMS population dynamics across the Mediterranean Basin and

130 Northern Atlantic Ocean through complementary population genetic and modeling approaches
131 based on demographic models, incorporating population structure and connectivity among
132 populations through time. Based on our results, we formulate recommendations that might
133 contribute to future evidence-based conservation strategies of the MMS.

Methods

Laboratory procedures and genotyping

We collected 383 samples from recent (N=314; 1989 – 2020) and historical (N=69; 1833 –
1975) specimens of MMS covering the extant populations' and historical species' range (Figs 1,
S1-3). The delineation between recent (from 1989 on) *vs*. historical (prior to 1975) is based on
the extinction period of the last resident MMS groups from the western and parts of the central
Mediterranean Sea during the 1970s [19], which suggested that after this period the species
distribution fragmented into North Atlantic and eastern Mediterranean populations.
Genomic DNA was extracted from fresh tissue and skin samples, hairs, feces, bones,
and tanned skin using dedicated protocols (Method S1). We amplified 524 bp of the
hypervariable region I of the mitochondrial control region (CR1) following [41] for the modern
samples (N = 121), and reconstructed 484 bp fragment encompassing all the variable sites of
CR1 for the historical samples (N = 3), as already described elsewhere [35] and detailed in
method S2. We complemented our original dataset with 209 already published CR1 sequences
[35,41]. Final CR1 alignment included 299 modern, 23 dated historical samples, and four
historical samples with unknown dating, for a total of 326 sequences (Figs 1, S1-3, Tables S1a ,
S1b & 2).
We genotyped 383 samples at 19 nuclear microsatellite loci ([38], Method S3) from 314
modern samples and 69 historical samples. In order to mitigate scoring errors, PCRs of DNA
extracts from hairs, feces, and museum material were systematically replicated 2-to-5 times
([38], Method S3). We controlled markers for linkage disequilibrium and departure from Hardy –
Weinberg equilibrium (Method S3, Table S3; Fig. S4-7), and applied a two-fold sample selection
procedure to optimize the number of retained individuals genotyped at informative loci. We
relied on the minimum number of loci necessary to discriminate among individuals and the
region-based discriminant analysis of principal components [DAPC] contribution of loci (Method
S3, Fig. S8). The final microsatellite data set comprised 253 samples (including 14 historical
and 239 modern samples) with 10.18 % missing data (Tables S1a & S1b, Figs 1 & S1-2).

Genetic diversity and structure

To assess the overall, per sampling site, and per locus microsatellites genetic diversity,
we estimated the allelic richness (A_R [42]) with the R package *hierfstat* [43]. We also assessed
the number of alleles (A), the observed (H_O) and Nei's unbiased expected heterozygosities (H_E
[44]) with the R package *adegenet* [45]. To evaluate the effect of drift in the small colonies of
MMS, we estimated the average individual inbreeding coefficient F [45] in *adegenet*, from 100
iterations. We assessed the level of genetic differentiation among localities, populations, and

time periods using Nei's F_{ST} [46]. Additionally, we investigated patterns of genetic variance with
a principal component analysis (PCA) of allele frequencies, with a discriminant analysis of
principal components (DAPC, considering sampling regions as groups), and using the
SnapClust clustering approach [47] for K = 1-15. To assess how the geographic distance alone
explains the genetic diversity [48,49], we investigated individual- and population-based patterns
of isolation by distance (IBD) using Mantel tests [50], and assessed which genetic distance
metric and spatial scale best fitted the data [51,52]. To assess how dispersal is distributed
geographically we used Mantel correlograms [53,54].
For CR1, we estimated the number of haplotypes (Nh) and polymorphic sites (S),
haplotype diversity (h [55]), and nucleotide diversity (π [56]) per sampling area using the R
package *pegas* [57]. CR1 haplotype relationships were reconstructed with a maximum
parsimony network using *pegas*. All the above-mentioned methods are detailed in Method S4.

Demographic history

We used Approximate Bayesian Computation (ABC; [58]) to compare data simulated
under several alternative scenarios to the real data (microsatellites & CR1), and estimate
parameters of interest from the best-supported scenario [59–61]. We first tested scenarios
assuming a single panmictic population and different histories of population size change (Figs
S9-10). Second, we modeled stable-size structured populations with constant migration rates
among populations (Figs S9, S11). These models included unsampled ghost populations
mimicking extinct populations and a change in connectivity, mimicking the loss of gene flow
among extant populations (Figs S9, S11, S12). Third, we combined population size change,
structure, ghost populations, and changes in connectivity, to model populations that suffered
one or several events of decline and fragmentation (Figs S9, S13). Structured scenarios also
compared n-island, stepping-stone, and spatially-explicit models of connectivity (Table S4, S5).
To overcome the potential overfitting of parameters, we reduced the summary statistics
dimension [62], both using partial least square regression (PLS) and minimizing the sample
entropy [63]. We retained models with low marginal density (MD), high proportions (MD *p-value*)
of retained simulations showing a lower or equal likelihood under the inferred GLM as compared
to the observed genetic data [59], and with high centrality of the observed data within the
multidimensional cloud of retained simulations (Tukey *p-value* [64]). We assessed the ABC's
ability to distinguish between the proposed models, with 1000 pseudo observed data sets
(pods) randomly selected from simulated data sets under each model. All the above-mentioned
methods are detailed in Method S6.

Results

Genetic diversity and structure

The microsatellite population-structure analyses exhibited, at best fitting K of 2-4 (Figs 1, S14-
15), a strong East-West pattern of differentiation (F_{ST} > 0.4, Table S6-7) separating the Atlantic
and Mediterranean populations. This pattern appeared as a continuous cline of differentiation

driving the first component of the PCA (Fig. 1) and was also the first revealed by the DAPC (Fig.
S16). Second, our analyses showed clear segregation of Madeira from Cabo Blanco individuals,
distinguishing the two Atlantic populations (Figs 1, S15-16). The Western-Mediterranean
population formed a group relatively distinct from the Eastern-Mediterranean and was
genetically intermediary between the former and Atlantic populations (Figs 1, S15-16). The only
historical sample genotyped from the Black Sea was assigned to the Eastern-Mediterranean
population (Figs 1, S15). Further subdivision of the MMS resulted in erratic clustering results
(Figs S14-15, S17).
Overall, all the populations showed relatively low nuclear genetic diversity (H_O = 0.12 -
0.42 & H_E = 0.04 - 0.36) and a wide range of inbreeding levels (Fig. 2). Madeira had the lowest
levels of genetic diversity across all estimated indexes and among the highest estimates of
inbreeding (Fig. 2, Tables S6-7). The historical Western-Mediterranean population showed a
relatively high allelic richness, and the Cabo Blanco and Eastern-Mediterranean populations
exhibited the highest number of private alleles (Fig. 2). The historical sample from the Black Sea
did not harbor private alleles (Fig 2). Similarly, mtDNA (CR1) diversity was low in all
populations, with Madeira exhibiting only one haplotype (Table S2, Fig S18).

Figure 2. *Monachus monachus* nuclear genetic diversity.
Boxplot of the microsatellites (**A**) allelic richness (A_R), (**B**) observed (H_O), and (**C**) expected (H_E) heterozygosity, (**D**)
mean private alleles (P_A) of 500 resampling of the smallest sample size across each sampling area, and barplot of
the number of private alleles (P_A), and (**E**) boxplot of the average individual inbreeding coefficient (F), per sampling
area, expressed for modern (>1979) and historical (<1980) samples. In A-D, historic samples were removed from the
analyses for all extant populations apart from the W-Med where no population is clearly identified and for which we
almost exclusively have historic samples. Letters and box colors in A-D illustrate the Tukey posthoc group
assignment. W-Sahara: Cabo Blanco, W-Med: Western Mediterranean Sea, CE-Med: merged Central and Eastern
Mediterranean Sea, C-Med: Central Mediterranean Sea, E-Med: Eastern Mediterranean Sea.

The likelihood estimates of individual homozygosity (F) suggested higher levels of
inbreeding in modern samples than in historical ones (per population; Fig. 2). In addition, the
overall estimates [all historical *vs* all modern samples] of allelic richness (A_R) and private allelic
richness (P_A) were higher historically than in modern samples (Fig. S19). However, the overall
observed (H_O), and expected (H_E) heterozygosity, as well as the overall evolution of inbreeding
levels through time (Fig. S19), did not exhibit a pattern of diversity loss across the timespan

covered by our dataset (1840-2020). Across modern samples (>1975), and for the two major
populations (Cabo Blanco and Eastern-Mediterranean), we could not see any clear pattern of
diversity loss between 1990 and 2020 (Figs S20-21). At the mtDNA level, one historical CR1
haplotype was not recovered in modern samples (MM07) and, reciprocally, one modern
haplotype was not recovered from historical samples (MM06; Figs S18, S22-23).

<mark>In 1997 the Cabo Blanco colony underwent a massive die-off (>⅔ of the population),</mark>
<mark>from different potential causes</mark> [28]. Our results based on various population and individual-
based genetic diversity indexes (A_R, H_O, H_E, F), apart from private alleles (P_A), were not
consistent with the genetic diversity decrease expected after a bottleneck despite a large
sampling before and after the event took place (except for private alleles [PA]; Figs S24-25).
Furthermore, we did not record signals of rare allele loss, or decrease in frequency, nor of major
allele gain in frequency (Fig. S26-27), expected after a reduction of the population size [65,66].

In line with the PCA showing an East-West cline of differentiation, we found an isolation
by distance (IBD) pattern explaining up to 69.6% of the among-individuals genetic distance
variability (Figs 3, S28-29). Such a strong IBD signal was sustained by Mantel correlograms
exhibiting high values for the first 500 km classes, progressively declining up to ~1,500 km,
above which values were no longer significant or negative (Fig. 3). Females exhibited significant
positive values up to higher distance classes (1500 km) than males (700 km), but this signal
may have resulted from sample size differences among males and females for each class.

Figure 3. Isolation by distance in *Monachus monachus*.
A: isolation by distance (IBD) relationship between individual-based genetic and maritime distances. **B** & **C**: Mantel
correlogram of spatial correlation, for 59 classes of 100 Km, in all, modern, and historical samples (**B**), and all, males
and females samples (**C**). Significant values, over 1000 permutations, are represented by filled symbols.
Complementary analyses using alternative genetic distances are presented in Figs S26-27.

Demographic history

The ABC procedure supported models including population structure, ghost populations (local extinctions), and an effective population size decline (Figs. S9-13). The most supported population structure used a custom stepping stone framework, realistically modeling extant and extinct population connectivity, based on their respective position in space (Table S4, S8-9, & Figs S12-13b). Among the models including these features, the most supported ones (M180 & M181, Fig. S13b) suggest that all sampled populations underwent one to two major declines, most of which were of at least one order of magnitude (Table S10, Figs 4, S30). Furthermore, the best models reveal that the declines, and the loss of connectivity among sampled populations, most likely occurred during Antiquity, and at the onset of the Middle-Age (Figs 4, S30-32).

Figure 4. The antique decline of MMS.
ABC-GLM posterior estimates of effective population size (N_e; **A**) and of the time (T_1 and T_2; **B**) of the major demographic event(s) - Population decline + local extinction + change in connectivity - estimated from the most likely scenarios (M181 & M180), within the ABC framework. **A**: mode (symbol) and 50 % quantiles (dotted arrows) of the effective population size (N_e) under the models 181 and 180, represented after (N_0: current population) and before (N_1 and N_2: ancient populations) the demographic events occurring at T_1 and at T_2. **B**: scaled-density distributions of the time of the demographic event(s) T_1 and at T_2. WS: Cabo Blanco, MAD: Madeira, WM: Western-Mediterranean, CM: Central Mediterranean, EM: Eastern-Mediterranean, Other N1 and Other N2: cumulated ancient effective population sizes of other populations before T_1 and T_2 respectively.

Discussion

Genetic structure of Mediterranean monk seal through time and space

Our study is based on an unprecedented genetic sampling, covering the entire distribution range of the Mediterranean monk seal and encompassing historical (extinct) and extant populations. This enabled us to provide the most comprehensive diversity and structure assessment of the species to date, in comparison to previous studies limited in their geographic

and temporal representation and/or genetic information. Our results illustrate how the inclusion
of historical samples helps to understand the dynamics of species' genetic diversity through
time and space [67]. The global analysis showed that MMS are composed of four populations,
including the previously delineated eastern Mediterranean (reaching the Ligurian Sea at its
westernmost location) and two northern Atlantic populations (Madeira and Cabo Blanco) [35–
38,41], and a newly identified, historical population in the Western Mediterranean. The latter
ranged from the northern Maghreb to southern France, Sardinia Isl. and the Adriatic Sea, partly
overlapping at its eastern fringe with the extant eastern Mediterranean population (Fig. 1).
Because the western Mediterranean population holds both North Atlantic and Mediterranean
mitochondrial haplotypes, it could not be clearly delineated in the previous, single-locus studies
integrating historical samples [35,36]. The modern sample from northern Morocco (1993) raises
the long-standing question as to whether the western Mediterranean population still exists
today, notably on the western Algerian coast where MMS were present until the recent past
[68]. Although sightings of MMS across the western Mediterranean basin have been reported
within the last c. 20 yrs [69], their actual relevance is difficult to assess unless proper monitoring
of potential breeding sites is systematized.
The status of the extinct population in the Black Sea [70] remains unsolved. The sole
successfully genotyped historical sample included in the microsatellites analyses did however
carry information attributing it to the eastern Mediterranean population (Figs 1, S15). Together
with the sharing of a unique CR1 haplotype in the Black Sea with individuals from the Aegean
Sea, our and previous results [35,36] do not support the straits of Çanakkale (Dardanelles) and
Istanbul (Bosporus) as a putative barrier to MMS movements, despite its effect on other species
at a wide taxonomic scale, including marine mammals [71,72].
Although substantial differentiation was previously reported between Cabo Blanco and
Madeira [38], Cabo Blanco and Eastern Mediterranean [37], and Aegean and Ionian seas [40],
our results demonstrate that this differentiation is clinal and likely a consequence of isolation by
distance (R^2 = 0.7, Figs 1, 3). At the level of the Mediterranean basin, similar continuous
differentiation is found in several marine organisms with varying mobility [e.g. dolphins, Sea star
[73,74]], suggesting that oceanic distance alone can mold species genetic diversity across a
broad taxonomic range. Because the inclusion of historical samples did not disrupt the genetic
covariation with geography, the isolation by distance pattern might have been long-standing
over the past two centuries. Interestingly, this pattern, previously unveiled at the eastern
Mediterranean Sea scale [40], appears driven by prevailing short distance (< 500 km) and
infrequent long distance dispersal (< 1,500 km, Fig. 3). MMS may tend to establish in proximity
to their place of birth, eventually dispersing within their local water basin, and, on rare
occasions, to further-away places. This is in sharp contrast with the lack of structure of
Neomonachus schauinslandi, the Hawaiian Monk Seal [75], which shows long distance
dispersal across a ~2,700 km linear oceanic distribution. On the contrary, our results suggest
that natal philopatry, exacerbated by the rarity of suitable habitat and breeding sites, and by the
weaker and circular Mediterranean Sea currents, are likely contributors to the current MMS
genetic structure [76,77].

A long depauperate and decreasing genetic diversity

Combining data from all extant populations, our genetic survey confirms that MMS harbors the most depauperate genetic diversity (H_E = 0.04-0.36) of all seal species [29]. Furthermore, it reveals Madeira's dramatic levels of genetic diversity (H_E = 0.04, Fig. 2), which echoes its decline to less than 10 individuals in the 1980s [26]. As a matter of comparison, Madeira's diversity is lower than that of the ringed seal population (*Pusa hispida*) landlocked since the LGM in Saimaa Lake, Finland [78].

We uncovered a decrease of allelic richness (A_R) and private allelic richness (P_A) from historical to modern samples, and an increasing trend in inbreeding at the population level (Figs 2, S19). This pattern is consistent with the trend exhibited by CR1 [35] and mitogenome [36] studies, confirming that MMS genetic diversity has decreased over the past two centuries. In all cases, the decrease in genetic diversity started from remarkably low historical levels, which could explain the relatively stable distribution of heterozygosity that we observed over time (Fig. S19). This contrasts with the sharp drop in diversity caused by commercial exploitation in other seal species [79,80], implying an older exploitation of MMS populations.

Remarkably, the MMS genetic diversity has not been decreasing in the past 30 years (Figs S20-21), at least in the two populations with sufficient temporal sampling (eastern Mediterranean and Cabo Blanco). This encouraging signal can be interpreted as the incipient results of conservation efforts and population recovery over that period [12]. However, this pattern is surprising in the Cabo Blanco, where we could not trace any substantial genetic diversity or minor allele frequency drop - except private allele loss - following the massive 1997 die-off that decimated more than two thirds of the Cabo-Blanco colony (Figs S24-25 [28]). The originally low diversity and high inbreeding in the population, a higher death rate of most inbred individuals at the time of the die-off, or an insufficient resolution of the studied loci, may have blurred declining pattern signatures. Additionally, individuals from unsurveyed nearby localities (e.g. Guerguerat) at the time of the mass die-off may have participated in maintaining the genetic diversity of the Cabo-Blanco colony (pers. obs. AA).

Human-seal interactions in the Mediterranean basin: a history of overexploitation

The low diversity and high levels of inbreeding in all MMS populations (Fig. 2) are strong signals for the species undergoing a major decline with limited gene flow among populations. Furthermore, the marked IBD pattern (Fig. 3) reveals that MMS populations were connected by gene flow in the recent past. These patterns are confirmed by our demographic modeling analyses, which suggest that populations were organized in a stepping-stone manner, connected by intermediary (extinct) populations. These populations underwent one to two major demographic declines during Antiquity [~800BC-600AD], most of which were at least of one order of magnitude (Figs 4, S30).

Surprisingly, the LGM (~20 ka ago) that dramatically affected the sea level (~120-130 m below current level) and the Mediterranean basin area, did not leave an identifiable signature in the MMS genetic data. One of our two best models also shows weak support for a population size decrease during the Neolithic [~12,000 - 800 BC; Fig. 4]. The Neolithic transition allowed rapid human population growth and the development of complex civilizations [30,81], mastering

increasingly sophisticated seafaring and fishing techniques [82,83]. By the rise of Antiquity,
large human populations had spread across the entire Mediterranean basin and its islands [83–
85]. Although the extent of hunting pressure on MMS is hard to accurately gauge at this time,
historical sources relate hunting, meat consumption, oil use in lamps and skin use [86], conflict
with fishermen [87], use in circus shows [88] and use of body parts to produce medicines [86].
In Antiquity, the species was reported to be common, widespread, and of 'naive' behavior
towards humans, with rookeries of large size, and using open environments such as beaches,
outcrops, or promontories [86,89,90]. Such descriptions echo the large ancient effective
population sizes (between 100,000 [N_1-M181] and 356,000 [N_2-M180] individuals, Table S10)
inferred from our demographic reconstructions, and are consistent with the ancestral population
size estimates of *Neomonachus tropicalis* [91]. However, it is in sharp contrast with the current
elusive nature of the species, its habits of resting and giving birth in caves and remote islets, all
of which are probably a result of long-lasting persecution [8].

Whether Romans, as previously posited [10], and/or other ancient civilizations played a
significant role in the abrupt demographic decline of MMS remains unsolved, given the lack of
direct evidence. However, the Roman Empire's large-scale wildlife exploitation that led to local-
fish stock depletion and local megafauna extinction [88,92,93], and the emerging evidence of
their whaling activities [94] are clues pointing towards their potential role in the demise of MMS.
Importantly, humans have been hunting MMS long before Antiquity [20,33,95,96] and have
continued afterward [11], together with increasing competition for habitat use and marine
resources. Therefore, the MMS decline was likely a continuous process, with a peak during
Antiquity that dramatically sealed the genetic impoverishment of the species. Furthermore,
sealed animals were undoubtedly processed on the beach, and this explains that only very few
bones reached inland sites [96]. Thus, systematic sealing (like whaling [94]) is unlikely to have
produced large archeological accumulations, as is the case for smaller marine resources, like
tuna [97]. The resulting paucity of ancient MMS remains [95,96] limits our ability to accurately
conclude a precise period of MMS overexploitation.

Conservation outcomes

The existence of a western Mediterranean MMS population, likely on the verge of extinction,
calls for the urgent identification and protection of breeding areas. Prior sightings suggest that
several sites potentially hosting individuals belonging to this population should be surveyed,
monitored and adequately protected on priority (e.g. from Al Hoceima to Cap des Trois
Fourches - Morocco, Balearic Islands - Spain, Corsica -France, Sardinia, Tuscan Archipelago,
Sicily - Italy, La Galite - Tunisia [69,98]).

The clinal pattern of genetic diversity across the entire species distribution supports the
absence of locally manageable conservation units. It, therefore, calls for an ambitious
integrative conservation plan that comprehensively includes all countries (30) of the MMS past
distribution range. Such concerted actions should be strengthened by governmental and
intergovernmental agencies' efforts and policies [99].

The low genetic diversity of extant populations calls for the development of an ambitious
program that includes (i) increasing the size of colonies and populations and (ii) restoring
genetic diversity and connectivity among extant populations. Ensuring population growth should
be at the core of the MMS conservation program, as it will directly improve the resilience of

populations to stochastic events [28]. Ways of re-establishing dispersion among populations
should be considered, through, for example, the conservation of historical breeding sites (see
[69]).
However, the dramatic levels of genetic diversity reported from our study indicate that
rescuing the genetic diversity of MMS through the translocation of individuals among extant
populations [100,101] is a scientifically-backed option to promote the long-term recovery of the
species and should not be disregarded. Translocation should balance the benefits of
counteracting the present limited gene flow among populations with any potential detrimental
effects, such as the introduction into the receiving MMS community of pathogens to which it had
not been previously exposed.
Furthermore, our results demonstrate that all extant MMS populations were connected
until recently, and that their differentiation is explained by their frequent short distance (< 500
km) and rare long distance dispersal (< 1,500 km). Theory predicts that gene flow reduces local
adaptation [102], and small populations governed by strong genetic drift (e.g. Madeira) are less
likely to have fine-scale local adaptations [103]. Therefore, translocation among inbred nearby
populations presents limited risks of outbreeding depression [104,105]. The success of such
genetic rescue is however conditioned by long-term conservation of extant populations, their
habitat and resources, and actions to re-establishing natural dispersion among populations
[105].
Finally, our broad scale genetic survey provides a new turnkey cost-efficient tool to
accurately trace the origin of vagrant MMS individuals and monitor the genetic diversity of MMS
populations and colonies. The combination of the 19 MMS microsatellites markers with the CR1
mitochondrial sequence may indeed serve not only to select the MMS individuals for potential
translocation, but also to assess translocation success over time [101].

Data accessibility

445 Microsatellite data is available in table S1. All additional data, scripts and materials are available
446 to readers at https://doi.org/10.5281/zenodo.6511499.

Acknowledgments

We thank all the collectors and museums listed in Table S1 for providing access to genetic
samples. We are grateful to Sophie Courjal and the staff of the "Plateau technique - Biologie
moléculaire et microbiologie" at EDB for their assistance during lab work, to P. Kyritsis
[Archipelagos], for his logistic help, to I. Carvalho for early comments on the manuscript and two
anonymous reviewers who significantly helped to improve the manuscript. This work was
funded by the Fondation Prince Albert II de Monaco (project "Génétique de la conservation du
phoque moine de Méditerranée"). The Genotoul bioinformatics (Bioinfo Genotoul) platforms
provided computing resources. JS was supported by PANGO-GO (ANR-17-CE02-0001), and
LABEX TULIP (ANR-10-LABX-0041).

Conflict of interest disclosure

The authors of this article declare that they have no financial conflict of interest with the content of this article.

Authors' contribution

PG, EL and JS designed the study. AA, AAK, PFL, RP, GM, AP, TA, SA, AB, ED, BO, AMT, LMG and PD collected the genetic samples and supplied curated databases. JD and PG did the lab work. JS conducted data analyses. JS and PG drafted a first version of the manuscript. All co-authors participated in the writing of the manuscript and agreed with its last version.

References

1. Koch PL, Barnosky AD. 2006 Late Quaternary extinctions: state of the debate. *Annu. Rev. Ecol. Evol. Syst.* **37**, 215–250.
2. Shafer AB *et al.* 2015 Genomics and the challenging translation into conservation practice. *Trends Ecol. Evol.* **30**, 78–87. (doi:10.1016/j.tree.2014.11.009)
3. Fordham DA, Brook BW, Moritz C, Nogués-Bravo D. 2014 Better forecasts of range dynamics using genetic data. *Trends Ecol. Evol.* **29**, 436–443.
4. Fenderson LE, Kovach AI, Llamas B. 2020 Spatiotemporal landscape genetics: Investigating ecology and evolution through space and time. *Mol. Ecol.* **29**, 218–246.
5. Bicho N, Haws J. 2008 At the land's end: marine resources and the importance of fluctuations in the coastline in the prehistoric hunter–gatherer economy of Portugal. *Quat. Sci. Rev.* **27**, 2166–2175.
6. Davidson AD, Boyer AG, Kim H, Pompa-Mansilla S, Hamilton MJ, Costa DP, Ceballos G, Brown JH. 2012 Drivers and hotspots of extinction risk in marine mammals. *Proc. Natl. Acad. Sci.* **109**, 3395–3400.
7. Hofman CA, Rick TC, Erlandson JM, Reeder-Myers L, Welch AJ, Buckley M. 2018 Collagen fingerprinting and the earliest marine mammal hunting in North America. *Sci. Rep.* **8**, 1–6.
8. Karamanlidis AA, Dendrinos P. 2015 *Monachus monachus*. The IUCN Red List of Threatened Species: e.T13653A117647375.
9. Spalding MD *et al.* 2007 Marine ecoregions of the world: a bioregionalization of coastal and shelf areas. *BioScience* **57**, 573–583.
10. Johnson WM, Lavigne DM. 1999 Monk seals in antiquity. *Mediterr. Monk Seal* , 1–101.
11. González LM. 2015 Prehistoric and historic distributions of the critically endangered Mediterranean monk seal (*Monachus monachus*) in the eastern Atlantic. *Mar. Mammal Sci.* **31**, 1168–1192.
12. Karamanlidis AA, Dendrinos P, De Larrinoa PF, Gücü AC, Johnson WM, Kiraç CO, Pires R. 2016 The Mediterranean monk seal *Monachus monachus*: status, biology, threats, and conservation priorities. *Mammal Rev.* **46**, 92–105.
13. Karamanlidis AA, Adamantopoulou S, Tounta E, Dendrinos D. 2019 *Monachus monachus* Eastern Mediterranean subpopulation. *IUCN Red List Threat. Species 2019* , e.T120868935A120869697. (doi:https://dx.doi.org/10.2305/IUCN.UK.2019-1.RLTS.T120868935A120869697.)

14.   Pires R, Aparicio Garcia F, Pereira S, Cedenilla M, Fernandez De Larrinoa P. 2019
Conservation status of the Mediterranean monk seal *Monachus monachus* population at
Madeira archipelago. Barcelona, Catalonia, Spain.
15.   Hernández M *et al.* 1998 Did algal toxins cause monk seal mortality? *Nature* **393**, 28–29.
16.   Osterhaus A *et al.* 1998 Monk seal mortality: virus or toxin? *Vaccine* **16**, 979–981.
17.   Scheel D-M, Slater GJ, Kolokotronis S-O, Potter CW, Rotstein DS, Tsangaras K,
Greenwood AD, Helgen KM. 2014 Biogeography and taxonomy of extinct and endangered
monk seals illuminated by ancient DNA and skull morphology. *ZooKeys* , 1–33.
18.   CEPF. 2010 Mediterranean basin biodiversity hotspot.
19.   Johnson WM. 2004 Monk seals in post-classical history: The role of the Mediterranean
monk seal (*Monachus monachus*) in European history and culture, from the fall of Rome to
the 20th century. *Mededelingen* **39**, 1–91.
20.   Stringer CB *et al.* 2008 Neanderthal exploitation of marine mammals in Gibraltar. *Proc.*
Natl. Acad. Sci. U. S. A. **105**, 14319–14324.
21.   Trantalidou K. 2011 From Mesolithic fishermen and bird hunters to Neolithic goat herders:
the transformation of an island economy in the Aegean. *Cave Cyclops Mesolith. Neolit.*
Netw. North. Aegean Greece **2**, 53–150.
22.   Brito C. 2012 Portuguese sealing and whaling activities as contributions to understand
early northeast Atlantic environmental history of marine mammals. In *New approaches to*
the study of marine mammals (eds A Romero, EO Keith), pp. 207–222. Rijeka, Croatia:
InTech.
23.   Panou A, Jacobs J, Panos D. 1993 The endangered Mediterranean monk seal *Monachus*
monachus in the Ionian Sea, Greece. *Biol. Conserv.* **64**, 129–140.
24.   Borrell A, Cantos G, Aguilar A, Androukaki E, Dendrinos P. 2007 Concentrations and
patterns of organochlorine pesticides and PCBs in Mediterranean monk seals (*Monachus*
monachus) from Western Sahara and Greece. *Sci. Total Environ.* **381**, 316–325.
25.   Kovacs KM *et al.* 2012 Global threats to pinnipeds. *Mar. Mammal Sci.* **28**, 414–436.
26.   Pires R, Neves HC, Karamanlidis AA. 2008 The critically endangered Mediterranean monk
seal *Monachus monachus* in the archipelago of Madeira: Priorities for conservation. *Oryx*
42, 278–285.
27.   Silva MA, Brito C, Santos SV, Barreiros JP. 2009 Historic and recent occurrences of
pinnipeds in the Archipelago of the Azores. *Mammalia* **73**, 60–62.
28.   Forcada J, Hammond PS, Aguilar A. 1999 Status of the Mediterranean monk seal
Monachus monachus in the western Sahara and the implications of a mass mortality
event. *Mar. Ecol. Prog. Ser.* **188**, 249–261.
29.   Stoffel MA *et al.* 2018 Demographic histories and genetic diversity across pinnipeds are
shaped by human exploitation, ecology and life-history. *Nat. Commun.* **9**, 1–12.
30.   Benjamin J *et al.* 2017 Late Quaternary sea-level changes and early human societies in
the central and eastern Mediterranean Basin: An interdisciplinary review. *Quat. Int.* **449**,
29–57.
31.   Thunell RC. 1979 Eastern Mediterranean Sea during the last glacial maximum; an 18,000-
years BP reconstruction. *Quat. Res.* **11**, 353–372.
32.   Mikolajewicz U. 2011 Modeling Mediterranean ocean climate of the last glacial maximum.
Clim. Past **7**, 161–180.
33.   Colonese AC, Mannino MA, Mayer DB-Y, Fa DA, Finlayson JC, Lubell D, Stiner MC. 2011
Marine mollusc exploitation in Mediterranean prehistory: an overview. *Quat. Int.* **239**, 86–
103.
34.   Cunliffe B. 2017 *On the Ocean: The Mediterranean and the Atlantic from prehistory to AD*
1500. Oxford University Press.
35.   Gaubert P *et al.* 2019 Insights from 180 years of mitochondrial variability in the
endangered Mediterranean monk seal (*Monachus monachus*). *Mar. Mammal Sci.* **35**,

1489–1511.

36. Rey-Iglesia A *et al.* 2021 Mitogenomics of the endangered Mediterranean monk seal
(*Monachus monachus*) reveals dramatic loss of diversity and supports historical gene-flow
between Atlantic and eastern Mediterranean populations. *Zool. J. Linn. Soc.* **191**, 1147–
1159.

37. Pastor T, Garza JC, Aguilar A, Tounta E, Androukaki E. 2007 Genetic diversity and
differentiation between the two remaining populations of the critically endangered
Mediterranean monk seal. *Anim. Conserv.* **10**, 461–469.

38. Dayon J, Lecompte E, Aguilar A, de Larrinoa PF, Pires R, Gaubert P. 2020 Development
and characterization of nineteen microsatellite loci for the endangered Mediterranean
monk seal *Monachus monachus*. *Mar. Biodivers.* **50**, 1–7.

39. Pastor T, Garza JC, Allen P, Amos W, Aguilar A. 2004 Low genetic variability in the highly
endangered Mediterranean monk seal. *J. Hered.* **95**, 291–300.

40. Karamanlidis AA, Skrbinšek T, Amato G, Dendrinos P, Gaughran S, Kasapidis P, Kopatz
A, Stronen AV. 2021 Genetic and demographic history define a conservation strategy for
earth's most endangered pinniped, the Mediterranean monk seal *Monachus monachus*.
Sci. Rep. **11**, 1–10.

41. Karamanlidis AA, Gaughran S, Aguilar A, Dendrinos P, Huber D, Pires R, Schultz J,
Skrbinšek T, Amato G. 2016 Shaping species conservation strategies using mtDNA
analysis: the case of the elusive Mediterranean monk seal (*Monachus monachus*). *Biol.*
Conserv. **193**, 71–79.

42. Hurlbert SH. 1971 The nonconcept of species diversity: a critique and alternative
parameters. *Ecology* **52**, 577–586.

43. Goudet J. 2005 Hierfstat, a package for R to compute and test hierarchical F-statistics.
Mol. Ecol. Notes **5**, 184–186.

44. Nei M. 1978 Estimation of average heterozygosity and genetic distance from a small
number of individuals. *Genetics* **89**, 583–590.

45. Jombart T. 2008 adegenet: a R package for the multivariate analysis of genetic markers.
Bioinformatics **24**, 1403–1405.

46. Nei M. 1973 Analysis of gene diversity in subdivided populations. *Proc. Natl. Acad. Sci. U.*
S. A. **70**, 3321–3323.

47. Tonkin-Hill G, Lees JA, Bentley SD, Frost SD, Corander J. 2019 Fast hierarchical
Bayesian analysis of population structure. *Nucleic Acids Res.* **47**, 5539–5549.

48. Wright S. 1943 Isolation by distance. *Genetics* **28**, 114–138.

49. Slatkin M. 1993 Isolation by distance in equilibrium and non-equilibrium populations.
Evolution **47**, 264–279.

50. Mantel N. 1967 The detection of disease clustering and a generalized regression
approach. *Cancer Res.* **27**, 209–220.

51. Cayuela H, Boualit L, Laporte M, Prunier JG, Preiss F, Laurent A, Foletti F, Clobert J,
Jacob G. 2019 Kin-dependent dispersal influences relatedness and genetic structuring in a
lek system. *Oecologia* **191**, 97–112.

52. Salmona J *et al.* 2021 How ancient forest fragmentation and riparian connectivity generate
high levels of genetic diversity in a micro-endemic Malagasy tree. *BioRxiv*
20201125394544 Ver 4 Peer-Rev. Recomm. Peer Community Evol. Biol.
(doi:https://doi.org/10.1101/2020.11.25.394544)

53. Oden NL, Sokal RR. 1986 Directional autocorrelation: an extension of spatial correlograms
to two dimensions. *Syst. Zool.* **35**, 608–617.

54. Sokal RR. 1986 Spatial data analysis and historical processes. *Data Anal. Inform. IV* , 29–
43.

55. Nei M, Tajima F. 1981 DNA polymorphism detectable by restriction endonucleases.
Genetics **97**, 145–163.

601 56. Nei M. 1987 *Molecular Evolutionary Genetics*. Columbia University Press.

602 57. Paradis E. 2010 pegas: an R package for population genetics with an integrated–modular
603   approach. *Bioinformatics* **26**, 419–420.

604 58. Beaumont MA. 2019 Approximate Bayesian computation. *Annu. Rev. Stat. Its Appl.* **6**,
605   379–403.

606 59. Wegmann D, Leuenberger C, Neuenschwander S, Excoffier L. 2010 ABCtoolbox: a
607   versatile toolkit for approximate Bayesian computations. *BMC Bioinformatics* **11**, 1–7.

608 60. Csilléry K, François O, Blum MG. 2012 abc: an R package for approximate Bayesian
609   computation (ABC). *Methods Ecol. Evol.* **3**, 475–479.

610 61. Nunes MA, Prangle D. 2015 abctools: an R package for tuning approximate Bayesian
611   computation analyses. *R J.* **7**, 189–205.

612 62. Blum MG, Nunes MA, Prangle D, Sisson SA. 2013 A comparative review of dimension
613   reduction methods in approximate Bayesian computation. *Stat. Sci.* **28**, 189–208.

614 63. Nunes MA, Balding DJ. 2010 On optimal selection of summary statistics for approximate
615   Bayesian computation. *Stat. Appl. Genet. Mol. Biol.* **9**.

616 64. Kousathanas A, Duchen P, Wegmann D. 2018 A guide to general-purpose ABC software.
617   In *Handbook of approximate Bayesian computation*, pp. 369–413. Chapman and
618   Hall/CRC.

619 65. Cornuet JM, Luikart G. 1996 Description and power analysis of two tests for detecting
620   recent population bottlenecks from allele frequency data. *Genetics* **144**, 2001–2014.

621 66. Garza JC, Williamson EG. 2001 Detection of reduction in population size using data from
622   microsatellite loci. *Mol. Ecol.* **10**, 305–318.

623 67. Slatkin M, Racimo F. 2016 Ancient DNA and human history. *Proc. Natl. Acad. Sci. U. S. A.*
624   **113**, 6380–6387.

625 68. Boutiba Z, Squabria B, Robineau D. 1988 Etat actuel de la population du phoque-moine
626   (*Monachus monachus*) sur le littoral ouest algérien (région d'Oran). *Mammalia* **52**, 549–
627   556.

628 69. Bundone L, Panou A, Molinaroli E. 2019 On sightings of (vagrant?) monk seals, *Monachus
629   monachus*, in the Mediterranean Basin and their importance for the conservation of the
630   species. *Aquat. Conserv. Mar. Freshw. Ecosyst.* **29**, 554–563.

631 70. Kiraç C, Savaş Y. 1996 Status of the monk seal (*Monachus monachus*) in the
632   neighbourhood of Ereğli, Black Sea coast of Turkey. *Zool. Middle East* **12**, 5–12.

633 71. Kalkan E, Karhan SÜ, Bilgin R, Hemond EM. 2016 The Turkish Straits System as a
634   phylogeographic boundary – a literature review. In *The Sea of Marmara; Marine
635   Biodiversity, Fisheries, Conservation and Governance.* (eds E Özsoy, NM Çağatay, N
636   Balkıs, N Balkis, B Öztürk), pp. 550–569. Istanbul, Turkey: TUDAV Publ.

637 72. Tonay AM *et al.* 2017 Is there a distinct harbor porpoise subpopulation in the Marmara
638   Sea? *Mitochondrial DNA Part A* **28**, 558–564.

639 73. Zulliger DE, Tanner S, Ruch M, Ribi G. 2009 Genetic structure of the high dispersal
640   Atlanto-Mediterranean sea star *Astropecten aranciacus* revealed by mitochondrial DNA
641   sequences and microsatellite loci. *Mar. Biol.* **156**, 597–610.

642 74. Gaspari S, Scheinin A, Holcer D, Fortuna C, Natali C, Genov T, Frantzis A, Chelazzi G,
643   Moura AE. 2015 Drivers of population structure of the bottlenose dolphin (*Tursiops
644   truncatus*) in the Eastern Mediterranean Sea. *Evol. Biol.* **42**, 177–190.

645 75. Schultz JK, Baker JD, Toonen RJ, Harting AL, Bowen BW. 2011 Range-wide genetic
646   connectivity of the Hawaiian monk seal and implications for translocation. *Conserv. Biol.*
647   **25**, 124–132.

648 76. Poulain P-M, Bussani A, Gerin R, Jungwirth R, Mauri E, Menna M, Notarstefano G. 2013
649   Mediterranean surface currents measured with drifters: From basin to subinertial scales.
650   *Oceanography* **26**, 38–47.

651 77. Wren JL, Kobayashi DR, Jia Y, Toonen RJ. 2016 Modeled population connectivity across

652   the Hawaiian archipelago. *PloS One* **11**, e0167626.

653 78. Valtonen M, Palo JU, Aspi J, Ruokonen M, Kunnasranta M, Nyman T. 2014 Causes and
654   consequences of fine-scale population structure in a critically endangered freshwater seal.
655   *BMC Ecol.* **14**, 1–15.

656 79. Weber D, Stewart BS, Garza JC, Lehman N. 2000 An empirical genetic assessment of the
657   severity of the northern elephant seal population bottleneck. *Curr. Biol.* **10**, 1287–1290.

658 80. Cammen KM *et al.* 2018 Genetic diversity from pre-bottleneck to recovery in two sympatric
659   pinniped species in the Northwest Atlantic. *Conserv. Genet.* **19**, 555–569.

660 81. Bocquet-Appel J-P. 2008 Explaining the Neolithic demographic transition. In *The Neolithic
661   demographic transition and its consequences*, pp. 35–55. Springer.

662 82. Zilhão J. 2014 Early prehistoric navigation in the Western Mediterranean: Implications for
663   the Neolithic transition in Iberia and the Maghreb. *Eurasian Prehistory Isl. Archaeol. Orig.
664   Seafar. East. Mediterr.* **11**, 185–200.

665 83. Simmons AH. 2016 *Stone Age sailors: Paleolithic seafaring in the Mediterranean*.
666   Routledge.

667 84. Zeder MA. 2008 Domestication and early agriculture in the Mediterranean Basin: Origins,
668   diffusion, and impact. *Proc. Natl. Acad. Sci. U. S. A.* **105**, 11597–11604.

669 85. Phoca-Cosmetatou N. 2011 *The first Mediterranean islanders: initial occupation and
670   survival strategies*. University of Oxford School of Archaeology.

671 86. Plinius Secundus G. 77 AD*Naturalis Historiae*. 1940th-The Loeb Classical Library edn.
672   Cambridge, Massachusetts,: Harvard University Press.

673 87. Oppianus Corycius. 169 AD*Halieutica*. Harvard University Press.

674 88. Scullard HH. 2010 *From the Gracchi to Nero: a history of Rome 133 BC to AD 68*.
675   Routledge.

676 89. Όμηρος αποσαφήνιση. BC750 *Οδύσσεια (Odyssey)*.

677 90. Αριστοτέλης. BC340 *Τ⬚ν περ⬚ τ⬚ ζ⬚α ⬚στορι⬚ν (Historia animalium)*.

678 91. McClenachan L, Cooper AB. 2008 Extinction rate, historical population structure and
679   ecological role of the Caribbean monk seal. *Proc. R. Soc. B Biol. Sci.* **275**, 1351–1358.

680 92. Trakadas A. 2006 'Exhausted by Fishermen's Nets' Roman Sea Fisheries and their
681   Management. *J. Mediterr. Stud.* **16**, 259–272.

682 93. Kidd E. 2012 "Beast-Hunts" in Roman Amphitheaters: The Impact of the Venationes on
683   Animal Populations in the Ancient Roman World. *Eagle Feather* **9**.

684 94. Bernal-Casasola D. 2018 Whale Hunting in the Strait of Gibraltar during the Roman
685   Period? *SAA Archaeol. Rec.*

686 95. Steele TE, Álvarez-Fernández E. 2011 Initial investigations into the exploitation of coastal
687   resources in North Africa during the Late Pleistocene at Grotte des Contrebandiers,
688   Morocco. In *Trekking the Shore*, pp. 383–403. Springer.

689 96. Morales-Pérez JV, Ripoll MP, Pardo JJ, Álvarez-Fernández E, González AM, Tortosa JA.
690   2019 Mediterranean monk seal hunting in the regional Epipalaeolithic of Southern Iberia. A
691   study of the Nerja Cave site (Málaga, Spain). *Quat. Int.* **515**, 80–91.

692 97. Bernal-Casasola D, Expósito JA, Díaz JJ. 2018 The Baelo Claudia Paradigm: The
693   Exploitation of Marine Resources in Roman Cetariae. *J. Marit. Archaeol.* **13**, 329–351.

694 98. Mo G, Bazairi H, Bayed A, Agnesi S. 2011 Survey on Mediterranean Monk Seal
695   (*Monachus monachus*) Sightings in Mediterranean Morocco. *Aquat. Mamm.* **37**.

696 99. Reilly SB. 2005 Report of the scientific research program under the International Dolphin
697   Conservation Program Act.

698 100. Batson WG, Gordon IJ, Fletcher DB, Manning AD. 2015 Translocation tactics: a framework
699   to support the IUCN Guidelines for wildlife translocations and improve the quality of applied
700   methods. *J. Appl. Ecol.* **52**, 1598–1607.

701 101. Pacioni C, Wayne AF, Page M. 2019 Guidelines for genetic management in mammal
702   translocation programs. *Biol. Conserv.* **237**, 105–113.

703   102. Lenormand T. 2002 Gene flow and the limits to natural selection. *Trends Ecol. Evol.* **17**,
704   183–189.
705   103. Leimu R, Fischer M. 2008 A meta-analysis of local adaptation in plants. *PloS One* **3**,
706   e4010.
707   104. Edmands S. 2007 Between a rock and a hard place: evaluating the relative risks of
708   inbreeding and outbreeding for conservation and management. *Mol. Ecol.* **16**, 463–475.
709   105. Bell DA, Robinson ZL, Funk WC, Fitzpatrick SW, Allendorf FW, Tallmon DA, Whiteley AR.
710   2019 The exciting potential and remaining uncertainties of genetic rescue. *Trends Ecol.*
711   *Evol.* **34**, 1070–1079.